NMFS Honolulu Library
00000000002688

QL625
J
v. 3

JORDAN, DAVID STARR
THE FISHES OF NORT
AND MIDDLE AMERICA.

1896

LIBRARY
BUREAU OF COMMERCIAL FISHERIES
U. S. FISH AND WILDLIFE SERVICE
P. O. BOX 3830
HONOLULU 12, HAWAII

LIBRARY
JUL 9 1964
FISH & WILDLIFE SERVICE



Digitized by the Internet Archive
in 2025

https://archive.org/details/fishes-of-north-and-middle-america_1898_47-part-3

This reprint is the first in a publication program inaugurated pursuant to an agreement between the Smithsonian Institution and T.F.H. Publications, Inc. The proceeds derived from the sale of this and other such reprints will be used to establish the T.F.H. Fund "for research, collection or purchase of fish specimens, explorations, and publication of scientific reports related to aquarium fishes."

This edition is reprinted from the original without change except that the plates have been printed on both sides of a page. The four parts may be collated with the present four volumes as follows:

Part 1. Vol. 1, pp. i-lx, 1-954.
Vol. 2, pp. 955-1240.

Part 2. Vol. 2 (continued) pp. i-xxx, 1241-1936.
Vol. 3, pp. 1937-2183.

Part 3. Vol. 3 (continued) pp. i-xxiv, 2183a-2860.
Vol. 4, pp. 2861-3136.

Part 4. Vol. 4 (continued) pp. i-Ci, 3137-3313, pls. 1-392.

LIBRARY
BUREAU OF COMMERCIAL FISHERIES
U. S. FISH AND WILDLIFE SERVICE
P. O. BOX 3830
HONOLULU 12, HAWAII

QL625
J
V.3
3151 64

THE FISHES OF NORTH AND MIDDLE AMERICA

by

David Starr Jordan

and

Barton Warren Evermann

Reprinted 1963, for the Smithsonian Institution by T.F.H. Publications, Inc., Jersey City 2, New Jersey, U.S.A.

For sale by the Smithsonian Institution, Washington 25, D.C.
Price per set of four volumes, cloth bound, $25.00

2309. CALYCILEPIDOTUS SPINOSUS, Ayres.

Head $2\frac{3}{4}$; depth 5. D. III, VIII, 19; A. 15; scales 60; eye $4\frac{1}{3}$ in head; maxillary $2\frac{1}{3}$; fourth dorsal spine 4; pectoral $1\frac{1}{2}$; ventral 2; caudal $1\frac{2}{3}$. Body rather elongate, depressed; head broad, somewhat concave between the occipital ridges; 2 sharp radiating ridges behind upper posterior margin of each orbit, somewhat broken up into irregular spines; posterior part of supraorbital rim broken up into spines; top of head covered with loose skin, and with thick-set mucous tubes; interorbital space narrow, concave, $\frac{1}{2}$ diameter of eye; preopercle with 2 strong, shortish, diverging spines above; fleshy slips above opercle, near upper posterior part of eye, and at occiput; a long fleshy slip on maxillary, and 4 on lower jaw; many scales on sides with small flaps; skin, where not scaly, thin and lax; dorsal band of scales with about 7 rows at its widest part, anteriorly much wider than the space between it and the lateral band; isthmus rather broad, the membranes not forming a fold across it; dorsal fins considerably connected, spines very low, the highest about $\frac{2}{3}$ the height of the soft rays; pectorals broad, shortish, about reaching vent; distance from spinous dorsal to snout greater than length of pectoral. Color brown, mottled and obscurely barred, often tinged with red; top of head usually with brick red; fins all, except ventrals, mottled with blackish and reddish; skin joining bones of jaws unspotted; belly whitish, immaculate. Here described from a specimen from Monterey Bay, California, 10 inches in length. Coast of California, in rather deep water; not common; known only from about Monterey and San Francisco. (*spinosus*, spiny.)

Calycilepidotus spinosus, Ayres, Proc. Cal. Ac. Sci., I, 1855, 76, San Francisco.
Hemilepidotus spinosus, Girard, Proc. Ac. Nat. Sci. Phila. 1856, 134; Girard, U. S. Pac. R. R. Surv., X, Fish., 68, 1858; Jordan & Gilbert, Synopsis, 715.

727. ENOPHRYS, Swainson.

(Stone Sculpins.)

Enophrys, Swainson, Class'n Fishes, II, 271, 1839 (*claviger*).
Aspicottus, Girard, Proc. Ac. Nat. Sci. Phila. 1854, 130 (*bison*).
Clypeocottus, Ayres, Proc. Cal. Ac. Sci. 1854, 12 (*robustus*).

Body short and thick, depressed anteriorly. Head very large, mailed above with rugose, bony plates; a series of large, rough, bony plates along lateral line; no scales. Teeth in villiform bands in jaws and on vomer, none on palatines; preopercle with strong, straight spines; suborbital stay broad, externally bony; gill membranes joined to the isthmus, not forming a fold across it; a slit behind fourth gill. Dorsal fins separate, the anterior short, not notched; anal short. Intestinal canal elongate. Herbivorous, feeding chiefly on algæ. (*ἐν*, on; *ὀφρύς*, eyebrow.)

Aspicottus (*ασπίς*, shield; *Cottus*):
a. Preopercular spine about $\frac{1}{3}$ head, reaching end of opercle or a little beyond; bony plates of side without keel or spine. bison, 2310.

Enophrys:
aa. Preopercular spine very long, reaching middle of spinous dorsal; bony plates of sides each with a serrated keel or spine. claviger, 2311.

Subgenus ASPICOTTUS, Girard.

2310. ENOPHRYS BISON (Girard).

Head $2\frac{1}{2}$; depth 4. D. VIII, 11; A. 9; lateral plates 29; eye 5 in head; maxillary $2\frac{1}{2}$; pectoral $1\frac{3}{5}$; ventral $2\frac{1}{4}$; caudal 2; snout blunt; maxillary reaching beyond pupil; external bones of head rough granular; interorbital space elevated and concave, the orbital ridge without spine; suborbital stay covering most of cheek; a ridge extending backward from each eye, the two connected by a cross ridge at occiput; the ridges large and rough, and the space between them concave; preopercle with 4 spines, the upper very long, straight, and rough, usually reaching past opercle, a little more than $\frac{1}{3}$ head; opercular ridge very broad; subopercle with 2 diverging spines; a single series of large, rough, granular plates along sides, from opercle to base of caudal, the plates without keel or spine and growing smaller behind; spinous dorsal small, much lower than soft rays; anal short. Olivaceous above, variegated with blackish and reddish, yellowish below; fins olivaceous, marked with black; ventrals pale. Length 12 inches. San Francisco to Sitka; very abundant, especially about Puget Sound; a coarse species, not used for food. When disturbed it extends its long preopercular spines, making the head flat and very wide. Here described from a specimen, 4 inches in length, from Puget Sound, Washington. (*Bison*, the American buffalo, from its horns.)

Aspicottus bison, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 130, Fort Steilacoom (Type, No. 322, U. S. Nat. Mus. Coll. Dr. Suckley); Fort Point (No. 323. Coll. Lieutenant Trowbridge); San Francisco (No. 324. Coll. Dr. Heermann); Tomales Bay (No. 325. Coll. E. Samuels.)

Clypeocottus robustus, AYRES, Proc. Cal. Ac. Sci. 1854, 12, San Francisco.

Aspicottus bison, GIRARD, U. S. Pac. R. R. Surv., x, Fishes, 66, 1858.

Enophrys bison, JORDAN & GILBERT, Synopsis, 710, 1883.

Subgenus ENOPHRYS.

2311. ENOPHRYS CLAVIGER (Cuvier & Valenciennes).

D. VI–13; A. 11; V. I, 3. Top of head everywhere rough; 2 strong nasal spines; superciliary margins much elevated, with a deep groove between them; 4 preopercular spines, the upper exceedingly long and strong, extending nearly to the vertical from end of first dorsal, serrated and coarsely toothed on its upper margin, but without antler-like processes; "occiput with a very long cuneiform process on each side;" lateral line with bony plates, rougher than in *Enophrys bison*, each with a serrated keel and spine; skin subvillous above, with small rough warts; a series of small cutaneous appendages above the anal; suborbital stay spinous; vomerine teeth present; isthmus broad; slit behind last gill large. Dark brown, with 3 or 4 vertical bands; belly white. Length $2\frac{1}{2}$ inches. Bering Sea. Here described from the original type in the British Museum.

Dr. Gilbert mentions also: A single specimen, 52 mm. long, from Albatross Station 3233, Bristol Bay, Alaska; depth $7\frac{1}{4}$ fathoms. The upper preopercular spine is long and very slender, extending to below middle

of spinous dorsal, bearing small serrations on anterior and posterior edges of basal half, but without larger teeth or accessory spinules. Area above lateral plates thickly beset with minute prickles; posterior part of abdominal region and the area above anal fin similarly beset with prickles which scarcely project beyond the small tubercles in which they occur. A few larger postaxial prickles and a small number of white filaments scattered along middle of sides. Filaments also at base of preopercular spine and 2 or 3 at tip of maxillary. Dorsal VIII, 14; anal 11; lateral line 35.

A specimen, 151 mm. long, from *Albatross* Station 3645, off Robben Island, in 10 fathoms, is entirely similar to the one reported on by Gilbert. It may be described as follows: Preorbital with 2 strong spinous projections, which overlap the premaxillary in closed mouth. Interorbital space deeply channeled, the orbital rim raised posteriorly into a blunt spinous tubercle. A small, spinous, occipital tubercle, behind which rises a high, sharp nuchal ridge, which is highest posteriorly and has its upper edge finely toothed. No cirri on top of head. Upper preopercular spine long, simple, reaching beyond head to fourth or fifth plate of lateral line; below it 3 short, strong spines, the lowermost directed downward and forward; outer surface of the upper spine with 3 or 4 low, finely serrated ridges, its inner edge smooth, without accessory cusps or spinules; opercular ridge high, serrate; two sharp spines on anterior angle of subopercle; top and sides of head rough, with minute spinous points; preopercle and lower jaw with numerous short filaments, a longer one on end of maxillary. Body entirely covered with minute prickles, which invest also the abdominal region; those above lateral line longest and most thickly placed; lateral line with a series of plates similar to those in *E. bison*, each surmounted by a sharp spine; lateral line with 2 curves, approaching back most nearly at end of spinous and at end of soft dorsal; many conspicuous white filaments scattered over sides below lateral line. Dorsals entirely separate, the free interspace as wide as pupil. Head $2\frac{2}{3}$ in length; depth $3\frac{3}{5}$. Eye larger than interorbital width, $4\frac{1}{4}$ in head. D. VIII–14; A. 12 (11 in previously noted specimen); P. 16; lateral line with 35 plates. Dusky above, with faint, darker cross bars; light below; two black blotches on cheek; some faint dusky V-shaped prolongations of the coloration of the back down toward base of anal fin; fins indistinctly cross-banded; a dark area at base of pectoral, a narrow oblique dusky cross bar on base of caudal fin. (*clavis*, key; *gero*, I bear.)

Cottus claviger, CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 195, 1829, Kamchatka (Coll. M. Collée); GÜNTHER, Cat., II, 167.

Cottus elegans, GRAY, in CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 195, 1829; same type.

Enophrys claviger, JORDAN & GILBERT, Synopsis, 711, 1883; GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 426; JORDAN & GILBERT, Fishes Bering Sea, 1897.

728. CERATOCOTTUS, Gill.

Ceratocottus, GILL, Proc. Ac. Nat. Sci. Phila. 1859, 165 (*diceraus*).

This genus is very close to *Enophrys*, from which it differs in having the long preopercular spine armed above with recurved hooks or antler-like processes. Bering Sea. (κέρας, horn; *Cottus*.)

a. Transverse occipital ridge not developed; interorbital groove deep and narrow; preopercular spines small. LUCASI, 2312.

aa. Transverse occipital ridge well developed; interorbital groove less deep and wider; preopercular spines larger. DICERAUS, 2313.

2312. CERATOCOTTUS LUCASI, Jordan & Gilbert, new species.

Head $2\frac{4}{5}$ in length, measured to end of opercular flap; depth $3\frac{1}{5}$. D. VII, 13; A. 12; P. 17 or 18. Eye $4\frac{3}{4}$ in head; maxillary $2\frac{1}{2}$; preopercular spine $2\frac{1}{2}$; pectoral fin $1\frac{1}{2}$. Maxillary reaching nearly to the vertical from posterior margin of eye; villiform teeth on jaws and vomer, the outer series in jaws enlarged; no teeth on palatines; interorbital space very deeply channeled, its least width $\frac{2}{3}$ diameter of eye; upper edge of orbital rim sharp and beset with a single series of small teeth; occiput abruptly depressed behind the eyes to below the floor of interorbital space, as in *C. diceraus*, nearly flat both transversely and longitudinally, the occiput ridges being very low, and the transverse ridge so conspicuous at back of occiput in adult *C. diceraus* is here not developed; nuchal ridges high, elevated, and compressed into a minutely serrated edge, a very small cusp-like elevation at their base anteriorly; nasal and preorbital spines as in *C. diceraus;* upper preopercular spines slender and somewhat decurved toward tips, minutely roughened on the outer surface, and bearing on their upper edge 3 retrorsely hooked spines, resembling the spines on a rose bush; below this 3 short, strong spines, the first immediately below the upper spine and diverging from it, the second directed nearly vertically downward, the third downward and forward; 2 strong diverging spines at anterior angle of subopercle; opercular ridge elevated; all exposed bones of head roughened with radiating series of lines which are beset with minute prickles; lateral line with 36 bony plates decreasing in size posteriorly, each plate bearing small slender spines, those on the middle of plate longer than the others and directed backward. From fragments of skin left on snout and side of head in 1 specimen, it is evident that this species is colored much as in *C. diceraus*, the ground color light olive, thickly covered with small dusky spots, around which the ground color forms narrow reticulating lines. When taken the bones of the head were a bright vitriol green. Bering Sea; 2 specimens, 135 and 132 mm. long; 1 taken from the stomach of a cod, the other from a halibut, both of which were captured near St. Paul Island. The skin is digested off from both specimens, so that the color can not be determined, but they are otherwise in good condition. Differing from *C. diceraus* in the deeper narrower interorbital groove and the smaller size and different armature of the preopercular spines. (Named for Mr. Frederick Augustus Lucas, Curator of Comparative Anatomy in the U. S. National Museum and member of the U. S. Fur Seal Commission in 1896 and 1897.

Ceratocottus lucasi, JORDAN & GILBERT MS., Fishes Bering Sea, 1896, 1897 MS., St. Paul Island (Type, No. 48234; Cotype, No. 5661, L. S. Jr. Univ. Mus. Coll. Jordan.)

2313. CERATOCOTTUS DICERAUS (Pallas).

D. VII, 14; A. 10; C. 12; P. 17; V. I, 3. Form of *Enophrys bison*, the head large, wider than deep or long; top of head nearly as in *Enophrys bison*, the ridges higher and very rough; orbital ridges elevated, continued

backward toward the nape, the occipital ridges sharp behind; upper preopercular spine very long, rough, nearly $\frac{2}{3}$ as long as head, with strong recurved hooks or serrations on the upper edge; lower preopercular spines strong; opercle with a longitudinal rib and no distinct spine; lateral line with a row of rough bony scutella, each with a minute central spine; skin above more or less villous or prickly, elsewhere smooth. Isthmus wide; a slit behind last gill; vomer with teeth. Bering Sea, rather common from Alaska to Kamchatka and Saghalin. Here described from a specimen in the British Museum, about 6 inches long.

Dr. Gilbert has this note on several adults taken with seine at Herendeen Bay, Alaska Peninsula:

There are 6 or 7 strong barbs inclined forward on upper surface of preopercular spine, which is very long, slender, and straight, reaching to below middle of spinous dorsal. Filaments arranged as in *Enophrys claviger*, but those on posterior part of body much more numerous. Anal papilla very large, 0.27 mm. in a specimen 195 mm. long. There seems to be no prickles on region above lateral line, or on abdomen. Stomachs filled with molluscan shells, mainly limpets. D. VIII. 13 or 14; A. 10 to 12.

One specimen from Robben Island, collected by Mr. Barrett-Hamilton, and 3 young specimens from Petropaulski, show the following characters: The depth of the occipital depression, the height of the different ridges and spines, and the amount of the irregularity in the cusps of the preopercular spine is subject to great individual variation. We find no important differences between this specimen and those from the Alaskan Peninsula reported on by Gilbert (Report Commissioner of Fish and Fisheries, 1896, 426). The occipital depression is less and the cross ridge behind it much lower in the adult from Robben Island, the occiput comparatively flat and without cross ridge in the young. We do not venture to base any distinctions on these differences which may be due in part to age and in part to individual variation. (*δίς*, two; *κεραός*, horned.)

Cottus diceraus, PALLAS, Nov. Act. Petropol. 1783, 354, pl. 10, fig. 7, Petropaulski, Kamchatka (Coll. Steller); CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 189, 1829; GÜNTHER, Cat., II, 189.

Synauceia cervus, TILESIUS, Mém. Ac. Pertersb., III, 1811, 278, pl. 13, Petropaulski. (Coll. Steller.)

Cottus stelleri, BLOCH & SCHNEIDER, Syst. Ichth., 63, 1801; after STELLER.

Ceratocottus diceraus, GILL, Proc. Ac. Nat. Sci. Phila. 1859, 165, and 1861, 167.

Enophrys diceraus, JORDAN & GILBERT, Synopsis, 711, 1883; GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 426.

729. COTTUS (Artedi) Linnæus.

(MILLER'S THUMBS.)

Cottus, ARTEDI, Genera Piscium, 49, 1738.

Cottus, LINNÆUS, Syst. Nat., X, 264, 1758 (*gobio* *).

Pegedictis, RAFINESQUE, Ichth. Ohiensis, 85, 1820 (*ictalops*).

* *Cottus gobio*, the common Miller's Thumb of Europe, belongs to the same group as *Cottus pollicaris*, *philonips*, etc., having no palatine teeth, the skin smooth, and the ventral rays I, 4. The rays are D. VI to VIII, 16 or 17; A. 12. The name *Cottus* is restricted to this type by Cuvier & Valenciennes, who remark, "Ce genre (*Cottus*) avait pour type primitive, un petit acanthopterygien de nos rivières à tête large," etc.

Cottopsis, GIRARD, Proc. Bost. Soc. Nat. Hist., III, 1850, 303 (*asper*).
Potamocottus, GILL, Proc. Bost. Soc. Nat. Hist., VIII, 1861, 40 (*punctulatus*).
Tauridea, JORDAN & RICE, Man. Vert. E. U. S., Ed. 2, 255, 1878 (*ricei*).

Fresh water sculpins. Body fusiform. Head feebly armed; skin smooth or more or less velvety, its prickles, if present, not bony or scale-like; villiform teeth on jaws and vomer, and sometimes on palatines. Gill openings separated by a wide isthmus, over which the membranes do not form a fold; no slit behind fourth gill. Branchiostegals 6. Dorsals nearly or quite separate, the first of 6 to 9 slender spines; ventrals moderate, each with a short concealed spine and 4 soft rays. Lateral line present, usually more or less chain-like, sometimes incomplete. Preopercle with a simple spine at its angle which is usually curved upward, its base more or less covered by skin, very rarely obsolete; usually 2 or 3 spines turned downward below this; subopercle usually with a concave spine turned downward. Vertebræ 10+23=33; pyloric cæca about 4. Fishes of small size, inhabiting clear waters in the northern parts of Europe, Asia, and America. The species are extremely numerous, and are very difficult to distinguish, all being very similar in form, coloration, and habits. It is probable that these are descended from some Asiatic marine type, as *Trachidermus* (*Centridermichthys*) rather than from *Myoxocephalus*. In this case *Cottus asper* would approach more nearly to the ancestral form. From it are descended *Cottus semiscaber* and *Cottus ictalops*, with their multitude of variations. *Cottus gobio, aleuticus, philonips*, and *pollicaris* indicate the possible descent of *Uranidea*. In like manner *Triglopsis* seems to be descended through *Oncocottus* from *Myoxocephalus*. The Miller's Thumb, or Blob, is found in most streams and lakes where trout occur, and it is one of the most destructive enemies of the trout, devouring its eggs in great numbers. (*Cottus; κόττος*, an old name of the European Miller's Thumb, *Cottus gobio*, Linnæus, from *κόττα* head.)

a. Palatine bones each with a band of teeth (these rarely few in number or wholly absent).
PEGEDICTIS (*πηγή*, fountain; *ἰχθύς*, fish):
b. Preopercular spine short, hooked upward, partially or wholly concealed by skin; 2 or 3 smaller spines directed downward below it.
c. Anal rays 15 to 20; back and sides smooth or variously prickly.
d. Body robust; head rounded anteriorly.
e. Vent midway between base of caudal and tip of snout.
ASPER, 2314.
ee. Vent nearer base of caudal than tip of snout.
GULOSUS, 2315.
dd. Body long and slender; head narrowing rapidly forward, the snout acutely rounded. EVERMANNI, 2316.
cc. Anal rays 11 to 15.
f. Head narrowed anteriorly, the snout long; lateral line complete; back and sides coarsely prickly, sometimes smooth.
RHOTHEUS, 2317.
ff. Head more or less blunt and rounded in outline anteriorly; the snout short.
g. Anal rays 13 to 15, usually 15; caudal peduncle stout; skin smooth. SHASTA, 2318.

gg. Anal rays 11 to 13, usually 13.

h. Caudal peduncle very slender; its least depth not much greater than diameter of eye; body and head profusely speckled. PUNCTULATUS, 2319.

hh. Caudal peduncle deep, its least depth equal to length of snout; back and sides less distinctly speckled.

i. Head blunt, low, rounded anteriorly; body with vague dark clouds or specks, not very distinctly barred; skin smooth or variously prickly, sometimes very rough, sometimes perfectly smooth. SEMISCABER, 2320.

ii. Head less rounded, with a median depression; body usually with broad, oblique, dark bars; axil usually with some fine prickles, the back smooth. ICTALOPS, 2321.

TAURIDEA (*ταύρα*, cow; *εἶδος*, appearance):

bb. Preopercular spine very large, as large as eye, and spirally hooked; back with stiffish prickles.

j. Dorsal VIII, 17; anal 12; head very broad and flat; body olivaceous, finely speckled and motled. BICEI, 2322.

COTTUS:

aa. Palatine teeth wanting (rarely a few developed in males).

k. Preopercle with a spine at its angle, below which are 2 others directed downward.

l. Back and sides more or less prickly; dorsals connected; lateral line complete; preopercle with a long claw-like spine; dorsal VIII, 17; anal 13. ONYCHUS, 2323.

ll. Back and sides smooth or prickly behind axil only.

m. Preopercular spine long, curved upward in a spiral; dorsal fins low, connected; dorsals VII–19; anal 13. POLLICARIS, 2324.

mm. Preopercular spine curved upward; pectoral as long as head; dorsal VIII, 18; anal 14. COGNATUS, 2325.

mmm. Preopercular spine very short, hooked upward; dorsals well connected, the spines low, the soft rays high; lateral line incomplete. Lower preopercular spines well developed; D. VII, 21; A. 15. PERPLEXUS, 2326.

kk. Preopercle with a single short spine at its tip only, and no distinct spines below it, the lower spines reduced to angles of the bone.

n. Posterior nostrils in short, but conspicuous tubes;* dorsals connected; skin smooth or nearly so; dorsal VII to X, 18 or 19; anal 13 or 14.

o. Lateral line very incomplete. KLAMATHENSIS, 2327.

oo. Lateral line complete or nearly so.

p. Dorsal spines IX or X. ALEUTICUS, 2328.

pp. Dorsal spines VII. MINUTUS, 2329.

nn. Posterior nostrils without tubes; skin nearly smooth; D. VII or VIII, 16 to 18; anal 12 to 14.

q. Head moderate, $3\frac{1}{2}$ to 4 in length.

r. Mouth rather large, the maxillary $2\frac{2}{3}$ in head, reaching to opposite middle of eye.

s. Depth $4\frac{1}{2}$ to 5 in length. BELDINGII, 2330.

ss. Depth 6 in length. PHILONIPS, 2331.

rr. Mouth very small, the maxillary $3\frac{2}{3}$ in head, reaching front of pupil. ANNÆ, 2332.

qq. Head larger, $3\frac{1}{4}$ in length; depth 5; body much spotted. SPILOTUS, 2333.

* Not described in *Cottus minutus*.

kkk. Preopercle without spine, its edge covered by smooth skin, a trace of the upper spine rarely present; skin smooth.
t. Dorsals separate. LEIOPOMUS, 2334.
tt. Dorsals connected, long and low, a shallow notch between spinous and soft parts. PRINCEPS, 2335.

Subgenus **PEGEDICTIS**, Rafinesque.

2314. COTTUS ASPER, Richardson.

(PRICKLY BULLHEAD.)

Head 3¼ to 3⅙; depth 4½. D. IX or X, 19 to 21; A. 17 or 18, rarely 15 or 16; eye 5 in head; maxillary 2½; pectoral 1⅓; ventral 2; caudal 1¾. Body deepest at shoulders; caudal peduncle moderately slender, its depth a little less than length of snout. Head wedge-shaped as viewed from above; mouth quite at lower profile of head; maxillary to below posterior margin of pupil; palatine teeth in a very narrow band on front of palatines only; preopercular spine long and sharp; interorbital rather wide, about equal to vertical diameter of eye; pectoral barely reaching front of anal; ventrals not reaching vent; dorsals slightly connected at base. Prickles coarse and stiff, not closely crowded; lateral line complete; vent in the middle of the body in the large majority of specimens, sometimes slightly nearly tail. Color grayish olive, much spotted and mottled with blackish; all fins, except anal and ventrals, with dark wavy cross lines. The coarser prickles, more anterior position of vent, and flatter interorbital space separate *Cottus asper*, at least subspecifically, from the Sacramento River form, *Cottus gulosus*. Streams of the Cascade Range from Vancouver Island to Oregon, abundant in cold streams; extremely variable, almost every variation in the roughness of the skin being found. Length a foot; the largest species of *Cottus*. Here described from a specimen from Walla Walla.

Dr. Gilbert notes:

Five specimens taken in a small stream emptying into Departure Bay, Vancouver Island. The head is naked in all of these, and the prickles absent on belly, along bases of spinous dorsal and anal fin, and on caudal peduncle. D. VIII or IX, 21 or 22; A. 16 to 18.

Still other specimens from Vancouver Island and about Port Townsend have a band of small prickles extending along lateral line to middle of second dorsal. (*asper*, rough.)

Cottus asper, RICHARDSON, Fauna Bor.-Amer., Fish., 295, 1836, Columbia River at Fort Vancouver. (Coll. Dr. Gairdner.)
Trachidermis richardsoni, HECKEL, Ann. Wiener Mus. 1840, 162, Columbia River; after RICHARDSON.
Centridermichthys asper, RICHARDSON, Voyage Sulphur, Ichth., 74, 1845; GÜNTHER, Cat., II, 170.
Cottopsis asper, GIRARD, U. S. Pac. R. R. Surv., X, Fish., 51, 1858.
Uranidea aspera, JORDAN & GILBERT, Synopsis, 694, 1883.

2315. COTTUS GULOSUS (Girard).

Head 3; depth 4½; D. VIII or IX, 19 to 21; A. 16 to 18; eye 5 to 6 in head; maxillary 2½; pectoral 1⅓; caudal 1⅓. Body not much compressed, the

caudal peduncle about as wide as length of snout, head broadly rounded at snout as viewed from above; mouth large, the maxillary reaching to the posterior margin of pupil; bands of palatine teeth variable in specimens from different localities, varying from a moderate band to a wide one; interorbital equal to width of eye; preopercular spine rather sharp and long. Pectorals reaching to front of anal; ventrals not reaching to vent; dorsals slightly connected at base. Skin prickly or more or less smooth, the prickles, when developed, small, crowded, and flexible. Vent nearer base of caudal than tip of snout. Color grayish olive, mottled with darker; ventrals light, anal light or slightly dusky; other fins crossed with wavy lines of light and dark. Streams of the Coast Range in California south to Point Concepcion, very common; the back and sides usually closely prickly (*parvus*), but the skin often wholly or partly smooth (*gulosus*), the smooth specimens usually with axillary prickles; the prickly form more common coastwise, the other perhaps more abundant in streams of the interior, the two not distinguishable by any permanent character. Here described from specimens from San Francisquito Creek, Santa Clara County, California, 3 to 7 inches in length. (*gulosus*, big-mouthed.)

Cottopsis gulosus, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 129, San Mateo Creek * (Coll. R. D. Cutts. Type, No. 290, U. S. Nat. Mus.); San Joaquin River (Coll. Dr. Heermann, No. 291, specimens prickly in axils only); GIRARD, U. S. Pac. R. R. Surv., x, Fish., 53, 1858.

Cottopsis parvus, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 54, Monterey, Presidio, Fort Reading, Petaluma; prickly specimens.

Cottus semiscabra centropleura, EIGENMANN & EIGENMANN, West Amer. Scientist, Vol. VI, No. 49, November, 1889, 149, Allen Springs, Lake Co., Cal. (Coll. D. Cleveland); specimens with sides prickly.

Centridermichthys gulosus, GÜNTHER, Cat., II, 170.

Uranidea gulosa, JORDAN & GILBERT, Synopsis, 695, 1883.

2316. COTTUS EVERMANNI, Gilbert.

Head $3\frac{2}{5}$ in length; depth 5; depth of caudal peduncle $2\frac{4}{5}$ in greatest depth. D. VII, 21; A. 18; P. 16; V. I, 4. Head small, depressed, narrowing rapidly forward, the snout more acutely rounded than in *C. punctulatus*. Mouth with distinct lateral cleft, the maxillary reaching a vertical immediately in advance of pupil, $2\frac{4}{5}$ in head. Mandible slightly protruding. Teeth in narrow bands on jaws, vomer, and palatines, the latter very weak, apparently concealed in part beneath the skin. Total interorbital width about $\frac{2}{3}$ diameter of eye, shallowly concave; occipital area flat or gently convex. Eye small, $1\frac{1}{6}$ in snout, 5 in head. Pores on head unusually large, the most conspicuous occurring on suborbital ring, along mandible and preopercle, and in a horizontal line above opercle; 3 pores form a straight transverse line behind the orbit; a short nasal tube; upper preopercular spine represented by a short triangular process, the margin of the bone below it being smoothly rounded. Spinous dorsal short and comparatively very high, the longest spine slightly more than

* The young individual from Upper Pitt River (Coll. Dr. J. S. Newberry), mentioned by Dr. Girard, probably belongs to *Cottus shasta*.

$\frac{2}{3}$ the longest soft ray; last spine higher than the first and about $\frac{2}{3}$ the longest, the least height of the membrane joining last spine to first soft ray exceeding length of snout; longest ray of soft dorsal slightly more than $\frac{1}{2}$ head; all the rays of dorsal and anal fins simple, unbranched; caudal long and narrow, nearly truncate when spread, $\frac{6}{7}$ length of head; 9 caudal rays are branched at tip for about $\frac{1}{3}$ length of rays. The pectoral reaches the vertical from fourth ray of soft dorsal, the upper ray simple, the next 6 or 7 forked, the remaining rays being thickened with incised membranes; ventrals with 1 spine and 4 rays, not reaching vent, $1\frac{3}{4}$ in head. Lateral line conspicuous anteriorly, running high, interrupted under eleventh or twelfth ray of soft dorsal, a mere trace visible thence to base of caudal. Sides of body thickly covered with coarse prickles, the head, the breast, belly, and a narrow strip along base of anal fin naked. Color light brownish, faintly vermiculated with darker, with traces of 5 irregular cross bars from back, and a narrow distinct bar at base of caudal; pectorals, dorsal, and caudal cross-barred. Characterized by the long slender body entirely covered with coarse prickles, the short spinous dorsal very broadly united to the very long soft dorsal, the long anal fin, the incomplete lateral line, the very large pores on head, the branched pectoral rays, and the absence of any distinctly projecting preopercular spine. One specimen, $2\frac{1}{2}$ inches long, from Lost River, near Klamath Falls, Oregon. (Named for Barton Warren Evermann.)

Cottus evermanni, GILBERT, Bull. U. S. Fish Comm. 1897, 11, with figure, Lost River, Lostine, Oregon. (Type, No. 48228. Coll. Gilbert, Cramer, and Otaki.)

2317. COTTUS RHOTHEUS, Rosa Smith.

Head $2\frac{5}{6}$; depth $4\frac{1}{2}$. D. VII or VIII-16 or 17; A. 11 or 12; lateral line 32 to 34; eye $5\frac{1}{2}$ in head; maxillary $2\frac{1}{3}$; pectoral $1\frac{1}{2}$; caudal $1\frac{5}{6}$. Body rather deep at shoulders, tapering into a slender caudal peduncle; head pointed, somewhat wedge-shaped; mouth large, horizontal, at lower profile of head; maxillary reaching to posterior margin of pupil; palatine band of teeth very broad and long; interorbital concave, about as wide as eye, preopercle spine sharp, with 2 smaller ones below it. Pectorals reaching to front of anal; ventrals not reaching vent; dorsals scarcely connected; lateral line complete. Skin of head smooth, that of back and sides usually rough, with short coarse prickles, shorter and stiffer than in *Cottus asper*. Color dark gray, spotted and mottled with blackish; all fins, except ventrals, mottled with blackish; belly white in the smaller specimens, dusky, with small black points, in the larger ones. Columbia River Basin, generally common; a well marked species. Here described from specimens from Hangman Creek, Tekoa, Washington, from 3 to 4 inches in length.

Concerning this species Gilbert & Evermann remark: "This strongly marked species is abundant in the Spokane region, and was taken at the following stations: Little Spokane River at Dart's Mill near Spokane, and at Chattaroy, Washington; Columbia River at Colville, Washington; Cœur d'Alene Lake near Cœur d'Alene, Idaho; Clearwater River near

Lewiston, Idaho; Walla Walla River at Wallula, Oregon; Hangman Creek at Tekoa, Washington; Natchess River at North Yakima, Washington; Newaukum River near Chehalis, Washington; Snoqualmie River at Snoqualmie Falls, Washington."

The salient features of this species are: (1) the pointed, wedge-shaped profile of head, as viewed from above, this contrasting strongly with the usually broadly rounded contour of other species; (2) the wide horizontal mouth, quite at lower profile of head; (3) the noticeably concave interorbital and occipital regions; (4) the very broad and long palatine band of teeth; (5) the rather slender body and the extremely slender caudal peduncle, the latter expanding fan-like at base of caudal fin; (6) the dorsals usually separate; when united, at extreme base only; (7) lateral line complete; (8) sides usually well invested with prickles, which are triangular and coarse, and less closely placed than in *asper*. They are arranged more or less definitely in oblique series. There is considerable variation in the completeness of the investment, and in one specimen from Chehalis an axillary patch only is present.

The specimens from Chattaroy and from Snoqualmie Falls are referred to this species with doubt as to their identity. The following table will give an idea of the amount of variation in number of fin rays found in this species:

Locality.	Spinous D.		Soft D.		Anal.			Pectoral.		Lateral line.
	VII.	VIII.	16	17	11	12	13	15	16	
Little Spokane River	4	6	6	4	3	6	1	....	10	32 to 34
Natchess River	.....	2	2	....	....	1	1	....	2	32 to 33
Newaukum River	.....	2	1	1	1	....	1	2	.....	.....
Walla Walla River	1	2	1	2	1	2	....	....	3	.....
Colville River	.....	1	1	....	....	1	....	1	.....	.....
Tekoa	1	5	5	1	1	4	1	....	.....	.....
Cœur d'Alene	1	4	3	2	3	2	....	....	5	.....
Lewiston	.....	1	1	....	....	1	....	....	1	.....

(*ῥόθος, ῥοθιάς*, rushing of the torrent.)

Cottus rhotheus, Rosa Smith (Mrs. Eigenmann), Proc. U. S. Nat. Mus. 1882, 347, Spokane Falls, Washington. (Type, No. 30737.)

Uranidea rhothea, Jordan & Gilbert, Synopsis, 953, 1883.

2318. COTTUS SHASTA, Jordan & Starks.

Head 3 to $3\frac{1}{3}$; depth $4\frac{1}{2}$. D. VIII or IX, 17 or 18; A. 13 to 15; eye nearly 5 in head; maxillary $2\frac{1}{7}$; third or fourth dorsal spine $3\frac{3}{7}$; highest soft ray about 2; pectoral $1\frac{1}{10}$; caudal $1\frac{1}{4}$. Body not much compressed; caudal peduncle rather wide, about equal to snout; mouth rather large, the maxillary reaching to posterior margin of pupil; teeth in a moderate band on jaws and vomer, in an exceedingly narrow band on front of palatines; interorbital space not much over $\frac{1}{2}$ eye; upper preopercular spine short,

not much hooked up and not very sharp, a shallow concave space between it and the second, scarcely a notch; the second small and sharp, the third but slightly developed; pectoral reaching to below the fourth ray of soft dorsal; ventral not reaching the vent; dorsals scarcely connected, the soft dorsal high, the highest rays equal to eye and snout; vent slightly nearer tail than tip of snout. Skin smooth, except a few scattered prickles under pectorals. Color very dark brown or blackish; sides mottled; top of head uniform blackish; all the fins more or less mottled, ventrals white or dusky. The following is the fin formula of 4 specimens: Dorsal, IX, 17; VIII, 18; IX, 17; IX, 18. Anal, 14; 15; 15; 13. Upper Sacramento Basin about Mount Shasta. Here described from 4 specimens from McCloud River, at Baird, Shasta County, California, about 4 inches in length. The species is very close to *Cottus semiscaber*, but it has a longer anal, which in turn is shorter than in *Cottus gulosus*. (Named for Mount Shasta.)

Cottus shasta, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1896, 224, McCloud River, at Baird, California. (Coll. E. C. Starks. Type, No. 4196, L. S. Jr. Univ. Mus.)

2319. COTTUS PUNCTULATUS (Gill).

Head 3; depth 4⅔. D. VII or VIII, 17 or 18; A. 11 or 12; eye 6 in head; maxillary 1⅔; third dorsal spine 3; highest ray of soft dorsal 2½; highest anal ray 2¼; pectoral 1½; ventral 1⅓; caudal 1¾. Body tapering into a slender caudal peduncle which is not much wider than eye; teeth in wide bands on jaws, vomers, and palatines; interorbital space rather wide, about equal to eye; maxillary reaching to posterior margin of eye; preopercular spine rather sharp; vent slightly nearer tail than tip of snout. Pectoral reaching to below base of fourth ray of soft dorsal; ventrals not reaching vent; dorsals scarcely connected, the soft dorsal extending farther back than anal; skin in these specimens entirely smooth. Color olivaceous, everywhere punctulate with black spots, more conspicuous on top of head; tip of snout and lower jaw dark, dusky with dark points under lower jaw; dorsals, pectorals, and caudal with wavy streaks and series of spots; anal and ventrals white, anal sometimes dusky; 4 or 5 dark blotches on back indicating cross bars; a dark bar at base of caudal. Length 4 or 5 inches. Headwaters of Green River, Wyoming. Here described from many specimens from Green River, Wyoming, collected by Evermann & Rutter. This species will prove to be a local or desert variation of the widely distributed *Cottus semiscaber*, from which it differs only in the slender caudal peduncle and the speckled coloration. The name *punctulatus* has priority over *semiscaber*, if the forms are to be united, but in this case it would hardly be worth while to separate either from *Cottus ictalops*.

Gilbert & Evermann refer to the specimens here described as—

Numerous specimens from Green River, Wyoming. Comparing these with typical *semiscaber* from the vicinity of Pocatello, Idaho, we can appreciate no difference whatever, except in the matter of armature. None of our Green River specimens shows any prickles whatever. About half the Pocatello specimens are also naked and are indistinguishable from typical

punctulatus; in the others more or less prickles are developed, varying from a few in axil of pectorals to a band covering more than ½ of the sides. As this is not an unusual amount of variation, we do not consider *Cottus semicaber* worthy of recognition. The relations of *Cottus punctulatus* with the eastern species have not been carefully worked out, and it seems best to recognize it for the present as distinct. Specimens from Green River, Wyoming, and from Mink Creek, Ross Fork, and Port Neuf River, Pocatello, Idaho. The collection contains also 2 specimens from Thompson Falls and 3 from Flathead Lake, which seem to be this species. By the Fort Hall Indians this fish is called *ahwe,* a word meaning horns.

(*punctulatus,* speckled.)

Potamocottus punctulatus, Gill, Proc. Bos. Soc. Nat. Hist. 1861, 40, Bridgers Pass, Wyoming (Coll. Capt. Simpson); Gill, Ichth. Capt. Simpson Expl., 402, 1876, with plate.
Cottus punctulatus, Gilbert & Evermann, Bull. U. S. Fish Comm. 1894, 202.
Uranidea punctulata, Jordan & Gilbert, Synopsis, 697, 1883.

2320. COTTUS SEMISCABER (Cope).

(Rocky Mountain Bullhead.)

Head 3; depth 4¼; D. VIII, 17 or 18; A. 12 or 13; eye 5 in head; maxillary 2¼; third dorsal spine 4 in head; highest dorsal rays 2⅔; pectoral 1¼; ventral nearly 2; caudal 1⅔. Body not much compressed, caudal peduncle wide, equal to length of snout; head broad and rounded anteriorly as viewed from above; mouth large, the maxillary reaching about to posterior margin of eye; band of teeth broad on jaws, vomer, and the palatines; interorbital space (bone only) narrow, not over ¼ eye; preopercular spines rather stout and blunt. Pectorals reaching to front of anal; dorsals not connected, scarcely meeting in some specimens; ventrals not reaching to vent; last rays of soft dorsal reaching almost to base of caudal, highest dorsal rays equal snout and ½ eye; vent much nearer tail than tip of snout. Color slaty gray in spirits, with mottlings on sides which form inconspicuous cross bars; pectorals, dorsals, and caudal, with wavy cross lines; ventrals and anal white or sometimes dusky and mottled; a dark spot on spinous dorsal posteriorly, fin edged with white. Rocky Mountain region, in clear streams on both slopes. Abundant in Colorado, northern New Mexico, Wyoming, Montana, Idaho, Utah, eastern Washington, and Oregon. Found in the upper basins of the Fraser, Columbia, Missouri, Colorado, Platte, Arkansas, and Lake Bonneville, its eastern and northern limits not well ascertained. It is subject to very great variations in color, size, and roughness of skin. Most specimens are smooth, or rough only in the axil. Some, especially from southern Idaho and Provo River, Utah, are quite rough, some of them as rough as *Cottus asper* or *Cottus rhotheus.* These rough specimens differ in no other respect from smooth ones from the same localities. The specimens here especially described are from Eagle River at Gypsum, Colorado, the largest 4½ inches in length.

Rough specimens from Provo are thus described by Jordan & Gilbert (Synopsis, 695): Head 3¼; depth 4⅓. D. VII, 18; A. 14; V. I, 4; eye 4½ in head. Form of *Cottus asper,* but slender. Skin of top of head and entire

body, except lower part of caudal peduncle, thickly covered with slender papillæ; those of anterior and upper parts of body each tipped with a spine, many of those on head with median pores; top of head with a median lengthwise depression. Head narrowed anteriorly, the maxillary extending to beyond pupil; opercular and preopercular spines as in *Cottus asper*; lateral line not complete; spinous dorsal low; soft dorsal and anal high; caudal long; pectorals about reaching anal; ventrals rather broad, reaching about halfway to anal. Olivaceous, barred and spotted as in other species; fins mottled; spinous dorsal pale at base and tip, with a median broad black band; 2 dark blotches at base of caudal. Length 4 inches. The specimens here described from Provo River at Provo, where it is associated with the smooth form called *wheeleri*, which differs only in its smooth skin. In specimens from the Yellowstone Park the band of palatine teeth is very broad; there are no prickles on the skin. The head is $3\frac{1}{2}$ in length and the rays are D. VII, 17; A. 13; V. I, 4. Comparing these (Gibbon River, Wyoming) with specimens of *Cottus ictalops*, from Mammoth Spring, Arkansas, the differences seem well marked. *Cottus semiscaber* has the head blunter, lower, and more rounded, the cheeks more tumid and the top of the head without median longitudinal depression. *Cottus ictalops* has the axil prickly, the outline of the head angular, the top of the head with a median longitudinal depression from snout to nape, and the body has broad distinct black cross bars. These 2 forms seem like distinct species, but other specimens are intermediate; specimens from Torch Lake, Michigan, agree with *semiscaber* in color, and are intermediate in form; specimens from White River, Indiana, are colored like *ictalops*, but are intermediate in form. Apparently *semiscaber* should be recognized as a species, but its range and distinctive characters are yet to be made out in detail. (*semiscaber*, half rough.)

Cottopsis semiscaber,* Cope, Hayden Survey of Montana, 476, 1871 (1872), Fort Hall, Idaho; Jordan & Gilbert, Proc. U. S. Nat. Mus. 1880, 459; Jordan, Bull. U. S. Fish Comm. 1889, 53.

Uranidea semiscabra, Jordan & Gilbert, Synopsis, 695, 1883.

Uranidea vheeleri, Cope, Proc. Amer. Phil. Soc. Phila. 1874. 138, Bear River, Utah. (Coll. Lieut. G. M. Wheeler.)

Uranidea vheeleri, Cope & Yarrow, Zool. Wheeler's Expl. W. 100th mer., v, 696, pl. 32, figs, 3, 3a, 3b, 1876; Jordan & Gilbert, Synopsis, 697.

Cottus bairdi punctulatus, Jordan, Bull. U. S. Fish Comm. 1889, 29, 36, 53; with plate.

2321. COTTUS ICTALOPS (Rafinesque).

(Miller's Thumb; Blob; Muffle-jaw; Bullhead; Springfish.)

Head $3\frac{1}{3}$; depth 4 to 6. D. VI to VIII, 16 or 17; A. about 12; V. I, 4. Body slender or stout, tapering regularly backward to the tail; vertex

* *Cottopsis semiscaber* is thus described by Professor Cope:
"Radii, D. VII, 18; A. 13; V. I, 4; first ray of anal below third of second dorsal. Skin prickly above the lateral line, smooth below it posteriorly. Body compressed, profile rising rather steeply to the basis of first dorsal fin. Eye 4.5 times in head, .75 time in interorbital space. Muzzle contracted, maxillary bone reaching to below middle of pupil. Two spines on preoperculum; 1 on inferior angle of operculum. Lateral line discontinued on last fourth of caudal peduncle. Head $\frac{1}{3}$ length, without caudal fin. Below yellow; dorsal line with a series of dark spots; sides with large dark clouds. Three specimens from Fort Hall, Idaho." (Cope.)

somewhat depressed; interocular space with a groove; preopercle with a short, sharp spine, little hooked, directed backward and upward, mostly covered by the skin; below this are 2 smaller concealed spines; subopercle with a stoutish spine, directed forward. Skin smooth, except the region immediately behind the pectorals, which is beset with very small, sharp prickles, which are sometimes obsolete; lateral line conspicuous, continuous, or interrupted behind; first dorsal low and feeble; pectoral fins large, their length nearly equal to that of the head, their tips usually reaching beyond the origin of the soft dorsal; ventral fins moderate; isthmus very broad, the gill membranes not forming a fold across it. Olivaceous, more or less barred and speckled with darker; fins mostly barred or mottled. Length 3 to 7 inches. Middle and Northern States, abounding in all clear, rocky brooks and lakes east of the Dakotas and Kansas to New York and Virginia, extending southward along the Alleghanies to North Carolina and northern Alabama, especially abundant in limestone springs and entering caves. Extremely variable; very destructive to eggs of trout. (*Ictalurus*, the catfish; ὤψ eye; "the name means cat's eye; eyes like those of the catfishes with oblong eyes." Rafinesque.)

NOTE.—As here understood, *Cottus ictalops* is a widespread and abundant species, varying in different regions, as is the case with most nonmigratory species. In this, as in others of similar range, the inhabitants of each stream may show local peculiarities. A number of these forms have received from Dr. Girard specific names, which are accompanied by detailed descriptions. Large collections of these fishes show that numerous similar "species" still exist undescribed, as it is a rare thing to find a specimen which exactly agrees in all respects with any of the species in Dr. Girard's "Monograph of the Fresh-Water Cottoids." In this work the figures and long descriptions give characters of individuals, not of species.

Pegedictis ictalops,* RAFINESQUE, Ichth. Ohiensis, 85, 1820, spring near Lexington, Kentucky.

Cottus richardsoni,† AGASSIZ, Lake Superior, 300, 1850, north shore of Lake Superior (Coll. Louis Agassiz and Dr. C. T. Jackson); GIRARD, Monograph Fresh-Water Cottoids N. A., 39; GÜNTHER, Cat., II, 158; not *Trachidermis richardsoni*, HECKEL.

Cottus bairdii,‡ GIRARD, Proc. Amer. Assoc. Adv. Sci., II, 1850, 410, and Monograph Cottoids, 44, with plate, Mahoning River, Poland, Ohio. (Coll. S. F. Baird.)

Cottus meridionalis,§ GIRARD, Proc. Amer. Assoc. Adv. Sci., II, 1850, 410, and GIRARD, Monograph Cottoids, 47, James River, Virginia. (Coll. S. F. Baird.)

* In spite of the errors in description, there can be no doubt that Rafinesque's "cat's eye Springfish." *Pegedictis ictalops*, is the present species; the dorsal rays are counted wrongly, and the scales are said to be small. Nevertheless, "in the distorted perspective of his mental vision" Rafinesque could have had nothing else. In other words, his description was drawn up hastily and in part from memory.

† The form called *Cottus richardsoni*, Agassiz, is rather slender, with the vent rather more posterior than usual, placed midway between the snout and the tip of the caudal; in the others it is nearly midway between the snout and the middle of the caudal. Lake Superior. A specimen taken by Dr. Jordan in Lake Superior at Marquette, Michigan, agrees with *Cottus ictalops* from southern Missouri. It is a little more slender, the color a little darker and less definite, the cross bars fainter, the dark punctulations more conspicuous. The vent has the usual position, although Dr. Girard figures the vent in his Lake Superior specimens, types of *Cottus richardsoni*, as farther back than in *Cottus ictalops*. In his figure of *Cottus richardsoni* the vent is a shade nearer base of caudal than posterior margin of eye. In all specimens we have seen it is nearly midway between nostril and caudal. D. VIII, 18; A. 12, in the specimen from Marquette.

‡ The form called *Cottus bairdi*, Girard, is small and slender, with the spinous dorsal very low, and the palatine teeth less developed than in the other forms. Cayuga Lake, New York, to Ohio.

§ *Cottus meridionalis*, Girard, is rather robust, with the dorsal fins scarcely connected, and the mouth larger, the maxillary extending to opposite posterior border of eye; the preopercular spine is sharp and directed well upward. Pennsylvania to North Carolina, along the Alleghanies; abundant.

Cottus wilsoni,* GIRARD, Monograph Cottoids, 42, 1851, Pittsburg, Pennsylvania. (Coll. Jacob Green.)

Cottus alvordii,† GIRARD, Monograph Cottoids, 46, 1851, Fort Gratiot, Lake Huron. (Coll. Major Benj. Alvord.)

Potamocottus zopherus,‡ JORDAN, Ann. Lyc. Nat. Hist. N. Y. 1876, 320, Etowah River and tributaries, Rome, Georgia. (Coll. Jordan & Gilbert.)

Uranidea richardsoni. JORDAN & GILBERT, Synopsis, 696, 1883.

Uranidea richardsoni, JORDAN & GILBERT, Synopsis, 696, 1883; but the name *richardsoni* is preoccupied in *Cottus*.

Cottus bairdii, JORDAN, Bull. U. S. Fish Comm. 1889, 29.

Potamocottus carolinæ,§ GILL, Proc. Bost. Soc. Nat. Hist. 1861, 40, Carolina; GILL, in Simpson Rept. Ichth. Utah, 403, 1877.

Subgenus TAURIDEA, Jordan & Rice.

2322. COTTUS RICEI, Nelson.

Head $3\frac{2}{5}$; depth $5\frac{1}{3}$. First dorsal VIII, second dorsal and anal destroyed; V, I, 4; P. 15. Eye $4\frac{1}{4}$, $1\frac{1}{2}$ in interorbital space and equalling snout. Body short and stout, abruptly contracted opposite base of anal. Head much depressed, very broad and flat, broader than body, breadth greater than length; depth $\frac{1}{2}$ length. Palatine teeth present. Tail very small, subterete. Outline tadpole-like. Jaws about equal; mouth rather narrow; jaws contracted and somewhat produced. Eyes on upper surface near together. Preopercular spine extremly large, 3 times as large as in any other fresh-water Cottoid known, as long as eye, hooked backward and upward, giving a buffalo-like appearance. Three spines hooked downward below the larger spine; the lower concealed; a strong spine hooked forward at base of opercles. Branchiostegals 6. Isthmus as wide as from snout to middle of orbit. Base of pectorals crescentic, their tips just short of anal; rays all simple; ventrals reaching $\frac{2}{3}$ distance to vent. Profile rising rapidly to dorsal, which runs along a sort of carina. Dorsal beginning a trifle behind ventrals, just behind head, about midway between snout and anal. Vent midway between snout and base of caudal. Depth at first ray of anal less than $\frac{1}{2}$ length of head, thickening at same point over $\frac{1}{3}$. Head smooth. Space above lateral line behind head covered with small stiff prickles hooked backward, readily visible as small black specks when skin is dry. Color pale brown, irregularly spotted and mottled with darker brown somewhat as in *Lota;* pectorals mottled; belly white; spines spirally curved, forming $\frac{1}{2}$ a spiral. The most peculiar characters are the strong spines of the preopercle and the smaller ones below,

* *Cottus wilsoni*, Girard, is rather stouter, with stronger palatine teeth, and with some of the uppermost of the pectoral rays bifurcate, these being entire in the others. Ohio Valley, Pennsylvania, and Indiana.

† *Cottus alvordi*, Girard, is short and chubby, with the first dorsal rather high, and joined by membrane to second more than in the other forms. Common in Wisconsin and Michigan.

‡ *Cottus zopherus*, Jordan, is slender and very dark in color, and more conspicuously variegated; the first dorsal high; the palatine teeth well developed; probably worthy of varietal recognition. Alabama Basin.

§ *Cottus carolinæ*, Gill, is a very large form, rather robust, reaching a length of nearly 6 inches, without axillary prickles, and with the palatine teeth well developed. The lateral line, as in the other forms, is sometimes continuous and sometimes interrupted. It abounds in the limestone region from Indiana to Tennessee, and is frequently found in caves.

the carinated back and abruptly contracted body, forming the subterete caudal peduncle. The prickles of the skin are coarser than in any other species. (Nelson.) Length of type $2\frac{5}{6}$ inches. Great Lakes, the 2 type specimens from Lake Michigan, near Evanston, Illinois, in rather deep water; remains of others occur among Girard's types of *Triglopsis thompsoni* from Lake Ontario. The group *Tauridea*, of which this species is type, may be worthy of generic rank. (Named for its discoverer, Mr. F. L. Rice, then a student in zoology in Northwestern University at Evanston.)

Cottus ricei, NELSON, Bull. Ills. Mus. Nat. Hist., Vol. I, No. 1, 40, 1876, Lake Michigan, at Evanston (Coll. F. L. Rice).
Uranidea spilota, JORDAN & GILBERT, Synopsis, 694; not of COPE.
Uranidea ricei, JORDAN & GILBERT, Synopsis, 953.

Subgenus COTTUS.

2323. COTTUS ONYCHUS, Eigenmann & Eigenmann.

Head $3\frac{3}{4}$; depth $5\frac{1}{2}$. D. VIII, 17; A. 13; V. I, 4; P. 13; eye $1\frac{1}{3}$ in snout, $\frac{1}{2}$ in interorbital, 5 in head. Teeth on vomer, none on palatines. Width of head equaling its length to end of preopercular spine, its depth 2 in length. Preopercle with an upturned claw-like spine, below which are 2 others, much smaller, the anterior one having its point turned downward and forward. Maxillary not reaching orbit; lateral line complete. Sides above lateral line, with stiff prickles from below first spine to below last dorsal ray; prickles below lateral line confined to the abdominal part of the sides. Dorsal connected by a low membrane, the rays much higher than the spines, $3\frac{1}{2}$ in head. Pectoral reaching past vent, its rays not branched. A dusky spot on breast just behind anterior end of gill slits; ventral surface including the ventrals otherwise plain; anal with a few dusky specks on its rays; other fins barred; sides and upper surfaces olive with darker spots; 3 dark bands below soft dorsal; a dark band just in front of caudal. Length about $3\frac{1}{2}$ inches. Saskatchawan Basin. This species is evidently closely related to *Cottus pollicaris*, from which it differs chiefly in having many prickles. (Eigenmann.) Not seen by us and perhaps not different from *Cottus pollicaris*. (ὀνυχός, clawed.)

Cottus onychus, EIGENMANN & EIGENMANN, Am. Nat., Nov., 1892, 963, Bow River, at Calgary, a tributary of the South Saskatchawan, Alberta Territory. (Coll. C. H. Eigenmann.)

2324. COTTUS POLLICARIS (Jordan & Gilbert).

Head $3\frac{3}{7}$ in length to base of caudal; depth $4\frac{3}{4}$; eye $5\frac{1}{8}$ in head. D. VII, 19; A. 13; V. I, 4; P. 17. Lateral line complete. Body robust; nape prominent, the profile of head steeply declined, thence to tip of snout in a straight or slightly concave line; head much depressed, broad and flat above, evenly narrowed forward to the broad, much depressed, bluntly rounded snout; eyes small, with extensive vertical range, their diameter less than snout or the flat interorbital width; mouth rather small, anterior, with but little lateral cleft, maxillary reaching vertical from front of orbit; teeth villiform on jaws and vomer, none on palatines; preoper-

cular spine large and strong, spirally curved upward and inward, wholly invested with membrane; a single sharp concealed spinous point below angle of preopercle; isthmus broad, without fold, its width equaling distance from snout to middle of pupil. Spinous dorsal rather low, nearly uniform in height, connected with second dorsal by a low membrane; longest spine equaling length of snout; soft dorsal long, its longest ray $2\frac{1}{5}$ in head; origin of anal fin under third dorsal ray, its last ray under sixteenth of dorsal; highest anal ray $2\frac{1}{2}$ in head; ventrals I, 4, reaching $\frac{2}{3}$ distance to vent; pectoral rays all simple, unbranched, the longest reaching vertical from vent, and contained $1\frac{1}{3}$ times in head. Vent equidistant between tip of snout and base of caudal fin. Skin everywhere smooth. Color olivaceous above, little punctulated, lower $\frac{2}{3}$ of sides and whole under side of head and body uniform whitish above, head and body with irregular spots and blotches of black; these in finer pattern on head, and not forming bands on back; dorsals, caudal, and pectorals with black spots arranged in more or less distinct series; anal, ventrals, and lower rays of pectorals translucent, unmarked. Lake Michigan; a single specimen $4\frac{1}{2}$ inches in length taken off Racine, Wisconsin, by Dr. Philo R. Hoy. (*pollex*, the thumb, suggested by the name, Miller's Thumb.)

Uranidea pollicaris, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 222, Lake Michigan, off Racine, Wisconsin (Type, No. 29663. Coll. P. R. Hoy); JORDAN & GILBERT, Synopsis, 954, 1883.

2325. COTTUS COGNATUS, Richardson.

(BEAR LAKE BULLHEAD.)

This species is thus described by Richardson: The *Cottus* which forms the subject of this article was taken in considerable numbers in the clear waters of Great Bear Lake during the month of May, at which period it resorts to the stony shallows to spawn. Specimens which we sent to Baron Cuvier were returned with the remark that they belonged to a species of *Cottus* and were "très semblables aux *C. gobio*," and they, indeed, correspond in most particulars with the extended description of the latter in the Histoire des Poissons. I have since compared these specimens minutely with an English *gobio*, kindly lent to me by Mr. Yarrell, and the principal difference that I have been able to detect in the American fish is the greater height of its dorsal and anal fins. There is also a discrepancy in the number of rays, but this can scarcely be accounted a specific distinction, since different individuals of *gobio* show equally extensive variations. The Bear Lake specimens having been long in spirits have lost much of their color, but the following particulars may still be noted: In *Cottus gobio* the rays of the pectorals are variegated with rings alternately dark and light; in *cognatus* the color is almost uniform, but varies in intensity in different individuals. In 1 small female specimen of the latter, however, there is a slight indication of these rings, and its body likewise is marbled in a more lively manner than the rest, having a greater resemblance to *gobio*. The males are darker than the females in the American as well as the European species. The specimen which we have described in detail was the largest we obtained, and was selected for

description from its being exactly in the same length with the *gobio* with which it was compared. Great Bear Lake. Length 4 inches. (Richardson.) Not seen by recent naturalists, probably allied to *Cottus pollicaris*. (*cognatus*, related, to *Cottus gobio*.)

Cottus cognatus, RICHARDSON, Fauna Bor.-Amer., III, 40, 1836, Great Bear Lake (Coll. J. Richardson); GIRARD, Mon. Cot., 41; GÜNTHER, Cat., II, 157.
Uranidea cognata, JORDAN & GILBERT, Synopsis, 953, 1883.

2326. COTTUS PERPLEXUS, Gilbert & Evermann.

Head $3\frac{1}{3}$; depth $4\frac{1}{5}$. D. VII, 21; A. 15; P. 16; V. I, 4; eye 4; snout 4; interorbital width $5\frac{1}{2}$. Least depth of caudal peduncle greater than snout, $3\frac{3}{8}$ in head; interorbital space rather broad, about $1\frac{1}{2}$ in eye. Body deeper and more compressed than in any other species known to us, this being especially noticeable posteriorly; caudal peduncle very short and deep, and entirely overlapped by posterior dorsal rays which extend beyond base of caudal fin. Length of caudal peduncle from base of last dorsal ray about $\frac{3}{4}$ depth of same; depth of body at origin of anal fin $\frac{3}{8}$ length of head. Interorbital space slightly concave; occiput flat or transversely convex. Mouth oblique, the maxillary reaching vertical from posterior margin of pupil, $2\frac{1}{2}$ in head. Teeth in a very narrow crescentic band on vomer, none on palatines. Upper preopercular spine short and broad, curved or simply directed upward; below this are 2 stout, blunt spines directed downward. Body entirely naked, lateral line incomplete, not reaching end of soft dorsal. Spinous dorsal low, the longest spines not greater than length of snout; soft fins all high, the fifteenth dorsal ray equaling snout and eye; a broad membrane always connecting the 2 dorsals, the notch inconspicuous; last rays of anal, as well as dorsal, extending beyond base of caudal; first anal ray under third ray of soft dorsal; ventral spine and rays slender and weak. Anus midway between base of caudal fin and front of eye. Color in alcohol, back and sides with vermiculations of light and dark, the back with 5 or 6 ill-defined black cross bars, which usually reach the lateral line; the usual black bar at base of caudal, emarginate posteriorly; below the lateral line a number of small, quadrate dark blotches, arranged in 2 irregular series; lower part unmarked, except with fine dark punctulations; dorsal, pectoral, and caudal fins cross-barred with dark; anal and ventrals with numerous small dark specks. Length about $3\frac{3}{8}$ inches. Shookumchuck and Newaukum rivers, near Chehalis, in western Washington. (Gilbert & Evermann.) (*perplexus*, perplexed.)

Cottus perplexus, GILBERT & EVERMANN, Bull. U. S. Fish Comm. 1894, 202, pl. 20. (Type, No. 45387, U. S. Nat. Mus.; Cotypes, Nos. 1324 to 1343. L. S. Jr. Univ. Mus.); Shookumchuck River, near Chehalis, Washington (No. 45388, U. S. Nat. Mus.), and Newaukum River, near Chehalis. (Coll. Gilbert & Jenkins.)

2327. COTTUS KLAMATHENSIS, Gilbert.

Head $2\frac{9}{10}$ to $3\frac{1}{10}$ in length; depth $3\frac{9}{10}$ to $4\frac{1}{5}$. D. VII, 19; A. 14; P. 15; V. I, 4. Body heavy and deep, the head narrowed and wedge-shaped anteriorly, the snout rather acute and the mouth with much lateral cleft. Max-

illary broadly exposed, its tip reaching vertical from behind front of pupil, its length 2⅔ or 2⅞ in head. Broad bands of teeth on jaws and vomer; palatines toothless. Anterior nostril with a distinct tube. Eye of moderate size, 1⅔ in snout, 4⅞ to 5 in head. Interorbital space and occiput gently concave in adults, the total interorbital width 1⅛ to 1½ in orbit, the bony septum narrower. Upper preopercular spine robust, straight, directed backward, or backward and slightly upward; below this the margin of the bone is without evident spines, but bears 1 or 2 slight prominences which may be rounded or acute; anterior angle of the subopercle with a short spine directed forward; opercle ending in a short flat spine. Head with large pores, 2 pairs above front of orbit, those of the posterior pair nearest together; distant from these, a single median pore on the posterior portion of interorbital space, from which diverge 2 lines of pores around the back of the orbits; spinous dorsal short and low, the longest spine usually less than ⅔ the longest soft ray; the 2 fins very broadly joined; distance from base of last dorsal ray to base of caudal slightly less than depth of caudal peduncle; caudal short and broadly rounded, its length 1⅔ in head; pectorals very short, usually not reaching vertical from front of anal, 1⅔ in head; ventrals large, sometimes reaching vent but usually shorter, 1¾ in head; caudal with 9 (sometimes 8 or 10) forked rays; rays of other fins simple, unbranched. The variation in fin rays is shown in the following table:

	Number of specimens.	Spinous dorsal.		Soft dorsal.			Anal.			Pectoral.	
		VII	VIII	18	19	20	13	14	15	14	15
Upper Klamath Lake	21	19	2	1	18	2	....	16	5	3	18
Klamath River below lake	4	3	1	....	4	....	1	3	....	....	4
Lost River	7	7	.....	3	3	....	2	4	1	2	5
Scott River	1	1	.....	....	1	....	....	1	....	....	1

Skin mostly naked, the young with a narrowly oblong patch of prickles below the lateral line and under the posterior half of pectorals, these becoming gradually absorbed with age, adults being nearly or quite naked. Lateral line very incomplete, the last pore under some portion of the anterior half of soft dorsal in all our specimens from the lake. From the last pore, a shallow open groove or trace follows the course of the obsolete portion of the canal. In 4 specimens from Klamath River below the falls, and in 1 collected by Mr. R. C. McGregor in Scott River, Siskiyou County, California (a tributary of Klamath River), the lateral line is much more nearly complete, ending under the last fifth of soft dorsal. Color brownish olive, with 4 or 5 indistinct dark bars downward from back, breaking up below into narrow bars which may unite to form V-shaped markings, or often into mere irregular blotches; a narrow bar at base of tail; caudal with broad dark bars alternating with much narrower light ones; dorsal and anal with somewhat narrower oblique bars;

pectorals very conspicuously checkered, the dark and light spots on the rays arranged in vertical series. A large, strongly marked species, very abundant in Upper Klamath Lake. It is characterized by its short spinous dorsal, broadly joined to the long soft dorsal, the unbranched pectoral rays, the very incomplete lateral line, the weak development of prickles, the lack of palatine teeth, and the distinctive coloration. It is most nearly related to *C. perplex*. Klamath Lakes, Oregon.

Cottus klamathensis, GILBERT, Bull. U. S. Fish Comm. 1897, 10, figure, Upper Klamath Lake, Oregon. (Type, No, 48226, U. S. Nat. Mus. Coll. Gilbert, Cramer, and Otaki.)

2328. COTTUS ALEUTICUS, Gilbert.

Head small, $3\frac{1}{4}$ to $3\frac{1}{2}$; depth 5 to $5\frac{1}{5}$. D. IX or X, 18 or 19; A. 13 or 14; P. 13 to 15; V. I, 4; C. 8 or 9 (forked rays); 35 to 37 pores in lateral line. In appearance resembling *Cottus philonips;* the head small, its width but $\frac{1}{4}$ greater than its depth; the body low and but little compressed, the depth at shoulders but little greater than the width. Caudal peduncle moderate, not slender, its length from base of last anal ray $1\frac{2}{3}$ to $1\frac{1}{2}$ in head, measured from last dorsal ray but little more than $\frac{1}{2}$ as long. Depth of caudal peduncle $4\frac{1}{3}$ to $4\frac{1}{2}$ in head. Mouth small, variable, maxillary reaching vertical from front or middle of pupil, $2\frac{2}{3}$ to 3 in head. Vomerine patch of teeth small, the palatines toothless. Preorbital as wide as eye, produced anteriorly into a convex lobe, which conceals all but the widened tip of the maxillary; both pairs of nostrils opening in short but evident tubes, a character not known by us to exist in other species of the genus. Eye $4\frac{1}{3}$ to $4\frac{2}{3}$ in head; interorbital space narrow, the least width of the bone slightly less than $\frac{1}{2}$ the vertical diameter of the orbit in adults 160 mm. long, much narrower in younger specimens. In this respect our specimens differ conspicuously from the types of *U. microstoma*, in which the interorbital width is said to equal vertical diameter of eye. As in *Cottus philonips* we have but a single preopercular spine, which is straight and directed obliquely upward; preopercular margin below the spine evenly rounded; subopercular spine well developed. Lateral line complete, following outline of back to opposite last dorsal ray, where it abruptly declines to middle of caudal peduncle. Body naked, or with a narrow band of prickles extending from upper axil of pectorals along under side of lateral line. Dorsals more or less joined at base in all our specimens, but varying in the height of the connecting membrane, which usually joins well up on first ray of soft dorsal, sometimes at its extreme base. The spinous dorsal is long, with 9 spines in 13 specimens examined, 10 spines in 5 specimens. The first 2 spines are very closely approximated, and spring from a single, wide interspinal; the first spine is easily overlooked, as has been done by Gilbert and Evermann (Investigations in the Columbia River Basin, 1894, p. 54), and possibly also by Lockington in his description of *Uranidea microstoma*. Spinous dorsal low; the soft dorsal higher, the longest rays equaling length of snout and $\frac{1}{2}$ eye; the last rays when depressed not quite reaching base of caudal. Anal fin much shorter than soft dorsal, its last ray under the fourth or fifth before the last ray

of dorsal; caudal truncate, slightly rounded when spread, its length 1½ to 1⅞ in head, its rays twice forked; pectorals reaching to or nearly to front of anal; ventrals varying in length, not quite reaching vent in any of our specimens; pectoral rays all simple; dorsal and anal rays all simple, except the last, which is usually divided to the base in the former, and sometimes divided in the latter. Head and body mottled or spotted above, uniformly light brown, the darker markings on back often arranged as 6 cross bars, of which 2 are below spinous dorsal, 3 below soft dorsal, and 1 on caudal peduncle, these usually broken up into spots or reticulations, and often obscure, sometimes wanting; usually a light bar downward and backward from eye; rays of dorsals, pectorals, and caudal crossed with series of dark blotches; ventrals and anal light, very obscurely barred with darker. This species was very abundant in the small streams passing through the village of Iliuliuk, Unalaska, living both in the upper, strictly fresh-water portion of the stream, and in the lower, more or less brackish part. A specimen transferred to the salt-water aquarium on the *Albatross* seemed to suffer no inconvenience from the change of water, and lived for several days. This is probably the *Uranidea microstoma* of Lockington, based on specimens collected near St. Paul, Kadiak, and considered by the describer to be identical with other specimens examined by him from the Aleutian Islands. Four specimens were collected by us May 26, 1889, in a small stream entering Departure Bay, Vancouver Island. These exhibit perfectly the differences separating *Cottus aleuticus* from its nearest ally, *Cottus philonips*. (Gilbert.)

Uranidea microstoma, Lockington, Proc. U. S. Nat. Mus. 1880, 58, St. Paul, Kadiak Island (Coll. W. J. Fischer); not *Cottus microstomus*, Heckel, a European species.

Cottus aleuticus, Gilbert, Rept. U. S. Fish Comm. 1893 (1896), 418, streams of Unalaska; also in Departure Bay, Vancouver Island. (Coll. C. H. Gilbert).

2329. COTTUS MINUTUS, Pallas.

Head 3½. D. VII, 18; A. 13; C. 13; P. 14; V. 2. Head large, conic, the upper jaw longer, the teeth very slender; eyes small, near together; preopercle with an incurved spine on each side, its lower margin subdentate; the head otherwise unarmed; body terete, thick anteriorly, tapering behind; the lateral line unarmed. Color pale, dotted with large dusky, irregular black spots on the back; fins all variegated, the pectorals translucent, with rows of dusky spots; ventrals white, with dusky rings; second dorsal variegated; anal dotted with dusky, caudal with dusky bands; body smooth. Island of Talek, near Tauisk, Sea of Okhotsk; known from a dried specimen 3 inches long, and a drawing. (Pallas.) Evidently close to *Cottus aleuticus*, but not to be identified without new material from Siberia. (*minutus*, very small.)

Cottus minutus, Pallas, Zoogr. Rosso-Asiat., III, 145, 1811, Talak Island, Sea of Okhotsk. (Coll. D. D. Merk).

2330. COTTUS BELDINGII, Eigenmann & Eigenmann.

Head 2¾ to 4; depth 4 to 5. D. VI to VIII, 15½ to 18; A. 11 to 13; V. I, 4. Head rather short and broad, the profile convex, more steep from eye for-

ward; eye large, orbit 4 to 5 in head; interorbital concave, 2 in orbit; mouth large, maxillary reaching at least to below the pupil, about 2 in head. Preopercle with a simple backward-directed spine, very slightly curved upward. Teeth on jaws and vomer, none on palatines. Skin smooth. Pectorals reaching vent, or further in the young; ventrals $1\frac{1}{2}$ to 2 in head. Distance of anal from caudal $1\frac{1}{2}$ in its distance from snout. Anus nearer insertion of caudal than to end of snout. Mottled with black and white; about 6 blackish cross bars on back, the first across head just behind eyes, next at origin of dorsal; first dorsal tinged with rusty, the second less so; all the fins except the ventrals spotted with dark. The ground color varies greatly with the bottom over which these fishes live. (Eigenmann.) Streams of Columbia River Basin, south to Lake Lahontan; abundant east of the Cascades. We find this species abundant in shallow water in Lake Tahoe. Except by the stouter body, we are unable to separate it from *Cottus philonips*, and perhaps Gilbert & Evermann are right in identifying it with the latter. We refer to *Cottus beldingii* most of the specimens called *Cottus philonips* by Dr. Gilbert and Dr. Evermann, with the remark that the name *Cottus beldingii* is in any event prior to *Cottus philonips*.

Cottus beldingii is a small-headed form, typically with perfectly smooth skin and unarmed palatines. Like most other species of the genus it occasionally develops a band of postaxillary prickles, which are often accompanied in the same specimens by a small patch of teeth on the palatine bones. The head is less strongly armed than usual, the single preopercular spine being short, the preopercular margin otherwise wholly unarmed. In this respect *C. beldingii* differs from all other western species of *Cottus*, except the Alaskan form above mentioned. The dorsal varies from VII or VIII, 16 to 18; the anal from 12 to 14. The nostrils are without tubes, and the preorbital little produced, exposing the greater part of the maxillary in closed mouth. Specimens were obtained in the Port Neuf River near Pocatello, at Snoqualmie Falls, and in a spring branch emptying into the South Fork of the Cœur d'Alene River, near Wardner, Idaho. We have also seen specimens taken from Birch Creek, in western Idaho, by Merriam and Bailey. (Named for Lyman Belding, of Stockton, California, well known as an ornithologist.)

Cottus beldingii, Eigenmann & Eigenmann, Amer. Nat., xxv, 1891, 1132, Lake Tahoe (Coll. Lyman Belding and C. H. Eigenmann), and Donner Lake (Coll. Eigenmann).

2331. COTTUS PHILONIPS, Eigenmann & Eigenmann.

Head $3\frac{1}{4}$; depth 6. D. IX–17; A. 13; eye 4 in head; maxillary $2\frac{2}{3}$; highest dorsal spine 3; highest soft ray $2\frac{1}{4}$; pectoral $\frac{7}{8}$; caudal $1\frac{1}{3}$. Body extremely slender, not much compressed; caudal peduncle moderate. Head small, wedge-shaped, as viewed from above; from the side the upper profile is broadly rounded from the dorsal to the tip of the rather sharp snout, lower profile straight; mouth moderate, the maxillary reaching to middle of eye; teeth on jaws and vomer, palatines toothless; interorbital space equals $\frac{1}{2}$ eye; eye about equal to length of snout; preopercle with only 1 small spine above, below which its edge is entire; pectoral reach-

ing front of anal or past; dorsals not connected, the highest spine about $\frac{2}{3}$, the highest soft dorsal $1\frac{1}{3}$, in head; ventrals reaching to vent. Color light gray, covered with many spots forming reticulations on sides and top of head; ventrals white, other fins covered by wavy, dark bars; dorsal entirely dusky, with a narrow white edge; belly and lower parts white. Fraser River Basin. Here described from one of the type specimens, 3 inches in length; collected at Field, British Columbia, in Kicking Horse River, by Dr. Eigenmann. This has been compared with many specimens of *Cottus beldingii* from localities in Washington and Idaho and from Lake Tahoe. All of these are less slender than the types, the depth being $4\frac{1}{2}$ to 5 inches in length, but they do not differ otherwise, and later investigations will probably show the entire identity of *Cottus philonips* with *Cottus beldingii*.

This name (*Cottus philonips*) was proposed as a substitute for *Cottus minutus*, Pallas, supposed to be preoccupied, and *Cottus microstomus* (Lockington), not of Heckel; but the original description was taken from a specimen from Kicking Horse River. The first mentioned is perfectly available, but was applied to a specimen from the island of Talek, near Tauisk, in the Okhotsk Sea. It is very doubtful, therefore, whether *Cottus minutus* should be used for any American species in advance of comparison with the Siberian form. From the Aleutian Island species (*C. microstomus* Lockington, *C. aleuticus* Gilbert), *C. philonips* differs in many important respects, and is undoubtedly distinct. Thus the Alaskan form has the posterior nostrils in short, but conspicuous tubes, the preorbital produced into a lobe which conceals all of the maxilary except the extreme tip, and the dorsal fin with 8 or 9 spines and 18 to 20 soft rays. (*φιλέω*, to love; *νίψ*, snow, *νίπτω*, to wash.)

Cottus philonips,* Eigenmann & Eigenmann, Amer. Nat., xxvi, 963, 1892, Kicking Horse River, Field, British Columbia; Eigenmann & Eigenmann, Bull. U. S. Fish Comm. 1894, 118; Gilbert & Evermann, Bull. U. S. Fish Comm. 1894, 204, in part, includes *Cottus beldingii*.

2332. COTTUS ANNÆ, Jordan & Starks.

Head $3\frac{1}{2}$ to $3\frac{2}{3}$; depth 5. D. VII or VIII, 17 or 18; A. 12; eye 5 in head; maxillary $3\frac{3}{8}$; highest dorsal $3\frac{1}{2}$; highest soft ray 2; pectoral 1; ventral $1\frac{2}{5}$; caudal $1\frac{1}{4}$. Body elongate, not much compressed; caudal peduncle wide, wider than length of snout. Head small, broadly rounded anteriorly as viewed from above, snout blunt as viewed from the side; mouth very small, without so much lateral cleft as *Cottus philonips;* the maxillary reaching to front of pupil; teeth in moderately wide bands on jaws and vomer, palatines toothless, or with a few teeth in a narrow band on front; interorbital (bone only) equals $\frac{1}{2}$ eye; eye smaller than length of snout; preopercle with only 1 small blunt spine, below which the edge is entire.

* This species is thus described by Eigenmann & Eigenmann: Head about $3\frac{3}{4}$ to 4. D. VIII or IX, 16 to 18; A. II, 13; V. I, 4. Pectoral reaching anal or past vent even in largest specimens. Anal equidistant from tip of snout and base of caudal or nearer tip of snout. Ashy gray with blackish blotches. No well-defined cross bars excepting sometimes near the tail. Frequently a dusky blotch on anterior part of spinous dorsal and another near its posterior end; the fin sometimes wholly dusky, margined with white. Pectorals, soft dorsal, and caudal more or less barred. The types taken in the icy waters of the Kicking Horse, at Field, British Columbia, with *Coregonus coulteri*. (Eigenmann.)

Pectoral barely reaching front of anal; spinous dorsal very low, from ½ to ⅔ as high as soft dorsal, its base from its first spine to first ray of soft dorsal 1¼ in head; dorsals barely meeting, not at all connected. Color light gray, somewhat mottled; ventrals and anal colorless, other fins crossed with wavy lines; a soft spot on each end of spinous dorsal. Eagle River, a tributary of Grand River, Colorado, Colorado River Basin. Here described from 4 specimens from 2½ to 3½ inches in length, collected at Gypsum, Colorado, from the Eagle River. It has hitherto been confounded with *Cottus semiscaber*, which was taken in abundance at the same place and recorded as "*Cottus bairdii punctulatus*." We have compared this species with specimens of *Cottus beldingii* from Birch Creek, Idaho, and with a cotype of *Cottus philonips* from Field, British Columbia. It differs from both of these in having the mouth and eyes smaller, and from the Field specimen in having a deeper body. (Name for Miss Anna Louise Brown, artist of the Hopkins Seaside Laboratory.)

Cottus annæ, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1896, 223. with plate, Eagle River, Gypsum, Colorado. (Coll. Jordan, Evermann, Fesler & Davis. Types, Nos. 1305, 1308, 1309, and 1310, L. S. Jr. Univ. Mus.)

2333. COTTUS SPILOTUS (Cope).*

This species is thus described: "D. VIII, 17; A. 13; V. I, 4; P. 15; branchiostegals 6. Entering the section with 5 ventral rays, and with an elongate body, resembling apparently the *bairdii*, except in its short and anteriorly situated ventral fins. In *wilsonii* the eye is smaller, and frontal width greater; the pectoral rays are branched, in the present species simple. In *richardsonii* the vent is said to be the median point of distance from the muzzle to the caudal fin; here it is much nearer the muzzle. In *cognatus* we are informed that the anal fin has a more posterior position. The length of the head is contained 3 times plus 1 orbital diameter from end of muzzle to base of caudal fin; said diameter enters 4½ times length of head, and is ⅓ greater than interorbital width. The head is slightly contracted laterally, and not so depressed as in *C. alvordii*, giving the orbits less vertical range. One preopercular spine. Insertion of pectorals oblique, rays undivided, reaching anus and anterior rays of second dorsal; ventrals below middle of pectoral, insertion in advance of dorsal, extending halfway to vent. Width of isthmus equal from border of (closed) premaxillary to opposite hinder margin of pupil. Dorsal outline low, regularly descending to near end of second dorsal. Greatest depth enters 5 times from end of muzzle to base of caudal. Lateral line disappears between middle and end of caudal. First dorsal low, first ray ⅔ of second, third, and fourth; anal begins opposite fourth ray of second dorsal. Caudal fin rather small, rays once divided. No trace of palatine teeth. Above brown, below yellowish, everywhere densely punctulated with darker, except between the vent and anterior to ventral fins; dorsal, caudal, and pectoral fins barred; anal yellowish; base of caudal and dorsal spots

* Dr. Bean records a fish from York Factory, Hudson Bay, under the name *Uranidea spilota*, Cope, "D. IX, 18; A. 12; V. I, 4. One specimen measuring 4 inches without tail, which is wanting. Vomerine teeth only." (Bean, Proc. U. S. Nat. Mus. 1881, 127.)

blackish; large lateral round spots of the same color sometimes in 7 or fewer cross-bars. Length 3 inches. Several specimens from Grand Rapids, on the Grand River, which flows into Lake Michigan." (Cope.) A little-known species apparently close to *Cottus beldingii*. On the types of this species we have the following notes: It has now no evident teeth on the palatines and the ventral rays are I, 3.* The skin is smooth, and the preopercular spine, although prominent and directed upward, is not hooked. The spots on the body are less sharply defined than in *Cottus ricei*, with which Jordan & Gilbert first identified it. (Jordan, Cat. Fish. N. A., 111.) (σπιλωτός, spotted.)

Uranidea spilota, COPE, Proc. Ac. Nat. Sci. Phila. 1865, 182, Grand River at Grand Rapids, Mich.; JORDAN & GILBERT, Synopsis, 954, 1883.

2334. COTTUS LEIOPOMUS, Gilbert & Evermann.

Head $3\frac{1}{2}$; depth $5\frac{1}{2}$. D. VII–17; A. 12; P. 13; V. I, 4. Eye $4\frac{1}{3}$; snout $3\frac{1}{6}$; interorbital width $7\frac{1}{4}$. Least depth of caudal peduncle about equaling length of snout; interorbital space very narrow, much less than diameter of eye; mouth small, maxillary reaching vertical from middle of eye, a wide strip visible laterally in the closed mouth. Preopercular spines entirely absent, the preopercular margin evenly rounded throughout, without prominence, and without the least trace of a spine. Vomer with a very narrow band of teeth; palatines naked; skin wholly naked; lateral line complete. Dorsal fins not joined unless at extreme base; fins all low, the pectorals barely reaching front of anal under third ray of second dorsal, its last ray under fourth from last ray of latter. Free portion of caudal peduncle (behind last anal ray) $1\frac{3}{8}$ in head; portion behind base of last dorsal ray 3 in head; neither dorsal nor anal reaching base of caudal when depressed. Color in alcohol, head on sides rather finely vermiculated with light and dark; plain whitish below; not coarsely spotted or blotched as in *C. philonips;* dorsal bars indistinct; 2 narrow black lines downward and backward from eye; an evenly convex dark bar at base of caudal; dorsals, pectorals, and caudal faintly cross-barred. Length about $3\frac{1}{4}$ inches. Upper Little Wood River, Shoshone, Idaho. Closely related to *Cottus philonips*, but remarkably distinguished by the lack of preopercular spines. (Gilbert & Evermann.) (λεῖος, smooth; πῶμα, opercle.)

Cottus leiopomus, GILBERT & EVERMANN, Bull. U. S. Fish Comm. 1894, 203, pl. 20, Upper Little Wood River, Shoshone, Idaho. (Type, No. 45389, U. S. Nat. Mus.; Cotype, No. 1151 L. S. Jr. Univ. Mus. Coll. H. H. Kinsey.)

2335. COTTUS PRINCEPS, Gilbert.

Head $3\frac{1}{2}$ to $3\frac{5}{6}$ in length; depth 5 to $5\frac{1}{3}$. D. VI or VII, 21 to 23. A. 16 to 18; V. I, 4; P. 15. A slender form with small, narrow head, which is nearly quadrate in cross section, the opercles and cheeks being subvertical, the greatest width of head but $\frac{1}{8}$ or $\frac{1}{6}$ more than its depth at occiput. Mouth small, oblique, the gape slightly curved, the maxillary reaching a vertical crossing eye in front of pupil, $2\frac{3}{5}$ to 3 in head. Eye equaling snout, $4\frac{1}{8}$ in

* This is perhaps an error in our notes. The type can not now be found in the collection at Philadelphia.

head. Teeth small, uniform, in narrow bands in the jaws; vomer with a narrow patch; palatines smooth. Eyes small, separated by a narrow, flat interspace, as wide as pupil. Margin of preopercle evenly rounded, without developed spine, a minute spinous point sometimes occupying the position of the upper preopercular spine; opercle without spine; tubes and pores of head extraordinarily developed. A series of 6 very large pores across cheek and on lower edge of preorbital; a large median pore at symphysis, and a series of 7 occupying each ramus and extending onto edge of preopercle; similar somewhat smaller pores form the supraorbital series. Branchiostegals 6. Gill membranes broadly united to the isthmus, without free fold; no pore behind last gill. Dorsal and anal fins very long and low, the dorsal spines very slender, the notch shallow between spinous and soft portions; pectorals reaching beyond front of anal; ventrals usually to vent. About ⅔ of our specimens have the back and sides completely invested with minute close-set prickles, the head and belly and a narrow area along base of anal naked; caudal peduncle also naked in varying degree. In the remaining third (possibly males) the body is smooth except for a postaxial band of prickles, and in 1 specimen these are absent, leaving the body entirely naked. Lateral line variously incomplete, interrupted at some point under posterior half of second dorsal. Color light olive with darker markings, which may, on the head, take the form of vermiculating lines; 7 quadrate dark blotches along base of dorsal fin, the first and third usually narrower than the others, an eighth on back of caudal peduncle; very distinctly marked individuals show a series of blotches along middle of sides, which may be connected with the dorsal series by broad dusky bars; dorsal, caudal, and pectoral with faint bars; ventrals and anal unmarked. We subjoin table of fin rays in 12 specimens:

	Dorsal spines.		Dorsal rays.			Anal.			Pectoral.	
	VI.	VII.	21.	22.	23.	16.	17.	18.	14.	15.
Specimens	3	9	7	4	1	2	7	3	1	11

Numerous specimens were obtained in shallow water along the shore of Upper Klamath Lake on a bottom of fine sediment and vegetable débris. This differs widely from any other species of *Cottus* in the very narrow, slender form, the long fins, and especially in the extreme development of the mucous tubes and pores. Klamath Lakes, Oregon. (Gilbert.) (*princeps*, chief.)

Cottus princeps, Gilbert, Bull. U. S. Fish Comm. 1897. 12, figure, Upper Klamath Lake, Oregon. (Type, No. 48227. Coll. Gilbert, Cramer, and Otaki.)

730. URANIDEA, De Kay.

Uranidea, De Kay, New York Fauna: Fishes, 61, 1842 (*quiescens*=*gracilis*).

This genus is very close to *Cottus*, from which it differs in the reduction of its ventrals to a concealed spine and 3 soft rays, a step further in the degeneration characteristic of fresh-water types. The skin is smooth, or

very nearly so, the preopercular spines small, and there is usually no trace of teeth on the palatines. Cold streams and springs of the United States from New England and the Great Lakes to the Pacific Coast. (*οὐρανός*, sky; *εἶδον*, I looked; *i. e.* stargazer.)

a. Palatines with teeth.
b. Dorsal VIII, 16; preopercle armed with 4 spines, the longest about 2 in eye: least depth of caudel peduncle 12 in length of body. BENDIREI, 2336.
bb. Dorsal VI, 19; preopercle with a single short spine; least depth of caudal peduncle 8⅓ in length of body. GREENEI, 2337.
aa. Palatines toothless.
c. Anal rays 14 or 15; dorsal spines 7 or 8.
d. Preopercular spine short, bluntish; first dorsal dark, with pale margin. MARGINATA, 2338.
dd. Preopercular spine well developed, broad, rather sharp, curved upward. TENUIS, 2339.
cc. Anal rays 12 or 11.
e. Preopercular spine curved upward more or less abruptly.
f. Head large, about 3⅓ in length.
g. Mouth small, the maxillary barely reaching past front of eye; pectorals short, not reaching anal; preopercular spine short, hooked abruptly upward, body very robust. FRANKLINI, 2340.
gg. Mouth larger, the maxillary reaching middle of eye; pectorals usually reaching past front of anal; preopercular spine stoutish, little hooked.
h. First dorsal nearly as high as second; anal prickly. KUMLIENII, 2341.
hh. First dorsal much lower than second; axil not prickly, so far as known; pectorals variable, usually long. GRACILIS, 2342.
ff. Head small, 4½ in length. FORMOSA, 2343.
ee. Preopercular spine sharp, nearly straight, directed backward rather farther than upward; head narrowed forward. HOYI, 2344.

2336. URANIDEA BENDIREI (Bean).

Branchiostegals 6; D. VIII, 16; A. 12; V. I, 3; P. 15; C. 15. The length of the type is 2.7 inches (69 mm.) to the base of the caudal. The greatest height of the body at the origin of the dorsal equals its greatest width over the base of the pectorals, and is contained nearly 4 times in its length without caudal. The least height of the tail is $\frac{1}{12}$ of the length without caudal. The length of the middle caudal rays is contained 4½ times in the length of the body. Width of the mouth equals ½ the length of the head. The maxillary extends a little behind the front margin of the eye. The snout is as long as the eye. The distance between the eyes is less than their diameter, and is contained 5 times in the length of the head; the diameter of the eye 4 times. The length of the lower jaw equals ½ the length of head. Preoperculum armed with 4 spines, the largest at the angle, and is about ½ as long as the eye; the 3 anterior ones are very small and hidden by the skin. The distance of the spinous dorsal from the snout is ⅓ of the length of the body; the length of the longest spine of the first dorsal is about ⅛ of that of the head and about ⅜ of the longest dorsal ray. The longest dorsal and anal rays are equal. The length of the anal base almost equals the length of the head. The length of the

second dorsal base is a little greater than the length of the head. The pectorals reach to the origin of the anal; ventrals do not reach to the vent. Vomerine and palatine teeth present. Skin smooth. Color uniform dark brown above, lighter on the throat and belly. (Bean.) Eastern Oregon and northern Idaho, in the Columbia Basin. ("I take pleasure in dedicating this species to Capt. Charles E. Bendire, an excellent collector and observer, who has contributed so much to the Museum.")

Potamocottus bendirei, Bean, Proc. U. S. Nat. Mus. 1881, 27, Walla Walla, Washington. (Type, No. 24196. Coll. Capt. Chas. Bendire.)
Uranidea bendirei, Jordan & Gilbert, Synopsis, 696, 1883.

2337. URANIDEA GREENEI, Gilbert & Culver, new species.

Head 3⅛ in length; depth 4½; eye 3¾ in head; snout 3¾; interorbital width (bone only) 9; caudal peduncle very deep, its least depth 2½ in head. Dorsal VI, 19; anal 12; pectoral 14 or 15, ventral I, 3. Head broad with rounded profile, viewed from above, the snout and interorbital region slightly convex, the occiput flat or slightly concave. Mouth large, horizontal, the maxillary extending as far back as vertical from center of pupil, 2⅓ in head. Vomer and palatines with broad bands of teeth. A single preopercular spine, rather short, directed somewhat obliquely upward. Lateral line incomplete, ending under fifteenth ray of second dorsal; a small patch of prickles behind the axil, and a few along base of second dorsal. Spinous dorsal high, its longest spine exceeding length of snout, 2⅓ in head; longest ray of second dorsal about 2 in head; last dorsal spine joined by membrane to base of first soft ray; last dorsal rays reaching with their tips to base of caudal; pectorals reaching to or beyond vertical from first anal ray; pectoral rays all unbranched; ventrals not quite reaching vent. Olive brown above and on sides, light below; 2 short dark bars under spinous dorsal, both, or the hinder one only, continued upward to form a black blotch on the fin; a conspicuous dark bar under anterior, and 1 under posterior third of second dorsal, with a shorter fainter one midway between them; a vertical bar at base of caudal; area below lateral line with a series of 6 or 7 vertically oblong dark blotches, more or less distinctly confluent at the upper limit, to form a wavy streak, which is also joined by the darker markings of the back; a blackish streak at base of pectoral; second dorsal, caudal, and pectorals rather broadly cross-barred with dusky; ventrals and anal colorless. Length 58 mm. This species is characterized by its short spinous dorsal and its very deep caudal peduncle. Snake River Basin, Idaho; only the type known. Type locality, Thousand Springs, Snake River, Idaho, near mouth of Salmon Fall River. Collected by C. H. Gilbert, C. W. Greene, and K. Otaki, August 9, 1894. ("Named for Prof. Charles Wilson Greene, of Stanford University, to whose energy was due much of the success of the expedition.")

2338. URANIDEA MARGINATA, Bean.

Branchiostegals 6; D. VII or VIII, 18 or 19; A. 14 or 15; V. I, 3; P. 13 or 14; C. 14 or 15. Head broad, slightly depressed, its greatest length contained a little more than 3 times in length of body without caudal (4

times in total length); distance from tip of snout to eye equals length of eye, which is 4 in head. Vomerine teeth present; none on the palatines. Body stout anteriorly, moderately compressed posteriorly, its height at origin of first dorsal equal to its width at the same place, and contained $4\frac{1}{2}$ times in length of body without caudal; the least width of the caudal peduncle less than $\frac{1}{2}$ its height. The distance between the eyes equals $\frac{1}{2}$ of their long diameter. The maxilla extends to the vertical through the end of the anterior third of the orbit. The preoperculum has a short, broad, rather blunt spine at its angle and a much smaller one on its lower limb; between these 2 the margin in some specimens is crenulated, sometimes forming an additional blunt spine. The preopercular spines are all hidden under the skin. The distance of the first dorsal from the snout is contained $2\frac{3}{5}$ times in length of body without caudal; its longest spine is $\frac{3}{5}$ as long as the head; the length of its base is contained 5 times in length of body without caudal, 6 times in total length; the length of the second dorsal base is $\frac{1}{5}$ of total length. The length of the anal base is $\frac{1}{5}$ of length without caudal. The length of the ventral is contained from 6 to $6\frac{1}{2}$ times in the total length. The length of the pectoral is $\frac{1}{4}$ of length of body without caudal. The colors have faded, but the ground color seems to be plumbeous, with occasional blotches of darker; the fins are more or less distinctly punctulated; the first dorsal is darker than the body and has a distinct white margin. (Bean.) Eastern Oregon, in the Columbia River basin. Well distinguished from related species by its long anal.

Gilbert & Evermann have the following notes on this species:

Six small specimens from Mill Creek at Walla Walla (the type locality of *marginatus*) agree with Bean's description and differ from all other western specimens of *Cottus* which we have seen, in having but 3 soft rays in the ventral fins. So far as can be ascertained from our very immature specimens, *marginatus* strongly resembles *perplexus*, with which it agrees in fin rays, naked skin, the incomplete lateral line, and the absence of the palatine teeth. *Cottus perplexus* has constantly 4 soft rays in the ventral fins, and other differences may appear when compared with adult specimens. In our specimens of *marginatus*, the anus varies in position, being sometimes nearer base of caudal fin than snout, sometimes nearer snout. Twenty-two small specimens, collected by Bean & Woolman at Sand Point, Idaho, are for the present referred to this species, though we are not certain that this identification is correct. The ventrals seem to be I, 3; but the body is more or less covered with prickles. (*marginatus*, edged.)

Uranidea marginata, Bean, Proc. U. S. Nat. Mus. 1881, 26, Walla Walla. (Coll. Capt. Charles E. Bendire.)

Cottus marginatus, Gilbert & Evermann, Bull. U. S. Fish Comm. 1894, 204.

2339. URANIDEA TENUIS, Evermann & Meek.

Head $3\frac{4}{5}$; depth 7; eye $4\frac{1}{2}$ in head; snout 4; V. 1, 3. D. VI–I, 17; A. 15; vomer with teeth; palatines toothless. Head long, contracted from eyes forward; snout rather long; body much compressed, very slender, greatest width of caudal peduncle $2\frac{1}{2}$ in eye; least depth of same slightly greater

than eye; preopercular spine well developed, broad, rather sharp, partly covered by skin, curved upward; below this 2 other spines, the anterior one blunt, the other sharp, directed toward lower base of pectoral; post-temporal spine well developed. Body smooth, wholly without prickles or scales; lateral line complete except on caudal peduncle. Color dark above and on upper ¾ of sides, pale below; dorsal fins barred with series of dark dots or blotches; caudal similarly marked but rather darker; anal light with a few dark blotches; pectorals same as caudal; ventrals plain; under side of head profusely covered with small round black specks; muciferous pores on head well developed. Klamath Lakes, Oregon. Besides the specimen described above we have 2 others of the same slender style from Pelican Bay, Upper Klamath Lake, and many others from the lake near Klamath Falls, the majority of which are much smaller and less slender. The slender ones are apparently entirely smooth; a ripe female 3 inches long has a few prickles on anterior part of body, while all the smaller ones are pretty well covered with small prickles. (Evermann & Meek.) (*tenuis*, slender.)

Uranidea tenuis, Evermann & Meek, Bull. U. S. Fish Comm. 1897, 83, fig. 6, Upper Klamath Lake, Klamath Falls, Oregon. (Type, No. 48229. Coll. Meek & Alexander.)

2340. URANIDEA FRANKLINI (Agassiz).

Head 3½; depth 4½. D. VIII, 17; A. 12; V. I, 3. Body rather short and stout; snout not very obtuse; maxillary reaching about to pupil; eye 4 in head, twice the interorbital space; preopercular spine hook-like, very acute; paired fins rather short, the pectorals not reaching vent; first dorsal nearly as high as second; dorsals contiguous; anal inserted under fourth ray of second dorsal; caudal 6 in length; lateral line incomplete; vent nearer base of caudal than tip of snout. Length 3 inches. Lake Superior. (Girard.) Perhaps not distinct from *Cottus gracilis*. (Named for Sir John Franklin, the Arctic explorer.)

Cottus franklini, Agassiz, Lake Superior, 303, 1850, north and east shores of Lake Superior (Coll. James Hall and Louis Agassiz), Girard, Monograph Cottoids, 53, 1851.
Uranidea franklini, Jordan & Gilbert, Synopsis, 954, 1883.

2341. URANIDEA KUMLIENII, Hoy.

D. VI, 17; A. 12; P. 14; V. I, 3. Body slender and elongated, its depth 6 times in length. Head large and long, 3⅓ in length of body, its width a little more than ½ its length, its depth a little less. Eye large, 3½ in head, about equal to snout, more than twice interorbital space. Base of pectorals crescentic, the fin about as long as head; the lower rays rapidly shortened, reaching second or third dorsal ray, and falling just short of anal, fourth and fifth rays longest. Vent equidistant between base of caudal and front of eye. Mouth pretty wide and oblique, the maxillary reaching to middle of eye; lower jaw projecting. No palatine teeth. Preopercular spine large, directed upward and backward, not strongly hooked. First dorsal rather high, ⅝ the height of the second, the second spine longest and filamentous, the 2 fins connected by a membrane. Caudal peduncle

slender. Caudal fin narrow, more than ¾ length of head. Second dorsal and anal high. Lateral line disappearing under middle of second dorsal. Color obliterated, traces of a black spot on spinous dorsal. Length of specimen 3 inches. One of Dr. Hoy's types, in very bad condition, examined by us. It has grown soft in weak alcohol, and its remarkable slenderness of body is perhaps, in part at least, due to this fact. Lake Michigan, in deep water. (Named for A. L. Kumlien, the ornithologist.)

Uranidea kumlienii, HOY, Nelson, Bull. Ills. Mus. Nat. Hist., Vol. I, No. 1, 1876, 41, Lake Michigan (Coll. Dr. Hoy); JORDAN, Proc. Ac. Nat. Sci. Phila. 1877, 64.

2342. URANIDEA GRACILIS (Heckel).

Head 3½; depth 5. D. VIII, 16; A. 12. Body rather slender, fusiform; preopercular spine moderate, concealed. Mouth rather large, the maxillary reaching to the pupil; pectorals reaching front of anal; ventrals about to vent. Color olivaceous, mottled, upper edge of spinous dorsal red in life. Streams of New England and New York, recorded from tributaries of the Connecticut, Lake Champlain, Hudson, Delaware, and Susquehanna; common; probably a variable form, including several of Dr. Girard's nominal species, as Dr. W. O. Ayres vigorously insisted in 1845, before the latter had been described. (*gracilis*, slender.)

Cottus gracilis, HECKEL, Ann. Wien Mus., II, 1839, 148, New York.
Cottus gobio, AYRES, Bost. Jour. Nat. Hist., V, 1845, 121.
Uranidea quiescens, DE KAY, New York Fauna: Fishes, 61, 1842, Lake Pleasant, Adirondack region, New York.
Cottus viscosus,* HALDEMAN, Suppl. Mongr. Limnea, 3, 1840, Susquehanna River (Coll. S. S. Haldeman); GIRARD, Monograph Cottoids, 51, 1851.
Cottus gobioides,† GIRARD, Proc. Amer. Assoc. Adv. Sci. 1850, 41, and *l. c.*, 55, Lamoille River, Johnson, Vermont. (Coll. Ransom Colberth.)
Cottus boleoides,‡ GIRARD, Proc. Amer. Assoc. Adv. Sci. 1850, 41, and *l. c.*, 56, Windsor, Vermont (Coll. Ed. Cabot); GÜNTHER, Cat., II, 156, 1860.
Cottus copei, ABBOTT, Proc. Ac. Nat. Sci. Phila. 1861, 15, no locality given, probably New Jersey.
Cottus gracilis, GIRARD, Monograph Cottoids, 49, 1851.
Uranidea gracilis, JORDAN & GILBERT, Synopsis, 699.
Uranidea viscosa, JORDAN & GILBERT, Synopsis, 698.
Uranidea gobioides, JORDAN & GILBERT, Synopsis, 699.
Uranidea boleoides, JORDAN & GILBERT, Synopsis, 699.

* *Cottus viscosus* is thus characterized:
Head 3½ in length; depth 4¾. D. VI, 18; A. 12; V. I, 3. Body rather stout; mucous pores on head usually numerous. Mouth small; maxillary reaching middle of eye; preopercular spine acute, extremely short, directed obliquely upward; isthmus rather narrow; first dorsal low, slightly connected with second; pectorals shorter than head, reaching second dorsal. Color olivaceous, variegated, the spinous dorsal edged with orange in life; fins mostly barred. Eastern Pennsylvania and Maryland, in tributaries of the Schuylkill, Susquehanna, and Potomac rivers; not rare, entering caves.

† *Cottus gobioides* is thus described:
Head 3¾; depth 4¾. D. VII, 17; A. 12. Similar to *Uranidea gracilis*, but larger and more robust, the mouth larger, reaching to beyond the line of the pupil, and the pectorals short, not quite reaching anal; preopercular spine stout, curved suddenly upward. Length 4 inches. Tributaries of Lake Champlain. (Girard.)

‡ *Cottus boleoides* is thus characterized:
Head 3½; depth 5½. D. VIII 17; A. 11. Slender, subfusiform. Mouth rather large, the maxillary extending to opposite the pupil; preopercular spine acute, directed obliquely upwards; isthmus narrow. Fins larger than in any other species; pectorals reaching fourth ray of anal. Length 3¾ inches. Connecticut Valley in Vermont. (Girard.)

2343. URANIDEA FORMOSA (Girard).

Head $4\frac{1}{4}$. D. VIII–16; A. 11; V. I, 3. Body slender and graceful; head small, depressed above; eye moderate; preopercular spine short, stout, acute, curved upward, a small spine below it; subopercular spine well developed. Dorsals well separated; anal beginning under third ray of soft dorsal; pectorals not reaching to posterior margin of spinous dorsal; ventrals not nearly to vent. Length $3\frac{1}{4}$ inches. Deep water in Lake Ontario. (Girard.) A doubtful species, known only from 1 mutilated specimen, the head shorter than in *Uranidea gracilis*. (*formosus*, pretty.)

Cottus formosus, GIRARD, Monograph Cottoids, 58, 1850, Lake Ontario off Oswego, in stomach of *Lota maculosa*. (Coll. S. F. Baird.)
Uranidea formosa, JORDAN & GILBERT, Synopsis, 955.

2344. URANIDEA HOYI, Putnam.

D. VI, 15; A. 11; V. I, 3; C. 12; the first dorsal low and small, $\frac{2}{3}$ as long as soft part, and connected to it by membrane. Fins all low. (Hoy.) Body small, rather short and thick, the depth about $5\frac{1}{2}$ in length. Head $3\frac{1}{8}$ in length to base of caudal. Eyes large, high up, and close together, about equal to snout, $3\frac{2}{3}$ in head, 3 times the interorbital space. Head rather narrow and contracted; the lower jaw narrowed and slightly projecting. No palatine teeth; maxillary reaching anterior border of eye, or a little beyond. Preopercular spine prominent, sharp, very nearly straight, directed backward, but not strongly upward, its form and direction peculiar in this genus. Below this is another sharp, prominent spine, also nearly straight, directed partly downward; one or 2 minute concealed spines still lower. Isthmus well developed. Pectoral fins long, longer than head, reaching beginning of anal; ventral fins long, reaching almost to vent. Vent midway between snout and base of caudal, in the female farther back, owing to the distended abdomen. Ventral rays I, 3 (on one side of each specimen apparently I, 4; perhaps a soft ray has been split). Color nearly obliterated. Lower parts profusely punctate. Length of specimens $2\frac{1}{5}$ inches. Lake Michigan in deep water. Female specimen taken 12 miles off Racine, Wisconsin, in 12 fathoms, June 4, 1875, by Dr. Hoy; the male off Milwaukee, June 15. The specimens are now in bad condition from rough handling. The female is distended with ripe eggs, so that the width of the body is $\frac{1}{3}$ the total length. This species seems to be quite distinct from all those described by Girard. The peculiar characters are the number and form of the preopercular spines, the contracted mouth, the large eyes, the small size of the body, and the length of the ventral fins. The characters first mentioned are the most striking. (Named for Dr. Philo R. Hoy, of Racine.)

Uranidea hoyi (PUTNAM MS.) NELSON, Bull. Ills. Mus. Nat. Hist., Vol. I, No. 1, 1876. 41, Lake Michigan (Coll. Philo R. Hoy); JORDAN, Man. Vert., 244, 1876; JORDAN & COPELAND, Bull. Buff. Soc. Nat. Sci. 1876, 41; JORDAN, Proc. Ac. Nat. Sci. Phila. 1877, 63; JORDAN & GILBERT, Synopsis, 700, 1883.

731. MYOXOCEPHALUS (Steller) Tilesius.

(Great Sculpins.)

Myoxocephalus, Steller MS. 1741.
Myoxocephalus, Tilesius, Mém. Acad. Sci. Petersb., iv, 1811, 273 (*stelleri*).
Acanthocottus, Girard, Proc. Bost. Soc. Nat. Hist., iii, 1849, 185 (*grœnlandicus*).
Cottus, Putnam, Bull. Mus. Comp. Zool., I, No. 1, 3, 1863 (*scorpius*).
Boreocottus, Gill, Proc. Ac. Nat. Sci. Phila. 1859, 166 (*axillaris*).

Body slender or robust, subfusiform, covered with thick skin, in which are sometimes embedded prickly plates; deciduous, granular, or stellate tubercles also sometimes present, but no true scales. Head large. Mouth terminal, large, the lower jaw always included, the uppermost the longer; villiform teeth on the jaws and vomer, none on the palatines; suborbital stay strong; preopercle with 2 strong straight spines above directed backward, and 1 below directed downward and forward; opercle, nasal bones, orbital rim, and shoulder girdle more or less armed; gill membranes forming a fold across the rather narrow isthmus; slit behind last gill small or wanting, if present, reduced to a mere pore; vertebræ about 28. Branchiostegals mostly 6. Dorsal fins 2, separate, the first short, its spines rather slender; ventral rays I, 3; caudal fin moderate, fan-shaped; pectoral fin broad, its lower rays procurrent. Lateral line well developed, its tubes sometimes provided with bony or cartilaginous plates, never chain-like nor reduced to separated pores. Species numerous, in the seas of northern regions; coarse fishes, little valued as food. (*μυοξός* the dormouse; *κεφαλή*, head; the allusion not evident).

Acanthocottus (*ἄκανθα*, spine; *κόττος*, *Cottus*):

a. Top of head covered with thin, smooth skin, having a very few warty prominences or none, and not hiding the bony occipital ridges; lateral line continuous, with concealed cartilaginous plates; Atlantic species.

b. Lateral line with regularly arranged, small bony-keeled plates, most distinct anteriorly; no stellate tubercles; a small cirrus above eye. D. VIII, 12; A. 9. bubalis, 2345.

bb. Lateral line unarmed or with concealed cartilaginous plates or with prickles which are not keeled or are not regularly arranged.

c. Anal fin very short, of 10 or 11 rays only; dorsal rays IX, 13 or 14; upper preopercular spine short, sharp, nearly twice the length of the next; third spine directed downward; no slit behind last gill. æneus, 2346.

cc. Anal fin of 12 to 14 rays.

d. Upper preopercular spine moderate, about as long as eye, and about twice length of next spine.

e. Last gill arch without slit or pore. D. X, 17; A. 12. scorpioides, 2347.

ee. Last gill arch with a small slit or pore behind it. D. X, 17; A. 14.

f. Interorbital space narrow, 1⅔ in eye in adult; spinous dorsal low, the longest spine 8 in body. scorpius, 2348.

ff. Interorbital space broader, as wide as eye; spinous dorsal higher, the longest spine 6 to 7 in body. grœnlandicus, 2349.

dd. Upper preopercular spine very long, longer than eye, reaching tip of opercular spine, 4 times length of the spine below it; a sharp postocular spine, and a similar occipital spine; no slit behind last gill. D. IX, 15; A. 14. octodecimspinosus, 2350.

MYOXOCEPHALUS:

aa. Top of head covered with small warty prominences.

g. Supraocular spine with 2 or 3 diverging digitate ridges behind it; skin of top of head thin, not concealing the bony occipital ridges; lateral line continuous; upper preopercular spine long; no stellate plates on sides below lateral line; nasal spines sharp; no cirri on head.

h. Distance from supraorbital to occipital tubercle about equal to space between supraorbital tubercles; anal rays usually 12; membrane of maxillary more or less mottled. POLYACANTHOCEPHALUS, 2351.

hh. Distance from supraorbital to occipital tubercle 1½ times the distance between supraorbital tubercles; head very long and flat; dorsals well separated; anal rays usually 14; adult with round rough plates above; throat white; membrane of maxillary unspotted. JAOK, 2352.

gg. Supraocular spine without digitate spines behind it; preopercular spines short.

i. Warts on top of head not ending in cirri; nasal spines and occipital ridges not wholly concealed in the thick skin.

j. Skin of sides of back in adult covered with round stellate tubercles; spines of preopercle short, more or less hidden in the skin.

k. Supraocular spine blunt, with a blunt tubercle behind it, the cirri small or obsolete; crests moderate; soft dorsal very long, of 15 to 19 rays. VERRUCOSUS, 2353.

kk. Supraocular spine small, blunt, without tubercle behind it; supraocular and occipital spines each with a small flat cirrus; occipital crests very low; soft dorsal of 15 or 16 rays. AXILLARIS, 2354.

jj. Skin of back and sides perfectly smooth (except in very old individuals which may have hidden rough plates below the lateral line).

l. Occipital crests very low, almost concealed by the thick warty skin; body robust, the depth 4 in length; membrane of upper jaw with sharply defined black spots; throat always spotted; suprascapular spine sharp; humeral spine obscure.

m. A pale bar at base of caudal and 1 behind anal. STELLERI, 2355.

mm. No pale bar at base of caudal and none behind anal. MEDNIUS, 2356.

ll. Occipital crests strong, the space between them concave, the skin thin and with rather few warts; body rather slender, the depth 4⅓ in length.

n. Sides with round white spots. NIVOSUS, 2357.

ii. Warts on top of head numerous, each ending in a small cirrus; occipital crests obsolete, hidden in the thick skin; nasal spines also hidden in the thick skin; preopercular spines small, almost concealed, the second tubercle-like; spines on shoulder girdle obsolete; head broad, mouth large; skin entirely smooth, color very dark. NIGER, 2358.

Subgenus ACANTHOCOTTUS, Girard.

2345. MYOXOCEPHALUS BUBALIS (Euphrasen).

(FATHER-LASHER: LUCKY PROACH.)

Head 2⅔; depth 3⅔. D. VIII, 12; A. 9; vertebræ 12+17. Head broad, depressed, covered with soft skin, in which are many mucous pores; maxillary reaching past middle of orbit; upper preopercular spine straight,

slightly longer than eye; opercular spine granulated at base; small cirri above eye and elsewhere on head, usually 1 on end of maxillary; cranial bones mostly covered by skin; lateral line with some bony plates, which are most distinct anteriorly, these very much smaller than in *Enophrys bison*, but somewhat similar in structure; interocular space very narrow, its ridges continuing backward, serrated, each ending in a sharp spine; no trace of slit behind last gill; spinous dorsal low; anal small; pectorals reaching front of anal; ventrals moderate. Colors variegated, the dark markings sometimes red, corresponding to the hue of red algæ. Coasts of northern Europe; abundant in rock pools; said to stray to Greenland, but there is no sure evidence of its occurence in any American waters. Lütken refers the Greenland records to *M. scorpius*. (Eu.) (*βούβαλις*, buffalo.)

Cottus bubalis, Euphrasen, Köng. Vet. Nya Abhandl. 1786, 65, Taf. 3, fig. 2, 3, Sweden; Günther, Cat., II, 164; Day, Fish. Gt. Brit. Ireland, 51, 1880; Jordan & Gilbert, Synopsis, 701; Bean, Proc. U. S. Nat. Mus. 1885, 166.

Cottus maculatus, Fischer,* Jahrb. Wissensch. Aust., Hamburg, II, 78, taf. 2, fig. 8, 1885, Barbados.

2346. MYOXOCEPHALUS ÆNEUS (Mitchill).

(Grubby.)

Head $2\frac{2}{4}$; depth 4. D. IX, 13 or 14; A. 10 or 11; V. I, 3. Head rather broad, covered with smooth thin skin; no cirri; a few very small warts between occipital ridges; maxillary $2\frac{1}{3}$ in head, reaching to just beyond pupil; supraocular and occipital ridges prominent, each with a low, bluntish spine; the region between the supraocular spines rather convex, the space before and behind it concave; nasal spines moderate; upper preopercular spine shorter than eye, nearly twice length of next spine, about reaching middle of opercle. Lateral line complete; each pore with a concealed cartilaginous plate, scattered, concealed asperities on skin of sides;

* Concerning the identity of *Cottus maculatus*, Fischer, with *Acanthocottus bubalis*, Dr. Tarleton H. Bean has the following pertinent remarks:

"In a recent extract from the annals of the Scientific Association of Hamburg, Dr. J. G. Fischer has described and figured a species of *Cottus* from Barbados. The anomaly of finding a species of this genus within the tropics induced me to examine the description very carefully, to ascertain, if possible, the source of Dr. Fischer's specimen. There is no reasonable doubt that the type of his new species is the common Father-lasher of Europe, and it is a source of wonder that the species should have been misinterpreted. After a study of our examples of *Cottus bubalis* from Bergen and Christiania, in Norway, and Leeds, England, I have no hesitation in stating that they agree perfectly with the description and figure of *Cottus maculatus*. In some unknown way the locality of Dr. Fischer's specimen has been incorrectly given, and thus the describer of the supposed new species has been completely misled. The differential characters claimed by Dr. Fischer for *Cottus maculatus* do not serve at all to separate his species from *C. bubalis*. The arrangement and number of the spines on the preoperculum are precisely the same in our examples of *Cottus bubalis* as represented in the figure of *C. maculatus*. The ventral and pectoral are not longer in our specimens than they are made to appear in the figure of the alleged new species. The notion seems to have got abroad among the European ichthyologists that North America is a comparatively benighted and barbarous country, whose natural history is still in its infancy; it is perhaps owing to this impression that we are startled by information concerning the supposed occurrence of *Cottus* within the tropics, and of a species of *Platycephalus* in the Potomac River. (*Platycephalus americanus*, Sauvage, Nouv. Archiv. Mus. (2), pl. 2, fig. 3 (head only). Potomac River.) Our common little *Cottus œneus* of Mitchill has also been redescribed, from a New York specimen, under the name of *Cottus* (*Acanthocottus*) *anceps*. (*Cottus* (*Acanthocottus*) *anceps*, Sauvage, Nouv. Archiv. Mus. Hist. Nat., Paris (2), I, 1878, p. 145, pl. 1, fig. 13.) As a general rule it will be safe to intrust the novelties of fish distribution in our country to its resident ichthyologists."

dorsal spines rather low, higher than the soft rays; pectorals reaching anal; no trace of slit behind last gill. Grayish olive, much variegated with darker; no distinct paler spots; back and sides with broad, dark, irregular bars; all the fins barred; mandible mottled; throat and belly pale; membrane of maxillary unspotted. Length 6 to 8 inches. Coast of southern New England and New York; our smallest species; common in seaweeds near shore, but having a very narrow range. (*æneus*, brassy.)

Cottus æneus, MITCHILL, Trans. Lit. and Phil. Soc. N. Y., I, 1815, 381, New York; GOODE & BEAN, Bull. Essex Inst., XI, 13, 1879; JORDAN & GILBERT, Synopsis, 702.

Cottus scorpio, MITCHILL, Trans. Lit. and Phil. Soc., I, 1815, 381, New York.

Cottus mitchilli, CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 188, 1829, New York; after *Cottus scorpio*, MITCHILL; GÜNTHER, Cat. Fish., II, 164.

Cottus (*Acanthocottus*) *anceps*, SAUVAGE, Nouv. Archiv. Mus. Paris. (2), 1, 1878, 145, pl. 1, fig. 13, New York.

Cottus nigricans, LE SUEUR, in VAILLANT, Bull. Soc. Philom., serie 8, tome VIII, No. 1, 1895, 12; no locality, with unpublished plate of LE SUEUR, 1819.

2347. MYOXOCEPHALUS SCORPIOIDES (Fabricius).

(ARCTIC SCULPIN.)

Head 3; depth 4½. D. IX or X, 17; A. 12; V. I, 3; lateral line 38. Body stoutish, tail slender. Head moderate, not depressed; jaws short, the maxillary reaching middle of the large eye; lower jaw included; top of head concave between the 2 occipital ridges; these very low, with obtuse tubercles instead of spines, these ending each in a small tentacle, which is often obsolete; preopercular spines 3, short, the upper equal to eye; the second about ½ as long, the lower directed downward; opercular spine obscure; a short humeral and a short suprascapular spine; nasal spines sharp; top of head with smooth skin. Skin nearly smooth, sometimes a few warts above lateral line. Isthmus narrow, the fold across it very narrow; no trace of slit or pore behind last gill. Dorsal fins slightly joined, the spines slender; pectorals reaching past front of anal. Coloration, dark olive, finely mottled with paler; fins dusky, with paler spots; anal with 2 oblique dark bars; a pale blotch at base of caudal. Arctic regions of America; Greenland and neighboring waters; common about Disco Bay, from which locality we have numerous specimens, received from Prof. D'Arcy W. Thompson (Coll. Lohmann). (*σκορπίος, scorpius; εἶδος*, likeness.)

Cottus scorpioides, FABRICIUS, Fauna Grœnl., 157, 1780, Greenland; BEAN, Bull. U. S. Nat. Mus., XV, 122; JORDAN & GILBERT, Synopsis, 702; LÜTKEN, Aft. Vidensk. Meddels. Kjöbenh., 12, 1876.

Cottus pachypus,* GÜNTHER, Cat. Fish., II, 161, 1860, Port Leopold. (Haslar Coll.)

* *Cottus pachypus*, GÜNTHER, is thus described:

"D. IX, 16; A. 12 to 14. Two small spines above the snout, an obtuse one above each orbit, and a pair of obtuse prominences on the occiput. The impression on the crown becomes narrower posteriorly, and is nearly twice as long as broad. Three preopercular spines, 2 of which are at the angle; the upper is the longest, its length being equal to the diameter of the eye, but less than the width between the orbits. The height of the first dorsal is somewhat more than the length of the maxillary bone; the ventral terminates at a great distance from the vent, and the pectoral does not reach to the anal. Skin above the lateral line with smooth warts. Brownish, spotted with darker (colors faded). Port Leopold.

"Description of the specimen: The greatest height of the body, in front of the dorsal, is 4⅔ in the total length; its greatest width, behind the pectorals, 4 times. The tail tapers

2348. MYOXOCEPHALUS SCORPIUS (Linnæus).

(European Sculpin; Ulke.)

Head 2¼; depth 4¼. D. X, 16; A. 14. General characters of *Myoxocephalus grœnlandicus*, from which it differs chiefly in the smaller size, the narrower interorbital space, which is ⅝ diameter of eye, and in the lower spinous dorsal, the highest spines being about ⅓ the length to base of caudal; pore behind last gill usually very small, but evident. Dark olivaceous, mottled with paler; fins dusky, with paler spots. Northern Europe and Arctic regions, not common on our coasts; recorded by Dr. Lütken from the Baltic, Finland, Spitzbergen, New Zembla, coasts of England, and Northern Asia; recorded from Eastport, Maine. (Eu.) (*σκορπίος*, scorpion, the root of our word sculpin.)

Ulka. Linnæus, Iter Scand., 325.

Cottus scorpius, Linnæus, Syst. Nat., Ed. x, 265, 1758, after Iter Scand.; Günther, Cat. Fishes, II, 159; Bean, Bull. U. S. Nat. Mus., xv, 116; Jordan & Gilbert, Synopsis, 702; Lilljiborg, Sveriges och Norges Fiskar, 135, 1891.

2349. MYOXOCEPHALUS GRŒNLANDICUS (Cuvier & Valenciennes).

(Daddy Sculpin.)

Head 2½; depth 4½. D. X, 17; A. 14; V. I, 3; P. 18. Head large. Eye large, equal to least interorbital width (in specimens a foot long). Mouth large, the lower jaw included; maxillary reaching posterior edge of orbit 2⅓ in head; the supraorbital and occipital spines blunt, tubercle-like, without cirrus; a small tubercular spine on front of occipital ridge; upper preopercular spine short, only reaching the middle of opercular spine, its length equaling eye, not twice that of the spine below it; opercular spine

much posteriorly, and its height, before the caudal, is 21 times in the total length. The head is moderately broad and depressed, its length is 3⅜ in the total. The cleft of the mouth is moderate, the maxillary reaching to the vertical from the center of the eye; the snout is not very obtuse, and the upper jaw slightly overhangs the lower; there is a pair of small spines above the snout. The space between the eyes is slightly concave, and its width rather more than the horizontal diameter of the eyes; there is an obtuse protuberance above the posterior angle of the orbit, from which a slight ridge proceeds to the occipital protuberance; the latter is very little prominent, and furnished with small skinny tentacle. The impression between these 2 pairs of protuberances is shallow, and becomes narrower posteriorly, being nearly twice as long as broad. The preoperculum is armed with 3 spines, 2 of which are opposite the infraorbital, the third being situated at the inferior extremity of the bone, pointing downward. The suboperculum has a single spine anteriorly, directed downward; throat without spines. The spinous dorsal begins at a distance from the head which is rather less than the length of the impression of the crown; it is not continuous with the soft. The fourth spine is the longest, its length being somewhat less than that of the maxillary bone, and a little less than that of the ninth ray. The caudal has the posterior margin slightly convex, and its length is 6⅓ in the total. The anal is formed by 12 rays, and begins in the vertical from the fifth dorsal ray, and terminates a little before the opposite fin; its height equals that of the spinous dorsal. The pectoral is composed of 16 simple rays, and reaches to the vent, which is situated nearly in the middle of the total length. The root of the ventrals falls behind that of the pectoral; they terminate at a great distance from the vent, and are composed of 1 spine and 3 rays. The spine is enveloped in the same membrane with the first ray; the middle ray is the longest, flat and compressed like the interior. The lateral line is continued to the caudal, and is composed of elongate bony tubes, 38 in number. This specimen is nearly entirely smooth, exhibiting, however, some flat and smooth warts above the lateral line; it is a female. The young specimen differs from the adult one in having 14 anal rays. The colors have somewhat faded; the upper parts are brown, with some darker spots; the pectoral, dorsal, and caudal fins appear to have been spotted, the spots being arranged in bands; the lower part of the sides with several white spots."

sharp; nasal spines sharp; suprascapular spine rather strong, shortish. Sides of body above lateral line with a series of embedded prickly plates, below which are numerous scattered spines and prickles. Dorsal and anal fins high; spinous and soft dorsals about equal in height, their height more than ⅓ length of body; ventrals long; pectorals about reaching vent. Dark brown above, with broad darker bars; below yellowish, the belly in the male with large pale spots; back and top of head with grayish blotches; fins brown and yellow, all of them spotted and barred. Sexual differences large, the males more brightly colored; the round white spots strongly marked, the females with rough crests on the head. Length 25 inches. New York to Greenland, common; one of the largest sculpins. Here described from specimens from Cape Cod. Very close to *Myoxocephalus scorpius*, which it replaces on our coast and of which Dr. Lütken regards it as a variety. Lilljiborg regards the 2 as identical, a view not unlikely correct. According to Ensign H. G. Dresel, *M. grœnlandicus* differs from *M. scorpius* (1) in its larger size; (2) in the greater interorbital width which in *M. scorpius* seldom exceeds ⅛ of the longest diameter of the eye; and (3) in the higher spinous dorsal, the longest dorsal spine in *M. grœnlandicus* being contained 5 to 6 times in the total length of the caudal base, while in *M. scorpius* it is contained as much as 7 to 8 times in the same length.

According to Fabricius this species is abundant "in all the bays and inlets of Greenland, but prefers a stony coast clothed with seaweed. It approaches the shore in spring and departs in winter. It is very voracious, preying on everything that comes in its way, and pursuing incessantly the smaller fish, not sparing the young of its own species, and devouring crustacea and worms. It is very active and bold, but does not come to the surface unless it be led thither in pursuit of other fish. It spawns in December and January, and deposits its red-colored roe on the seaweed. It is easily taken with a bait, and constitutes the daily food of the Greenlanders, who are very fond of it. They eat the roe raw."

According to Lütken, the Greenland Sculpin (*grœnlandicus*) is not separated from *scorpius* by any character trenchant or constant. In Greenland it inhabits slight depths up to 12 fathoms; the young in 40 to 100 fathoms. There is great variation, especially in fin rays. In counting extreme cases both forms have the following range: D. IX to XI, 14 to 19; P. 15 to 19; A. 11 to 15. Normally, however, the range is D. X, 16 or 17; P. 17; A. 13 or 14. Vertebræ usually 14+25=39, in a rare case 13+23=36. Pyloric cæca 8, rarely 10 or 11.

Cottus scorpius, FABRICIUS, Fauna Grœnlandica, 156, 1780, Greenland.

Cottus grœnlandicus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 156, 1829, after FABRICIUS; GÜNTHER, Cat. Fish., II, 161, 1860; GOODE & BEAN, Bull. Essex Inst., XI, 13, 1879.

Cottus scorpius grœnlandicus, LÜTKEN, Vid. Medd. Kjöb., 12, 1876; JORDAN & GILBERT, Synopsis, 703; DRESEL, Proc. U. S. Nat. Mus. 1884, 252.

Cottus porosus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., VIII, 498, 1831, Baffins Bay. (Coll. Guédon and Ducrost.)

Acanthocottus mucosus, AYRES, Proc. Cal. Ac. Nat. Sci. 1854, 12.

Ananthocottus variabilis, GIRARD, Boston Journ. Nat. Hist., VI, 1850, 248; D. H. STORER, Hist. Fish. Mass., 26.

Acanthocottus ocellatus, H. R. STORER, Bost. Journ. Nat. Hist. 1850, 253; male.
Cottus glacialis, RICHARDSON, in BELCHER, Last Arctic Voyage H. M. S. Assistance, 1853, App. 3, pl. 24, Northumberland Sound. (Coll. Sir Edward Belcher.)

2350. MYOXOCEPHALUS OCTODECIMSPINOSUS (Mitchill).

(LONG-SPINED SCULPIN.)

Head $2\frac{1}{2}$; depth $5\frac{1}{4}$. D. IX, 15; A. 14; V. I, 3. Body very slender, tapering backward to the long and slender caudal peduncle. Head long and narrow; mouth moderate, the lower jaw included; maxillary not extending past eye, $2\frac{1}{4}$ in head. Upper preopercular spine extremely long, longer than in any other species of the genus, $3\frac{1}{8}$ in head; longer than eye, extending to or beyond tip of opercular spine, its length more than 4 times that of the short spine below it; third spine turned downward; nasal spines strong; a strong spine at upper posterior margin of orbit directed upward and backward; occipital ridges long, low, converging behind, each ending in a similar spine, sharp and directed backward; a single sharp suprascapular spine; a sharp spine on shoulder girdle at tip of the sharp opercular spine; the usual downward-directed spines on preopercle and subopercle. Vertex nearly flat, bounded by low ridges. No cirri on head; skin of head smooth. Eye very large, as long as snout, $4\frac{1}{2}$ in head, much wider than the concave interorbital space. Skin without rough tubercles; lateral line with a series of partly concealed cartilaginous plates. Dorsal fins well separated; spinous dorsal higher than soft dorsal, the spines strong, the longest nearly $\frac{1}{2}$ head; pectorals reaching anal; ventrals not to vent; no trace of slit behind last gill. Olivaceous, with 4 obscure transverse dark bars, paler below; fins barred and mottled; jaws unspotted; lower side of head plain white; belly white. Length about a foot. Atlantic coast, from Labrador south to Virginia, common about Cape Cod; a strongly marked species, easily known by its long spine and by the sharp spine at the occiput, there being 18, really 20, distinct spines about the head. (*octodecim*, eighteen; *spinosus*, spined.)

Scorpius virginianus, WILLUGHBY, Hist. Pisc., App., 25, pl. 10, fig. 15 (nonbinomial).
Cottus octodecimspinosus, MITCHILL, Trans. Lit. and Phil. Soc. N. Y., I, 1815, 380, New York; GÜNTHER, Cat., II, 163.
Cottus virginianus, STORER, Rept. Fish. Mass., 18, 1838, coast of eastern Massachusetts; DE KAY, New York Fauna: Fishes, 51, pl. 5, fig. 13, 1842; JORDAN & GILBERT, Synopsis, 701, 1883; GIRARD, Proc. Bost. Soc. Nat. Hist., III, 187.
Acanthocottus virginianus, STORER, Hist. Fish. Mass., 28, pl. 4, fig. 2, 1867.

Subgenus MYOXOCEPHALUS.

2351. MYOXOCEPHALUS POLYACANTHOCEPHALUS, Pallas.

(GREAT SCULPIN; KALOG.)

Head $2\frac{1}{2}$; depth $4\frac{1}{4}$. D. IX or X, 13 to 16; A. 11 to 13. Body rather elongate. Head long and narrow, somewhat compressed, concave between the orbits, the orbital rim being elevated; lower jaw included; nasal spines strong; a strong ridge above eye, with a blunt, compressed spine behind it; behind this a small digitate cluster of ridges ending in low

spines; behind these an irregular broken ridge on each side of the vertex, extending to the occiput; outside of this another ragged ridge; suborbital stay strong, striate; upper preopercular spine very long, straight, simple, striate at base, longer than eye; a similar but shorter spine below it, not ½ as long, and the usual downward-directed spine at lower edge of preopercle and subopercle; opercular spine moderate. Skin of top of head thin, with small, smooth warts, not hiding the occipital ridges; no cirri. Mouth rather large, the maxillary reaching beyond eye, ½ length of head; skin of body with some scattered rough tubercles, usually nearly smooth. Dorsals not very high; dorsal spines slender; pectorals reaching anal; ventrals moderate, I, 3. A minute pore usually present behind last gill, this sometimes wholly wanting. Lateral line complete. Dark olive above, much variegated with darker and reddish; belly mostly whitish; sides and belly (in males) with numerous blackish reticulations surrounding large round white spots; jaws dusky, mottled with whitish; membrane joining maxillary to preorbital black, with round white spots in the adult, more or less plain in the young; fins, all but the ventrals, mottled and barred with blackish and yellowish. Length 1½ to 2½ feet. Alaska to Kamchatka; abundant throughout Bering Sea, and southward along the islands to Puget Sound; one of the largest sculpins and everywhere familiar to fishermen. Dr. Gilbert found it abundant about Unalaska and in Bristol Bay. Mr. Scofield found it at Chignek Harbor, and we obtained it about the Pribilof* and Commander Islands and at Petropaulski. (*πολυς*, many; *ἄκανθα*, spine; *κεφαλή*, head.)

Cottus polyacanthocephalus, PALLAS. Zoogr. Rosso-Asiat., III, 133, 1811, Aleutian Islands; no definite locality; GÜNTHER. Cat., II, 166; JORDAN & GILBERT, Synopsis, 704, 1883.

2352. MYOXOCEPHALUS JAOK (Cuvier & Valenciennes).

D. VIII to X, 15 to 17; A. 13 to 15 (usually D. IX, 16; A. 14). This species has a very slender body and an extremely wide, flat head, the latter strikingly triangular when viewed from above, on account of the regular way in which it tapers toward the snout. The species is further distinguished by possessing but 9 dorsal spines and by the presence in the adult of an irregular series of circular spinous plates above the lateral line, these plates wanting in very young individuals. They begin to make their appearance in specimens 6 inches long, and are invariably present in larger specimens. In adults, the region below the lateral line contains strong spinous prickles mostly concealed in the skin and directed backward. Some of the anterior ones may be broader and may have more than one point, but none is circular with a rosette of short spinous points, as is the case with the dorsal series. Lower jaw included; top of head covered with small warts; scapular spine short and sharp; humeral spine obscure; upper preopercular spine very long, nearly as long as eye, low, sharp, 3 times length of next spine, not quite reaching tip of opercular

* Several specimens from Unalaska, where the species is abundant, and 1 from Robben Island, the latter perfectly typical in all respects and giving us the first Kamchatkan record for the species. In the Robben Island specimen, the pectoral rays are roughened on their inner surface with horny tubercles, as is usual with adult males of this species. The fin rays are: D. X, 14; A. 12; P. 18.

spine. Occipital crests long, gently converging behind, suddenly diverging near their posterior ends. Distance from supraorbital to occipital tubercle 1½ times the distance between the 2 supraorbital tubercles (the 2 measurements about equal in *M. polyacanthocephalus*); 2 or 3 low digitate ridges behind supraocular spine; a sharp ethmoidal ridge extends backward from level of small spines to above front of pupil; mouth very large, the maxillary extending to posterior border of eye; the pore always present behind last gill arch; spinous dorsal low, the interval between dorsals unusually long; fins moderate; pectoral barely reaching anal; ventrals not to vent. Color olive grayish, mottled with darker, paler than in related species; back with 4 dark cross bands, made up of blackish spots; lower side of head and belly plain white; membrane of upper jaw unspotted; fins, all except the ventrals, with oblique dark bars, fainter than in most species. In the adult, the dark cross bands break up into sharply defined black spots, with vermiculating blotches and lines which closely cover the back. Length 12 to 18 inches. Bering Sea, in shallow water; everywhere common on both coasts, extending into the Arctic and south to the Amur River and Unalaska. Our specimens from Unalaska, Bristol Bay, Petropaulski, Robben Reef, Port Clarence, and Grantley Harbor; one of the most characteristic fishes of Bering Sea. (*jaok*, the vernacular name in Kamchatka.)

Cottus jaok,* CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 172, 1829, Kamchatka (Coll. Pallas; specimen in Mus. Berl. called *Cottus scorpius* by PALLAS); GÜNTHER, Cat., II, 165, 1860; JORDAN & GILBERT, Synopsis, 705, 1883.

Myoxocephalus jaok, JORDAN & GILBERT,† Rept. Fur Seal Invest. 1898.

* *Cottus jaok* is thus described: "D VII, 15; A. 14. M. Lichenstein has kindly confided to us the individual which served Pallas for the description of the great *Cottus* of Kamchatka, which he believed the same species as our *scorpius*. We have compared it carefully with our individuals from Europe and we have found that it has, in fact, several characters, notably, that the spines of its preopercle are similarly disposed and of the same proportions, but its differences are many. Instead of tubercles it has behind the eye, behind the cranium, and on the temple, some light granulations. Along its back, above the lateral line, is a row of scales quite unlike those of *scorpius*. They are round, a little concave; their surface is rough, and their edge surrounded with small slender points, but above the lateral line there are some like those of *scorpius*. The first dorsal is lower, shorter, and I have been able to find but 7 rays. Pallas counted but 6. Its size is much greater than that of the Sculpins in Europe. The individual before us is 21 inches long. Pallas mentions them of 2 feet. He thus gives the coloration: Back reddish, with scattered brown spots, small, irregular, disappearing by degrees below the lateral line; belly white; 5 brown, irregular transverse bands on the pectoral; spinous dorsal spotted with brown; soft dorsal with 4 vertical brown bands; caudal with 3; 3 bands on the anal. The fish is very active in life. Pallas was told that it would live for 2 days out of water; even after being eviscerated and being in the smoke to dry, it would wave for several hours. It is called by the Kamtschadales *Jaok*; by the Koriakes, *Ilaal*; by the Kouriles, *Susiatki* and *Keischag* (?). The Russians of Kamchatka call it *Ramscha*, and the Lamutes, *Takfchi*. (Cuvier & Valenciennes.)"

† Concerning the synonymy of this species, Jordan & Gilbert observe: In a report on the ichthyological collections of the *Albatross* in Alaska (Report U. S. Commissioner of Fish and Fisheries for 1893, p. 421), Dr. Gilbert writes as follows: "*A. humilis* closely resembles the description of *A. jaok*, with which it may well be identical. We do not venture to make this identification, as *A. jaok* is said to have but 7 dorsal spines, a number we have not found in *A. humilis*." On further consideration we have decided that the 2 must be identical. The type of *jaok* was a large dried specimen, the same which had served Pallas for his account of *Cottus scorpius*. In such a dried specimen it would be very difficult to enumerate correctly the low feeble spines, of which the first 2 are very closely approximated, and the last 1 often minute and hidden in the membrane. *Humilis* is abundant along the coast of Kamchatka and agrees with the account of *jaok* in having the upper parts covered with small brown spots, the back with a series of round spinous plates, and the sides below the lateral line with posteriorly directed spines; it also agrees in reaching a very large size. In the description of *jaok* the fin formula, except the number of dorsal spines, is that most frequently found in *humilis*. Specimens are before

Cottus humilis, BEAN, Proc. U. S. Nat. Mus., IV, 1881, 149, Chamisso Island, Bering Straits (Coll. T. H. Bean); D. X, 16; A. 13; JORDAN & GILBERT, Synopsis, 705, 1883.
Cottus polyacanthocephalus, KNER, Sitzb. Akad. Wiss. Wien, LVIII, 1868, 21, taf. 4, fig. 11, Decastris Bay; not of PALLAS.
Cottus tæniopterus, BEAN & BEAN, Proc. U. S. Nat. Mus. 1896, 384; not of KNER.

2353. MYOXOCEPHALUS VERRUCOSUS (Bean).

D. X or XI, 15 to 19; A. 12 to 17 (usually D. X, 16; A. 13). Top of head strongly verrucose, the preopercular spine short; supraorbital and occipital filaments present. The adults have spinous plates along the sides, extremely high vertical fins, and very large supraorbital and occipital tubercles, from the summits of which the tubercles arise. Spinous plates above the lateral line similar to those found in *M. jaok*, the anterior ones at least being circular with a depressed center, and having the margin provided with a series of short, strong spines, sometimes interrupted for a short distance anteriorly. Under the soft dorsal and on the caudal peduncle the plates are smaller and less regular, being often reduced to very small plates bearing 2 or 3 prickles directed backward from the margin. Below the lateral line are plates similar to those above, but fewer. The cephalic tubercles are undeveloped in the young, but become very conspicuous in half-grown and adults. One above posterior margin of orbit and 1 at hinder edge of occiput are the largest and bear short filaments. These rise very abruptly without evident connection with ridges. Behind the supraorbital tubercle is a smaller one, sometimes accompanied by 1 or 2 still smaller elevations, recalling in their arrangement the digitate postocular ridges of *M. jaok* and *M. polyacanthocephalus*. A smaller tubercle is present also immediately in advance of the principal occipital one. Lower jaw included. Preopercular spine short, the upper one not exceeding, sometimes much less than, longitudinal diameter of orbit. A well-developed pore behind last gill. Dorsals very high, without appreciable interspace, the longest spine sometimes equaling length of snout and eye, 2⅓ in head, equaling the longest rays of soft dorsal. The largest specimen seen by us is nearly uniform in the coloration of the upper parts, showing but faint traces of the dark bars usually found in this group. In all other specimens these are distinctly marked, though more irregular and less sharply defined than in *M. sellaris* and *M. polyacanthocephalus*. As usual, there is a broad bar under spinous dorsal, 2 narrower ones under soft dor-

us from Petropaulski and from Stations 3646 and 3648, off Robben Island, in 18 and 20 fathoms. All of these have the supraocular and occipital crests higher and sharper than in those from the eastern portion of Bering Sea, and the preopercular spines are longer, usually reaching in young examples to or beyond opercular margin. These are, however, characters subject to much variation within this group, in which it will always be unsafe to recognize subspecies, unless based on very extensive collections. Our 21 specimens show the following fin formulæ:

	First dorsal.		Second dorsal.			Anal.			Pectoral.	
Rays	IX.	X.	15.	16.	17.	13.	14.	15.	17.	18.
Specimens	18	3	10	10	1	1	15	5	7	14

sal, and a fourth on end of caudal peduncle; ground color unusually pale. In a highly colored male the lower part of sides is blackish, provided with roundish, large, white spots, the margins of which are often made conspicuous by a series of minute black specks; fins conspicuously barred. In most specimens a broad band of the light ground color crosses occipital region and extends backward and downward, including margin of preopercle above the spines and the greater portion of opercle. Arctic Ocean and Bering Sea; Unalaska and Bristol Bay; taken by the *Albatross* at depths of 5 to 30 fathoms, and taken by Mr. Scofield at Kings Island, Port Clarence, and Grantly Harbor; originally recorded from Plover Bay. (Gilbert.) Here described from adult specimens 10 inches long. (*verrucosus*, warty.)

Cottus verrucosus, BEAN, Proc. U. S. Nat. Mus., IV, 1881, 152, Plover Bay, Siberia (Coll. T. H. Bean); JORDAN & GILBERT, Synopsis, 707, 1883.

Acanthocottus verrucosus, GILBERT, Bull. U. S. Fish Comm. (1893) 1896, 421; SCOFIELD, in Jordan & Gilbert, Rept. Fur Seal Invest., 1898.

2354. MYOXOCEPHALUS AXILLARIS (Gill).

Head $2\frac{3}{4}$; horizontal diameter of orbit 5 in head and $1\frac{1}{3}$ in snout; interorbital space 6 in head; depth $4\frac{1}{2}$; maxillary reaching vertical with posterior edge of eye, $2\frac{1}{3}$ in head; D. IX or X, 15 or 16; A. 11 or 12; pectoral 15 or 16; caudal with 9 branched rays; lateral line 40. Head wide and depressed; mouth horizontal; lower jaw included; nasal spine well developed, but completely covered by the skin. Preopercle with a straight spine at its upper angle almost covered with skin and equal in length to vertical diameter of orbit; a second spine immediately below this, completely covered by the skin and $\frac{1}{2}$ as long as upper spine; at the lower angle of the preopercle there is a tubercle; opercle with a strong horizontal spine at its upper angle, completely embedded in the membrane and not reaching edge of gill flap. At the lower angle of the opercle there is a small, downward-directed spine, also completely covered by skin; suprascapular spine well developed, but completely embedded; humeral spine blunt and covered; occipital ridges scarcely elevated, a slight pineal elevation; 4 broad, conspicuous tentacles corresponding to the positions of the supraocular and occipital tubercles; orbital rims considerably elevated, having a flat, depressed space between them; top of the head covered with small, wart-like elevations. Lateral line inconspicuous, without plates, its pores distant and small; above the lateral line a row of osseous plates, smaller and more closely placed beneath the second dorsal; a similar scattered row below the lateral line just beneath the second dorsal; longest ray of first dorsal (the fourth or fifth) $2\frac{1}{2}$ times in head, first 6 rays about equal in length; second dorsal higher and about the same shape as the first, the longest ray $1\frac{5}{6}$ in head; caudal truncated, the corners about square; pectorals large and reaching to second ray of anal; ventrals scarcely reaching vent. Color above quite dark, strongly marked with black and white; a saddle of black under the anterior $\frac{3}{4}$ of first dorsal; 2 similar but smaller saddle markings over the back below the second dorsal, 1 beneath the anterior, the other beneath the posterior end; a black blotch on the side

of the caudal peduncle, a characteristic marking in several of the members of this genus; ventral surface lighter; sides with large, ovate white spots; 4 or 5 bright cream spots at edge of the black and hidden by the pectorals; mandibles mostly black; lower lip with black mottlings; angle of mouth light; tip of maxillaries black; a black blotch at isthmus just back of membrane; first dorsal mostly black; a white blotch covering lower ½ of membranes between fourth and sixth spines; from the upper anterior angle of this blotch the white extends across the fourth spine and runs into the spot at the edge of the membrane and between the third and fourth spines; at the edge of the membrane and between each 2 spines there is a white spot, very small between first and second, but increasing in size posteriorly and between the seventh, eighth, ninth, and tenth, running together, leaving a tip of black at the end of the eighth spine; second dorsal black with a few circular white spots, each spot with its center on a fin ray and its edge reaching to middle of membrane; these spots inclined to run together; caudal mostly black, partially divided by narrow interspace of white into 2 heavy vertical bands; anal colored similar to the second dorsal; tips of rays and edge of membrane white; ventrals with several black spots; pectorals black with oval white spots on membranes, which join each other imperfectly across the rays to form 4 or 5 interrupted vertical bars of black; 2 white spots at base of rays. (Scofield.) Arctic Ocean and Bering Sea; chiefly northward. The specimens here described from Port Clarence, Chignik Bay, and Herendeen Bay. Also known from Bering Straits and St. Michaels. (*axillaris*, pertaining to the axil, or armpit.)

*Boreocottus** *axillaris*, GILL, Proc. Ac. Nat. Sci. Phila. 1859, 166, Bering Straits.
Cottus axillaris, JORDAN & GILBERT, Synopsis, 706, 1883.
Myoxocephalus axillaris, SCOFIELD, Rept. Fur Seal Investigations, 1898.

2355. MYOXOCEPHALUS STELLERI, Tilesius.

D. IX (rarely VIII), 15 (rarely 16); A. 12 (11 to 13). Resembling in shape *M. polyacanthocephalus*, the head less depressed and the snout deeper than in *M. jaok*. Characteristic features are the thickened papillose lips, the presence of a supraocular tentacle, and the peculiar coloration. Skin naked or with a few scattered small plates in adult males, the interorbital deeply concave, and the occipital and parietal ridges heavy and more or less broken or rugose. Head 2⅔ to 2¾ in length; depth 4; least depth of caudal peduncle 1½ in snout; greatest width of head equaling

* This species is the type of the nominal genus *Boreocottus*. Gill, thus described by Gill: "Body subcylindrical, rapidly declining to the caudal fin. Skin mostly naked, or with small and distinct tubercles above the lateral line. Lateral line opening on the sides through small distant pores. Head large, depressed, and subrhomboidal; nasal spines small; preopercle with 2 simple spines near the angle and 2 tubercles below; opercle with a longitudinal rib terminating in a spine and with a round membranaceous margin; subopercle with a spine directed downward. Postorbital longitudinal crests little developed. Mouth moderate. Teeth villiform, present on each jaw and on the front of the vomer. Branchiostegal membrane continuous under the throat, but attached along the middle of its length almost to its margin, and thus nearly restricting the branchial apertures to their respective sides. Branchiostegal rays 6. Dorsals separate, the first low, highest at the middle. Ventrals small, moderately approximated and behind the pectorals, each with a spine and 3 soft rays."

distance from tip of snout to base of preopercular spine; depth of head at occiput equaling ½ its length. Mouth large, the lower jaw included, but less conspicuously overlapped than in *M. jaok*, the maxillary reaching beyond the eye, 2¼ in head. Lips very thick and fleshy in adults, the inner margin of each with a dense band of fine papillæ; lower lip may also bear externally a few papillæ or short filaments; a fleshy slip or filament often present on upper posterior angle of maxilla. Nasal spines pungent, rather short; preopercle with 2 diverging spines at angle and a third remote one below directed downward and forward; the upper spine varying in length, but extending usually about halfway to tip of opercular spine; opercle with a strong rib and spine; humeral and subopercular spines strong; interorbital width 5½ to 6 in head, gently concave, its floor usually with traces of 2 low ridges; a definite supraorbital tentacle borne on the anterior end of the occipital ridge, its basal tubercle never conspicuous; slender occipital tentacle often present, especially in the young, but not infrequently absent; ridges on occiput strong, often irregular or partly interrupted, their surface roughened with lengthwise lines or with clusters of granules; occiput more deeply concave than in *M. polyacanthocephalus;* usually a cluster of short digitate ridges behind the eye; top and sides of head with small, warty protuberances. A minute pore behind last gill, to be detected with difficulty in the young. Dorsals with short interspace or none, the membrane from last spine usually joining base of first soft ray; spinous dorsal very high in adult males, the fifth spine highest, ½ as long as head, longest soft ray 2½ in head; pectorals reaching front of anal, the ventrals not to vent; vertical fins much lower in the young. Skin smooth, without plates or spines in young 7 or 8 inches long; 1 adult male of 14 inches with scattered small subcircular spinous plates, all but a few of which are below the lateral line. In the young the maxillary and mandibular membranes are whitish, very conspicuously marked with irregular jet-black spots and blotches; branchiostegal and gular membranes and the membrane behind the preopercle crossed with narrow dark streaks; entire under side of head faintly dusky, mottled and maculated with white "like a frog's belly;" iris with small black spots and blotches; these colors fainter in our adult specimen, where the under side of head is nearly uniform whitish; the maxillary membranes, however, conspicuously black spotted; body brownish, with 3 light-gray saddles, the most conspicuous crossing the back of the caudal peduncle immediately behind the dorsal fin; the second below the dorsal notch, and the third, often obscure or wanting, forming a V-shaped area on top of head, the 2 arms diverging from interorbital space toward the base of opercular spine; the dark areas often lighter centrally, and variously blotched and mottled with brown or dusky; dorsals very irregular in the marking; anal usually with 3 or 4 oblique dark bars; caudal usually with a basal translucent bar followed by varying alternations of translucent and black; ventrals showing 2 black cross bars; pectorals with no definite color pattern on their outer face, but crossed on their inner face by a few irregular black bars. Males show the usual round white spots on sides of abdomen. It is well distinguished by the speckled throat and belly, aptly compared

by Steller to the speckled color of a frog. Western shore of Bering Sea and Okhotsk Sea; not recorded from Alaska or the Aleutian Islands. This description from 11 specimens from Petropaulski and Bering Island, taken by the *Albatross*. This species is now recorded from Bering and Medni islands, Petropaulski, and the mouth of the Amur River. It is evidently abundant in western Bering Sea, but probably does not occur among the Aleutian Islands or on the Alaskan coast. There seems to be no doubt that this species is correctly identified with *Myoxocephalus stelleri*, with which it agrees in fin rays and in the peculiar and characteristic coloration. It agrees also with the description of *Cottus decastrensis*, from which the figure, however, diverges in several important details. It is highly improbable that *Cottus mertensii* and *Cottus marmoratus* can ever be satisfactorily identified, as we have only very brief accounts of them, based on colored drawings. (Named for Georg Wilhelm Steller.)

Myoxocephalus stelleri, TILESIUS, Mém. Acad. Petersb., IV, 1811, 273 with plate, not referred to in text, Petropaulski (Coll. G. W. Steller); JORDAN & GILBERT, Rept. Invest. Fur Seal Islands, 1898.

Cottus decastrensis, KNER, Denk. Kais. Akad. Wissen., XXIV, 1865, 2, taf. 2, figs. 1, 1*a*, Decastris Bay, near mouth of the Amur.

? *Cottus marmoratus*,* CUVIER & VALENCIENNES, Hist. Nat. Poiss., VIII, 497, 1831, Petropaulski; on a drawing by MERTENS.

Cottus platycephalus, BEAN & BEAN, Proc. U. S. Nat. Mus. 1896, 240, 384; not of PALLAS.

Cottus niger, BEAN & BEAN, Proc. U. S. Nat. Mus. 1896, 240, 384 (in part: Nos. 33899, 33872, 33833, 33850, 33908, 33844, and 33879); not of BEAN.

2356. MYOXOCEPHALUS MEDNIUS, B. A. Bean, new species.

Head $3\frac{1}{2}$; depth $4\frac{1}{2}$; eye 4 in head. D. VIII, 17; A. 12; V. I, 3; P. 14; C. 14. Profile of head and body gradually ascending from tip of snout to sixth dorsal spine, thence tapering to caudal peduncle; ventral line almost straight, slightly tapering to caudal peduncle; longest dorsal spine almost as long as longest ray, 3 in head including flap. Mouth moderate, maxillary reaching about to vertical through middle of eye. Pectorals large, reaching slightly beyond anal origin, the middle rays being $\frac{3}{4}$ as long as the head; ventrals moderately well developed, reaching anal; anal origin under third ray of dorsal, ending under fourteenth ray of that fin. Gill membranes united, forming a fold across the isthmus. Preocular spines moderate; opercular spines but moderately developed; 2 flattish tubular

* *Cottus marmoratus* is thus described:
"D. 8-14; A. 12; C. 14, etc. Une seconde espèce, observée dans ces parages par les naturalistes de la même expédition, ne paraît pas non plus se rapporter à aucune de celles que nous avons décrites. Ce petit chaboisseau a deux fortes épines au-devant de l'œil. Deux autres peu alongées au bord du préopercule; une petite dirigée vers le bas au bord horizontal du même os, et une assez forte à l'angle de l'opercule. Les rayons épineux de la première dorsale sont assez robustes; le cinquième est le plus long. Ce poisson a la tête et le dos bruns, de grandes marbrures brunes et blanches sur les côtés, et rougeâtres sur le ventre. La première dorsale est jaune, avec une grande tache brune, qui part de la pointe du quatrième rayon et descend obliquement jusqu'au pied du septième. La seconde dorsale est rouge. La caudale jaune, marbrée de roux. La pectorale brune à la base et jaune ensuite. Le jaune est traversé par trois raies circulaires brunes près de l'extrémité des rayons. L'anale est blanchâtre et tachetée de nombreux points rouges. Nous ne connaissons aussi cette espèce que par un dessin long de quatre pouces, communiqué par M. de Mertens." (Valenciennes.)

pores, 1 on each side of front of eye; numerous pores on head; 2 rows of pores, 1 above and 1 below, the raised ridge running laterally on dorsal half of body. General color dark reddish brown, mottled, barred, and spotted with white; under parts whitish; a wide whitish bar from opercles across nape; posterior part of interorbital space whitish; the dark color on front of snout and under lower jaw relieved by bars and mottlings of whitish; pectorals and ventrals barred; rays of caudal finely mottled; several (5) small white spots on body immediately behind pectoral origin, and several larger white blotches on lower posterior half of body. Bering Sea. A single example, 2 inches long. It is allied to *M. stelleri*, from which it differs greatly in form and coloration. (Type, No. 33863, U. S. Nat. Mus. Collected at Medni (Copper) Island, Bering Sea, spring of 1883, by Dr. Leonhard Stejneger.) (B. A. Bean.) (*Medni*, the Russian name, meaning Copper, of the island where the species was found.)

Myoxocephalus mednius, B. A. BEAN MS., in JORDAN & GILBERT, Rept. Fur Seal Investigations, 1898, Medni Island. (Type, No. 33863. Coll. L. Stejneger.)

2357. MYOXOCEPHALUS NIVOSUS* (Herzenstein).

Head 2¼; depth 4½. D. IX, 15; A. 13; P. 17; lateral line 33 to 35. Head comparatively deep and compressed, with large mouth, narrow, deeply concave interorbital space, and depressed concave occiput, which is bounded by strong lateral crests; at the anterior end of these crests they are each accompanied on the outer side by a short ridge, and on the inner side by a still shorter ridge or a small tubercle. The occipital crests converge strongly toward the nape; temporal ridges also strong; a short filament above posterior edge of orbit and 1 at posterior end of occipital crest, each surmounting a very low tubercle. Upper preopercular spine straight, directed toward opercular flap, scarcely reaching middle of opercle, its length ⅔ diameter of orbit; the second spine is ⅔ length of upper, directed downward and backward; the third points downward and forward, the long interval between it and the second being smooth, without spine or tubercle. The contiguous angles of sub- and inter-opercles are

* Allied to *M. nivosus* is the following species from the Okhotsk Sea:
Myoxocephalus brandti (STEINDACHNER).
Head 2⅔; depth 5. D. IX, 13; A. 11; P. 17. Head narrowed forward, the lower jaw included. Eye 6 in head, a little larger than snout; interorbital space strongly concave, 1½ in eye. A rather long tentacle above eye posteriorly; behind this a ridge, low and rather sharp, converging with its fellow, and inclosing a depressed quadrangular area; no spine at its posterior end; nasal spines prominent; preopercle with 3 spines, the upper shorter than eye, the second not ½ its length, the third short, blunt, and turned downward; opercular spine blunt, concealed; suprascapula with a single long spine. Top of head covered with small rounded warts. Lateral line with thin bony concealed plates, its pores sending numerous simple branches above and below; below the lateral line 2 rows of small cross plates, tube-like, apparently connected with the system of tubes of the lateral line; no rounded bony plates or scales. Longest dorsal spine 3 in head; pectoral moderate, a little more than ½ head; ventral 2⅓ in head. Head bluish violet above; sides of body reddish yellow; violet marblings on the lower lip and on the sides of the lower jaw, leaving the ground color to form ocellated spots; fins violet with yellow spots, the ventrals uniform yellowish. Known from 1 specimen 13 inches long from the mouth of the Amur River. (Steindachner.) (Probably named for the natural history collector, Brandt.)

Oottus brandti, STEINDACHNER, Ichth. Notizen, v, 6, Taf. 3, figs. 1, 2, 1867, mouth of the Amur River. (Type, in Vienna Mus.)

provided with prominences which are not spine-like; opercle and supra-scapula each with a strong ridge ending in a spine; no scapular spine. Nasal spines small, not projecting; top of head, nape, and suborbital ring with small warts, many of which contain openings of the mucous canals; both pairs of nostrils with short, wide tubes. Jaws and vomer with wide bands of cardiform teeth of equal size. A round pore, nearly as large as nostril, immediately behind last gill; gill membranes with a short free border mesially. Least interorbital width $\frac{3}{5}$ diameter of orbit, which is $\frac{1}{5}$ length of head to tip of opercular spine; greatest width of head $1\frac{1}{3}$ in its length. Mouth large, the lower jaw included, the maxillary reaching the vertical immediately behind the orbit, its length $2\frac{2}{3}$ in head. Third, fourth, and fifth dorsal spines nearly equal, the fifth longest, equal to length of snout and $\frac{1}{2}$ eye; a very short interspace between the 2 dorsals; second dorsal very high, the longest rays equaling length of snout and eye; caudal gently rounded when spread, its length $\frac{1}{3}$ that of head to end of opercular spine; ventrals reaching halfway to front of anal, equaling height of second dorsal; pectorals scarcely to vent, the length of the longest rays equaling distance from eye to tip of opercular spine; vent midway between base of caudal and base of lower pectoral ray; skin everywhere smooth. Coloration very dark on back and sides, white below, with traces of blackish cross bars below the dorsal fins; along lower part of sides a number of large roundish white spots, which are present also in a band along base of anal, becoming smaller posteriorly and toward middle of sides; sides of head and body, and especially the dorsal, caudal, and pectoral fins with scattered small spots of pearly white; spinous dorsal with large, roundish, transparent spots, the anal margined with white and marked with scattered white spots of various sizes and shapes; the thickened pectoral rays largely white; ventrals with 3 dusky cross bars; lower lip and mandible with white areas surrounded by dusky reticulations. Okhotsk Sea, the type from St. Olga Bay. Here described from a large specimen 39 cm. long from Iturup Island, one of the Kurils. (Coll. *Albatross.*) Our specimen agrees well with the description of the much smaller type (185 mm), apparently differing in the shorter pectorals and more deeply concave interorbital space. The white spots also show no tendency to run together to form streaks either along back or on the bases of the fins. (*nivosus*, snowy.)

Cottus nivosus, Herzenstein, Mélanges Biologiques du Bull. Acad. Imp. Sci. Petersb., xii, 113, 1890, St. Olga Bay.
Myoxocephalus nivosus, Jordan & Gilbert, Rept. Fur. Seal Invest., 1898.

2358. MYOXOCEPHALUS NIGER (Bean).

(Black Sculpin.)

Head $2\frac{1}{2}$. D. IX or X, 16; A. 11 or 12; V. I, 3; P. 16; B. 6. Nasal spines blunt, covered with skin; no spines above orbits or on occiput; no sharp spines on head except the 2 upper preopercular spines, which are almost concealed; the uppermost as long as eye, 6 in head, and equal to snout or interorbital space; top of head with fine dermal granulations, the

vertex with numerous slender tentacles; a soft tentacle of moderate length above eye; jaws equal, the lower slightly included; maxillary longer than fourth dorsal spine, 2 in head, reaching hinder margin of orbit. Caudal and ventrals each ½ length of head; pectoral not reaching vent; skin smooth. A small slit behind last gill. Color typically, almost black; sides mottled with lighter brown, sometimes with round white blotches on belly and sides; pectoral plain or with white blotches. Specimens from granitic rocks have sanded coloration corresponding to the hue of the surroundings. Pribilof Islands; very abundant in the crevices of black lava rocks of St. Paul and St. George. Also obtained in some abundance about the Komandorski Islands, Medni and Bering.

This strongly marked species can be readily distinguished by its peculiar coloration, the greatly thickened naked skin which partially conceals the short nasal and opecular spines, and the numerous tentacles surmounting the warty tubercles on crown and occiput. The fins may be uniformly black with a narrow white tip to the soft rays, or may be more or less variegated with white. On the soft dorsal, these marks are in the form of white spots which may become confluent to form 1 or 2 streaks. In some specimens, the caudal membranes are white in their middle portion the rays remaining black. The under side of head and the maxillary membranes are sometimes marked with large blackish spots with ill-defined edges.

In addition to the minute pores which lie at intervals along the course of the lateral line, the latter gives off pairs of lateral branches, each of which opens in 3 or more small pores. The sides of the head are also thickly studded with pores. Owing to the thickened integument, the pectorals and ventrals are more largely adnate to the body than in other species. The fin rays are as follows, in 10 specimens:

	First dorsal.		Second dorsal.			Anal.		Pectoral.	
Rays	IX.	X.	15.	16.	17.	11.	12.	16.	17.
Specimens ..	6	4	1	8	1	3	7	2	8

(*niger*, black.)

? *Cottus mertensii*,* CUVIER & VALENCIENNES, Hist. Nat. Poiss., VIII, 496, 1831, Petropaulski; on a drawing by MERTENS.

Cottus niger, BEAN, Proc. U. S. Nat. Mus., IV, 1881, 151, St. Paul Island, Pribilof Island (Coll. H. W. Elliott); JORDAN & GILBERT, Synopsis, 707, 1883.

Myoxocephalus niger, JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

* *Cottus mertensii*, is thus described: Il se rapproche du petit chaboisseau de Groënland par le nombre de rayons de l'anale; mais par ceux des dorsales il est plus voisin du *Cottus octo-decim-spinosus*, et il a les épines de la première dorsale tout aussi fortes. Voici les nombres comptés par le naturaliste russe. D. 8, 15; A. 12; C. 10; P. 18, etc. La tête paraît lisse; sans épines près des narines. La pointe du préopercule n'atteint pas l'angle de l'opercule. Les couleurs indiquées sur le dessin qui nous a été communiqué par M. de Mertens, étaient de grandes marbrures de bistre foncé sur un fond jaunâtre. Ce dessin représente un individu long de huit pouces.

732. MEGALOCOTTUS, Gill.

Megalocottus, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 166 (*platycephalus*).

This genus includes large sculpins, with broad flat heads, differing from *Myoxocephalus* chiefly in the sharply projecting lower jaw; the preopercle has 3 spines, the upper short and straight, the lowermost turned downward; dentation as in *Myoxocephalus*; suprascapula with 2 spinous projections; top of head warty; sides with rounded tubercles in the adult; lateral line of separated pores; a large pore behind last gill. North Pacific. (μέγας, great; κόττος, cottus.)

a. Interorbital space moderate, about equal to eye in adult; ventrals reaching anal papilla. PLATYCEPHALUS, 2359.

aa. Interorbital space very broad, 1⅔ diameter of eye in adult; ventrals not reaching vent. LATICEPS, 2360.

2359. MEGALOCOTTUS PLATYCEPHALUS (Pallas).

Head 3. D. VIII, 13; A, 12; V. I, 3; P. 16 or 17; lateral line 40. Lower jaw somewhat longer than upper; eye 5 in head, as long as snout; nasal spines distinct; orbital rim moderately prominent; a low ridge behind it toward the occiput, without spines or points. Top of head between these ridges forming a quadrangular depression, as broad as the eye in front, but narrower behind. Top of head covered with naked, warty skin. Preopercle with 4 spines, the upper as long as eye, directed upward and backward, the second much shorter, the others directed downward. Opercle with a long forked ridge, ending in 2 short diverging spines; a spine directed downward on subopercle; two sharp spines on scapular region, the lower the longer and touching the first tube of the lateral line. Lateral line complete, above, it numerous rounded bony plates armed with spinules as in *M. laticeps;* these in 2 rows anteriorly and 1 posteriorly; a few similar plates below lateral line. Fins all high, the soft dorsal especially so. Color dark, with many dark shades and spots on throat and belly, especially on gill membranes; sides and belly (in males) with irregular, round white spots; first dorsal with round pale spots, the membranes otherwise largely dusky; soft dorsal with 3 or 4 dark oblique cross shades, between which are rows of pale spots; anal with 4 dark oblique cross shades; caudal with 3 narrower dark stripes; pectoral with 4; ventrals dusky with pale spots. Okhotsk Sea and western part of Bering Sea. This description (after Kner) from a single specimen, 7½ inches long, from Decastris Bay, near the mouth of the Amur. The species is close to *M. laticeps*, which has been recorded from Alaska as *Cottus tæniopterus*, but the 2 species are different, and the type of *M. platycephalus* has the narrower interorbital space of *M. tæniopterus*. So far as known *M. laticeps* is found on the eastern shores of Bering Sea, and *M. platycephalus* on the western. Dr. Hilgendorf has, at our request, reexamined Pallas's type of *platycephalus*, a specimen a foot long. He notes: Lower jaw projecting; fin rays not easily counted, the skin being dried and varnished. D. VIII–14; A. 11. A sketch of the head shows the armature of *M. laticeps*, but the eyes closer together, 1 diameter apart. There is a blunt tubercle behind

each eye and 1 at the occiput; the spine on the suprascapula is forked, and the lower fork is twice the length of the other. The upper preopercular spine is about ½ longer than the second. (πλατυς, broad; κεφαλή, head.)

Cottus platycephalus,* PALLAS, Zoogr. Rosso-Asiat., III, 135, 1811, Kamchatka; after STELLER; CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 177, 1829.

Cottus tæniopterus, KNER, Sitzber Ak. Wiss. Wien, LVIII, 1868, 18, Taf. 14, fig. 10, Decastris Bay, near mouth of Amur River. (No. 5574, Mus. Wien.)

Megalocottus platycephalus, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 166.

2360. MEGALOCOTTUS LATICEPS (Gilbert).

This species differs from *M. platycephalus* (*tæniopterus*, as shown in Kner's figure) in the following respects: The interorbital width is greater, 1⅘ times the diameter of the eye in *laticeps*, equaling the eye in *platycephalus*. Similarly the distance between the anterior ends of occipital ridges is 1⅔ times diameter of eye in *laticeps*, equal to eye in *platycephalus*. Conspicuous supraorbital and occipital tubercles in *laticeps*, each surmounted by a fleshy filament; in *platycephalus* "crown and occiput are without prominences or spines, covered only with naked warty skin." In *laticeps* the preopercle has 2 diverging spines, a single downwardly directed spine below them. In *platycephalus* are 2 downwardly directed spines below the angle. The ventrals are much shorter, not reaching vent when depressed, while in *platycephalus* they reach the anal papillæ. The dorsals are also much lower in *laticeps*. The principal features of this species are the very broad flat head, short wide snout, projecting lower jaw, the 2 pairs of cephalic tubercles provided with tubercles, the verrucose head, the very short preopercular spines, the large pore behind the last gill, the presence of circular spinous plates above the lateral line and prickles below it. Head and anterior part of body broad, depressed, the depth of head, at occiput, 1⅘ in its greatest width, its length 2¾ in body. Body tapering to a slender caudal peduncle, whose least height is equal to diameter of eye. Depth of body 4 to 4⅓ in length. Interorbital space very wide, shallowly concave, its width 1½ times diameter of orbit, and 4 to 4½ in head. The low supraocular ridge ends in a blunt tubercle above hinder margin of eye which grows higher with age, depressed, bounded by 2 low ridges which converge very strongly toward the nape, where they curve out again in low rounded tubercles; the inclosed depressed area is twice as wide anteriorly as it is at the narrowest posterior part. A strong temporal ridge, less dis-

*The following is the substance of Pallas's account of *Cottus platycephalus*. D. VII–12; A. 11; C. 10; P. 15; V. I, 3. Head large, much depressed, as though crushed, as broad as body. Tail slender; mouth large, lower jaw projecting; teeth on vomer, none on palatines. Belly much inflated. Nostrils tubular. Eyes moderate, near together; vertex flat behind the eyes, with a ridge on each side, in front of which, near the orbit, is a bony tubercle; behind at the nape, oblong tubercles, each ending in a short spine; preopercle with 2 very strong, diverging spines above; opercle with a concealed spine; lateral line with concealed, elongate plates; between it and the dorsal some round very rough warts, smaller ones below; pectorals large; dorsal spines feeble. Olivaceous, much variegated with greenish lines on sides of body below; spots large below the tail; dorsals translucent, varied with brown; caudal yellowish with faint bands; pectorals and anal banded. Length 12 inches. Kamchatka. (Pallas.) The type in the museum at Berlin has no teeth on the palatines. The generic name *Megalocottus* was originally based on the supposed presence of palatine teeth, a fallacious character. The character of the projecting chin has, however, generic value. Pallas says: "In arca lunata palatio medii et utrinque lineari," which, if we translate it correctly, is not true.

tinct in young specimens. Nasal spines very small. Mouth wide, transverse, oblique, the maxillary reaching vertical from hinder edge of pupil, 2⅛ in head; mandible with its triangular tip protruding well beyond upper jaw in adults, the jaws nearly equal in young; length of snout equal to interorbital width. Bands of short cardiform teeth on the jaws, and a broad patch on vomer; none on palatines. A conspicuous pore behind last gill. Gill membranes almost wholly joined to the isthmus, the free border scarcely noticeable, its width less than ⅓ diameter of the small pupil. Eye small, shorter than snout, 5½ to 6½ in head. Preopercle with 2 short strong spines diverging backward, and a strong concealed point below directed downward and forward; upper preopercular spine about equal to eye, but little longer than the lower and reaching ⅓ the distance from its base to the tip of the opercular flap; opercle with a well-marked longitudinal rib ending in a sharp point; scapular and subopercular spines present. Entire top and sides of head, nape, and anterior dorsal region covered with small dermal warts; supraorbital and occipital tubercles with short filaments. Space above the lateral line with an irregular series or double series of large round spinous tubercles; a few scattered plates on sides below the lateral line; axil smooth; plates of lateral line concealed in skin. Spinous dorsal low, its longest spine 1½ in second dorsal and 3 in head. In some specimens a single line of small sharp tubercles, resembling spines, extends each side of the rays of the second dorsal. The 2 dorsals are separated by a narrow space; pectorals large, reaching front of anal, the lower rays much thickened; caudal rounded; ventrals reaching ⅔ distance to vent. Dorsal IX, 14; anal 13; pectoral 18; ventral I, 3; lateral line 36 to 40. Color dark olive brown above, with faint traces of blackish bars; sides spotted or marbeled with whitish; belly and lower parts generally white; a blackish blotch on cheeks, 1 on opercle, and a third on front of mandible; pectoral rays dusky, the membrane whitish, the fins crossed by 3 or 4 wavy black bars, which sometimes join, inclosing oblong or roundish white areas; spinous dorsal not banded, the dusky and translucent areas variously arranged; soft dorsal with 5 oblique broad dusky bars; anal with 4 bars sometimes uniting to inclose white spots; caudal similar to pectoral and anal; brilliantly colored males are largely black on sides and below, with many large, rounded, partially confluent pearly white spots. Length 6 to 11 inches. (Gilbert.) East shore of Bering Sea; known from 13 specimens from the Nushagak River, near its mouth; 1 from Herendeen Bay on the northern side of the Alaskan Peninsula and from Port Clarence (Scofield*). (*latus*, broad; *-ceps*, head.)

Cottus tæniopterus, Bean, Proc. U. S. Nat. Mus. 1881, 248; Bean, in Turner's Nat. Hist. Alaska, 94, pl. 6, 1886; not of Kner.

Acanthocottus laticeps, Gilbert, Rept. U. S. Fish Comm. 1893 (1896), 422, pls. 26 and 27, Nushagak River and Herendeen Bay, Alaska. (Coll. *Albatross*.)

Megalocottus laticeps, Scofield, in Jordan & Gilbert, Rept. Fur Seal Invest. 1898.

* Mr. Scofield observes: "We took 4 specimens of this species at Port Clarence. They differ from the types slightly in coloration. Our specimens are dark, almost black; below the anterior portion of the second dorsal is a darker saddle-like marking on the body; the caudal is black at the base and has an undulating dark band on its posterior third; the pectorals vary, being either banded or mottled. (One specimen has distinct bands on 1 pectoral while the others are simply mottled with black.) There is no slit behind last gill arch."

733. ZESTICELUS, Jordan & Everman.

Zesticelus, JORDAN & EVERMANN, Check-List Fishes, 443, 1896 (*profundorum*).

Deep-sea sculpins, closely allied to *Porocottus*, but with the skeleton little developed, the head soft and spongy, filled with mucous channels, the skin perfectly smooth, the lateral line reduced to a series of separate open pores, the vertical fins few-rayed and weak. Preopercular spine slender, curved upward. Deep seas; 2 species known; probably degraded from *Porocottus*, the soft skeleton and feeble structure being results of deep-sea life. (ζεστός, soft-boiled, *Icelus*, a son of the god of sleep.)

a. Dorsal rays VI or VII, 10 to 13; anal rays 8 to 11; pectoral 20; upper preopercular spine usually not reaching opercular margin. PROFUNDORUM, 2361.

2361. ZESTICELUS PROFUNDORUM (Gilbert).

Head 2⅔; depth 5. D. VI or VII, 10 to 13; A. 8 to 11; pectoral 20; ventral I, 3; lateral line with 17 pores. From above the head appears smooth and evenly rounded, without projecting spines or ridges. The occipital depression is very shallow, the occipital ridges depressed, scarcely noticeable, ending in depressed spines, which are made out with difficulty. Nasal spines undeveloped, the nasal bone small, posteriorly pointed, but not furnished with a projecting spine. Upper preopercular spine strongly compressed, curved upward, not reaching opercular margin, its length equaling diameter of eye; no spine at its base in front; below it 2 short, strong spines directed downward and backward, and 1 more slender downward and forward; opercle with a longitudinal rib ending in a short spinous point; a short spine on angle of opercle, and 1 below it on interopercle. Mucous canals everywhere greatly enlarged, giving a spongy texture to the entire head; series of very conspicuous pores on the preopercle, the mandible, and below suborbital chain. Mouth broad, oblique, the maxillary reaching middle of pupil, 2⅓ in head; mandible slightly protruding. Minute teeth in upper jaw, anteriorly in 2 rather distinct rows, laterally in narrow bands; teeth on vomer, none on palatines. Eye longer than snout, 3¾ to 4 in head. Interorbital width ½ diameter of orbit. Gill membranes widely joined, with a wide free posterior edge; no slit nor pore behind last gill. Body smooth, without plates, granulations, or filaments; no plates developed in connection with the lateral line; pores of lateral line in a double series, the 2 closely approximated, those of the lower series much the larger; no evident tubercles by which these communicate with the main canal. Longest rays of second dorsal ½ length of head, twice the longest dorsal spine; pectoral reaching front of anal or slightly beyond; ventrals short, not nearly reaching vent. Upper parts very light brownish, the belly and sides below lateral line dark brown; fins blackish; mouth and gill cavity dark. A deep-sea form, characterized by the obsolescence of the occipital and nasal spines, the absence of the accessory spine in advance of upper preopercular spine, and in the more numerous rays of dorsal and anal fins. Length about 2

inches. Bering Sea in deep water, 4 specimens known; from off Unalaska in 399 fathoms, and north of Bogoslof Island at *Albatross* Station 3634, in 664 fathoms, where a single specimen 58 mm. was obtained. This specimen closely resembles the type of *Zesticelus profundorum*, but the spines are somewhat stronger than in the latter and the fins shorter. In the Bogoslof specimen the upper preopercular spine extends to margin of opercle, and the occipital spines are more evident, terminating in short strong ridges. D. VII, 10; A. 8; P. 20. The lower parts, up to and including the lateral line, are blackish, the line separating them from the lighter under parts very sharply defined. Fins all black. Otherwise essentially as in the type of *Z. profundorum*, with which we now consider it identical. The variation in the fin rays in this species is considerable, if the figures given by Dr. Gilbert are correct. One of the cotypes, however, has P. 20, not 17 or 18, as stated by Gilbert. The figure also shows 20 rays in the pectoral. It also shows 11 rays in the anal, while our specimen has but 8. (*profundorum*, of the depths.)

Acanthocottus profundorum, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 423, pl. 26, Bering Sea north of Unalaska Island at Albatross Station 3329, in 399 fathoms.
Zesticelus profundorum, JORDAN & GILBERT, Rept. Fur Seal Investigations, 1898.

734. DASYCOTTUS, Bean.

Dasycottus, BEAN, Proc. U. S. Nat. Mus. 1890, 42 (*setiger*).

Head large, with large, bony tubercles and numerous filaments. Mouth wide, jaws equal, or lower jaw slightly projecting. Teeth minute, in villiform bands in jaws; vomerine teeth in 2 patches, separated by an interspace; palate toothless. Preopercle with 2 spines at its angle and 2 rudimentary ones on each edge of its lower limb. Gills $3\frac{1}{2}$, no slit behind the last; gill rakers short, tubercular, in moderate number; gill membranes free from isthmus; pseudobranchiæ present. Skin smooth, except on nape and along dorsal base, where it bears a few small, bony tubercles. Spinous dorsal well developed and separated from the soft dorsal by a deep notch; pectorals moderately long, their rays procurrent below; ventrals with a spine and 3 rays; caudal rounded. North Pacific. Very close to *Cottunculus*, apparently differing in having cirri on the head and the preopercular spines not hidden by skin. (δασύς, woolly; *Cottus*.)

2362. DASYCOTTUS SETIGER, Bean.

Head 2; depth 3. D. IX, 15; A. 13; V. I, 3; lateral line (pores) 11 or 12. Eye nearly equal to snout, $\frac{2}{3}$ length of head. Maxillary expanded behind, its length equaling that of snout and eye combined, extending to a little behind middle of eye. Pectoral extending to above anal origin; ventral $\frac{1}{2}$ as long as postorbital part of head; longest dorsal spine about equaling eye in length; spinous dorsal with 2 dusky bands extending down on the

sides; soft dorsal with 2 bands; a similar one at caudal base and 3 narrow bands on second half of caudal; pectoral indistinctly banded; head speckled with brown dots. (Bean.) North Pacific; the types from *Albatross* Station 2855, north latitude 57°, west longitude 153° 18′, at a depth of 69 fathoms, off Sitkalidak Island; our specimens from Puget Sound and off Karluk. Concerning this species Dr. Gilbert observes:

"Taken at *Albatross* Stations 3216, 3257, 3310, 3311, and 3334, located north and south of the Alaskan Peninsula, and north of Unalaska Island, in 50 to 85 fathoms. Tubercles on head definitely placed; 1 in front of eye; 4 above orbit, the posterior 2 the largest; a pair on middle of suborbital stay, with a smaller one above them; 1 on temporal region, and 1 on shoulder; by far the largest pair on occiput, where they are high, compressed spines, directed vertically upward, as long as diameter of pupil; nasal spines obsolete. Cirri are generally distributed over upper part of head and body, the longer ones being specially numerous on maxillary, under surface of mandible, and on the opercle and preopercle; of the larger ones, 2 often proceed from 1 base; a series of short filaments along upper edge of pupil. Mucous pores large, those of the mandibular and buccal series slit-like. In adults the dorsal bands break up into series of spots and become inconspicuous." (*seta*, bristle; *gero*, I bear.)

Dasycottus setiger, Bean, Proc. U. S. Nat. Mus. 1890, 42, off Sitkalidak Island, Alaska (Type, No. 45370. Coll. *Albatross*); Jordan & Starks, Proc. Cal. Ac. Sci. 1895, 810, pl. 83; Gilbert, Rept. U. S. Fish Comm. 1893 (1896), 411.

735. COTTUNCULUS, Collett.

Cottunculus, Collett, Norges Fiske, 20, 1875 (*microps*).

Body tadpole-shaped, the head extremely large, the body tapering rapidly from the shoulders to the slender tail; mouth rather large, terminal, oblique, the jaws about equal; villiform teeth in the jaws; a double patch on vomer; no teeth on the palatines; no spines on the head, the tubercular surface of the skull covered by skin; skull thin, its bones not firm. Gills 3½, no slit behind the last arch; gill membranes broadly joined to the isthmus, their union extending to above the lower edge of the base of the pectorals. Pseudobranchiæ very small; no cirri, scales, or prickles; the skin thin and movable, smooth, or roughened with small warts. Spinous dorsal little developed, the 2 fins usually continuous; spines very slender, flexible, embedded in the skin; pectorals short, procurrent below; ventrals very short, well separated, their rays I, 3; caudal rounded. Deeper parts of the Atlantic. (A diminutive of *Cottus*.)

a. Anal rays 10. MICROPS, 2363.
aa. Anal rays 13. THOMSONII, 2364.

2363. COTTUNCULUS MICROPS, Collett.

Head 2⅝; depth 3½. D. VI, 19; A. 10. Head very large, its length, breadth, and depth nearly equal; the greatest depth at the nape; 4 bony tubercles on top of head and some at the sides, all covered by the skin;

lower jaw included; maxillary extending to below the middle of the eye, which is equal to the snout, and about 4½ in head; suborbital stay and the roughish edge of the preopercle both covered by smooth skin; chin and preorbital with pores; skin everywhere thin, somewhat movable, its surface roughened by small blunt warts; dorsal fin continuous, the feeble spines lower than the soft rays; pectorals barely reaching anal; caudal long. Color pale, with 3 broad, dusky cross bands on body and fins, 1 on head, 1 through spinous dorsal and pectoral, and 1 through second dorsal and anal, besides a small band at base of caudal. Length 8 inches. Deep water off coasts of Norway and Rhode Island; taken by the United States Fish Commission in the expeditions of the *Blake*, *Fish Hawk*, and *Albatross*. Here described from specimens taken off Rhode Island. (*μικρός*, small; *ὤψ*, eye.)

Cottunculus microps, COLLETT, Norges Fiske, 20, pl. 1, figs. 1–3, 1875, Hasvig, near Hammerfest, Norway, in 200 fathoms (Coll. Prof. G. P. Sars); COLLETT, Meddelelser om Norges-Fiske, 1875-78, 1879; COLLETT, Forh. Vid. Selsk., Christiania, 1880, 11; COLLETT, Norsk. Nordh. Exped., 18, pl. 1, figs. 5, 6; Nyt. Mag. f. Naturvid., XVIII, 53, 1884; STRÖM, Norsk. Vid. Selsk. Skrift. 1880, 75; GOODE, Proc. U. S. Nat. Mus. 1880, 470; GOODE & BEAN, Bull. Mus. Comp. Zool., X, 212, 1883; LILLJEBORG, Sverig. och. Norg. Fiske, 113; JORDAN & GILBERT, Synopsis, 688, 1883; GÜNTHER, Challenger Report, XXII, 60, pl. 9, fig. A, 1887; GOODE & BEAN, Oceanic Ichth., 269, figs. 257 and 261, A, B, 1896.

2364. COTTUNCULUS THOMSONII (Günther).

Head 2½; depth 3½ to 4; eye 4½ in head; snout 4½; D. VI, 17; A. 13; P. 22; V. I, 3; C. 10. Head and body smooth, scaleless, covered with a tough, lax skin; length of head nearly ⅓ extreme length of body, including caudal, its greatest height ¼ body without caudal; greatest width of head twice length of maxillary; distance of vent from insertion of anal equals length of maxillary; eye close to dorsal profile; intermaxillary long and slender, slightly more than 3 times in distance from tip of snout to insertion of first dorsal, 3⅓ in head; maxillary very slender, except in posterior third, where it is considerably expanded; mandible very stout, posteriorly widened, its length contained nearly 2⅓ times in head. Teeth in broad, villiform bands on intermaxillary and mandible; 2 short, separate, similar bands on vomer; none on palatines. Head armed with blunt spines as in *C. microps*. Distance of dorsal from tip of snout nearly equal to ⅓ total length, caudal included; anal fin midway between tip of snout and end of caudal; length of upper pectoral rays equal to that of postorbital portion of head, its rays diminishing rapidly in size, the lowest being exceedingly short; distance of ventral from tip of snout ⅓ of total length without caudal; length of free portion of ventral equals eye. Color light brown, the fins somewhat darker. Northern Atlantic, in deep water. Length 6 inches. Specimens obtained by the *Blake* from Station 306, in 41° 32′ 50″ north latitude, 65° 55′ west longitude, etc.; also by the *Albatross* from Station 2181, in 39° 29′ north latitude, 71° 46′ west longitude, etc., and by the *Fish Hawk* from Station 1029, in 39° 57′ 6″ north latitude, 69° 16′ west longitude, etc. (Named for Sir C. Wyville Thompson, Regius professor of Natural History in the University of Edinburgh, first Director of the

Civilian Staff on board H. M. S. *Challenger* during her magnificent scientific explorations.) (Goode & Bean.) (Eu.)

Cottus thomsonii, GÜNTHER, Proc.Royal.Soc. Edinburgh, XI, 1882, 679, Challenger Station 4.
Cottunculus torvus, GOODE, Proc. U. S. Nat. Mus., III, 1880, 479, Gulf Stream, off Rhode Island; GOODE & BEAN, Bull. Mus. Comp. Zool., X, 212, 213, 1883; JORDAN & GILBERT, Synopsis, 688, 1883.
Cottunculus thomsonii, GÜNTHER, Challenger Report, XXII, 61, pl. 9, fig. B, 1887.

736. MALACOCOTTUS, Bean.

Malacocottus, BEAN, Proc. U. S. Nat. Mus. 1890, 42 (*zonurus*).

Shape similar to that of *Cottunculus*. Body tapering rapidly to the slender tail; head large; mouth terminal; jaws subequal; minute villiform teeth in broad bands on maxillary and mandible; vomer and palate toothless; preoperculum armed with short, stout, simple spines. Bones of the skull thin. Gills 3½, no slit behind the last. Gill openings wide, the membranes broadly attached to the isthmus. Gill rakers tubercular, in moderate number. Spinous dorsal low, separated by a deep notch from the soft dorsal. Pectorals procurrent in front; ventrals small; caudal rounded; vent distant from the anal origin. Head and body naked. Lateral line consisting of a series of large pores. North Pacific. (μαλακός, soft; *Cottus*.)

2365. MALACOCOTTUS ZONURUS, Bean.

Eye 3½; snout 3½; depth equaling length of head without snout. D. IX, 14; A. 11; V. I, 3; lateral line (pores) about 14. Interorbital spaces less than ½ length of eye. Maxillary extending to below middle of eye. Pectoral reaching to a vertical through the anal origin or a little beyond; ventral scarcely as long as eye; spinous dorsal low, its base ½ as long as head, its longest spine ⅔ as long as eye; origin of dorsal immediately above upper angle of gill opening; least height of tail scarcely ⅔ length of eye; four spines on preopercle, the largest less than ½ as long as eye and with a supplementary spine at its base. A dark-brown saddle-shaped band over the end of spinous dorsal and 2 on soft dorsal; a brown band at base of tail and 3 on caudal; pectoral with several indistinct dark bands intermingled with pale areas; tips of pectorals in their lower half milky white. Length 4¾ inches. Coasts of Alaska, about the peninsula. Concerning this species Dr. Gilbert observes:

Several specimens were taken north of Unalaska Island and south of Unimak Island, in 138 and 351 fathoms. Nasal spines obsolete; supraorbital rim low, slightly elevated in front but not behind, the interorbital space wide, shallowly concave. Occiput with 2 blunt conical protuberances in lieu of ridges, and without spines; a slight occipital depression; preopercular angle with 3 radiating spines of nearly equal length, a smaller spine directed outward in advance of the middle one of the 3; below these a partially concealed spine directed downward and forward. Opercular rib very strong, sharp anteriorly, broadening behind and provided with 3 low ridges, not ending in a definite spine; a spinous point on subopercle and 1 on interopercle; none on shoulder. Anterior nasal tube short, the posterior margin prolonged into a laciniate flap; head well provided with slender cutaneous filaments, 3 on upper portion of eyeball, 4 in a transverse line behind

occiput, a very long one on opercular angle, and numerous shorter ones on opercle, jaws, and along anterior portion of lateral line. Branchiostegals 7. Body without plates or prickles; the head, including upper part of eye, and the upper anterior part of body, with sparsely distributed stellate granulations, visible only in large specimens. In our specimens the brown bar at base of caudal is followed by a wide white bar, sometimes more or less broken, the terminal half blackish, narrowly margined with white.

(*ζώνη*, band; *οὐρά*, tail.)

Malacocottus zonurus, BEAN, Proc. U. S. Nat. Mus. 1890, 43, off Trinity Islands, Alaska, at Albatross Station 2853, at 56° N., 154° W., in 159 fathoms (Type, No. 44643. Coll. *Albatross*); GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 411.

737. ARGYROCOTTUS, Herzenstein.

Argyrocottus, HERZENSTEIN, Mélanges Biol. Ac. Imp. Sci., XIII, 1892, 219, St. Petersburg.

Spinous dorsal short; ventrals extremely long, their tips extending beyond front of anal fin; gill membranes scarcely united to the isthmus, forming a broad fold across it, no slit behind the last gill; teeth on the vomer, none on the palatines; skin entirely naked, without scales or bony plates; preopercles with 3 small spines. North Pacific. (*ἄργυρος*, silver; *Cottus*.)

2366. ARGYROCOTTUS ZANDERI, Herzenstein.

Head $3\frac{1}{3}$; depth $4\frac{5}{6}$. C. VIII–15; A. 13; P. 14; V. 3; C. 18; lateral line with 35 tubes. Eye $3\frac{5}{9}$ in length of head; interorbital space $1\frac{1}{9}$ in eye; 2 well-developed crests at the vertex; maxillary reaching middle of eye; lower jaw slightly projecting; 2 anal spines; 3 small spines on preopercle, the upper half as long as the eye, turned upward, the others directed downward; longest dorsal spine $\frac{1}{2}$ length of head; pectoral reaching ninth ray of soft dorsal; ventrals reaching to the next to the last ray of anal. Color brownish, dark above, with numerous silvery spots bordered with darker, a silvery stripe from below the eye to the base of the lower jaw; another from eye to angle of preopercle; a row of 6 large irregular spots along middle of sides, many smaller ones below these; 2 smaller irregular spots behind these; a bell-shaped spot with a point directed upward on the belly between the ventrals; the first dorsal with its upper margin blackish, with milk-white spots dotted with black; a transparent spot at base of fourth and sixth rays; a small transparent spot near base of fourth ray; another behind sixth; about these spots the coloration is darker; soft dorsal almost uniformly dark; anal colorless, its border dark, with 2 or 3 dark spots on each ray; caudal uniformly dusky, the upper and lower rays blackish, with clear spots; ventral with darker cross streaks; pectorals with dark rings and streaks. Ventral in female probably much shorter than in male, the color less marked. Three specimens of this beautifully marked Cottoid taken in Shana Bay, Iturup Island, show the following characters: The branchiostegal membranes are widely joined across the throat, narrowly united in front of the middle line of isthmus, with a rather wide free margin behind; lateral line without plates; nasal spines small; 4 short spines on preopercular margin. These specimens answer well to the detailed description of the type, but are

still more ornate, in that they possess along the back a number of broad dark bars alternating with lighter bars, the former confluent below, with the ground color of the sides. In the largest specimen, 7 cm. long, the ventral fins extend only to base of third anal ray. There are no tubercles on the rays, and the membranes extend nearly to tips of the 2 outer rays, and ⅔ length of the inner ray. The ventral spine is slender, nearly as long as the inner ray, and is firmly adnate to outer ray. The smaller specimens are, respectively, 4 cm. and 3.5 cm. long, the ventrals reaching in one to front of anal, in the other to vent; fins are finely cross barred, more variegated than in the type. Known only from Saghalien Island, 1 specimen 92 mm. long (Herzenstein), and Iturup Island, where 3 specimens were obtained by the *Albatross* in 1896, our description from the latter. (Named for its discoverer, Dr. Zander.)

Argyrocottus zanderi, Herzenstein, Mélanges Biol. Ac. Imp. Sci., XIII, 1892, 219, Korsakow, Saghalien. (Type, No. 9679, Mus. St. Petersburg. Coll. Dr. Zander.)

738. POROCOTTUS, Gill.

Porocottus, Gill, Proc. Ac. Nat. Sci. Phila. 1859, 166 (*quadrifilis*).

This genus contains small sculpins, distinguished from *Myoxocephalus* by the presence of 4 spines on the opercle, the uppermost being curved or hooked upward. Lower jaw included; 1 suprascapular spine; lateral line modified, giving off pairs of diverging branches, with pores at their ends. Cirri above eye and on nape. Northern Pacific; the species not well known and possibly reducible to 1 or 2. (πόρος, pore; κόττος, *Cottus*.)

a. Dorsal rays VIII, 13 or 14; anal rays 11 or 12.
  b. Opercle with 3 ridges, the upper ending in a sharp spine; pores of lower jaw small; fins low. SELLARIS, 2367.
  bb. Opercular spines well developed; fins high; pores well developed. QUADRATUS, 2368.
  bbb. Opercle undescribed. POLARIS, 2369.
  bbbb. Opercle without rib or spine. QUADRIFILIS, 2370.
aa. Dorsal rays VI, 16; anal rays 14; lower jaw with large pores; opercular spines obsolete. TENTACULATUS, 2371.

2367. POROCOTTUS SELLARIS (Gilbert).

Head 2⅘ to 3 in length; depth 3⅗. D. VIII, 14 (13); A. 10 to 12 (usually 11); P. 16. Pairs of pores in lateral line 32 to 34. Branchiostegals 6. Head cuboid, the anterior profile of snout subvertical, the greatest width a trifle more than the depth at occiput. Cheeks subvertical. Interorbital region elevated, the supraorbital rim furnished posteriorly with a low tubercle which usually bears an inconspicuous cirrus. The interorbital space is rather wide, transversly concave, its least width 2½ in eye. Occipital depression well marked, the ridges unbroken, straight, converging rapidly backward, the distance between their tips but ⅓ that between their anterior ends. From the latter, 2 low ridges converge for a short distance onto the floor of occipital depression. The ridges do not terminate

in spines, but bear small cirri similar to those above eye. Like the latter, they may be indistinct or wanting. Maxillary reaching vertical from posterior margin of pupil, $2\frac{1}{3}$ in head; eye $3\frac{1}{8}$, slightly exceeding snout. Nasal spines very strong. Two short spines diverge from angle of preopercle, the upper one slightly curved, directed upward and backward, $\frac{1}{2}$ the diameter of orbit, the lower directed straight backward, $\frac{2}{3}$ the length of the upper one. Below these a short spinous projection concealed in the skin, and a longer spine directed downward and forward; opercle with 3 lengthwise ridges, the uppermost ending in a definite sharp spine; subopercular spine well developed; 1 also at posterior end of interopercle, and 1 at shoulder. Gill membranes broadly joined to isthmus, with a wide free fold posteriorly; a few small prickly plates behind axil of pectorals; skin otherwise smooth; occiput and nape thickly covered with minute dermal papillæ, interspersed with very small mucous pores, which are distributed also over the anterior part of the head; pores on mandible and preopercular margin small, not better developed than in other species of *Porocottus*. Lateral line giving off pairs of short diverging branches, at the tips of which are the pores. Dorsals connected at extreme base; soft dorsal terminating slightly behind the anal; fins all low; ventrals reaching vent; pectorals to opposite third ray of anal. Color in spirits: Upper parts brownish, with 2 very conspicuous white saddle-shaped bars extending downward and forward from back, the first, about as wide as orbit, extending from below the dorsal notch in a straight line toward lower axil of pectorals, including above, the posterior margin of spinous dorsal, and the extreme basal portion of first rays of soft dorsal; the second bar including dorsally the terminal portion of soft dorsal onto which it extends, and the anterior portion of caudal peduncle; like the anterior bar, it is variable in width, extending forward and downward to below caudal peduncle; the ground color becoming intensified along the margins of these light bars; caudal with a light basal bar which becomes widened and pure white in its lower half, adding $\frac{1}{8}$ to the series of conspicuous light markings; in some specimens, these white bars are tinged with brownish, hence less conspicuous; under parts whitish; lips and lower jaw with light and dark cross bars, which are often indistinct; branchiostegal membranes sometimes with indistinct cross bars; the darker interspaces below soft dorsal and on caudal peduncle usually encircle the body below, but are sometimes (in males) broken on middle of sides with spots and blotches of white (see figure in Nelson's report cited below); axil of pectorals usually with 2 round white spots, most distinct in males; basal portion of anterior face of pectorals dusky or black, in males with 2 round white spots; fin distally barred with light and dark; caudal cross barred; anal and ventrals with faint cross bars, or plain; dorsal blackish. Taken abundantly in Bristol Bay, Bering Sea, and neighboring parts of the Arctic, in depths 5 to 17 fathoms. We are unable to follow Dr. Bean in identifying this fish with *Porocottus quadrifilis*, Gill (Proc. Ac. Nat. Sci. Phila. 1859, 166). The latter is described as having a single hooked preopercular spine, an opercle without rib or spine, and large pores on head and lower jaw. None of these applies to

the present species. *P. quadrifilis* is also said to have 5 branchiostegal rays, but this is possibly an error. It is also described as having a slender superciliary filament, and 1 on each side of the nape. We do not find, to offset these, any details in the original description which agree strikingly with our fish, even the color being inapplicable. (Gilbert.) (*sellaris*, saddled.)

Cottus quadrifilis, BEAN, in NELSON'S Report Nat. Hist. Coll. in Alaska, 309, pl. 18, 1887; not *Porocottus quadrifilis*, GILL.

Acanthocottus sellaris, GILBERT, Report U. S. Fish Comm. 1893 (1896), 419, Bristol Bay, at Albatross Stations 3229, 3231, and others, in 5 to 17 fathoms.

2368. POROCOTTUS QUADRATUS, B. A. Bean, new species.

Head $2\frac{2}{3}$; depth $3\frac{3}{4}$; eye $3\frac{1}{2}$ in head; mandible $3\frac{1}{2}$; maxillary 3; interorbital width 2 in eye. D. VIII, 14; A. 12; V. I, 3; P. 16. Head rather short and broad, quadrangular; profile from tip of upper jaw ascending and thence gently sloping upward to origin of dorsal, from which point almost vertically to end of nasal spine, rounding over orbital ridge, the body gradually tapering to the tail, the depth of the caudal peduncle being contained about $3\frac{1}{3}$ times in greatest depth of body. Gill membranes united and forming a fold across the isthmus. Opercular spines well developed, the lowermost on opercle and that on preopercle being curved downward and upward. Cranial ridges ending in spines of small size. Numerous pores on head, those in front visible to the naked eye; tubular pores on body, especially above anal base, where they appear to the naked eye as raised white specks. Fins all well developed, large; length of first dorsal base little more than $\frac{1}{2}$ length of second; anal fin origin under third and fourth rays of second dorsal and ending opposite its last ray; length of longest dorsal spine about $\frac{3}{4}$ as long as longest ray, or equal to length of longest anal ray; pectorals large and broad; ventrals reaching past anal origin. Color reddish brown, relieved by much white; under parts whitish; head above and below brown, flecked with white; lips pale; a broad white half bar on body extending from end of spinous dorsal to fourth ray of second dorsal, another extending from sixth to ninth ray, and a third from last ray to near end of caudal peduncle, which it encircles in connection with a dark bar, the latter extending on the caudal fin; pectorals and caudal barred; ventrals with round black blotches forming rows on the rays. Bering Island; only the type known. (B. A. Bean.) (Type, No. 33875, U. S. Nat. Mus., a single example 3 inches long, Bering Island, 1883; collected by Dr. L. Stejneger.) (*quadratus*, four-angled.)

Porocottus quadratus, B. A. BEAN MS., in JORDAN & GILBERT, Rept. Fur Seal Investigations, 1898, Bering Island.

2369. POROCOTTUS POLARIS (Sabine).

This very imperfectly described species is no doubt allied to *Porocottus sellaris* and *quadrifilis*. It is thus characterized: D. VI, 13; P. 15; V. 5; A. 14; C. 14 (Sabine). D. VIII, 13; P. 15; V. 5; A. 15; C. 12 to 14 (Ross). This species is compared by Captain Sabine to *Cottus gobio*, but that has

the head almost unarmed. It agrees with *Cottus* (*Enophrys*) *claviger* in the number of rays in the dorsal, but its ventrals are described as having more rays than usual in this genus. Capt. J. C. Ross informs us that it seldom exceeds 2 inches in length, and that it is very abundant on the east side of the peninsula of Boothia, affording a plentiful supply of food to the numerous waterfowl which breed there. "A species of *Cottus*, similar in habits to *C. gobio*, very adundant on the shores of north Georgia (lat. 75°), inhabiting the pools of water left by the tide, and the mouths of small rivulets by which the snow on melting found its way to the sea; the largest individual did not equal 2 inches in length; the head is more compressed and not so much flattened as in the *Cottus quadricornis*, and is armed with 2 strong spines placed before and between the eyes; the gill covers are also armed with 4 strong spines; the pectoral fins are larger in proportion than those of *C. gobio*, and the upper jaw rather exceeds the lower; the lateral lines are furnished with a series of small tubercles directed backward; color light, with clusters of minute dusky spots." (Richardson; quoted from Sabine.) Not recognized by recent writers. Lütken compares it with *Icelus bicornis*, but it must be different, and it may be identical with *Porocottus quadrifilis;* it is at least a *Porocottus*. (*polaris*, polar.)

Cottus polaris, Sabine, App. Parry's First Voyage, ccxiii, north Georgia; J. C. Ross, App. liii; Richardson, Fauna Boreali-Americana, iii, 43, 1836; Jordan & Gilbert, Synopsis, 706.

2370. POROCOTTUS QUADRIFILIS, Gill.

This species is known only from Dr. Gill's account, as follows:

"D. VIII, 13. The color is purple, irregularly spotted with black. There is a dark spot under the eye, and another on the maxillary. The dorsals, caudal, and pectorals are irregularly variegated with black. On the pectorals there is a dark spot at the upper axilla, and another larger one toward the middle of the base. There is a slender superciliary filament and 1 on each side of the nape. It is to the presence of these that the specific name is designed to draw attention. Specimens were obtained in Bering Straits at the same time as *Boreocottus axillaris*, Gill."

The genus *Porocottus*, Gill, is thus described:

"Body anteriorly subcylindrical, rapidly declining to the caudal. Skin naked. Lateral line opening by pores in raised papillæ, under a cutaneous keel. Head large and depressed, subrhomboidal. Nasal spines small. Preopercle with a single hooked spine. Opercle without a longitudinal rib or spine. Large pores under the lower jaw and on various parts of the head. Mouth moderate. Teeth on the jaws and front of the vomer. Branchiostegal membrane continuous under the throat, but attached along the middle to the throat, almost as far as the margin. Branchiostegal rays 5. Dorsals separate; the first low, the second oblong and opposite the anal. Ventrals small, behind the pectorals, each with a spine and 3 soft rays. The *Porocotti* have a strong resemblance to the *Boreocotti*, but are distinguished by the naked skin, hooked preopercular

spine, the absence of a longitudinal rib on the opercle, numerous pores, and the presence of only 5 branchiostegal rays. (Gill.)"

Not recognized by recent collectors. (*quatuor*, four; *filum*, filament.)

Porocottus quadrifilis, GILL, Proc. Ac. Nat. Sci. Phila., 1859, 166, Bering Straits.
Cottus quadrifilis, JORDAN & GILBERT, Synopsis, 708, 1883.

2371. POROCOTTUS TENTACULATUS (Kner).

Head 3½. D. VI–16 or 17; A. 14 or 15; V. I, 13, P. 14 or 15. Head small, mouth small, the lower jaw included; eye large, 3 in head, longer than snout; interorbital space channel-like, less than ½ diameter of eye; a sharp spine before each eye, on which is a fringed tentacle; supraocular ridge ending in a blunt, forked, bony knob, on which is a small thread-like tentacle; the quadrangular interspace between these two sets of tentacles is excavated; preopercular spine long, curved upwards, ⅔ diameter of eye; the second spine much shorter and turned backward; 2 spines lower, turned downward; subopercle with a spine turned downward; a similar one on opercle; maxillary reaching to middle of eye; lower jaw with large pores. Dorsal fins low, the rays flexible, the two close together; ventrals reaching vent; pectorals past beginning of anal. Skin of body wholly naked; lateral line complete, bending downward on caudal peduncle. Color clear brown, darker above, the head above with a few large dark brown spots, which form obscure bands, 1 of these from front of eye across upper lip, second broader from eye across subopercle; lower jaw speckled; throat and breast plain yellowish; back with about 6 dark cross bands, those most anterior the broadest; sides of body with a network of brown streaks around pale spots; a large bright yellow spot at base of caudal; fins all finely spotted with whitish and dotted with dark, the caudal with 5 or 6 dark cross bands; pectorals faintly barred (Kner.) A single specimen, 2 inches long, said to be from Singapore, which is of course an error. It probably came from the Pacific coast of Asia, perhaps from Yezo or Decastris Bay. As Kner has indicated, this species has much in common with *Porocottus quadrifilis*, but the fin rays are more numerous. It is, however, not impossible that the two are identical and both may be identical with *P. sellaris* and *P. polaris*. (*tentaculatus*, bearing tentacles.)

Cottus tentaculatus, KNER, Sitzber. Akad. Wiss. Wien, LVIII, 1868, 22, taf. 5, fig. 12, Singapore; evidently an error. (No. 5591a Wien Mus.)

739. ONCOCOTTUS, Gill.

Oncocottus, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 13 (*quadricornis*).

This genus contains small sculpins allied to *Myoxocephalus*, but with 4 preopercular spines, the uppermost being straight; a rather large slit behind the last gill arch; the lateral line chain-like in form. Mouth small, the lower jaw included, the dentition as in *Myoxocephalus*. Isthmus

narrow, the gill membranes forming a broad fold across it, males with rough stellate tubercles and with the crest on top of head rough and formed like the comb of a cock, these little developed in the female. Bones of head somewhat cavernous, but less so than in *Triglopsis*, from which genus *Oncocottus* is scarcely distinct. Species circumpolar, probably reducible to one. (ὄγκος, hook; κόττος, *Cottus*.)

a. Atlantic species; maxillary short; pectoral moderate; caudal rounded.
QUADRICORNIS, 2372.

aa. Arctic American species; maxillary longer; pectoral longer; caudal truncate.
HEXACORNIS, 2373.

2372. ONCOCOTTUS QUADRICORNIS (Linnæus).

Head 3¼; depth 4; eye 4 in head. D. VIII or IX–14; A. 13 or 14; cæca 7; vertebræ 40; lateral line 45. Body rather slender, the caudal peduncle very slender. Head long, tapering forward; mouth large, the maxillary reaching to below posterior margin of eye; lower jaw included; bones of head below eye cavernous, as in *Triglopsis thompsoni*, but less so; preopercle with 2 long, diverging spines, the upper and longer not quite reaching opercular margin; opercular and scapular spines quite short; adult male with a rugose spine, broader and expanded at tip like a cock's comb above each eye posteriorly, and a similar one on each side of occiput; these spines smaller in the female and the young. Males with irregular series of round, rough, wart-like scales above the lateral line, these wanting in the female; lateral line chain-like, with small embedded plates; head naked. First dorsal convex, of slender spines, well separated from second, which is rather high, the longest ray 1⅔ in head; pectorals reaching anal; ventrals moderate; caudal rounded. A moderate slit behind last gill. Color little variegated; olivaceous above; the fins faintly spotted. Arctic regions, south to the Baltic Sea, westward to eastern Greenland. If it is identical with *O. hexacornis*, its range extends westward to Siberia, and its distribution is circumpolar. It is abundant in the eastern Baltic and in Lakes Ladoga and Onega where it is dwarfish in size, without the horns on front and nape;* north to the White Sea and Nova Zembla. It is said to be rare in England and eastern Greenland; unknown in western Greenland. Dr Lütken has compared specimens from eastern Greenland with others from the Baltic and finds no difference except that the interorbital space in European examples is more concave. *O. hexacornis* is found in the waters of Arctic America and may not be different. (*quatuor*, four; *cornu*, horn).

Cottus quadricornis, LINNÆUS, Syst. Nat., Ed. x, 264, 1758, Baltic Sea; GÜNTHER, Cat., II, 166, 1860; LÜTKEN, Vid. Med. Kjöb., 87, 1876; DAY, Fish. Great Brit., 53, 1880; JORDAN & GILBERT, Synopsis, 705, 1883.

Oncocottus quadricornis, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 13.

? *Acanthocottus labradoricus*, GIRARD, Bost. Journ. Nat. Hist., VI, 1850, 247 pl. 7. fig. 3., coast of Labrador; female.

* These dwarf specimens may not be separable from *Triglopsis*, which genus is evidently derived from the lacustrine degradation of *Oncocottus*.

2373. ONCOCOTTUS HEXACORNIS (Richardson).

D. VIII or IX, 14 or 15; A. 14 or 15; P. 17; V. I, 3; Br. 6. Head $3\frac{1}{3}$ (to end of opercular spine); eye 5; snout $3\frac{1}{2}$; interorbital space $6\frac{1}{2}$; maxillary 2. Body slender, tapering gradually from the rather narrow depressed head to a very slender caudal peduncle; mouth large and horizontal; the maxillary reaching past the orbit; lower jaw shorter than the upper but not quite included within it. The spine on the head smooth and without tubercles or warts. In place of the supraocular and occipital spines there are 4 large bony elevations, each much resembling the comb of a cock, their upper surfaces rough and scabrous. (In the young these 4 big bones are very small or warty.) The occipital ridges scarcely elevated and inclosing an oblong and slightly concave area narrowest posteriorly and running into the concave interorbital space anteriorly; nasal spines well developed and in the older fish inclined to be club-shaped and scabrous; upper angle of the superclavicle prolonged posteriorly into a strong spine, its upper surface scabrous; just at the base and immediately in front of it a much smaller spine or tubercle on the posterior end of the post-temporal bone; a spine almost concealed in membrane just above base of pectoral; preopercle with 4 spines, the upper equaling the orbit in length and extending upward and backward, in some curved slightly inward, covered with membrane for about $\frac{1}{2}$ its length; the second spine immediately at the base of the first and is $\frac{1}{2}$ as long, straight, and extending outward and backward; the third, found farther down, curved downward; the fourth, at the lower angle of the preopercle, directed forward and downward and sometimes entirely covered by membrane; opercular spine rather strong, its point raised from the membrane and not reaching end of opercular flap. There is a slender sharp spine at the lower angle of the opercle, lateral line running rather high and composed of elongated rectangular plates, each with a concave depression at either end, thus making a row of elliptical depressions along the lateral line; the plates not present on the posterior half of the caudal peduncle; above the lateral line a row of rough circular osseous plates, beginning on the nape and running to the base of the caudal, gradually becoming smaller and nearer together; under the second dorsal this row is double; below the lateral line and beneath the second dorsal about 3 irregular rows of similar but smaller plates.

Adult female.—First dorsal lower than in male, scarcely rounded posteriorly but sloping almost gradually toward the base of the second dorsal; third and fourth spines longest and equal to length of snout; second dorsal about twice as high as first and varying a great deal in shape, the membrane scarcely incised, the first 3 or 4 rays with rough, prickly edges; anal oblong and slightly rounded, the middle rays longer than snout, the membrane between the rays not incised; pectoral scarcely reaching front of second dorsal, first 2 or 3 rays rough on their outer edge, the membrane not incised; caudal not rounded but truncated with rather sharp corners, 10 developed rays, 8 of them branched; ventrals reaching $\frac{3}{5}$ to the vent, the inner ray no longer than the middle one.

Adult male.—The first dorsal higher than in the female, the third and fourth rays longest and equal in length to the distance from snout to middle of eye, posterior end of fin more rounded; second dorsal more than

twice as high as first, the membrane between the rays deeply incised, the anterior rays projecting for nearly ½ their length beyond the membrane, the membrane following around the edge of the projecting rays, the rays scabrous on their sides; anal oblong, but more angular than in female, the longest ray reaching from snout to posterior edge of orbit; the membrane between the first 4 rays quite deeply incised, outer edge of rays scabrous; ventrals reaching ⅓ to vent, the inner ray longest.

Color.—In the young the color is gray above and white beneath, a dark saddle marking over the back at the center of the first dorsal; another similar marking under the anterior, and another under the posterior end of the second dorsal; a dark spot the size of the eye on top of the caudal peduncle and midway in its length (the young may be distinguished at a glance by this marking on top of the caudal peduncle); a V-shaped marking at the base of the caudal with the angle of the V on the lateral line and directed forward; this marking in the older specimen extends forward into the peduncle spot; the caudal is marked by 2 vertical wavy bands on its posterior half, the first one the wider, the last one at the tips of the rays; these bands are solid, extending across the rays and membrane; the pectorals are dark at the base, the rest marked by 3 vertical black bars; the first dorsal is slightly dusted with black, the second is slightly mottled with black, but with no bars; the anal has 3 or 4 vertical dark bars. In the adults the general appearance of the fish is very much darker, almost black; the lower parts are light; the markings are not so distinct; the fins except the ventrals are all dark, and the markings on the pectorals, caudal, and anal run together into solid black; the lower parts, especially in the males, brilliantly colored with red. (Scofield.) Arctic waters of America, from western Greenland to Bering Straits. This fish seems to differ from *Oncocottus quadricornis* from the Baltic Sea, having a longer maxillary, longer pectorals, and a square-cut caudal fin. All these characters are of doubtful value, and the species may be the same as the European *O. quadricornis*. We can find no good difference between our specimen and a small one from Arctic Bay, western Greenland (received from Prof. D'Arcy W. Thompson). The fish figured by Nelson as *Cottus quadricornis* (Nelson, Natural History of Alaska) is probably the same as the one here described, but according to the figure there are several differences. Scofield and Seale found this fish very abundant along the shores of Herchel Island. It was about the only fish to be found where the bottom was muddy. A few young were taken at Point Barrow, and at Port Clarence, both young and old; 3 of the young were found in the river back of Grantly Harbor. (ἕξ, six; *cornu*, horn.)

*Cottus hexacornis.** RICHARDSON, Franklin's Journal, 1823, 726, mouth of Tree River near Coppermine River (Coll. John Franklin): RICHARDSON, Fauna Bor.-Amer., III, 44, 1836; GÜNTHER Cat., II, 166, 1860.

Acanthocottus labradoricus, GIRARD, Bost. Journ. Nat. Hist. VI, 1850, 247, pl. 7, fig. 3, Coast of Labrador.

Oncocottus hexacornis, SCOFIELD, in Jordan & Gilbert, Rept. Fur Seal Invest., 1898.

* *Cottus hexacornis*, Richardson, was thus originally described:
D. VII, 13; A. ; C. 12; P. 16; Br. 6. Head large and depressed. Eyes large. Six club-shaped, or rather nail-shaped, processes stand erect on the top of the head, their summits flattish, minutely cancellated, and scabrous; the smallest pair stand between

Cottus labradoricus,* GÜNTHER, Cat., II, 163, 1860; BEAN, Proc. U. S. Nat. Mus., IV, 1881, 128; JORDAN & GILBERT, Synopsis, 704, 1883.

the nares; the largest over the posterior angles of the orbits, and the third, of intermediate size, on the occiput. The mouth is very capacious; its margins are formed by the intermaxillaries and lower jaw; the maxillaries have an elongated wedge form, and lie in a membrane behind the intermaxillaries. Both jaws and vomer are set with bands of 9 teeth, en velours. Tongue obtuse and smooth, as are the palate and maxillaries. The preoperculum is armed beneath with 3 strong, devaricated spines, the posterior one, which measures ½ inch, being the longest. The gill covers are composed of several bones connected by membrane, and armed on the exterior edges with 4 or 5 small, spinous teeth. The bones which support the pectoral fins are also armed with small spines, and have sharp, rough edges. The branchiostegous membrane contains 6 slender, cylindrical, curved rays. The body is much narrower than the head, and tapers to the insertion of the caudal fin. The anus is situated midway between the mouth and the caudal. The lateral line is rough, and runs near the back; above it there is a row of small, orbicular, scabrous, bony plates, the row being doubled opposite to the second dorsal. There are no other perceptible scales. The pectoral fins are suborbicular and contain 16 rays, none of them branched; the upper ray is scabrous throughout; the others are scabrous only near their middles. The ventrals, soft and whitish, have 3 rays, of which the first is the strongest, but none of them are spinous. The first dorsal commences posterior to the pectorals, and terminates opposite to the anus; it has 7 simple rays; the second dorsal is larger, and has 13 rays; its commencement and termination correspond with those of the anal, and most of its rays are scabrous; both dorsals are rounded or arched. The anal fin occupies about ⅔ of the space between the anus and caudal, commencing near the former; this fin becomes slightly lower or less deep posteriorly. The caudal is cuneiform, and has 12 rays, most of them forked. In the form of the bony processes on the top of the head this species approaches closely to *C. quadricornis* of the Baltic; but it does not appear from the descriptions I have consulted that there is a distinct pair on the nasal bones of the latter. There are also differences in the form of the spines of the preoperculum, those of *C. hexacornis* being quite simple, while in the other they are truncated, or divided at the point. In the *C. quadricornis*, also, there is a thick spine on the suprascapular bone, which is likewise truncated; while in *C. hexacornis*, that bone, the humerals, and the gill covers are merely armed with small, spinous teeth; and the rows of scales on the body are different. It appears to me likely that the *C. quadricornis*, Sabine (Zool., App. to Captain Parry's First Voyage, p. ccxiii), may be really the *C. hexacornis*. Capt. J. C. Ross, who considers it to be the same with the *C. scorpioides* of Fabricius, says that though very abundant on the Greenland Coast it is more rare in the higher latitudes, but several were taken on both sides of the peninsula of Boothia. The natives prize it highly as an article of food, preferring it to codfish or salmon. The Esquimaux of Boothia call it Kaneeok, the same name which the Greenlanders give to *C. Grœnlandicus*.

Color.—Of the upper aspect a clouded admixture of brocoli-brown and olive-green tints; of the belly white. The fins are streaked with bluish black. Irides tinged with red.

Size.—About 7 inches in total length.

Numerous specimens of this fish were caught in a net set in the mouth of a small river near the Coppermine, and the above description is drawn up from notes written on the spot. The subsequent calamities which befell that expedition having occasioned the loss of all the specimens, no actual comparison has been made with other species; but after an inspection of the Cotti brought home by Captain Beechey, and an attentive perusal of the Histoire des Poissons, I am satisfied that it differs from all other described species in the form of the horns or processes which arm the head, and in other particulars. From the peculiar shape of these horns in our species it might bear the name of *claviger* still more appropriately than the one so termed by *M. valenciennes*. The individuals that we caught retained life long after they were drawn from the water, leaping vigorously over the sands, and when touched inflating the head. In this operation the branchiostegous membrane is distended, and the several pieces composing the gill covers are separated by the extension of the intervening membranes. Our Canadian voyageurs were both astonished and alarmed by these fish, and termed them Crapauds de mer, probably from a kind of croak they uttered when first handled. (Richardson.)

In 1849, having again visited the same region, Richardson procured more specimens of this form which he took home with him. He saw then that he had misunderstood his notes, and wrongly ascribed to *C. hexacornis* rostral horns of the same nature as the characteristic horns of the head and nape, and that the form in question was no other than *C. quadricornis*, an opinion all must accept who, recalling this correction, will read the old description of Sir John. (Lütken.)

*Specimens of *Oncocottus labradoricus*, Girard, are thus described by Dr. Bean:

Eye 5; snout nearly 5. D. X, 14; A. 14; V. I, 3; P. 17; C. 11 (developed). Two small spines above the snout; a rough, irregular prominence above each orbit and 2 similar ones on the occiput. The slight depression on the crown becomes narrower posteriorly where its width is about ½ the length of the space included between the supraorbital and occipital prominences. Four preopercular spines, 2 of which are at the angle, the uppermost and longest is ⅔ as long as the eye; the 2 lower spines are short and extend downward and slightly forward. The length of the longest preopercular spine equals the distance between the eyes measured on the bone. The maxillary is twice as long as

740. TRIGLOPSIS, Girard.

Triglopsis, GIRARD, Proc. Bost. Soc. Nat. Hist., IV, 1851, 18 (*thompsoni*).

Ptyonotus, GÜNTHER, Cat., II, 175, 1860 (*thompsoni;* substitute for *Triglopsis* on account of the prior and similar name, *Triglops*).

Body and head slender. Skin naked; lateral line chain-like. Teeth on vomer, none on the palatines. Eyes large, the interorbital area concave; bones of lower part of head extensively cavernous; a small but distinct slit behind last gill; gill membranes almost free from the isthmus, forming a broad fold across it; preopercular spines straight, simple, 4 in number, the lower turned downward; fins large. Fresh-water fishes, closely related to *Oncocottus*, from which they have doubtless become degraded through fresh-water life. There is no tangible difference in structure in any part of the body. (*τρίγλα, Trigla; ὄψις*, appearance.)

2374. TRIGLOPSIS THOMPSONI, Girard.

Head 3; depth 6. D. VII, 18; A. 15; V. I, 3. Body elongate, very slender. Head long, depressed above; snout long and pointed; eye quite large, nearly as long as snout, much wider than interorbital space, 4 in head; jaws subequal; mouth large, the maxillary extending rather beyond middle of eye; preopercle with 4 sharp spines, the upper much shorter than pupil; cavernous structure of skull highly developed; upper surface of head smooth; gill membranes not broadly united, nearly free from isthmus. Dorsal fins well separated; spinous dorsal short and low, its height little more than length of snout; second dorsal very large, 3 times height of first, its longest rays about as long as head; anal high, $\frac{1}{2}$ as high as second dorsal; pectoral long, reaching past front of anal; ventrals well developed; lateral line chain-like, conspicuous; skin perfectly smooth. Pale olivaceous, with darker blotches; upper fins faintly banded. Length 3 inches. Deep waters of the Great Lakes; not common; known from Lake Michigan and Lake Ontario; doubtless a relic of a former Arctic marine fauna, and descended from a species of *Oncocottus*. (Named for Rev. Zadock Thompson, the accomplised author of the History of Vermont.)

Triglopsis thompsoni, GIRARD, Proc. Bost. Soc. Nat. Hist., IV, 1851, 19, off Oswego, Lake Ontario (Coll. S. F. Baird); GIRARD, Monograph Cottoids, 65; JORDAN & GILBERT, Synopsis, 709, 1883.

Triglopsis stimpsoni, HOY, Trans. Wis. Ac. Sci. 1872, 98, Lake Michigan. (Coll. Dr. William Stimpson.)

Ptyonotus thompsonii, GÜNTHER, Cat., II, 175, 1860.

---

the eye, about $\frac{1}{3}$ as long as the head to the end of the opercular spine, and extends about to the vertical through the hind margin of the eye. Teeth on the vomer, none on the palatines. Dorsal spines slender; the first twice as long as distance between eyes; third and longest $\frac{2}{3}$ as long as maxillary and $\frac{1}{2}$ as long as distance from tip of snout to end of occipital prominences. Base of spinous dorsal $\frac{1}{3}$ total length without caudal; distance between dorsals 2 in eye. Longest soft dorsal ray (ninth) about as long as middle caudal rays, or $\frac{1}{8}$ total length with caudal; pectorals reaching a little beyond origin of anal (to second ray of anal); ventrals as long as postorbital part of head and not reaching vent. Skin above the lateral line with a few spiny tubercles, none of them more than $\frac{1}{4}$ as long as eye. A small slit behind fourth gill. Coast of Labrador and Hudson Bay. This description by Dr. Bean from specimens taken in 1880 near York Factory, Hudson Bay, by Robert Bell, M. D. This nominal species is apparently the female of *Oncocottus hexacornis*.

741. GYMNOCANTHUS, Swainson.

Gymnocanthus, SWAINSON, Class. Fish., etc., II, 271, 1839 (*ventralis*).
Phobetor, KRÖYER, Naturh. Tidschr., I, 263, 1844 (*tricuspis*).
Elaphocottus, SAUVAGE, Nouv. Arch. Mus. Paris (2), I, 1878, 142 (*pistilliger*).

General characters and appearance of *Acanthocottus*, but with no teeth on the vomer, the slit behind the last gill small or wanting; upper preopercular spine stout, armed with 2 or 3 antler-like processes; fins all very large, the ventrals notably so. Arctic Seas. (*γυμνός*, naked; *ἄκανθα*, spine.)

a. Interorbital space not armed with bony granulations; males with a round white spot on belly; last gill arch without slit or pore.
b. Dorsal rays X, 14 or 15; anal 16; a cirrus behind eye in young; occipital ridge with 3 bony prominences; axial region in male with fringed filaments, white at tip. PISTILLIGER, 2375.
bb. Dorsal rays XII, 16; anal 18; no cirrus behind eye; occiput without bony prominences; axial region without fringed filaments in either sex. TRICUSPIS, 2376.
aa. Interorbital region covered with bony granulations; males without white spots on belly; dorsal rays XI, 16; anal 19. GALEATUS, 2377.

2375. GYMNOCANTHUS PISTILLIGER (Pallas).

Dr. Gilbert describes this species as follows:

"Compared with *Gymnocanthus tricuspis*, *G. pistilliger* has a different fin formula, the spines and rays of dorsal and anal fins being fewer in number. This is shown by the following table, based on an examination of 40 specimens:

	Spinous dorsal.		Soft dorsal.				Anal.		
Rays	IX.	X.	13.	14.	15.	16.	15.	16.	17.
Specimens	10	30	1	17	21	1	2	28	10

"The normal formula may therefore be given: D. IX or X, 14 or 15; A. 16 or 17. The type of *G. pistilliger* is said to have D. IX, 13; A. 16; and its synonyms, *G. ventralis*, Cuvier & Valenciennes, and *G. intermedius*, Temminck & Schlegel, have, respectively, D. IX, 13; A. 17; and D. IX, 13; A. 14. In *G. tricuspis* the formula is D. XI or XII (rarely X), 15 to 17; A. 16 to 19. The dorsal fins are more widely separated in *G. pistilliger*, where the interspace is equal to $\frac{1}{2}$ or more than $\frac{1}{2}$ the diameter of the pupil. The vertebræ are fewer in number, 12 + 24 instead of 12 + 28. In *G. pistilliger* an obtuse prominence above hinder margin of orbit bears in young individuals a slender cirrus, which frequently disappears in adults. Behind the eye, a continuous occipital ridge bears 3 smaller bony prominences, the first immediately behind the eye, the second and third approximated at posterior end of ridge; these bear no cirri. *G. tricuspis* has no tubercles on occiput. In males of *G. pistilliger*, the post axial region is furnished with a number of very slender filaments, each of which is expanded at tip

into a compressed froud-like lamina, having the free edge more or less laciniate or fringed. These expanded tips are bright white and very conspicuous. No trace of them is present in females, but they develop in males at a very early age. These agree with the structures described by Pallas, on which he based the name *pistilliger*. They seem not to be present in *G. tricuspis*. The upper preopercular spine is sharply bifurcate in even our smallest specimens (50 mm.), but in these no trace of a second medial upwardly directed spine is present. The latter is evident in specimens 70 mm. and more in length, and a small concealed prominence representing a third spine is exceptionally present. No trace of slit behind last gill. The following measurements give the depth of body and length of head in millimeters, as compared with total length, in 8 specimens:

Total length.	Length of head.	Depth.
mm.	*mm.*	*mm.*
156	44	27
145	39½	25½
142	38½	24½
135	38	24½
135	40	25
134	37	24
125	34½	21
93	25½	16

Very young examples show no groups of granulations on head or nape, these being usually wanting in specimens less than 100 mm. long. In older examples they are variously developed, the degree of armature dependent neither on age nor sex. They are never armed on interorbital space, being unlike *G. galeatus* in this respect, the granulations being confined to the occipital and nuchal regions, with an additional elongate patch on the upper part of the opercle. In highly developed males, the dorsal and ventral rays are accompanied with series of tubercles. The color is brown above, with very narrow vermiculating lines of lighter; a black blotch on cheek, more conspicuous in males, and 4 inconspicuous cross bars on back; the darker dorsal area is bounded below lateral line by an irregular series of dark streaks or blotches. In males, the lower jaw and preopercle is cross-banded with black and light yellow; the abdomen, lower half of sides in front of anus, and prepectoral region, with large roundish white spots, separated by vermiculating areas, rendered dusky by aggregations of coarse black dots; ventrals dusky and silvery, the latter frequently forming cross bands; spinous dorsal dusky or black, with irregular series of white spots not confined to basal parts of fin. In both sexes the pectorals, second dorsal, and caudal are translucent or yellowish, crossed by narrow black bars. The females are more numerous than the males in our collection, but the disparity in number is not so great as has been found by other writers. In 45 specimens examined

as to this, 17 are males, 28 females. Among specimens obtained at Petropaulski and at *Albatross* Station 3646, off Robben Island, in 18 fathoms, no males are included. The females differ from those we have examined from Bristol Bay in having the top of head more extensively plated, the rough plates extending onto middle of interorbital space, or in 1 specimen onto snout. The preorbital ridges are less regular and have lower tubercles. The specimens indicate an approach therefore to *G. galeatus*. The fin rays are as previously given. In 8 specimens they are as follows:

	Dorsal spines.		Dorsal rays.			Anal rays.	
Ray	IX.	X.	14.	15.	16.	16.	17.
Specimens	1	7	1	6	1	3	5

Coasts of Alaska; taken abundantly in Bristol Bay in 4½ to 26 fathoms." (*pistilla*, pistil; *gero*, I bear; in allusion to the axillary papillæ of the male.)

Cottus pistilliger, PALLAS, Zoogr. Rosso-Asiat., III, 43, 1811, Unalaska; CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 193, 1829; GÜNTHER, Cat., II, 167, 1860.

Cottus ventralis, CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 194, 1829. Kamchatka (Coll. Mr. Collée. Type in Brit. Mus.); GÜNTHER, Cat., II, 167, 1860.

Cottus cephaloides, GRAY, in CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 194, 1829, Kamchatka. (Type of *C. ventralis*.)

Elaphocottus pistilliger, SAUVAGE, Nouv. Archiv. Mus. 1878, 142.

Gymnocanthus pistilliger, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 424; SCOFIELD,* Rept. Fur Seal Investigations, 1898.

2376. GYMNOCANTHUS TRICUSPIS (Reinhardt).

Head 3⅓; depth 4⅔. D. XII, 16; A. 18; V. I, 3; P. 16; vertebræ 12 + 28 = 40. Nasal spines small; occipital ridges low, the space between them

*Mr. Scofield has the following note on this species:
Three specimens taken at Grantly Harbor, Port Clarence, Alaska.

Fin formulæ.

	D.	A.	P.	C. branched rays.
♂	14	16	18	9
♂	15	17	18	9
♀	13	15	17	9

I have compared them with specimens taken by the *Albatross* in southern Bering Sea and find them the same. In the female taken at Port Clarence the crown (not between the eyes), nape, and upper half of opercles are covered with scabrous bony plates. These plates are present in some of the females taken by the *Albatross*, but are not so evident. The female has a few scattered rough prickles back of the origin of the pectorals. The males have the mushroom-like filaments covered by the pectorals, and the middle rays of the pectorals have papillæ along their inner edge. This fish has been reported from the Atlantic side of North America, but it is probably a different species, as the mushroom-like filaments are not mentioned in their description. Sir John Richardson took a single specimen, a female 5 or 6 inches long, at Hudson Bay, which appears to be the same as these from Bering Sea. He identified it as *Phobetor tricuspis* (Kröyer), and his plate of it agrees perfectly with our specimens, but in his description he does not mention the prickles back of the origin of the pectorals, and he mentions a few scabrous bony plates just below the second dorsal which our specimens do not have.

concave, with rough plates; supraorbital ridge ending in a blunt tubercle; interorbital area with smooth skin. Eyes very large; maxillary extending to pupil; upper preopercular spine broad; much shorter than eye, with about 3 points; skin mostly smooth; axil prickly, but without fringed filaments; inner edge of middle pectoral rays papillose (male). Spinous dorsal very high and long, the longest spines in the male ¾ length of head; second dorsal a little lower; anal very long, rather low; pectorals very broad, reaching past front of anal, the lower rays rapidly shortened; ventrals extremely long, the rays long, exserted, reaching past front of anal; anal papilla large. Dark brown above, with traces of darker vertical bars; belly pale; males with the axillary region dusky, with many large round white spots; first dorsal blackish, with pale blotches; second dorsal with alternating oblique bands of white and blackish; anal and caudal nearly plain; pectorals and ventrals yellowish, with black cross bars; mandible barred with black. Arctic seas, south to Norway and Labrador; not very common on our coasts. Here described from a specimen from Greenland. Dr. Günther gives the following anatomical details: "The liver is large, round, not divided into lobes, and situated principally on the left side of the stomach. The stomach is very spacious and curved; the pylorus with 6 appendages; the intestines appear to make 1 complete circumvolution. The ovaria are separated from each other to their posterior extremity. The urine bladder is narrow, elongate, situated above the right-hand ovarium. Skeleton: The configuration of the skull is much more similar to *C. gobio* than to *C. scorpius* or *bubalis*. The space between the orbits is very slightly concave, very narrow, its width being nearly ⅓ the distance between the upper posterior angles of the orbits. The crown is flat, without any longitudinal or transverse ridges, but with very slight impression in the middle. The frontal bones, the preoperculum, the mandibula, and the infraorbitals have very distinct muciferous channels; the turbinals are provided with a minute spine. The number of the caudal vertebræ is increased, there being 12 in the abdominal portion and 28 in the caudal." (Eu.) (*tres*, three; *cuspis*, cusp.)

Cottus gobio, FABRICIUS, Fauna Grœnlandica, No. 15, 1780, Greenland.

Cottus tricuspis, REINHARDT, Vidensk. Selsk. Nat. Math. Afhandl., VII, 1838, 117, Greenland; GÜNTHER, Cat., II, 168, 1860.

Cottus fabricii, GIRARD, Monograph Cottoids, 59, 1851, Greenland; after *Cottus gobio*, FABRICIUS.

Cottus ventralis, COLLETT, Christiania Vid. Selsk. Forh. 1878, 151; not of CUVIER & VALENCIENNES.

Acanthocottus patris,* H. R. STORER, Bost. Jour. Nat. Hist.. VI, 1857, 250, Labrador. (Coll. Dr. Horatio Robinson Storer.)

Phobetor tricuspis, KRÖYER, Natur. Tidskr., I, 263, 1844.

Gymnocanthus pistilliger, JORDAN & GILBERT, Synopsis, 709, 1883; not of PALLAS.

* This southern form, named "*patris*" by Dr. H. R. Storer, for his distinguished father, needs comparison with *Gymnocanthus tricuspis*.

2377. GYMNOCANTHUS GALEATUS, Bean.

Head 3¾ (including caudal); depth 7½. D. XI, 16; A. 19; V. I, 3. Body elongate. A small tubercle above each eye; 4 preopercular spines, the longest about as long as eye, and with 2 or 3 processes. Space between eyes deeply concave, completely covered with bony granulations, as are the crown and neck; similar granulations on hinder margin of orbit, on suborbital stay and on opercle. Skin of body naked. Pectorals, and in males the ventrals also, reaching beyond vent; maxillary reaching to below eye. Slit behind last gill not described. Olivaceous; back with 4 distinct brown spots, the longest nearly twice as long as eye, and extending a little below lateral line, there blending with a wavy lateral stripe; dorsals and pectorals with interrupted black band; lower fins plain whitish. Dr. Gilbert describes a single male specimen, 210 mm. long, from Chernofski Harbor, Unalaska Island:

The sexual peculiarities are less strongly marked than in much smaller males of *G. pistilliger*. The ventrals extend but little beyond the front of the anal fin, and are unmarked. The abdomen is also plain, without the round white spot characteristic of male specimens of *G. pistilliger* and *G. tricuspis*. These marks are apparently absent also in the type, as no mention is made of them. The spinous dorsal is not greatly elevated, its longest spine being contained 2⅓ times in the head. It is without distinctive markings, being colored like the soft dorsal, light yellowish, cross-banded with darker. Anal papilla short, 5 mm. in length. Plates on head as in description, covering the interorbital region, crown, occiput, and nape as far back as front of dorsal; present also on upper part of opercle, and in a vertical streak immediately behind eye. A very prominent tubercle over hinder margin of eye, a constriction behind it. Occipital ridge broadly rounded, without tubercles or conspicuous prominences of any kind, but with 3 smooth areas corresponding in position with the tubercles of *G. pistilliger*, perhaps present as such in the young. The anterior one is slightly elevated. No superciliary filaments. Preopercular spines massive, short, with a single fork at tip, no accessory spines developed.* Axil with prickly scales, but without filaments. Dorsal XI, 16; anal 19. Lateral line 43. Head 52 mm. long; depth 32 mm. The depth is there 6½ in the total length, not 7½ as described for the type.

Aleutian and Pribilof islands; known from Unalaska and St. Paul, and Point Barrow. (*galeatus*, helmeted.)

Gymnocanthus galeatus, Bean, Proc. U. S. Nat. Mus. 1881, 153, Unalaska (Type, No. 28097. Coll. Sylvanus Bailey); Jordan & Gilbert, Synopsis, 707, 1883; Gilbert, Rept. U. S. Fish Comm. 1893 (1896), 425.

742. LEIOCOTTUS, Girard.

Leiocottus, Girard, Proc. Ac. Nat. Sci. Phila. 1856, 133 (*hirundo*).

Body elongate, covered with thick, smooth skin. Head compressed, narrowed above, not externally bony. Mouth small, horizontal, low; villiform teeth in both jaws and on the vomer, none on the palatines; upper

* Mr. Scofield has the following note on a small specimen (3½ inches) collected at Point Barrow, Alaska:

Fin formula: ♀ D. X, 15; A. 17; P. 18; V. 3. Head 3¼ or 3$^{6}_{10}$, including caudal; eye 3⅓ to end of opercular spine; snout 4; maxillary 2¾ or past posterior edge of pupil; interorbital 2 in length of orbit, depth 5¼; Lat. line 43 or 44. Fifth or sixth ray of pectoral longest and reaching to third anal ray. Ventrals reaching ⅔ to vent. Mouth slightly oblique. Upper preopercular spine with 3 sharp points on its upper side. No papillæ or sharp points under pectorals.

preopercular spine short, strong, with bifurcated tip; suborbital stay narrow; gill membranes partly free from the isthmus, over which they form a broad fold; a slit behind fourth gill. First dorsal with its upper margin somewhat S-shaped, the first 2 spines elongate, the middle ones of nearly equal length, and the posterior ones rapidly shortened; ventrals I, 3. Shore fishes of the Pacific. (*λεῖος*, smooth; *κόττος*, *Cottus*.)

2378. LEIOCOTTUS HIRUNDO, Girard.

Head $3\frac{1}{2}$; depth $4\frac{1}{2}$; eye large, $4\frac{1}{2}$ in head. D. IX, 17; A. 15; V. I, 3. Body elongate, fusiform, the caudal peduncle slender; profile of snout decurved; maxillary reaching front of orbit; 3 small cirri at the end of each maxillary; preopercular margin with several similar cirri; opercle and shoulder girdle without spines; upper preopercular spine as long as pupil. Top of head smooth; supraorbital ridges little elevated; nasal spines distant from tip of snout. Spinous dorsal elevated in front, the first 2 rays much longer than the others, nearly $\frac{2}{3}$ length of head; soft dorsal and anal fins rather long; pectorals reaching past front of anal; ventrals to vent. Olivaceous, shaded with light blue and reticulated with brownish red, the latter color predominating on the head; sides with 4 broad, oblique, brownish-red bars, the first 3 running from dorsal forward and downward, the fourth from caudal peduncle backward to base of caudal; abdomen orange brown, with pale spots; caudal orange brown, with yellow bars near the tip; fins with bars and spots of dark brownish red; breast and ventrals whitish; 3 dark blotches at base of pectorals; spinous dorsal with oblique dark streaks; a dark blotch on each eye above, and a light streak forward and downward from eye. Length 10 inches. Santa Barbara Islands, in shallow water; not rare, but extremely local. (*hirundo*, swallow.)

Leiocottus hirundo, Girard, Proc. Ac. Nat. Sci. Phila. 1856, 133, San Miguel Island, near Santa Barbara (Coll. Lieutenant Trowbridge), Girard, U. S. Pac. R. R. Surv., x, Fishes, 62, 1858.

Cottus hirundo, Günther, Cat., II, 166, 1860.

Liocottus hirundo, Jordan & Gilbert, Synopsis, 712, 1883.

743. LEPTOCOTTUS, Girard.

Leptocottus, Girard, Proc. Ac. Nat. Sci. Phila. 1854, 130 (*armatus*).

Body elongate, depressed, covered with perfectly smooth skin. Head depressed, oblong, not very broad, without cirri; lower jaw included; suborbital stay narrow; teeth on jaws, vomer, and palatines. Gill membranes rather narrowly joined to the isthmus, not forming a fold across it; a slit behind fourth gill; preopercular spine strong, with 2 or 3 points hooked upward. Dorsal fins separate; the spinous dorsal short and small, entire; ventrals I, 4. Shore fishes of the Pacific. (*λεπτός*, slender; *κόττος*, *Cottus*.)

2379. LEPTOCOTTUS ARMATUS, Girard.

(Smooth Cabezon.)

Head 3 in length; depth 6. D. VII, 17; A. 17; V. I, 4. Head long and depressed: mouth large, the maxillary reaching beyond eye; interorbital space broad, scarcely concave; nasal spines concealed; top of head flattish, covered with rugose skin; upper preopercular spine with 3 or 4 spinules hooked upward; suborbital stay slender, not reaching preopercle; eye very small, less than interocular width, 7½ in head; lateral line complete; skin everywhere smooth; no prickles nor cirri. Dorsal spines very slender and low; pectorals reaching or extending beyond vent; ventrals halfway to vent. Grayish olive above, becoming abruptly white and silvery below; sides creamy; pectoral fins creamy yellow, with 5 or 6 black cross bars; spinous dorsal dusky, with an ink-like blotch on tip of last rays, and an oblique white band below; soft dorsal dusky with several oblique white bands; caudal banded; ventrals and anal plain. Length 12 inches. Pacific Coast from Kadiak to San Diego; everywhere very common, the most abundant of the Cottoids of our west coast; living near shore. It flattens the head and expands the preopercular spines when disturbed. (*armatus*, armed.)

Leptocottus armatus, Girard, Proc. Ac. Nat. Sci. Phila. 1854, 131, Cape Flattery, Fort Steilacoom, Willapa Bay, Humboldt Bay, San Francisco, Monterey, San Pedro, Fort Point, San Diego, Tomales Bay (Coll. Dr. Suckley, Dr. Cooper, Lieut. Trowbridge, etc.); Girard, U. S. Pac. R. R. Surv., x, Fish., 60, 1858; Jordan & Gilbert, Synopsis, 714, 1883.

Centridermichthys armatus, Günther, Cat., II, 171, 1860.

744. CLINOCOTTUS, Gill.

Clinocottus, Gill, Proc. Ac. Nat. Sci. Phila. 1861, 166 (*analis*).

This genus differs from *Oligocottus* in having the skin of the sides of body mesially, covered more or less thickly with minute prickly scales embedded in the skin; anteriorly, forked cirri take the place of scales; mouth with lateral cleft; preopercular spine short, bifurcate. Small fishes of the California rock pools. (*Clinus; Cottus.*)

2380. CLINOCOTTUS ANALIS (Girard).

Head 3½; depth 4½. D. IX, 16 to 18; A. 13 or 14; V. I, 3; P. 15. Head narrower anteriorly and rather pointed; mouth with lateral cleft, the maxillary reaching beyond pupil; band of palatine teeth short and narrow, often absent on one side or on both; eye large, 5 in head, about twice the width of the deeply grooved interorbital space; nasal spines distant from snout; preopercular spine short, bifurcate; cranium plane above; upper part of body mesially covered with minute, embedded, nonimbricated, pectinate scales; anteriorly, small cirri, mostly bifid or trifid, take the place of the scales. Head, especially above, with very many similar cirri; a fringe of cirri on edge of preopercle. Dorsal fins contiguous, rather low; pectoral reaching past front of anal; anal papilla very large.

Olivaceous, sometimes very pale, sometimes nearly black, occasionally dashed with reddish, much mottled, and with numerous small black and white spots; about 5 irregular darker bars; a dark bar at base of caudal; fins all spotted; cirri very numerous, mostly whitish, giving the fish a woolly appearance in life. Length 3 to 7 inches. Coast of California; abundant in rock pools from Monterey southward to Lower California, farther south than any other of the California *Cottidæ*. (*analis*, from the large anal papilla.)

Oligocottus analis, GIRARD, Proc. Ac. Nat. Sci. Phila. 1857, 201, Monterey (Coll. A. S. Taylor. Type, No. 486, U. S. Nat. Mus.); GIRARD, U. S. Pac. R. R. Surv., x, Fishes, 57, 1858; JORDAN & GILBERT, Synopsis, 718, 1883.
Cottus criniger, GÜNTHER, Cat., II, 522, 1860, Monterey, California
Centridermichthys analis, GÜNTHER, Cat., II, 171, 1860.
Clinocottus analis, JORDAN & EVERMANN, Check-List Fishes North and Middle America, 444, 1896.

745. OLIGOCOTTUS, Girard.

Oligocottus, GIRARD, Proc. Ac. Nat. Sci. Phila. 1856, 133 (*maculosus*).

Body rather elongate, the skin smooth; preopercular spine slender and sharp, simple or with a single antler-like process above; small tentacles on head and anterior parts of body; gill membranes broadly united, free from the isthmus; slit behind fourth gill small or obsolete; dorsal spines slender, the fin short and not emarginate; anal papilla of male large; ventrals I, 3. Small fishes of the North Pacific, inhabiting rock pools between tide marks, the very smallest of the *Cottidæ*. (*ὀλίγος*, small; *Cottus*.)

a. Preopercular spine bifid, having a process directed upward in addition to the pointed spine; D. VIII, 16; A. 12 to 14.
  b. First dorsal without ocellate spot; head with many cirri; male with the anterior rays of anal enlarged and detached. MACULOSUS, 2381.
  bb. First dorsal with a black ocellus; cirri on head few; anal fin in males with the anterior rays not especially enlarged or detached. BOREALIS, 2382.

2381. OLIGOCOTTUS MACULOSUS, Girard.

(JOHNNY.)

Head 3; depth 4. D. VIII–16; A. 12 to 14. Head slender, narrowed above, the snout rather pointed; maxillary reaching pupil; top of head with several scattered cirri; a few on sides of head; cirri on lateral line anteriorly and on front of back, also along base of dorsals. No scales or prickles anywhere; preopercular spine forked; front rays of anal enlarged and detached in males; anal papilla large (in males); pectorals reaching well beyond front of anal. Usual color, reddish brown, varying to gray, intense green or crimson according to surroundings, the vivid colors developed in the presence of similarly colored algæ; fins all barred; belly usually livid bluish or greenish; lower side of head with white mottlings. Length 2 to 3 inches. Pacific coast, from Cape Mendocino to Point Concepcion, rather common; one of the smallest of the marine *Cottidæ*, a bright-colored inhabitant of rock pools. (*maculosus*, spotted.)

Oligocottus maculosus, GIRARD, Proc. Ac. Nat. Sci. Phila. 1856, 153, Tomales Bay (Coll. E. Samuels); San Francisco (Coll. Lieut. Trowbridge); Fort Steilacoom (Coll. Dr. G. Suckley); GIRARD, U. S. Pac. R. R. Surv., X, Fish., 56, 1858; JORDAN & GILBERT, Synopsis, 718, 1883.

Centridermichthys maculosus, GÜNTHER, Cat., II, 171, 1860.

2382. OLIGOCOTTUS BOREALIS, Jordan & Snyder.

Head $3\frac{1}{5}$; depth $4\frac{1}{5}$; orbit 4 in head; snout 4; maxillary $2\frac{4}{7}$. D. VIII or IX-16 or 17; A. 12 to 14; P. 14; highest dorsal spine 3; dorsal ray $2\frac{1}{5}$; caudal ray $1\frac{1}{2}$; first anal ray, male $1\frac{1}{2}$, female 3; ventrals $1\frac{5}{6}$; caudal peduncle $3\frac{1}{2}$; pectorals 3 in length. Body compressed, elongate; back somewhat elevated, deepest below middle of spinous dorsal. Head almost as wide as long, tapering from behind to the somewhat pointed snout; profile of head rounded above, straight below; mouth terminal, nearly horizontal; maxillary extending to vertical through posterior part of pupil; lower jaw included; jaws, vomer, and palatines with villiform teeth; snout as long as orbit; eye high in head; interorbital space narrower than width of orbit, its concavity angular; opercle with a triangular flap; angle of preopercle produced into a forked spine, which is covered with skin, except on the sharp points; prongs of preopercular spine $\frac{1}{2}$ as long as orbit; nasal spines prominent; the long premaxillary processes forming a sharp ridge between the latter; branchiostegal membranes forming a fold across the isthmus; gills $3\frac{1}{4}$, the slit behind the fourth arch much reduced; pseudobranchiæ present; gill rakers represented by a few protuberances on the arch. Skin smooth; filaments on free end of maxillary, on inferior edge of preopercle, and first dorsal to the bases of the pectoral fins; also a row of filaments extending along the supraorbital crest, over the back of the head and along the lateral line for about $\frac{1}{2}$ the length of the body, the filaments usually paired, i. e., 2 grow from the same place; anal papilla prominent; large mucous pores scattered about the top and sides of head; pores of lateral line 36 to 40. Dorsal fins 2, separate; first dorsal 4 in head and body, curving from distal end of first spine to posterior part of base; spines rather feeble; second dorsal 2 in head and body, a little higher than first, its middle rays longest; anal fin about $3\frac{1}{4}$ in head and body; in the male the first and second rays longest, the third, fourth, and fifth each a little shorter than the preceding one, the last shortest; in the female the first ray is the shortest; caudal fin somewhat rounded; pectoral rays below the sixth ventrally free from the connecting membrane for a portion of their length; ventral fins reaching to vent, in some specimens to anal. Color, usually reddish brown, varying to gray, intense green or crimson, according to surroundings, the colors developed in the presence of similarly colored algæ; dorsals, pectorals, and caudal barred; anal sometimes barred; front of spinous dorsal with an ocellated black spot. Pacific Coast, from Prince William Sound to Oregon; the types collected in tide pools at Neah Bay, Puget Sound, by Mr. E. C. Starks. Others were collected at Neah Bay by Mr. Henry St. Clair, and still others in Departure Bay, Vancouver Island, by Dr. C. H. Gilbert. This species is closely related to *Oligocottus maculosus*, but may be distinguished from the latter by having fewer filaments on the head and body, an ocellated spot

on front of first dorsal, and by having the rays of the anal fin in the male all connected by membrane; the first ray of anal is much shorter and weaker in *O. borealis*. (*borealis*, northern.)

Oligocottus borealis, JORDAN & SNYDER, Proc. Cal. Ac. Sci., Series 2, vol. VI, 1896, 225, Neah Bay. (Coll. E. C. Starks. Type, No. 3396. L. S. Jr. Univ. Mus.)

746. OXYCOTTUS, Jordan & Evermann, new genus.

Oxycottus, JORDAN & EVERMANN, new genus (*acuticeps*).

This genus is allied to *Oligocottus*, differing in the absence of upward process on the sharp, upwardly curved preopercular spine. No slit behind last gill. (ὀξύς, sharp; *Cottus*.)

a. Snout sharp; anal rays 12 or 13. ACUTICEPS, 2383.
aa. Snout bluntish; anal rays 10. EMBRYUM, 2384.

2383. OXYCOTTUS ACUTICEPS (Gilbert).

Head 3½ to 3⅔; eye 4 in head; snout 4. D. VII or VIII, 15 or 16; A. 12 or 13. Head slenderer and with sharper snout than in *O. maculosus*. Interorbital space slightly concave, its width ½ eye; maxillary region vertical just in front of the pupil, 3 in head. Cardiform teeth on jaws, vomer, and palatine bones. Preopercular spine slender, sharp, curved upward and inward, neither notched nor forked. Preopercular margin unarmed below it; nasal spines sharp; occiput without ridges or spines; opercle thickened above, ending behind in a rounded lobe; without definite ridge or spine. Branchiostegals 6; no evident pores behind last gill; gill membranes broadly united, free from the isthmus. A cirrus at inner base of nasal spines, 3 pairs evenly spaced on top of head, 1 above orbits, 1 posteriorly on occiput, and 1 midway between these 2; a cirrus at angle of opercle; 1 above each pore on anterior portion of lateral line; sides of body otherwise smooth, without further cirri and without axillary and other prickles. Dorsal fin usually slightly joined at base; pectorals reaching to or slightly beyond front of anal fin; ventrals short, equaling length of snout and eye, extending halfway to front of anal; anus anterior in position, thus differing from *O. maculosus* and *C. analis*, its distance from base of ventrals but ½ its distance from front of anal fin; pores of lateral line 33, each of the anterior 15 usually accompanied by a cirrus. Fin rays in 7 specimens are as follows:

	Spinous dorsal.		Soft dorsal.		Anal.	
Rays	VII.	VIII.	15.	16.	12.	13.
Specimens...	2	5	3	4	3	4

Color varying with the surroundings, often nearly uniform bright green. When dark markings are present, we usually find 6 short wedge-shaped dorsal bars, widening rapidly below, and joining one another by their extreme tips; below these a dusky wavy lengthwise streak, and another wider one below lateral line, the latter marked posteriorly by round white spots, the size of pupil; occiput dusky; a black bar from eye to snout; 1 from eye to behind maxillary; 1 from eye to base of preopercu-

lar spine; the interval between these subocular bars may be silvery white; spinous dorsal often showing 2 dark bars, as in *B. globiceps;* ventrals plain; fins otherwise finely mottled or indistinctly barred; some or all of these dark markings sometimes absent. Prince William Sound (Arthur White Greeley) to Vancouver Island. Four specimens, the largest 47 mm. long, from tide pools at Unalaska, are the types of the species. Three additional specimens were taken in Departure Bay, Vancouver Island. (Gilbert.) (*acutus,* sharp; *-ceps,* head.)

Oligocottus acuticeps, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 432, Unalaska (Coll. *Albatross*).

2384. OXYCOTTUS EMBRYUM (Jordan & Starks).

Head 4; depth $4\frac{1}{4}$. D. IX, 15; A. 10; orbit 4 in head; snout 4; maxillary $2\frac{5}{6}$; highest dorsal spine $2\frac{1}{2}$; dorsal ray $1\frac{3}{4}$; anal ray $1\frac{3}{4}$; length of caudal fin $1\frac{3}{5}$; ventrals $1\frac{1}{6}$; pectorals $2\frac{1}{3}$ in body. Body elongate, compressed, back slightly elevated, deepest under spinous dorsal; caudal peduncle moderately slender. Skin without scales or prickles; head small, tapering rapidly forward to the rather sharp snout as viewed from above; profile of head straight below, acutely and evenly rounded above; mouth terminal and horizontal; maxillary reaching to the vertical from the middle of pupil; lower jaw included; teeth on jaws, vomer, and palatines in narrow villiform bands; process of premaxillary prominent, extending slightly above nasal spines, giving the appearance of 3 spines above snout; eye set high in head, the orbit as long as snout; preopercular spine short, blunt, and triangular, entirely covered with the skin, edge of preopercle below, entire; opercle ending in a short flap; top of head with 2 rows of "mossy" cirri, running from the superior orbital margin, curving over head and continuing on lateral line, they disappear on its anterior third. Dorsal spine rather stout, the fin lower than soft dorsal, rounded in outline; soft dorsal well separated from spinous, the front of fin the highest; pectorals long, the eighth ray the longest, giving the fin a point behind which reaches to the base of about the seventh dorsal ray; the same rays below the eighth swollen and posteriorly free from the membrane; anal about as high as soft dorsal, the rays somewhat swollen and more or less free; ventrals long, reaching about to front of anal, their insertion behind base of pectoral a distance equal to the snout and eye; caudal fin slightly rounded. Color variable from light green to a rich maroon; traces of 5 or 6 dark cross bars on back; lower parts dusky with small light spots; belly white; a dark bar from eye to side of snout, 1 from eye to edge of preopercle behind end of maxillary, and another from eye to below preopercular spine; lips black; lower rays of pectorals crossed with black and white bars which fade out above; ventrals light with some dusky mottlings; dorsal dark above, light at base, no markings; anal with black and white bars running across the rays, caudal fin mottled. In this species the preopercular spine is very short, sharp, turned upward, and covered by skin. It has a more acute snout than *B. globiceps,* mouth with a lateral cleft, body more compressed, and the premaxillary process extending above nasal spines. Puget Sound to Mon-

terey, scarce; 2 specimens collected in the tide pools left in the sand on a beach near Neah Bay, the largest 2¾ inches in length. A third specimen, darker in color, obtained at Point Lobos, Carmelo Bay, near Monterey, by Mr. John O. Snyder. (*ἐν*, in; *βρύον*, sea moss.)

Oligocottus embryum, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 808, pl. 82, Neah Bay, Washington. (Type, No. 3128 L. S. Jr. Univ. Mus. Coll. E. C. Starks.)

747. BLENNICOTTUS, Gill.

Blennicottus, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 166 (*globiceps*).

This genus is very close to *Oligocottus*, differing in the form of its preopercular spine, which is short, blunt, and simple. Head short and blunt, the cleft of the mouth largely or wholly anterior. Last gill slit obsolete. North Pacific. (*Blennius*; *Cottus*.)

a. Head nearly globular, very blunt; the cleft of mouth not extending on its side; preopercular spine with a short point; ventrals 1⅛ to 1⅓ in head; D. IX, 16; A. 11.
b. Head with rather few cirri. GLOBICEPS, 2385.
bb. Head with very many cirri. var. BRYOSUS, 2385*a*.

2385. BLENNICOTTUS GLOBICEPS (Girard).

Head 3¾; depth 4½. D. IX, 16; A. 11; V. I, 3. Body little compressed; head extremely short and blunt, narrowed above, nearly everywhere convex. Mouth anterior, short and broad, almost without lateral cleft; lower jaw shortest; maxillary reaching past front of the small eye; interorbital space very narrow, grooved, about ½ width of eye; preopercular spine with a single point turned upward; top of head with 2 series of cirri; none on sides of head, except a few on upper part of opercle; a series of cirri along anterior half of lateral line; skin without scales or prickles; pectorals reaching beyond front of anal. Dark olive, with obscure darker cross bars; first dorsal with 2 dark streaks above; fins generally barred with greenish, orange, or blue; cirri mostly black. Length 3 to 4 inches. Pacific Coast, from Kadiak to San Diego; the typical form from San Diego northward to Oregon; in rock pools, rather rare, more common southward. (*globus*, globe; *-ceps*, head.)

Oligocottus globiceps, GIRARD, U. S. Pac. R. R. Surv., Fish., 58, 1858, South Farallones. (Coll. Lieut. Trowbridge. Type, No. 300, U. S. Nat. Mus.); JORDAN & GILBERT, Synopsis, 718, 1883.
Centridermichthys globiceps, GÜNTHER, Cat., II, 171, 1860.
Blennicottus globiceps, JORDAN & STARK, Proc. Cal. Ac. Sci. 1895, 808.

Represented in Puget Sound and northward to Kadiak by

2385*a*. BLENNICOTTUS GLOBICEPS BRYOSUS, Jordan & Starks.

Northern specimens are larger in size, 4 to 6 inches, with paler and more variegated coloration and red markings; the preopercular spine is stronger and slightly hooked upward, and the opercle has many more cirri above than in the southern form. These differences are very slight and perhaps unworthy of notice. (*βρύον*, sea moss, from the cirrous head.)

Blennicottus globiceps bryosus, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 808, Point Orchard, Seattle. (Coll. E. C. Starks.)

748. HISTIOCOTTUS, Gill.

Peropus, Lay & Bennett, Beechey's Voy., Zool., Fish., 59, 1839 (*bilobus*); name preoccupied.
Histiocottus, Gill, Proc. U. S. Nat. Mus. 1888, 573 (*bilobus*).

This genus is very close to *Blepsias*, from which it differs in the absence of smooth areas on the body; the fins are lower, the dorsal not emarginate, and the pectoral much larger, perhaps capable of being used for flight. North Pacific. (*ἱστίον*, sail; *Cottus*.)

2386. HISTIOCOTTUS BILOBUS (Cuvier & Valenciennes).

D. IX, 21; A. 18; P. 16. Body shorter and deeper than in *Blepsias cirrhosus*, with thicker caudal peduncle and heavier head, the bones less firm; snout short, obtuse, interorbital space very wide, concave, ½ wider than the small eye; short occipital ridges present, besides several bluntish tubercles. Mouth broad, oblique, the maxillary reaching pupil; teeth small, distant; barbels as in *cirrhosus*. Head and body covered with prickles, larger and blunter than in *cirrhosus;* no naked patches on body; fins less developed than in *cirrhosus;* first 5 or 6 spines of dorsal subequal, the last 3 abruptly shorter; pectorals longer and much broader than in *cirrhosus*, reaching seventh anal ray; caudal short, much shorter than head. Olivaceous, paler below; dorsal region with 4 or 5 black bars, reaching ⅓ the distance to the lateral line, and somewhat continued on the fins; caudal with a black bar at base, otherwise plain; pectorals and anal blotched with black. Coast of Alaska and Kamchatka, not very common; the specimen here described from Kadiak. (*bilobus*, two-lobed.)

Blepsias bilobus, Cuvier & Valenciennes, Hist. Nat. Poiss., iv, 379, 1829, Kamchatka (Coll. M. Collée, Brit. Mus.); Günther, Cat., ii, 153, 1860; Jordan & Gilbert, Synopsis, 720, 1883.
Peropus bilobus, Bennett, Beechey's Voy., Zool., Fish., 59, 1839.

749. BLEPSIAS, Cuvier.

Blepsias, Cuvier, Règne Animal, Ed. 2, 1829 (*cirrhosus*).

Head and body compressed; skin hispid with stiffish villiform prickles, and with definitely naked areas; snout and chin with several rather long barbels; mouth small; teeth villiform, on jaws, vomer, and palatines; preopercle with 2 short blunt spines; gill membranes free from the isthmus; gills 4, a slit behind the fourth; top of head with bony ridges; suborbital stay narrow; first dorsal short, elevated in front, emarginate, the spines slender; second dorsal large; anal similar, lower; ventrals very short, I, 3; pectorals long. North Pacific. (An old name of some fish; from *βλέπω*, look.)

2387. BLEPSIAS CIRRHOSUS (Pallas).

Head 3¾; depth 3¾. D. V–III, 23; A. 20; P. 12; scales 50. Supraocular ridges well developed; interorbital space concave, with 2 low ridges extending backward from nasal spines, diverging toward occipital ridges; the latter large, partly interrupted; suborbital stay conspicuous; tempo-

ral ridge present; maxillary reaching to beyond middle of pupil; snout with 5 barbels, chin with 6, the longest about as long as the eye, which is $3\frac{2}{3}$ in head; a small cirrus on each interorbital ridge; interorbital space as wide as eye; sides with 3 or 4 pale, well-defined, naked areas behind pectorals, the 2 anterior much the largest; behind these a long naked strip along the lateral line, colored like the rest of the body; first 4 spines of dorsal elevated, $\frac{2}{3}$ to $\frac{3}{4}$ length of head; the fifth much shortened; membrane deeply notched between the fifth spine and the sixth, which is longer than the one before it; ventrals as long as eye; pectorals reaching much beyond front of anal; first rays of soft dorsal short; the others gradually lengthened to near the last, the highest higher than the dorsal spines; caudal longer than head. Dark olivaceous, sides of back with 4 to 6 vertically oblong black blotches edged with paler, not reaching nearly to lateral line; belly and naked areas on sides whitish; black bands radiating from eyes; fins dark, with large pale blotches and dark spots; caudal with light and dark bars; ventrals plain; a small white spot on front of spinous dorsal. Adults from Unalaska show the following color: Olive green of varying shades, the belly bright coppery yellow, the cross blotches on back nearly black, with paler margins; naked patches on sides white or brassy, those on tail colored like body, those on head silvery; first dorsal light olive, with 2 translucent patches; second dorsal mottled olive with dark spots and translucent patches; caudal with 3 blackish and 4 translucent bands; anal yellowish olive with numerous spots and translucent patches; radiating blackish bands running out from eye; the upper barbels black, the lower olive. Length 6 inches. Alaska to San Francisco, in shallow water; not rare northward; common at Unalaska; known also from Petropaulski and Iturup Island; not often seen south of Puget Sound. (*cirrhosus*, bearing cirri.)

Trachinus cirrhosus, PALLAS, Zoogr. Ross.-Asiat., III, 237, 1811, Avatcha Bay, Kamchatka (Coll. Merk); Gulf of Penshin, Ochotsk Sea.

Blepsias trilobus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 375, 1829; same types.

Blepsias cirrhosus, GÜNTHER, Cat., II, 153, 1860; STEINDACHNER, Ichth. Beit., V, 128, 1876; JORDAN & GILBERT, Synopsis, 719, 1883.

750. NAUTISCUS, Jordan & Evermann, new genus.

Nautiscus, JORDAN & EVERMANN, new genus (*pribilovius*).

This genus is closely allied to *Nautichthys*, having the same gill structures and dentition, but the first dorsal is only moderately elevated, its height being less than depth of body. There are no cirri on the head, and there is no marked depression at the occiput, the base of the dorsal being little raised above the nape; preopercle with four obsolete spines; skin rougher than in *Nautichthys*, the anal fin shorter. Bering Sea. (a diminutive of ναυτής, a ship, from the low first dorsal.)

2388. NAUTISCUS PRIBILOVIUS, Jordan & Gilbert, new species.

Differing from *N. oculofasciatus* in the shorter lower fins, the lower cranial ridges, and the coloration. Head 3 to $3\frac{1}{3}$; depth $3\frac{2}{3}$. D. VIII, 23; A. 15; V. I, 3; P. 15; eye $3\frac{1}{4}$ in head; maxillary $2\frac{2}{3}$; lateral line with 39 spines.

Shape of head and body much as in *N. oculofasciatus;* head short, the snout rather sharp, the anterior profile steep; nasal spines prominent, a short occular cirrus, much smaller than in *N. oculofasciatus,* shorter than pupil; interobital space narrow, about ½ eye, deeply concave; a blunt triangular ridge above each orbit, with a deep cross furrow behind it which deepens to a pit at the vertex; nuchal ridges lower than in *N. oculofasciatus,* each with a coarse tubercle, lower and larger than in the other species. From the nuchal depression, the base of first dorsal spines rises much less abruptly. Preopercle with 4 blunt prominences, the upper often longer and more spine-like. Mouth nearly horizontal, the lower jaw included; a slender filament at end of maxillary; teeth small, a few on vomer and a narrow band on palatines. Gill membranes broadly united to isthmus, the gill opening extending a little below the lower edge of pectoral. Skin covered with close-set villous prickles, among which large ones are frequently seen arranged in rather definite longitudinal series, of which there may be 2 or 3 parallel with the back, and 1 running near lower line of tail. No smooth areas on sides. Lateral line conspicuous, the plates with short spines directed backward. Dorsals separate, the first not notched, comparatively low, the first spine highest, 1⅔ in head in type; in other specimens 1¼ to 1⅓ in head; soft dorsal and anal also low, none of the rays reaching base of caudal when depressed; pectoral longer than head; ventrals 1½ to 2 in head. Color, dull light olivaceous, mottled with darker; 3 or 4 dark bands below soft dorsal, 1 below spinous dorsal; a black band through eyes and across cheeks, extending onto branchiostegal membranes; 7 dusky spots along lateral line, a conspicuous pink blotch, rather larger than pupil, between first and second blotch; first dorsal dusky; second dorsal, anal, and pectoral dotted and checked; caudal with faint finely checked cross lines which deepen to form a dark bar at its base and a broader one toward its tip; ventrals pale; belly mottled. Bering Sea. One specimen, 6 cm. long, from *Albatross* Station 3635, off St. George Island, in 23 fathoms. Another specimen barely an inch long was dredged in 7 fathoms in the harbor of Unalaska. Very numerous specimens were obtained by the *Albatross* in 1890 in Bristol Bay and south of the Alaskan Peninsula. It was at that time incorrectly identified by Dr. Gilbert with *N. oculofasciatus.* In 5 specimens of those from Bristol Bay the dorsal contains VIII or IX, 23 or 24 rays, the anal 16 or 17, the pectoral 15 or 16. (*pribilovius,* from the Pribilof Islands, named for their discoverer, Gerassim Pribilof, 1786.)

Nautichthys oculofasciatus, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 434; not of GIRARD.
Nautichthys pribilovius, JORDAN & GILBERT, Fishes Bering Sea, MS., 1898, off St. George Island. (Type, No. 48237, U. S. Nat. Mus. Coll. Dr. Jordan.)

751. NAUTICHTHYS, Girard.

Nautichthys, GIRARD, U. S. Pac. R. R. Surv., x, Fishes, 74, 1858 (*oculofasciatus*).

Body rather elongate, compressed, but not elevated, the skin evenly covered with short, close-set, villiform prickles. Head short, strongly compressed, the cheeks subvertical; orbital ring much elevated above, with

several blunt protuberances behind; nape much depressed, with the base of the dorsal fin rising abruptly above it; mouth small; teeth on jaws, vomer, and palatines; preopercle with small, bluntish spines above; gill membranes broadly united to the isthmus; a slit behind fourth gill; branchiostegal 6; first dorsal falcate, the spines very flexible, the anterior much elevated, much longer than head, the posterior rapidly shortened; second dorsal long and rather high; anal fin similar to second dorsal, but shorter and lower; pectorals moderate; ventrals I, 3, well developed, nearly as long as head. Pacific Coast. Small fishes, of singular appearance. (*ναύτης*, sailor; *ἰχθύς*, fish; in allusion to the development of the first dorsal fin.)

2389. NAUTICHTHYS OCULOFASCIATUS (Girard).

Head 3½; depth 3¾. D. IX, 28; A. 19; P. 14; V. I, 3; scales 44. Maxillary reaching to opposite pupil; eye very large, longer than snout; nasal spines conspicuous, curved; maxillary and edge of preopercle with minute cirri; orbit with a branched cirrus, as long as pupil; the narrow interorbital space deeply channeled; supraocular ridge posteriorly with 3 blunt tubercles; 2 compressed tubercles on each side of first dorsal; a deep pit at nape, extending under origin of first dorsal; spinous dorsal smooth; rays of soft dorsal and caudal roughened with prickles; dorsal spines nearly twice length of head; ventrals ¾ length of head, a little shorter than caudal; pectorals long. Grayish above; sides with dark marblings and obscure dusky bands; a very conspicuous black band through eye and across cheek; first dorsal blackish; other fins barred with light and dark; ventrals plain. Length 6 inches. Pacific Coast, Kadiak to San Francisco, chiefly northward; not rare in Puget Sound; a most singular-looking fish. (*oculus*, eye; *fasciatus*, banded.)

Blepsias oculofasciatus, GIRARD, Proc. Ac. Nat. Sci. Phila. 1857, 202. Fort Steilacoom, Washington. (Coll. Dr. Geo. Suckley. Type, No. 512, U. S. Nat. Mus.)

Nautichthys oculofasciatus, GIRARD, U. S. Pac. R. R. Surv., x, Fish., 74, 1858; GÜNTHER, Cat., II, 157, 1860; STEINDACHNER, Ichth. Beitr., v, 130, with plate; JORDAN & GILBERT, Synopsis, 727, 1883; JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 810.

752. ULCA, Jordan & Evermann.

Ulca, JORDAN & EVERMANN, Proc. Cal. Ac. Sci. 1896, 227 (*marmoratus*).

The genus is close to *Hemitripterus*, from which it differs in its shorter first dorsal of about 14 spines, the first 4 of which are not differentiated, the fin therefore not emarginate. Alaska. (*Ulke*, the Danish-Norwegian name for all *Cottidæ*.)

2390. ULCA MARMORATA (Bean).

Head 2⅔; depth 3⅔. D. XIV, 12; A. 13; V. I, 3; lateral line (tubes) 44. Eye about as long as snout and ¼ as long as head. Pectorals reaching to the vertical through the anal origin; ventral scarcely longer than eye. Gill membrane free from isthmus. Maxillary reaching beyond end of eye. First dorsal base as long as head without snout, the first 4 spines less

elevated and differentiated than in *Hemitripterus*. Sides dark gray, intermingled with whitish reticulations. Type 2¾ inches long. North Pacific and Bering Sea, in deep water. (Bean). Dr. Gilbert notes also: "Several small specimens were secured in Bering Sea, to the north of Unalaska Island, depths 70 to 121 fathoms. They agree with the types in having but 14 dorsal spines, the first 4 of which are not noticeably differentiated. The second dorsal contains 11 or 12 rays, and the anal fin 13. The last 2 rays of the anal fin are approximated at base, but do not evidently constitute a divided ray." (*marmoratus*, marbled.)

Hemitripterus marmoratus, BEAN, Proc. U. S. Nat. Mus. 1890, 43, off Sitkalidak Island, at Albatross Station 2855 in 69 fathoms; GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 410.

? *Cottus villosus*, * PALLAS, Zoogr. Ross.-Asiat., III, 129, 1811, Cape of Kronok and mouth of the Itscha; after STELLER; CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 196, 1829.

753. HEMITRIPTERUS, Cuvier.

(SEA RAVENS.)

Hemitripterus, CUVIER, Règne. Anim., Ed. 2, vol. 2, 164, 1829 (*americanus*).

Body moderately elongate, scaleless, but the skin covered with prickles and bony protuberances of various sizes and forms. Head large, with numerous bony humps and ridges and fleshy slips above; orbital rim much elevated, the interorbital space deeply concave; a depressed area at the occiput, behind which are 2 blunt spines on each side. Mouth very wide; jaws, vomer, and palatines with broad bands of teeth; no slit behind last gill; gill membranes broadly united, free from isthmus; preopercle with stout, blunt spines; suborbital stay very strong, forming a sharp ridge. Spinous dorsal much longer than the soft part, of 16 to 18 spines, of which the first 2 are the highest, and the fourth and fifth shorter than the succeeding ones, the fin thus deeply emarginate; pectoral fins very broad, much procurrent; ventrals I, 3. Large fishes of singular appearance, inhabiting the North Atlantic and Pacific. Dr. Gill makes of them a distinct family on account of the great length of the spinous dorsal and the peculiar development of the myodome. The genus is, however, related to *Blepsias* and *Nautichthys*, and the spinous dorsal is as long in *Jordania* as in *Hemitripterus*, while the two genera stand as extremes in the Cottoid group. (ἡμι—, half; τρεῖς, three; πτερόν fin.)

a. Dorsal spine about 16; the upper surface of head moderately uneven. AMERICANUS, 2391.

aa. Dorsal spines 18; top of head excessively uneven. CAVIFRONS, 2392.

* Some unknown fish allied to *Ulca marmorata* is *Cottus villosus*, Pallas. Size, form, and dimensions of *Oncocottus quadricornis*, the same intestines, but with a disagreeable smoke-like odor. Its skin is soft, loose, sand-colored, covered with villosities like a calf's tongue. On the lateral line these prickles are stronger and ½ of a line to a line in diameter. Eight soft cutaneous flaps each 3 lines long, and bifid or trifid, at equal intervals on the lower jaw; belly white, back varied and marked with brown. Cape of Kronok and mouth of the river Itscha. Known only from notes of Steller. (Pallas). (*villosus*, hairy.)

2391. HEMITRIPTERUS AMERICANUS (Gmelin).

(Sea Raven.)

Head $2\frac{2}{3}$; depth $3\frac{3}{4}$. D. IV, XII-I, 12; A. 13; scales 40. Body villous, the prickles enlarged and tubercle-like along back and lateral line; nasal spines strong; supraocular ridge much elevated, with dermal flaps and 2 blunt spines; 3 pairs of fleshy slips on nasal bones, and 2 on supraocular ridges; smaller cirri on maxillary, on preorbital, and several on lower jaw; interocular space very deeply concave; 2 blunt occipital spines on each side, outside of which are 2 or 3 others; opercle small, with a bony ridge; preopercle with 2 blunt spines, below which are 1 or 2 others; lower jaw slightly projecting; maxillary reaching beyond eye, about $\frac{1}{2}$ head; pectorals nearly reaching anal; highest dorsal spine $1\frac{2}{3}$ in length of head, as long as caudal; ventrals reaching halfway to anal. Reddish brown, marbled with darker brown, and much variegated; yellowish below; fins variegated with light and dark. Length 8 inches. Atlantic coast of America, chiefly northward, from New York to Labrador, not rare, common at Woods Hole in October and November; a most remarkable looking fish.

Scorpæna americana. Gmelin, Syst. Nat., 1220, 1788, after "Diable en crapand de Mer d'Amérique" of du Hamel de Monceau, Pêche, III, 2, 93.
Cottus acadian, Pennant, Arctic Zool., III, 371, **Nova Scotia**.
Cottus acadian, Walbaum, Artedi Pisc., 392, 1792, **Nova Scotia**; **Newfoundland**.
Cottus hispidus, Bloch & Schneider, Syst. Ichth., 63, 1801, **New York**.
Cottus tripterygius, Bloch & Schneider, Syst. Ichth., 63, 1801, **Nova Scotia**; after Pennant.
Hemitripterus americanus, Cuvier & Valenciennes, Hist. Nat. Poiss., IV, 268, 1829; Günther, Cat., II, 143, 1860; Jordan & Gilbert, Synopsis, 685, 1883.
Hemitripterus acadianus, Storer, Hist. Fish. Mass., 35, 1867.

2392. HEMITRIPTERUS CAVIFRONS, Lockington.

D. IV, XIV, 13; A. 14; P. 20; V. I, 3; scales 44. Very similar to *Hemitripterus americanus*, but with the upper surface of the head more uneven, the concavity of the interocular area more marked, and the dorsal fin with more spines. Head equaling depth, about $3\frac{1}{2}$ in total length with caudal. Length 16 inches. Coast of Alaska, not common; perhaps not distinct from the preceding. (*cavus*, concave; *frons*, forehead.)

Hemitripterus cavifrons, Lockington, Proc. Ac. Nat. Sci. Phila. 1880, 233, **Kadiak Island, Alaska**. (Coll. W. J. Fisher.)

754. SYNCHIRUS, Bean.

Synchirus, Bean, Proc. U. S. Nat. Mus., XII, 1889, 641 (*gilli*).

Body slender and moderately elongate, resembling that of *Triglops;* covered with thin, tough skin. Lateral line armed with spiny tubercles. Spiny scales in a series along the dorsal base. Head subconical, with moderately pointed snout. Mouth small, very slightly oblique; the rami of the mandible a little concave beneath. Premaxillaries protractile;

jaws with slender, villiform teeth, in bands; teeth on vomer and palatines. Pseudobranchiæ present. Gills 3½, no slit behind the last; gill openings wide, extending above the median line, the membrane free from the isthmus. Suborbital connected by a bony stay with the preopercle, which bears a strong bifid spine at its angle. Pectorals completely united around the breast, the membranes free at the margin. Ventrals distant from the gill opening, the pubic bones being remarkably long, the fins diverging widely, and consisting of a rudimentary spine and 3 rays. Dorsal long, the spinous portion low, with slender spines, and the soft portion twice as long as the spinous. Anal long. Caudal moderately elongate, its middle rays somewhat produced. Genital papilla of male received in a pit in front of anal. Pacific. (*σύν*, together; *χείρ*, hand.)

2393. SYNCHIRUS GILLI, Bean.

Head 3½; depth 5½; Br. 6; D. VIII to IX, 19 to 21; A. 20; V. I, 3; P. 22. Eye about as long as snout, 4 in head. Maxillary extending to about below middle of eye. The interorbital space not quite equal to length of eye; a pair of strong nasal spines; preopercle with a short and very sharp bifid spine; lateral line with about 41 spiny tubercles, and most of the specimens have a single series of spiny scales along the dorsal base; pectorals nearly as long as the head, and extending to about below the fourth ray of the soft dorsal; ventrals nearly under the middle of the pectorals, their length varying greatly; in some specimens scarcely ½ as long as the head; in others as long as postorbital part of head. In some males the anal papilla is ⅔ as long as the ventral fin of the same individual. This papilla can be received into a pit in front of the anal fin. Spinous dorsal beginning over axil of pectoral, length of its base a little greater than postorbital part of head; none of its spines much longer than eye; distance of anal origin from head about ⅔ length of head. Rays of soft dorsal and anal not much longer than dorsal spines. Caudal about ⅔ as long as head, its middle rays somewhat the longest. Color in spirits pale yellowish brown; the sides showing traces of several small, pale blotches, and the caudal and pectoral with a few very small dark blotches, those on the caudal forming interrupted bands; across the back are faint indications of about 5 pale cross bands. Length 2 inches. Barclay Sound, British Columbia. (The species is dedicated to Dr. Theodore Gill, in appreciation of his researches upon the mail-cheeked fishes.)

Synchirus gilli, Bean, Proc. U. S. Nat. Mus., XII, 1889, 642, Barclay Sound, British Columbia. (Type, No. 41820. Coll. *Albatross*.)

755. ASCELICHTHYS, Jordan & Gilbert.

Ascelichthys, Jordan & Gilbert, Proc. U. S. Nat. Mus. 1880, 264 (*rhodorus*).

Body plump, fusiform, tapering backward, covered with loose smooth skin; head broad, depressed, its upper surface evenly and weakly convex; a hooked preopercular spine; no nasal spines; a supraorbital cirrus; teeth on jaws, vomer, and palatines; no slit behind fourth gill; gill membranes broadly united, free from the isthmus; spinous dorsal low; the spines

slender, almost concealed in the membrane; pectorals broad; ventral fins wanting. Small fishes of the rock pools of the North Pacific. (ἀ-, privative; σκέλος, leg; ἰχθύς, fish.)

2394. ASCELICHTHYS RHODORUS, Jordan & Gilbert.

Head 3; depth 5. D. IX, 19; A. 15; P. 16. Head low, rounded anteriorly; maxillary reaching to posterior border of eye. No scales, prickles, or barbels anywhere, except a fringed cirrus over the eye; preopercle with a short, simple, strongly hooked spine; spines directed downward and forward on subopercle and interopercle; lateral line continuous. Dorsal fins connected by membrane, the spines low, weak, nearly uniform; soft dorsal nearly twice as high as spinous; pectorals about reaching vent. Dark olivaceous, sometimes with saddle-like pale blotches; lips rosy; spinous dorsal dusky, edged in life with bright crimson; other fins dusky, edged with paler, the pectoral slightly barred. Length 15 inches. Pacific Coast, Sitka to Cape Mendocino; very abundant among rocks between tide marks at Neah Bay, the only locality where numbers of specimens have been found, one of the most remarkable of the *Cottidæ*. (ῥόδον, rose; ὅρος, margin.)

Ascelichthys rhodorus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 264, Waadda Island, Neah Bay, near Cape Flattery (Coll. Jordan & Gilbert); JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 811, pl. 84.

756. PSYCHROLUTES, Günther.

(SPINELESS SCULPINS.)

Psychrolutes, GÜNTHER, Cat., III, 516, 1861 (*paradoxus*).

Body tadpole-shaped, tapering from the head to the very slender tail, covered with very loose, naked, movable skin. Head large, depressed, flattish above; snout obtuse, rounded; interocular space very broad, the ocular ridges obsolete; mouth very large; mandible short, little cavernous, its forms broadly U-shaped; maxillary entirely adnate to the skin of the preorbital; jaws with bands of villiform teeth; no teeth on vomer or palatines; no spines or cirri about the head; suborbital stay narrow. Gill membranes united to the isthmus; gills 3½, no slit behind the fourth. Branchiostegals 7. Fins connected; spinous dorsal of short, slender, flexible spines entirely embedded in the skin and not visible without dissection as the spines do not rise above level of the muscles; soft dorsal short, high, the rays close together, the total number 12 to 24; anal low, of 9 rays; caudal separate; pectoral fins long, with a broad, procurrent base. Ventrals I, 3, close together, distinct, the inner edge adnate to the body. Small fishes, very closely allied to *Cottunculus* and *Malacocottus* on the one hand and to the *Liparididæ* on the other, their characters, like those of the latter family, arising from degeneration of the *Cottidæ*. The extension of the lax skin over the spinous dorsal and the bones of the head afford the only tangible diagnostic character of the subfamily

Psychrolutinæ. From the *Liparididæ,* their separate ventrals distinguish them sufficiently. Small shore fishes of the North Pacific; only a single species known. (ψυχρολουτης, one who bathes in cold water.)

2395. PSYCHROLUTES PARADOXUS, Günther.

Head 3⅜; depth 4½, equal to head without snout. D. IX, 15 to 17, only 12 rays seen without dissection; A. 12 or 13; V. I, 3; branchiostegals 7. Head large, very broad, tapering suddenly to caudal peduncle. Eye 4¼ in head, a little shorter than snout, a little less than width of interorbital space. Maxillary extending to below middle of eye. Interocular space flat, not concave; jaws equal, the lower ⌒-shaped, its arch not narrowed behind. Pectoral reaching to slightly above origin of anal, 2⅜ in body; ventrals very small, ⅓ length head. Vent about midway between ventral and anal origin, the second dorsal beginning nearly over tip of pectoral; first dorsal entirely buried in the skin, its short, stiffish spines to be found only by dissection. Color light chocolate brown above, with minute spots of dark; whitish below; a pale stripe from eye downward and forward, between 2 dark ones; body with 3 very irregular dark cross bands, the third extending on dorsal and anal, the first 2 largely confluent and all very irregular in form; an oblique dark band on base of caudal, a narrow dark band behind it; pectoral with 2 or 3 cross bands; all fins, except the ventral, with traces of bands. Length 2½ inches. A specimen, 50 mm long, taken by the *Albatross* off St. Paul Island, Bering Sea, in 1896, shows the following characters: Head 2⅔; depth 3. D. IX, 15; A. 13; P. 19; eye 4 in head; width of mouth, from angle to angle, 1½; snout 3⅛; interorbital 3⅕. Body short, broad, thick, tadpole shape, the texture soft like that of a Liparid, especially about the head; the skin is limp and smooth, covered with little soft dermal warts, that of head especially lax, the cheeks tumid and translucent. No trace of spines on head, the bones all thin and weak; nostrils each in a short tube; mouth broad, its cleft chiefly anterior, the jaws equal; teeth very minute, none on vomer or palatines; lower jaw with 8 large open pores. Gill membranes broadly united to the isthmus, the gill opening extending to slightly below base of pectoral. Lateral line obsolete. Dorsals united, with a slight notch between, the first buried in a ridge of skin so that its delicate spines can not be counted from without; second dorsal low, similar to anal, both of them free from the caudal; lower pectoral rays progressively shortened, the longest 1⅓ in head; ventrals moderate, I, 3, reaching vent, 2⅓ in head; caudal rounded. Color creamy white, with blackish cross bands, irregular in form and broken by whitish patches; 5 black spots on lower jaw; top of head blackish; a narrow blotch at shoulder; a wider one across first dorsal; a broad one on second dorsal abruptly broadened on body, then narrowed, extending across anal; an irregular bar at base of caudal; a narrow bar and some spots and streaks on the fin; pectoral with 2 curved bars, the inner concave, the outer convex backward, the two inclosing a rounded pinkish or deep orange area. Alaska to Puget Sound, rather common in water of moderate depth; a remarkable little fish, evidently a degraded Cottoid. Here described

from specimens from Unalaska. This species is taken abundantly in shallow water about Kadiak south of the Alaskan Peninsula, thence westward through Unimak Pass, along the northern shore of Unalaska Island to the Kuril Islands, and the Pribilof Islands and in Bristol Bay. The depths range from 38 to 121 fathoms. (παράδοξος, unaccountable.)

Psychrolutes paradoxus,* GÜNTHER, Cat., III, 516, 1861, Gulf of Georgia, Vancouver Island (Coll. H. M. S. *Plumper*); JORDAN & GILBERT Synopsis, 687, 1883.

Psychrolutes zebra, BEAN, Proc. U. S. Nat. Mus. 1890, 43, Aleutian Islands, between Unga and Nagai Islands, at Albatross Station 2848, 55° 10′ N., 160° 18′ W., in 110 fathoms; JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 811, pl. 85; GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 410.

757. GILBERTINA, Jordan & Starks.

Gilbertina, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 810 (*sigolutes*).

This genus differs from *Psychrolutes* chiefly in the long continuous dorsal and anal, the former having 24 rays, the latter 14. Anterior rays of spinous dorsal not bound down by skin of the body, the first apparent ray being nearly above gill opening. Skull extremely soft and cavernous, the long oblique lower jaw especially so. (Named for Dr. Charles Henry Gilbert, "who has contributed more than anyone else to our knowledge of the fishes of the North Pacific.")

* The following is the original description of *Psychrolutes paradoxus*, Günther:

"Branchiostegals 7; D. 9; A. 9; C. 12; V. 2. Head large, not quite so high as broad, its length being ⅔ of the total. The snout is obtuse, although considerably longer than the eye, rounded, with the upper profile steeper than the lower. The cleft of the mouth is oblique, of moderate width with the jaws equal anteriorly; a narrow band of minute teeth in the jaws; the palate appears to be smooth. The maxillary, which is widened at its posterior extremity, extends to below the middle of the eye. The nostril is situated very near to the extremity of the snout, and provided with a very short tentacle. The eye is immediately below the upper profile; its diameter is about ⅓ of the length of the head, ⅓ of that of the snout, and nearly ⅓ of the width of the interorbital space. The infraorbital emits a slender process across the cheek, which, however, does not reach to the preopercular margin. None of the opercles are armed; the operculum alone is produced into a flexible short posterior process; the gill opening commences above that process, and is not continuous with that of the other side, the gill membranes being attached to the isthmus. The branchiostegals are exceedingly slender, 7 in number. The body is subcylindrical anteriorly and compressed posteriorly; its greatest depth, behind the head, is ⅛ of the total length; the skin is perfectly smooth and rather loose. The pectoral is pointed, the upper rays being the longest and extending nearly to the dorsal; the ventrals rather short, close together, situated below the middle of the base of the pectoral, and composed of 2 rays, the inner of which is bifid. Dorsal and anal fins opposite each other, situated far backward on the tail, terminating at some distance from the caudal, and nearly entirely enveloped in skin. Caudal slightly rounded, of moderate length. The vent is situated about midway between the root of the ventral and the origin of the anal. The upper parts are brownish gray, minutely dotted with black, and with 2 very large deep brown blotches of irregular size. The caudal and pectoral fins are dotted and spotted with black.

	Lines.
Total length	21
Length of head	6
Heighth of the body	4

"This fish resembles in its general habit somewhat the genius *Liparis*. Gulf of Georgia (Vancouver Island), voyage of H. M. S. *Plumper*."

Dr. Boulenger says, October 24, 1895: "I have examined the type of *Psychrolutes paradoxus*. I count 12 soft rays, the anterior embedded in the thick skin, so that Dr. Günther's statement is to be accounted for. But there are no spines." Later Dr. Boulenger writes us that he has compared Dr. Günther's type with the figure of *Psychrolutes zebra*, published by Jordan & Starks, and finds that it agrees fully with the latter. There is, therefore, little doubt that the species *zebra* is identical with *paradoxus*.

2396. GILBERTINA SIGOLUTES, Jordan & Starks.

Head 3; depth 4. D. VIII, 18; A. 14; eye $4\frac{1}{2}$ in head; interorbital $2\frac{1}{2}$; maxillary $2\frac{1}{5}$; ventrals 2; pectorals 1; caudal $2\frac{1}{4}$; base of dorsal $1\frac{3}{5}$ in length of body; base of anal 3. Body rather slender, robust anteriorly, compressed posteriorly, the greatest breadth and depth at shoulders. Head large, the nape slightly produced; mouth large and broadly rounded, oblique, the jaws about equal, maxillary extending to posterior margin of eye, its end buried under the skin of the cheek; eyes placed high, the interorbital space very wide and slightly convex, its width about $1\frac{1}{2}$ times that of the eye; the posterior end of mandible very prominent, mandible ∩-shaped, its rami approaching each other posteriorly; bones of head cavernous, largely made up of cartilage, the mandible especially so; anterior end of preorbital forming a blunt spine over mouth; process of premaxillary prominent; a couple of blunt projections behind each eye; upper part of shoulder girdle projecting, forming a blunt spine on nape above gill slit, a row of large pores around suborbital ring, and along under part of mandible no opercular spines. Head and body covered with a very loose, naked, movable skin. Dorsal fin continuous, no notch between spines and soft rays, the spines very slender, the first one inserted over end of opercular flap; dorsal spines covered by skin, but even the first visible without dissection as they rise above the general integument of body, the last reaching to base of caudal fin; anal lower than dorsal, its origin midway between base of caudal fin and posterior margin of eye, ending at about the same vertical that dorsal does, but not reaching so far; pectorals long and slender, reaching past front of anal and over halfway between their bases and base of caudal fin, adnate to the body for the anterior third or fourth of their length; ventrals long, not quite reaching to vent, adnate to the body for $\frac{1}{2}$ their length; caudal fin rounded. Color, light olivaceous; body and head with innumerable dark points, giving the fish a dusky appearance; large dark blotch across body at the posterior ends of dorsal and anal, a similar spot under pectoral; head uniform dusky, lighter below; belly white; middle of pectoral dark; dorsals dark; lower fins white. Puget Sound; known from a single small specimen $1\frac{1}{4}$ inches in length. (*σιγή*, quiet; *λούτης*, bather.)

Gilbertina sigolutes,* Jordan & Starks, Proc. Cal. Ac. Sci. 1895, 811, pl. 86, Point Orchard, near Seattle. (Type, No. 3129, L. S. Jr. Univ. Mus. Coll. Edwin C. Starks.)

Note.—Remotely allied to the *Cottidæ* is the Old World family *Platycephalidæ*, characterized by the greatly depressed head and body. Its principal genus is characterized as follows:

PLATYCEPHALUS, Bloch & Schneider.

Platycephalus, Bloch & Schneider, Syst. Ichth., 58, 1801 (*scaber*).

Head broad, very depressed, more or less armed with spines; body depressed anteriorly, subcylindrical posteriorly, covered with ctenoid scales. Lateral line present. Two dorsal fins, the first spine isolated from the others. Ventrals thoracic, but rather remote from the root of the pectorals; no pectoral appendages. Jaws, vomer, and palatine bones with bands of villiform teeth. Air bladder none; pyloric appendages in moderate numbers. Polynesia and the Australian seas, through all the Indian seas to the eastern coast of Africa. (Günther.) Two species of this genus have been described from American waters, in both cases apparently by error, as no American ichthyologist has found any specimens of either. (*πλατύς*, broad; *κεφαλή*, head.)

*Misprinted *sigalutes* in the orignal description.

The following are the species in question:

Platycephalus americanus, SAUVAGE, Nouv. Archiv. Mus., 148, pl. 14, fig. 3, 1878, Potomac River.

This specimen, of course, never came from the Potomac River. The following is the substance of Sauvage's description: Head nearly 3½; width of head nearly 2 in its length. No spines at end of snout; a strong spine before eye; rim of orbit armed with spines directed backward, those in front smaller; ridges of vertex slightly prominent, with few spines; cheek with 4 or 5 strong spines; edge of opercle soft; preopercle with 2 spines, the upper a little the longer, about half space between it and eye; maxillary reaching middle of eye; snout 2 in eye; lateral line spiny throughout its whole length; second dorsal spine shorter than third, as long as maxillary. Ventrals inserted far from anal; pectoral a little shorter than snout and eye. Body reddish, deep red at caudal peduncle; second dorsal and anal with 2 oblique bands of yellowish; pectoral with similar band.

Platycephalus angustus, STEINDACHNER, Sitzgber. Akad. Wiss. Wien 1866, 213, taf. 1, fig. 4, Surinam.

This specimen may not have come from Surinam. The following is the substance of Steindachner's description: Head 3⅔; breadth of head 1⅛ in its length. D. I–VIII, 13; A. 13; scales 108. Eye 7, somewhat greater than breadth of forehead. Caudal with 3 deep, blackish-brown, lengthwise stripes; body with small spots and some larger light-brown spots; 3 brown cross bands on posterior half of body. Numerous little raised ridges on preorbital rim, and a short spine; no tentacles; 2 spines on preopercle, the lower ⅔ eye. Greatest height of first dorsal ⅓ of second or $\frac{6}{13}$ of head. Caudal rounded.

Family CLXXX. RHAMPHOCOTTIDÆ.

Body short, elevated. Head very large, its greatest depth greater than that of the body; skull with 2 strong bony ridges from above the front of the eye, continuous with 2 large occipital ridges, leaving the interorbital space and middle line of the top of the head strongly concave; snout slender, narrow, and abruptly protruding; mouth very narrow, ∩-shaped, its gape longer than wide; teeth villiform, none on the vomer or palatines; gill opening confined to the region above the base of the pectorals, the membranes below completely united to the shoulder girdle and isthmus; apparently no slit behind last gill; a stout, straight, preopercular spine; nasal spines present; no other spines on head. Skin everywhere on head and body firm, immovable, densely covered with stiff, bifid or trifid spinous prickles; spinous dorsal very small; pectoral with procurrent base. North Pacific; a single species known; a small shore fish. The following account of the skeleton of *Rhamphocottus* is given by Mr. Starks: The posterior end of the prominent ridge, which runs backward from the superior orbital rim on each side, is formed by the epiotic process. It ends in the form of a long "occipital spine;" almost directly under it is the short parotic process. The post-temporal is short, wide, and flat; its upper end is attached to the inner side of the epiotic spine, and for the whole length of its anterior edge, to the skull between the epiotic and parotic processes. From its lower inner surface it sends a wide, thin bone, which is firmly fastened to the base of the skull. It bears a backward projecting spine on its lower end, inside of which the supraclavicle is attached. Actinosts large, wide and thin, without an opening between them. Subopercle absent; preopercle large, sending a spine backward; opercle triangular on its lower inner angle; the interopercle is developed and strongly coossified with it; it sends a slender process forward under the preopercle; a projection downward from the posterior end of the articular; suborbital wide, thin, and concavo-convex, its convex surface outward. Skull without basal chamber; vertebræ 10+14.

758. RHAMPHOCOTTUS, Günther.

Rhamphocottus, Günther, Ann. Mag. Nat. Hist., XIV, 1874, 370 (*richardsoni*).

Characters of the genus included above. (*ῥάμφος*, snout; *Cottus*.)

2397. RHAMPHOCOTTUS RICHARDSONI, Günther.

Head 2; depth 2. D. VII, 13; A. 6; P. 14; orbit 6 in head; maxillary 4; snout 3; highest dorsal spine 6⅔; dorsal ray 4; anal ray 4; pectoral 2⅙; ventral 2; caudal 3; vertebræ 10+14=24. Body short, compressed, the back elevated, its greatest depth just in front of spinous dorsal. Head large, as long as rest of body; snout long and narrow; mouth U-shaped, its gape longer than wide; lips thick, their surface broken up into papillæ; maxillary reaching the nasal spine; lower jaw included; teeth in villiform bands on jaws and vomer, none on palatines; eye placed high, its diameter contained twice in the snout, 1½ in the interorbital; a branched dermal flap, as long as pupil, at tip of snout; head with 2 large bony ridges above, continuous with the orbital rim and ending in strong blunt spines at occiput; head deeply concave between these ridges; nasal spine sharp and recurved; a pair of strong spines over the eyes; a sharp spine just above opercle, a blunt one on opercle below flap, and a long sharp one at angle of preopercle, a low bony ridge leading to each of these spines; a long sharp spine on clavicle just behind gill opening; a blunt bony knob at posterior end of mandibles; gill openings extending upward from upper pectoral ray, their length equal to snout. Entire head and body covered with multipointed spines, those on head much smaller than the ones on sides, a few simple spines along cephalic ridges; the first dorsal spine covered with spines, and each dorsal ray with a row on its side; a few spines on the base of the pectoral and anal rays. Dorsal spines very weak, fitting in a groove in back; soft dorsal higher than spinous, the tips of the rays reaching base of caudal fin; anal short, few-rayed, reaching slightly beyond soft dorsal; pectorals pointed, their lower rays entirely free, reaching about to base of third anal ray; ventrals reaching to ends of pectorals, their origin behind lower part of pectoral base a distance equal to length of snout; caudal rounded behind. Body creamy yellow, with conspicuous irregular dark stripes, edged with black, running obliquely across the body; similar stripes radiating from eye in all directions, 1 to end of snout, a triangular 1 downward, 1 running backward and downward to middle of preopercle, then turning upward and running nearly to occipital spine, 2 or 3 short ones above, each of these involving the membrane of eye; 2 or 3 black-bordered dark spots on edge of opercle; a light yellow streak surrounded by black across caudal peduncle, behind which all is bright cherry red to the end of caudal fin; 2 similar spots on base of pectoral; top of head crossed with wavy black-edged dark bars; top of lower jaw black, a line of black spots running along under parts of mandible; fins all bright red, each ray of dorsal with a sharp black spot at its base, a few spots on dorsal spines; anal, pectorals, and ventrals dark at base. Here described from a specimen 3

inches in length, collected in a rock pool on Channel Rocks, near Point Orchard, Puget Sound, by Miss Adella M. Parker, of Seattle. North Pacific, from Sitka to Monterey; scarce, but not infrequent in Puget Sound, at about 2 to 10 fathoms; a most singular fish. (Named for John Richardson, naturalist and explorer.)

Rhamphocottus richardsoni, GÜNTHER, Ann. Mag. Nat. Hist., XIV, 1874, 370, Fort Rupert, Vancouver Island, British America; BEAN, Proc. U. S. Nat. Mus. 1881, 252; JORDAN & GILBERT, Synopsis, 722, 1883; JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 813, pl. 87.

Family CLXXXI. AGONIDÆ.*

(THE SEA POACHERS.)

Body angular, commonly 8-angled, the caudal peduncle 6-angled, covered with 8 to 12 longitudinal rows of imbricated, radially striated plates, the anterior edge of each plate overlying the posterior edge of the plate next in front of it; plates spinous or not. Teeth small, even, in villiform bands on jaws, and in most species on vomer and palatines, sometimes wholly obsolete; gills 3½, no slit behind the last; pseudobranchiæ large, extending down the inner side of opercle; gill rakers small; gill membranes united, free, or joined to isthmus; ventral fins thoracic, narrow, their rays I, 2; vent usually close behind ventrals; spinous dorsal large, small, or absent; anal without spines; caudal rounded, about 3 times as long as wide at base, with 10 to 12 long rays; base of pectorals usually broad, the lower rays sometimes produced; all rays of all fins simple; branchiostegal rays 6; myodome (tube of *recti* muscles) with membranaceous roof; basisphenoid absent; post-temporal not bifurcate, continuously articulated with epiotic and pterotic; pyloric cæca few, about 4 to 7; vertebræ numerous, 35 to 50. Fishes of the cold seas, living among rocks or kelp, most of them of small size and fantastic form, not valuable as food. Genera 20; species about 40. The species are extremely varied, and must be placed in very many genera, or else reduced to a single one in each sub-family. The plates vary somewhat in number in all parts of the body in most if not all of the species, although not to the same extent in all the genera. Even the pattern on the breast, which is definite for the species of all the genera except *Hippocephalus*, varies in the different individuals of the same species, so that it is probable that even where no variations are indicated by the numbers given in the following descriptions they would be found by comparing large numbers of individuals. The plates in the dorsal series vary from 1 to 3 in number in most species; they correspond closely with the number of vertebræ, there being usually 1 or 2 fewer of the former than of the latter. It seems probable that the vertebræ vary a little in number within the species, and the rings of plates correspond in number with them at least throughout most of the length of the body. There is no definite proof, though the balance of evidence seems to indicate, that the superior lateral series of plates in *Aspidophoroides*, bearing the lateral line, corresponds with the median lateral series

* The account of the *Agonidæ* is contributed by Mr. Frank Cramer, a graduate student in Leland Stanford Jr. University, some additions having been made by Jordan & Evermann.

of plates of *Agonus*, etc. In the following descriptions of species, when the diameter of the eye is not specified, the longitudinal diameter is meant. The rays of the fins arise regularly in the intervals between the successive pairs of plates alternating with them, and the membrane behind the last spine of the first dorsal fin ends at variable distances from the base of the spine; hence in giving the number of plates between the dorsals, the number between the last spine and the first ray is meant, unless otherwise stated. "Length of body," as the standard of measurement, means the length from tip of premaxillary to base of caudal, unless "total length" is mentioned. The rostral spines, whether 1 or 2 pairs, belong to the nasal bones, and are properly "nasal spines" in the species in which there is no free terminal plate. The latter, with its spines, may be the freed anterior part of the nasal bones. The hypural bone is included in the number of vertebræ, and these were counted in nearly all cases in only 1 individual. The young of several species have been described, some of them in detail, and so far as known they all differ from the adults in the same way. The spines and ridges are more prominent; the body is relatively shorter, its anterior end abruptly broader, forming with the head a more or less definite rhombic figure; the tail is more compressed; the head is broader, the snout more blunt, and the lower jaw relatively longer; the vent lies farther back from the base of the ventral fins, and there are usually pairs of small plates between them; the dorsal and ventral series of plates on the tail, which are single in the adults of nearly all the species, are double in the young, or at least have the double row of serrations. Nearly all the species are found within the limits assigned to this work. The few not so found are included, for completeness' sake, in footnotes. (F. C.)

a. Spinous dorsal present.
  b. Gill membranes free from the isthmus.
  PERCIDINÆ:
    c. Body compressed; lower jaw not projecting; plates of body spinous; first dorsal at nape.
      d. Body elongate; dorsals well separated.
        e. Teeth on vomer; no occipital spines or barbel on snout.
          PERCIS, 759.
        ee. No teeth on vomer; snout with a long barbel; occipital spines present.
          AGONOMALUS, 760.
      dd. Body short and high; dorsals close together; vomer without teeth; occipital spines present; lower rays of pectorals free.
        HYPSAGONUS, 761.
  BRACHYOPSINÆ:
    cc. Body more or less depressed; lower jaw projecting; plates of body spinous or not; first dorsal behind nape.
      f. Chin without terminal barbel.
        g. Snout short (not produced in form of a tube); plates of body spinous.
          h. Bones of snout short, not forming a tubular nose.
            i. Vomer and palatines without teeth; breast simply prickly.
              STELLERINA, 762.
            ii. Vomer and palatines with some teeth; breast with large plates.
              OCCA, 763.
          hh. Bones of snout produced into a long tube which bears the short jaws at the end; body rather robust, the plates with spines.
            BRACHYOPSIS, 764.

ff. A long barbel at tip of chin; snout long (produced into form of a tube, as in *Syngnathidæ*); body subterete, very slender, its plates not spinous. PALLASINA, 765.

AGONINÆ:

bb. Gill membranes joined to isthmus, with or without a narrow free fold behind.

j. Tip of snout without free median plate or spine; mouth inferior, lower jaw short.

k. Vomer without teeth.

l. Lower side of snout with no barbels; one pair of spines on tip of snout. LEPTAGONUS, 766.

ll. Lower side of snout with barbels.

m. Gill membranes without barbels; a pair of complex barbels under tip of snout; 1 pair horizontal and 1 pair recurved spines at tip of snout. Lower jaw very weak; teeth feeble, sometimes wanting; plates of body mostly with spines. PODOTHECUS, 767.

mm. Gill membranes with scattered barbels.

n. Plates of body without prominent spines; no preocular spines. AGONUS, 768.

nn. Plates of body largely spinous; a pair of preocular spines. STELGIS, 769.

kk. Vomer and, often, palatines with teeth.

o. Dorsal fins rather long, the rays shortened behind, the last one attached by membrane to the body; no large life-like spine above eye.

p. Plates on body largely spinous; gill membranes with cirri, 1 or more on each branchiostegal; teeth on palatines; lower jaw short beneath the long snout.* AVERRUNCUS, 770.

pp. Plates on body largely unarmed; no teeth on palatines; gill membranes without cirri. SARRITOR, 771.

oo. Dorsal fins rather short, the rays scarcely shortened behind, the last one not adnate by membrane behind; a large knife-like spine over each eye; plates on body largely spinous; no cirri on gill membranes. XYSTES, 772.

jj. Tip of snout with a free terminal plate or spine; teeth on jaws, vomer, and palatines.

q. Occiput without distinct pit; 2 pairs of occipital spines; dorsal and anal well developed.

r. Lower jaw strongly projecting; lower pectoral rays not much produced; terminal rostral plate with 3 spines. BATHYAGONUS, 773.

rr. Lower jaw not projecting; lower rays of pectorals much produced in adults; anal fin short, of 6 or 8 rays. XENOCHIRUS, 774.

qq. Occiput with a deep pit; occipital spines none, or reduced to mere traces; dorsal and anal very short.

s. Body slender, elongate; tail not compressed. ODONTOPYXIS, 775.

ss. Body short and broad in front; tail compressed teeth. BOTHRAGONUS, 776.

ASPIDOPHOROIDINÆ:

aa. Spinous dorsal absent; body not compressed, its plates not spinous; mouth small, terminal; gill membranes free from the isthmus. ASPIDOPHOROIDES, 777.

759. PERCIS,† Scopoli.

Percis, SCOPOLI, Int. Hist. Nat., 454, 1777 (*japonicus*).
Hippocephalus, SWAINSON, Nat. Hist. Fishes, etc., II, 272, 1839 (*superciliosus*).

* Jaws subequal in *Agonopsis*, a South American genus very close to *Averruncus*.
† Dr. Gill has called our attention to the identity of *Percis* with *Hippocephalus*.

Body moderately elongate, compressed throughout; back elevated behind nape; 2 rows of strong, curved spines along whole length of each side (spines of the other series smaller); first dorsal fin beginning behind nape; dorsals far apart; anal fin long. Vent far back from base of ventrals. Head narrow anteriorly, abruptly broader behind; mouth terminal; median rostral plate none; nasals united in front of maxillary pedicles. Interorbital space broad, the supraocular ridges very prominent, with large, flat, triangular supraocular spine or shelf; no occipital spines. Teeth on jaws and vomer, none on palatines. Gill membranes united, free from isthmus. North Pacific. (*περκίς*, a synonym of *περκη*, perch.)

2398. PERCIS JAPONICUS* (Pallas).

Head $4\frac{1}{3}$; Br. 6; D. VI, 7; A. 8; P. 12; V. 3 (I, 2); C. 2–13–2. Body compressed throughout, the shoulder girdles prominent; width of body at base of pectorals a little more than greatest height and 5 in body length. Immediately behind pectorals the width a little less than greatest height; between dorsals and on peduncle it is nearly 2, and under second dorsal a little more than 2 in the height at the same points. Abdomen moderately swollen in front of vent. The back rises at an angle of 45° behind occiput, slopes downward under first dorsal, leaving a hump under its front end, rises in front of second dorsal and slopes downward again under it; ventral outline nearly straight; vent between $\frac{1}{3}$ and $\frac{1}{2}$ distance from ventrals to anal. Breast and area between ventrals and vent almost completely occupied by nearly flat, radially striated plates, with slightly raised centers and of variable size, with innumerable, excessively minute plates scattered among them; vent surrounded by prickles. Plates of ventrolateral series small, beginning as distinct rows of spinous plates about opposite vent, with an imperfect row of smaller spinous plates between them and the vent on each side, and converging toward anal. They pass along sides of anal, remain distinct to about the eighth pair of plates behind anal, and unite in a single median plate with a double spine; from this plate to caudal the rows are again distinct, the plates alternating instead of standing opposite each other; the superior and inferior lateral series begin as distinct rows of spinous plates about opposite middle of first dorsal, diverge to about front of second dorsal, and converge toward base of caudal; the plates elongated vertically, their spines, rising abruptly from their centers, are strong, thick, blunt, curved; about 37 plates in inferior and 35 in superior longitudinal row; between them lies

* Diagnosis: Body moderately slender, compressed, its width between the dorsals nearly 2 in depth at same place; 2 rows of strong and 2 of weak spines along each side of body, and a median lateral row of spineless plates. Head $4\frac{1}{5}$. Preopercular spine large, suborbital spine tubercular. Supraocular ridge expanded into a triangular shelf projecting laterally far beyond eye and ending bluntly; no occipital spines. Back elevated behind nape. First dorsal short, beginning at nape. Plates in dorsal series 43 to 45; between dorsals (from last spine to first ray) 14 pairs (9 pairs between end of membrane of first dorsal and first ray of second dorsal. Nasal spines sharp, far apart, at tip of snout. No median rostral plate. Gill membranes united, free from isthmus. Teeth in broad bands on jaws and on vomer; none on palatines. Vent between $\frac{1}{3}$ and $\frac{1}{2}$ distance from ventral to anal. Color (in spirits) "old ivory," with brown patches on sides, 1 under first dorsal, 1 between dorsals, 1 under second dorsal, 2 or 3 on peduncle. B. 6; 1 D. 6; 2 D. 7; A. 8; P. 12; V. 3 (I–2); C. 2–13–2.

the median lateral series, bearing the lateral line, with a nearly continuous row of 38 or 39 spineless plates from head to caudal (a few of them with small, blunt tubercles) dorsolateral series composed of small, indistinct plates from occiput to beyond middle of first dorsal, where they become larger and spinous; plates of sixth, seventh, eighth, and ninth pairs behind last ray of second dorsal closely approximated, but not forming single median plates; from ninth pair to caudal the plates of the 2 rows alternate as on ventral surface; plates in dorsal series 43 to 45. There are almost complete series of small plates, alternating with the large ones, between the dorso and superior lateral series, between the superior and median lateral, between the median and inferior lateral, and between the inferior and ventro lateral series; this multiplication by intercalation of small plates is evident all over the body. There are no large, but numerous minute, plates in front of and on base of pectorals. Gill membranes posteriorly and medially with several rather weak plates and many minute plates and prickles, anteriorly and laterally with a few small plates; underside of lower jaw with many imperfect plates or strong prickles; weaker prickles along the branchiostegal rays. Head depressed throughout, narrow to posterior border of orbits, behind which it widens rapidly. Depth behind orbits $\frac{7}{9}$, and over opercles $\frac{3}{4}$, of width at same points; orbits large, oval, $4\frac{1}{2}$ in head, far forward; interorbital space a little concave, very broad; at anterior border of orbits wider than the orbit itself, $3\frac{1}{2}$ in head; at posterior border of orbit $2\frac{3}{8}$ in head. Supraorbital ridges expanded into a flat, triangular shelf projecting upward and outward over eye and ending bluntly. Occipital ridges heavy, low, spineless, forming low domes behind; space between them concave. Preorbital with about 2 ridges radiating from the anterior inferior border of orbit, each ending in a short, broad, plate-like blunt spine; supraorbital with a broad, dome-like tubercle; preopercle with a dorsoventrally compressed spine; below this 3 successively smaller blunt spines; cheek below orbit and suborbital with 4 or 5 fairly developed and many minute plates; nasal spines far apart, near tip of snout sharp, nearly upright. A large membraneous nasal tube; a short, flap-like barbel at tip of each maxillary; mucous pores of lower jaw with flap-like borders. Mouth small, terminal; no median rostral plate; maxillary reaching a little beyond front of orbit; angle of jaw prominent, tubercular. Dorsals far apart (about 14 pairs of plates between last spine and first ray; 9 pairs from end of first dorsal membrane to first ray), the first a short distance behind occiput, its spines very rough (like the rays of all the fins), with minute prickles or plates; its leathery membrane also rough; spines 6, transversely broad at base, the fourth longest, 2 in head; base of second dorsal about $1\frac{1}{3}$ in that of first dorsal; rays 7, the third longest, about $1\frac{1}{2}$ in head; distance between dorsals about equal to base of first dorsal; anal beginning about 3 plates in front of second dorsal; rays 8, the fifth longest; the membrane notched, the distal third of the 3 anterior rays exserted, with a narrow border of membrane; caudal rounded, its base about 2 in its length; pectorals close to gill openings, $4\frac{1}{5}$ in length of body, their base about 3 in their length, rays a little exserted; ventrals (male) less than 2 in pectorals, the inner ray

longer, 8 in body. Lateral line with about 11 pores on the anterior and about 9 on posterior part of body. Gill membranes united, free from isthmus. Teeth on jaws and vomer; none on palatines. Color, pale brownish or yellowish ("old ivory"); a light-brown cross bar across the back in front of and under first dorsal, narrowing to upper end of base of pectoral; another, of irregular outline, passing downward and forward to posterior side of base of pectoral; a larger one, mostly below the superior lateral series, behind first dorsal; 1 under second dorsal, and 3 on peduncle, the last just in front of base of caudal; dorsal fins dusky, with small darker patches; first dorsal with a pale oblique bar near its base; distal half of caudal dark, its tip edged with lighter; posterior half of anal dusky; pectorals marbled with yellowish and brown, the membrane of the distal third dusky, with paler edge; ventrals pale; 4 or 5 brown spots on nape; a curved band of brown connecting the posterior ends of the occipital ridges, another between the latter and the temporal ridges, and a streak extending backward from posterior border of orbit; brown areas on opercle and preopercle. Length 14½ inches. North Pacific; Kuril Islands (Steller), Gulf of Patience, Island of Saghalien (Tilesius), Okhotsk Sea (Dr. Krause); not known from Japan. Here described from a specimen from Okhotsk Sea in Coll. Cal. Ac. Sci., the only one seen since the time of Pallas and Tilesius. (*japonicus*, from Japan, Japonia, because, according to Steller, it is more common toward the south, i. e., in Japan.)

Cottus japonicus, PALLAS, Spicilegia Zoologia, VII, 30, pl. 5, figs. 1–3, 1772, dry specimen, Kuril Islands (Coll. G. W. Steller); TILESIUS, Krusenstern's Reise um die Welt, IV, pl. 87, 1813.

Agonus curilicus, TILESIUS, Mém. Acad. St. Petersb., IV, 1811, 416; after PALLAS and STELLER.

Agonus stegophthalmus, TILESIUS, Memoirs Soc. Naturalists, Moscow, II, 219, 1809, Gulf of Patience, Saghalien Island; TILESIUS, Mém. Acad. St. Petersb., IV, 1811, 427, pl. 12; GÜNTHER, Cat., II, 214, 1860.

Aspidophorus lisiza, LACÉPÈDE, Hist. Nat. des Poiss., III, 1802; after PALLAS.

Aspidophorus superciliosus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 215, 1829, Kuril Islands; after PALLAS and TILESIUS.

Percis japonicus, SCOPOLI, Int. Hist. Nat., 454, 1777.

Agonus japonicus, BLOCH & SCHNEIDER, Systema Ichthyologia, 105, 1801.

Phalangistes japonicus, PALLAS, Zoog. Rosso-Asiat., III, 112, 1811.

Hippocephalus superciliosus, SWAINSON, Nat. Hist. Fishes, etc., II, 272, 1839.

Hippocephalus japonicus, JORDAN & GILBERT, Synopsis, 723, 1883; CRAMER, Proc. Cal. Ac. Sci. 1894, 194, with figure.

760. AGONOMALUS, Guichenot.

Agonomalus, GUICHENOT, Mém. Soc. Sci. Nat. de Cherbourg, IX, 1866, 252, pl. 9 (*proboscidalis*).

Head and body strongly compressed, the latter entirely armed with angular osseous plates. Teeth excessively small on both jaws, none on vomer or palatines. Dorsal fins separate; a long, fleshy barbel on tip of snout; mouth small; a very high, broad, somewhat recurved spine or bony projection above each orbit; occipital spines not prominent; origin of first dorsal above nape, the profile from nape to fin very steep; pectoral rays somewhat exserted; gill membranes probably united and free from isthmus. Closely related to *Hypsagonus*. (*Agonus*; ὁμαλός, level, even, or flat; i. e., compressed.)

2399. AGONOMALUS PROBOSCIDALIS (Valenciennes).

Height of body $5\frac{1}{3}$ in total length; greatest width of abdomen $\frac{5}{6}$ of height. Br. 6; D. VIII, 6; A. 12; P. 11; V. 3 (I, 2); C. 2–11–2. Body slender, compressed. Side of body throughout its length with 2 rows of spinous plates, 1 above and 1 below lateral line, the plates oval, radially striated, the spines high, curved, compressed upon middle of plates. Behind pectoral girdle, between first plates of superior and inferior spinous series, a circular striated plate with a small spine; a small spine upon a plate on the pectoral girdle above insertion of pectoral fin. Sides between lateral spinous series a little concave. The rounded, somewhat prominent abdomen with small striated plates armed with low blunt tubercles. Profile of body rising vertically behind nape and under first spine of first dorsal, running in a straight line to second dorsal, rising a little under this fin, and descending gradually to caudal. Inferior profile nearly rectilinear. Head small, compressed, with a thick long median barbel at tip of snout; supraocular ridges high, passing down anterior border of orbit; the supraocular spines large, flat, thin; interorbital space deeply concave, with a small longitudinal ridge; nasal spines present; a pair of occipital spines present; also a suprascapular tubercle; none on opercle, but a large, oblong, compressed spine on preopercle and 1 on suborbital; eye large, circular, 4 in head. The preorbital with ridges, ending at its lower border in strong denticulations. Mouth small; lower jaw shorter than the upper; teeth excessively small; none on palatines. Spines of first dorsal large, curved, and higher than the body beneath them; second dorsal lower, its rays strong and curved; anal very long; caudal narrow and rounded; pectoral very large, its longest rays equal to longest spines of first dorsal; rays simple and rough; ventrals small. Lateral line indicated by a series of small longitudinal "traits" set off on naked space between the 2 longitudinal series of large spinous plates (after Valenciennes). Guichenot says the total length is scarcely $6\frac{3}{5}$ inches (16 cm.). His figure, which seems to be good, furnishes the following facts: About 7 plates of the superior lateral series between the adjacent rays of first and second dorsals, and about 2 between posterior edge of membrane of first dorsal and first ray of second dorsal; 10 plates from last ray of second dorsal to base of caudal. A small but distinct tubercle or spine posteriorly at base of supraocular spine. Occipital spine apparently very close to elevated part of back, but distinct; in front of its base a small but distinct tubercle, as in *Hypsagonus quadricornis*. Spines of first dorsal all exserted, the first for about $\frac{1}{4}$ of its length; all rays of anal exserted, the anterior 3 or 4 for about $\frac{3}{4}$ of their length; posterior anal rays (except the last) longest; all rays of pectoral considerably exserted (about $\frac{1}{3}$ of their length); 27 spinous plates in the series above lateral line, 30 in the series below lateral line. One specimen known, taken in 1 of the coves of the Port of the Emperor Nicolas, Gulf of Tartary (Nicolaevsk, mouth of Amur River, west of the Island of Saghalien). (Guichenot.) (*proboscidalis*, bearing a proboscis; in allusion to the prominent barbel on snout.)

Aspidophorus proboscidalis, VALENCIENNES, Comptes Rendus de l'Acad. des Sciences, XLVII, 1040, 1858, Port of Emperor Nicolas (Nicolaevsk), Gulf of Tartary.

Agonomalus proboscidalis, GUICHENOT, Mém. Soc. Sci. Nat. Cherbourg, 1865, 254, pl. 9; SAUVAGE, Nouv. Arch. Museum Hist. Nat., Paris (2), I, 1878, 157.

761. HYPSAGONUS, Gill.

Hypsagonus, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 259 (*quadricornis*).

Cheiragonus, HERZENSTEIN, Bull. Acad. Imp. Sci. de St. Petersb., XIII, 1890 116, (*gradiens* = *quadricornis*).

Body compressed, elevated, depth greater than length of head, more than $\frac{1}{3}$ of body; head small, separated from first dorsal by a very deep nuchal depression; top of head very uneven, 1 pair of large supraocular and 1 pair of large occipital spines, strong, blunt; mouth terminal, jaws about equal, a large barbel or none at tip of snout; teeth on jaws, none on vomer or palatines; gill membranes united, free from isthmus; scales or plates large, radially striated, with a central spine or tubercle; 2 rows of strong and 2 of weaker spines along side of body; dorsal fin long, high, beginning immediately behind nape, the spines strong, the first serrated; pectorals short, procurrent, the lower 8 or 9 rays free; ventrals small; vent nearly halfway between ventrals and anal. (*ὑψι-* high; *Agonus*.)

2400. HYPSAGONUS QUADRICORNIS* (Cuvier & Valenciennes).

Br. 6; D. IX to XI (rarely IX, usually X), 6 or 7; A. 9 or 10; P. 13 or 14; V. 3 (I, 2); C. 13; vertebræ $\frac{15}{21}$ (36). Pyloric cæca 5 (2 individuals). Body strongly compressed throughout its length. Abdomen swollen (much less in male than female) about $\frac{7}{10}$ of the depth, the latter 3 to $4\frac{1}{3}$ (usually about 4) in the length; body much narrower under second dorsal, only about $\frac{1}{3}$ of depth. Back rises almost perpendicularly behind the nape, slopes downward to end of first dorsal, rises gently to base of second dorsal, then sinks and rises again to base of caudal; ventral outline gently convex. Plates distributed in general as in the other Agonoids, but with many deviations in detail. Interspaces membranaceous, studded with a variable number of small plates; vent nearly halfway between ventrals and anal, the membrane between it and ventrals with scattered plates; plates of ventrolateral series small, the spines short, blunt, strongest from vent to anal; those of the inferior lateral series elongated vertically, beginning behind middle of base of pectorals, with long, strong, curved spines, which rise abruptly from the center of the plates, growing smaller posteriorly, minute at base of caudal, their number varying from 28 to 30. The series of the lateral line consists of 7 to 9 plates

*Diagnosis: Body short, much compressed throughout its whole length; its greatest width 3 to $4\frac{1}{2}$ in depth; 2 rows of strong and 2 of weaker spines along side of body; a strong spine above base of pectoral; head $3\frac{1}{2}$; preopercular spines present, large; 1 pair large supraocular and 1 pair large occipital spines, strong and blunt. Body abruptly elevated behind nape; first dorsal long, high, beginning immediately behind nape; 8 or 9 lower rays of pectoral free; plates in dorsal series about 33 to 35; between dorsals about 5; usually a single barbel at tip of snout; no median rostral plate; gill membranes broadly united, free from isthmus; no teeth on vomer or palatines; vent nearly halfway between ventrals and anal. Color, front half of body blackish violet (variable), hinder half pale, yellowish, with 2 or 3 narrow dark bands; caudal with dark band distally.

(sometimes varying on the 2 sides of the same animal), the first 2 (occasionally the first 3) very large, supplying the lack of plates in the superior lateral series anteriorly and sometimes apparently forming a part of this series, their spines long, strong, and curved; remaining 5 or 6 plates distributed at varying distances along side of body, small, but sometimes provided with small spines; the superior lateral series begins under about the seventh spine of first dorsal and runs in a straight line to caudal, the spines a little smaller than those of the inferior lateral series, and vary from 25 to 27 in number (plus the 2 or 3 plates of the lateral line when they appear to belong to this series). The dorsolateral series begins behind occipital spine and follows outline of back close under dorsals, 33 to 35 pairs; plates small, with tubercles or blunt spines. Between the dorsolateral series and the base of each dorsal fin is a supplementary series of very small tuberculated plates or prickles; those under the anterior fin sometimes become apparently continuous with the dorsolateral series, crowding the anterior plates of the latter downward; about 5 pairs between dorsal fins. Behind pectoral, between the inferior lateral series and the large anterior pore plates there are usually 2 or 3 plates, of which 1 or 2 sometimes bear minute spines; 1 plate with a large, strong spine just above base of pectoral; none in front of the fin, which is close to gill opening. No plates on branchiostegal membranes or lower jaw. Head compressed, snout $3\frac{1}{2}$; nearly as wide as high across preopercle, much narrower under orbits. Orbits large, circular, high up, 3 in head. Interorbital space wide behind, much narrower in front, concave; supraorbital ridges high, shelving outward over eye, ending in strong spines directed outward, upward, and a little backward, in front of the base of which there is a very small one, usually in the form of a tubercle of variable size (most distinct in the young). Preorbital with 2 or 3 ridges radiating from anterior inferior border of orbit, its edge scalloped; suborbital with a strong, compressed spine projecting nearly at a right angle; preopercle with a strong, compressed, curved spine, and below it 2 or 3 flattened smaller ones; nasal spines strong, curved, separate, somewhat diverging; 1 or 2 or no plates on lower part of cheek; no median rostral plate. Barbel at tip of snout, above premaxillaries, varying much in length, frequently entirely absent. Mouth small, terminal; maxillary reaching front of orbit. Teeth on jaws; none on vomer or palatines. Angle of jaws prominent, tubercular. Gill membranes broadly united, free from isthmus. First dorsal about $2\frac{1}{2}$ times as long as second, high, its spines strong, nearly upright, the first very, and some of the others somewhat, granulated or beset with prickles; the anterior spines but little exserted; membrane stretched. Second dorsal low, its rays stiff; caudal nearly truncate, its width at base about $1\frac{3}{8}$ in length; anal long, beginning under posterior end of first dorsal and ending opposite posterior end of second dorsal; 2 or 3 of the posterior rays (except the last) longest, all the rays much, the anterior more than half, exserted; ventrals of female 8 or 9 in length of body; in male longer, 6 to $7\frac{1}{3}$ in body length. Abdominal region much swollen in female. Lateral line, pores 15 to 19, in pairs, except the first. Vent nearly halfway between ventrals and anal. Color

blackish or grayish violet, paler below; breast and belly in front of vent marbled with darker; anterior region to opposite posterior end of first dorsal dark, the dark region limited by a still darker band which runs up on the fin; posterior part of body paler, with narrow dark cross bands, 1 under the anterior and 1 under the posterior part of the second dorsal, both passing down onto the anal; caudal peduncle darker, sometimes with irregular patches more intensely dark; caudal with a dark band at base, and another near distal end, the extreme tip white-edged; rays of anal with black spots near their tips; pectorals indefinitely cross-barred by series of short black streaks on rays; ventrals of both sexes with about 2 narrow bars of black. A variable species, widely diffused. Length 2½ inches. North Pacific; Kamchatka, Bering Sea, and coasts of Alaska, south to Bristol Bay and Puget Sound. (Gilbert.) Here described from specimens taken by Dr. Gilbert (*Albatross* Coll.) about the Aleutian Islands. On these specimens Dr. Gilbert has the following notes:

Taken north and south of the Aleutian Islands, in the shallow water; also at 1 station in Bristol Bay, in depths from 34 to 56 fathoms. Our specimens agree perfectly with the description of *Hypsagonus* (*Cheiragonus*) *gradiens*, Herzenstein (Bull. Acad. Imp. des Sci. de St. Petersb., XIII, 116, May 29, 1890), described from the Gulf of Avatcha in Kamchatka. Dr. G. A. Boulenger, of the British Museum, has kindly compared one of our specimens with the type of *Aspidophorus quadricornis*, C. & V., and states that they are undoubtedly identical. Body short, much compressed, the head also narrow and compressed, especially above and in front. Nasal spines short and strong, a slender barbel, of varying length, in front of them on middle line of snout. Ocular region abruptly rising above the short, slender snout, the eyes vertical, overarched by the supraorbital rim, which bears posteriorly a strong vertical spine. Interorbital space with a deep median groove and without ridges, the occipital region depressed below the bottom of the groove. No deep pit on occiput, the space being gently concave transversely, bounded laterally by moderate ridges, which bear posteriorly a spine preceded by a lone tubercle. A strong spine at lower inferior border of orbit; a strong spine at upper preopercular angle and 3 smaller ones below it; a strong spine above the base of pectoral, behind and above which on sides are 2 strong spines nearly in line with the upper lateral series of plates; surface of opercles with a few short spinous processes, but without definite spine. Vomer and palatines toothless. Branchiostegal membranes broadly joined, forming a free fold across the isthmus. Mouth narrow, horizontal, terminal, the lower jaw included. Maxillary reaching vertical slightly behind front of orbit, 3¾ in head. Eye large, 3 in head; snout 4; interorbital width over middle of orbit ⅓ head behind snout. Nape rising very abruptly from occiput to front of dorsal, the outline thence descending to near front of second dorsal when it again ascends. The points of origin of the 2 dorsals are therefore prominent, the profile concave behind them. Body deepest under first dorsal spine, 3⅞ in length. Greatest width of body near ventral outline immediately behind ventral fin, 5½ in length. A series of small prickles along base of spinous and anterior portion of soft dorsal, bearing 1 spinule to each ray. Plates all concealed, the spines alone projecting; upper lateral series of plates very small, bearing each a minute spinous point; this series is narrowly separated from base of spinous dorsal, and runs along base of soft dorsal, the 2 series uniting immediately behind the latter, bearing each a pair of diverging spines; lateral series with very strong spines, the largest being the anterior ones of the lower lateral series. The upper lateral series is incomplete anteriorly, ending under the eighth dorsal spine. It is apparently completed by 2 very strong spines, which belong, however, to the series of the lateral line. Ventral series of spines small, running along immediate base of anal fin, the pairs uniting behind the anal, the resulting plates bearing a pair of spines. As in the case of the dorsal series, this union is more or less irregular and incomplete, the corresponding plates sometimes failing to unite, and then either maintaining their opposite position or alternat-

ing. Lateral line with few widely spaced pairs of pores, those of each pair approximated and horizontally instead of vertically placed. The interspaces bear in the posterior part of the body a minute prickle each; in the case of the 2 or 3 anterior pairs, these become very strong spines, nearly on a line with the incomplete upper lateral series of plates. Anus anterior, nearly midway between base of ventrals and front of anal. First dorsal spine vertically over upper axil of pectoral. The fin is rigidly spread in alcoholic specimens; the third and fourth spines are longest and about equal length of snout and eye; interspace between dorsals equaling ½ length of orbit; anal much longer than second dorsal, its first ray under last dorsal spine, its last ray slightly in advance of the last ray of soft dorsal; anal membranes deeply incised, especially anteriorly. Caudal short, rounded; pectoral of 2 distinct divisions, the upper portion consisting of 4 or 5 rays joined by membrane; the lower part of 8 entirely disconnected rays. These upper and lower portions of the fin are used alternately in pushing the fish forward on the bottom, the upper lobe being downward and forward for the purpose. In the aquarium, the fish appears to walk, resting alternately on the upper and lower pectoral rays and on the front rays of the anal; the longest pectoral rays reach to or just beyond front of anal; ventrals short, not reaching vent in females, reaching to or slightly beyond vent in males. D. IX to XI, 6 or 7; A. 9 or 10; P. 13 or 14; V. I, 2; C. 13. Vertebræ 8 + 28. Lateral line 7 to 9. Color: blackish or greyish violet, paler below; breast and belly in front of vent marbled with darker; anterior part of sides, to opposite last dorsal spines dark, the darker region limited by a still darker band which runs up on the fin; posterior part of body paler, usually with 3 darker cross bars, the last of which often broadens out to occupy all of the caudal peduncle; caudal with a dark bar at base and another at posterior margin, the extreme edge white; the dark vertical bars continued onto the anal fin, the rays also spotted with back near tips; ventrals with a black bar or spot at base; sometimes a second on middle of fin; pectorals indefinitely cross-barred, largely pale on basal portion; color varying greatly, the lighter areas on body and fins often with dusky marblings. (Gilbert.) (*quatuor*, four; *cornu*, horn.)

Aspidophorus quadricornis, CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 221, 1829, **Kamchatka.** (Coll. Dr. Collée. Type in British Museum.)

Hypsagonus (*Cheiragonus*) *gradiens*, HERZENSTEIN, Bull. Acad. Imp. des Sci. de St. Petersb., tome XIII, 116, May 29, 1890, **Kamchatka, Sinus Awatska, Port Petropaulski.**

Agonus quadricornis, GÜNTHER, Cat., II, 215, 1860.

Hypsagonus quadricornis, GILL, Proc. Ac Nat. Sci. Phila., XIII, 1861, 167; JORDAN & GILBERT, Synopsis, 722, 1883; GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 439.

762. STELLERINA, Cramer.

Stellerina, CRAMER, in JORDAN & EVERMANN, Check-List Fishes, 447, 1896 (*xyosternus*).

Body moderately elongate, tapering uniformly from head to caudal, 8-hedral, subterete, a little depressed anteriorly, depth about 7, width about 6 in length; head 4½, tapering but little forward; profile of head and snout in straight line with back. Plates in dorsal series about 35; between occiput and first dorsal 6 or 7 pairs. Plates of body spinous; breast covered with radially striated prickles. Mouth very oblique, the snout short, not tubular, lower jaw protruding, entering profile; teeth small, in few rows on jaws, none on vomer or palatines. A barbel at tip of each maxillary. No median rostral plate. Nasal spines small; no supraocular or occipital spines. Gill membranes united, free from the isthmus. Anal long. (Named for Georg Wilhelm Steller, naturalist and explorer, the first to study the fishes of Bering Sea.)

2401. STELLERINA XYOSTERNA* (Jordan & Gilbert).

B. 6; D. VI or VII–6; A. 8 or 9; P. 17 to 19; V. 3 (I, 2); C. 1, 11, 1; lateral line 36. Body slightly depressed in front, as high as wide under first dorsal, 8-hedral, subterete, the faces but little concave; tail strongly depressed, hexagonal. The sharp, slightly curved spines present on all the plates of each series, but minute on the median, dorsal and ventral plates of caudal peduncle; the superior lateral series beginning opposite front of first dorsal, being replaced anteriorly by the series of the lateral line. Unlike the species of *Phalangistes* (in which there are several), only 1 pair of plates between last ray of second dorsal and first median plate and 1 between last anal ray and first median plate. Breast, unlike that of other species, without plates, but covered with innumerable radially striated prickles; edge of breast in front of lower end of base of pectoral with 3 to 5 plates. Branchiostegal membrane and under side of mandible naked; membrane between the ventrolateral series behind and around vent with minute scattered prickles; 2 or 3 spinous plates in front of pectoral; the inferior lateral series, unlike that of *O. dodecaedron*, running directly toward base of pectoral. Head tapering but little forward, as high as wide, 4½ in length. Orbits moderate, nearly circular, longer than snout, a little more than 3 in head; interorbital space nearly flat, nearly 2 in orbit; supraorbital and occipital ridges but little developed, spineless; temporal ridge moderate; a small suprascapular spine. Cheeks vertical; 2 minute spines at border of preorbital, 1 on suborbital, 2 rather large and sometimes a third small one on preopercle; no plates on lower part of cheek; nasal spines minute, converging behind the maxillary pedicles. Median rostral plate absent; mouth very oblique, lower jaw protruding, entering profile; maxillary nearly reaching pupil, enlarged at tip. Teeth few and small in few rows on jaws; none on vomer or palatines. One barbel at tip of each maxillary. Gill openings large, membranes united, free from isthmus. Dorsals well separated; anal long, beginning under posterior end of first dorsal; caudal about 3 times as long as wide; pectoral 4⅔ in length of body, the base more than 3 in the length, with 17 to 19 rays (19 rays in 1 pectoral of a small specimen, 18 in the other), rounded, the lower rays rapidly shortening; ventrals of female 2 in pectorals, 1⅔ in male; tips of rays exserted. Color, light olive brown, pale below; top of head and back with minute spots and sinuous streaks of darker; lower jaw and lower edge of cheek silvery; pectorals dark, with a broad pale patch below, near base, and a broad pale tip; ventrals pale in both sexes (?); rays of dorsals dark; caudal dark; posterior part of anal dark, as in B. *dodecaëdron*. Length 6½ inches. Coast of California and Oregon, rather rare, south to Monterey. Here described from specimens from coast of Oregon, taken in 24 fathoms. (ξύω, to scrape; στέρνον, breast.)

* Diagnosis: Body moderately elongate, tapering uniformly from head to caudal, 8-hedral a little depressed anteriorly, the depth at base of pectorals ⅔ of width, the latter 6 in length; profile of head and snout nearly in a straight line with back; head but little tapering forward, 4⅔ in body; plates in dorsal series 35; between occiput and first dorsal 6 or 7; between dorsals 4 to 6; from ventrals to anal about 10 pairs. One barbel at tip of maxillary; gill membranes free; median rostral plate wanting; nasal spines small; no supraocular or occipital spines. Color light olive brown, pale below; mandible and lower part of cheek silvery.

Brachyopsis xyosternus, JORDAN & GILBERT, Proc. U. S. Nat. Mus., III, July 2, 1880, 152, Santa Cruz, California; (Coll. Dr. C. L. Anderson); JORDAN & GILBERT, Synopsis, 727, 1883; JORDAN, Cat. Fishes N. A., 113, 1885.

Agonus (*Brachyopsis*) *annæ*, STEINDACHNER, Ichth. Beiträge, IX, 254 (17), pl. 6, figs. 1, 1a, and 1b, Sitzb. der K. Acad. der Wiss., LXXXII, July 15, 1880, San Francisco, California.

763. OCCA, Jordan & Evermann, new genus.

Occa, JORDAN & EVERMANN, new genus (*verrucosus*).

This genus differs from *Brachyopsis* in the short snout which is not tubular, and is formed as in *Stellerina*. From the latter genus it is separated by the large plates on the breast, and by the presence of at least a few teeth on the vomer and palatines. North Pacific. (*occa*, a harrow.)

a. Anal rays 10 to 12; plates of belly bluntly spinous; ventrals pale. VERRUCOSA, 2402.
aa. Anal rays 14 to 16; plates of belly nearly smooth; ventrals dark.
DODECAEDRON, 2403.

2402. OCCA VERRUCOSA* (Lockington).

Br. 6; D. VII to IX–7 or 8; A. 11 or 12 (10 or 11, Lockington); P. 14 or 15; V. 3 (I, 2); C. 1, 11, 1; vertebræ 13 + 24 = 37 (37). Pyloric cæca 6 (1 individual); lateral line 36 or 37. Body and tail depressed throughout, the former rounded octagonal, the faces little concave; caudal peduncle hexagonal, short, with about 7 median dorsal and 5 or 6 median ventral plates; the sharp curved spines better developed than in *O. dodecaedron*, present on all ridges from head to caudal, except 2 or 3 plates of median ventral series of peduncle; plates of median lateral (lateral line) series large and bearing small spines from head to near posterior end of second dorsal; 4 or 5 pairs of plates between last ray of second dorsal and first median plate, and 5 pairs between last anal ray and first median plate; about 18 fairly large and tuberculated plates on breast, some of them separated by narrow intervals of membrane with numerous prickles. Branchiostegal membrane naked posteriorly. Under side of mandible and anterolateral parts of branchiostegal membrane with minute plates or prickles; 3 or 4 large spinous plates in front of, and a number of small plates and prickles on base of pectoral fin; the ventrolateral series separated, from ventrals nearly to anal, by membrane studded with 6 or 8 small striated plates behind and a few around vent; the latter a little farther than usual from base of ventrals (as in *O. dodecaëdron*). Top of head, dorsal, and 2 adjacent faces of body granulated; space behind pectoral as far back as its tip naked. Head depressed, elongate, broad behind, its depth $\frac{3}{4}$ of its width, the latter about 6 in length of body. Orbits large, oval, the longitudinal diameter longer than snout, $3\frac{1}{2}$ in head. Interorbital space wide, nearly equal to orbit, flat between supraocular ridges; the latter spineless, turning out almost at right angles, forming a thin saw-like transverse ridge along

* Diagnosis: Body moderately elongate, tapering, 8-hedral, depressed; depth about $\frac{3}{4}$ width, the latter 6 in length; head broad, much depressed, tapering by concave outlines to a blunt snout, $4\frac{2}{3}$ in length of body. Plates in dorsal series 35 or 36; between occiput and first dorsal, 7; between dorsals, 4; from ventral to anal, 10 or 11 pairs. One barbel at tip of each maxillary; gill membranes united, free from isthmus. Median rostral plate wanting; nasal spines minute; no supraocular or occipital spines. Olivaceous, banded with darker; pectorals with 3 or 4 black areas.

posterior border of orbit; occipital ridges spineless, temporal ridge broken into 2 or 3 tubercles, sometimes ending in a small suprascapular spine. Cheeks nearly vertical, the suborbital ridge but slightly developed; usually a spine at lower border of orbit, 3 or 4 small ones on inferior border of preorbital, a moderate one on suborbital, a rather large one, and beneath it, a smaller one on the preopercle; 3 or 4 moderately developed plates, or as many smaller ones with numerous minute plates or prickles, on cheek below ridge; nasal spines minute, converging behind maxillary pedicles; median rostral plate absent. Mouth oblique, a little less so than in *O. dodecaëdron*, the lower jaw protruding; maxillary reaches to opposite front of pupil. Teeth present on jaws, few and small on vomer, few or none on palatines. One barbel at tip of each maxillary. Dorsals long, moderately separated; anal long, beginning under posterior end of first dorsal; pectorals $4\frac{1}{3}$ in length of body, their width at base 3 in their length, are rounded, the lowermost rays rapidly shortening; ventrals of female a little less than 2 in pectorals, those of male much longer, reaching beyond front of anal, longer than pectorals; the membrane broad. Caudal 3 times as long as wide at base. Color dark grayish or brownish, pale below; the young with distinct cross bars, a broad one in front of first dorsal, 1 under the front end and 1 behind middle of first dorsal, 1 between dorsals, 1 under second dorsal, and 2 on peduncle; bars indistinct in adults; dorsals dusky, faintly barred with darker, caudal dark; anal dark on membrane, white-edged (dark only on posterior end in young, as in *O. dodecaëdron*); pectorals washed with orange, 2 dark blotches near base, the terminal part with 2 (sometimes confluent) or 1 large dark patch (these patches in the young appearing as partially confluent bars, more like *O. dodecaëdron*); ventrals of female pale, of male bright orange yellow, with a large black spot inside of first ray and 2 small ones at tip. Length 8 inches. Coast of California and Oregon, south to San Francisco, in 11 to 36 fathoms. Here described from specimens from off the Oregon coast. (*verrucosus*, warty.)

Brachyopsis verrucosus, Lockington. Proc. U. S. Nat. Mus., III, May 6, 1880, 60, Drakes Bay, near San Francisco, California; Jordan & Gilbert, Synopsis, 726, 1883; Jordan, Cat. Fishes N. A., 114, 1885.

Agonus (*Brachyopsis*) *barkani*, Steindachner, Ichth. Beiträge, IX, 253 (16), pl. 5, Sitzb. der k. Akad. der Wiss., LXXXII, July 15, 1880, San Francisco, California.

2408. OCCA DODECAEDRON* (Tilesius).

Br. 6; D. IX or X, 7 or 8; A. 14 to 16; P. 14 or 15; V. 3 (I, 2); C. 1, 11, 1; vertebræ 13 + 26 = 39. Pyloric cæca 5 (1 individual). Lateral line with 37 or 38 pores. Body as far as middle of first dorsal depressed, becoming gradually compressed behind. Caudal peduncle short, 5 to 7 dorsal and

* Diagnosis: Body moderately elongate, tapering nearly uniformly from head to caudal, depressed, 8-hedral; peduncle 6-hedral, compressed posteriorly; depth of body $\frac{3}{4}$ of width, the latter at base of pectorals 6 in length. Head broad, much depressed, $4\frac{1}{4}$. Plates in dorsal series 40; between occiput and first dorsal 8 or 9; between dorsals 3 to 5; from ventrals to anal 13 to 15 pairs. One barbel at tip of each maxillary; gill membranes free. Median rostral plate wanting; nasal spines minute, usually absent; no supraocular or occipital spines. Brownish olivaceous, with several indistinct cross bars; dorsal fins with 2 black bars.

3 or 4 ventral median plates, compressed, hexagonal. Short blunt spines present on the dorsolateral series from a little in front of first dorsal to middle of second dorsal, none on median dorsal plates of peduncle; sharper spines on the superior lateral series from near head to near caudal; the very low ridges of the inferior lateral and ventrolateral series bear a few spines near middle of body; plates of the median lateral (lateral line) series nearly as large as those of the adjoining series, about a dozen of the most anterior bearing small spines; 5 or 6 pairs of plates between last ray of second dorsal and first median plate, and about 5 pairs between last anal ray and first median plate; a strong sinuous ridge of plates from lower end of base of pectoral to isthmus; a small median ridge on breast, of 2 small plates in front of bases of ventrals, followed anteriorly by a median series of 3 or 4 larger single plates; between this and the edge the breast is covered with small, thin, distinct plates with numerous small tubercles interspersed; the branchiostegal membrane posteriorly and medially and anterolaterally covered with small plates and prickles; whole underside of mandible with small plates and some prickles or tubercles; 4 or 5 large plates in front of, and a number of minute plates or prickles on base of pectoral fin. Membrane behind and around vent and between vent and ventrals with a variable number of small plates, but mostly with numerous prickles or tubercles. Vent a little farther removed from base of ventrals than usual in the Agonoids. An elongate triangular space behind pectoral, between the inferior and ventrolateral series, naked. Top of head, median dorsal, and 2 adjacent faces of body granulated. Head much depressed, broad behind, tapering, its depth $\frac{2}{3}$ its width, the latter about 6 in length of body. Orbits small, nearly circular, the longitudinal diameter a little more than 4 in head; interorbital space concave, wide, equal to snout, nearly equal to orbit; supraocular, occipital, and temporal ridges low, spineless; cheeks nearly vertical, suborbital ridge hardly developed, spineless; a large, heavy tubercle or spine projecting from preopercle at an angle of 45°; below this 1 or 2 flat small ones; no plates on cheeks; mouth very oblique, maxillary reaching front of pupil; teeth in bands on jaws, vomer, and palatines. Dorsals long, their membranes sometimes continuous; anal very long, beginning a little behind middle of first dorsal; pectorals a little more than 4 in length of body, their base $3\frac{1}{2}$ in their length, with 14 or 15 graduated rays (a few of the uppermost shorter); ventrals of female about $\frac{7}{10}$ those of male; the latter less than 2 in pectorals, about $\frac{5}{8}$ of head; caudal 3 times as long as wide at base. Color brownish olivaceous, pale below; a much-interrupted, indefinite dark band from angle of mouth, under orbit, across preopercles, above pectorals, and along lateral line; a few indistinct cross bars on back; dorsals with a broad dark band along the edge and a parallel one across the middle, otherwise white; caudal dusky; anal dusky near the posterior end; pectorals transversely barred by about 6 series of black points or short streaks on rays; ventrals pale in female, the membrane black in male; head and lips with numerous minute black spots and streaks. North Pacific, Kamchatka, south to Bristol Bay, coast of Alaska, in 10 to 15 fathoms. (Gilbert.) Here

described from specimens from Bristol Bay. (δωδεκάεδρος, having 12 surfaces or sides.)

On these specimens Dr. Gilbert has the following notes:

"A few specimens taken in Bristol Bay in 4½ to 11 fathoms. In males the colors are much brighter than in females, recalling *O. verrucosa*. The bars on dorsal fins are, in males, intense black and bright white, instead of olive brown and whitish as in females; males show also a larger black patch on last anal rays, and have the interradial membrane of ventral jet-black; in none of our specimens does the caudal fin show transverse lines of brown points, as described by Cuvier; in both males and females the caudal is dusky or black, the median rays lighter, the outer ones white. D. IX or X, 7 or 8; A. 14 to 16." (δώδεκα, twelve; ἕδρα, surface, side.)

Agonus dodecaëdron, Tilesius, Mém. Acad. Petersb., iv, pl. 13, 1810, Kamchatka (Coll. W. T. Tilesius); Günther, Cat., ii, 214, 1860.

Phalangistes loricatus, Pallas, Zoog. Rosso-Asiat., iii, 114, taf. 19, 1811, Kamchatka.

Aspidophorus dodecaëdrus, Cuvier & Valenciennes, Hist. Nat. Poiss., iv, 209, 1829.

Brachyopsis dodecaëdrus, Jordan & Gilbert, Synopsis, 723, 1883.

764. BRACHYOPSIS, Gill.

Brachyopsis, Gill, Proc. Ac. Nat. Sci. Phila., xiii, 1861, 167, 259 (*rostratus*).

Siphagonus, Steindachner, Ichth. Beiträge. v, 140, Sitzb. Acad. Wiss. Wien, lxxiv, July, 1876 (*segaliensis*).

Body moderately elongate, tapering nearly uniformly from head to caudal; depressed, 8-hedral (6-hedral on peduncle); depth about 8, width about 6 in length; head broad, depressed, about 4½ to 5 in standard length. Snout long, almost tubular, bearing the short jaws at the end. Plates in dorsal series about 35 to 40 or more; a barbel at tip of each maxillary; median rostral plate none; nasal spines minute or absent; supraocular and occipital spines none; gill membranes united, free from isthmus; anal fin long, with 12 or 13 rays, first dorsal usually long; mouth oblique, lower jaw projecting; teeth present on jaws, vomer, and palatines; at least some of the plates on body spinous; plates on breast usually with interspersed small prickles or tubercles. (βραχύς, short; ὄψις, face.)

a. Anal rays 13; dorsal spines 8; body fusiform, broad at the breast, the tail very slender. ROSTRATUS, 2404.

aa. Anal rays 12; body slender; eye behind middle of head; 2 spines on suborbital. SEGALIENSIS, 2405.

2404. BRACHYOPSIS ROSTRATUS* (Tilesius).

Br. 6; D VIII, 8; A. 13; C. 10; P. 14; V, I, 2. Body more elongate than in *Agonus cataphractus*, the tail very slender, the body distinctly fusiform.

* Diagnosis. Lateral line 36. Body more fusiform than in other species of this genus, very robust in front of the middle, tapering to the slender tail. Head triangular and pointed, as seen from above, the mouth quite narrow; eyes small, well forward; head considerably depressed, snout without spine; a short flap at angle of mouth; scales not very rough; gill membranes free from isthmus; ventrals quite short; breast with a median row of raised convex plates; a row of similar plates bordering edge of gill opening, the three series forming a Λ-shaped figure, the interstices filled with very small plates. This diagnosis from Bering Sea specimens obtained by Dr. Bean. (Jordan & Gilbert.)

widest between pectorals; mouth short oblique, the maxillary not reaching to below eye; profile straight, cranium flat above, the eyes prominent; no spines on the orbit; preopercle with 3 spines on its vertical margin; 2 on the lower border of the large suborbital; trunk 8-angled; tail 6-angled. Plates more numerous than in *Agonus*, 40 on each upper range from nape to point of their union; numbers in the other rows in proportion; upper series with their ridges rather sharp, lower with obtuse ridges; 2 series of abdomen separated by dilatable skin; no barbels on gill membranes. Pectorals longer and dorsals farther back than in *O. dodecaedron* or *Podothecus acipenserinus*. Lower rays of pectoral larger than the others; anal longer than second dorsal, beginning under middle of first dorsal; length 10 inches. (Tilesius, as quoted by Cuvier & Valenciennes.) The figure of Tilesius shows a tubular compressed snout, with the short mouth at the end and a very wide back, mesially concave. Specimens from Petropaulski have the eye $2\frac{1}{3}$ in snout; mandible $2\frac{1}{8}$ in head; maxillary $4\frac{3}{5}$; snout 3; pectoral nearly as long as head; breast with large plates in 3 dividing rows, with smaller ones between. Tail very slender, not spinous; body chubby, the greatest width $2\frac{1}{4}$, greatest depth $2\frac{1}{8}$ in head; head nearly 5 in body. Scales 44. Teeth minute. The species is intermediate in form between *Occa dodecaedron* and *Pallasina barbata*. Several specimens from Shana Bay, Iturup Island, show the following characters: Dorsal face wider than in either and deeply concave; snout elongate, depressed, its width taken at middle of its length $\frac{1}{2}$ greater than its depth at same point, and $\frac{1}{2}$ its length, measured from tip of lower jaw; lower jaw much longer than upper, the symphysis entering upper profile of snout, vertically furrowed at tip; maxillary not reaching orbit, 4 in head; preorbital elongate, with a lengthwise ridge which divides anteriorly, the branches not terminating in spines, the edge of preorbital entire; anterior nostril in a short tube. Teeth all minute, present on jaws and vomer, often absent on palatines, sometimes present in a small patch on extreme anterior end. Suborbital stay without spine, forming a gibbous striated protuberance on middle of cheek, between which and the horizontal edge of preopercle is a series of three or four small plates; 2 strong diverging spines at angle of preopercle; a shorter spine below them; orbital margins elevated superiorly and posteriorly; interorbital space very narrow, grooved and longitudinally striated, its width equalling $\frac{1}{2}$ diameter of orbit, which is 6 in head; no spines on top of head, the ridges low and rounded. Head $4\frac{3}{5}$ to $4\frac{5}{6}$ in length; width of body $8\frac{1}{2}$ to $8\frac{1}{2}$; length of caudal peduncle $3\frac{1}{2}$ to 4. Body anteriorly hexagonal, the upper lateral ridge becoming obselete immediately in front of spinous dorsal; lower lateral ridge also becoming rounded and obsolescent anteriorly; dorsal face widening rapidly from occiput to front of spinous dorsal where its width equals snout; it gradually narrows posteriorly, the dorsal ridges becoming confluent at a point much nearer base of caudal than end of second dorsal; ventral ridges spineless, the lateral ridges with short spinous points, often distinguishable with difficulty; dorsal series anteriorly with stronger spines which rapidly diminish posteriorly; branchiostegal and gular membranes without plates; plates on body without the minute prickles

so characteristic of *Occa verrucosa* and *O. dodecaedron;* breast covered with polygonal plates, a series elevated to form a median ridge, the marginal plates also prominent; prepectoral area wide, with 4 prominent plates, the uppermost bearing a short spine posteriorly; in the dorsal series of plates, 10 lie in advance of first dorsal, 11 between origins of first and second dorsals, 9 or 10 along base of second dorsal, 6 to 9 between second dorsal and the point of confluence of the dorsal series, and 5 to 7 between the latter point and base of caudal; total number of plates in dorsal series 43 to 45, in 6 specimens examined. Pectorals long and narrow, 1¼ in head; dorsal with 8 (rarely 9) spines and 8 soft rays; anal with 13 (rarely 14) rays; pectoral with 14 rays. Color dusky above, marked with small black spots and lines; white below, growing dusky posteriorly; caudal blackish; ventrals white; anal white, with the last rays dusky; dorsals and pectorals with the rays finely dotted with black. North Pacific, recorded from Saghalien, Gulf of Aniva, Petropaulski, and the Kuril Islands. (*rostratus*, pertaining to the *rostrum* or snout; "not because it has a beak, but because its head and snout are more contracted than in the others.")

Agonus rostratus, TILESIUS, Mém. Acad. Petersb., IV, 1810, pl. 14, Saghalien, Gulf of Aniva. (Coll. Tilesius.)

Phalangistes fusiformis, PALLAS, Zoog. Rosso-Asiat., III, 116, 1811, Saghalien, Gulf of Aniva; Kuril Islands. (Coll. Steller & Merk.)

Agonus rostratus, GÜNTHER, Cat., II, 214, 1860.

Aspidophorus rostratus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 212, 1829.

Brachyopsis rostratus, JORDAN & GILBERT, Synopsis, 726, 1883; JORDAN, Cat. Fishes N. A., 113, 1885.

2405. BRACHYOPSIS SEGALIENSIS (Tilesius).

D. VII, 8; A. 12; C. 10; P. 14; V. I, 2. Form resembling *Brachyopsis rostratus*, but the tail shorter; body depressed; eye behind middle of head; 2 spines on suborbital, and some others about eyes; dorsals contiguous; anal longer than second dorsal; pectorals large; no barbels under throat; gill membranes and barbels at chin undescribed, probably as in *Brachyopsis rostratus*. Yellowish brown; fins with black bands. Length 7 inches. Island of Saghalien. (Cuvier & Valenciennes.) Not seen by recent writers. (Name from Saghalien.)

Syngnathus segaliensis, TILESIUS, Mém. Soc. Imp. Nat. de Moscow, II, 216, pl. 14, 1810, Bay of Patience, Saghalien. (Coll. Krusenstern.)

Siphagonus segaliensis, STEINDACHNER, Ichth. Beiträge, V, 140, and Sitzb. der k. Acad. der Wiss., LXXIV., 1876; JORDAN & GILBERT, Synopsis, 723, 1883.

Agonus lævigatus, TILESIUS, Mém. Acad. Petersb., IV, 436, 1810, Saghalien; CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 214, 1829.

Phalangistes lævigatus, PALLAS, Zoog. Rosso-Asiat., III, 116, 1811.

765. PALLASINA,* Cramer.

Pallasina, CRAMER, Proc. Cal. Ac. Sci. 1895, 815 (*barbata*).

Form of *Syngnathus* or *Siphostoma;* body slender, depressed; 4-hedral anteriorly, 8-hedral under dorsals; 6-hedral on peduncle; snout produced

* The genus *Siphagonus*, Steindachner, was expressly based on *Agonus segaliensis*. The name can not be used for this group.

in a tube; lower jaw projecting beyond upper, turned upward at tip, a long barbel at the symphysis; teeth on jaws and vomer, a single row on palatines; gill membranes free from isthmus, united; both dorsals present; ventrals very short; plates of body slightly keeled, without spines; vertebræ about 45. (Named for Petrus Simon Pallas, naturalist and explorer, the accomplished author of Zoographia Rosso-Asiatica, 1811.)

a. Three plates in front of ventrals; barbel long. BARBATA, 2406.
aa. Two plates in front of ventrals; barbel short. AIX, 2407.

2406. PALLASINA BARBATA (Steindachner).

Br. 5; D. VI to IX, 7; A. 9 to 12; P. 12; V. 3 (I, 2); C. 11; lateral line 45 or 46; vertebræ 15 + 30 = 45. Pyloric cæca 6 (1 individual). Body slightly depressed in front, depth at base of pectorals $\frac{5}{6}$ or $\frac{7}{7}$ of width, slender, width about 11 in length. Ridges of the dorso and ventro lateral series strong, the dorsal and lateral halves of the plates form a right angle; no ridge on the inferior lateral row and the plates of the superior lateral row absent anteriorly, so that the body is 4-hedral in front of first dorsal, with dorsal and ventral faces flat or a little concave and the lateral convex. Under first dorsal, the superior lateral series begins with keeled plates, the ridge of the inferior lateral series becoming more prominent, so that under the dorsals the body is 8-hedral; dorsal and ventral faces grooved, and depth equaling length. Caudal peduncle strongly depressed, nearly 4-hedral (median dorsal and ventral ridges extremely low); 45 or 46 plates in the dorsal series, 4 pairs between dorsals, 12 to 13 pairs from ventrals to anal, 5 to 9 pairs between last ray of second dorsal and first median ventral plate; 3 or 4 large plates in a median longitudinal row on breast, with about 1 row of 4 or 5 small ones between it and the series forming the edge of breast; plates radially striated and a little elevated at the center; none between ventrals and vent. Membrane behind vent without plates (or as many as 19, Steindachner's excellent figure). Branchiostegal membrane naked posteriorly, 2 or 3 plates anterolaterally. Narrow nude surface of lower jaw with a series of several plates; 2 or 3 plates in front of pectoral. Head very long and narrow, gently tapering, nearly as high as wide. Orbits nearly circular, the longitudinal diameter about $5\frac{1}{3}$ in head and 2 in snout. Interorbital space moderately concave, nearly 2 in orbit. Supraocular ridges moderate, occipital ridges scarcely defined, temporal ridges moderate, all spineless; no suborbital ridge; suborbital bone spineless; a sharp spine at posterior angle of preopercle and 2 smaller flat ones below this; a longitudinal series of 4 or 5 poorly developed plates on lower part of cheek, between the long horizontal limb of preopercle and orbit. Snout long, tubular, about $2\frac{1}{2}$ in head. Frontal bones much elongated forward, an additional bony plate in front of preorbital and overhanging the maxillary; several small plates in membranaceous interval between preorbital and frontal in front of orbit. Posterior inferior angle of maxillary produced backward, reaching a little more than halfway to orbit. Median rostral plate absent. Lower jaw long, curved upward in front, projecting beyond the upper and entering profile; mouth oblique. Teeth in narrow bands on jaws and vomer, about 1 row on pal-

atines (Steindachner apparently saw none on palatines). A single barbel of variable length (sometimes twice as long as, sometimes less than diameter of orbit) at tip of lower jaw. Gill membranes united behind, free from isthmus. First dorsal with 12 pairs of plates between it and the occiput; anal long, beginning under middle of first dorsal; caudal about 3 times as long as wide at base; pectorals long, 5¾ to 5⅘ in body, width at their base about 4 in their length; ventrals about 3 in pectorals and 4 in head in female, 2 in pectorals and 2⅜ in head in male. Color reddish or grayish brown with innumerable minute black spots; ventral surface pale, from front of anal to caudal progressively darker with minute black spots; a dark band extends along the side of snout, across orbit and preopercle, and along the side of body, dying out posteriorly; anterior dorsal dusky, darker behind; second dorsal indefinitely cross-banded with short streaks of darker on rays; caudal dusky; pectoral pale, indefinitely banded with short streaks of darker on rays; ventrals pale in both sexes; anal pale. Vertebral column (of a single specimen) with 15 abdominal and 30 caudal (including hypural) vertebræ. North Pacific, south to Japan and Oregon. The specimens here described from Bristol Bay (Coll. *Albatross*); Japan (Steindachner); Arctic Ocean near Bering Strait (W. J. Fisher, Steindachner); Bering Sea, Bristol Bay * (Gilbert); (Tareinsky Bay (Barrett-Hamilton); Port Clarence, Alaska (Scofield).† A variable species. Length of longest specimen examined by us 5½ inches. (*barbatus*, provided with a barbel.)

Siphagonus barbatus, STEINDACHNER, Ichth. Beiträge, v, 140, taf. 5, Sitzb. der k. Acad. der Wiss., LXXIV, July, 1876; JORDAN & GILBERT, Synopsis, 725, 1883; JORDAN, Cat., 113 1885.

2407. PALLASINA AIX, Starks.

Head from tip of snout, 4 in body; depth 3⅓ in head. D. VII, 7; A. 11; P. 11; eye 5⅓ in head. Mouth rather oblique, the lower jaw much project-

* Dr. Gilbert notes specimens from numerous stations in Bristol Bay in 4½ to 70 fathoms. The dark band on sides is often very strongly marked; the barbel varies much in length. D. VII to VIII, 6 to 8; A. 9 to 12.

† Mr. Scofield gives the following note on 6 specimens taken at Port Clarence, Alaska, averaging about 6 inches in length. We have compared them with specimens of *P. barbata* from Bristol Bay, Alaska, and find our specimens have a much longer barbel and slightly longer pectorals. In all other points they appear to be the same. The barbel is ⅓ distance from tip of lower jaw to edge of gill membrane. The 2 dorsals vary in their distance from each other. In 1 specimen they touch, in the others they vary in distance the width of 1 or 2 plates. Following is the fin formula of the specimens. These specimens show a sexual difference. The females have shorter ventrals and a lower smaller first dorsal than the males. The first dorsal, too, is without color. The males have larger ventral fins and a large darkly colored first dorsal:

Dorsal.	Anal.	Pectoral.	Sex.	Remarks.
V, 7	10	12	♀	
VIII, 7	10	12	♂	
VI, 7	8	12	♀	The anal injured.
VII, 7	9	12	♂	
VIII, 6	10	12	♂	
V, 6	10	13	♀	

ing; upper edge of maxillary slipping under preorbital for its whole length, its posterior end midway between tip of snout and middle of eye; villiform teeth on jaws and vomer, none apparently on palatines; lower jaw with a fleshy tip, scarcely produced into a barbel, scarcely as long as diameter of pupil; supraorbital rim prominent, making the interorbital space deeply concave; width of the latter about $\frac{2}{3}$ eye; 2 ridges from the inner edges of supraorbital rim run backward, and are continuous with dorsal ridges of body; edge of preopercle with 3 spines, the middle one the largest. Two large median plates in front of ventrals on breast, a row of plates along each lateral ridge of breast, a large plate on each side of the union between first and second median plates, and behind them a row of small plates irregular in size and position, sometimes continuous and interposed between median and lateral plates, and sometimes allowing the edges of median and lateral plates to touch; 11 or 12 plates in front of dorsal; spinous dorsal on 9 or 10 plates, counting to end of membrane; lateral line 43. Space between dorsal ridges strongly concave, the ridges coming together on caudal peduncle, but not uniting, continuing parallel for a short distance and then becoming obsolete; upper lateral ridge ending anteriorly on about the twelfth plate from head. Last rays of dorsal and anal connected to the body by a membrane; space between dorsals about equal to the width of a plate; front of anal midway between posterior end of maxillary and base of caudal; pectoral reaching just past front of spinous dorsal; vent distant an eye's diameter from base of ventrals; length of caudal equal to head behind anterior ridge of pupil. Color blackish, with fine punctulations; belly white; a light streak running backward from eye to upper edge of gill opening; below this an area scarcely so wide as eye, darker than the rest of body, its lower edge sharply defined against the white under parts of head; chin black; dorsals and caudal dusky; pectorals light, the rays with many black spots which do not involve the membrane; ventrals and anal white. This species differs from *Pallasina barbata* in having a much shorter barbel on chin, in having 2 median plates in front of ventrals in place of 3, and in having the plates between the median and lateral plates much smaller and less regular in arrangement. In *P. barbata* these plates are about as large as the median plates and always interposed between them and lateral plates, the arrangement being constant. The abdominal ridges are generally closer together in *P. aix* than in the northern species. Puget Sound to the Aleutian * Islands; taken with the seine in great abundance in Puget Sound, near Port Ludlow; the largest nearly 5 inches in length, the others about 3. (αἴξ, a goat, from the pointed beard, or ἄξ, a darter, from the slender form.)

Pallasina aix, STARKS, Proc. Cal. Ac. Sci. 1896, 558, pl. 75, Puget Sound, near Port Ludlow, Washington. (Type, No. 5040, L. S. Jr. Univ. Mus. Coll. E. C. Starks.)

* Two small specimens were taken by Dr. Gilbert at Chignik Bay, Alaska. Concerning these he remarks: "We have compared these specimens with the types of *Pallasina aix* and can find no good differences. One of our specimens has 3 median plates in the front of the ventrals, the other 2. (Our specimens of *P. barbata* have either 3 or 2 median plates in front of the ventrals.) A few of the types have 12 pectoral rays, but the majority have 11. Fin formula: D. VII, 6; A. 10; P. 12. D. VII, 6; A. 9; P. 12."

766. LEPTAGONUS, Gill.

Leptagonus, GILL, Proc. Ac. Nat. Sci. Phila., XIII, 1861, 167, 259 (*spinosissimus* = *decagonus*).

Body tapering, depressed; head about 5½; depth 7½ to 9; width 6½ to 8 in total length. Plates in dorsal series 40 to 44; 5 to 6 pairs between occiput and first dorsal; both dorsals present; mouth inferior, snout much projecting; teeth on both jaws, probably none on vomer or palatines; 5 pairs of barbels about mouth; none under tip of snout; 1 pair of recurved spines at tip of snout; 1 pair of supraocular and 1 pair of occipital spines. Gill membranes joined to isthmus. (λεπτός, slender; *Agonus*.)

2408. LEPTOGONUS DECAGONUS* (Bloch & Schneider).

Br. 6; D. V to VII, 5 to 8; A. 6 to 8; P. 14 to 16; V. I, 2; C. 24. Body distinctly more slender than in *A. cataphractus;* its greatest breadth, over base of pectorals, about 6½ to 8 times, and greatest height, at the third pair of dorsal plates, about 7½ to 9 in total length, hence but little depressed. Vent nearer base of ventrals than in *A. cataphractus;* about 2 pairs of plates from vent to ventrals (no plates medially between them), and 9 pairs from vent to anal fin. Distance from tip of snout to vent about 4 in total length. Body anteriorly nearly oval, angular; abdominal region relatively shorter and tail, relatively longer than in *A. cataphractus*. Body octagonal; tail hexagonal, depressed, its ventral keel quite prominent. Ridges on plates of body and their backwardly directed spines more prominent than in *A. cataphractus*. Keels of the superior and inferior lateral series especially sharp on tail; 5 or 6 pairs of plates from head to first dorsal, and 10 to 13 pairs between ventrals and anal; from anterior end of first dorsal to point of union of the dorsal pairs behind second dorsal about 22 pairs, thence to base of caudal 14 to 17 median plates. Head about as broad as deep, the breadth about 1½ in its length; head about 5½ or somewhat more in total length, much narrowed anteriorly. Snout projecting beyond premaxillaries, lower jaw included; a pair of short upwardly and backwardly directed spines near tip of snout; a pair of strong supraocular and 1 of occipital spines; interorbital space considerably less than longitudinal diameter of orbit; preorbital not covering upper jaw; suborbital without prominent spine; no barbels under tip of snout; mouth somewhat larger than in *A. cataphractus*, maxillary nearly reaching anterior edge of orbit; 5 pairs of barbels, 1 near and 2 at tip of maxillary, 1 near angle of mouth, and 1 (bifurcate) out on side on lower jaw; none on branchiostegal membrane. Orbits quite large, their longitudinal diameter about 4 in head. Preopercle with 2 quite small spines, 1 at its posterior angle, the other below this, and 1 or 2 smaller ones below these; opercle unarmed. Branchiostegal rays 6. Teeth fewer than in

*Diagnosis: Body tapering, octagonal anteriorly, hexagonal posteriorly, a little depressed, its height 7½ to 9, and its breadth 6½ to 8 in total length. Head 5½; pectorals equal to head. Plates in dorsal series 44, sometimes 41–43; from occiput to first dorsal 5–6; between ventrals and anal 10–13; between dorsals 6 or 7; 5 pairs of barbels, 3 on maxillary, 1 near angle of mouth, and 1 (bifurcate) on lower jaw; 1 pair recurved spines near tip of snout; 1 pair supraocular and 1 of occipital spines. Teeth on jaws. Gill membranes joined to isthmus. Yellowish gray, with 2 or 3 cross bands, pale below. B. 6; 1 D. 5–7; 2 D. 5 to 8; A. 6 to 8; P. 14 to 16; V. 3 (I–2); C. 2–9–11 + 2.

A. cataphractus, present on jaws. First dorsal, with 6 or 7 plates between it and occiput, with 5 to 7, usually 5 or 6, spines, its length considerably more than its height; third and fourth rays longest; second dorsal distant from first about ½ the length of either fin, with 5 to 8, usually 6 or 7, rays, a little shorter but somewhat higher than first dorsal; anal fin with nearly ½ its length in front of first ray of second dorsal and ending a little behind the middle of the latter is somewhat higher than this and with 6 to 8, usually 7, rays; pectoral fin, about as long as head, with 14 to 16, usually 14 or 15, simple rays; ventral fin short, a little more than ¼ of pectoral, rays 3 (I, 2), the 2 soft rays of about equal length; caudal narrow and long, about 8 in total length, with 9 long rays, and on each side 2 or 3 short ones. Lateral line, pores 23 to 25 (Collett), 30 (Krøyer). Color yellowish gray, with 2 or 3 large grayish brown patches forming indefinite cross bands, first above base of pectorals, second under posterior part of first dorsal, and third under middle of second dorsal; between these smaller indefinite patches and cloudings; fins, especially pectorals and caudal, brownish black toward their tips; ventral surface grayish yellow; on each side of head a quite broad black band from tip of snout across orbit and preopercle. Males rarer than females (Collett), distinguished by having outer soft ray of ventral fin about twice as long as inner, more than ½ of pectorals. Young (according to Steenstrup and Lütken) with a shorter and thicker body, its breadth about 7 in total length, tail compressed. Head broader and anteriorly blunter, with somewhat oblique mouth and lower jaw projecting beyond upper. Keels of plates strongly developed, with sharp backwardly directed spines; longitudinal keels on upper and under side of tail distinctly double. Pectoral and ventral fins proportionately longer. Vent lies farther back, between fifth or even eighth pair of ventral plates. Arctic Ocean, south to Newfoundland and Norway; recorded from northern coast of Norway, east and west of North Cape, but not south of 70° N. lat., Spitzbergen and west of these islands, between them and Beeren Island, and between this and Norway (Collett); Faröe Islands and Iceland (Lütken); Greenland, spreading as far south as Newfoundland with the cold ocean currents. Dr. Reinhardt has shown that the fish described as *Cottus cataphractus* from Greenland by Fabricius belongs to this species, and that Bloch's statement that the type of the species came from the East Indies is erroneous. It is a deep-water species, found in 123 to 260 fathoms, at a temperature of + 1.6° C. to —1.1° C. (Nordhavs expedition); 50 to 120 fathoms, in the Varangerfiord. Here described from papers of Collett and Lütken, and from small specimens from Upernavik. (Eu.) (*δέκα*, ten; *γωνία*, angle.)

Cottus cataphractus, Fabricius, Fauna Grœnl., 155, 1780, Greenland; not of Linnæus.

Agonus decagonus, Bloch & Schneider, Syst. Ichth., I, 105, pl. 27, 1801, erroneously recorded from the East Indies, the type came from Greenland; Günther, Cat., II, 215, 1860; Collett, Norges Fiske, 40, 1875; Lütken, Forelöb. Meddel. Nord. Ulkefiske; Vidensk. Meddel. fra den Naturhist. Foren. Kjöb., 381, 1876; Collett, Fiske fra Nordhavs Expeditionens, Sommeren 1878, Christ. Vidensk. Selsk. s. Forh., No. 14, p. 28, 1878; Collett, Norske Nordhavs Expedition, 1876-1878, Zool. Fiske, 44, pl. 2, figs. 11-12, 1880; Jordan, Proc. Ac. Nat. Sci. Phila. 1883, 293; Lilljeborg, Sveriges och Norges Fiskar, Första Häftet, 193, 1883–84.

Aspidophorus spinosissimus, KRÖYER, Ichth. Bidrag. Naturhist. Tidskr., I, 250, 1844–45, Greenland; KRÖYER, Gaimard, Voyages en Scand., etc., Zool. Atlas, Poiss., pl. 5, figs. 2a–2d, 1845.

Aspidophorus malarmoides, DES LONGCHAMPS, Mém. Soc. Linn. de Normandie, tome IX, 167, 1853.

Aspidophorus decagonus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 223, 1829; J. REINHARDT, Sr., Overs. over det. Konigl. Danske Vid. Selsk. s. Forh., 53, 1830–1832; KRÖYER, Ichth. Bidrag. Naturhist. Tiddskr., I, 243, 1844–45; KRÖYER, Gaimard, Voyages en Scand., Lapnonie, etc., Zool. Atlas, Poiss., pl. 5, figs. 1a–1d, 1845.

Agonus spinosissimus, GÜNTHER, Cat., II, 214, 1860.

Leptagonus spinosissimus, GILL, Proc. Ac. Nat. Sci. Phila., XIII, 1861, 167.

Brachyopsis decagonus, JORDAN & GILBERT, Synopsis, Fishes N. A., 727, 1883.

767. PODOTHECUS, Gill.

Podothecus, GILL, Proc. Ac. Nat. Sci. Phila. XIII, 1861, 77, 259 (*peristethus*=*acipenserinus*).

Paragonus, GILL, Proc. Ac. Nat. Sci. Phila. XIII, 1861, 167, 259 (*acipenserinus*).

Body tapering nearly uniformly from head to caudal, about as high as or higher than wide anteriorly; depth about 6 to 8 inches in length; head about 4, compressed; plates in dorsal series about 35 to 40, 4 or 5 pairs between occiput and first dorsal fin; gill membranes joined to isthmus, without free fold; no scattered barbels under lower jaw or on branchiostegal membranes; 2 complex groups of barbels on under side of tip of snout, another group at each angle of mouth; usually a few barbels at sides of pores under lower jaw; tip of snout with usually 2 pairs of sharp slender spines, the anterior directed forward, the posterior outward and backward. Mouth small, inferior, snout projecting far beyond it. Teeth on both jaws few and weak, sometimes wanting on 1 or both jaws; none on vomer and palatines; plates of body spinous; 1 pair of supraocular and 1 pair occipital spines; both dorsals present. This genus is very close to *Agonus*, differing mainly in the spinous plates of the body. The numerous species differ much among themselves. (*πους*, foot; *θηκη*, box; from the groove for the receptacle of the ventrals which appears through the shrinking of the naked skin in preserved specimens.)

a. Plates on caudal peduncle all or nearly all armed each with a spine.

b. Barbels below snout very numerous and large; angle of mouth with many barbels. ACCIPITER, 2409.

c. Dorsal rays VIII–9; teeth well developed; fins all very high, the pectoral emarginate in the adult, its longest rays 1¼ in head; ventral long; anal rays 10.

cc. Dorsal rays XI–8; teeth present; fins moderate; pectoral not emarginate; ventral short; anal rays 9. HAMLINI, 2410.

bb. Barbels comparatively few and slender.

d. Dorsal rays VIII–8; teeth in upper jaw almost obsolete; fins moderate; pectoral not emarginate, 1⅓ in head; ventral very short; anal rays 9. GILBERTI, 2411.

dd. Dorsal rays IX–6; fins small; ventrals long; sides and top of head with very large crests; anal rays 6. THOMPSONI, 2412.

aa. Plates on caudal peduncle mostly not ending in spines; fins rather low, the anal rays 7 or 8.

e. Teeth well developed; barbels below snout and at angle of mouth long and numerous. ACIPENSERINUS, 2413.

ee. Teeth wanting in the adult; barbels rather small and sparse; body not everywhere deeper than wide; the caudal peduncle very long, slender, and depressed. VETERNUS, 2414.

2409. PODOTHECUS ACCIPITER, Jordan & Starks.

Head $3\frac{3}{5}$ in length; depth $6\frac{1}{4}$. D. VIII, 9; A. 10; pectoral 15; lateral plates 36; eye $4\frac{1}{2}$ in head; snout $2\frac{1}{8}$; second dorsal spine $1\frac{5}{6}$; second dorsal ray $1\frac{1}{2}$; third anal ray $1\frac{3}{5}$; caudal $1\frac{4}{5}$; upper ray of pectoral $1\frac{1}{7}$; ventrals $2\frac{1}{3}$. Body elongate, not compressed; head triangular as viewed from above; the mouth wide, entirely inferior, ⌓-shaped, the lower jaw shutting behind the upper by a distance equal to $\frac{1}{2}$ eye; maxillary not reaching quite to anterior orbital rim; distance of anterior edge of upper lip from tip of rostral spines a little more than $\frac{1}{2}$ eye; teeth in upper jaw almost obsolete, villiform bands of teeth in lower jaw, wide in front, becoming narrow at sides; vomer and palatines toothless; a patch of thick barbels below snout in front of mouth, the longest equal to vertical diameter of eye, a similar patch at end of maxillary, about equal in length to the shortest on snout; 2 short barbels on each side of lower lip between symphysis and angle of mouth. A pair of short, sharp, rostral spines, pointing directly forward; at their base and much wider apart is a pair of spines which point upward, backward, and slightly outward; running backward from these are the ridges that bound the wide groove in which the maxillary process fits; these approach each other behind and end in sharp spines which point backward and upward, these spines midway between middle of eye and the spines behind rostral spines; a pair of large spines above posterior third of eye, and a pair of large ones at occiput, these continuous with the upper ridges; a curved bridge running from superior orbital rim and ending in a small spine just above opercle; a small ridge on opercle; preopercle with a large spine; a couple of spines below eye at lower edge of suborbitals; running from them to tip of snout a ridge along lower edge of preorbitals, somewhat irregular but without spines; interorbital space wide and deeply concave, a pair of ridges on each side, converging forward; supraorbital rim prominent; anterior nostril ending in a short, wide, conical papilla, with a small opening at the apex; no noticeable depression at occiput. Dorsal ridges converging from the occiput to behind the soft dorsal, uniting on the second plate behind the base of last dorsal ray, continued as a single ridge on about 8 plates where it becomes obsolete; the upper lateral ridge follows the course of the lateral line to about the middle of spinous dorsal, where it slants sharply upward and is continued to tail above lateral line; lateral line midway between upper and lower lateral ridges posteriorly; a single spine above base of pectoral indicating an obsolete ridge between the lateral ridges, lower lateral ridge becoming obsolete under pectoral on 2 or 3 plates behind its base; abdominal ridges widest apart behind base of ventrals, uniting directly behind anal base and running simply backward, becoming obsolete on caudal peduncle; all the ridges with sharp, recurved spines, with the exception of abdominal ridges behind part of anal; where the dorsal and anal ridges disappear the caudal peduncle assumes a quadrangular shape, the corners being framed by the spines of the lateral ridges; no row of spines around base of caudal or pectoral. Fins all high, origin of dorsal between the fourth and fifth dorsal plates, the fin to base of last spine covering 6 plates, the

membrane covering 2½ more; the second and third spines the longest, a membrane connecting the last spine to the body for its whole length, when fin is depressed the ends of the last spines reach to the front of second dorsal; the second dorsal to the end of last ray covers 8 plates, the membrane covers 1 more; the second and third rays are the longest, the last ray is connected to the body for about ⅓ of its length; base of anal covering 8½ plates, the rays very long and not differing much in length, the last ray not connected to body by a membrane; the fin begins in front of soft dorsal but is about coterminous with it, its rays when depressed reaching past ends of soft dorsal, reaching 6 plates past base of its last ray; pectorals barely reaching to tip of last dorsal spine, the fin pointed above, first and second rays the longest, the lower rays produced beyond the membrane, making a notch in posterior outline of fin; origin of ventrals directly below base of pectoral, their tips reaching 6 plates beyond their base; caudal long and truncated; vent directly behind base of ventrals. Color light brown above, white below; back with many narrow brown bars placed at irregular distances apart; head with many blended brown spots, 1 under eye, 1 on front margin of eye, 1 or 2 on top of head, 1 behind eye, 1 on preorbital, a similar spot on base of pectoral rays; pectoral dusky; first dorsal with 3 rows of spots across the rays, a very narrow brown border to fin, second dorsal with similar spots, not arranged in rows; anal light above, uniform brown below; ventrals white; caudal fin dark at base and 3 or 4 dark spots toward middle of fin. Ochotsk Sea; one specimen collected at Robben Island by Captain Blair. It is 8 inches in length. (*accipiter*, a hawk; in allusion to the large fins.)

Podothecus accipiter, Jordan & Starks, Proc. Cal. Ac. Sci. 1895, 816, pl. 88, Robben Island. (Coll. Capt. J. G. Blair. Type in L. S Jr. Univ. Mus.)

2410. PODOTHECUS HAMLINI, Jordan & Gilbert, new species.

D. IX to XI-8; A. 9 or 10; P. 15. Head 3¾ in length; depth, not including spines of dorsal plates, 8⅔; width at base of pectorals 7⅙; length of caudal peduncle, from base of last anal ray, 2⅔; snout long and slender, depressed, produced beyond the mouth for a distance equaling a little less than ⅓ its length, the tip formed of 2 spines, the space between which is covered with membrane; a pair of strong, nearly erect spines at their base, between which are 2 or 3 very small spines on the median line; a second pair of strong spines at posterior end of premaxillary fossa, the ridges bounding which may bear 1 or more pairs of small prickles; preopercle with a wide wing-like crest terminating in a bluntish spine; snout long and slender, its lateral profile concave as seen from above or below; suborbital crest with 3 very strong, backwardly hooked spines; interorbital space very narrow, deeply concave; fins all low; supraocular ridge strong and much elevated, the interorbital space deeper and narrower than in any other species of this group, its least width 1⅐ in orbit; postorbital spine small; ridges on sides of snout minutely serrate; an irregular group of small spines above and behind anterior nostril; lateral ridges of head

with 3 pairs of very strong backwardly directed spines in line with the greatly expanded preopercular ridge; the posterior portion of this ridge produced into a compressed, bluntish process which overlaps the subopercle and reaches margin of gill opening; opercular and temporal ridges low and sharp. Teeth present on jaws in the young, becoming nearly or quite obsolete in adults; in the type, 17 cm. long, a few weak teeth on one side of upper jaw, but none elsewhere; vomer and palatines toothless. A cluster of 14 barbels on each side of lower surface of snout in front of mouth, and a cluster of 14 occupying end of maxillary and angle of mouth; each side of lower lip with 2 barbels; gill membranes widely joined to the isthmus, without distinct free fold posteriorly. Plates on body with very strong spines, those of the dorsal series the largest, but decreasing rapidly backward, becoming much smaller than the lateral series under the second dorsal fin; a median series of short sharp spines still persists along entire back of tail after the confluence of the dorsal series; the lateral series lower anteriorly, the upper row persisting to gill opening, the lower row now becoming obsolete about 4 plates behind axil; the abdominal series bear short sharp spines (longer in the young), becoming obsolete along anterior portion of anal fin; of the dorsal series, 4 are in front of the first dorsal, 10 (or 9½) along base of first dorsal, 1 (or 1½) between dorsals, 9 along base of second dorsal, 15 along back of tail, the first of the latter being the plate in which the dorsal series first becomes confluent; anterior portion of lateral line running along upper lateral series, gradually descending to middle of sides where it runs on a special row of plates which bear no spines; where the lateral line begins to ascend, 3 of these become confluent with corresponding plates of the upper lateral series; in advance of these, 5 members of the series again appear distinct, considerably enlarged and bearing spines; 40 pores in the lateral line; fins all comparatively short and low; pectorals reaching twelfth plate of upper lateral series, the longest ray equaling length of snout and ½ eye; rays becoming rapidly shortened below, the lower 5 or 6 slightly thickened with exserted tips; ventrals very short, not exceeding length of snout before mouth, not received into longitudinal groove. We consider it very doubtful whether such a groove exists in any of the other species of this group. It has been described as existing in the types of *P. gilberti* (Collett) and *P. peristethus*, Gill. In both cases the type specimens were in a poor state of preservation, and the groove was probably due to a softening of that longitudinal strip of the abdominal wall, which includes the anal opening, and extends backward from the base of the ventral fins and is interposed between the firm outer series of ventral plates. That such a softening had occurred in the type of *Podothecus peristethus* is evident from Gill's statement that the ventrals had dropped out. This view is rendered more probable from the fact that *P. peristethus* is apparently identical with the common *P. acipenserinus* which contains no such groove. We have also examined 2 of the type of *P. gilberti* without being able to satisfy ourselves of the existence of any special groove. The dorsal fins are closely juxtaposed, the interspace including 1 or 1½ pairs of plates. The base of the last ray of second dorsal is midway

between base of caudal and origin of spinous dorsal. Color dark or brownish above, with irregular spots or dashes of darker, which do not form definite cross bars; a black streak from eye to tip of snout passing onto lower side of rostral spines; a dark blotch on expanded limb preopercle; a black spot on base of middle pectoral rays, the fin very obscurely marked with dusky; dorsal spines and rays with linear dark markings, 1 or 2 black spots near tip of spinous dorsal anteriorly; under parts including fins, unmarked. Two specimens from *Albatross* Station 3653, off Shana Village, Iturup Island, in 18 fathoms. A young individual from *Albatross* Station 3646, off Robben Island, 18 fathoms, seems to belong to the same species, but has the snout less produced and the dorsal VIII, 6; anal 8. (We take great pleasure in naming this species in honor of Hon. Charles Sumner Hamlin, late Assistant Secretary of the Treasury, under whose auspices the fur seal investigations of 1896 and 1897 were carried on by the United States Fur Seal Commission.

Podothecus hamlini, JORDAN & GILBERT, Rept. Fur Seal Invest. 1896, 1897 MS., off Shana Bay, Iturup Island, Kuril Group. (Type, 5662, L. S. Jr. Univ. Mus. Coll. *Albatross.*)

2411. PODOTHECUS GILBERTI (Collett).

Head about 4; depth 7½ or 8. D. VIII, 8; A. 9; P. 17; V. I, 2. Body deeper than wide; octagonal to end of dorsal and anal fins, hexagonal on tail. The curved spines of the dorsolateral, superior and inferior lateral ridges strong and sharp as far as base of caudal. Ventral series of plates spinous to origin of anal, uniting behind it to form a single series; dorsal series uniting into 1 behind second dorsal; about 12 to 15 well developed, radially striated plates without tubercles or spines on breast; a few median plates between ventrolateral series behind vent. Head 4-hedral, tapering, cheeks vertical, orbit large, high up, about equal to the concave interorbital space. Supraocular and occipital ridges high, with crenulate edges, the former ending in a sharp, the latter in a blunt, spine; suborbital and opercular bones radially striated; suborbital ridge low down on cheek, thin, with crenulate edge and a small sharp spine at its posterior end; preopercle with a high thin keel with serrated edge, ending in a strong blunt spine; rostrum projecting far beyond premaxillaries, with a pair of spines at tip directed forward, another pair, farther apart, curved backward and outward. Maxillary reaching about to front of orbit; lower jaw very short, broad, falling far short of upper. Teeth on lower jaw weak, few and excessively weak on upper jaw, none on vomer or palatines. A group of barbels under tip of snout, another at angle of mouth (specimen with mouth in bad state). Dorsals separated by about 3 pairs of plates; anal begins a little in front of origin of second dorsal; pectorals about 6 in body, their base nearly 3 in their length; ventrals (male?) about ⅔ as long as pectorals. Lateral line about 35. Color apparently brownish or yellowish, with well-defined dark-brown spots on sides of head, back, and sides; the spots largest along sides, some of them as large as the osseous plates, smaller on back; apparently a dark band from front of orbits to tip of snout; dorsals with dark patches; a dark

spot on base of pectoral. Kamchatka. This account taken from 1 of the type specimens presented * to the Leland Stanford Junior University Museum by Dr. Collett. (Named for Charles Henry Gilbert.)

* The following is Dr. Collett's description: First dorsal 8; second dorsal 8 or 9; anal 10 or 11; pectoral 15 to 17; lateral line 38. The body is very elongated and compressed, rather high in front, and tapering to the tail. Its height (from ventrals to first dorsal scale) equals the length of the snout, and is contained in the total length about 8 times. The tail is compressed, long and slender; its height everywhere greater than its breadth. The head: Its length is contained in full-grown specimens 4 times, in younger ones about 3¾ times in the total length (caudal included). Thus in 4 specimens of different sizes the proportions are the following: Total length 183 mm., head length 3-66 times; total length 254 mm., head length 3-73 times; total length 277 mm., head length 4-01 times; total length 290 mm., head length 4-02 times. Snout very long, 3 times longer (or more) than the interorbital space (between the bases of the supraocular spines). Posterior part of the head comparatively smooth, the interorbital space rather concave; no quadrangular pit on the occiput in front of the dorsal scales. Eye comparatively large; the horizontal diameter a little larger than the vertical. It is contained a little more than 2 times in the length of the snout, and rather more than 4 times in the length of the head. Cirri on lower side of snout (in front of the premaxillary) and at the angle of the jaws.[1] Their length equals that of the eye. Mouth entirely inferior; distance from premaxillaries to tip of rostril spines about equal to the length of the eye. Teeth in the jaws, vomerine or palatine teeth none. Armature of the head much like that of *A. acipenserinus*. The rostral spines 4, 2 projecting horizontally forward, 2 (behind the first) curved backward. A third pair on the snout (much nearer the eye than the rostral spines). Orbital ridge with a single spine (supraocular, no preocular); the lower ridge finely serrated. Occipital ridges, operculum, and preoperculum as in *A. acipenserinus*, but the spines more pointed and longer. Suborbital with a double ridge at its lower margin, the upper ridge with 2 distinct spines behind, and a third (sometimes indistinct) at equal distance from the eye and the tip of the snout. Head with about 18 distinct spines altogether. Armature of body: Plates on the back and sides of the same number as in *A. acipenserinus*, but the spines are longer and curved more backward, and strong everywhere from head to caudal. Between the 2 dorsal keels and between the 2 lateral keels there are no traces of another keel (as in *A. vulsus*). Breast with about 10 polygonal plates, 4 of which form a series on each side and 2 a median series; bases of pectorals and ventrals also surrounded with plates. All the plates have a short spine in their center. Dorsal plates numbering—

		Plates.
From occiput to first dorsal	(pair)	4
First dorsal extending over	do	8
Between the dorsals	do	3
Second dorsal extending over	do	9+1
From second dorsal to caudal	(single)	15

The dorsal keel (coalescing with the keel on the other side at the fifteenth plate in front of the caudal fin) is consequently composed of 38 or 39 plates; the lower lateral keel extending from lower base of the caudal to base of the tenth pectoral ray, contains 35 plates. Abdominal plates numbering—

		Plates.
From ventrals to anal	(pair)	11
The anal extending over	do	10
From anal to caudal	(single)	17

The abdominal keel (coalescing at the seventeenth plate in front of the caudal) is formed by a series of 38 plates. Lateral line distinct; 38 pores. Fins: In the 10 specimens at present preserved in the museum at Christiania, the fin rays are the following:

1 D.	2 D.	A.	P.
8	9	10	?-16
8	9	10	16-16
8	8	10	15-16
8	8	10	17-17
8	8	10	17-17
8	8	10	16-17
8 (+1)	8	11	16-17
8	8	10	15-16
8	9	10	17-17

First dorsal begins behind the fourth dorsal plate and has 8 rays (1 specimen has an additional slender ray in the space between the 2 dorsals). Its height equals its distance from the head. It extends over 8 scales; the first 2 rays in the space between the fourth and fifth plate. The dorsal fins are separated by 3, sometimes by 4 plates. Second dorsal has 8, sometimes 9, rays, and extends over 8 plates; behind the last rays is 1 pair of plates

[1] All the specimens are in a bad state of preservation and most of the barbels are lost.

Agonus gilberti, COLLETT, Proc. Zool. Soc. London 1894, 670, pl. 45, Kamchatka. (Coll. Henry Lund. Types, Mus. Christiania; cotype, 2783, L. S. Jr. Univ. Mus.)

2412. PODOTHECUS THOMPSONI, Jordan & Gilbert, new species.

D. VIII or IX–6; A. 6; P. 16. Head rather broadly triangular, its greatest width across preopercular ridges greater than distance from anterior end of preopercular ridge to tip of snout. Lateral ridge on head continuous from tip of snout along suborbital bones to base of preopercular crest, the lateral spines usual in this genus, being represented by triangular processes borne on the ridge; preopercular ridge produced posteriorly beyond gill opening, but not spine like; snout terminating anteriorly in 2 rounded processes, each bearing on its upper surface a vertical crest, and finely serrate along its margins; no terminal pair of strong spines as in other species; under side of snout with an acute median spine directed downward and backward; a pair of strong spines on upper side of snout behind terminal nostral ridge; a pair of coalesced spines behind the nostral groove; a semicircular series of spinelets below the eye; two small tufts of filaments on under side of snout, one on middle of maxillary, and one at its tip. A narrow band of sharp teeth in each jaw; vomer and palatines toothless. Gill membranes united to isthmus without evident free fold. Orbital rim much elevated; interorbital space narrow, deeply concave, its width $\frac{9}{10}$ diameter of orbit; occipital ridges strong, elevated posteriorly, ending in a backwardly directed spine which is much larger than those of the body plates; occipital area narrow, deeply concave, its central portion sunk somewhat below level of interorbital space, from which it is separated by a shallow transverse groove; a similar groove behind occipital spines; area between occipital and temporal ridges also deeply concave; a strong opercular ridge. All the plates with strong spines, including those of the ventral series; the weakest spines are on the anterior plates of the lateral series; plates on breast with central spine and radiating ridges; dorsal series with fewer plates than in other

before the unpaired series begins. Anal has commonly 10 rays (in 1 specimen 11); its height equals that of the second dorsal, and is rather less than that of the first dorsal. It commences between the eleventh and twelfth pairs of scales in the abdominal series. Ventrals short in the females, shorter than the vertical diameter of the eye; longer in the male, equalling the length of the snout. Each has 1 short spine and 3 articulated rays, 2 of which are divided at their base. They are received in a longitudinal common groove ("*Podothecus*"). Pectoral has 16 or 17, rarely 15 rays, some of which are sometimes branched in their upper half, but not always. The first ray is short, about equal to $\frac{1}{2}$ the second ray; the lowermost rays a little thicker than the rest. Its form is a little emarginate, the fifth lower ray being a trifle longer than the sixth and seventh. The tip extends to a distance from the anal of 2 or 3 plates. Color: grayish brown, with dark spots and shortish bands; belly whitish. On the upper part of the head the spots form longitudinal bands, 1 of these (single) running down in the median line of front, between the eyes. A second (and more distinct) band extends on each side of the snout from the tip to the anterior margin of the eye, hence running under the orbital rim; in some specimens it is continued as a narrow ring round the eye, but commonly this is broken and indistinct. On the opercles and the sides of the snout the spots are roundish and well marked. On the body also the spots are roundish, barely oblong, their size equalling that of the pupil. They are darkest and most distinct on the back, being sometimes almost obliterated on the sides. The color of the fins is rather indistinct in the badly preserved specimens before me. The pectorals have a dark oblong spot at their bases (from about sixth to eleventh ray); the dorsals have 2 dark bars separated by whitish, and with the tip in first dorsal also blackish (in second whitish). The caudal has a dark cross bar a little behind the base, and a dark margin. The anal is apparently whitish to the margin in the female; in the single male specimen at least the outer half is blackish. Kamchatka (type specimens in the Christiania Museum). (Collett.)

species; 3 in front of spinous dorsal, 11 (or 12) along base of spinous dorsal, 2 (or 1) between dorsals, 6 at base of soft dorsal, 14 (or 13) unpaired plates on back of caudal peduncle. The unpaired plates with the spine notched at tip; all the plates marked with strong radiating ridges and the surface of the spines minutely roughened. Color, light gray above, white below; top and sides of head with black dots and dashes; back crossed by six narrow black bars. Distinguished from all other species of the genus by the great development of the lateral ridge and spines on head. This gives the head and especially the snout a much broader outline, approaching in this respect *Agonus cataphractus*. The species is represented by several young specimens in rather poor condition, the type being 53 mm. long. The outline and armature of the head are not, however, essentially different in young and adults of such species as are known to us from specimens of different sizes. Off Shana Bay, Iturup Island, Kuril Group. (This species is named in honor of Prof. D'Arcy Wentworth Thompson, of the University at Dundee, the commissioner of Great Britain in the fur seal investigations in Bering Sea in 1896 and 1897.)

Podothecus thompsoni, JORDAN & GILBERT, Rept. Fur Seal Invest. 1896, 1897 MS., off Shana Bay, Iturup Island, at Albatross Station 3653, in 18 fathoms. (Type, No. 5667, L. S. Jr. Univ. Mus. Coll. *Albatross.*)

2413. PODOTHECUS ACIPENSERINUS * (Tilesius).

(COMMON ALLIGATOR FISH.)

B. 6; 1 D. 8 to 10; 2 D. 7 or 8; A. 7 or 8; P. 17 or 18; V. 3 (I, 2); C. 1–12–1; vertebræ 12 + 29 = 41. Pyloric cæca 7 (2 individuals). Total length reaches 10 inches. Body as deep as wide, ridges of dorso- and ventrolateral series strong, while those of the rest are but little developed, thus giving body a sharply quadrangular form in front; dorsal and ventral faces flat between ridges, the lateral convex. Behind middle of first dorsal the lateral ridges become more prominent and body more regularly 8-hedral; tail 6-hedral, strongly depressed. The strong, curved, minutely serrated spines of the dorsolateral series grow smaller from occiput backward and disappear at beginning of second dorsal; superior lateral series spinous from middle of first dorsal to middle of peduncle; inferior lateral series spinous for a short distance from a little in front of to a little behind anal; ventrolateral series keeled, not spinous. Two pairs of plates between last ray of second dorsal and first median plate, and 2 to 4 pairs between last anal ray and first median plate. About 20 well-developed

Diagnosis: Body tapering uniformly from head to caudal, as high as wide anteriorly, width at base of pectorals 7 in length; tail from front of second dorsal to caudal much depressed; head 4; plates in dorsal series 37 to 38; between occiput and first dorsal 4 pairs; between dorsals 2 or 3, from ventrals to anal 10 to 11 pairs. Gill membranes joined to isthmus without fold; no teeth on vomer or palatines. A pair of complex groups of barbels on under side of tip of snout, another group at angle of mouth, and a pair of double barbels at sides of lower jaw. Snout with 2 pairs of spines, 1 pair directed horizontally forward, the other curving backward and outward. One pair supraocular and 1 pair occipital spines. Color grayish brown, yellowish below; sides with narrow undulating vertical blackish streaks.

plates on breast. Branchiostegal membranes and lower side of mandible everywhere naked; two or 3 moderate plates in front of pectoral fin; membrane between ventrolaterals, behind vent, with numerous imperfectly developed plates; none between ventrals and vent. Head 4-hedral, tapering to a long snout, its width 6¾ in length of body, nearly vertical; orbits high up, about 4⅓ in head. Supraocular ridges high, with a very strong spine; occipital ridges long, rising gradually and ending in a single strong spine; temporal ridges moderate, ending in a small, suprascapular spine; suborbital, preopercle, and opercle radially striated; suborbital ridge low down on cheek, shelf-like in front on preorbital, with 2 small spines, a minute one on preorbital and a strong one on preopercle. Opercle with a slight ridge, minute spine or none; interorbital space concave, about 3½ in head, flat between the supraorbital ridges, covered with skin and with 2 minute ridges extending upward longitudinally from near the mesethmoidal spines; rostrum projecting far beyond premaxillaries, with a pair of spines directed horizontally forward, behind these another pair, farther apart, curved backward and outward. Snout from premaxillaries to tip of spine equal to orbit; maxillary reaching front of orbit; lower jaw very short, broad, falling far short of upper. Teeth weak, few, excessively few and weak on upper jaw, none on vomer and palatines. A pair of complex groups of barbels on under side of tip of snout; another group at angle of mouth, apparently made up of 4 subordinate groups—1 near tip and 2 at tip of maxillary, and 1 at angle of mouth; a pair of double barbels at sides of lower jaw. Dorsal fins close together; anal beginning a little in front of second dorsal; pectorals about 6 in body, their base 2½ in their length, nearly truncate. Ventrals of female equal orbit, in male twice as long; width of caudal at base 3 in the length. Lateral line 37 to 39. Color grayish brown above, yellowish below, sides with narrow undulating brownish black vertical streaks in intervals between plates; first dorsal with a broad band of darker along middle, black edged; pectorals with a dark bar near base; several more or less distinct cross bars of darker, formed by long dark streaks on rays; inferior third of fin nearly uniformly dark; ventrals light in female, dark in male; anal edged with black posteriorly; caudal olivaceous, tipped with darker. Length 10 to 12 inches. Kamchatka to Puget Sound, in moderate depths, everywhere the most abundant of the family. Very common in Puget Sound. The specimens here described from the Aleutian Islands. According to Dr. Gilbert, this is one of the most abundant species in Alaska, occurring everywhere in shallow water around the Aleutian Islands and in Bristol Bay. (*acipenserinus*, sturgeon-like; *Acipenser*, sturgeon.)

Agonus acipenserinus, Tilesius, Mém. Acad. Petersb., iv, 422, pl. 11, fig. 163. 1811, Unalaska; Günther, Cat. Fishes, ii, 212, 1860; Jordan & Gilbert, Proc. U. S. Nat. Mus. 1880, 332.

Podothecus peristethus, Gill, Proc. Ac. Nat. Sci. Phila. xiii, 1861, 77, 259, Simeahmoo, Washington Territory.

Phalangistes acipenserinus, Pallas, Zoog. Rosso-Asiat., iii, 110, pl. 1811.

Aspidophorus acipenserinus, Cuvier & Valenciennes, Hist. Nat. Poiss., iv, 207, 1829.

Podothecus acipenserinus, Jordan & Gilbert, Synopsis, 730, 1883.

Paragonus acipenserinus, Gill, Proc. Ac. Nat. Sci. Phila., xiii, 1861, 167.

2414. PODOTHECUS VETERNUS,* Jordan & Starks

Head $3\frac{2}{3}$ in length; depth $7\frac{1}{3}$. D. IX, 8; A. 8; scales in lateral line 38 or 39; pectoral 15; orbit $4\frac{1}{3}$ in head; snout $2\frac{1}{5}$; upper rays of pectoral $1\frac{1}{2}$; highest dorsal spine $2\frac{1}{3}$; highest dorsal ray $2\frac{2}{5}$; highest anal ray $2\frac{2}{5}$; caudal $2\frac{1}{2}$. Body elongate, about as wide as deep anteriorly, much wider than deep posteriorly; mouth inferior, the lower jaw shutting far behind the upper; teeth on jaws, vomer, and palatines obsolete; a few short barbels beneath snout in front of mouth and at angle of mouth, their length about equal to pupil, lower jaw very cavernous. A pair of short, blunt, rostral spines pointing directly forward; at their base and wider apart is a pair of sharp spines curving outward, backward, and upward; at the posterior end of the rather wide rostral groove is a pair of small spines pointing upward and backward; from their base a pair of diverging ridges run through the interorbital to above posterior margin of orbit; a strong spine over eye, and a longer one at occiput; a low, sharp ridge on side of head, running from ocular spine and ending in a low spine at upper end of gill opening; a very low ridge on opercle not ending in a spine; preopercle with a strong spine with a wide keel-like base; a hooked spine below eye on suborbital, from which a ridge runs along lower edge of preorbital to end of snout, below posterior end of rostral

* Allied to *Podothecus veternus* is a Chinese species:

PODOTHECUS STURIOIDES (Guichenot).

D. IX, 8; A. 10; P. 15; V. I, 2. Depth, at nape, 8 in total length. Body elongate, octagonal in front, much attenuated, and becoming pentagonal behind; everywhere covered with osseous radially striated, keeled plates which are higher than long; the keels with denticulate edges and terminating in backwardly directed spines, forming 8 longitudinal series, 4 on each side. Breast in front of pectorals and median line of belly armed with smaller polygonal plates with keels or tubercles, but without backwardly directed spines. Snout conical (not divided at tip), forming a strong projection in front of mouth, and somewhat resembling that of the sturgeon, armed at its tip with 2 pairs of spines, the anterior directed forward (?) and the posterior upward and backward; the bony pieces of which it is composed with their edges denticulate, as are the sharp keels of the suborbitals; a strong backwardly directed spine (closely approximated nasal spines) on middle of snout. Lateral line marked by an interrupted series of small tubulous and osseous elevations, and lies between the 2 adjacent lateral series of plates; commencing at height of shoulder it describes a slightly convex curve and bends a little to run in a straight line to base of caudal. Mouth a little protractile, with fleshy lips, and situated far behind tip of snout, cleft as far back as front of eye. Lower jaw with extremely fine, velvety teeth; none on upper jaw nor on vomer or palatines. A group of long, unequal, fleshy filaments or barbels at each angle of upper jaw and under tip of snout; under each branch of lower jaw 2 others, which are excessively small. Eyes large, oval, on line with profile; interorbital space a little concave, equal to "transverse" (vertical?) diameter of orbit. A projecting tubercle or spine on posterior upper part of orbit, and another on each side of posterior part of occiput. Dorsals separated by an interval equal to eye; the anterior longer than posterior; their form elliptical, their height nearly equal to that of body; their rays slender, flexible at tips, especially those of posterior fin. Pectorals large, nearly as long as head; upper rays a little longer than those next below, decreasing gradually to tenth, whence they increase a little and again decrease by degrees to last; 5 lower rays thickest, their tips a little exserted; rays all simple (not branched). Anal commencing under end of first dorsal, which it equals in length and height. Origin of ventrals under base of pectorals; they are small (tips broken in the single individual), apparently of 3 rays. Caudal long, its border rounded, about 8 in total length. Color yellowish brown, paler below, with a very light black punctulation and more or less regular, round brown spots on head and body; spots smaller on back than elsewhere; a spot of same form and color on base of pectorals; pectoral fins uniform yellow; dorsals with brown upon rays; a dark border on spinous (dorsal); a dark, narrow band from anterior border of eye to tip of snout. A single specimen, $10\frac{1}{2}$ inches long. China. (Guichenot.) Guichenot's figure gives 1 more ray in anterior dorsal than the description, and gives the caudal fin as strongly concave (*sturio*, sturgeon; εἶδος, likeness.)

Paragonus sturioides, GUICHENOT, Nouvelles Archiv. Museum, 202, pl. XII, fig. 3, China.

groove; on this ridge is a triangular spine pointing backward; between this and the suborbital spine is an acute outward-pointing spine not much widened at its base; interorbital concave, its width equal to the length of the eye, 2 in snout; supraorbital rim prominent. Dorsal ridge of body continuous with occipital and supraorbital spines, joining its fellow of the opposite side posteriorly directly behind the second dorsal, and continued simply on caudal peduncle; the spines large and strongly hooked back anteriorly, becoming nearly obsolete posteriorly, only traceable on caudal peduncle by the center of each plate on the median line being slightly produced; spines on lateral ridges with stronger spines near middle of body than anteriorly or posteriorly; 2 or 3 blunt spines above base of pectoral, indicating an obsolete ridge between lateral ridges; lateral line at end of pectoral fin running along the upper lateral ridge a short distance, and becoming obsolete anteriorly; spines of abdominal ridge low and blunt, nearly obsolete posteriorly, the ridge joining its fellow of the opposite side directly behind base of anal fin and continuing as a single low ridge on caudal peduncle; a small plate before base of each ventral, a median row of 3 running forward to gill membrane, 3 on each side of these, a row around base of pectorals. Origin of dorsal behind the fourth dorsal plate, including the membrane behind, it covers 9 plates; 1 plate between dorsals, the second dorsal covers 8 plates, behind which are 14 plates; the last ray of first and second dorsal and anal connected to the body by a membrane; upper ray of pectoral the longest, reaching to below the ninth or tenth spine of dorsal ridge, the lower rays slightly produced beyond the membrane. Color in spirits, reddish brown above, light below; narrow, irregular, transverse streaks across back and sides, a longitudinal dark bar along each side of base of both dorsals; a dark streak forward from eye; margin of spinous dorsal blackish, soft dorsal with a small spot behind, a dark spot on pectoral rays near their base, and some dark bars behind it across rays; anal and ventrals colorless; caudal dusky. Sea of Ochotsk. A single specimen, about 8 inches in length, collected by Captain Blair at Robben Island. This species is related to *P. acipenserinus* and *P. gilberti*, differing from the former in having fewer and shorter barbels, teeth on jaws obsolete, keel and preopercle larger, dorsal ridges without spines posteriorly, and the spines on the preorbital ridge different in shape. From the latter in having the body different in shape, not everywhere deeper than wide, but the reverse posteriorly; anal much shorter and lower, no teeth on jaws, and the spines on preorbital ridge better developed and different in shape. (Jordan & Starks.) (*veternus*, an old man, veteran, in allusion to the want of teeth.)

Podothecus veternus, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 819, pl. 89, Robben Island. (Coll. Captain Blair. Type presented by Alaska Commercial Company to L. S. Jr. Univ. Mus., No. 4823.)

768. AGONUS, Bloch & Schneider.

Agonus. BLOCH & SCHNEIDER, Syst. Ichth., 104, 1801 (*cataphractus*).
Aspidophorus, LACÉPÈDE, Hist. Nat. Poiss., III, 221, 1802 (*cataphractus*).
Phalangistes, PALLAS, Zoog. Rosso-Asiat., III, 1811, 113 (*cataphractus*).

Closely allied to *Podothecus*. Body tapering, depressed anteriorly, its depth 6 to $6\frac{2}{3}$, its width $4\frac{1}{4}$ to $4\frac{1}{2}$ in its length. Head $3\frac{2}{3}$ to 4, depressed, about as broad as long. Plates in dorsal series about 31 or 32; between occiput and first dorsal 4 pairs. Plates of body keeled, without spines. Mouth small, subinferior, lower jaw included; teeth on jaws, none on vomer or palatines. Gill membranes joined to isthmus, with a narrow free fold behind. Numerous scattered barbels on chin, lower jaw, and branchiostegal membrane; a single pair under tip of snout. Snout with 2 pairs of upwardly directed spines. Fins short. Northern Europe; ranging to Iceland and Greenland. (*ἀ-*, without; *γωνία*, joint, i. e., rigid.)

2415. AGONUS CATAPHRACTUS, Linnæus.

(Sea-Poacher; Pogge.)

Br. 6; D. V or VI, 6 to 8; A. 5 to 7; P. 15 to 17; V. 3 (I, 2); C. 2(3)–7–2(3). Pyloric cæca 5 or 6 (Day). Greatest height, over base of pectorals about $7\frac{1}{2}$, and greatest depth, over preopercles, about 5 in total length; greatest height about $1\frac{1}{2}$ in greatest breadth. Vent lies at about $\frac{1}{4}$ distance from base of ventrals to anal, the distance between it and tip of snout $3\frac{1}{2}$ in total length. Body anteriorly depressed, appearing less so because of prominence of dorsal ridges; posteriorly much narrowed, height about equal to breadth; anteriorly, from head to about end of first dorsal, hexagonal; caudal peduncle hexagonal, with dorsal and ventral keels; middle of body octagonal. Plates of body keeled, 4 pairs between head and first dorsal; about 10 pairs from ventrals to anal; 10 median plates on peduncle; plates in dorsal series about 32. Dorsal and anal fins lie in broad, shallow grooves. Vent between plates of third pair behind ventrals. Head broad, short, its length about $4\frac{3}{4}$ in total length; anteriorly much narrowed, dorsally somewhat concave, ventrally flat. Occipital and temporal ridges each with a blunt tubercle or spine. Interorbital space concave, considerably more than longitudinal diameter of orbit. Nasal bones large, fused together, covering the mesethmoid, and armed anteriorly with 2 pairs of upwardly directed strong spines. Under these, near tip of snout, 1 pair of small barbels. Preorbital scalloped or tuberculated at its free edge, covering maxillary. Suborbital armed with a strong, pointed, outwardly and backwardly directly spine. Lower jaw included, maxillary extending much behind the short premaxillary. Mouth rather small, almost semicircular. Numerous barbels on chin, angles of mouth, and gill membranes. Eyes small, their longitudinal diameter about 5 in head, that of orbit considerably less than snout. Bones of top and sides of head striated. Preopercle with a strong backwardly and upwardly directed spine; the subtriangular opercle without spine. Branchiostegal rays 6, the membranes united and joined to isthmus, with a fold behind (?). Teeth small on jaws; none on vomer or palatines. First dorsal somewhat rounded, with 5 or 6, usually 5, spines; second dorsal with 6 to 8, usually 8, soft rays, a little higher than first; (about 3 pairs of plates between the adjacent rays of the 2 dorsals); anal begins a little

behind beginning of second dorsal, and ends under the posterior end of the latter, rays 5 to 7, usually 6 or 7; pectorals large, about as long as head, broadly oval in form, with 15 to 17, usually 15, rays, of which the lower are somewhat thickened and have their tips exserted; ventrals about ½ of pectorals, rays 3 (I, 2); caudal rounded, with 7 long rays, and on each side 2 to 3 small ones. Scales with 37 to 40 pores. Color of back and sides grayish brown; sometimes with olive-brown markings and somewhat marbled with darker; ventral surface pale; 3 or 4 blackish or blackish-brown cross patches, of which 1 lies about midway on peduncle and 1 at base of caudal; ventral side of tail pale, with grayish-brown spots; dorsals, pectorals, and anal grayish, with blackish-brown patches forming indefinite bands. Males are distinguished, according to Nilsson, by a genital papilla and broader interorbital space, which equals 1½ diameter of eye. The young, about 1½ inches or more, differ in many respects from the adults. According to Steenstrup and Lütken they have a less elongate body form, its breadth about 3¾ in total length; snout less prominent; diameter of orbit about equal to snout; the 4 nasal spines smaller than in adult, but the keels on head and the post-temporal spines more prominent and sharper, as are also the keels and spines of the plates of the body, especially those on anterior part of back and on tail. Median dorsal keel of tail, which is single in the adult, is double almost to base of caudal. Barbels on under side of head rudimentary. Ventrals somewhat longer. Vent farther back, between the plates of fifth pair behind base of ventrals. Sharper contrasts between the light and dark colors. In young of about 17 mm. (about ¾ inch) Collett found head 4 in total length. Between second dorsal and caudal fins mere traces of the embryonic dorsal fin. Upper and lower jaws equal. Plates of body with very sharp and high spines; supraorbital ridge with a high, sharp spine; both the dorsal and ventral median keels of tail double. Only anterior part of lateral line apparent. In a young one of 39 mm. (about 1⅜ inches) the upper jaw projected considerably beyond the lower, and the double row of plates along dorsal face of tail was nearly fused into one (after Lütken). Total length about 8 inches, the more usual length being 6 inches. Northern Europe to western Greenland;* recorded from Baltic Sea; southern, western, and northern coasts of Norway, as far as Russian Lapland; Arctic Ocean, White Sea (Pallas); Iceland and Faröe Islands (Lütken); Great Britain; northern coast of France (Gervais and Boulart). Found in 6 to 16 fathoms (except in winter), mostly on sandy bottoms; till lately not known from American waters, it having been confounded with *Leptagonus decagonus*. (Eu.) (*cataphractus*, κατάφρακτος, mailed.)

Cottus cirris plurimis, ARTEDI, Ichthyologia, Part IV, 87, 1738.

Cottus cataphractus, LINNÆUS, Syst., Nat., Ed. X, 264, 1758; after ARTEDI; Syst., Nat., Ed. XII, Part I, 451, 1766; FABER, Naturgesch. der Fische Islands, 117, 1829, Jutland; Iceland.

Cottus brodamus, BONNATERRE, Encycl. Meth., 67, northern seas, after OLAFSSEN, Ist. I, 589.

* The Museum of Stanford University has recently received 2 fine examples of this species from Davis Straits, western Greenland, through the kindness of Prof. D. W. Thompson of the University of Dundee.

Aspidophorus armatus, LACÉPÈDE, Hist. Nat. des Poiss., III, 222, 1802, Atlantic Ocean.
Cataphractus schoneveldii, FLEMING, British Animals, 216, 1828, Great Britain.
Aspidophorus europœus, CUVIER & VALENCIENNES, Hist. Nat. des. Poiss., IV, 201, 1829, sandy shores of Northern Europe; RICHARDSON, Fauna Boreali-Amer., III, 48, 1836.
Agonus cataphractus, BLOCH & SCHNEIDER, Syst. Ichth., I, 104, 1801; GÜNTHER, Cat., II, 211, 1860; GILL, Proc. Ac. Nat. Sci. Phila., XIII, 1861, 167; LÜTKEN, Forelöb. Meddel. Nord. Ulkefiske, Vidensk. Meddel. Naturhist. Foren. Kjöb., 381 (27), 1876; COLLETT, Norges Fiske, 38, 1875; MALM, Göteborgs och Bohusläns Fauna, 406, 1877; DAY, Fishes of Great Britain, 67, pl. 28, fig. 1, 1881; LILLJEBORG, Fauna, Sveriges och Norges Fiskar, 1883-84.
Aspidophorus cataphractus, KRÖYER, Danmarks Fiske, I, 143, 1838; STEENSTRUP & LÜTKEN, Vidensk. Meddel. fra den. Naturhist. Foren. Kjöb. 1861, p. 277, 1862.

769. STELGIS, Cramer.

Stelgis, CRAMER, Proc. Cal. Ac. Sci. 1895, 821 (*vulsus*).

Body tapering, depressed throughout, its height about 8, its breadth about 7 in its length; head 4, depressed. Plates of body spinous, about 37 in dorsal series; only about 8 median plates on peduncle; 7 pairs between occiput and first dorsal. Mouth inferior, snout projecting beyond premaxillaries; teeth on jaws, none on vomer or palatines. One pair of horizontal and 1 pair of recurved spines at tip of snout. Scattered barbels or cirri on maxillary, mandible, and branchiostegal membranes; a few on under side of snout, or none. One pair preocular, 1 pair supraocular, and 2 (4) pairs occipital spines. A transverse pit at occiput. Branchiostegal membranes joined to isthmus. (*στελγίς*, or *στλεγγίς*, a scraper.)

2416. STELGIS VULSUS * (Jordan & Gilbert).

B. 6; D. IX, 7; A. 9; P. 14; V. 3 (1, 2). Body elongated, tapering rapidly backward, depressed throughout, height $\frac{6}{7}$ of breadth, the latter about 7 in length of body. All plates of body armed with spines except those in anterior and posterior parts of the ventrolateral series; spines strong and recurved. Only about 8 plates in median dorsal series of tail, but the spines of the ridge continue, double. At base of tail on each side is a spine, between the lateral ridges; vent just behind middle of length of ventrals. Lateral line about 40. Ventral surface flat throughout. Head acutely triangular (seen from above), depressed, depth about $\frac{3}{5}$ of width, the latter about $5\frac{1}{4}$ in length of body; length of head about 4; profile irregularly sigmoid, depressed at front of eyes, thence nearly straight to tip of rostral spines. Mouth V-shaped, inferior; maxillary reaching front of pupil; snout projecting anteriorly beyond premaxillaries. Teeth in bands on jaws, none evident on vomer or palatines. Maxillary, mandible, and branchiostegal region with small, scattered cirri; under side of snout with few barbels or none. Eye large, nearly

* Diagnosis: Body tapering, depressed throughout, height about $\frac{6}{7}$ of breadth, the latter 7 in length of body. Body spinous; only about 8 median plates on peduncle. Head 4; plates in dorsal series about 37 (number between last ray of second dorsal and first median plate not recorded; all others 34); between occiput and first dorsal 7; between membranes of dorsal 4. Gill membranes joined to isthmus. No teeth on vomer or palatines. Scattered barbels on maxillary, mandible, and branchiostegal membranes; a few on under side of snout, or none. One pair of horizontal and 1 pair of recurved spines on snout. One pair preocular, 1 pair supraocular, and 2 (or 4 ?) pairs of occipital spines. Dark brown, with 7 to 9 cross bars; fins mostly dark.

as long as snout, 3⅜ in head, the orbital bones forming a ridge around it; interorbital space concave, straight longitudinally, with a longitudinal groove and 2 slight ridges. Spines on head highly developed; 2 pairs rostral spines, 1 pair directed horizontally forward and the other pair behind these, recurved upward, backward, and outward; behind these a smaller pair projecting upward (mesethmoid spines); orbital ridge serrated, with a peculiar spine anteriorly and a supraocular spine posteriorly; occipital ridges with 2 pairs (or 4 pairs) of spines, space between them roughish, somewhat concave, with traces of a median keel; between these ridges, at occiput, a conspicuous transverse pit, broader than long, longer than deep; temporal ridges more prominent, each with 4 spines, last one very strong; a median row of minute spines on back and top of head; suborbital ridge extending from front of preorbital across cheek, with 2 or 3 irregular series of spines and tubercles, and ending in a strong preopercular spine; below this on preopercle 3 or 4 other bluntish spines; opercle with ridge and 1 or more spines; between 70 and 80 more or less developed spinous processes on head. Branchiostegal rays 6 (?); membranes united, joined to isthmus. Plates between occiput and first dorsal 7; between dorsals about 4 (number between the adjacent rays of the 2 fins not recorded). Anal region beginning under end of first dorsal; pectorals reaching eleventh plate, with minute spines at their base; ventrals short. Color dark brown, with 7 to 9 darker cross bars, extending on fins; one bar across caudal, 1 across posterior part of second dorsal and anal, 1 across posterior part of first dorsal and front of anal, and 1 across front of first dorsal; pectorals black with whitish edging and a pale blotch near base; other fins chiefly black, anal with whitish edge; ventral surface pale. Total length of known individuals reaches 4⅛ inches. (Jordan & Gilbert.) Deep water off San Francisco, California; known only from the original type, taken in a parranzella or sweep net, near Point Reyes, 1880. (*vulsus*, beardless.)

Agonus vulsus, JORDAN & GILBERT, Proc. U. S. Nat. Mus., III, 1880, 330, Point Reyes, near San Francisco. (Coll. Jordan & Gilbert.)

Podothecus vulsus JORDAN & GILBERT, Synopsis, 730, 1883; JORDAN, Cat. Fishes N. A., 114, 1885.

Stelgis vulsus, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 821, pl. 90.

NOTE.—Allied to *Stelgis* and *Averruncus* is the single Antarctic genus, which may be thus defined:

AGONOPSIS, Gill.

Agonopsis, GILL, Proc. Ac. Nat. Sci. Phila., XIII, 1861, 167, 259 (*chiloensis*).

Body 8-hedral, peduncle 6-hedral; head depressed, its length a little more than 4, its width 5, and its depth in front of nape a little less than 7 in length of body. Jaws nearly equal, teeth on jaws and vomer, and an imperfect row on palatines. Two pairs of spines on snout, 1 pair preopercular, 1 pair supraocular, and 2 pairs occipital spines; a deep transverse depression at nape. Plates in dorsal series about 37; 7 pairs between occiput and first dorsal; plates of body with curved spines. Barbels under tip of snout, and scattered on chin and lower jaw, and sparingly on branchiostegal membrane. Gill membranes united, joined to isthmus. Antarctic. (*Agonus*; ὄψις, appearance.)

AGONOPSIS CHILOENSIS (Jenyns).

B. 6; 1 D. 8; 2 D. 7; A. 8; P. 14; V. 3 (1, 2); C. 2-11-2. Interorbital space concave, a little less than diameter of eye, with a pair of sharp minor longitudinal ridges terminating at a transverse groove behind eyes. Snout does not project beyond mouth; larger margin of suborbital with an irregular ridge formed of bluntish tubercles, the last terminating in a minute backwardly directed spine. Limb of preopercle with 3 diverging smooth ridges dilating at their extremities into 3 flattened blunt points; opercle with a smaller spineless ridge. Jaws about equal, the upper a little longer, with narrow bands of teeth; teeth on front of vomer and a short imperfect row on each palatine. Gill membranes not notched, attached to isthmus; gill openings large. Barbels short, on chin, lower jaw, and branchiostegal membrane (a pair also on under side of tip of snout in fig. 1, pl. 7). Temporal and occipital ridges present, formed of granulated tubercles (the occipital "tubercles" very plainly in the form of 2 pairs of spines in the figure); between occipital ridges a slightly raised longitudinal line; a deep transverse depression at nape. Temporal ridges end in a sharp point not prolonged into a spine. The carinated plates more sharply serrate than in *A. cataphractus*, the spines curved. (All the plates spinous from head to caudal in fig. 1, pl. 7.) Six slightly serrated scales forming on breast a somewhat triangular patch, 2 single ones standing first, the 4 others in pairs. Second and third rays of second dorsal largest; anal under second dorsal; pectorals rounded, 5 in length; ventrals scarcely more than ½ of pectorals. Vent a little anterior to a line connecting the extremities of ventrals. Color in spirits, dusky gray above and on sides, paler below; a blackish transverse band under front of first dorsal, 1 under anterior and 1 under posterior end of second dorsal, 1 in middle of peduncle, and a trace of a fifth behind this; body mottled in places with blackish, and the fins, except ventrals, are of the same hue. (Jenyns.) Coasts of Chile and Patagonia. (*chiloensis*, living at Chiloe, an island off Patagonia.)

Aspidorphorus chiloensis, JENYNS, Zool. Voyage of the Beagle, Fish., 30, pl. 7, figs. 1, 1a, 1b, 1840, Chiloe, west coast of northern Patagonia (Coll. Darwin); GAY, Hist. Chil. Zool., II, 174.

Aspidophorus niger, KRÖYER, Ichth. Bidrag. Naturhist. Tidsskr. I, 238, 1844, Valparaiso.

Agonus chiloensis, GÜNTHER, Cat., II, 216, 1860, Valparaiso.

Agonopsis chiloensis, GILL, Proc. Ac. Nat. Sci. Phila., XIII, 1861, 161.

Agonus niger, GÜNTHER, Cat., II, 215, 1860.

770. AVERRUNCUS, Jordan & Starks.

Averruncus, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 821, pl. 91 (*emmelane*).

This genus is very close to *Stelgis*, with which it agrees in the rough armature, the presence of cirri on the gill membranes, and in other respects; differing in the presence of teeth on vomer and palatines; anal fin very long, of 11 rays; lower rays of pectoral exserted. North Pacific. (*Averruncus*, an averting demon, from the armed head.)

a. Snout rather long, the rostral spines protruding. EMMELANE, 2417.
aa. Snout shorter, the rostral spines scarcely protruding. STERLETUS, 2418.

2417. AVERRUNCUS EMMELANE, Jordan & Starks.

Head from tips of rostral spines, 4 in length of body; depth 7½. D. VIII or IX, 8; A. 11; P. 14; lateral line 35; orbit 4 in head; snout, to tips of rostral spines, 3⅛; maxillary 3¼; interorbital 6½; pectoral 1⅔; second dorsal spine 2⅔; third dorsal ray 2⅓; longest anal ray 2⅔; caudal fin 2. Body elongate, subcylindrical, the caudal peduncle long and slender, very slightly depressed, about 3 plates in front of caudal fin it widens slightly

and is compressed; belly somewhat prominent, breaking the otherwise straight vertical outline from chin to caudal fin; dorsal outline straight from occiput to caudal fin. Head, as viewed from above, almost regularly triangular, the prominent preopercular spines and the snout forming the angles, its dorsal profile irregular, much broken by spines. Mouth inferior, broadly U-shaped, maxillary reaching just past the vertical from front of orbit; lips thick, covered with short, fine papillæ; upper jaw protractile; teeth small, in villiform bands on the jaws, vomer, and palatines; the distance from the anterior edge of premaxillary to end of the rostral spines is less than ½ the length of snout; 2 cirri as long as pupil under rostral spines, anterior lower edge of preorbitals with cirri, a group of 3 cirri on end of maxillary, and a group of 4 or 5 on posterior end of mandible, 1 on the middle of each branchiostegal ray, these forming a line from isthmus to opercle, an area on chin from the mouth to the hyal bones that is "woolly" with short cirri; 2 or 3 on lower edge of opercle and interopercle. A pair of parallel rostral spines pointing forward, their tips covered with skin; behind them is a deep oval pit, on the anterior outer edge of which are a pair of spines that point upward and outward and are slightly hooked backward; at the posterior end of the pit are 2 spines, smaller than those above and slightly curved backward; a group of 4 short spines around anterior edge of eye and 1 large triangular spine over posterior edge; the interorbital space is deeply concave, with a low, sharp ridge on each side of the median line; preopercle very rough, with irregular spines and tubercles; middle of suborbital stay with a strong, hooked spine; below the stay, on the naked area, are 2 or 3 plates with spines on their centers; angle of preopercle with a large, sharp spine; along the lower edge of preopercle are 3 or 4 blunt spines; a ridge of 4 spines running back from each eye corresponding with the dorsal keels of body; below this on each side is a ridge, somewhat irregular, but not broken into spines, terminating in a spine that points between the dorsal and upper lateral keel of body; a small ridge on upper edge of opercle which does not end in a spine; a few small spines around posterior edge of opercle; a few minute spines along median line of top of head, the upper part of the eye covered with minute prickles. At the occiput is a deep pit, broader and deeper than long, divided by a low ridge through its middle. Body with 4 ridges on each side formed by the body plates, each plate ending in a strong recurved spine, except those of the abdominal ridge, which are smooth; a row of minute spines along median dorsal line from first dorsal to occiput; small spines following the lateral line; no trace of keels or spines in front of ventrals. The abdominal ridges widest apart on the belly, uniting on tenth plate in front of caudal fin; dorsal ridges uniting on ninth scale in front of caudal fin, but the spines continuing double to the tail; a row of sharp, small spines around the base of the pectoral and caudal fins; dorsal spines slender, the fins highest in front, the second spine the longest, its tip reaching to the base of the next to the last spine, where fin is depressed; last ray adnate by membrane to the back; third dorsal ray the highest, its tip reaching nearly to the last ray, where fin is depressed; the last ray

is very short and adnate to the body for the whole of its length. Lower rays of pectoral fins produced, extending beyond the membrane, the longest extending beyond the upper ray of the fin; anal longer and lower than soft dorsal, ending at the same corresponding place, last ray reaching to the fifteenth plate before caudal fin; ventral differing in length in the different sexes; vent anterior, situated on the tenth plate in front of anal. Color dark brown, belly white; sides crossed with irregular white bars, giving the fish a mottled appearance; snout black; a black streak along lower edge of preopercle; a black spot on iris above; dorsals light, mottled with black; anal white, with dark mottlings, a dark bar across the posterior rays, the tips of all the rays white; ventrals black, abruptly white at tips; pectoral and caudal dark, with a white border, a light spot in the center of fins and many white spots on the rays; a black spot at base of pectoral. Puget Sound; known from 2 specimens, collected with a seine near Point Orchard, the largest 7 inches in length. (Jordan & Starks); another since taken at Port Ludlow. (ἐν, in; μελάνη, ink, from the dark colors.)

Averruncus emmelane, Jordan & Starks, Proc. Cal. Ac. Sci. 1895, 821, pl. xci, with plate, Port Orchard, near Seattle. (Coll. E. C. Starks. Type, 3135, L. S. Jr. Univ. Mus.)

2418. AVERRUNCUS STERLETUS, Gilbert, new species.

Head $4\frac{3}{5}$ in length; eye $3\frac{1}{4}$ in head, width at base of pectorals 7; depth $8\frac{1}{3}$. D. VII, 8; A. 9; P. 12 on each side; 39 plates in dorsolateral series. Very closely related to *A. emmelane*, with which it agrees in coloration and general appearance as well as in most details of structure. It differs in the following respects: The snout is shorter, the rostral spines scarcely protruding beyond the premaxillaries. No barbels on snout below rostral spines, or on margin of preorbital, or at mandibular joint. Region between rostral spine and front of premaxillaries wholly occupied by a triangular movable plate, with rough granular surface; breast with 3 parallel series of sharply keeled plates; no spine at posterior end of premaxillary fossa; ventrals very long and slender, wholly white; spinous dorsal, anal, and pectorals with fewer rays; rostral projection shorter than in *A. emmelane*, with 2 short forwardly directed spines, behind which is a pair more widely separated directed downward and backward; supraocular ridge elevated, not sharp, finely granular, with preocular and postocular spines; ridges and spines on head as in *A. emmelane*, but none of the former rough serrate. Eye large, longer than snout and more than twice the interorbital width. Mouth little overpassed by the rostral spines, the maxillary reaching slightly behind front of orbit, $3\frac{1}{5}$ in head. Teeth present on jaws, vomer, and palatine. Barbels fewer than in *A. emmelane;* 3 present on maxillary, 2 of which are at its posterior end, the upper much the longer; the third inserted more anteriorly behind the middle of the maxillary; 8 shorter barbels are present on each mandibular ramus, the posterior only near the joint; several short barbels on gular region, and a cross series on each branchiostegal membrane, usually 1 barbel for each ray. Plates on body as in *A. emmelane*, all with sharp

spines, which are present though small in the ventral series; middle of breast with 1 median and 2 lateral series of plates, all of which bear distinct longitudinal keels; the 2 lateral ridges on breast being the anterior continuation of the ventral ridges of the trunk; spinous dorsal beginning at the seventh dorsal plate, the last dorsal spine articulating with the thirteenth plate; the first and last rays of the second dorsal articulate respectively with the eighteenth and twenty-fourth plates; the dorsal series unite at the thirtieth plate, the median series of 9 plates thus formed bearing double or bifid spines throughout; the first and the last anal rays articulating respectively with the sixteenth and twenty-third plates of ventral series (excluding the anterior 3 on breast); the ventral series coalescing immediately opposite the union of the dorsal series; the anus is opposite the interspace between the third and fourth plates; ventral spines long and slender, equaling length of snout and eye; pectorals equaling length of head in advance of opercular joint; five lower pectoral rays with incised membranes, the tips projecting. Color, similar to *A. emmelane*, the back and sides with 7 or 8 narrow black cross bars, the posterior of which extend faintly on the under surface; the interspaces on back are somewhat dusky, with lighter vermiculating lines and spots, a few of which extend on the bars; the dorsals have a speckled appearance and are darker when they lie above the black dorsal bars; head blackish above, the head and body light or slightly dusky below; ventrals white; anal white, with some black markings along the base of the rays; pectorals with a wide black bar at base succeeded by a wide white bar; then follow a narrower black bar and a narrow terminal white bar; caudal with a narrow basal bar of black, then a narrow white bar followed by a broad black bar and edged with white. Coast southern California, Coronado Island. The type, a specimen 4½ inches long, from *Albatross* Station 3662. (*sterletus*, sturgeon; a modern Latin word, probably derived from *Sturio*.) (Type in U. S. Nat. Mus. Coll. *Albatross*.)

Averruncus sterletus, GILBERT MS., off Avalon, Coronado Island, in 47 fathoms.

771. SARRITOR, Cramer.

Sarritor, CRAMER, in Jordan & Evermann, Check-List Fishes, 448, 1896 (*frenatus*).

Body tapering uniformly to base of caudal; head 4 to 4½, depth 6 to 8 in standard length. Plates on body nearly all without spines. Plates in dorsal series 38 to 45, 5 to 6 pairs between occiput and first dorsal. No large knife-like plate over eye. Both dorsal fins present, rather long, the rays growing shorter behind the last adnate to back. Four to 6 pairs of barbels about mouth, 1 pair under tip of snout. A pair of recurved spines near tip of snout. One pair of supraocular and 1 or 2 pairs of occipital spines. Teeth on jaws and vomer, none on palatines. Gill membranes joined to isthmus, without free fold; no barbels on gill membranes. Lower rays of pectorals with free exserted tips. Vertebræ 40 to 46. This genus is very close both to *Podothecus* and to *Odontopyxis*, differing from the former in the presence of vomerine teeth, and from the latter in the absence of a free median plate at the tip of the snout. Its relations to

Averruncus are still closer, the chief difference being in the smooth plates of the body and in the absence of cirri on the gill membranes. (*sarritor*, one that scrapes.)

a. Snout moderate, about ⅓ length of head; ventrals dark in male. FRENATUS, 2419.
aa. Snout very long, produced in a flat, triangular piece exserted for a distance equal to ⅔ orbit; ventrals pale. LEPTORHYNCHUS, 2420.

2419. SARRITOR FRENATUS,* Gilbert.

Head 4½ to 4⅓. D. VI to VIII, 7 or 8; A. 6 or 7; P. 15; V. I. 2; C. 11, with a rudiment; lateral line 40. Body slightly depressed, tapering regularly backward from occiput, the depth about ⅞ of the width at base of pectorals. The ridges are prominent, the dorsal and dorsolateral ridges provided with strong spines, the ventral and ventrolateral series with weak or scarcely discernible spines, all decreasing in size backward, becoming obsolete on caudal peduncle. Dorsal face deeply concave anteriorly, its ridges coalescing, from 3 to 4 plates behind the dorsal fin; other faces much less concave, the ventral ridges coalescing, 3 or 4 plates behind the anal fin; plates in dorsal series 44 or 45; 5 or 6 pairs between occiput and first dorsal, 9 or 10 under first dorsal, 2 or 3 between dorsals, 7 or 8 under second dorsal, 17 or 18 behind dorsals; about 25 plates on breast, consisting of a strong median series which bear a well-marked rounded ridge, a strong lateral series at edge of breast also projecting, and a number of small plates occupying the concave intermediate areas. In young specimens the breast plates have central elevations, and bear each a backwardly directed spine; these disappearing in adults; a number of small irregular plates in front of and on base of pectorals; membranaceous intervals behind and around vent smaller than in *Podothecus acipenserinus*, occupied by 8 or 9 irregularly arranged plates, not in pairs and not corresponding to those of the ventral series between which they are intercalated; medial part of branchiostegal membrane and the gular region covered with roundish plates, the whole forming a halberd-shaped patch; lateral line running on a series of small plates occupying the middle of the lateral face; anteriorly these entirely disappear, the lateral line ascending and running on the upper lateral series; about 5 large plates, sometimes bearing spines, lie behind upper axil of pectorals, between this ascending portion of the lateral line and the inferior lateral series or plates; head depressed, tapering rapidly to the snout; depth of head at occiput ⅔ its width at preopercular spine, the latter contained 6⅓ times in length of body. Eye large, the orbit about equaling length of snout behind the serrated rostral ridge, 3½ to 3⅔ in head; snout somewhat vari-

* Diagnosis: Body tapering uniformly from occiput to caudal, a little more slender than *Podothecus acipenserinus*, depth about ⅞ of width at base of pectorals, the latter 7⅔ to 7⅞ in length; head 4⅓. Plates in dorsal series 44 or 45; between occiput and first dorsal 5 or 6; between dorsals 6 or 7; from ventrals to anal 14 pairs. Six pairs of barbels, 1 on under side of snout in front of premaxillaries, 1 near and 2 at tip of maxillary, 1 at angle of mouth, and 1 at side of lower jaw. A pair of perpendicular serrated plates at tip of snout, behind these a curve nasal spine. One pair mesethmoidal, 1 pair supraocular, and 1 pair occipital spines. Teeth on jaws and vomer, none on palatines. Gill membranes joined to isthmus without free fold. Last ray of each dorsal shortened and adnate to the back. Light grayish with a few indefinite blotches and bars of darker on sides and back; pectorals with several narrow dark cross bands near base and broader ones toward tip.

able in length, averaging ⅓ head. Supraocular ridges strong, rugose, ending in a short strong spine; an inner pair of ridges occupying floor of interorbital groove, very broad and closely joined anteriorly; the triangular space included between these latter ridges flat and opening posteriorly onto the depressed occipital area; occipital ridges low, rounded, ending in very strong spines which form the first of the dorsal series; 2 or 3 small rounded projections may occur on anterior ridges of the occipital spines; top and sides of head more or less closely beset with very fine prickles, which are most thickly clustered on occiput, interorbital area, upper part of opercles, suborbital chain, and sides of snout; a row of prickles on eyeball just above pupil; temporal ridge uneven, sometimes interrupted with a long, strong posterior spine; a strong spine, sometimes with an accessory tubercle on middle of cheek; preorbital with 2 pairs of spines placed vertically, the upper ones directed outward and upward, the lower spines directed downward and backward; rostral ridges rough, usually terminating posteriorly in a pair of spinous projections, which are located midway between tip of snout and front of pupil; anteriorly, at tip of snout, these ridges expand to form such a vertically projecting rounded spinous lobe, the posterior spine of which is much the strongest, and points backward and outward. In the very young the last-mentioned spines alone are present on snout, and are directed very obliquely backward; anteriorly ridges converge from them to tip of snout, and are very minutely serrulate; these ridges afterward increase in height and in strength of serrations, and become the spinous lobes already described; two strong diverging spines at angle of preopercle, and 2 rounded lobes below them. Three large plates and a number of smaller ones occupy cheek below suborbital stay. Posterior portion of mandible expanded into a rough projecting bony prominence. Mouth horizontal, overpassed by the snout in adults for a distance equaling ½ or less than ½ diameter of orbit; snout not noticeably projecting in the very young. Maxillary reaching slightly beyond front of orbit, equaling ½ length of snout and eye. Teeth in broad bands in jaws; a distinct patch on front of vomer, none on palatines. Branchiostegal membranes broadly joined, with a very narrow free fold posteriorly, or with none. Six pairs of barbels on under side of head; 1 on under side of snout in front of premaxillaries; 2 at end of maxillary; 1 near middle of maxillary; 1 on lower lip just below angle of mouth; 1 forked for ½ its length at middle of side of lower lip. In their distribution, relative lengths, and in the constantly bifid character of the last described, they correspond exactly with the barbels of *L. decagonus*, but the latter has apparently none on under side of snout. Interspace between dorsals somewhat variable, ⅔ to ⅞ diameter of orbit. Anal beginning 2 plates in advance of second dorsal; pectorals 5½ in length of body, the lower rays graduated, 4 to 6 of the lower ones thickened, with exserted tips; ventral fins nearly twice as long in males as in females, in the latter less than diameter of orbit; caudal slender, 1¾ in head. Color light grayish or brownish, pale below; a bluish black stripe from ventral spines to front of orbit; suborbital, preopercle, and opercle with numerous dark spots;

a dark blotch on the side opposite middle of first dorsal; a faint dark bar under the anterior and a similar one under posterior part of second dorsal; both dorsals with indefinite oblique dark bands; pectorals dark except the lower proximal part, with narrow bars of black; ventrals light in female, dark in male; anal dark posteriorly; caudal dark, faintly barred. Coast of Alaska; known from *Albatross* Stations 3219, 3225, 3226, 3255, 3256, 3257, 3258, 3263, 3269, 3279, 3282, 3309, and 3330, located on both sides of the Peninsula of Alaska and both north and south of the Aleutian chain; depth 16 to 350 fathoms. (Gilbert,); also obtained at Provostnaya, Kamchatka. (*frenatus*, bridled.)

Odontopyxis frenatus Gilbert, Rept. U. S. Fish Comm. 1893 (1896), 435, pl. 35, fig. 3. Alaska and Aleutian Islands, at Albatross Stations 3215, 3219, and others, in 16 to 351 fathoms.

2420. SARRITOR LEPTORHYNCHUS (Gilbert).

Head 4 in length; snout 2⅔ in head in specimen 100 mm. long. Eye 3⅔, equaling length of maxillary; interorbital with ⅔ eye. Branchiostegal membranes broadly united, extensively free laterally, joined to isthmus mesially to extreme posterior margin, or leaving a very narrow margin free. Teeth present on jaws and on vomer, none on palatines. Dorsal VI to VIII, 6 or 7; anal 6 or 7; pectoral 14; ventral I, 2; pectorals long, reaching to or beyond middle of spinous dorsal, as long as snout and eye; ventrals equaling length of snout. Very close to *S. frenatus*, with which it agrees in arrangement of plates, spines on head, and barbels; distinguishable at once by the elongate slender snout and differing in the following numerous details: The body is somewhat broader and more depressed, its greatest depth a little less than ⅔ its greatest width, which occurs across preopercular spines. The body narrows rapidly backward to below spinous dorsal, as in young *S. frenatus* of the same size. Compared with *S. frenatus* of the same size, the plates on body are much less spinous, the superior and inferior lateral and the ventral series in some specimens bearing spines on a few of the anterior plates only, and the spines of dorsal series are lower. Five plates before dorsal, 10 under spinous dorsal, 2 between dorsals, 7 under second dorsal, and 16 on caudal peduncle. The inferior lateral ridges rise anteriorly, greatly constricting the lateral face under anterior part of spinous dorsal. It then descends slightly and becomes almost or quite obsolete, the series of plates ending behind the upper pectoral rays. In *S. frenatus*, the constriction of the lateral face does not occur, the ridge is strongly marked anteriorly, and ends below middle of pectoral base. In *S. leptorhynchus* we have, therefore, a much narrower interval between the anterior ends of the upper and the lower lateral series. This interval is occupied by but 3 plates, arranged in a series, decreasing in size backward. The upper preopercular and the humeral spines are much larger than in *S. frenatus*, the former greatly overpassing the second spine. The rostral spines are similar, but the terminal plate roughened but not serrate, the posterior spine not detached. Snout greatly produced into a narrow triangular piece which overpasses the mouth, for a distance equaling ⅔ diameter of orbit in a specimen 100 mm. long. In specimens of *S. frenatus* of this length the

ends of the rostrum can barely be seen from below. A few prickles present on upper side of rostrum, and the usual series above pupil. Minutely serrated ridges on sides of snout, and 1 below eye. No prickles on top or sides of head. Plates on branchiostegal membranes and on gular region smaller and more numerous than in *S. frenatus*. Twenty plates on breast, without spines, or the young with very small ones. Color darker than in *S. frenatus*, the under parts unmarked anteriorly, dotted posteriorly with brown; upper parts dark brown in spirits, with 6 or 7 more or less distinct black bands, which are margined narrowly with lighter; a black streak forward from eye, and several black spots and blotches on sides of head; caudal blackish; soft dorsal dusky, obscurely marked with lighter; spinous dorsal black, sharply blotched with pure white; ventrals and lower pectoral rays white, the upper part of pectorals with obscure bars of black. Coast of Alaska. A few specimens from *Albatross* Stations 3215, 3219, 3259, and 3267, north and south of the Alaskan Peninsula, in 32 to 59 fathoms. (Gilbert.) (*λεπτός*, slender; *ῥύγχος*, snout.)

Odontopyxis leptorhynchus, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 437, Bering Sea, north of Alaska Peninsula, at Albatross Station 3267, lat. 55° 23′ 30″ N., long. 163° 29′ W. (Type, No. 48727. Coll. *Albatross*.)

772. XYSTES, Jordan & Starks.

Xystes, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 824 (*axinophrys*).

This genus is allied to *Averruncus*, differing in its shorter vertical fins, the rays of both dorsals being subequal, the last of each not much shortened and not adnate by membrane to the skin of the back; the lower rays of the pectoral progressively shortened; plates on body very rough, the spines strong; a strong knife-like spine above eye; no barbels; no pit at the occiput. North Pacific. (*ξύστης*, one that scrapes.)

2421. XYSTES AXINOPHRYS, Jordan & Starks.

Head 3½ in length of body; depth 5. D. IX, 8; A. 10; pectoral 15; lateral line 38; orbit 4 in head; snout to tip of rostral spines 3⅙; maxillary 3⅙; interorbital 3½; pectoral 1⅓; highest dorsal spine 2½; highest dorsal ray 2; highest anal ray 2¾; length of caudal fin 1⅔. Body elongate, subcylindrical, deepest and broadest at shoulders; belly prominent; dorsal outline straight from first dorsal spine to caudal fin, curved up anteriorly to occiput. Head very irregular, much broken by large spines; mouth inferior, rather broad, maxillary reaching to the vertical from front of orbit; lips thin, not broken up into papillæ; upper jaw protractile; teeth small, in villiform bands on jaws, vomer, and palatines; the anterior edge of premaxillary directly under the base of rostral spines; a few very small blunt papillæ behind chin, a barbel at end of maxillary, not ½ so long as diameter of pupil. A pair of sharp rostral spines pointing forward and upward; behind these a pair of curved spines pointing upward, outward, and backward; between these and behind the rostral spines an almost circular pit which is entirely occupied by the upper end of the premaxillary process; interorbital wide and concave, a slight median ridge, running from the rostral pit to a point above pupil, each

side of which is an outwardly curved ridge ending in a minute spine; over each eye is the largest spine on the head or body; the large triangular orbital spine, its base occupying nearly the whole space above eye, sharp and strongly hooked back; on the anterior part of its base a small, sharp, preorbital spine, pointing upward; a series of minute spines, running medially along the top of the head and body, from a point between the orbital spines to the first dorsal spine; on each side of these are 2 large blunt spines with traces of a smaller one between them, continuous with the dorsal keels of body; farther down and continuous with the upper lateral keel of body is a ridge broken up into 4 irregular spines, larger than the body spines; 4 triangular spines on edge of preopercle, the upper one the largest; a very irregular ridge running from upper preopercular spine, under eye, to snout; a ridge on upper part of opercle. Body with 4 ridges on each side, formed by the scales, each of which ends in a spine; traces of a ridge between lateral ridges, the spines on abdominal ridges as sharp as those on rest of body; a Y-shaped ridge of spines in front of ventrals, the forks toward the ventrals and the base ending at gill membrane; a raised area between ventral fins, running from their base to their tips, which is entirely covered with small prickles, the anus in the posterior end of this; dorsal and abdominal ridges coalescing with their fellows of the opposite side, but coming together so gradually that it is impossible to tell exactly where they unite as the spines continue distinct to the caudal fin. Small spines covering the outer part of the base of the pectoral; a ring of spines around caudal base; occiput abruptly higher than body, but scarcely forming a pit as body is about level behind it. Spinous dorsal highest in front, the second spine reaching to base of last spine when fin is depressed; the dorsal rays subequal in length, the last not shortened and not adnate to body, last ray reaching to the tenth plate before caudal fin; pectoral fin posteriorly rounded in outline, the lower rays not produced, reaching to second plate before anal fin; ventrals small, reaching just past vent; anal longer and lower than soft dorsal; dorsal and anal ending at the same corresponding place; caudal fin rounded behind. Color in spirits, gray with 7 or 8 dark cross bars; head uniform gray with the exception of a dark spot at occiput; belly dusky; dorsals somewhat mottled; anal black with a white spot near its middle; pectorals white with a large black spot on base of rays; ventral black, abruptly white at tips; caudal black, edged with white. Puget Sound. One specimen, 1¼ inches in length, dredged at Port Orchard, near Seattle. (Jordan & Starks.) Another since taken at Port Ludlow. (*ἀξίνη*, an ax; *ὀφρύς*, eyebrow.)

Xystes axinophrys, Jordan & Starks, Proc. Cal. Ac. Sci. 1895, 824, pl. 92, Port Orchard, Admiralty Inlet. (Coll. E. C. Starks.)

773. BATHYAGONUS, Gilbert.

Bathyagonus, Gilbert, Proc. U. S. Nat. Mus., xiii, 1890, 89 (*nigripinnis*).

Spinous dorsal developed; lower jaw the longer; plates of body spinous; gill membranes united to isthmus, not forming a fold across it; teeth well developed on jaws, vomer, and palatines; pectorals not notched, the

upper rays longest, the lower becoming regularly shortened; bones of head thin and yielding, the system of mucous canals very strongly developed. Deep-sea Agonoids. (*βαθύς*, deep; *Agonus.*)

2422. BATHYAGONUS NIGRIPINNIS,* Gilbert.

B. 6; 1 D. 6 to 8; 2 D. 6 or 7; A. 7 to 9; P. 15 or 16; V. 3 (I, 2); C. 1–11–1; vertebræ 11 + 34 = (45). Pyloric cæca 5 (1 individual). Body depressed throughout, depth $\frac{1}{3}$ of width, the latter $9\frac{2}{3}$ in body length; sharply octagonal, tail strongly depressed hexagonal, its lateral faces very narrow. The ridges compressed; the small sharp spines present on all the ridges from head to caudal except on the median ventral series of peduncle and the first few plates behind the pectoral in the imperfect lateral series. One to 3 pairs of plates between last ray of second dorsal and first median plate, and 1 or 2 pairs between last anal ray and first median plate; 1 or 2 plates in front of and 4 or 5 small, sharply spinous plates on base of pectorals; about 22 plates, radially striate, elevated at the center, some of the strongest minutely spinous, on breast in a median and 2 lateral rows (the latter continuous with the ventrolateral series) and in the ridge-like rows forming the edges of breast; no plates medially on posterior part of the branchiostegal membrane or laterally on the anterior part; from 1 to 3 or 4 well-developed plates on front part of lower side of lower jaw; the membranaceous interval behind vent, between the ventrolateral series, occupied by about 5 to 9 plates, partly arranged in pairs; none between ventrals and vent. Head $4\frac{1}{4}$ to 5 in body, much depressed, its greatest depth $\frac{4}{5}$ of its greatest width, the latter equaling width of body; snout broad, flat, shovel-like, nearly equal to long diameter of orbit; orbit oblong-oval, its vertical diameter $\frac{2}{3}$ of the longitudinal, the latter 3 or a little less in head; interorbital space concave, about as in *triacanthus*, but broader, with a pair of minor longitudinal ridges, about 9 in head, about 2 in vertical diameter of orbit; the moderate supraocular ridges ending in a sharp spine, the scarcely developed occipital ridges with 2 pairs, of which the anterior is low and blunt; the well-developed temporal ridge with 2 or 3 small spines and terminating in a larger suprascapular spine; bridge across cheek close under orbit with about 1 spine on preorbital, 3 on suborbital, and ending in 1 on preopercle; 1 of about the same size below the last on preopercle. Cheek below ridge with 2 to 4 (usually 3) spinous plates; nasal spines small, sharp, directed backward; movable median rostral plate small, varying in size and shape (disappearing with the prolongation of lower jaw), its lateral expansions usually free, as spines; 3 small upright diverging spines (only 2 systematically placed in 1 specimen). Mouth oblique, lower jaw somewhat protruding, maxillary scarcely reaching front of orbit. Teeth moderate, present on jaws, vomer, and palatines. Two barbels, 1 large and the other small, at tip of maxil-

* Diagnosis: Body slender, tapering uniformly, depressed throughout, its depth $\frac{4}{5}$ of its width at base of pectorals, the latter $9\frac{2}{3}$ in length; head $4\frac{1}{4}$ to 5 in length. Plates in the dorsal series 43 or 44; between occiput and first dorsal 7 or 8; between dorsals 6, sometimes 7; between ventrals and anal 13 to 15 pairs. Lower jaw protruding; 2 barbels at tip of each maxillary; median rostral plate present, with 3 small diverging spines; nasal spines small, separate; 1 pair supraocular and 2 pairs occipital spines. Gill membranes joined to isthmus. Under side dusky, lighter above; all the fins blue black.

lary, none on lower jaw, the mucous pores bordered by lobed flaps. Gill membranes broadly united, joined to isthmus, without free fold. Dorsal fins well separated; anal beginning about 3 plates in front of beginning of second dorsal; pectorals $5\frac{5}{8}$ in body, their base $3\frac{1}{4}$ in their length, with 15 or 16 graduated rays, the uppermost longest; ventrals of male nearly 3 in head, about equal to long diameter of orbit, those of female $\frac{5}{7}$ to $\frac{5}{6}$ as long as those of male; caudal about 3 times as long as wide at base. Lateral line with 43 or 44 pores; plates spineless, rather large. In a young male 2 inches long the median rostral plate is absent, all the plates on breast sharply ridged and spinous; vent near tip of ventrals, $\frac{1}{4}$ of the distance from ventrals to anal, with 5 pairs of minute spinous plates between ventrals and vent. Body not tapering uniformly; broader in front, narrowing abruptly near front of first dorsal; tail not so much depressed; head $3\frac{1}{2}$ in body length. Color light above, blue black on under side of head; breast and belly dusky; all the fins intensely blue black. Pacific; known from the Aleutian Islands, Bering Sea, south to coast of Washington; abundant in 350 to 477 fathoms. Here described from Dr. Gilbert's types. Dr. Gilbert observes:

In adult specimens the lower pectoral rays show a tendency to elongate, as in *Xenochirus*, but the fin is never distinctly notched. The lower jaw always strongly protrudes, and the genus differs further in the very thin cranial bones and the inordinate development of the mucous system. In addition to the specific characters mentioned in the original description, we note that the eyeball does not exhibit the usual row of prickles and that 2 barbels are usually present at end of maxillary, either black or white in color. The majority of the specimens were obtained north of Unalaska Island, but it was also taken south of the Islands, and off the coast of Washington, at *Albatross* Stations 3210, 3316, 3324, 3325, 3329, 3330, 3331, 3337, and 3343; depths 109 to 483 fathoms.

(*niger*, black; *pinna*, fin.)

Bathyagonus nigripinnis, GILBERT, Proc. U. S. Nat. Mus. 1890, 89, Aleutian Islands and coast of Washington. (Coll. *Albatross*.)

774. XENOCHIRUS, Gilbert.

Xenochirus, GILBERT, Proc. U. S. Nat. Mus. XIII, 1890, 90 (*triacanthus*).

Spinous dorsal present. Jaws equal or the upper the longer. Plates of body spinous. Gill membranes united to isthmus, the posterior edge sometimes forming a very narrow free fold across throat. Tip of snout with a movable median plate or spine. Teeth well developed on jaws, vomer, and palatines. Pectorals divided by a deep notch into 2 portions, the lower composed of greatly thickened rays which are simple and frequently longer than those of the upper lobe, these lengthened rays not usually developed in the young. A series of small spines on eyeball above pupil. North Pacific. This genus is very close to *Bathyagonus* which represents it in deeper water. Both are near *Odontopyxis*, but the relation to *Averruncus* and *Sarritor* is more remote. (ξενός, strange; χείρ, hand.)

a. Rostral plate with 3 upright spines.
  b. Breast with numerous plates. PENTACANTHUS, 2423.
  bb. Breast smooth; plates of body rougher. ALASCANUS, 2424.
aa. Rostral plate with 1 upright spine.
  c. Gill membrane with posterior free fold. LATIFRONS, 2425.
  cc. Gill membrane without posterior free fold. TRIACANTHUS, 2426.

2423. XENOCHIRUS PENTACANTHUS,* Gilbert.

B. 6; 1 D. 5 to 7; 2 D. 5 to 7; A. 6 to 8; P. 14 or 15; V. 3 (I, 2); C. 1–12–1; vertebræ 12 + 28 = (40). Pyloric cæca 5 (1 individual). Body 8-hedral, slightly depressed; peduncle considerably depressed, hexagonal, the lateral faces narrow. Spines of body small and sharp, projecting backward nearly parallel with axis of body, decreasing gradually in size from head to caudal in the dorsolateral and superior lateral series, poorly developed on the inferior lateral series only from behind the pectoral to opposite front of anal, entirely absent from the ventrolateral series. Plates on breast, 2 pairs in front of ventrals, followed anteriorly by 3 median single plates; on each side of these a longitudinal row of 4 or 5 smaller plates, continuous with the ventrolateral series; outside of these the plates forming the edge of the breast. Branchiostegal membrane posteriorly with 3 or more very weak plates medially, or none at all; none on the membranes laterally; 2 or 3 fairly developed ones in front part of under side of lower jaw; the membrane behind vent studded with several poorly developed plates, none between ventrals and vent. Head as in *X. latifrons*, but more depressed and slender; snout not quite so short, blunt; greatest depth of head $\frac{2}{3}$ of its width, the latter 7 in length of body; orbits large, oval, their longitudinal diameter $2\frac{1}{2}$ in head; a slightly curved row of 4 or 5 backwardly directed small spines on eyeball just above pupil; interorbital space deeply concave, with minor ridges very narrow, about $3\frac{1}{3}$ in long diameter of eye, 9 in head; a single pair of supraocular spines; the poorly developed occipital ridges with 2 pairs of spines, of which the anterior is very low and blunt; temporal ridges moderate, terminating in a blunt spine; the interrupted ridge across cheek less prominent and shelflike than in *X. latifrons*, bearing 1 spine (sometimes 2) in front of anterior inferior angle of orbit, 1 or 2 on suborbital, and a single broad flat spine (sometimes preceded by a very small one) at posterior edge of preopercle; cheek below ridge entering into ventral surface of head with a longitudinal series of 3 or 4 plates; nasal spines sharp, far apart; median rostral plate small, movable, variable, its lateral expansions exposed as spines and with 3 (in 1 specimen 4) diverging spines directed upward and backward; not projecting beyond premaxillaries; maxillary reaching very little beyond front of orbit, entirely covered by preorbital. Jaws equal; teeth on jaws, vomer, and palatines. Two barbels at tip of each maxillary; those on under side of mandible inconstant; occasionally a very small pair near tip of jaw, at the 2 ends of the terminal mucous pore; another at edge of next mucous pore nearly constant, but very variable, long, short, or only a small pimple, single, double, or even a pair on each side; occasionally a very small one at edge of third pore. Dorsal fins separated by a moderate

* Diagnosis: Body slightly depressed, its depth a little less than its width, tapering moderately, and the tail tapering more gradually toward the caudal; width of body at base of pectorals 7 in length; head $4\frac{1}{3}$; plates in dorsal series 39; from occiput to first dorsal 7 or 8, between dorsals 5 or 6, from ventrals to anal 12 or 13 pairs; barbels 2 at tip of maxillary, those on lower jaw inconstant, usually 2 pairs. Median rostral plate small, nearly vertical, with 3 upright spines; nasal spines prominent, separated; 1 pair of supraocular and 1 pair of occipital spines. Teeth on jaws, vomer, and palatines; gill membranes joined to isthmus without free fold. Brownish olive above, pale below; 4 or 5 dusky cross bars on back.

space; first dorsal with 5 to 7, usually 6 spines; anal begins 2 or 3 plates in front of second dorsal; pectorals $6\frac{1}{3}$ in length of body, their base 3 in their length, the lower 5 or 6 rays exserted, separated from the upper by a notch in the adult, the 2 or 3 uppermost of the exserted rays sometimes longer than the longest upper rays; ventrals of female a little less than long diameter of orbit, $\frac{2}{3}$ as long as in male; caudal 3 times as long as wide at base; lateral line 39 to 41; pore plates thin, weak, spineless. Color olive brownish, pale below, with 5 faint cross bars, 1 beneath first dorsal, 1 between dorsals, 1 beneath posterior part of second dorsal running up on fin, 2 on peduncle; these bars often indistinct or absent; a blackish blotch in axil; caudal black, narrowly edged with white; dorsals dusky, darker toward tips, second dorsal narrowly white-edged; pectorals dusky; short, black streaks on rays forming indefinite cross bars, those at base aggregated into a blotch; ventrals light in both sexes. Length 7 inches. Deep waters of North Pacific, Bering Sea to San Diego, in 70 to 339 fathoms. Here described from Dr. Gilbert's types. (*πέντε*, five; *ἄκανθα*, spine.)

Xenochirus pentacanthus, GILBERT, Proc. U. S. Nat. Mus. 1890, 91, Bering Sea and off San Diego. (Coll. *Albatross*.)

2424. XENOCHIRUS ALASCANUS, Gilbert.

Head $4\frac{1}{3}$ to $4\frac{5}{6}$ in length; width of head equaling or slightly exceeding length of snout and eye. Depth of body equaling length of snout and $\frac{1}{2}$ eye. Fin rays in 8 specimens as follows:

	Spinous dorsal.			Soft dorsal.		Anal.	
Rays	V.	VI	VII.	6.	7.	7.	8.
Specimens	1	4	3	6	2	7	1

Pectoral 15 or 16. Ventral I, 2. Lateral line 39 or 40. A decided pit behind the eyes, and a deep transverse nuchal depression, the two separated by the prominent occipital region. Snout of moderate length, much depressed behind the spines, $3\frac{2}{3}$ to $3\frac{3}{4}$ in head. Eye 3 in head in adults. Interorbital space wider, much more deeply concave, the supraocular ridges very heavy, minutely roughened, ending posteriorly in robust spines. Rostral spines as in *X. pentacanthus*. A small apical plate bearing 3 small diverging spines, behind which are 2 longer ones. Preorbital with a small spinous point directed backwards. A spine posteriorly on bony bridge across cheek. Below this bridge, the cheek is entirely mailed by 3 rounded plates which bear no spines, except in the young, and so intimately joined that the sutures are difficult to discover. In *X. pentacanthus* the plates are much smaller and do not entirely cover the cheeks, leaving soft areas surrounding them; the 2 posterior plates in this species also bear spines. Maxillary $3\frac{3}{4}$ in head, barely reaching front of orbit. Teeth on jaws, vomer, and palatines, the usual row of 5 or 6 prickles on eyeball. Preopercle with 3 diverging spines at angle, a rounded lobe beneath them; spines and ridges otherwise as in *X. penta-*

canthus, but stronger and rougher; 3 or 4 strong plates present on gular membrane; a few weak ones or none on branchiostegal membrane mesially. Two barbels at tip of each maxillary, and a pair, often double, on under side of mandible arising from the margin of the anterior pair of mandibular pores. The symphyseal pore has its margin sometimes provided with very short barbel-like elevations. Space between dorsal ridges very deeply concave in front of dorsal fins; the single ridge behind dorsal fins provided with very short, scarcely preceptible, double spines; lower lateral series of plates continued forward to axil of pectoral, becoming indistinct in *X. pentacanthus;* ventral series anteriorly with few short spines or none, this series strongly spined in *X. pentacanthus;* plates on breast arranged alike in the 2 species, but in *X. alascanus* they are more finely striate, and bear neither spines nor raised centers, except in very young examples. In *X. pentacanthus* the elevated centers may or may not bear short spines. Seven plates before dorsal, 8 or 9 under spinous dorsal, 2 or 3 between dorsals, 7 or 8 under soft dorsal, 13 or 14 behind dorsals. Distance from snout to nape equaling or slightly exceeding that from nape to first dorsal. Front of anal under end of spinous dorsal or slightly behind that point, more anteriorly placed than in *X. pentacanthus.* Ventrals $2\frac{1}{8}$ to $2\frac{2}{3}$ in head. Lower pectoral rays produced, with incised membranes as long as head behind postral spines. Color lighter than in *X. pentacanthus*, more or less finely speckled above, usually with 5 or 6 dusky cross bars on back; a series of linear dark blotches below the lateral line; head often finely speckled with brown, and showing traces of a brown bar forward from eye to snout; dorsals, caudal, and upper half of pectoral light, finely speckled with brown, the caudal shaded with dusky; ventrals and anal white. Most nearly related to *X. pentacanthus* with which it agrees in having a rostral plate bearing 3 spines and in the absence of a free fold to branchiostegal membranes. It differs conspicuously in the broader head, with its much heavier spines and ridges, in the presence of deep postocular and nuchal pits, in the smooth breast and cheeks, in adults in the different coloration and fin rays, and in many other different details. Aleutian Islands; taken rather abundantly in the vicinity of Unimak Pass, both north and south of the islands, at depths of 56 to 138 fathoms. (Gilbert.) Taken by us off Karluk. (*alascanus*, pertaining to Alaska.)

Xenochirus alascanus, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 438, Unimak Pass Aleutian Islands. (Coll. *Albatross.*)

2425. XENOCHIRUS LATIFRONS,* Gilbert.

B. 6; 1 D. 6 or 7; 2 D. 7; A. 7 or 8; P. 14 or 15; V. 3 (I, 2); C. 1-11-1; vertebræ 12 + 29 = (41). Body as deep as wide, 8-hedral; caudal pe-

*Diagnosis: Body slender, tapering uniformly from head to caudal; depth equaling width, the latter at base of pectorals $8\frac{1}{3}$ in length; head $4\frac{1}{3}$. Plates in dorsal series 37 to 40; between occiput and first dorsal 6, occasionally 7; between dorsals 4 or 5; from ventrals to anal 13 or 14 pairs. One large barbel at tip of maxillary, 1 pair near tip of lower jaw, at edges of terminal mucous pore, and 1 at edge of next pore. Median rostral plate present, with a single upright spine. Nasal spines prominent, separated; 1 pair supraocular and 2 pairs occipital spines. Teeth in jaws, vomer, and palatines. Gill membranes joined to isthmus, with narrow free fold behind. Dusky olive, paler below; 5 or 6 indistinct dark cross bars; spinous dorsal with black margin.

duncle depressed, 6-hedral. The thin, sharp, curved spines present on all the ridges; on the dorsolateral and superior lateral series, from head to caudal; on the inferior lateral series, from opposite front end of first dorsal to base of caudal; and on ventrolateral series, from base of pectorals to a short distance in front of anal; 2 pairs of plates between last ray of second dorsal and first median plate, and 2 pairs between last anal ray and first median plate; about 23 oval, striated plates with raised centers on breast; no plates on branchiostegal membrane; 2 or 3 anteriorly on under side of mandible; a small plate or none in front of pectorals. Membrane behind vent with 6 to 9 small plates arranged in pairs, none between ventrals and vent. Head depressed, its depth $\frac{7}{9}$ of width, the latter (across preopercles) 7 in length of body; snout short, blunt. Orbits large, oval, the vertical diameter $\frac{3}{8}$ of the longitudinal diameter, the latter $2\frac{1}{2}$ in head; 3 to 5 small, backwardly directed spines in a slightly curved line on eyeball just above pupil. Interorbital space very narrow, 7 in head, deeply concave, with 2 minute longitudinal ridges. Supraocular ridges with 1 spine, the poorly developed occipital ridges with 2, of which the anterior is much the smaller; the interrupted suborbital ridge close under orbit, well developed, with 2 spines on suborbital and 1 on preopercle; no plates on lower part of cheek, which enters ventral surface of head; nasal spines sharp, far apart; median rostral plate small, its lateral expansions embedded in the skin, a single, small, median, curved, upright spine. Maxillary reaching slightly beyond front of orbit; lower jaw a very little included; well developed teeth in several rows on jaws, vomer, and palatines. A single long barbel at tip of maxillary; a small pair at ends of the terminal mucous pore, near the tip of lower jaw, and another just behind this at the edge of the next pore, the former sometimes, the latter frequently, bifid or a pair on each side. Dorsals fairly separated; second dorsal higher than first; anal beginning about 1 plate in front of second dorsal; pectorals $6\frac{1}{3}$ in length of body, their base nearly 3 in their length, with 14 or 15 rays, the lower exserted, separated by a notch from the upper rays in adult, the 3 or 4 uppermost of these as long as or longer than the longest upper rays; ventrals of female a little less than long diameter of orbit, $\frac{3}{4}$ as long as those of male; caudal broad and short, its width at base a little more than 2 in its length. Lateral line with 39 or 40 pores; the plates small, spineless. Color, dusky olive, light below, the back with 5 or 6 faint darker cross bars; axillary region blackish, soft dorsal and caudal with the rays black; spinous dorsal with a conspicuous jet-black margin; pectorals somewhat dusky above; ventrals pale in both sexes. Total length reaches $7\frac{1}{2}$ inches. Deep waters of the North Pacific; coast of Oregon to San Diego and outward; abundant in 50 to 204 fathoms. Here described from Dr. Gilbert's types. (*latus*, wide; *frons*, forehead.)

Xenochirus latifrons, Gilbert, Proc. U. S. Nat. Mus. 1890, 92, off Coast of Oregon and off San Diego. (Coll. *Albatross*.)

2426. XENOCHIRUS TRIACANTHUS,* Gilbert.

B. 6; 1 D. 5 or 6; 2 D. 6 or 7; A. 6; P. 13; V. 3 (I, 2); C. 1-11-1. Body depressed throughout its length, sharply octagonal, peduncle hexagonal; the sharp, delicate spines present on all the ridges from head to caudal, except the first few plates on the inferior lateral, and on the ventrolateral from a short distance in front of anal to caudal; one pair of plates between last ray of second dorsal and first median plate (the first 3 median plates sometimes each with a pair of closely approximated spines) and 1 pair between last anal ray and first median plate; about 25 radially striated plates on breast, 5 or 6 small ones on posterior part of gill membranes in front of breast, 1 on each side anteriorly, and 3 or 4 well-developed plates on front part of under side of mandible; intervals between ventrolateral behind vent with 12 to 15 small plates, mostly in pairs, none between ventrals and vent. Head more depressed and elongate than in *X. pentacanthus* and *latifrons*, its depth $\frac{4}{5}$ in width, the latter $8\frac{2}{3}$ in length of body; snout elongate triangular, nearly equal to long diameter of orbit. Orbits oval, the vertical $\frac{2}{3}$ of the longitudinal diameter, the latter 3 in head; a slightly curved row of 3 to 5 spines on eyeball above pupil. Interorbital space narrow, concave, without minor ridges, 9 in head. Supraocular ridges ending in a sharp spine; occipital low, with 2 pairs of spines, of which the anterior is low and blunt. Temporal moderate, ending in a sharp suprascapular spine; suborbital ridge moderate, close under orbit, with 1 spine on suborbital and ending in 1 at edge of preopercle; below the latter spine, another, broadly triangular; 3 or 4 well-developed spineless plates on lower part of cheek, which enters into ventral surface of head; nasal spines strong, sharp, far apart; median rostral plate small, movable, slightly overhanging, premaxillary (its lateral expansions as free spines) bearing a single upright spine; lower jaw a little included, maxillary not reaching orbit. Barbels probably varying much, for they differ markedly on the 2 sides of the jaw in the single specimen examined. Dorsals well separated; anal beginning about 1 plate behind the beginning of second dorsal; pectorals $5\frac{3}{4}$ in length of body, their base $\frac{3}{10}$ of their length, with 13 rays, the 4 lower exserted, separated from the rest by a notch, the 2 upper of these 4 longer than any of the other rays; ventrals of male $2\frac{1}{4}$ in head (in female probably shorter); width of caudal at base $2\frac{1}{3}$ in its length. Color olivaceous, with traces of darker cross bars on back, fins light or somewhat dusky, rays dark; ventrals pale in both sexes. Description taken from Dr. Gilbert's type. Total length reaches 7 inches. Coast of California to coast of Oregon, in 47 to 204 fathoms; a specimen also taken off Point Reyes in 75 fathoms. (τρεῖς, three; ἄκανθα, spine.)

Xenochirus triacanthus, Gilbert, Proc. U. S. Nat. Mus. 1890, 91, coast of California, at Albatross Station 2893, in 145 fathoms; Jordan & Starks, Proc. Cal. Ac. Sci. 1895, 827, pl. 93.

* Diagnosis: Body tapering uniformly from head to caudal, its depth $\frac{4}{5}$ of width at base of pectorals, the latter $8\frac{1}{3}$ in length; head 5 or a little less. Plates in dorsal series 39; between occiput and first dorsal 7; between dorsals 5; between ventrals and anal 15 pairs. Two barbels at tip of each maxillary, 1 pair at edge of the terminal mucous pore on under side of mandible, 1 at edge of next pore, and 1 at edge of third pore. Median rostral plate with a single upright spine. Nasal spines prominent; 1 pair supraocular and 2 pairs occipital spines. Teeth on jaws, vomer, and palatines. Gill membranes joined to isthmus, without free fold. Olivaceous, with traces of darker cross bars; fins light, the rays darker.

775. ODONTOPYXIS, Lockington.

Odontopyxis, LOCKINGTON, Proc. U. S. Nat. Mus., II, 1879, 328 (*trispinosus*).

Body very slender, depressed, tapering uniformly from head to caudal; head 3⅛ to 4½, the width of body 6 to 7, depth 6 to 8 in length of body. A deep pit behind occiput with a longitudinal ridge in its bottom; plates in dorsal series 35 to 40; both dorsal fins present; 4 (3) rays in first, 6 (7) in second; teeth numerous on jaws, vomer, and palatines. Gill membranes broadly joined to isthmus, without free fold; a single upright spine on median rostral plate; supraocular spines small, sharp; 2 pairs of blunt occipital spines or tubercles. (*ὀδούς*, tooth; *πύξις*, box.)

2427. ODONTOPYXIS TRISPINOSUS,* Lockington.

B. 6; 1 D. 4 (occasionally 3); 2 D. 6 (occasionally 7); A. 5 to 7; P. 14; V. 3 (I, 2); C. 1-11-1 (sometimes 1-10-1); pyloric cæca 5 to 7 (1 or 2 very small; 3 individuals). Body octagonal, the faces a little concave; spines small, sharp, present on the dorsolateral and superior lateral series nearly from head to caudal; a few very weak ones on inferior lateral; none on ventrolateral series; about 3 long, sharp spines on each side of base of caudal on the terminal plates. Two or 3 pairs of plates between last ray of second dorsal and first median plate, and 1 or 2 pairs between last anal ray and first median plate; 18 to 20 quite large, radially striated plates on breast, 6 or 7 of varying size in front of, and several minute spinous ones on base of pectorals; 3 or 4 strong plates on median part of gill membranes posteriorly, 1 to 3 on each side in front, 3 or 4 on under side of jaw, so that the whole of under side of head is well armored; 1 or 2 plates between ventrals and vent, the latter surrounded by small tubercles. Head somewhat depressed, its depth $\frac{7}{10}$ of its width, the latter 7 in length of body; orbits nearly circular, the longitudinal diameter about 3⅔ in head; a few minute spines on eyeball above pupil; snout pointed, equal to orbit; interorbital space concave, narrow, $\frac{5}{6}$ of longitudinal diameter of eye; occipital and temporal ridges weak. A deep transverse pit behind occiput, broader in front, its bottom divided longitudinally by the cranial ridge above the foramen magnum; suborbital ridge moderate, 1 spine on preorbital, and a very blunt flat one at posterior edge of opercle; 1 or 2 blunt flat projections below the latter spine; opercle with a ridge and 3 to 7 or 8 minute spines or prickles along its posterior border. Cheek below suborbital ridge entering into ventral surface of head, with 2 or 3 well-developed plates; several small plates in a longitudinal series behind orbit, above suborbital and preopercle, the highly movable median rostral plate with a single upright spine, its lateral expansions free, in

*Diagnosis: Body tapering uniformly from head to caudal, depressed, its depth about $\frac{2}{3}$ of its width (at base of pectorals), the latter 7 in length of body. Head 4½; a deep pit behind occiput, with a longitudinal ridge in its bottom. Plates in the dorsal series 35 to 37; between occiput and first dorsal 6 (occasionally 7) pairs; between dorsals 5 (occasionally 6); between ventrals and anal 12 or 13 pairs; teeth numerous on jaws, vomer, and palatines. One small barbel at tip of maxillary; gill membranes broadly joined to isthmus, without free fold. Median rostral plate with a single upright spine; nasal spines separated; supraocular spines small, sharp; 2 pairs of occipital in the form of low, blunt tubercles. Color olivaceous, with 6 or 7 darker cross bars.

form of small spines, slightly overhanging premaxillaries. Jaws equal, maxillary reaching front of orbit. Dorsals well separated; anal beginning about 1 pair of plates behind front of second dorsal; pectorals 5⅓ to 5½ in length of body, the base about 3 in their length, with 14 rays, of which the lower 5 to 7 are exserted, the depth of the notches in the membrane variable, the 2 or 3 uppermost of these exserted rays nearly as long (sometimes as long) as the longest upper rays; ventral fins of female about twice the interorbital space, about ⅔ of the ventrals of male; width of caudal at base 3 in the length. Lateral line 36. Color olivaceous, with 7 or 8 darker cross bars—1 in front of first dorsal, 1 under and extending into middle of first dorsal, 1 under front, and 1 under hind end of second dorsal, and 3 or 4 on caudal peduncle; a large bluish-black blotch on preopercle and opercle; first dorsal dark, the distal ⅓ white; second dorsal with 2 minute indefinite cross bars of darker; caudal dark at base, a broad dark band near its distal end, and the tip broadly white-edged; pectorals light, indistinctly cross-barred by series of dark streaks on the rays; ventrals pale in both sexes. Length about 3 inches. Coast of California, from Puget Sound to Santa Barbara; rather common in 11 to 57 fathoms. Here described from specimens taken off Point Reyes, where it is abundant. One specimen recorded by Lockington (loc. cit.) as having come from Alaska. (*tres*, three; *spinosus*, spined.)

Odontopyxis trispinous, Lockington, Proc. U. S. Nat. Mus., II, 1879, 328, San Francisco, California (Coll. W. N. Lockington); Jordan & Gilbert, Synopsis, 729, 1883; Jordan, Cat. Fishes N. A., 114, 1885.

776. BOTHRAGONUS, Gill.

Bothragonus, Gill MS., in Jordan & Gilbert, Synopsis, 728, 1883 (*swanii*).

Head and front of back rhombic as seen from above; body strongly compressed behind; length of head 3, width of head 2⅞, depth of body about 4⅔ in standard length. A deep pit at nape, several processes extending into it from its posterior edge. Mouth subinferior; posterior edge of opercle serrated. Both dorsals and anal weak, with short rays; anal under second dorsal; first dorsal 3; second dorsal 5; anal 3; no spines anywhere. Teeth on jaws and vomer (state of palatines not mentioned). Gill membranes broadly united to isthmus. (*βόθρος*, a pit; *Agonus*.)

2428. BOTHRAGONUS SWANII (Steindachner).

B. 5; D. III, 5; A. 5; P. 12; V. 3 (I, 2); lateral line 32. Head widening rapidly backward from the blunt snout, with 4 broad, low (occipital and temporal) ridges, between which the surfaces are concave. Body narrowing rapidly from head to beginning of first dorsal, from there to caudal strongly compressed. A deep transverse oval pit on mid-dorsal surface between occiput and nape; 3 delicate larger and several smaller processes extending into it from its posterior edge to about its middle. Length of head 3, its greatest breadth (over opercles) about 2⅞; depth of body under first dorsal about 4⅔, from the occipital pit to base of ventrals

a little more than 3 in length of body. Interorbital space slightly more than ¼ of head; snout 4; width of mouth about 3¼, orbit about 5⅜ in head. Snout rounded anteriorly, longitudinally arched, rising knob-like before the flat forehead, and extending a little beyond mouth, the latter broader at its angles than long, these a little in front of middle of eye. Teeth on jaws and vomer in several series, very small, conical, blunt at tips; plates on cheeks raised into tubercles; posteriorly the transversely flattened forehead is separated by a slightly curved, very low ridge from the occipital region; plates surrounding the occipital pit in part large and polygonal; last plate of head, above posterior end of opercle, extending outward as a broad, low pyramid; small bony plates embedded in skin on underside of head; preopercle spineless; edge of opercle scalloped and serrated. First dorsal beginning about middle of body; space between dorsals equaling distance from center of eye to tip of snout; spines of first dorsal but little shorter than rays of second dorsal; both dorsals and anal rounded; caudal about 2 in head, slightly rounded; base of large pectoral reaching to ventral surface, its longest ray 1⅜ in head; ventrals some distance behind pectorals, about 2 in the latter; distance between base of ventrals and front of anal a little more than the head; anal pit some distance behind base of ventrals. Plates of lateral line flat, those of remaining series considerably larger and elevated at their centers, forming 4 series of blunt processes or ridges; 14 pairs of plates between ventrals and anal, 5 between hinder edge of occipital pit and first dorsal (6 from occiput to first dorsal); 7 between second dorsal and caudal; ventral surface narrowing rapidly from head to anal and anteriorly convex, posteriorly flat. A blackish-gray cross band surrounds the head, wreath-like, its outlines indefinite in places, passing transversely across forehead and from lower edge of orbit completely around edge of mouth; band under first dorsal passing obliquely forward to near insertion of ventrals; the next band falling between the dorsals, and the third under second dorsal; both are vertical; pectorals with a narrow dark cross bar at base and a much broader marbled one occupying nearly the whole distal half of fin; ventrals with 3 or 4 bands of dark spots; on both dorsals and anal oblique cross bars of dark spots; caudal with numerous cross bars of dark spots, which increase in intensity of color distally and flow together and leave isolated spots of lighter; ground color of body light yellowish with a tinge of brownish. Length of the single specimen 1 inch 11 lines; found dead on the beach after a storm. (After Steindachner.) The following points may be added from the 4 excellent figures: The ventrolateral series lie entirely on ventral surface of body anteriorly and meet in front of vent; 2 pairs of plates between ventrals and vent; a single plate between bases of ventrals, 4 plates in pairs in front of them on breast, and these preceded by 2 single plates; 10 or 12 rather small plates in about 3 rows in front of base of pectorals; several plates on lower part of cheek and a longitudinal series of them behind orbit, above suborbital and preopercle. Puget Sound; known from single specimen. This species has been erroneously referred to the genus *Hypsagonus*, because of its short compressed body and peculiar form of head.

A very careful comparison of the description and the excellent figures with *Odontopyxis trispinosus*, Lockington, with which Steindachner was probably not acquainted, makes it evident that in spite of its aberrant form, the *Hypsagonus swanii* is more closely related to it than to any other known species. There is in reality no point of resemblance to *H. quadricornis* except the shortness of the body and the compressed condition of its posterior part. In all of the following points, some of which Steindachner mentioned as differences between his species and *H. quadricornis*, *B. swanii* agrees with *O. trispinosus:* Both dorsals and anal small; the distance of the first dorsal from nape and the distance between dorsals (measured by the number of plates); the deep pit at nape; position of vent, with 1 or 2 pairs of plates between it and the base of the ventrals; 2 or 3 series of small plates between base of pectoral and gill opening; small plates on gill membranes and lower jaw; the longitudinal series of small plates behind orbit; the coloration, etc. The only known specimen was less than 2 inches (50 mm.) long and was doubtless a young one. The most marked characters which distinguish *B. swanii* from *Odontopyxis* are the shortness of the body, the wideness of the body in front, and the compression of the tail; all are characters which distinguish the young from the adult in all species of the family in which the young have been seen and described, except that here these characters seem exaggerated. (Named for James G. Swan, of Port Townsend, Washington, a veteran observer and collector.)

Hypsagonus swanii, Steindachner, Ichth. Beiträge, v, 144, pl. 4, Sitzb. der Akad. Wiss., lxxiv, July, 1876, Port Townsend. (Coll. James G. Swan. Type in Coll. Mus. Vienna.)
Bothragonus swanii, Gill, in Jordan & Gilbert, Synopsis, 728, 1883; Jordan, Cat. Fishes N. A., 114, 1885.

777. ASPIDOPHOROIDES, Lacépède.

Aspidophoroides, Lacépède, Hist. Nat. Poiss., iii, 228, 1802 (*tranquebar=monopterygius*).
Canthirhyncus, Swainson, Nat. Hist. Fishes, etc., ii, 272, 1839 (*monopterygius*).
Anoplagonus, Gill, Proc. Ac. Nat. Sci. Phila., xiii, 1861, 259 (*inermis*).
Ulcina, Cramer, in Jodran & Evermann, Check-List Fishes, 449, 1896 (*olriki*).

Body and head more or less slender; head 4 to 6, width 5 to 8 in length of body; 8 longitudinal rows of plates, the lateral line in the upper lateral row; about 40 plates in the dorsal series. Terminal rostral plate present, unarmed; mouth terminal; teeth on jaws, vomer, and palatines. Supraocular and occipital spines absent; plates of body more or less keeled, without spines. First dorsal absent; second dorsal and anal small, opposite each other, each with 4 to 7 rays. Gill membranes united, narrowly joined to isthmus anteriorly, free behind. (ἀσπίς, shield; φορέω, to bear; εἶδος, form.)

Ulcina (*ulke*, a sculpin, in Danish):
a. Body rather robust, the depth about 5 in length, the head 4 in length; plates in lateral line usually less than 40.
b. Lateral line with 40 plates; nasal spines well developed. olriki, 2429.
bb. Lateral line with 36 to 38 plates; nasal spines very small or obsolete. guntheri, 2430.

aa. Body very slender, the depth about 8 in length, the head 5 to 6; plates in lateral line more than 40.

c. Nasal spines present; no median dorsal row of plates behind occiput.

ASPIDOPHOROIDES:

d. About 48 (or fewer) plates in dorsal series; temporal ridge ending posteriorly in a slight elevation. Atlantic coast.

MONOPTERYGIUS, 2431.

dd. 49 or 50 pairs of plates in dorsal series; temporal ridge not ending posteriorly in a slight ridge. Pacific coast. BARTONI, 2432.

ANOPLAGONUS (ἄνοπλος, unarmed; *Agonus*):

cc. Nasal spines absent; a median dorsal row of small plates from occiput halfway to dorsal. INERMIS, 2433.

Subgenus ULCINA, Cramer.

2429. ASPIDOPHOROIDES OLRIKI, Lütken.

D. 6 or 7; A. 6 or 7; P. 13 or 14; V. 3 (I, 2); C. 10 (Lütken). Form short and thick compared with *A. monopterygius*, depressed anteriorly, posteriorly narrower and somewhat angular. Nape and fore part of back concave; forehead strongly arched antero-posteriorly. Interorbital space concave, supraorbital ridges high. Ventral surface flat. Greatest breadth, over opercles, $\frac{3}{4}$ of length of head, the latter $4\frac{1}{5}$ to $4\frac{1}{2}$ in total length, its greatest height, at shoulder, $\frac{2}{3}$ of length of head. One pair of nasal spines; a single barbel at angle of mouth (at tip of maxillary, as in other *Aspidophoroides*,) overlooked in the original description. Pectorals large, about as long as head; other fins small; all rays simple; anal under dorsal, about midway between head and caudal; ventrals inserted a little behind base of pectorals. Eyes large, their diameter $\frac{1}{3}$ or a little less in length of head, longer than snout. Mouth small, its angle in line with anterior edge of orbit. Vent large, lying with its surrounding naked area about 2 pairs of plates distant from base of ventrals. Gill membranes united. Eight longitudinal rows of plates on body, 6 on tail. In dorsal series 36 to 38, the posterior 12 or 13 on tail, median unpaired; in front of dorsal, 14 to 16 pairs. Superior lateral (lateral line) series 36 to 38; inferior lateral series 35 to 37; on breast 12 to 14, paired and unpaired. Two unpaired plates between vent and ventrals (from ventrals to anal about 12 pairs). Teeth on jaws, vomer, and palatines. Fin rays, dorsal 6, anal 6, in 3 individuals, 5 in each fin in the fourth individual (6 or 7 in each fin in the Greenland specimens); pectorals 13 to 15 (in 2 specimens, 14 in 1 fin and 15 in the other); ventrals 3 (I, 2); caudal 10 or 11, the eleventh very short. Color more uniform dark gray on back than those previously described (a few very indefinite dark cross bands); a large dark patch on preopercle and opercle seems to be quite constant; the dark spots on pectorals and caudal form cross bands when the fins are folded; tips of dorsal, anal, and ventral fins in general white, but the white patch is wanting in 1 specimen, which is paler than the others, has a smaller head and shorter ventrals. This is undoubtedly a female; the others males. Holm's notes on the color of the newly caught specimens say, "Sides of back greenish, with brown markings, otherwise whitish." Arctic Ocean; known from west coast of Greenland, Davis Strait, at 32 fathoms; our specimens from Godhavn, Kara

Sea, at 55 fathoms; Barents Bay, Nova Zembla. (Eu.) The Dutch expedition took 15 specimens of this form in Barents Bay and probably confused it with *A. monopterygius*. Their length was 40 to 75 mm. Later, on the *Fylla* expedition, Holm took 9 specimens in Davis Strait on the whitefish banks north of Holstenborg at a depth of 32 fathoms. Their length varied from 62 to 86 mm. The dark color bands in these are more or less distinct, at least among the smaller specimens. The ventrals varied a little in length, those with the short ventrals apparently in a decided minority. The small barbels were present in all of them. Rays of dorsal and anal fins: D. 6; A. 6 (3 specimens); D. 6; A. 7 (1 specimen); D. 5; A. 6 (1 example); D. 6; A. 5 (1 example); D. 7; A. 6 (3 examples). That the white patch on the dorsal (and anal and ventral) is invariably present on the individuals with longer ventrals is undeniable. The assumption that it is a sexual distinction of the male thus gains probability; this was demonstrated in the specimens that were opened. Total length 63 to 69 mm. (4 individuals). (Lütken.)

Aspidophoroides olriki, LÜTKEN, Forelöb. Meddel. om Nord. Ulkefiske; Vidensk. Meddel. Naturhist. Foren. Kjöb. 386 (with 3 figures), 1876, Greenland; JORDAN, Cat. Fishes N. A., 11$\frac{3}{4}$, 1885; LÜTKEN,* Kara-Havets Fishe, 6, pl. 15, figs. 1–3, 1886; specimens from Kara Sea.

2430. ASPIDOPHOROIDES GUNTHERI, Bean.

D. 7; A. 7; V. I, 2; C. 10; P. 12. Body short, anteriorly very wide, somewhat depressed. Height of body 6, and its width 5 in its length; head 4$\frac{1}{8}$, its depth not more than $\frac{1}{2}$ of its width, the latter nearly 5 in length of body; triangular, very short, wide posteriorly. A small barbel at tip of each maxillary; teeth in jaws, vomer, and palatines; nasal spines almost invisible. Along sides of head inferiorly 4 large mucous pores in oblong depressions, the largest of which is nearly as long as snout. Eye 3 in head, considerably more than width of interorbital space. Maxillary not reaching beyond anterior border of orbit; mandible barely included; snout equaling interorbital space, the latter deeply concave. Greatest width of head nearly 5 in length of body, and 2 from beginning of dorsal to base of caudal. Gill membranes narrowly attached to isthmus anteriorly, free posteriorly; gill opening wide. A deep groove along anterior third of back. Pectorals 2$\frac{1}{2}$ times as long as ventrals and 4 in length of

* The following is the substance of Dr. Lütken's original account from specimens from Greenland: Body short, thick, total length 72 to 75 mm.; head 4$\frac{1}{2}$ in total length (including caudal fin); width of body (at base of pectorals) a little more than 5 in total length; angles of body not distinctly serrated; interorbital space and back concave; nasal spines 2; diameter of orbit 3 in head, nearly twice the width of interorbital space or length of snout; plates in dorsal series 35 to 37; in front of dorsal fin 14 to 17; superior lateral series 39; inferior lateral 37; ventrolateral 36; in front of anal 12 or 13. First dorsal fin absent; teeth present on vomer and palatines; only 8 longitudinal series of plates, the lateral line on the superior lateral series; the ventrolateral series not forming the inferior angles of the body, but confined to the ventral surface and meeting in a single plate between ventrals and vent.

The above diagnosis, made in 1875 from the Greenland specimens taken from the stomachs of flounders, was not accompanied by a detailed description. The description was made ten years later from specimens taken in the Kara Sea. The figures of the individuals from Kara Sea differ from the figures of specimens from Greenland in several respects. On the former the anterior part of the back just behind the nape is more elevated; the rostral spines are directed much more backward and seem much less prominent; and there is 1 less median plate on breast (this last difference an individual one in other species). The gill membranes are probably free, at least behind.

body; ventrals as long as head without postorbital part; vent nearer tips than bases of ventrals; beginning of dorsal nearly midway of total length; anal under dorsal but a little shorter; length of caudal 5½ or 6 in length of body; second dorsal ray longest, equal to postorbital part of head; longest anal ray less than ½ of head. Lateral line 40. Plates on breast about 14. Color dusky above, whitish below; a dark stripe on snout, continued on opercle; a few indistinct dark blotches on side of head; axillary region dusky; pectoral with 3 or 4 imperfect cross bars; body with 3 indistinct dusky cross bars, the middle one extending on middle of dorsal fin; tail with 2 dusky bars, 1 anteriorly, the other terminal, with a dirty yellowish area between them; ventrals and anal pale. Probably a young individual. Longest known individual, which served as the type, was 70 mm. (about 2¾ inches). (Bean.) Coasts of northern Alaska. (Named for Dr. Albert Günther, keeper of the British Museum of Natural History.)

Aspidophoroides güntheri,* BEAN, Proc. U. S. Nat. Mus. 1885, 74, northern Alaska (Type, No. 37032. Coll. Lieut. G. M. Stoney); JORDAN, Cat. Fishes N. A., 113, 1885.

2431. ASPIDOPHOROIDES MONOPTERYGIUS† (Bloch).

(SEA POACHER.)

Br. 6; D. 5 or 6; A. 5 or 6; P. 9; V. 3 (I, 2); C. 10 or 11; Pyloric cæca 4 or 5 (2 individuals). Width of body at base of pectorals 8 in length; dorsal face concave from occiput to dorsal; ventrolateral ridges fairly prominent, but these series lie entirely on ventral side of body anteriorly and converge to a single plate between ventrals and vent. The ridge of the inferior lateral series, instead of disappearing anteriorly, is lower down than usual, becomes prominent anteriorly and runs to lower end of base of pectorals, thus forming the inferior angle of body, which is markedly 4-hedral in front. Plates in dorsal series 46 to 48; between ventrals and anal 16 or 17 pairs; between last ray of dorsal and first median plate 1 to 3 pairs; 2 or 3 pairs between last anal ray and first median plate. A small plate between ventrals, a larger one between ventrals and vent; a variable number (3 to 8) of small plates behind and beside vent; plates of breast radially striate, the two median ones larger than the rest; a few small plates on hinder median part of gill membranes, and on narrow under side of mandible; 2 large plates with raised centers in front of and

* This species agrees with the measurements of *A. olriki*, Lütken, and with the coloration as far as it is possible to judge; it differs from *A. olriki* only in these recorded characters: The nasal spines are very small, the longest anal ray is shorter than in the figures of *A. olriki*, and the pectoral rays number 12, while they range from 13 to 15 in *A. olriki*. But in none of these respects does it differ more from Lütken's descriptions and figures of *A. olriki* from Greenland and from Kara Sea, in some respects apparently less, than these differ from each other. Each of the localities is about ⅛ of the earth's circumference from the other, and *A. guntheri* may be only a variety of *olriki*, which would then be a circumpolar species; if this view should ultimately prove untenable, Lütken's Kara Sea specimens will have nearly as good a claim to specific distinction as the type of *A. guntheri* has.

† Diagnosis: Body very slender, depressed in front, nearly 4-hedral anteriorly; depth ⅔ of width, the latter 8 in length; caudal peduncle very long, 6-hedral subterete; head long, tapering forward, 5⅞ to 6 in length of body; plates in dorsal series about 48; between occiput and dorsal fin 19 to 21; from ventrals to anal 16 or 17 pairs; first dorsal absent; barbels none; gill membranes free; median rostral plate small, unarmed. Nasal spines very strong, diverging; no other spines on head or body. Color, brownish, obscurely banded with dark.

4 to 8 small spineless plates in 1 or 2 series on base of pectoral. Head long, tapering, 5⅞ to 6 in length of body; orbits very large, nearly circular, the longitudinal diameter longer than snout, 3½ in head; interorbital space wide, less than orbit, deeply concave; supraocular ridges high, bulging; occipital ridges evident; temporal ridges low, terminating in an upwardly directed blunt elevation; suborbital ridge close under orbit, low, spineless, 2 or 3 plates below it on the nearly vertical cheek; median rostral plate small, immovable or slightly movable, spineless, overhanging premaxillary; mouth small, terminal, lower jaw a little included; maxillary reaching front of orbit, almost entirely covered by preorbital. Dorsal and anal fin small, the latter under the former; caudal 3 times as long as wide at base; pectorals 6 or less in body, their base about 5 in their length; ventrals of female ⅔ those of male, the latter ⅔ of pectorals. Lateral line 50. Color brownish, pale below, with indefinite cross bands of darker, 2 in front of, and 1 under dorsal, and 2 or 3 on caudal peduncle; rays of dorsal and upper rays of pectoral brownish, interrupted by lighter, giving an indefinite cross-banded appearance; caudal dark, ventrals in both sexes and anal pale. Total length 6 inches. Greenland to Cape Cod; common in rather deep water. This description based on specimens from Massachusetts Bay. (*μόνος*, single, *πτερύγιον*, fin.)

Cottus monopterygius, Bloch, Ichthyologia, II, 156, pl. 178, figs. 1, 2, 1786, Tranquebar; an error.

Cottus indicus, Bonnaterre, Tableau Encycl. Meth., 68, pl. 87, fig. 367, 1788, India; after Bloch.

Aspidophoroides tranquebar, Lacépède, Hist. Nat. Poiss., III, 228, 1802 Tranquebar; after Bloch.

Aspidophoroides grœnlandicus, Valenciennes, in Cuvier, Régne Anim., Ed. II, Vol. 2, pl. 21, fig. 3, 1829, Greenland.

Aspidophoroides monopterygius, Cuvier & Valenciennes, Hist. Nat. Poiss., VI, 554, pl. 169, 1830; Storer, Fishes of Mass., 22, pl. 1, fig. 1, 1867; De Kay, New York Fauna: Fishes, 62, pl. 3, fig. 5, 1842; Günther, Cat., II, 216, 1860; Jordan & Gilbert, Synopsis, 724, 1883.

Canthirhyncus monopterygius, Swainson, Nat. Hist. of Fishes, etc., II, 272, 1839.

2432. ASPIDOPHOROIDES BARTONI, Gilbert.

Very close to *Aspidophoroides monopterygius*. From this species it differs in the much lower ridges on body, which are slightly rounded rather than sharply carinate, and leave the intervening faces shallowly concave, instead of deeply so. The plates of the lower lateral ridge do not in the young bear backwardly directed spinous points as they do in *A. monopterygius*. The keel in front of the ventral fin, prominent in *A. monopterygius*, is here nearly or quite obsolete. The same is true of the keel below the eye, and the occipital ridge. The plates in front of the pectorals also protrude less, and in general the angles and prominences are less marked. The space between the dorsal ridges is less, its greatest width being ½ length of head behind middle of eye in adults, and behind posterior margin of eye in young. In *A. monopterygius* the same width equals ½ head behind front of eye in adults, behind middle of eye in young. These scarcely admit of quantitative statements, but are sufficiently evident on comparing specimens from the Atlantic and the Pacific. Vertebræ

11 + 39 = 50. General proportions, fin rays and color are the same in the 2 species. Aleutian Islands; taken very abundantly both north and south of the Aleutian Islands and in Bristol Bay, at depths of 17 to 121 fathoms. (Gilbert.) Also found about the Pribilof Islands. (Named for Mr. Barton A. Bean.)

Aspidophoroides bartoni, GILBERT, Rept. U. S. Fish. Comm. 1893 (1896), 434, Aleutian Islands; Bristol Bay, Alaska. (Coll. *Albatross*.)

Subgenus **ANOPLAGONUS**, Gill.

2433. ASPIDOPHOROIDES INERMIS,* Günther.

B. 6; 1 D. absent; 2 D. 5 or 6; A. 4 or 5; P. 10; V. 3 (I, 2); C. 11; vertebræ 11 + 32 = 43. Pyloric cæca 4 (1 individual). Only 8 longitudinal rows of plates, the lateral line in the upper lateral row. No spines anywhere on body; no ridges on the superior and inferior lateral series; plates of the dorso- and ventrolateral series bent upon themselves at right angles, forming the 4 prominent angles of the body; the ventrolateral series do not therefore lie entirely on underside of body, as in *A. monopterygius*, nor do they converge between ventrals and vent, but pass forward to lower end of base of pectorals; three pairs of plates between last dorsal ray and first median plate, and 2 to 4 pairs between last anal ray and first median plate; a median dorsal series of hexagonal plates about as large as those of the lateral series, on widest part of back, from occiput to about halfway to dorsal; these plates alternating, but not regularly, with still smaller ones; an inconstant number of small plates behind and around vent; 2 or 3 pairs between ventrals and vent, and a single small one between the bases of ventrals. Gill membrane medially on its posterior part and laterally on its anterior part and narrow under surface of mandible between the rami strongly plated; two or 3 large plates in front of and 7 to 10 small ones in 2 series on base of pectoral. Head depressed, its depth $\frac{3}{4}$ of its greatest width; orbit small, circular, less than snout, $4\frac{1}{2}$ in head; interorbital space narrow, $1\frac{2}{3}$ of orbit, moderately concave; supraocular, occipital, temporal, and suborbital ridges very low; 4 or 5 plates on lower part of cheek; median rostral plate small, vertical, movable, unarmed; mouth small, the lower jaw slightly projecting, maxillary scarcely reaching front of orbit, entirely covered by preorbital; teeth in bands on jaws, vomer, and palatines. Dorsal and anal fins small, the latter under the former, with usually 5 rays; caudal rounded, its width at base $2\frac{1}{2}$ in its length; pectorals 7 in length of body, their base $2\frac{1}{2}$ to 3 in their length, nearly truncate, with 10 rays, the 3 lowermost a little exserted. Ventrals of female nearly twice as long as orbit, $\frac{5}{6}$ to $\frac{3}{4}$ as long as those of male, the latter less than 2 in pectorals. Lateral line 41. Color dark (nearly black) grayish or brownish and lighter

* Diagnosis: Body slender, much depressed anteriorly, 4-hedral, the dorsal face concave, the ventral flat, the lateral convex; caudal peduncle 6-hedral, subterete; depth of body $\frac{3}{4}$ in width, the latter 8 in length; head long, tapering, $5\frac{1}{4}$ in length of body; plates in dorsal series 42; from occiput to dorsal 19 to 21; from ventrals to anal 17 pairs; a median dorsal row of small plates from occiput to opposite tip of pectorals; first dorsal absent; barbels none; gill membranes narrowly joined to isthmus with moderate free fold behind; median rostral plate small, unarmed; nasal and all other spines absent from head. Color brownish black, with several alternating cross bars of darker and lighter. Pectorals with black patch at base and 1 near tip.

gray, dusky below; a black band from mouth across orbit to pectoral and alongside of body, much interrupted; dark markings deepest on sides of body; a dark cross band halfway between occiput and dorsal, 1 in front of dorsal, 1 under posterior half of dorsal (running up on latter), and about 2 on caudal peduncle; posterior half of dorsal dark, marbled; a marbled patch at base of pectorals, another on their distal third; ventrals pale in both sexes; anal with dark patches on or behind the rays, darker posteriorly; caudal black at its base and on its distal half, the 2 patches connected by a black band along middle of fin, leaving 2 light patches on the dorsal and ventral thirds of the proximal half of the fin; the bands of color very much as in *A. monopterygius*, but better defined. Length 4 inches. Coast of Alaska, south to Vancouver Island; not abundant; recorded from Vancouver Island (Günther); eastern Aleutian Islands, 34 to 59 fathoms, and Bristol Bay. (Gilbert.) Here described from specimens from the Aleutian Islands and Bristol Bay. (*inermis*, unarmed.)

Aspidophoroides inermis, GÜNTHER, Cat., II, 524, 1860, Vancouver Island; LÜTKEN, Forelöb. Medd. om Nord. Ulkefiske: Vidensk. Meddel. Naturhist. Foren. Kjöb., 385, 1876; JORDAN & GILBERT, Synopsis, 725, 1883; JORDAN, Cat. Fishes N. A., 113, 1885.
Anoplagonius inermis, GILL, Proc. Ac. Nat. Sci. Phila., XIII, 1861, 167.

Family CLXXXII. CYCLOPTERIDÆ.*

(THE LUMP SUCKERS.)

Body short and thick, more or less elevated, covered with a thick skin, which is smooth, tubercular, or spinous; head short and thick; suborbital

* Dr. Gill thus defines the superfamily *Cyclopteroidea*, including the *Cyclopteridæ* and the *Liparididæ*, the group being equivalent to that called by other writers "*Discoboli*":

"Acanthoptergians with the third infraorbital bone developed as a stay obliquely crossing the cheek and connecting with the preoperculum, the myodome suppressed, the post temporal bifurcate and normally connected with the cranium, the actinosts enlarged and mostly connected with the inner ridge of the proscapula, the hypercoracoid being dislodged upward and the hypercoracoid downward on a row with the 4 actinosts; ribs sessile on the vertebral centra or hæmapophyses; pharyngials reduced to the large epipharyngeal (homologous with the third of typical Acanthoptergians), and ventrals modified to form a suctorial disk supported by 6 immovable rays on each side converted into osseous tissue and without articulations; typically suppressed. They appear to have the branchial apparatus constructed on the same plan as in the *Cottoidea*; 2 or 3 basibranchials ossified; hypobranchials of 3 pairs in line with the corresponging ceratobranchials of fourth arch suppressed; ceratobranchials of all and epibranchials (of all or 3) arches well developed; pharyngobranchials reduced to 1 pair of compressed epipharyngeals; hypopharyngeals divergent and rather compressed. There are 3½ gills, that is double branchiæ, on all the arches except the fourth, which has a single row of filaments. There is no fissure behind the fourth arch." (Gill.)

Mr. Garman (Monograph of the Discoboli, 1892, 19) has the following remarks on the *Cyclopteridæ*:

"Anteriorly the form of the lump fishes is stout, thick, and deep; behind the body cavity, which occupies the greater portion of the length, it rather abruptly becomes weak and slender. The head is short and broad, subquadrangular in transsection; the snout is short and blunt; the mouth is of moderate width, anterior, and opens slightly upward; the teeth are small, subconical, and arranged in a band or cord; the eyes are of medium size and have a lateral outlook. All of the members of the family have pseudobranchiæ, 3½ gills, 6 branchiostegal rays, small gill openings, numerous pyloric cæca, and an elongate intestine. In their lower portions the broad, rounded pectorals extend forward under the throat, along the sides of the disk. The vertical fins are not of large extent; the caudal and the 2 dorsals are quite separate. The disk is comparatively large. Early in life the skin is tender and naked; later it grows tough and is covered with roughened or spine-bearing osseous tubercles. Semicartilaginous describes the skeleton with tolerable accuracy, the small amount of bony matter lies in thin plates, often forming cells and chambers similar to those to be noticed in the bones of *Lophius*. The third

stay present, thin and flattish; mouth small, terminal; jaws with bands of slender, simple teeth; no teeth on vomer or palatines; gill openings narrow, restricted to the sides, the membranes being broadly joined to the isthmus and shoulder girdle; branchiostegals 6; gills 3½; pseudobranchiæ present; dorsal fins 2, the anterior part of flexible spines, which, in the adult, are sometimes hidden by a fleshy hump, in 1 subfamily entirely wanting; soft dorsal usually opposite the anal and similar to it; caudal fin rounded, free from the dorsal and anal; ventrals thoracic, rudimentary, forming the bony center of a sucking disk; pectorals short, placed low, their bases broad and procurrent; pyloric cæca numerous; intestine elongate; vertebræ 12+16, the skeleton feebly ossified. Genera 7; species 8; inhabiting the northern seas of both hemispheres. By means of the adhesive ventral disk these fishes are enabled to attach themselves very firmly to rocks or other objects. They feed on crustacea, worms, small fishes, and plants. The young of *Cyclopterus* bear a close resemblance to *Liparis*, an evidence of the common origin of the 2 groups, which is borne out by the anatomy. (*Discoboli;* group *Cyclopterina*, GÜNTHER, Cat., III, 154 to 158.)

CYCLOPTERINÆ:

I. Spinous dorsal present, sometimes concealed in adult.

a. Barbels none; disk anterior, below the head.

b. Skin armed with large tubercles.

c. Larger tubercles on body in rows well separated; spinous dorsal fin obsolete with age. CYCLOPTERUS, 778.

cc. Larger tubercles not in rows, crowded closely together; spinous dorsal fin usually not disappearing with age. EUMICROTREMUS, 779.

bb. Skin naked or with scattered, slender spines; no lateral line. LETHOTREMUS, 780.

aa. Barbels present about the mouth; disk below the abdomen; tubercles on sides minute. CYCLOPTEROIDES, 781.

LIPAROPSINÆ:*

II. Spinous dorsal wholly wanting.

e. Dorsal short, opposite anal, of 9 or 10 rays; skin wholly smooth, without bony tubercles. CYCLOPTERICHTHYS, 782.

ee. Dorsal long, beginning near middle of back; skin with bony tubercles. LIPAROPS, 783.

---

suborbital is thin and broadens as it extends back to the preopercle. The interopercle is more blade-like than in the *Liparididæ*. On 1 genus the pores around the mouth are tubular and form barbels. In this important paper are valuable notes on the osteology and embryology of the *Cyclopteridæ* and *Liparididæ*.

* The *Liparopsinæ*, regarded as a separate family (*Liparopsidæ*) is thus defined by Mr. Garman:

"The deterioration of the first or spinous dorsal seen in the Lumpfish, has apparently proceeded so far in the *Liparopsidæ* as to cause the disappearance of that fin. Each of the 2 genera in the family has but a single dorsal, the posterior. In one genus the fin is short and situated near the caudal, in the other the fin begins near the middle of the back and extends nearly to the origin of the caudal. The genera are further distinguished by dorsal tubercles in one case, and by a naked skin in the other. The shape is somewhat like that of the Diodons, bulky, thick, broad, and longer forward, in the section containing the visceral cavity, and short and greatly reduced in size behind it. The head is short, broad, and thick, the snout short and blunt, the mouth terminal, the teeth subconical, the eyes lateral, the branchiostegal rays 6 in number, the gill openings narrow, the gills 3½, the pseudobranchiæ small, all, with disk, pectorals, and caudal, as in the *Cyclopteridæ*. Until recently this family has been known only from the North Pacific. A short time ago a second species of *Cyclopterichthys* was named by Vaillant from a sketch of a fish taken in the Straits of Magellan, which would extend the distribution to the Antarctic regions." (*Liparopsidæ*, Garman, Discoboli, 40, 1892.)

778. CYCLOPTERUS (Artedi) Linnæus.

(Lumpfishes.)

Lumpus, Gesner, Hist. Anim., iv, 1284, 1588 (nonbinomial).
Cyclopterus, Artedi, Genera Pisc., 62, 1738 (nonbinomial).
Cyclopterus, Linnæus, Syst. Nat., Ed. xi, 260, 1788 (*lumpus*).
Lumpus, McMurtrie, in Cuvier, Anim. Kingd., 1831 (*vulgaris*).

Body more or less compressed toward the back, somewhat triangular in a transverse section at the first dorsal, covered with conical, rough, bony tubercles; head short, thick, subquadrangular in a cross section; snout blunt, rounded; mouth anterior, opening slightly upward; teeth simple, small, arranged in a band; eye moderate, lateral; dorsals 2; caudal distinct; disk moderately large, anterior below the head; no barbels about the mouth; first dorsal fin in the adult completely hidden by the skin, the larger tubercles of the flanks, though in regular series, having a scattered appearance. One species, reaching a considerable size, in the north Atlantic. (*κύκλος*, circle; *πτερόν*, fin.)

2434. CYCLOPTERUS LUMPUS, Linnæus.

(Lumpfish; Cock and Hen Paddle; Lump Sucker.)

Head 5 in entire length; depth 2; D. VI * to VIII, 11; A. 9 or 10; ventral 6; pectoral 20; caudal 12 to 14; branchiostegals 6; vertebræ 11 + 18 = 29. Body massive, compressed, subtriangular in transverse section through the middle, belly flattened, the portion behind the abdominal chamber much compressed, and less than ½ the length of the body proper. Head short, subquadrangular in transverse section, forehead broad, flattened; nape high; snout short, broad, blunt; mouth wide, anterior, opening with a slight upward direction, extending backward almost to a vertical from the front margin of the eye. Eyes lateral, near the top of the head, as long as the snout, ¼ as long as the head and ⅓ as wide as the interorbital space. Forehead broad, depressed, convex. Nostrils small, the hinder smaller, near the eyes on interorbital space, the anterior farther forward, halfway to the mouth, with a short tube. Gill opening moderately wide, its lower third in front of the base of the pectoral. Fins with rounded margins, rough, with small tubercles. First dorsal distinct on very young individuals, variable in shape, thick and fleshy, with weak rays in older stages; second dorsal distinct, broad, rounded; caudal broad, subtruncate or rounded posteriorly; anal moderate, opposite the second dorsal, which it resembles in shape; pectorals broad, rounded, fringed, not indented at the sides of the disk. Disk little longer than wide, about the width of the head, or nearly ¾ of the length of the latter. Skin thickly sown with small, irregular subconical tubercles, the sides of which are roughened with small, conical protuberances. On older individuals, larger, longitudinally compressed tubercles form a vertical series from the nape over the first dorsal; a series of 3 others stands at each side

* The following formula is given for British specimens: Dorsal IV to VII, 10 or 11; anal 9 or 10; pectoral 20 or 21, ventral 6; caudal 10 or 11.

of the space between the dorsals; a row of larger ones extends from the supraorbital region along the flank to the upper part of the tail; a series, starting a little above the pectoral, passes to the lower portion of the tail; and a third lateral series reaches along each line of the lower surface from the side of the disk to the anal. The fleshy ridge enveloping the first dorsal is subject to considerable variation; it usually continues forward on the nape and becomes indefinite at the occiput. In alcohol the colors are brownish or olive to grayish, the tubercles darker. In life the tints vary from yellowish or greenish in the young to more or less brilliant red in the males, or bluish to dark brown in females; spots, blotches, cloudings, or other markings are not infrequent. The young often take the color of their surroundings.

Mr. Dresel gives the following notes on a small example of this species, 31 mm. long, from Davis Straits:

The spinous dorsal is comparatively high, and is not enveloped in thick skin as in the adults. The abdominal tubercles are the most developed. The gill opening is as long as the base of the anal fin, which is as long as the disk. D. IV, 10; A. 10. Only 1 specimen, badly mutilated by the Eskimo dogs, was seen at Godhavn. The color was bright olive green, with the belly white. These fish are seldom caught in this harbor after May, during which month they are very abundant.

North Atlantic, on rocky shores of both coasts; south to Cape Cod and France; generally abundant, reaching a length of 20 inches or more, but usually much smaller; they are rarely used as food. According to Garman, the Lumpfish spawns near shore in March or April, after which the female retires to deep water, leaving the male to watch the eggs, which hatch among seaweed and eelgrass. (Eu.) (*lumpus*, from the English lump, the fish having been called *Lumpus anglorum* by Gesner.)

Cyclopterus lumpus, Linnæus, Syst. Nat., Ed. xi, 1, 260, 1788, Baltic and North Sea; Günther, Cat., iii, 155, 1861; Dresel, Proc. U. S. Nat. Mus. 1884, 250; Jordan & Gilbert, Synopsis, 747, 1883; Garman, Discoboli, 21, 1892, and of authors generally.
Cyclopterus minutus, Pallas, Spicilegia, Zool., vii, 12, 1769, Atlantic Ocean.
Cyclopterus pavoninus, Shaw, Nat. Misc., ix, pl. 310, 1797.
Cyclopterus pyramidatus, Shaw, Gen'l Zool., v, 390, 1804.
Gobius minutus, Müller, Zool. Dan., iv, 38, 1808, Denmark.
Cyclopterus cæruleus, Mitchill, Trans. Lit. and Phil. Soc. N. Y., i, 1815, 480, New York Harbor.
Cyclopterus coronatus, Couch, Nat. Hist. Cornwall, 47, 1823, Cornwall.
Lumpus vulgaris, McMurtrie, in Cuvier, Anim. Kingdom, 1831; after Linnæus.
Lumpus anglorum, De Kay, N. Y. Fauna: Fishes, iv, 305, 1842; after Linnæus.

779. EUMICROTREMUS, Gill.

Eumicrotremus, Gill, Proc. Ac. Nat. Sci. Phila. 1864, 190 (*spinosus*).

Body and head covered with spines and tubercles, not arranged in regular rows; first dorsal not hidden so early in life as in *Cyclopterus*. Eyes and disk larger than in *Cyclopterus*, the gill openings rather smaller. Dorsal rays VI or VII, 10 or 11. Size small. Northern seas. (εὖ, very; μικρός, small; τρῆμα, aperture.)

a. Body rather elongate, the depth 2¼ in length; tubercles large. SPINOSUS, 2435.
aa. Body short and deep, the depth ½ length; tubercles smaller. ORBIS, 2436.

2435. EUMICROTREMUS SPINOSUS (Müller).

Head $2\frac{3}{4}$; depth $1\frac{3}{4}$. D. VI, 11; A. 10; P. 21; branchiostegals 6. Body exceedingly robust, abruptly compressed posteriorly; the length of the head less than its height; crown convex, orbit large, less than $\frac{1}{3}$ of the length of the head. Mouth moderate, reaching a vertical from the foward margin of the orbit. Teeth small, simple, subconical, in a narrow band. On younger examples the anterior dorsal is quite distinct, and does not form a continuous arch with the outline of the back in front of it, as in the Lump, but aged specimens approach the latter more nearly in these respects. Between the 2 dorsals the space is less than base of first. Caudal subtruncate on its posterior margin, the angles rounded off; all the fins with rounded margins. Each of the conical tubercles with which the skin is covered is rough, with small projections on its sides and base; some of the tubercles in diameter of base measuring nearly or quite as much as width of orbit; the largest appearing in a group of 8 or 9 on the middle of the flank, and in several series from the crown to the base of the second dorsal; one of the large ones standing at each side of the space between the dorsals; a couple of moderate-sized ones seen in front of the shoulder, those on the entire caudal region smaller, as also those below the head and body. A young example, about $1\frac{1}{4}$ inches in length, has a similar outline in transverse section as wide as high, is abruptly compressed behind the abdomen, and shows the first dorsal as free and distinct as the second; back somewhat arched under the base of the anterior; interorbital space slightly concave, $\frac{1}{4}$ wider than the orbit, and $\frac{1}{4}$ narrower than the disk; disk nearly $\frac{1}{3}$ wider than the orbit. Entire body covered with irregular-sized spiny tubercles; on the sides of each tubercle the small spines slender and bristle-like, and their development comparatively greater than that obtaining among them on older examples. Later in life it is the central portion or cone of the tubercle that develops, while the lateral outgrowths remain small. On the young individuals the larger scales occupy the spaces behind the pectorals, above the opercles, and at the sides of the first dorsals. Dr. Günther, 1880, figures some young specimens an inch in length, some with tubercles, others without them, and shows the fins to be angular early in life. The color is olivaceous to brownish, very likely reddish in life, clouded with darker. From the material before us, this species appears to be less compressed than either *Cyclopterus lumpus* or *Eumicrotremus orbis*. In a transverse section, in front of the first dorsal, it is nearly round, whereas in either of the others such a section is considerably higher than wide. Compared with them in regard to length, it may be described as elongate. How much of this difference in form is to be credited to individual variation we can not at present determine. Bearing in mind the variations seen in a lot of specimens of *C. lumpus*, it does not seem at all impossible that *E. spinosus* and *E. orbis* may have to be united, as has been suggested by several authors, but unless other specimens show closer approaches than these, they are certainly better kept separate. (Garman.)

Mr. Dresel gives the following notes on a specimen, 90 mm. long, from the stomach of a halibut in Davis Straits:

D. VII, 11; A. ca. 10. The head and body suborbicular; the body posteriorly is abruptly compressed. The mouth is moderate, the jaws with narrow bands of villiform teeth. The maxilla reaches to below the anterior margin of the eye, its length being contained $2\frac{2}{3}$ times in that of the head. The gill opening is small, as long as the diameter of the eye, which is contained 3 times in the length of the head. The disk is about as long as it is broad, $\frac{2}{3}$ of the length of the head. The interorbital width is greater than $\frac{1}{2}$ the length of the head, and not quite twice the diameter of the eye. Length of head contained 3 times, greatest height of body 2 times, in total length. The body is covered with conical plates of various sizes, those of the pectoral region being the largest, about as large as the eye. The plates are studded with small tubercles, and the larger ones have the centers elevated and pointed. Color in spirits, light brown, with traces of punctulations on the skin between the plates.

North Atlantic and Arctic oceans, south to Maine and Denmark; the specimens above described (by Mr. Garman) from Eastport, Maine. (Eu.) (*spinosus*, spinous.)

Cyclopterus spinosus, MÜLLER, Prodr. Zool. Dan., IX, 1777, Denmark; FABRICIUS, Fauna Grœnlandica, 134, 1780; GÜNTHER, Cat., III, 157, 1861; JORDAN & GILBERT, Synopsis, 746, 1883.

Eumicrotremus spinosus, GILL, Proc. Ac. Nat. Sci. Phila. 1873, 190; DRESEL, Proc. U. S. Nat. Mus. 1884, 249; GARMAN, Discoboli, 34, 1892.

Lumpus spinosus, STORER, Synopsis, 482, 1846.

2436. EUMICROTREMUS ORBIS (Günther).

Head $2\frac{1}{2}$; depth $1\frac{1}{2}$. D. VII, 9; A. 10; caudal 10; branchiostegals 6. Body compressed, thick, short, and high; head short, higher than wide, flattened; supraorbital angles prominent. In the specimen described, badly shrunken by drying, the height of the head is contained in the distance from the snout to the base of the caudal nearly twice. Eye large, about $3\frac{1}{2}$ times in the length of the head. Teeth numerous, small, subconical, in pavement, 4 or 5 series. Anterior dorsal much as in a specimen of *C. lumpus* of 3 inches in length, the sides of the fin covered with spines of moderate size, the spinous rays 7 in number. The upper outline of this fin is much like that of the Lump, in being arched backward, or crescentic. The 10 rays of the second dorsal are very distinct; the membranes are not so tuberculate as those of the first. Entire body and head covered with spiny conical tubercles, the largest in a group of 7 or 8 on the flank behind the pectorals, another on the forehead, and those on the supraocular ridge. A single large one stands at each side of the space between the dorsals. Those on the flanks are in contact, and on an example $4\frac{1}{2}$ inches in length there are several that exceed $\frac{1}{2}$ inch in diameter. The sides of the tubercles are roughened by multitudes of fine sharp spines. Under chin and throat the tubercles are smaller, close together, and very numerous. The origin of the first dorsal is above and a little in front of the gill opening. The latter is above the base of the pectoral, behind the eye; its width is hardly as great as that of the orbit. Professor Collett gives a good figure of *C. spinosus;* to compare the specimen from which these notes are taken with it, the Pacific representative is higher and more arched in front of and over the first dorsal, the depth of the body is greater in proportion to the length, and the spines above the lateral line, from the space between the dorsals to the back of the skull, are a great deal smaller. These differences are such as may obtain on

individuals, it is true, but until direct comparisons of series from the Pacific are made with others from the Atlantic it is hardly worth the while to throw *C. orbis* in with *C. spinosus*. The 2 species are very closely allied, but with our specimens it does not seem best to bring them together as one. (Garman.) Northern Pacific, south to Vancouver Island; not rare. The specimen above described from St. George Island, Bering Sea.

The following notes are taken from a very large specimen, 5 inches long, from the Aleutian Islands. (Coll. Alaska Comml. Co.):

Head 3; depth nearly 2; D. VII, 11; A. 10; caudal 10. Body orbicular, cuboid anteriorly; base of spinous dorsal and body behind vent abruptly compressed; teeth in narrow bands; gill openings on a level with eye, slightly narrower than orbit. Spinous dorsal not hidden, covered with rough tubercles, similar to those on body, but smaller; width of ventral disk slightly less than its length, and somewhat less than length of head. Body covered with irregular roundish conical plates, varying much in size, some of the larger with a central point, turned backward; all the plates with small tubercles and slender flexible prickles; small plates along bases of all the fins; a series of 6 large plates extending backward from above eye to opposite the interval between dorsals; a series of 5 on each side of middle of interocular space from snout to nape, these increasing much in size posteriorly; a series of 3 along opercular margin; along base of spinous dorsal, a series of 4, not connected with that of interorbital space; a single large plate on each side of the interval between dorsals; 2 longitudinal series of 4 large plates each, beginning above base of pectorals and behind gill slits, terminating under the space between dorsals; the largest plate on body is immediately behind axil of pectoral; a smaller one below it, and 2 others between it and origin of anal; 2 series of smaller plates below eye; many small plates interposed between the series of large ones; very small plates on under side of head and middle of tail; no plates between ventral disk and vent; none in axil of pectorals. Color olivaceous, the skin between the plates thickly punctate.

"Three young examples, the largest 27 mm. long, were dredged south of Sannak Islands and in Bristol Bay (depth 19 to 70 fathoms). Fin rays in these vary as follows: D. VI or VII, 9 or 10; A. 8 to 10. We agree with Garman in considering it advisable to distinguish the north Pacific form from *E. spinosus*, until adequate series can be compared." (Gilbert.) (*orbis*, orb or sphere.)

Cyclopterus orbis, GÜNTHER, Cat., III, 158, 1861, Vancouver Island.

Eumicrotremus orbis, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 454; GARMAN, Disco. boli, 36, 1892.

Cyclopterus spinosus, JORDAN & GILBERT, Synopsis, 746, 1883; not of MÜLLER.

780. LETHOTREMUS, Gilbert.

Lethotremus, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 449 (*muticus*).

This genus differs from *Eumicrotremus* in the total absence of the bony plates, and of lateral line or pores on sides of head and body. The skin is either smooth or armed with scattered prickles. (λήθη, forgetfulness; τρῆμα, apertures.)

a. Dorsal rays VII, 11; anal 10; skin perfectly smooth. MUTICUS, 2437.

aa. Dorsal rays V, 7; anal 6; skin prickly. VINOLENTUS, 2438.

2437. LETHOTREMUS MUTICUS, Gilbert.

D. VII, 11; A. 10; P. 23. Depth 2 to 2½ in length; head 2⅛ to 2⅓. Eye very large, 2⅛ to 2½ in head, equaling interorbital width. In *Eumicrotremus orbis* of equal size, the length of the slit slightly less than its distance from upper base of pectorals. Diameter of ventral disk $\frac{5}{6}$ length of head, equal to width of mouth. No barbels or filaments. Origin of spinous dorsal slightly in advance of gill slit, distance between dorsals equaling half diameter of eye; origin of anal under that of second dorsal; no notch between upper and lower portions of pectoral fin, the lower rays thickened but not lengthened, the length of upper ray equaling that of snout and eye. Vent separated from disk by slightly more than ½ its distance from front of anal. Color in spirits, brownish above, white below, the upper parts finely freckled with small black specks. Length about 1¼ inches. Aleutian Islands. This species closely resembles in form, fin rays, and general appearance *Eumicrotremus orbis*, but differs, in addition to the generic characters already stated, in the much larger eye, the lower spinous dorsal, and the extreme reduction of the posterior nasal tube. In young examples of *E. orbis*, the posterior nasal tube is much longer than the anterior, and equals ½ the diameter of eye. In *L. muticus*, both tubes are short, and the posterior is little more than an elevated rim to the nasal opening. The naked specimens from the Atlantic, reported on by Dr. Günther under the name *Cycloptremus spinosus*, are probably referable to *Lethotremus*. (Gilbert.) (*muticus*, unarmed.)

Lethotremus muticus, Gilbert, Rept. U. S. Fish Comm. 1893 (1896), 449, pl. 31, Unimak Pass, Alaska, at Albatross Stations 3223 and 3258, in 56 and 70 fathoms.

2438. LETHOTREMUS VINOLENTUS, Jordan & Starks.

Head 2¼ in length; depth 2¼. D. V, 7; A. 6; eye 3 in head; snout nearly 4; maxillary 2½; interorbital 2⅛; ventral disk 1⅛; height of spinous dorsal 2⅛; length of pectoral 2¾. Body short and thick, broadest at head, deepest in front of first dorsal spine, abruptly compressed at vent; back somewhat elevated. Mouth terminal, oblique, the jaws about equal; snout very blunt; maxillary reaching slightly past the vertical from front of eye; teeth in narrow villiform bands; teeth on vomer (the specimen is so small, we can not be sure of the palatine teeth); eye large, set high in the head, its diameter greater than the length of the snout; interorbital wide and flat, the diameter contained 1½ times in the width; gill opening oblique, about as wide as eye and on a level with eye; disk ¼ longer than broad, its length about equal to distance from gill opening to anterior edge of eye. Skin thick; head and body nearly naked, a few spines scattered over it; spinous dorsal with 3 or 4 small spines, a minute simple spine in front of eye and 2 or 3 above it; 4 multifid spines following the curve of back, under spinous dorsal, and 1 under the front of second dorsal, 2 similar spines on each side of nape, just above opercles; 2 on edge of opercle and 3 on edge of preopercle; an irregular row of 6 running from above base of pectoral to front of anal fin, and a couple of small ones behind gill

opening; body otherwise naked. All the spines, with the exception of those noted as simple, are long sharp spines in groups of from 3 to 6 with a common base, generally the length of each spine exceeding length of the base. No lateral line. Spinous dorsal reaching to the first ray of soft dorsal when fin is depressed, higher than soft dorsal; anal and soft dorsal similar; caudal small, truncate or slightly rounded; pectorals very short, reaching to the posterior edge of ventral disk. Color bright wine red, slightly lighter below, without markings, sides dusted over with very small dark points; spinous dorsal dusky; other fins colorless. Colors disappear in alcohol. One specimen ½ inch long, dredged in Puget Sound. If the adult shows the same characters, it must become the type of a distinct genus. (*vinolentus*, wine-colored.)

Lethotremus vinolentus, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 827, pl. 94, Puget Sound, near Seattle, Washington. (Type, No. 3131, L. S. Jr. Univ. Mus. Coll. E. C. Starks.)

781. CYCLOPTEROIDES, Garman.

Cyclopteroides, GARMAN, Mon. Discoboli, 37, 1892 (*gyrinops*).

Body short, thick, slightly depressed anteriorly, compressed posteriorly. Head broad, short; snout short, obtuse; subordital produced to connect with the preoperculum, widening backward. The bases of both dorsals are thickly enveloped in skin and flesh. Ventrals united, forming a large adhesive disk. Chin with tubular pores, or barbels. Teeth small, subconical. Gill openings narrow; gill membranes united and attached to the isthmus. Gills 3½. Pseudobranchiæ present. Six branchiostegal rays. Intestine long. Like *Cyclopterus*, to which it is closely allied, the species of this genus appears to derive a portion of their food from vegetation. (*Cyclopterus*; *εἶδος*, resemblance.)

2439. CYCLOPTEROIDES GYRINOPS, Garman.

Head 3 in total length; D. 8 or 9; A. 9; pectoral 24; caudal 10; cæca 10 or 11; branchiostegal 6. Body oblong, compressed toward the dorsals, broad toward the belly, subtriangular in transverse section, abruptly compressed in the posterior portion between the second dorsal and the anal, deep near the abdomen, and tapering rapidly to the caudal fin. Belly flattened, ⅔ as wide as long. Head short, as long as high, wider than long; interorbital space concave transversely and slightly so longitudinally, steeply inclined on snout and sides; cheeks swollen; mouth anterior, somewhat oblique, little wider than interorbital space, not reaching a vertical from front of eye; lower lip interrupted for a short distance at the symphysis; chin with a series of 4 barbels on each side formed by tubular prolongations of the pores similar to those of the nostrils. Teeth very small, conical, slightly hooking backward, in 5 rows at the symphysis, some of which are shorter and do not extend so far toward the sides as the others. Nostrils small, tubular, the posterior between the eyes, and the anterior halfway between the posterior and the mouth. Eye moderate, lateral, as long as the snout, more than 4 in

head, prominent above the outline of the forehead. Gill opening very small, covered by the spine-like angle of the opercle, at a distance above the base of the pectoral equal to about twice its own width. Skin covered with mucus, with 4 series of very small, distant, 1- to 8-spined tubercles on each side; 1 series on each side of the median line of the forehead along the sides of the bases of the dorsals, another from each orbital ridge backward on the flank parallel with the first, another behind each eye, and the fourth at the lower edge of the gill opening; below the last series the skin is quite bare. Ventral disk as long as the head, as broad as long, or broader, nearly twice as far from the caudal as from the mouth. Dorsal fins very thick at their bases, enveloped in loose skin through which the rays may be distinguished, separated by an interspace, the first higher, originating above the gill opening; base of the second higher in front, and the posterior rays more free from the thick skin. (The extremities of the rays are all weak, and in most cases have been carried away. That the caudal fin was pointed is indicated by the strength of the median rays; in other words, the fin was longer in the middle.) Pectorals broad, rounded in the upper part posteriorly, reaching as far backward as the hinder edge of the disk; lower border but little indented, if at all, with ends of rays prolonged as a short fringe. Color brown, approaching a chocolate; belly light; a dark brown blotch between the eyes, another from each eye through anterior nostril to mouth, 1 below eye to throat, 1 or 2 back of eye on opercle, a larger one behind gill opening, several small ones close along bases of dorsals, and an elongate band on the hinder part of flank and on base of caudal portion; peculiar marks exist in a brown blotch, on each side of hinder part of abdomen, on which is a bunch of 3 white spots like a clover leaf, in a white band, reaching toward the flank at each side of hinder margin of disk, and in a white spot at posterior ends of branchiostegal rays. Total length a little less than 2 inches. St. Paul Island, Alaska (Garman); the young taken by us in the harbor of Unalaska. (*γυρινός*, tadpole; *ὤψ*, appearance.)

Cyclopteroides gyrinops, Garman, Mon. Discoboli, in Memoirs Mus. Comp. Zool., Cambridge, 37, 1892, St. Paul Island, Alaska.

782. CYCLOPTERICHTHYS, Steindachner.

Cyclopterichthys, Steindachner, Ichth. Beiträge, x, 14, 1881 (*glaber* = *ventricosus*).

Body short and thick, rounded, covered with thick, smooth skin, destitute of bony tubercles; tail slender, compressed, the body abruptly contracted to its base; head broad, obtuse; mouth oblique, the lower jaw prominent; teeth rather small, simple, hooked, sharp, in 2 rows anteriorly; pseudobranchiæ large; gills $3\frac{1}{2}$; suborbital connected by a bony stay with the preopercle; gill opening small, above the base of the pectoral, which is broad and procurrent; ventral disk moderate, fringed. Dorsal short and high, of soft rays only, opposite the short anal, both well separated from the small caudal. (*Cyclopterus*; *ἰχθύς*, fish.)

2440. CYCLOPTERICHTHYS VENTRICOSUS (Pallas).

Head $2\frac{2}{4}$; depth $3\frac{1}{2}$. D. 9; A. 7 (8 or 9 Pallas); P. 20; C. 11. Body stout, thick, short, smooth, abruptly compressed behind the body cavity. Caudal portion nearly $\frac{2}{3}$ of the total length. Head about $\frac{1}{3}$ of the length without the caudal, broad, depressed but convex on the crown, blunt and broadly rounded on the snout. Mouth wide, anterior, opening slightly upward, its corner before the middle of the eye. Teeth small, simple, in 2 series in front, and but 1 toward the sides. Pallas says of the dentition: "Maxillarum margines ossei, denticulis minutis, curvulis, obtusiusculis, ionordinatis et inæqualibus, passim geminatis asperi; quorum in superiore maxilla interiores majores." Eye lateral, its diameter contained 7 times in the length of the head. Gill slit 4 in head. Dorsal short, behind the abdominal chamber, its upper extremity extending behind a vertical from the base of the caudal, margin rounded; anal smaller than the dorsal and opposed to it, though originating a trifle farther back; caudal subtruncate, rounded. Vent almost directly in the middle of the total length. Disk subcircular, with a broad cutaneous margin, $1\frac{1}{2}$ in head. Brownish, with numerous scattered, small, rounded spots of darker; our specimens plain blackish olive. Bering Sea; Sea of Okhotsk (Steindachner); our specimens from St. Paul Island, Pribilof Group; also found at Petropaulski; occasionally taken in abundance. (*ventricosus*, large-bellied.)

Cyclopterus ventricosus, Pallas, Spicilegia, Zool., vii, 15, t. 2, 1769, Kamchatka.
Cyclopterichthys glaber, Steindachner, Ichth. Beiträge, x, 14, pl. 8, 1881, Sea of Okhotsk.
Cotylis ventricosus, Günther, Cat., iii, 498, 1861.
Cyclopterichthys ventricosus, Jordan & Gilbert, Synopsis, 745, 1883; Garman, Discoboli, 41, 1892.

783. LIPAROPS, Garman.

Liparops, Garman, Discoboli, in Mem. Mus. Comp. Zool., 42, 1892 (*stelleri*).

A little-known genus, apparently closely allied to *Cyclopterichthys*, but distinguished by the elongate dorsal, the tubercles, and the dentition. One species known. (*Liparis*; ὤψ, appearance.)

2441. LIPAROPS STELLERI (Pallas).

Body oval, smooth, the chest in the middle, the depth 4 inches; a single series of osseous tubercles along the middle of the back to the origin of the dorsal fin; lower jaw longest; teeth slender, blunt, unequal; eyes small, high; gill opening reduced to a round foramen; opercle large. Disk between pectoral fins. Dorsal fin commencing on the middle of the back and extending to near the base of caudal; anal ending opposite dorsal; anal slender. Petropaulski, Kamchatka, where it is rare (Pallas); not seen by any recent collector. (Named for Georg Wilhelm Steller, naturalist and explorer.)

Cyclopterus stelleri, Pallas, Zoogr. Rosso-Asiat., iii, 73, 1811, Kamchatka, Harbor of Peter-and-Paul; described from manuscripts of Steller.
Cotylis stelleri, Günther, Cat., iii, 499, 1861.
Cyclopterichthys stelleri, Jordan & Gilbert, Synopsis, 745, 1883.
Liparops stelleri, Garman, Discoboli, 42, 1892.

Family CLXXXIII. LIPARIDIDÆ.

(The Sea Snails.)

Body more or less elongate, tadpole-shaped, subcylindrical anteriorly, compressed behind, the head depressed; both head and body covered with smooth, thin skin, which is very lax. Head broad, obtuse, the snout short, wide, and blunt; third suborbital bone styliform behind, forming a bony stay articulating with the preopercle, as in *Cottidæ*; mouth moderate, anterior, terminal, the jaws equal, or the lower included; jaws with bands of small teeth, which are simple or more or less tricuspid, usually close set, forming a pavement; no teeth on vomer or palatines; premaxillaries protractile, little movable; opercular bones unarmed; interopercle slender, ray-like, overlying the branchiostegals; gill openings small, the membranes joined to the broad isthmus and to the humeral arch below. Branchiostegals 6. Gills 3½, no slit behind the last; pseudobranchiæ small or wanting; dorsal fin rather long, the spines feeble and flexible, low, similar to the soft rays; anal long, similar to the soft dorsal; ventral fins I, 5, the two completely united and forming the bony center of an oval sucking disk, or else sometimes entirely wanting; pectoral fin very broad, the base procurrent, extending forward under the throat, the outline usually emarginate, some of the lower rays being produced; tail diphycercal; caudal fin short, convex; vertebræ numerous, 35 to 50; pyloric cæca numerous; no air bladder; stomach siphonal, U-shaped, intestine elongate. Genera 9; species about 40. Small, sluggish fishes, nearly all of the Arctic seas, a few belonging to the Antarctic; found adhering to rocks at various depths. The group is evidently closely allied to the *Cottidæ*, and its origin must be sought in the ancestors of such types as *Psychrolutes* and *Cottunculus*, the *Cyclopteridæ* representing a coordinate phase of degradation.

The following additional characters are given by Mr. Garman:

Skeleton with a somewhat larger proportion of osseous matter than that of the *Cyclopteridæ*. As in that family, the skull is full on the back, as if truncate, and has no passage for muscles extending forward between the mastoid and the occipitals. Third suborbital very long, slender, and spine-like, posteriorly reaching toward the posterior margin of the preoperculum. Upper limb of preoperculum expanded. Operculum much reduced, its shape with some resemblance to that of a boot, the sole directed downward and the toe backward. Suboperculum slender, like an inverted V with the limbs curved outward, the anterior extending forward along the preoperculum, and the posterior reaching back under the lower border of the operculum. Interoperculum long, styliform, reaching from suboperculum to articular, frequently mistaken for one of the branchiostegal rays. Reasons for separating this family from the *Cyclopteridæ* and placing it farther from the *Cottidæ* exist in the continuous dorsal, the connection of dorsal and anal with the caudal, the more complete transformation of the ventral rays, the more slender and spine-like suborbital process, the expanded upper limb of the preoperculum, the styliform interoperculum, the shorter olfactory nerves, and the more elongate brain. (Garman.)

Discoboli liparidina, Günther, Cat., III, 158 to 165, 1861.

Liparidinæ:

a. Ventral disk present, large or small.

b. Ventral disk normal, composed of 13 lobes, a median one in front, and one corresponding to each of the fin rays, each lobe with a deciduous horny covering or papilla. Teeth close set in pavement-like bands, tricuspid in the

young, some or all becoming angular or bluntly arrow-shaped in the adult; caudal fin well developed, rather broad.

c. Dorsal fin divided by a deep notch, which separates the slender spines from the soft rays. NEOLIPARIS, 784.

cc. Dorsal fin continuous, the spines not separated from the soft rays. LIPARIS, 785.

bb. Ventral disk more or less reduced in size or degraded in structure, becoming anterior in position; teeth simple or tricuspid only in the very young; vertebræ numerous.

d. Caudal well developed; teeth slender, curved, in cardiform bands, the inner longer, depressible; ventral disk large; skeleton very soft. BATHYPHASMA, 786.

dd. Caudal fin slender, not distinct from other vertical fins; teeth simple in the adult; vertebræ numerous; dorsal rays 35 to 50; anal 33 to 46. Anterior nostrils usually without tubes; ventral disk very small, nearly under the eye; the vent close behind it.

e. Pectoral fins with the lower lobe well developed; gill openings moderate or small, not greatly reduced. CAREPROCTUS, 787.

ee. Pectoral fins not notched, the lower lobe obsolete, the rays progressively shortened; gill openings reduced to a small pore. GYRINICHTHYS, 788.

AMITRINÆ:

aa. Ventral disk entirely wanting; no ventral fins; teeth mostly simple in the adult; vertebræ and fin rays in large number.

f. Pseudobranchiæ present; lower lobe of pectoral produced. AMITRA, 789.

ff. Pseudobranchiæ wanting.

g. Tip of snout without barbel. PARALIPARIS, 790.

gg. Tip of snout with a pair of barbels; snout projecting. RHINOLIPARIS, 791.

784. NEOLIPARIS, Steindachner.

Neoliparis, STEINDACHNER, Ichth. Beitr., III, 54, 1875 (*mucosus*).

This genus differs from *Liparis* in having a deep notch in the dorsal fin anteriorly, separating the spines from the soft rays. The species approach more nearly to the Cottoid type, from which the Liparids are descended. In general the vertebræ are fewer, the fin rays fewer, the ventral disk larger, and the vertical fins better separated than in the more degenerate members of the family. The retention of the notch between the dorsals fully justifies the recognition of *Neoliparis* as a distinct genus. (*νέος*, new; *Liparis*.)

a. Gill opening very narrow, almost entirely above base of pectoral, the lower edge not below third pectoral ray.

b. Anterior nostrils with distinct tubes.

c. Dorsal spines in the male greatly elevated, not connected by membranes except at extreme base.

d. Dorsal rays about 25; anal 23; head small; ventral disk small. ATLANTICUS, 2442.

dd. Dorsal rays 26; anal 23; head larger; ventral disk comparatively large. RUTTERI, 2443.

cc. Dorsal spines not especially elevated; dorsal rays 34 to 36; anal rays 25 to 28; lower jaw included; form rather elongate, the head depressed; ventral disk $2\frac{1}{3}$ in head. Color pale, irregularly dotted with darker, sometimes plain brownish. CALLYODON, 2444.

bb. Anterior nostrils with a raised rim, and without distinct tubes; head short, blunt, 4 in length; ventral disk very large, 1⅓ in head; snout blunt; mouth very short, its cleft almost entirely anterior, the maxillary scarcely reaching eye; dorsal rays 32; anal 26. Color plain rosy or brownish, not spotted. MUCOSUS, 2445.

aa. Gill opening rather large, its base opposite 4 or 5 upper rays of pectoral; body deep posteriorly; nostrils with raised rim, but without distinct tubes; ventral disk moderate, 2¼ to 2½ in head; head about 3¾ in body, depressed above; cleft of mouth broader, partly lateral, nearly 3 in head. Color plain brownish or reddish.

e. Dorsal VI, 27; anal 21 to 23; pectoral 30; flesh firm. FLORÆ, 2446.

ee. Dorsal VI, 34; anal 30; pectoral 35; flesh lax. GREENI, 2447.

eee. Dorsal VI, 28; anal 26; pectoral 36. FISSURATUS, 2448.

2442. NEOLIPARIS ATLANTICUS, Jordan & Evermann, new species.

Head 4⅜; depth 4 to 4½. D. VI, 25; A. 23; P. 30. Eye 5 in head; snout 3; ventral disk 1¾ to 2. Body widest at gill opening, compressed posteriorly; deepest below third dorsal spine; head broader than deep, depressed above the eyes; mouth narrow, its cleft transverse and extending to anterior nostril; lower jaw included; teeth tricuspid, the middle cusps highest; gill openings very narrow, the lower border opposite first ray of pectoral; anterior nostril tubular, the tube ⅔ of eye; posterior nostril with a low flap; skin loose, lying in folds. Origin of dorsal not far behind pectoral, its distance from tip of snout ⅓ its distance to base of caudal. Dorsal with a very shallow notch, the spines nearly continuous with the soft rays; in the males much elevated, the tips thickened and membranes deeply incised; the first or longest spine as long as head; the sixth or last not quite ½ head; middle rays of soft dorsal 2 in head; dorsal and anal joined to base of caudal; caudal nearly as long as head; pectoral almost reaching anal, slightly longer than head; lower rays exserted, forming a slight lobe. Color reddish brown, with small scattered light or bluish dots over the body; fins darker, clouded with pale, the dorsal broadly edged with darker. The specimen here described, 37215 U. S. Nat. Mus., is about 5 inches long, from Godbout, Quebec. Numerous other specimens from Salem (20367) and Woods Hole, Massachusetts (40118), agree with this, the females having the first dorsal much lower. It is generally common along rocky shores from Newfoundland to Cape Cod. This species has been repeatedly recorded under the name *Liparis montagui,* Donovan, but it is not identical with the European species of that name. The published figures of *N. montagui* show a deeper fish with larger head, and with the spinous dorsal very low, and scarcely distinct from the soft rays. Day's figure looks remarkably unlike our fish. Unfortunately the European specimens of *Neoliparis montagui* in the National Museum are all too small for satisfactory comparison.*

Neoliparis atlanticus, JORDAN & EVERMANN, new species, Godbout, Quebec. (Type, No. 37215.)

Liparis montagui, CUVIER, Règne Anim. Ed. 1, vol. 2, 227, 1817; JORDAN & GILBERT, Synopsis, 743, 1883, in part; GARMAN, Discoboli, 47, 1892, with plate, and of authors generally.

* Dr. Boulenger has kindly sent us the following note on *Liparis montagui:* I have examined about a dozen of our British *Liparis montagui,* of all sizes up to 5 inches. The anterior dorsal looks very indistinct, has no detached portion, and none of its rays are ever produced into filaments.

Cyclopterus montacuti, TURTON, British Fauna, 115, 1807; after DONOVAN.

Liparis gobius, CUVIER, Règne Anim., I, 227, 1817; after *Gobius*, MÜLLER, Zool. Dan., IV, 16, 1806.

Cyclopterus liparoides, NILSSON, Prodr. Ichth. Scand., 62.

Lepadogaster cornubiensis, THOMPSON, Proc. Zool. Soc. Lond. 1835, 81, Cornwall.

Liparis reticulata, COUCH, Brit. Fish., II, 195, 1863.

Liparis ekströmi, LÜTKEN, Vid. Medd. Nat. For. Kjöbenh. 1865, 221.

Liparis maculatus, MALM, Förh. Skand. Naturf. 1865, 412.

Liparis montagui, with varieties *principalis*, *maculatus*, *annulatus*, *striatus*, *pictus*, *obscurus*, *liparoides* and *ekströmi*, MALM, Bohusläns Fauna, 103, 1877.

Prof. Robert Collett (Christiania Vidensk. Selsk. Förh., 1879, 38) has assigned names to the many color variations of *Neoliparis montagui* as shown in Norway, as follows:

Var. a. *principalis*, Collett, uniform colored; light red yellow with microscopic brownish dots. The common form from Christiania, Stavanger, Flor., onear Bergen, Komagsbjord in Finmarken. *Liparis montagui*, MALM, Götheborg og Bohusläns Fauna, 451, 1877.

Var. b. *maculatus*, Malm, dark grayish yellow, the whole body strewn with many rows of small brown spots, smaller than the eye and sharply defined. Bukn and Bohuslän. *Liparis maculatus*, MALM, Förh. Skand. Naturf., 9, 1863, 412.

Var. c. *annulatus*, Collett, reddish gray yellow, with large, irregular red-brown rings of the same length and breadth; on sides of head the rings becoming stripes; Christiansand.

Var. d. *striatus*, Collett, color of *c*, the rings drawn out into lines; the rings only seen in front of first dorsal; about 6 longitudinal red-brown stripes from gill opening to caudal. Bergen. *Liparis lineatus*, MALM, Förh. Sk. Förh., 9, 1863, 412.

Var. e. *pictus*, Collett, as in *striatus*, but the lines very numerous, close set, irregularly vermiculate, with much cross netting or partial rings. Bergen. Finmarken.

Var. f. *obscurus*, Collett, gray brown with many dark brown spots and specks on body and fins; irregular cross bands on caudal; back almost uniform gray brown. Hardanger Fjord.

Var. g. *liparoides*, Nilsson, uniform dark gray brown, the color given by innumerable dark points; trace of band at tip of caudal; paler below. Hardanger, Oefjord in Finmarken Tromsø. *Cyclopterus liparoides*, NILSSON, Prodr. Ichth. Scand., 62, 1832.

Var. h. *ekströmi*, Malm, dark brown, sides strewn with grayish yellow small spots which extend on vertical fins. Bohuslän. *Liparis ekströmi*, MALM, Förk. Sk., Naturf., 9, 1863, 412.

These are evidently mere color forms, expressing the range of variation. Similar color forms may be seen in most of the bright-colored species, *pulchellus*, *agassizii*, *aleuticus*. They are not in any proper sense subspecies, having no geographical bases or separate lines of descent.

2443. NEOLIPARIS RUTTERI, Gilbert & Snyder, new species.

Head $3\frac{2}{3}$; depth $4\frac{1}{3}$. D. VI, 26; A. 23; P. 30; eye 5 in head; snout $2\frac{2}{3}$; ventral disk $1\frac{2}{3}$. Body widest at gill opening, gradually tapering posteriorly, deepest below third dorsal spine; depth of caudal peduncle 4 in head. Head broader than deep, its depth at occiput 6 in body. Mouth narrow, its cleft almost wholly transverse, extending to vertical from anterior nostril; lower jaw included; teeth tricuspid, the middle cusp highest, arranged in 10 oblique series on each half of the upper and the lower jaws. Gill opening very narrow, restricted to the area above base of pectoral. Anterior nostrils tubular; posterior nostrils with a low rim, the anterior part of which is elevated into a triangular flap. Skin loose, lying in folds and wrinkles. Origin of dorsal at $\frac{1}{3}$ the distance from tip of snout to caudal; rays of first dorsal (in the types which are apparently all males) separate almost to base, each ray bordered with a thick mem-

brane, the margin thus formed sometimes incised, producing short, rounded lobes; first spine highest, its height contained 4⅔ in body, the succeeding spines gradually growing shorter, the sixth contained 11½ times in body. Height of second dorsal near its middle, 3 in head; dorsal and anal joined by membranes to base of caudal, the anal more broadly joined than the dorsal, attached to basal fifth of lowest caudal ray; caudal rounded posteriorly, its length 4⅓ in body; origin of anal below first ray of second dorsal; height of anal a little less than that of dorsal; pectoral extending to a vertical crossing body halfway between anus and origin of anal fin; length of pectoral 5½ in body, its upper part rounded posteriorly; the fifth, sixth, and seventh rays from below elongated, with exserted tips, forming a short lobe. Long diameter of ventral disk 6 in body. Vent slightly nearer to margin of disk than to origin of anal fin. Color bluish slate, lighter along middle of sides; belly and gill membranes whitish; pectorals, anal, and posterior half of dorsal very narrowly edged with white; rays of first dorsal indistinctly barred. In the following table the measurements of 3 specimens are expressed in hundredths of the length of the body:

	No. 1.	No. 2.	No. 3.
Length, in millimeters, to base of caudal fin	37	44	50
Length of head in body	28	26	27
Depth of body	20	20	23
Distance from snout to dorsal	33	31	34
Distance from snout to anal	50	47	48
Depth of caudal peduncle	7	7	8
Length of snout	10	11	11
Diameter of eye	5	5	5
Width of mouth	11	12	12
Width of gill opening	4½	5	5
Height of first dorsal ray	18	23	22
Height of dorsal fin near middle	8	7½	8
Height of anal	6	7	7
Length of pectoral	18	20	21
Longitudinal diameter of disk	15	16	16
Length of caudal	18	19	20
Number of dorsal rays	VI, 26	VI, 26	VI, 26
Number of anal rays	24	24	23
Number of pectoral rays	.....	30	30
Number of caudal rays	.....	12	13

Neoliparis rutteri is closely related to the Atlantic species, *Neoliparis atlanticus*. It differs from it in the longer head, and the comparatively larger ventral disk. The 2 species differ from all other known species of the genus in having the dorsal spines in the male greatly elevated and not connected by membrane except at extreme base. From *N. floræ* and *N. greeni*, *N. rutteri* differs also in having a much smaller gill opening and

fewer dorsal rays; and from *N. mucosus* in having the pectoral more deeply notched and the anterior nostrils with a tube. Kadiak Island, Alaska. Known from numerous specimens, the longest 61 mm. in length. (Gilbert & Snyder.) (Named for Cloudsley Rutter, now of the U.S. Fish Commission who collected in Kadiak in 1896.)

Neoliparis rutteri, GILBERT & SNYDER, MS., Fishes of Kadiak, Ugah Bay, Kadiak Island. (Coll. Cloudsley Rutter. Type, No. 5701, L. S. Jr. Univ. Mus.)

2444. NEOLIPARIS CALLYODON (Pallas).

Head $3\frac{4}{5}$; depth $4\frac{1}{3}$. D. VII, 33; A. 26; pectoral 31; caudal 14 to 16; eye $6\frac{1}{2}$; snout $2\frac{1}{2}$; pectoral $1\frac{2}{3}$; ventral disk $2\frac{1}{6}$; caudal 2. Body moderately elongate, compressed posteriorly; depth of head $1\frac{2}{3}$ in its length, breadth $1\frac{1}{3}$; mouth rather large, the maxillary buried under the skin, reaching slightly past anterior edge of eye; teeth small, tricuspid, in about 8 oblique rows; snout blunt and rounded; interorbital space wide and slightly convex; posterior nostril ending in a short wide tube, the anterior much smaller; behind the posterior nostril, over anterior edge of eye, is a mucous pore; length of gill opening twice the diameter of the eye, its lower third in front of pectoral, extending down to the second or third ray; vent about midway between posterior edge of ventral disk front of anal; pectorals short and broad, with the usual notch below, reaching slightly past vent; ventral disk a little longer than broad, its distance from tip of lower jaw equaling its diameter; origin of dorsal directly over vent, a shallow notch separating its spines from its rays; front of anal much nearer snout than base of caudal, directly under dorsal notch; dorsal and anal scarcely joined to caudal; caudal long and slender, rounded behind. Among our specimens are 2 types of coloration; the first is uniform light olive brown on back, white below; fins all colorless; the other is light brown on back and sides, everywhere spotted with blackish spots irregular in size; lower parts light; pectoral with indistinct dark cross markings; dorsal and anal mottled and spotted with blackish, darker posteriorly; caudal very dark, black posteriorly; lips dusky. Coasts of Alaska and Bering Sea, west to Kamchatka; generally common on rocky shores; recorded from St. Paul, Kamchatka, Plover Bay, Siberia, Unalaska, and Kadiak; our specimens from the 2 islands last named; the description from Unalaska specimens collected by the *Albatross*, the longest 5 inches long. This species is evidently the original *Cyclopterus callyodon* of Pallas, as is shown by the coloration, the small mouth, and the reduced size of the gill openings. It differs from *Neoliparis mucosus* in the much smaller ventral disk, the presence of nasal papillæ, as well as in proportions and coloration. (*καλλος*, beauty; *ὀδούς*, tooth, from the tricuspid teeth which Pallas had never before seen in a fish.)

Cyclopterus callyodon, PALLAS,* Zoogr. Rosso-Asiat., III, 75, 1811, Kamchatka and Aleutian Islands.

* The following is the substance of the account given by Pallas: Dorsal 33; anal 24; caudal 15; branchiostegals 6; dorsal and anal fins continued nearly to the base of the caudal. Body oblong, compressed; the head depressed; the snout much depressed, rounded; lower jaw included. Teeth with 3 distinct lobes; a slender bone from the eye to the preoperculum: gill openings reduced to lunate spiracles, on each side; ventral disk small

Liparis callyodon, GÜNTHER, Cat., III, 162, 1861; BEAN, Proc. U. S. Nat. Mus. 1881, 247, 271; JORDAN & GILBERT,* Synopsis, 743, 1883.

2445. NEOLIPARIS MUCOSUS (Ayres).

Head 4 in length; depth 4⅕. D. VI, 26; A. 26; pectoral 29; caudal 12; eye 7 in head; snout 3; ventral disk 1½; pectoral 1⅔; longest dorsal ray 2; highest anal ray 2; caudal 1¼. Body not greatly elongate, rather robust, compressed posteriorly, holding its width well past middle of body; head short and thick, broader than body, ⅛ longer than broad, its length 1⅔ times its depth; mouth small, truncate, its cleft almost entirely anterior, scarcely extending laterally; end of maxillary buried under the skin, barely reaching to eye; nostrils not ending in tubes; lower jaw slightly the shorter; teeth sharp, tricuspid, the middle cusp much the highest and longest, arranged in 9 oblique series in each jaw, becoming more and more oblique toward the sides; interorbital space moderately wide, about 3½ in head, a little convex; gill slit not extending below upper edge of pectoral, its length about 1⅓ times eye and 3 in ventral disk. Pectoral broadly rounded when spread, its notch comparatively very shallow, its tip reaching past vent but not to notch in dorsal; ventral disk large, slightly longer than broad, its posterior margin almost midway between its anterior and front of anal, its anterior margin ⅓ its length from chin; dorsal with a shallow notch; origin of fin over posterior margin of ventral disk, its longest rays in its posterior half; origin of anal a little nearer snout than base of caudal, the last 4 or 5 rays rapidly shortened, making the fin truncate behind; dorsal and anal scarcely joined to caudal; caudal long and slender, rounded behind. Color olive brown, light below; indistinctly mottled; dorsal and anal darker at their margins; pectorals uniform dark brown; caudal light, with indistinct cross lines; lips dark. Coast of California. Here described from the only specimen known to us, 5 inches in length, from near San Francisco. It is now in the collection of the California Academy of Sciences (No. 360). Collected by H. D. Dunn, off San Francisco. (*mucosus*, slimy.)

Liparis mucosus,† AYRES, Proc. Cal. Ac. Sci., I, 1855, 24, San Francisco (Coll. W. O. Ayres); GÜNTHER, Cat., III, 559, 1861; STEINDACHNER, Ichth. Beitr., III, 54, 1875; JORDAN & GILBERT, Synopsis, 744, 1883; description copied from STEINDACHNER.

Liparis mucosus, GARMAN, Discoboli, 52, 1892, with plates, figures and description in part; account of the gill openings apparently taken from *Liparis florœ*.

Neoliparis mucosus, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 832, pl. XCV.

2446. NEOLIPARIS FLORÆ, Jordan & Starks.

Head 3⅔ in length of body; depth at ventral disk 5½; depth under middle of soft dorsal 4½. D. VI, 27; A. 21 to 23; caudal 15; pectoral 30;

between the branchiostegals; dorsal commencing a little behind the gill opening. Yellowish brown above; skin everywhere with brown dots; a silvery streak from the upper jaw through the eye and opercles. Kamchatka; common along the shores and about the (Aleutian) Islands. (Pallas.)

* Described from a small specimen with injured fins lately reexamined by us, in the collection of the Alaska Commercial Company, Unalaska. (Coll. A. Greenebaum.)

† The description of Ayres is not very full and might apply almost equally well to *Liparis florœ*. The account of the head (4 in length) and the eye (nearly 6 in head) point rather to the present species as does also the description of the lax skin. Steindachner's account seems also to refer to the present species, although few of the really distinctive characters are mentioned by him.

eye 7 in head; interorbital space 2¾; maxillary 2½; pectoral 1⅔; ventral disk 2¼. Body moderately elongate, much compressed posteriorly, about as wide as deep anteriorly, its greatest depth under middle of soft dorsal where the back is elevated. Flesh very firm, the body retaining its form, the skin loose but not flaccid. Head small, the nape not produced; mouth moderate, the maxillary extending to below the anterior margin of orbit; jaws subequal; teeth tricuspid, arranged in series which are nearly transverse on middle of jaws, becoming more and more oblique toward the sides, the outermost series nearly parallel with the sides of jaws; nostrils ending in a short, wide tube; gill opening short, extending downward to about the fifth pectoral ray, its length about ½ interorbital space; opercle ending in a flap, which extends over middle of gill opening; ventral disk slightly longer than wide, its distance from tip of lower jaw 1⅓ times its length; vent equidistant from posterior edge of ventral disk and front of anal; skin thick and not very loose. Origin of spinous dorsal a little in front of the vertical from vent, its distance from snout 3 in length of body; anterior part of dorsal separated by a notch; origin of anal about equidistant from snout and base of caudal fin; some of the lower rays of pectoral produced forming a notch in the lower posterior margin of fin, the fourth to the tenth of the upper rays the longest, forming a rounded point behind, extending slightly past the vertical from snout; dorsal and anal scarcely connected with the caudal; caudal long and slender, rounded behind. Color, a uniform dark olive green, under parts white, a light streak medially along back from dorsal to occiput, a light spot over opercle; pectoral light at base, dusky behind; other fins colored like the body; lips white. The specimen described above, 4 inches long, taken in rock pool on Waadda Island, Neah Bay. A second specimen, about 5 inches long, is in the museum of the California Academy of Sciences. It was collected off San Francisco by Mr. H. D. Dunn. In this specimen, the dorsal rays are VI, 27; anal 25; pectoral 30; caudal 15; teeth blunt. Flesh firm; form and color of the young example above described. San Francisco to Puget Sound. (Named for Mrs. Flora Hartley Greene, then assistant curator of the museum of Leland Stanford Jr. University.)

Liparis cyclopus, JORDAN & GILBERT, Synopsis, 743, 1883; not GÜNTHER.
Neoliparis floræ, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 830, pl. 96, Waadda Island, Neal. Bay. (Type, No. 3133, L. S. Jr. Univ. Mus. Coll. E. C. Starks.)

2447. NEOLIPARIS GREENI, Jordan & Starks.

Head 3¾; depth 4; depth at disk 5. D. VI, 34; A. 30; pectoral 35; caudal 15; eye small, about 10 in head; snout 2¾; longest pectoral ray 1¾; disk 2½; longest dorsal ray 2⅙; longest anal ray 2⅙; caudal 1¾. Body elongate, posteriorly compressed; profile undulate, over snout blunt and rounded, depressed over eyes, well rounded from eyes over occipital region. Skin thin and exceedingly loose nearly to the end of the dorsal and anal rays. Jaws equal; maxillary extending to posterior margin of eye; teeth small, nearly simple, depressible and blunt, slightly hooked back, arranged in oblique series, those in the front running nearly straight in, but toward

the sides of the jaw they grow more and more oblique till they are nearly parallel with the jaw at the sides; superior pharyngeal teeth conical and sharp, slightly longer than the teeth in the jaws, arranged in a single round patch on each side; inferior pharyngeals separate, with small teeth. (Teeth probably tricuspid in the young.) Posterior nostrils in a short, wide tube; cheeks well rounded; gill rakers short and thick, no longer on the outer side of the first arch than on the other arches, fourth arch not free; gill slit short, its length contained about 3 times in head, its lower edge extending in front of pectoral to about the third ray; opercles with a blunt spine which is covered by the skin. Dorsals 2, connected by a low membrane; first dorsal about twice as high as anterior part of second dorsal; the first rays of pectorals inserted under eye and in front of disk; the anterior rays short, graduated to the sixth ray, which is about 4 times longer than the first, the next few rays again short and gradually lengthening posteriorly; posterior rays $\frac{1}{3}$ longer than anterior, fin broadly rounded behind; ventral disk nearly round, its posterior edge reaching the vertical from gill slit, its distance from tip of lower jaw $1\frac{1}{3}$ times its length; caudal truncate or slightly rounded; vent under ends of pectorals. Color, in alcohol, uniform brown, breast and lower parts of head creamy, fins slightly darker. When fresh the sides were blotched with pinkish. Esquimault Harbor, near Victoria, Vancouver Island; known only from the type, 10 inches long. (Named for Mr. Ashdown H. Green, of Victoria, president of the Natural History Society of that town.)

Neoliparis greeni, JORDAN & STARKS, Proc. Cal. Ac. Sci., 1895, 829, pl. 96, Victoria, Vancouver Island. (Type, No. 3010, L. S. Jr. Univ. Mus. Coll. Ashdown H. Green.)

2448. NEOLIPARIS FISSURATUS, Starks.

Head $3\frac{1}{4}$ in body; depth $4\frac{1}{4}$. D. VI, 28; A. 26; P. 36; C. 14; eye 7 in head; maxillary $2\frac{1}{6}$; ventral disk $2\frac{1}{2}$. Body moderately elongate, not produced at nape; mouth rather large, the maxillary extending to below middle of eye; jaws subequal; teeth tricuspid, arranged in about 10 oblique series in each jaw; nostrils ending in short, wide tubes; gill openings wider than in any other known *Neoliparis*, commencing a distance above pectoral about equal to the diameter of eye, ending about opposite the fourteenth ray; ventral disk a little longer than wide, its distance from chin equal to $1\frac{1}{2}$ its longest diameter, its posterior edge about the same distance from front of anal; vent nearer anal than ventral disk, its distance from anal equal to $\frac{1}{2}$ ventral disk. Origin of spinous dorsal at the vertical from midway between vent and ventral disk; dorsal scarcely joined to caudal, anal very slightly; front of anal nearer chin than base of caudal by a distance equal to ventral disk; pectoral rather short and wide, reaching to opposite front of anal, the lower lobe very narrow and long, much longer than upper lobe, but not reaching so far posteriorly on account of the oblique position of the fin; tip of lower lobe reaching to vent; length of caudal $1\frac{3}{8}$ in head. Color dusky, darker above, sides with fine punctulations; belly and under parts of head, except chin, white; lips dusky; dorsals and anal darker than body; pectoral dusky at base, the lower lobe dark; caudal crossed with wavy dark lines. This species differs from the other species in this

genus in having a wider gill opening, and in various minor characters. Length 2½ inches. Puget sound; only the type known. (Starks.) (*fissuratus*, split.)

Neoliparis fissuratus, STARKS, Proc. Cal. Ac. Sci. 1896, 560. (Type, No. 5044, L. S. Jr. Univ. Mus. Coll. E. C. Starks.)

785. LIPARIS* (Artedi) Scopoli.

(SEA SNAILS.)

Liparis, ARTEDI, Genera, 117, 1738 (nonbinomial).
Cyclogaster, GRONOW, Museum, 1763, 157 (nonbinomial).
Liparis, SCOPOLI, Introd. Hist. Nat., 453, 1777 (*liparis*).
Liparis, CUVIER, Règne Anim. Ed. 1, 1817 (*liparis*; not *Liparis*, Ochsenheimer, 1810, a genus of *Lepidoptera*).
Cyclogaster, GRONOW, Cat. Fishes, Ed. Gray, 40, 1854 (*liparis*; not of Macquart, 1854, a genus of flies).
Actinochir, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 193 (*major*).
Careliparis, GARMAN, Discoboli, 56, 1892 (*agassizii*).
Lyoliparis, JORDAN & EVERMANN, Check-List Fishes, 451, 1896 (*pulchellus*).

Body rather elongate, covered with smooth skin, which is usually freely movable; head short; flattened above; mouth horizontal, the jaws equal or the lower jaw included; teeth in several series, close set, always more or less tricuspid, the adult with the outer cusps often worn or obliterated; maxillary covered by skin of preorbital region; anterior nostrils tubular or not; ventral disk well developed on the breast, its front below or behind the middle of the head, its surface with 13 lobes; an anterior median lobe, and 1 corresponding to each of the 6 rays in each fin; each lobe with a horny papilla covering, which is sometimes lost; vent well behind the head, about midway between the sucking disk and anal fin; dorsal fin continuous, undivided, its spines not differentiated; caudal well developed; dorsal fin free from caudal or joined; pectoral broad, procurrent at base, emarginate and free at tips, some of the lower rays produced; vertical fins enveloped in the lax skin; vertebræ 35 to 55. Northern seas near the shores; the species less arctic in distribution and in general inhabiting shallower water than is the case with *Careproctus* and *Paraliparis*, a fact associated with the reduced number of vertebræ in *Liparis*. The species are numerous, but in general well defined, their characters varying with age. In most of the species color varieties occur, several (*pulchellus, liparis, aleuticus, agassizii*) having the body often marked everywhere with concentric curved stripes or rings. (λιπαρός, sleek-skinned.)

Concerning the species of *Liparis* (including *Neoliparis*), Mr. Garman observes:

> The fishes placed in this genus are mostly small; in general their outlines resemble those of larval anurous batrachians. The anterior portion of the body, containing the abdominal chamber, is usually short, broad, flattened beneath, and somewhat angular

* *Enantioliparis*, Vaillant, is based on *pallidus*, an Antarctic species with the pectoral very narrow and not emarginate; would seem to be distinct from *Liparis*.

toward the dorsal fin, while the hinder part is elongate, much compressed, and tapering. The more important of the distinguishing features may be summarized as follows: Head short, broad, rounded; snout short, blunt; mouth anterior; lips distinct; teeth small, numerous, in a cardiform band, tricuspid on intermaxillaries and dentary, simple on pharyngeals; eyes lateral, small to medium; nostrils, 2 on each side, with short, projecting tubes; olfactory nerve short; brain elongate; glossohyal rudimentary or absent; gills 3 double and 1 single; gill rakers with denticles; pseudobranchiæ present; gill opening small, above the pectoral; branchiostegal rays 6; branchial membranes not free; tubes and pores of the lateral system forward of the gill aperture; dorsal and anal elongate, more or less closely united with the caudal, with some anterior rays unsegmented; pectorals broad, procurrent under the throat; ventrals transformed into an adhesive disk, situated below the gills between the pectorals; skin thin, loosely attached, smooth, excepting a sexual outgrowth on males in breeding season; stomach siphonal; pyloric cæca numerous; intestine elongate; urinary bladder large, simple; kidneys long, fused a portion of their length; liver large, with partial divisions into lobes; gall bladder small, rounded; skeleton not heavily ossified, less in deep-sea forms; upper limb of preoperculum expanded backward; operculum small; suboperculum narrow; interoperculum elongate, styliform; third suborbital produced backward, in a long, slender process, to the hinder edge of the preoperculum.

The species of *Liparis* inhabit the colder waters of the north and of the south of both Atlantic and Pacific, ranging to great depths. Among those frequenting the shores, if not also the others, vegetation seems to form a portion of the food. The contents of the stomachs are miscellaneous lots of small marine animals (crustacea, worms, mollusks, small fishes, etc.), mixed with which are quantities of seaweeds. Some variation exists in regard to feeding habits among the different species. The teeth of some show no signs of wear from hard food, but in *L. mucosus* they are ground off and blunted, no doubt by contact with the hard-shelled crustacea and hard-scaled agonoid fishes, remains of which are found in the stomachs. Very likely *L. pulchellus* turns his peculiar snout to account by rooting in the mud or sand among the plants, or in turning over the pebbles. Certain species are fitted for a life in beds of seaweed, others are probably more abundant in rocky places.

LIPARIS:

a. Vertebræ in moderate number, about 39; dorsal rays about 35; anal rays 27 to 30.

b. Gill openings very narrow, entirely above base of pectoral; pectoral rays from 34 to 37; head a little shorter than broad, and a little longer than deep; dorsal and anal slightly joined to caudal; caudal narrow, its rays 12. LIPARIS, 2449.

bb. Gill openings broad, the lower part considerably below base of upper ray of pectoral.

c. Pectoral rays 30; head low, flattish, ⅓ longer than broad, ⅓ broader than deep; jaws subequal; dorsal free from caudal, which is slightly joined to anal; caudal narrow, of 12 rays. CYCLOPUS, 2450.

cc. Pectoral rays 41 to 43; head short, not quite as wide as long; caudal 15 to 20; the dorsal and anal slightly joined to its base. FUCENSIS, 2451.

CARELIPARIS (*καρά*, head; *Liparis*):

aa. Vertebræ about 46; dorsal rays 39 to 44; anal rays 33 to 36; dorsal and anal largely joined to caudal.

d. Pectoral rays 30 to 36.

e. Gill opening small, its lower edge not below first ray of pectoral; nostrils small, the tubes short or absent.

f. Posterior nostril scarcely tubular; fins plain, not distinctly mottled or barred; body robust; its color plain brownish or with dark spots.

g. Dorsal rays 39 or 40; anal 33; pectoral 30 to 34. TUNICATUS, 2452.

gg. Dorsal rays 44; anal 35; pectoral 37. HERSCHELINUS, 2453.

ff. Posterior nostril tubular, fins more or less mottled or barred, body moderately elongate; lower rays of pectoral rather short, not ½ head, not reaching beyond ventral disk; body mottled usually with concentric rings. AGASSIZII, 2454.

ee. Gill opening rather large, extending downward to about fourth ray of pectoral; nostrils with short tubes; lower lobe of pectoral long, reaching much beyond disk, nearly to vent; color brown, the body and fins mottled and clouded. DENNYI, 2455.

dd. Pectoral rays 42; gill opening large, its lower edge below upper part of pectoral; body robust, translucent, covered with round crimson spots. CYCLOSTIGMA, 2456.

aaa. Vertebræ about 52; dorsal rays 45 to 48; anal rays 38 to 40; pectoral rays 34 to 37; dorsal and anal largely joined to caudal; gill opening large, about ⅓ its length before pectoral; anterior nostril tubular.

LYOLIPARIS (λύω, to loosen; *Liparis*):

h. Head broad, flattened above; body rather elongate; skin usually with wavy concentric longitudinal streaks, sometimes spotted. PULCHELLUS, 2457.

ACTINOCHIR (ἀκτίς, ray; χείρ, hand; from the broad pectoral):

hh. Head high, boldly rounded, with prominent nape; color olivaceous, clouded and dotted, but without wavy streaks. MAJOR, 2458.

Subgenus LIPARIS (Artedi) Scopoli.

2449. LIPARIS LIPARIS * (Linnæus).

(SEA SNAIL.)

Head 3½; depth 3½; D. 33 to 35; A. 27 to 29; pectoral 34 to 37; caudal 12 to 14; vert. 38 or 39; cæca 10 to 13; branchiostegals 6. Body moderately stout, somewhat high on the shoulders, compressed behind the body cavity in the caudal portion, which is deep and rather thick anteriorly, tapering rapidly. Body proper about ½ of the entire length to the base of the tail. Length of the head little less than its width, a little more than its depth. Snout short, broad, blunt, nearly 3½ times in the head, very convex in the internasal region. Mouth moderate, maxillary reaching a vertical from the eye, lower jaw shorter. Teeth small, paved, alternating in the different rows, tricuspid. Lower lip interrupted in the middle, ⅓ or more of its length. The fold above the upper lip with 6 to 8 notches below the opening of the pores. A series of 6 pores on each side, from the middle of the fold above the upper lip passes backward and

* Prof. Robert Collett (Christiania Vidensk. Selsk. Förh., 1879, No. 1, 42), defines the chief color varieties of this species as seen in Norway, as follows:

Var. a. *assimilis*, Collett: uniform light reddish gray, the dots microscopic. Christiania, Bukn, Stavanger, Lofoten; scarce.

Var. b. *stellatus*, Malm: uniform light gray brown, the fins with vertical dark cross bands; dots larger. Christiania, Vadsø, etc. *Liparis stellatus*, Malm, Förh. Skand. Nat., 9, 1863, 412.

Var. c. *subfuscus*, Collett: almost uniform ground color, with dark points; light red cross bands on fins; body faintly marbled. Christiania, Farsund, Varanger.

Var. d. *scorpioides*, Collett: color pale, with large, irregular, dark spots. Varanger.

Var. e. *mixtus*, Collett: spotted and striped; dark spots on fins; along sides and on fins interrupted dark stripes: ring-like lines on head. Lofoten.

Var. f. *decorus*, Collett: striped and spotted; fins spotted; notable stripes on sides. Christiansund and Lofoten.

Var. g. *scriptus*, Collett: striped, the dark stripes larger and more conspicuous, the streaks of ground color fainter. Bukn, Christiania.

Var. h. *lineatus*, Kröyer: striped with 8 to 10 violet longitudinal stripes, alternating with pale streaks of the ground color. *Liparis lineatus*, Kröyer, Nat. Tidskr. 1847, 284.

Var. i. *arcticus*, Collett: uniform brown; grayish golden brown with many dark points, which make the surface dark. Tromsö, Spitzbergen, Magdalene Bay, in Greenland.

Var. k. *fuscus*, Collett: uniform blackish brown, very dark. North Cape, Spitzbergen, Greenland.

As noticed under *Neoliparis montagui*, these variations seem to have no taxonomic value, being products of the immediate surroundings.

curves up behind the eye; another series of 7 on each side passes from the middle of the chin back and upward toward the upper edge of the gill opening. Eye small, nearly as long as the snout, contained 4½ times in the length of the head, or less than twice in the convex interorbital space. Nasal tubules short; posterior smaller between the eyes; anterior farther forward, larger. A large pore is situated a short distance in front of the nostril toward the tip of the snout. Gill openings narrow, not as wide as the eye, above the base of the pectoral. Nape and shoulders high. Cheeks swollen. Skin smooth, thin, easily detached. Dorsal and anal fins rising gradually, the rays attaining their greatest height in the hinder third of the length, united with the caudal by membrane. The separation of the 3 fins is less marked than in *N. montagui*, so much so that it is difficult in some cases to determine whether certain rays belong to the caudal. Dorsal notch shallow or absent, most noticeable in the breeding season. First ray little farther back than the opercular angle, or the hinder edge of the disk; 5 or 6 of the anterior rays crooked and twisted, and without apparent segmentation. Evidently these rays are affected by a nuptial growth, which is not so patent above the skin, on the ends of rays, as in other species. In the anal the first ray is about opposite the seventh or eighth ray of the dorsal; the fin extends farther back than the latter, reaching under nearly ⅓ of the caudal. Caudal slightly rounded posteriorly, the longest rays being ½ to ⅔ of length of the head. Disk subcircular, ½ as long as the head, equal the distance from the chin, less than the distance to the anal. Vent nearly halfway from disk to first anal ray, which latter is halfway between the snout and the base of the caudal. The position varies in the sexes. The body cavity is longer in the females, and the vent is farther back; this applies to all the species. Pectorals broad, deeply notched below the middle of the hind border; rays of the lower portion extending beyond the membrane, a couple of the longer sometimes reaching almost as far back as the vent. A slight fold of the skin unites the fins in front of the disk. Olive to brown, light to dark, puncticulate and clouded, blotched, banded, vermiculate, or longitudinally striped with darker and with lighter. In life the tints vary from lilac to reddish or yellowish, with spaces on certain individuals often nearly white, and every variety of coloration may be taken in a single locality. This being the case, it is thought best to treat the striping, freckling, banding, etc., merely as marks of individual variation, and not as characters indicating established varieties. On the striped specimens the number of lines vary greatly, of the spotted ones no two are alike, and the same is true of every other pattern. Most often there are blotches on the posterior portions of the anal and the dorsal; frequently they take on the appearance of transverse bands, as is still more often the case on the caudal. (Garman.) North Atlantic, on both shores; north to Spitzbergen, south to Connecticut and France; most abundant in Northern Europe. Variable. "The specimens in the museum at Cambridge were secured from the Cheshire coast and other points in Great Britain, from Ostend, and from various localities off the coast of Massachusetts, Connecticut, and Maine. A small specimen, with the

locality 'Off Block Island, from a pecten,' that may be positively identified, bears peculiar markings: From the tip of the snout a white line passes above the eye to meet its fellow of the opposite side at the beginning of the dorsal, which is white; and a white spot as large as the orbit extends downward and slightly backward from the eye. These marks are very distinct, and they give the specimen quite a different appearance from that of another of the same size and origin, uniform in coloration." (Garman.) (Eu.)

Liparis nostras, WILLUGHBY, Hist. Pisc., App., 17, 1686, based on a specimen taken September 15, 1685, by Dr. Johnson.

Liparis,* ARTEDI, Descr. Spec. Pisc., 117, 1738, after WILLUGHBY.

Liparis cyclogaster, GRONOW, Mus., 157, 1768.

Cyclopterus liparis, LINNÆUS, Syst. Nat., Ed. XII, I, 414, 1766, Northern Ocean; after ARTEDI and GRONOW.

Cyclopterus lineatus, LEPECHIN, Nov. Comm. Petropol., XVIII, 1774, 522, White Sea.

Liparis liparis, CUVIER, Règne Anim., Ed. I, vol. 2, 227, 1817; GARMAN, Discoboli, 57, 1892.

Gobius smyrnensis, BONNATERRE, Encycl., Icht., 66, 1788; after LEPECHIN.

Cyclopterus musculus, LACÉPÈDE, Hist. Nat. Poiss., IV, 683, 1802, near Havre.

Liparis vulgaris, FLEMING, Brit. Anim., 190, 1828; GÜNTHER, Cat., III, 159; LÜTKEN, Naturh. Foren. Vidensk. Meddels, 1860 and 1861, 243; COLLETT, Norges Fiske, 1875, 65; JORDAN & GILBERT, Synopsis, 742, 1883.

Liparis communis, SABINE, App. Parry's Voy., 1824.

Liparis barbatus, EKSTRÖM, Vet. Ak. Handl., 168, pl. 5, 1832 (D. 32, A. 32, C. 9, perhaps a different species).

Liparis ophidoides, SWAINSON, Nat. Hist. Fishes, II, 339, 1839.

Liparis lineatus multistriatus, LÜTKEN, Naturh. Foren. Vids., 1861, pl. 7 (striped example).

Liparis stellatus, MALM., Förh. Skand., Naturh., 1865.

Liparis lineatus with variations, *assimilis*, *stellatus*, *subfuscus*, *scorpioides mixtus*, *decorus*, *scriptus*, *lineatus*, *arcticus*, and *fuscus* COLLETT; Vid. Selsk., Förh., Christ. 1, 41.

2450. LIPARIS CYCLOPUS, Günther.

Head $4\frac{1}{6}$; depth $4\frac{1}{2}$. D. 34; A. 29; pectoral 30; caudal 12. Body much depressed and rather broad anteriorly, deep and much compressed posteriorly; head $\frac{1}{3}$ longer than broad and $\frac{1}{8}$ broader than deep. Flesh much more firm and the skin less lax than in most species of *Liparis*. Opercles with a rather strong spine concealed by the skin; mouth rather large, terminal; jaws subequal; teeth small, tricuspid, in broad bands; eye small, 6 in head; snout 3, flattish and broad above; interorbital space $4\frac{3}{8}$ in head; ventral disk oval, $2\frac{1}{3}$ in head, its anterior edge $\frac{1}{3}$ the length of the eye behind postorbital margin; gill opening moderate, $3\frac{1}{3}$ in head, extending downward to the third or fourth ray of pectoral. Dorsal fin low, continuous, not joined to caudal, beginning slightly before anal, on a vertical with vent; vent midway between edge of ventral disk and front of anal. Pectoral fin emarginate, the upper lobe $1\frac{3}{4}$ in head, the lower 2, the shortest intervening rays 3. Anal long and low, barely joined to caudal. Caudal $1\frac{1}{2}$ in head. Color olivaceous, darker above; body and pectoral fin finely speckled with olive brown; fins dotted; bases of the fins paler than their tips; belly white. Bering sea to Puget Sound, not common; our description from two specimens $4\frac{1}{2}$ inches long, in excellent

* Garman gives detailed synonymy of this species in Discoboli, 57, 1892.

condition, taken in Elliot Bay, near Seattle, received from the Young Naturalists' Society. They are numbered 3126 in the register of the Leland Stanford Jr. University Museum. This species, not having been previously recognized since its original description, is recorded by Dr. Gilbert from Unalaska, and 2 specimens from Bristol Bay, Alaska (*Albatross* Station 3230, depth 3¼ fathoms). Garman places *L. cyclopus* in the synonymy of *L. callyodon*, but our specimens are undoubtedly distinct from the species described by Garman under this name (Discoboli, p. 54, pl. 6, figs. 1–5), from a specimen said to originate from San Francisco. Our Alaska specimens are much more slender, with wide depressed head, without nuchal elevation, with the dorsal fin beginning posteriorly slightly in front of the vertical from the vent, and the disk separated from the vent by a distance less than its own diameter. The mouth is also much smaller, not at all oblique, its angle in advance of vertical from front of eye. Not yet recorded from California. (*κύκλος*, rounded; *πούς*, foot.)

Liparis cyclopus,* GÜNTHER, Cat., III, 162, 1861, Esquimault Harbor, Vancouver Island; JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 834, pl. 97.

2451. LIPARIS FUCENSIS, Gilbert.

Head 3½ to 3⅘; depth 4 to 4⅔. D. V, 30; A. 28 or 29; C. 18 or 20; P. 38 to 39. Body moderately elongate, compressed; head depressed, with gibbous snout and occiput, separated by the depressed interorbital area, which forms a shallow transverse groove. Snout not blunt, the mouth terminal, nearly horizontal, with included mandible, the maxillary reaching to or nearly to the vertical from front of pupil, 3 in head. Teeth all trienspid. Eye of moderate size, contained 5⅔ in length of head, 1 to 1⅓ times in bony interorbital width, 1⅜ times in snout. Posterior nostril without tube, the anterior with a short tube, less than diameter of pupil. Gill slit comparatively wide, its width equaling length of snout and ⅓ eye, overlapped by a conspicuous triangular prolongation of the opercle. The slit extends down to opposite the upper third of the pectoral fin. Disk circular, of rather small size, distant from tip of snout 1⅓ times its own diameter, from vent 1¼ times. Diameter of disk 2⅔ times in head. Distance from tip of snout to vent 1⅔ to 1⅜ in distance from tail. Pectorals extending to a vertical midway between vent and front of anal. Lower rays produced, forming a narrow distinct lobe. First 5 dorsal rays spinous, unsegmented, shorter than the succeeding segmented rays, from which

* The following is the original description of *Liparis cyclopus:* Dorsal 32; anal 30; caudal 12. Caudal fin entirely free from dorsal and anal; the latter commences in the vertical from the seventh dorsal ray; pectoral margin with a notch, some of the lower rays being produced; ventral disk ovate, its length being less than ½ of that of the head. A fine specimen. Esquimault Harbor. Presented by Lord John Russell. This species agrees in nearly every respect with *Liparis vulgaris*, from which it differs in the following points besides those mentioned: The greatest depth of the body is nearly ¼ of the total, the length of the head nearly ⅕. The skin is tougher and more firmly adherent to the body. The head is flat above, the nape of the neck being scarcely prominent. The nostril is a short tube, situated nearly midway between the eye and the upper lip. The pectoral extends only to the vertical from the anal papilla, and the dorsal (anal) fin commences yet more backward. The teeth are small, with 3 lobes. Infraorbital extending to the preoperculum. The color is brownish olive, with darker spots, more conspicuous on the fins than on the body; belly reddish. The length of the specimen described is 45 lines. (Günther.)

they are not separated by notch. Dorsal and anal free from the caudal, the last rays being rapidly shortened, giving a rounded contour to the posterior portions of the 2 fins. Two styles of coloration are observed; 1 plain olive brown, with minute dark points, whitish below; the other, with numerous lengthwise streaks of light olive and dark olive brown, which extends forward on top and sides of head; in both cases the belly is whitish, and the fins dusky, mottled with darker, the mottlings forming indistinct cross bars on the caudal fin. Numerous specimens dredged by the *Albatross* in the Straits of Fuca (August, 1891), serve as the types of this species. It is probably the same as that described by Garman as *Liparis calliodon*, his description being based on specimens "said to have been taken near San Francisco." It is not evident from the text whether the same specimens served as basis for the figures (pl. 6, figs. 1 to 5), concerning which we have no independent data. This locality may be correct. The numerous types of Dr. Gilbert's examined by us are in a collection made by the *Albatross*, about Port Angeles in the Straits of Juan de Fuca. It has not been yet recorded from Alaska. Mr. Garman identifies this species with the *callyodon* of Pallas, but according to Pallas his species had the gill opening reduced to a lunate spiracle which is not the case in *Liparis fucensis*. (*fucensis*, from the Straits of Juan de Fuca.)

Liparis calliodon, GARMAN, *Discoboli*, 54, 1892, locality unknown, thought to be San Francisco; not *Cyclopterus callyodon* of PALLAS.

Liparis fucensis. GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 447, Port Angeles, Straits of Juan de Fuca, at Albatross Station 3451, in 106 fathoms. (Type, No. 48600.)

2452. LIPARIS TUNICATUS, Reinhardt.

D. 39; A. 33; P. 30; V. 7; C. 14. Body elongate, subterete; belly prominent. Head thick, obtuse, depressed behind, convex in front. Mouth narrow; the jaws equal; jaws, suborbital region, and gills with small, white points. Eyes small, lateral; nostrils 4, tubulate. Gill opening small, linear. Dorsal fin and anal united to caudal; caudal cuneate. Skin lax, slippery. Color dusky, the flesh reddish, showing through the skin. Length 1½ fingers. (Fabricius.) Coast of Greenland. This description differs from later ones in ascribing more rays to the caudal, and tubes to the nostrils. According to Mr. Garman, the description of *Liparis fabricii*, Kröyer, refers to the same species. The following is its substance: D. 13 + 27 = 40; A. 2 + 30 = 32; P. 34; C. 9; Vert. 46. Head 4 in length, higher than broad, the snout elevated, truncate, lower jaw shorter; anterior nostrils simple, the posterior with raised rim or with very short tube; eye 3 in head; teeth tricuspid; ventral disk elliptical, 9 in length of body; dorsal and anal connate with caudal, which is 6 to 8 in total length. Ashy above, shaded with bluish or orange, spotted with blackish, pale below. Greenland (Kröyer, description of *Liparis fabricii*). *Liparis arctica*, Gill, is apparently the same as *L. fabricii*. The following is the substance of the description: Head 4 in total length; depth 2¾; dorsal 42; anal 34; caudal 11; pectoral 35. Forehead depressed, snout rather high and decurved; the eyes are just within the anterior half of the head; the inter-

orbital is contained 3 times in the head; the anterior nostrils are simple, the posterior tubular; the dorsal and anal are connate with the caudal, the former united for about $\frac{1}{4}$ of the length of the caudal, the latter $\frac{2}{5}$; caudal convex behind and forms $\frac{1}{7}$ of the length, the pectoral $5\frac{1}{3}$; ventral disk 10. Color uniform, rusty brown or chestnut, without spots or bands. Port Foulke, Greenland (Gill). We have seen no specimens, and follow authority in regarding Fabricius's count of the caudal rays as probably an error. Garman uses the name *tunicatus* instead of *minor*, apparently regarding the original use of the latter name as simply that of the abjective meaning smaller, not as part of the binominal nomenclature. (*tunicatus*, bearing a cloak.)

Cyclopterus liparis minor, FABRICIUS, Fauna Grœnlandica, 135, 1780, Greenland; D. 39; A. 33; P. 30; V. 2; C. 14; not binomial, *minor* being merely an adjective, meaning smaller.

Liparis tunicata, REINHARDT, Overs. Kong. Danske Vidensk. Selsk., VI, CXI, 1836, Greenland; after FABRICIUS; GARMAN, Discoboli, 65, 1892.

Liparis fabricii, KRÖYER, Naturh. Tidsskr., II, 274, 1847, Greenland; D. XIII, 27; A. II, 30; P. 34; C. 9; Vert. 46; GÜNTHER, Cat., III, 161, 1861; GILL, Proc. Ac. Nat. Sci. Phila. 1864, 192.

Liparis arctica, GILL, Proc. Ac. Nat. Sci. 1864, 191, Port Foulke, Greenland; JORDAN & GILBERT, Synopsis, 742, 1883.

2453. LIPARIS AGASSIZII, Putman.

Head $3\frac{4}{5}$; depth 4. D. 39; A. 31; pectoral 34; caudal 12; eye 6 in head; interorbital $2\frac{1}{2}$; snout $2\frac{1}{2}$; ventral disk $1\frac{5}{6}$; pectoral $1\frac{1}{4}$; caudal $1\frac{4}{5}$. Body elongate, compressed posteriorly, about as wide as deep anteriorly; head as deep as broad, $\frac{1}{5}$ longer; mouth moderately large, the maxillary reaching slightly past front of eye; lower jaw included; teeth small, tricuspid, arranged in 8 or 9 oblique rows, which grow more oblique toward the sides; snout broad and blunt, slightly overhanging the mouth; nostrils ending in tubes, the anterior tube wider and longer than posterior; nape high and prominent, scarcely produced; gill opening short, its lower $\frac{1}{8}$ or $\frac{1}{4}$ in front of the pectoral, extending downward to about the second or third ray, its length a little greater than width of eye. Ventral disk slightly longer than broad, its distance from end of lower jaw $1\frac{3}{8}$ its length, its posterior margin scarcely midway between its anterior and front of anal; vent nearer posterior margin of disk than front of anal; pectoral broadly rounded behind, the notch made by the produced lower rays shallow, end of fin reaching to the vertical from front of anal; origin of dorsal just over posterior edge of ventral disk, its distance from snout about 3 in body, slightly joined to the caudal posteriorly; origin of anal nearer to snout by a distance equal to length of snout posteriorly joined to the caudal, the tips of the last rays reaching nearly to the middle of caudal rays. Color olive brown, irregularly marked with dark brown spots the size of pupil; belly white; top of head darker than body; dorsal spotted like the body; spots on anal in the form of indistinct bands; pectoral marked with interrupted irregular streaks; a wide dark band across middle of caudal rays, 2 or 3 other narrow streaks across caudal. Here described from a specimen collected by the *Albatross* at Station 3247, in Bristol Bay, Alaska, in 17 fathoms. Its length is nearly 3 inches.

The following is Garman's description of a specimen which was 10 inches in length:

Head 4½ in total length; depth over 4. D. 41 (41 to 44). A. 33 (32 to 35); pectoral 38 (35 to 38); caudal 12; vertebræ 46. Body elongate, rather less inflated anteriorly than the other Liparids, greatly compressed posteriorly. With the exception of the disk the appearance is very much the same as that of the Cottoids. Total length 2¾ times the distance from snout to vent. Head moderately broad, depressed anteriorly, slightly convex in transverse section through the frontal region; snout broad, blunt, rounded, convex, nearly 4 times the diameter of the eye, ⅛ of the length of the head. Mouth wide, maxilla extending behind a vertical from the anterior border of the eye; upper lip complete, lower separated for only about ½ the distance from the angle of the mouth to the middle. Teeth small, tricuspid, with slender, somewhat compressed bases, cusps compressed to sharp edges, outer cusp turned outward so as to give an inflated appearance at the top, arranged in about 72 rows, counting lengthwise on each jaw, or 10 rows transversely, from outer to inner. Pharyngeal teeth simple, in 2 bunches of 8 or 10 rows at the esophagus on the roof of the mouth, and below these on the floor in 2 smaller elongate groups. Nostrils small, tubes short or absent, posterior between the anterior halves of the eyes, anterior in front of these about 1 diameter of the orbit, and twice the same distance from each other. Eye small, less than ⅛ of head. Gills 3½. Gill opening a vertical slit, extending downward a short distance on 1 or 2 rays only of the base of the pectoral (not so wide as in *L. pulchellus*). Skin thin, loose. Backward from the vent the form is much compressed, the thickness being nearly ½ of the height, and both lateral and dorsal outlines taper regularly to the caudal. Dorsal, anal, and caudal fins confluent, anal extending farther on the caudal, i. e., more than ½ the length of the latter; dorsal commencing above the gill opening, at a distance from the snout that is contained in the total length without the caudal 3⅝ times; anal beginning a trifle in advance of the mid length, excluding the caudal; pectoral broad, not reaching the anal, upper portion rounded, lower fringed, a shallow indentation between the two portions; caudal subtruncate, rounded, with rays more than ½ as long as the head; ventral disk small, ¼ longer than broad, situated its length from the lips, occupying nearly ⅛ of the distance from the snout to the first ray of the anal, or about ⅛ of the total length. Color dark brown, irregularly marked with whitish or grayish; a series of 5 or 6 spots, each as large as the eye, along the middle of the flank; below these, near the lower edge, there are indications of 4 or 5 whitish blotches, resembling transverse bands; dorsal, anal, and caudal with irregular cloudings and blotches of darker and of lighter; base of pectoral with a couple of large blotches of light grayish, balance of the fin freckled with light color; head dark, somewhat reddened. Entire length 10 inches. (Garman.) North Pacific; common south to Unalaska; the original types above described were collected by Messrs. Pierce and Smith at Saghalien, Channel of Tartary; our specimens collected by the *Albatross* in Bristol Bay. Dr. Bean's types of *Liparis gibbus* came from Plover Bay, Siberia; others are

recorded from Unalaska, St. Paul Island, off Indian Point, Cape Chaplin, Eastern Siberia, Petropaulski, and Bering Strait.

The following is the substance of the account of *Liparis gibbus* which Garman identifies, apparently correctly, with *Liparis agassizii:*

Head 3½; depth 3¼. D. 42; A. 36; P. 85; C. 12. Body abruptly contracted near the vent, covered with lax skin; interorbital space shallow-concave, the vertex and nape somewhat elevated; snout depressed; head as wide as long, longer than deep; nostrils tubular, the tubes of anterior nostrils longest; eye small, 4 in head; ventral disk nearly circular, 8 in length; vertical fins confluent; dorsal continuous; longest dorsal ray ½ as long as head; pectorals reaching front of anal; caudal 6 in body. Head and body very pale brown or gray, paler below; head and anterior parts sometimes with concentric brown rings, much as in *Liparis pulchellus;* sides plain or marked with brown stripes and rings; tail sometimes with dark blotches; vertical fins usually with dark bands. Bering Sea, Aleutian Islands, and Siberia. (Named for Prof. Louis Agassiz).

Cyclopterus liparis, Bloch, Ausländ. Fische, I, 48, 1785, in part, Pacific specimens.
Liparis agassizii, Putnam, Proc. Amer. Assoc. Adv. Sci. 1874, 339, Saghalien, Channel of Tartary (Coll. Pierce and Smith); Garman, Discoboli, 62, pls. 1–3, 1892.
Liparis gibbus, Bean, Proc. U. S. Nat. Mus. 1881, 148, Unalaska, St. Paul Island, Indian Point, Cape Chaplin, and Plover Bay, Siberia; Jordan & Gilbert, Synopsis, 741, 1883.

2454. LIPARIS HERSCHELINUS, Scofield.

Head 3½ in body; depth 3¾. D. 42; A. 33; pectoral 35; caudal 10; eye 4½ in head, and 1½ in snout; interorbital space 3 in head; maxillary 2 in head. Body tadpole-like; head rounded and very little compressed; abdomen slightly distended; just back of the abdomen the body is suddenly compressed to a width equaling ⅛ its height, and from this point the body gradually tapers to the caudal, its height and width keeping the same proportions; the height of the base of the caudal equals the diameter of the eye; the maxillary extends to posterior edge of eye, and its end is concealed in the base of the skin of the head; upper jaw slightly longer than lower; teeth tricuspid; interorbital space flat; nape slightly elevated; gill openings small; the width of the slit equaling the interorbital space; the lower edge of the slit even with the first pectoral ray; the posterior nostrils end in very short, compressed tubules about ½ diameter of eye in front of eye; the anterior nostrils are simple and placed directly in front of the posterior nostrils a distance equal to ½ diameter of eye. The dorsal begins on a vertical line drawn from posterior edge of gill flap; the anterior rays are short, gradually lengthening till middle of fin is reached, where the rays equal 2½ times the diameter of the eye, the last rays scarcely shortened and not forming a notch at its junction with the caudal; last rays encroaching on the caudal for ¼ of its length; anal same shape as dorsal and of same height, its last rays encroaching on caudal for ⅓ its length; upper lobe of pectoral composed of 25 rays; the eleventh and longest ray 1½ in head; length of twenty-fifth ray 4 in head, lower lobe with 10 rays, of which the third from the last, or twenty-third, is the long-

est, being contained in the head $2\frac{1}{10}$ times; membrane between each 2 of the last 4 rays incised; caudal slender, rounded behind, its length a little more than twice in head. Diameter of ventral disk 8 in body. Skin very loose, attached only at opening and to ends of last rays of dorsal and anal and to base of caudal. Color dark, covered with fine punctulation; belly and underside of head free from markings; last half of dorsal darker than anterior half; anal mottled with black; caudal with 2 black vertical bands, the first bony and just back of tips of last dorsal and anal rays; the second band slender and faint and at top of fin. Arctic Ocean; several specimens were taken at Herschel Island. The longest specimen is $2\frac{1}{2}$ inches long. This species is nearest *L. tunicatus* from the coast of Greenland, and appears to be intermediate between it and *L. agassizii*.

Liparis herschelinus, Scofield, Fishes of Herschel Island, in Jordan & Gilbert, Rept. Fur Seal Invest., 1898, Herschel Island, Arctic Ocean. (Type, No. 5601, L. S. Jr. Univ. Mus. Coll. N. B. Scofield.)

2455. LIPARIS DENNYI, Jordan & Starks.

Head $3\frac{3}{5}$ in length of body; depth $4\frac{1}{2}$. D. 39; A. 30; pectoral 36; caudal 12; eye 8 in head; maxillary $2\frac{1}{5}$; snout $2\frac{3}{4}$; gill opening $2\frac{2}{5}$; upper pectoral lobe $\frac{1}{3}$; lower lobe $1\frac{1}{2}$; intervening rays $2\frac{1}{4}$; ventral disk $2\frac{1}{3}$; highest dorsal rays $2\frac{1}{5}$; highest anal rays $2\frac{2}{5}$; caudal rays $1\frac{3}{4}$. Body moderately elongate, much compressed posteriorly, slightly so anteriorly; head moderate, the cheeks and nape prominent. Mouth wide, with little lateral cleft; maxillary extending to below the anterior margin of eye, its end covered with the skin of the head; the lower jaw slightly the longer; the teeth tricuspid, those on the inner part of jaw largest, arranged in about 14 series in each jaw; series nearly transverse on middle of jaw, becoming more and more oblique toward the sides, where they are nearly parallel with the sides of the jaws; interorbital wide, slightly concave; nostrils ending in very short, wide tubes, the posterior over the anterior margin of eye, the anterior in front of it a distance equal to the diameter of eye; opercle ending in a short, wide spine covered with skin; it is situated slightly above the middle of gill opening; gill opening running from about the eleventh pectoral ray to a level with the eye. Origin of dorsal slightly behind base of pectoral, its distance from the snout $3\frac{1}{5}$ in length of body, its anterior rays short, gradually lengthening posteriorly, the rays from the anterior third to near the end about equal, the last ray abruptly shortened, forming a slight notch where the fin joins the caudal; posterior $\frac{3}{8}$ of caudal free above; anal similar to dorsal, about the same height, its origin nearer snout than base of caudal, about under the base of the tenth dorsal ray, posteriorly it is longer than the dorsal, joining the caudal at about $\frac{1}{2}$ its length; ventral disk nearly round, its distance from tip of lower jaw $1\frac{1}{8}$ in its diameter, 1 in distance from vent, 2 from first anal ray; vent midway between front of anal and edge of disk; upper lobe of pectoral broadly rounded, reaching to $\frac{3}{8}$ of the distance between vent and front of anal; lower lobe long, reaching nearly to vent; caudal long and slender, rounded behind. Skin very thin and loose on body and head, covering the anterior parts of dorsal and anal, attached at about the

middle of rays posteriorly, and covering the base of caudal rays. Color light brown, lighter below, thickly covered with minute brown points, which form spots and mottlings on sides; upper part of head dark, lips spotted with brown; dorsal and anal dark brown, slightly mottled with lighter; pectoral light, with irregular brown spots and bars running across it; caudal dark brown, mottled at base, 2 light bars crossing it toward its end, leaving a narrow posterior margin of brown. North Pacific, south to Puget Sound; numerous specimens in the *Albatross* collections from about Unalaska. (Named for Mr. Charles L. Denny, of Seattle, in recognition of his active and intelligent interest in the natural history of Washington.)

Liparis dennyi, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 835, pl. 98, Admiralty Inlet, Puget Sound. (Type, No. 3703, L. S. Jr. Univ. Mus. Coll. Young Naturalists' Society of Seattle.)

2456. LIPARIS CYCLOSTIGMA, Gilbert.

Head $3\frac{3}{5}$; depth $4\frac{1}{2}$. D. 44; A. 34; P. 42; C. 14. A robust, compressed species, with broad, gently convex head, the nape rather elevated, comparatively wide gill opening, a single continuous dorsal fin, the dorsal and anal broadly joined to the caudal, and the coloration peculiar. Profile gently and evenly declining from nape to end of premaxillary processes, thence descending more steeply to tip of snout. Interorbital space very wide, equaling length of snout and $\frac{1}{2}$ of eye, $2\frac{2}{3}$ in head. Distance from tip of snout to front of exposed portion of eye, $2\frac{9}{10}$ in head. Mouth terminal, broad and transverse, with but little lateral cleft, the 2 jaws equal, the lower not included. The maxillary is entirely bound down by skin of head, reaching vertical from front of pupil, the angle of mouth in advance of eye. Bands of teeth extremely broad, the teeth very small, all tricuspid, the outer ones minute, those toward inner margin of jaw increasing in size. The anterior series in each jaw are nearly transverse, the lateral series becoming successively more and more oblique, the uppermost nearly parallel with the jaw; about 20 series in each side of lower jaw, 30 on each side of upper jaw. The width of band in upper jaw equals $\frac{2}{3}$ diameter of exposed portion of eye, which is $\frac{1}{4}$ length of snout, $\frac{2}{7}$ interorbital width. Nostrils without tube. Lower lip distinct on lateral $\frac{3}{5}$ or $\frac{2}{3}$ of mandible. Gill opening wide, extending downward to opposite base of fifteenth pectoral ray, the length of the slit $2\frac{2}{3}$ in head. Upper jaw with a series of large pores; lower jaw short, with few pores. Disk large, oblong, its longitudinal diameter $2\frac{1}{3}$ in head, equaling its distance from anus and twice distance of latter from base of first anal ray. Pyloric cæca 28. Pectoral very broad, inserted low, its upper margin on a level with premaxillaries, much below the eye; the rays decrease but little in length from the first to the twentieth, and form a very broad, evenly rounded lobe. Below the twentieth the rays decrease gently and have exerted tips, until the shortest ray equals $\frac{2}{3}$ the long upper rays. There follow 3 or 4 somewhat longer rays, the tips still further exserted, then 4 or 5 rays which decrease rapidly, the shortest anterior one equaling diameter of eye; longest pectoral ray $1\frac{1}{7}$ in head; base of first dorsal ray in a vertical passing through axil of pectoral; longest dorsal ray $1\frac{2}{3}$ in

head, the last rays rapidly shortened so as to produce a notch at union with the caudal, the last ray less than ⅔ the longest, the membrane joining at end of basal third of caudal; the anal fin is equal in height to the dorsal, but the last rays are but little shortened, so that no notch exists posteriorly. It forms a much broader union with the caudal, which it joins at the end of its basal two-thirds; caudal broad, rounded, the outer rays ⅕ the length of middle rays, which equal the length of head without the snout. Body everywhere transparent, gelatinous in structure. Colors in life, olivaceous above, overlaid with light grayish; belly and lower side of head light yellow; body and fins with large brownish-red or blood-red spots and blotches, usually roundish, each having a darker margin surrounded with a light ring. Length 360 mm. Bering Sea; type from Bristol Bay, in 29½ fathoms; a second specimen from St. Paul Island, and a third still larger from Petropaulski; a most beautifully colored species, quite unlike the others. (Gilbert.) (*κύκλος*, round; *στίγμα*, spot.)

Liparis cyclostigma, Gilbert, Rept. U. S. Fish Comm. 1893 (1896), 446, Bristol Bay, Alaska, at Albatross Station 3252, in 29½ fathoms. (Type No. 48621.)

Subgenus LYOLIPARIS, Jordan & Evermann.

2457. LIPARIS PULCHELLUS, Ayres.

Head 4⅓ in length; depth 4½. D. 47 to 49; A. 40; pectoral 36; caudal 12; eyes 6 in head; interorbital space (flesh, not bone) 2; snout 3; pectoral 1⅓; ventral disk 2⅔; caudal 1⅘. Body elongate, somewhat compressed, tapering gradually to the caudal. The head is broader and longer than high; snout subtruncate, overhanging the mouth; teeth small, tricuspid, the median cusp longest, arranged in oblique rows as in the related species; upper lip complete, the lower interrupted on each side about midway between middle of jaw and angle of mouth; interorbital space wide and flat, posterior nostril ending in a short, wide tube, anterior smaller, closer together, about midway between the posterior and end of snout; gill opening moderate, wider than eye, its lower third in front of pectoral, extending downward to the third or fourth ray; skin thin and loose, as in the other Liparids; vent nearer to anal than ventral disk. Dorsal and anal confluent with the caudal, which seems to end in a point when not spread; origin of dorsal just behind base of pectoral, its distance from snout contained 4 times in the length of body, the anterior rays buried under the skin; origin of anal nearer snout than base of caudal fin; pectorals broadly rounded posteriorily, reaching to front of anal, the lower rays smaller and produced, forming a notch in lower part of fin. Color light olive brown, with numerous waving lines everywhere on head and body, running longitudinally, sometimes running together and forming irregular markings; belly and lower parts of head white; dorsal and anal dusky, with longitudinal stripes and mottlings; caudal dark at tip; pectorals with irregular cross streaks. Others are light brownish with spots of darker, not showing any trace of the longitudinal stripes; the head variously marked with irregular vermiculations. Here described from specimens from Point Reyes, California. The one from which the meas-

urements are taken is a little over 4 inches in length. Specimens vary greatly in depth of body. Length 10 inches. Northern Pacific, Alaska to Monterey.* Common; specimens are recorded from Puget Sound and from Tongass, St. Paul, Kadiak, Unalaska, and other localities in Alaska. The specimen from the northernmost locality examined by us is from Bristol Bay in 16 fathoms (Coll. *Albatross*). The species is not rare about San Francisco and Monterey, where the striped form is almost exclusively found. (*pulchellus*, pretty.)

Liparis pulchellus, AYRES, Proc. Cal. Ac. Nat. Sci., I, 1855, 23, San Francisco; GÜNTHER, Cat., III, 164, 1861; STEINDACHNER, Ichth. Beiträge, III, 53, 1875; JORDAN & GILBERT, Synopsis, 741, 1883; GARMAN, Discoboli, 67, pls. 4, 5, and 8, 1892.
Cyclogaster pulchellus, GIRARD, Pac. R. R. Rept., X, Fishes, 132, 1858.

Subgenus ACTINOCHIR, Gill.

2458. LIPARIS MAJOR (Gill).

Head 4 in length. D. 45 to 48; A. 38 to 40; P. 34 to 37; caudal 12 to 14; vertebræ 52 (10+42); cæca 26; branchiostegals 6. Body elongate, much compressed and tapering posteriorly, rather thin behind the abdomen, slender at the base of the caudal, broad and high between the pectorals. Head high and broad, prominent at nape, length a little more than depth, forehead depressed; snout broad, blunt, rounded, rather deep, as long as eye; mouth anterior, broad, maxillary subtending the eye, jaws about equal; a broad interruption in lower lip in middle of chin; teeth small, in pavement, tricuspid in younger specimens, simple in old. Some of those from which this description is drawn show the 3 pronged teeth in the outer rows, and an approach to the simple in the inner. A series of 5 or 6 pores on each side, just above upper lip, from snout to postorbital region; another of 6 or 7 pores from chin toward upper angle of gill opening; posterior nostril reduced, pore-like, on interorbital space; anterior in front of eye, tubular; between and a little in front of the tubes a couple of large pores. Eye moderately large, in anterior half of head, lateral, about equal to snout, once in interorbital space, $1\frac{1}{4}$ times in disk, and $3\frac{1}{2}$ in head; disk small, little longer than wide, $1\frac{1}{4}$ times as long as eye, distant from mouth $1\frac{1}{3}$ times the length, which is equal to $\frac{1}{2}$ distance to anal fin; vent about midway from disk to first ray of anal; gill opening little wider than eye, $\frac{1}{3}$ of its extent in front of base of pectoral; 1 single and 3 double gills; pseudobranchiæ small; opercular spine rather broad; skin thin, loose, easily carried away, that of the males in breeding season roughened with small, spine-bearing papillæ; dorsal and anal continuous with the caudal, the union occupying nearly $\frac{1}{2}$ length of last, anal extending a little farther back than dorsal. Caudal rays less than $\frac{2}{3}$ as long as head, hinder margin of the fin slightly rounded, narrow; pectorals broad and rounded in the upper portion, reaching anal fin; in the lower portion fringed; several of the rays at sides of disk, much longer than those immediately above them, form a notch in the margin, a fold uniting the fins in front of disk; no apparent notch in dorsal; like the

* Our Monterey specimens have D. 48; A. 39; P. 36; C. 12. Head 5; depth 5.

anal, it rises gradually, and attains its greatest extent in the posterior third of its length. Professor Lütken gives the following as the formula: Dorsal 43 to 49; anal 36 to 40; pectoral 32 to 38; caudal 9 to 11. The specimens described here, and from which the numbers placed at the head of this description were taken, were furnished this Museum by him. From his large series he finds a greater range of variation in all cases except that of the caudal, where 12 to 14 rays appear on these examples. Color olivaceous to dark brown, darkest about the head and body. On close examination the skin is seen to be thickly puncticulate with brown, in cases forming cloudings, blotches, or transverse bands on the fins and hinder parts of the body. In life there is no doubt of the presence of tints of lilac, reddish, or yellowish. One of these specimens has 6 bands of brownish on the dorsal, and 4 on the anal. Our largest is 5 inches in length. (Garman.) Coasts of Greenland, east to the White Sea; not seen by us; the description from a specimen in the Museum Comp. Zool., sent by Dr. Lütken. (Eu.) (*Major*, larger.)

Cyclopterus liparis major, Fabricius, Fauna Grönland., 136, 1780, Greenland; Walbaum, Artedi Piscium, III, 489, 1792; after Fabricius; not a binomial name, *major* being simply an adjective.
Liparis tunicata, Kröyer, Naturh. Tidsskr., I, 236, 1862; not of Reinhardt.
Actinochir major, Gill, Proc. Ac. Nat. Sci. Phila. 1864, 193, Greenland.
Liparis fabricii, Lütken, Kara-Haves, Fiske, 146, 1887.
Liparis major, Jordan & Gilbert, Synopsis, 741, 1883; Garman, Discoboli, 72, 1892.

786. BATHYPHASMA, Gilbert.

Bathyphasma, Gilbert, Rept. U. S. Fish Comm. 1893 (1896), 447 (*ovigerum*).

Teeth long, slender, simple, without basal cusps. Disk large, its surface even, with evident lobes or horny papillæ under posterior part of the head; the anus not immediately behind it. Gill opening comparatively wide. Caudal long, well developed, not acuminate, containing 12 rays, the dorsal and anal confluent with its basal $\frac{2}{3}$; pectoral broad and continuous, as in *Liparis*, the lower rays exserted. Suborbital process strong. Vertebræ numerous. The typical species has much the shape and general appearance of *Liparis agassizii*. The character of the simple teeth, which this genus shares with *Paraliparis* and others, has been independently acquired. (Gilbert.) Skeleton soft, little ossified. Deep-sea Liparids of large size, the dentition unlike that of the other genera, and apparently not acquired by the same line of descent as the simple teeth of *Paraliparis*, which seems to be a degradation of the form seen in *Liparis*. (*βαθύς*, deep sea; *φάσμα*, apparition.)

2459. BATHYPHASMA OVIGERUM, Gilbert.

Head $3\frac{3}{5}$; depth $3\frac{3}{4}$; eye 7. Occipital region greatly elevated, the upper profile of head strongly decurved above the orbits, a line from occiput to end of premaxillary processes forming an angle of 45 degrees with axis of body. In front of tips of premaxillary processes the snout descends almost vertically. Posteriorly the body tapers uniformly and slowly, the width of base of tail equaling $\frac{1}{2}$ diameter of eye. Mouth large, horizontal,

not overpassed by the snout, the lower jaw slightly shorter than the upper, not distinctly included; maxillary reaching a vertical from posterior edge of orbit, its length 2⅓ in head, its width 1⅝ in head; upper lip complete, the fold of lower lip extending halfway from angle of mouth to symphysis. Bands of teeth very wide in the front of each jaw, becoming narrow laterally, where the series are few in number and nearly parallel with the jaw; anteriorly the series grow more and more oblique, until at front of jaw they are nearly transverse; teeth all simple and slender, without cusps, directed very obliquely backward, and movable, implanted so as to admit of still further depression; outer teeth in both jaws very short, the inner ones growing gradually longer and becoming acicular, 16 series of teeth on each side of lower jaw, 22 series on each side of upper jaw. Posterior nostril in a short wide tube. Eye large, the diameter of its exposed portion 2½ in total interorbital width. Cheek and temporal region swollen, the suborbital stay running in a notable depression between the two. Gill slit wide, extending downward to opposite upper pectoral rays, longer than snout, 2⅜ in head. Mucous pores minute, on sides of snout, mandible, and preopercle, none visible on top of head. Disk large, nearly round, its center slightly in advance of gill slit, its length 3⅓ in head, distance from its posterior margin to vent equaling ⅓ its own diameter. A small anal papilla. Pyloric cæca 19. Pectoral with 34 rays, not notched, the lower rays regularly diminishing in length to the fifth or sixth before the last, the next 2 or 3 abruptly lengthened and exserted; longest ray of upper lobe equaling snout and eye; longest ray of lower lobe equaling snout and ½ eye; dorsal fin beginning slightly behind upper axil of pectorals and with 43 rays, the longest equaling distance from tip of snout to front of pupil; origin of anal fin with ½ diameter of eye in front of middle of body, with 34 rays; caudal ray long and slender, with 12 rays, its basal ⅔ confluent with dorsal and anal, its length equal to that of pectoral fin. Skin thin, not conspicuously lax. Head, body, and fins white, inconspicuously mottled with light brown. A single specimen, 315 mm. in total length, from Station 3342, off Queen Charlotte Island, British Columbia, depth 1,588 fathoms. The type is a male, with well-developed testes, and contained in its mouth, when captured, a spherical mass of eggs apparently of the same species. The eggs measured 4½ mm. in diameter, and were well along in their development, the embryos distinctly visible through the very tough egg membranes. The general form of head and body can be made out, as well as the long continuous dorsal and anal fins running backward into the tail. It seems not improbable that the male fish protect the eggs in this manner until after hatching. (D. 43; A. 34; P. 34; C. 12.) (Gilbert.) (*ovum*, egg; *gero*, I bear.)

Bathyphasma ovigerum, Gilbert, Rept. U. S. Fish Comm. 1893 (1896), 448, off Queen Charlotte Island, at Albatross Station 3342, in 1,588 fathoms. (Type, No. 48622.)

787. CAREPROCTUS, Kröyer.

Careproctus, Kröyer, Naturh. Tidsskr., 1, 257, 1862 (*reinhardi*).
Caremitra, Jordan & Evermann, Check-List Fishes, 452, 1896 (*simus*).
Allochir, Jordan & Evermann, Check-List Fishes, 452, 1896 (*melanurus*).
Allurus, Jordan & Evermann, Check-List Fishes, 452, 1896 (*ectenes*).

Body more or less elongate, semitransparent, covered with thin, lax skin; mouth terminal; teeth simple in the adult, hooked; ventral disk very small, situated far forward, under front of eye, sometimes reduced to a mere point; vent a little behind it, far in front of anal fin; pectoral fin broad, notched, the lower lobe well developed. Vertebræ and fin rays numerous; nostrils with or without tube. Species of rather large size, found in deep or cold waters. These form a transition from *Actinochir* to the *Amitrinæ*, in which group the ventral disk is wholly lost.

Concerning *Careproctus*, Mr. Garman has the following remarks:

"This genus and *Liparis* are brought very close together by *Careproctus liparis major*. The simple form of tooth in the one case and the tricuspid form in the other were formerly considered sufficiently distinctive for separating these genera. Lütken, 1887, pointed out that the young of the mentioned species has the tricuspid teeth of *Liparis*, and that with age it acquires the simple teeth of *Careproctus*. It is possible that a similar change takes place in the other species we have brought together in the latter. Indeed, assuming derivation from species of *Liparis* of the shoal waters, it is just what we should expect in all these *Careproctí* of the deep sea.

"Besides the dentition, other features may be selected that in comparisons may serve to distinguish this genus from the preceding: The head is higher at the back; the body is deeper in front of the dorsal fin; the caudal region is longer, more slender and tapering; the vertebræ are more numerous; the dorsal and anal fins have more rays, and are completely fused with the caudal; the disk is reduced in size; the suborbital process is less developed, and apparently the intestine is shorter. Among these differences there is none that may not be a consequence of the modifying influences of great depths upon species of *Liparis* such as now exist along the coasts." (καρά, head; πρωκτός, anus.)

CAREMITRA (καρά, head; μίτρα, stomacher):

a. Body short and deep, with blunt head, the inferior mouth below the prominent snout; disk nearly as long as eye; pectoral rays 33. SIMUS, 2460.

aa. Body more or less elongate.

b. Body not excessively attenuate, its median depth much more than length of eye.

CAREPROCTUS:

c. Pectoral fin distinctly notched, its middle rays shorter than upper or lower.

d. Lower pectoral rays extremely long, reaching front of anal; disk moderate, about as large as eye. COLLETTI, 2461.

dd. Lower pectoral rays not reaching anal.

e. Mouth horizontal or moderately oblique, not nearly vertical.

f. Disk nearly or quite as long as eye, not excessively reduced.

g. Disk about ½ longer than eye; anterior nostril with a tube; gill slit reduced; mouth oblique; D. 53; P. 34. PHASMA, 2462.

gg. Disk not larger than eye, which is about 4 in head; anterior nostril (so far as known) without tube.

h. Depth 4 in length; mouth large; D. 52; A. 47. SPECTRUM, 2463.

hh. Depth 5½ to 6.

i. Pectoral rays 32; D. 54; A. 45. REINHARDI, 2464.

ii. Pectoral rays 38; D. about 48; A. about 48. RANULA, 2465.

ff. Disk very minute. OSTENTUM, 2466.

ee. Mouth nearly vertical; body elongate. D. 50; A. 45; C. 6; P. 30. GELATINOSUS, 2467.

ALLOCHIR (ἄλλος, other; χείρ, hand):

cc. Pectoral fins not notched, the middle rays not shorter than upper or lower; gill openings reduced; depth 4¼ in length. D. 58; A. 50; P. 30. MELANURUS, 2468.

ALLURUS (ἄλλος, other; οὐρά, tail):

bb. Body excessively elongate, its depth at the middle not more than diameter of eye; disk smaller than eye; gill openings reduced; snout shovel-shaped. D. 51. ECTENES, 2469.

Subgenus CAREMITRA, Jordan & Evermann.

2460. CAREPROCTUS SIMUS, Gilbert.

A species with a very heavy head and body, an inferior transverse mouth, overlapped by the thick, rounded snout. Appearance much that of *Rhinoliparis barbulifer,* but the anterior parts much heavier, nape more elevated, snout shorter and blunter, without barbels, extending beyond front of eyes for a distance (measured axially) equal to ⅔ diameter of orbit, and projecting beyond mouth for ½ that distance. Mouth transverse, its width nearly twice the distance from symphysis of lower jaw to angle of mouth, the latter reaching a vertical midway between front of eye and pupil. Teeth indistinctly tricuspid, in narrow bands, 11 or 12 series in each half of lower jaw, 8 or 9 in the upper. Nostrils without tube. Eye large, 3 in head, equaling snout, 1½ in total interorbital width. Gill slit moderate, ⅔ of it opposite upper pectoral rays, its width ⅔ diameter of eye. Opercle prolonged into an acute lobe overlapping middle of slit. Disk rather small, round, under posterior part of eye, its diameter ⅓ that of eye. Vent immediately behind it, equidistant from front of anal and angle of mouth. Pectorals inserted high, the upper edge on a level with the middle of eye, continuous, the median rays greatly shortened, the lower ones again longer, with exserted free tips; upper lobe extending beyond front of anal, and equaling length of head behind snout, lower rays ⅔ length of upper ones; P. 33; origin of dorsal fin just behind axil of pectorals, continuous posteriorly with the caudal fin, which is very narrow and not distinct; origin of anal at end of first third of length. Whitish or bright brown, dusky posteriorly on body and fins; mouth, gill cavity, and peritoneum white. One specimen, 3½ inches long, from Bering Sea north of Unalaska Island. (Gilbert.) (*simus,* σιμός, snubnosed.)

Careproctus simus, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 444, Bering Sea north of Unalaska, at Albatross Station 3331, in 350 fathoms.

Subgenus CAREPROCTUS, Kröyer.

2461. CAREPROCTUS COLLETTI, Gilbert.

Head 5, blunt and heavy, with subvertical cheeks and bluntly rounded snout, the latter not projecting beyond the mouth; greatest depth (at occiput) 6; eye equals length of snout, 3¾ in head, 1⅔ in interorbital width; nape not elevated; mouth terminal at lower side of snout, slightly oblique,

its lateral cleft about ⅔ its width; lower jaw shutting within the upper; angle of mouth reaching a vertical midway between front of eye and front of pupil; teeth lanceolate, acute, without trace of basal cusps; gill opening a narrow slit, entirely above base of pectorals, its length ¾ diameter of eye; opercle produced posteriorly into a rounded lobe, which overlaps the gill opening; disk oblong, of moderate size, placed under the posterior part of head behind the eyes, forming, in alcoholic specimens, a very deep cup-shaped depression with incurved edges; diameter of disk about equaling that of eye; disk separated from vent by ½ its diameter; upper pectoral lobe reaching origin of anal fin, the rays of lower lobe elongate, extensively free, longer than upper lobe, and reaching to or nearly to front of anal; intermediate rays not as short as in *C. reinhardi*, hardly forming a separate division of the fin, the rays being gradually and uniformly shortened from above downward to origin of lower lobe; pectoral rays 29; dorsal beginning immediately behind the head; distance from tip of snout to origin of anal 3⅔ in length; series of conspicuous mucous pores on head, as in *C. reinhardi*. Color in spirits, dusky, the tip of snout, under side of head, opercles, abdomen, and posterior portion of vertical fins black; inside of mouth and gill cavity dusky; peritoneum black. Closely related to *C. reinhardi*, from which it differs principally in the elongation and exsertion of the lower pectoral rays. These extend in all specimens to or nearly to the origin of the anal fin, and are always free for the greater part of their length. Length 3½ inches. North Pacific, south of the Alaskan Peninsula. ("Named in honor of Prof. Robert Collett, the distinguished author of the Fishes of the Norwegian North Atlantic Expedition.")

Careproctus colletti, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 442, south of the Alaska Peninsula, at Albatross Station 3338, in 625 fathoms. (Type, No. 48698.)

2462. CAREPROCTUS PHASMA, Gilbert.

Eye 4 in head, 2 in total interorbital width. D. 53; P. 34. Head broad and flat above, subquadrate, with nearly vertical cheeks. Snout very obtuse, broadly rounded, much blunter than in *C. spectrum*, very slightly overlapping the mouth; width of snout 1⅓ in length of head; mouth very broad, somewhat oblique, reaching vertical slightly behind front of eye, its width more than twice the amount of lateral cleft taken axially. Teeth minute, acute, in a moderate band in each jaw, arranged in oblique series within the band. Nostrils opening in a short but conspicuous tube (the tube absent and the pore smaller in *C. spectrum*). Mucous pores small. Gill slit short, slightly less than diameter of orbit, its inferior end attached to base of upper pectoral ray, overlapped for almost its entire extent by the broadly rounded opercular flap. Sucking disk comparatively large, much larger than in *C. spectrum*, 1⅓ times diameter of eye, 3⅓ in head, very nearly round, the transverse diameter equaling or slightly exceeding the longitudinal diameter. Vent immediately behind edge of sucking disk; anal papilla slender, ½ as long as diameter of eye. Pectoral fin very broad, barely reaching front of anal, the lower rays equaling the upper,

extensively free at tip; lower lobe of pectoral containing 8 or 9 rays; dorsal beginning behind gill opening at a distance equaling diameter of eye. Skin exceedingly soft, thick, and lax, in alcoholic specimens forming folds on head and body and concealing the rays of the fins. Color uniform white in spirits. Closely related to *Careproctus spectrum*, Bean, from the same region, differing in the much larger sucking disk and the narrower gill slit, the latter confined to area above base of pectorals, its anterior margin formed of the broadly and evenly rounded opercular lobe. Length $3\frac{1}{2}$ inches. Bristol Bay, Alaska, in deep water. (*φάσμα*, a spectre.)

Careproctus phasma, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 443, Bristol Bay, Alaska, at Albatross Stations 3254 and 3256, in 46 and 49 fathoms.

2463. CAREPROCTUS SPECTRUM, Bean.

Head 4; depth 4; eye 3. D. 52; A. 47; ventral disk small, $\frac{2}{3}$ as long as eye. Mouth large, the maxilla extending to about below the middle of eye. Pectoral reaching to above anal origin; dorsal beginning over the axil of pectoral. Color along back of some examples light brown, elsewhere uniformly pale. Length of type $3\frac{9}{10}$ inches. Twenty-six specimens were obtained, July 31, 1888, between Unga and Nagai islands. (Bean.) Garman refers this species to the synonymy of *Careproctus gelatinosus*, which is probably not correct. (*spectrum*, spectre.)

Careproctus spectrum, BEAN, Proc. U. S. Nat. Mus. 1890, 40, between Unga and Nagai islands, 55° 10′ N., 160° 18′ W., at Albatross Station 2848, in 110 fathoms.

2464. CAREPROCTUS REINHARDI (Kröyer).

Head $4\frac{1}{2}$; depth $5\frac{1}{2}$; eye 5 to 6. D. 54; A. 45; C. 12; P. 32. Body semitransparent, covered with viscid skin. Head short and globular. Caudal fin very slender, joined to the dorsal and anal; mouth nearly horizontal; lower jaw scarcely projecting. Ventral disk little larger than eye. Pectoral deeply emarginate, nearly as long as head, reaching anal, the lower rays exserted, appearing as a series of short twisted filaments. Distance from ventral disk to vent not more than diameter of eye; head with conspicuous mucous pores. Color pale reddish gray, unmarked. (Collett.)

Mr. Garman adds to this: Body translucent, jelly-like; skin lax, viscid; teeth simple; ventral disk very small, situated far forward, almost hidden by the anterior portions of the pectorals; vent near the disk, about midway from snout to anal; tail long, slender, tapering to a point; caudal continuous with dorsal and anal; fin rays soft, slender; pectorals semicircular, below the head, beginning far in front and close together near the symphysis of the lower jaw, first 8 or 10 rays projecting, larger median portion with short ones, upper elongate; eyes comparatively large, more than $\frac{1}{4}$ as long as the head, which latter is $\frac{1}{4}$ to $\frac{1}{5}$ of the total length. Only 1 pair of nostrils was detected, and they were not prolonged in tubes. Color pale reddish gray, or whitish; no bands or marks. An Arctic, deep-sea form, descending to 700 fathoms. Known from Greenland, Jan Mayen, and Beeren Island. (Named for Professor J. Reinhardt, of the University of Copenhagen, who wrote largely on the fauna of Greenland.)

Liparis gelatinosus, REINHARDT, Oversigt, etc., 1844, 77; not of PALLAS.
Liparis (Careproctus) reinhardi, Kröyer, Naturh. Tidskr., I, 252, 1862, Greenland.
Careproctus reinhardi, COLLETT, Norske Nord-Havs Exped., 57, 1880; JORDAN & GILBERT, Synopsis, 957, 1883; GARMAN, Discoboli, 78, 1892.

2465. CAREPROCTUS RANULA (Goode & Bean).

Head 4; depth 6. D. 48±; A. 48±; pectoral 15 + 12 or 13. Body thick, subcylindrical anteriorly, rapidly tapering to the tail; skin thick, lax. Head swollen at the nape; width little greater than depth, twice that of disk; length of disk 2½ interorbital width. Snout broad, about ¼ as long as head; mouth cleft not reaching a vertical from orbit; upper jaw the longer; eye lateral, not interfering with upper profile, about ¼ of head; nostril near the eye; from snout to first dorsal ray ⅓, and from snout to first anal ray about ⅔ of length of body; origin of anal below eighth ray of dorsal; disk slightly longer than its distance from the snout, or than its width. Color uniform, whitish. Off Halifax Harbor, in 52 fathoms. (Goode and Bean.) Garman refers this species to the synonymy of *Careproctus reinhardi*, which may be correct. (*ranula*, tadpole, diminutive of *rana*, frog.)

Liparis ranula, GOODE & BEAN, Proc. U. S. Nat. Mus. 1879, 46, Halifax; JORDAN & GILBERT, Synopsis, 742, 1883.

2466. CAREPROCTUS OSTENTUM, Gilbert.

The snout is longer and more pointed than in *Careproctus phasma* or *Careproctus spectrum*, its width little greater than its length, 2¼ in head. Mouth with very distinct lateral cleft, its width less than its length. Maxillary reaching a vertical line crossing orbit behind pupil, 1$^{9}_{10}$ in head; teeth minute, in narrow bands, indistinctly tricuspid; eye equaling length of snout, 1½ in total width of interorbital area. The true bony interorbital width is much narrower than this. In the specimen examined the epidermis of the head is largely lost, and the width of the gill slip can not be determined. A short nostril tube. Fins as in *C. spectrum* and *C. phasma*. Skin loose, thinner than in *C. phasma*. Color white or slightly brownish, minutely punctulate with black. Differing from *C. spectrum* in the minute size of the sucking disk, which is reduced to a mere rudiment entirely concealed by the anterior (lower) lobes of the pectoral fins, about one millimeter in diameter in a specimen 78 mm. long. Bering Sea, north of Unalaska Island, in deep water. (*ostentum*, a spectre.)

Careproctus ostentum, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 444, Bering Sea, north of Unalaska Island, at Albatross Stations 3324 and 3331, in 109 and 350 fathoms. (Type, No. 48619.)

2467. CAREPROCTUS GELATINOSUS (Pallas).

D. more than 50; A. about 45; C. 6; cæca 48; vertebræ 64. Vertical fins contiguous. Ventral disk small, supported by a bilamellated cartilage, which descends from the throat. Body oblong, compressed, thicker toward the head and thinner toward the tail, semitransparent,

exceedingly soft, like jelly. Head thick, somewhat depressed, flat on the vertex, and convex between the orbits. Eyes directed upward; nostrils in middle between margin of mouth and orbit, with a small tubular appendage; a series of pores along upper lip toward gill opening. Lips thick; cleft of mouth directed upward, as in *Uranoscopus*, but with the lower jaw less prominent; jaws, palate, and pharynx rough with teeth; 4 gills on each side; 7 branchiostegals. Gill opening reduced to a slit above base of pectorals. Pectorals large, very broad, commencing at the throat, the inferior part of their bases being parallel, ascending to the gill opening, composed of about 30 rays, the inferior very short and not connected by a membrane. Dorsal and anal fins continuous with the caudal, commencing a little behind middle of body, and composed of feeble rays; jugular disk situated between the pectorals in front of the vent, very small, soft, supported by a bilamellated cartilage which descends from the throat; vent between pectorals; caudal small, 6-rayed. Rose-colored; vertical fins violet; gill cavity black. Steller adds the following anatomical details: Ovarium orbicular, containing eggs of the size of a pea. Liver large, divided into 4 lobes, of a whitish color; gall bladder absent; spleen triangular, brown; stomach 3 times as wide as the œsophagus; 48 pyloric appendages, 2 inches long and as thick as the wing feather of a pigeon, in a specimen 18 inches long; remainder of intestinal tract about as long as the fish; urine bladder the size of a hazelnut; kidneys united into one cuneiform mass, commencing near the gills and extending to anterior portion of ovarium; ureter single, very wide, flexous, becoming narrower before its insertion into the bladder. The *nervi optici* and *olfactorii* have one common ganglion, from which, first, the former arise, emitting the latter from the angle formed by the nerves and the ganglion. Skeleton very slightly osseous. Coast of Kamchatka. Specimens (1 of 18 inches long) were found by Steller in Peter and Paul Harbor, Kamchatka. (Pallas.) The species has not been recognized by recent writers. Not seen by us. (*gelatinosus*, jelly-like.)

Cyclopterus gelatinosus, PALLAS, Spicilegia, VII, 19, 1769, Peter and Paul Harbor, Petropaulski Kamchatka. (Coll. Steller.)

Liparis gelatinosus, GÜNTHER, Cat., III, 163, 1861.

Careproctus gelatinosus, JORDAN & GILBERT, Synopsis, 740, 1883; GARMAN, Discoboli, in Memoir M. C. Z., No. 2, XIV, 76, 1892.

Subgenus ALLOCHIR, Jordan & Evermann.

2468. CAREPROCTUS MELANURUS, Gilbert.

Head 4½ to 4⅔ in length; depth 4⅓ to 4½. D. about 58; A. 50; P. 30. Head heavy, with vertical cheeks, short bluntly rounded snout, and very broad, flat interorbital space. Width of interorbital space slightly more than ½ length of head. Mouth very broad, horizontal, with short lateral cleft, the maxillary reaching to below middle of eye; width of mouth equaling length of snout and eye. Teeth short and strong, in narrow bands in both jaws; except at symphysis, the teeth of the bands are arranged in regular series running from inner edge of jaw forward and

sideward. Tongue and roof of mouth papillose and toothless. Gill rakers tubercular. Nostrils single, in a very short wide tube, which is nearer eye than cleft of mouth. Gill opening oblique, entirely above base of pectorals, the length of slit equaling snout and ½ eye. A series of 6 pores along under side of mandible and interopercle; none on preopercle; 3 above premaxillary, and 2 pairs on snout; no pores along side of body. Sucking disk extremely small, little more than ½ diameter of orbit, its center and the middle of pupil equidistant from tip of snout; disk separated from vent by a trifle less than its own diameter. Pectorals forming a continuous lamella, following the margin of gill flap and lower jaw, the 2 fins becoming closely approximated in front of sucking disk; fin not notched, the rays decreasing regularly forward, 8 to 10 of the anterior rays exserted and elongated, the anterior 4 or 5 becoming again shorter and thicker, and wholly free from the membrane; pectorals scarcely reaching front of anal, ⅔ to ¾ length of head; dorsal beginning over gill slit, the distance from origin of anal to snout, 2½ in length; fins all enveloped in thin lax skin, so that it is impossible to count rays accurately without dissection; posterior dorsal and anal rays well overlapping base of caudal, their membranes joining middle of caudal rays. Color light rose-red on body and fins, abdomen usually dusky; inner face of pectorals, caudal, posterior portions of dorsal and anal, and inside of mouth and gill openings black; peritoneum silvery, with more or less black specking, sometimes nearly black. Several specimens, the longest about 6 inches long, from *Albatross* Stations 2840, 2891, 2892, 2925, and 3076, in depths from 178 to 339 fathoms off the coast of California and Oregon. (Gilbert.) (*μέλας* black; *οὐρά* tail.)

Careproctus melanurus, Gilbert, Proc. U. S. Nat. Mus., 1891, 560, off coast of California and Oregon, at Albatross Stations 2840, 2891, 2892, 2925, and 3076, in 178 to 339 fathoms. (Type in U. S. Nat. Mus.)

Subgenus ALLURUS, Jordan & Evermann.

2469. CAREPROCTUS ECTENES, Gilbert.

Head 5½ in total length, its depth ⅔, and its width ¾ its length. An extremely elongate form, depressed but narrow anteriorly, the head as seen from above appearing shovel-shaped, with truncate snout; nape not elevated and the cheeks not gibbous; width anteriorly everywhere exceeding depth; mouth inferior and transverse, overlapped by the short, depressed snout for a distance equaling diameter of pupil; width of mouth nearly twice distance from symphysis of lower jaw to angle of mouth, the latter reaching vertical from snout to front of pupil. Teeth small, weakly tricuspid, in narrow bands, lower jaw containing 10 series in each half, the upper 11. Eye large, 1¼ in total interorbital width, equaling length of snout, 3⅓ in head. Nostrils with a very short tubular rim. Mucous pores large; texture of head and body firmer than in most deep-sea Liparids. Gill opening reduced to a narrow slit entirely above the pectorals, its width equaling ½ diameter of orbit. Opercle produced into a narrow spinous process, forming, with its membranaceous flap, a

quadrate projection over middle slit. Disk small, under the opercles and posterior part of cheeks, round in shape, slightly smaller than eye, its diameter ¼ length of head; vent separated from disk by about ⅜ diameter of disk. Distance from vent to front of anal ½ its distance from tip of snout. Body extremely slender, its depth at middle of total length equaling diameter of eye, at base of caudal equaling ⅜ diameter of pupil. Dorsal beginning slightly behind axil of pectorals; distance from origin of anal to tip of snout ½ its distance from base of caudal; upper lobe of pectoral extending slightly past front of anal, the lower lobe to opposite the vent; rays of lower lobe partly free; dorsal with about 51 rays, pectorals with 29. Color nearly uniform dusky brownish, lighter on snout, belly, and under side of head; mouth, gill cavity and peritoneum white. Length 3 inches. Bering Sea north of Unalaska Island, in deep water. (*ἐκτενής*, drawn out.)

Careproctus ectenes, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 442, Bering Sea north of Unalaska Island, at Albatross Station 3331, in 350 fathoms.

788. GYRINICHTHYS, Gilbert.

Gyrinichthys, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 444 (*minytremus*).

Teeth simple, not tricuspid. Body attenuate posteriorly, as in *Paraliparis*, the tail scarcely distinct. Disk small, under the posterior part of the head, the vent immediately behind it. Pectorals without anterior lobe, the rays progressively shortened, none of them exserted or with free tips. Gill openings reduced to a minute round pore, well above base of pectorals. One species known. (*γυρῖνος*, tadpole; *ἰχθύς*, fish.)

2470. GYRINICHTHYS MINYTREMUS, Gilbert.

Body in the type greatly distended with eggs, the original shape difficult to ascertain, tapering posteriorly into an extremely slender, compressed tail. Head slender, not greatly depressed, the depth and width about equal. Nuchal region not elevated, the profile rising but little behind the snout, which is blunt, with almost vertical profile. Mouth very small, almost entirely transverse at end of snout, with but little lateral cleft, the angle of mouth scarcely reaching vertical from nostril; jaws even, the snout very slightly protruding beyond premaxillaries. Teeth slender, acicular, without cusps, the inner teeth longest; bands of teeth very narrow, with but 5 or 6 oblique series in each half of each jaw; the teeth directed backward, but scarcely depressible. Nostrils without tube. Eye large, 1½ in total interorbital width, equaling length of snout. Gill opening a small pore, scarcely larger than nostril, well separated from upper margin of pectoral. Disk of moderate size, round, its diameter equaling ½ length of head; vent close behind disk, separated from it by ⅙ its distance from front of anal fin. Pectoral small, its upper edge on a level with lower margin of eye, the 2 fins converging under the throat, the anterior rays progressively shortened, all included within the membrane; dorsal without any detached anterior portion, beginning well behind the head, at a distance from gill opening equaling ½ length of

head; like the anal, it is continuous with the very narrow caudal fin, there being no notch or evident separation between them; distance from tip of snout to front of anal $1\frac{2}{3}$ in distance of latter from base of caudal; dorsal with about 45 rays; caudal with a very narrow base, containing, apparently, 14 rays, its length equaling that of snout and eye. Color light brownish, everywhere dusted with minute black specks, which are largest on back and tail; lining of mouth and gill cavity and peritoneum white. Eggs large, visible through the abdominal wall, about $3\frac{1}{2}$ mm. in diameter. Length 3 inches. Bering Sea, north of Unalaska Island, in 350 fathoms; only the type known. (*μικρός*, reduced; *τρῆμα*, aperture.)

Gyrinichthys minytremus, GILBERT, Rept. U. S. Comm. 1893 (1896), 444, Bering Sea, north of Unalaska Island, at Albatross Station 3337, in 350 fathoms. (Type, No. 48617.)

789. AMITRA, Goode.

Amitra, GOODE, Proc. U. S. Nat. Mus. 1880, 478 (*liparina*).

Monomitra, GOODE, Proc. U. S. Nat. Mus. 1883, 109 (*liparina*); name changed on account of the prior use of the name *Amitrus*.

Body elongate, attenuate backward, covered with lax, smooth, slimy skin, which is separated from the body and fins by a filmy mucous intertissue. Head small, thick, convex between eyes; snout convex, protruding; mouth small, horizontal, the lower jaw included; teeth small; eyes lateral; gill openings restricted to small slits above the base of the pectorals, covered by the very small opercle; pseudobranchiæ present, very small, no barbels; dorsal and anal fins many-rayed; dorsal continuous, almost concealed by the skin, the spines scarcely different from the soft rays; anal similar to dorsal; both connected with the caudal, which is pointed; no trace of ventral fins or sucking disk; pectoral fin very broad, procurrent, its lower rays inserted nearly under the eye, the fin emarginate; vent well behind head. This genus is of very doubtful value, the presence of pseudobranchiæ alone separating it from *Paraliparis*.* (*ἀ*, without; *μίτρα*, stomacher.)

2471. AMITRA LIPARINA, Goode.

D. 67; A. 54; pectoral 23; caudal 6. Body elongate, compressed posteriorly, very thin at the tail, covered by a thick, lax, slimy skin. Head small, thick, convex between the eyes, length $4\frac{1}{3}$ times in width, $\frac{3}{20}$

*The following words of Professor Putnam, written before the discovery of *Psychrolutes* and *Liparididæ* without ventral disk, are of interest in this connection: "I should put the family of *Gobiesocidæ* far away, at least a suborder off, from the *Cyclopteridæ* and *Liparididæ*, which are far more closely united to the true *Cottidæ*, represented by *Cottus* and *Hemitripterus*, than to either the *Gobiesocidæ* proper or to the Gobies and Blennies. In fact, *Liparis* has as close affinities, as shown by its skeleton, with *Cottus* and *Hemitripterus* as with *Cyclopterus;* and we have in the three groups, represented by *Cottus*, *Liparis*, and *Cyclopterus*, well-marked families of the same suborder. The only character by which the *Cyclopteridæ* and *Liparididæ* are closely united consists in the peculiar formation of the ventral disk by the union of the ventral fins; but as this structure is simply brought about by the modification of the rays in a manner common to the several genera, and not by any marked anatomical difference in the structure of the same fins in *Cottus*, I can only look upon it as a generic character common to the known representatives of both families of *Cyclopteridæ* and *Liparididæ;* and the discovery of a representative of either family with ventral fins of the ordinary form would not necessitate the establishment of a family for its reception, as in that case we should simply consider the structure as of generic value." (Putnam, Proc. Am. Ass. Adv. Sci. 1873, 337.)

of the total length without caudal; snout convex, protruding; mouth under the snout and far back from its tip; eyes lateral, ½ as wide as the interorbital area; nostrils in front of the eye. Operculum very small, strap-shaped. Gills 3½. Pseudobranchiæ present. Teeth weak, paved. Dorsal beginning over end of pectoral, the rays and outline, as of the anal, hardly visible through the skin; anal beginning below eighth to tenth dorsal rays; dorsal and anal rays closely connected with those of caudal, which are larger and extend in a pencil-like point; ventrals absent; pectoral broad, lower base almost below posterior margin of orbit, lowest 6 rays prolonged. The jugular disk can not be found. Color yellowish white, dusky toward the tail and blackish upon the anterior part of the head; abdominal cavity showing black through the skin. Length 5 inches. Atlantic Ocean, off Rhode Island, in 487 fathoms. (Goode.) (*liparina*, like *Liparis*.)

Amitra liparina, GOODE, Proc. U. S. Nat. Mus. 1880, 487, off Rhode Island (Type, No. 26184. Coll. *Fish Hawk*); JORDAN & GILBERT, Synopsis, 739, 1883.
Monomitra liparina, GOODE, Proc. U. S. Nat. Mus. 1883, 109.
Paraliparis liparinus, GÜNTHER, *Challenger*, Report, XXII, Deep-sea Fishes, 68, 1887; GARMAN, Discoboli, 52, 1892; GOODE & BEAN, Oceanic Ichthyology, 278, 1896.

790. PARALIPARIS, Collett.

Paraliparis, COLLETT, Vid. Selsk. Forh. Christiana, 14, 32, 1878 (*bathybii*).
Hilgendorfia, GOODE & BEAN, Oceanic Ichthyology, 280, 1896 (*membranacea*).
Amitrichthys, JORDAN & EVERMANN, Check-List, 453, 1896 (*cephalus*).

This genus differs from *Careproctus* chiefly in the total absence of the ventral disk. The teeth, so far as described, are simple, as in the older individuals of *Careproctus*. Body long, slender in the caudal portion, covered with a loose, thin skin. Vent remote from the anal. Pectorals deeply notched or divided. Dorsal and anal long, and confluent with the caudal. Vertebræ numerous. Gill rakers 3½. Pseudobranchiæ entirely wanting. Deep-sea fishes of the Arctic. The species are all of very recent discovery. They vary considerably among themselves, and are perhaps divisible into 2 or 3 genera. Among such degenerate forms, the characters usually taken as generic lose their value. (παρά, near; *Liparis*.)

a. Insertion of pectoral low, its upper ray below the level of the eye.
PARALIPARIS:
b. Gill openings rather large, the gill slit extending downward to opposite middle of pectoral; head large, 5 in length; depth 6. D. 60; A. 54; pectoral divided to the base, its lobes wholly separate, its rays 24; lower jaw included. HOLOMELAS, 2472.

AMITRICHTHYS (*a*, without; μίτρα, stomacher; ἰχθύς, fish.):
bb. Gill openings restricted to the region above pectorals.
c. Head very large, 4½ in length, its depth at occiput 4⅔ in length, twice depth of body at front of anal; lower jaw projecting; pectoral rays 14, the fin simply notched. CEPHALUS, 2473.
cc. Head rather small, 5 to 6¼ in length.
d. Pectoral fin divided to the base, its lobes separate; lower jaw included; head 5⅔ in length; depth 4⅔. ROSACEUS, 2474.

dd. Pectoral fin more or less deeply notched, not divided to base.

e. Lower jaw projecting. P. 16; D. 57; A. 43; head 5⅔ in length; depth 5⅔. MENTO, 2475.

ee. Lower jaw included.

f. Pectoral rays 31; head 6¼ in length; depth 6. D. 66; A. 55. COPEI, 2476.

ff. Pectoral rays 30; head 5 in length; depth 6¼. D. 56; A. 46. DACTYLOSUS, 2477.

HILGENDORFIA:*

aa. Insertion of pectoral very high, its upper rays above pupil; lower jaw included; pectoral fin notched; gill opening small; head 5; depth 5. D. 65; A. 60; P. 25. ULOCHIR, 2478.

Subgenus **PARALIPARIS**, Collett.

2472. PARALIPARIS HOLOMELAS, Gilbert.

Head about 5; depth about 6; eye 3⅔. D. 58 to 61; A. 54. Head very large and heavy, with very broadly rounded snout, and much swollen occipital and nuchal regions, the highest point over upper opercular angle, from which point the profile descends rapidly backward, though much less so than in *P. cephalus*. Snout very blunt, evenly rounded, very slightly projecting beyond the mouth, its width equaling length of snout and eye, ½ length of head; eye ⅔ interorbital width; mouth large, horizontal, quite at lower side of snout, entirely below the eye; maxillary reaching a vertical slightly behind posterior margin of orbit, 1⅔ in head; teeth acute, arranged in oblique series in each jaw, forming a very narrow band in mandible, a broader band in upper jaw; very large mucous slits on head, 5 forming a series from tip of snout below eye and across cheek, 6 along mandible and preopercle; gill slit wide, extending from above opercular flap nearly to middle of base of pectorals, its length 2¼ in that of head; opercle forming posteriorly a narrow angular flap, projecting above base of pectorals; no pseudobranchiæ; vent below opercular flap, or somewhat in advance of that point, nearer to base of pectorals anteriorly than to first anal ray; pectorals inserted very low, the base of upper lobe vertical, the base of notch and lower lobe horizontal, the upper end of base below the level of the eye; pectorals with 2 wholly distinct lobes, the interspace without free membranaceous margin, the skin of abdomen directly continuous at this point with that of shoulder girdle. On dissecting off the integument, however, the interspace between the lobes is seen to be provided with 2 or 3 short, widely spaced rays, as in all other species examined by us; upper lobe long, reaching beyond front of anal, the rays close set, 18 in number, included in the membranes to their tips; lower lobe consisting of 5 or 6 slender, almost filamentous rays, the longest reaching front of anal, all of them free to the base, without connecting membrane; anterior (lower) ends of shoulder girdle approximate, the rays separated by a distance equaling ½ diameter of pupil; dorsal beginning above base of pectorals. Color uniformly black, including fins and lining of mouth and gill cavity. Closely allied to *P.*

* Named for Dr. Franz Hilgendorf, ichthyologist of the University of Berlin.

cephalus and *P. mento*, differing in its uniform coloration, its more inferiorly placed horizontal mouth, and the distinctly included lower jaw. Length 4 inches. Bering Sea, north of Unalaska Island, in deep water; only 2 specimens known. (Gilbert.) (ὅλος, entirely; μέλας, black.)

Paraliparis holomelas, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 441, Bering Sea, north of Unalaska Island, at Albatross Stations 3308 and 3332, in 406 and 1,625 fathoms. (Type, No. 48637.)

Subgenus AMITRICHTHYS, Jordan & Evermann.

2473. PARALIPARIS CEPHALUS,* Gilbert.

Head very large, high, and compressed, the upper profile descending in a strong convex curve behind the occiput; body tapering posteriorly to a very narrow thread-like tail. Sides of head vertical or inclining inwards below. Height of head twice height of body opposite origin of anal. Interorbital space transversely rounded, its width slightly less than length of snout and eye. Eye of moderate size, 4 in head, without vertical range. Mouth oblique, the premaxillaries but little below lower margin of eye; lower jaw longer than upper, the tip protruding; maxillary reaching vertical from posterior margin of eye, slightly more than length of head. Gill slits narrow, confined to a region above the base of the pectorals; membrane connecting branchiostegal rays with shoulder girdle very delicate and easily ruptured; vent distant less than a diameter of the orbit from this point, being below the front margin of the preopercle. Teeth in narrow bands in both jaws, the teeth of the bands arranged in oblique series running outward and forward; palate toothless. Head 4½ in length, greatest depth (at occiput) 4¾. Pectoral small, with about 14 rays, the upper ones closest and forming a projecting lobe, which extends backwards to beyond origin of anal, the succeeding rays shortened and wide set, and with the tips free from the membrane, the fin somewhat mutilated, but apparently notched, not, however, to the base, the median portion having wide-set rays; upper margin of pectorals on a level with tip of lower jaw; dorsal beginning slightly behind head; first ray of anal under eighth of dorsal. Color light reddish, the abdomen blue black; inside of mouth and gill opening white. In the smallest specimens, 2 inches long, the color is dusky on head and body, and on inside of mouth. Longest specimen 3½ inches. This species differs from *Paraliparis (Hilgendorfia) membranaceus* in the structure of the pectoral fins and their much fewer rays, in the position of the vent (below the posterior margin of the orbit in *membranaceus*), the different outline of head, and in the larger, more oblique mouth. In none of the specimens of *cephalus* are there fin folds extending forward from front of dorsal and anal, as described in *membranaceus*. Alaska to California, in deep water;

* Dr. Gilbert later records several more or less mutilated specimens afterwards taken north of Unalaska Island, and near Point Reyes, California, in depths of 351 and 455 fathoms. In this species the pectoral is inserted very low, its upper edge entirely below the eye. The lower jaw shuts within the upper, but the symphysis protrudes. The disproportion in size of head and body is more exaggerated in the young than in adults. In very small examples the head is almost spherical, diminishing abruptly to the very slender tail. No pseudobranchiæ. (Gilbert.)

several specimens in 284 to 685 fathoms off the coast of California and Oregon, and north of Unalaska. (*κεφαλή*, head.)

Paraliparis cephalus, GILBERT, Proc. U. S. Nat. Mus. 1891, 561, off California and Oregon, at Albatross Stations 2839, 2892, 3070, and 3071, in 284 to 685 fathoms.

2474. PARALIPARIS ROSACEUS, Gilbert.

Head small, $5\frac{2}{3}$; depth $4\frac{2}{3}$. D. 58; A. 53. Body slender, covered with lax skin, which invests also the fins; mouth transverse, nearly horizontal, the maxillary scarcely vertical from front of pupil, equaling length of snout, $3\frac{1}{4}$ in head; eye $\frac{1}{3}$ interorbital width, 5 in head. Teeth in lower jaw pointed, in a single close-set series, their even tips forming an almost uniform cutting edge; teeth in upper jaw apparently paved, the jaw presenting a smooth, rounded surface, on which the teeth are little evident, but occasionally presenting sharp, projecting points; lower jaw included. A series of conspicuous mucous pores on snout, mandible, and on preopercle. Gill opening a narrow slit above base of pectorals. Pectoral fin divided into 2 wholly distinct lobes not connected by intervening rays, a space without rays present, equaling diameter of pupil; lower lobe of 4 rays, the longest $\frac{2}{3}$ head, inserted under preopercular margin; main portion of the fin with about 15 rays and but slightly longer than the lower lobe; dorsal beginning behind middle of pectorals, its distance from tip of snout $4\frac{1}{6}$ in length; distance of origin of anal from tip of snout equaling $\frac{1}{3}$ standard length; vent under base of upper pectoral rays, but wholly behind base of lower lobe of pectorals, its distance from tip of snout equaling length of head and equaling also its distance from origin of anal; no trace of ventral disk or fins. Color light rose-red, the fins and head largely jet black; abdomen behind vent blackish; peritoneum and lining membranes of mouth and gill cavity jet-black. Length $5\frac{1}{3}$ inches. Off coast of Southern California. (*rosaceus*, rosy.)

Paraliparis rosaceus, GILBERT, Proc. U. S. Nat. Mus. 1890, 93, off coast of California at Albatross Station, 2919, 984 fathoms; GARMAN, Discoboli, 80, 1892.

2475. PARALIPARIS MENTO, Gilbert.

Head $5\frac{2}{4}$ in length without caudal, equaling the greatest depth. D. 57; A. 43; pectoral 16. Eye a trifle less than interorbital width, greater than snout, 3 in head; mouth oblique, with lateral cleft, the maxillary reaching slightly beyond the middle of the orbit, $\frac{1}{3}$ as long as the head; chin very prominent, the mandible with a wide, membranaceous border, which forms anteriorly a broad free fold over anterior portion of base of pectorals. Teeth in both jaws comparatively long and slender, not tricuspid, arranged in bands, those in upper jaw at least in oblique rows as usual in Liparids. Nostril with a single opening and without tube. Gill opening but little wider than diameter of eye, confined to region above base of pectorals. Pectorals inserted very low, their upper rays below level of angle of mouth, the base almost horizontal, and the anterior ends of shoulder girdle form prominent projections below tips of mandibles, fin very deeply notched, the upper lobe extending slightly

beyond origin of anal, the shortest middle ray about ¼ that length; a few of the lower rays elongate and partly free from the membrane, the longest equaling the length of the upper lobe; middle pectoral rays wide set, but no rayless interval between the lobes; pectorals converging to immediately behind symphysis of lower jaw, where their bases meet; anal opening vertically below middle of orbit; dorsal beginning a trifle behind the upper angle of gill opening, the anal origin under its eighth ray. Skin loosely investing the head and body. Light reddish, made dusky by minute dark dots; abdomen blue black; mouth and gill cavity light or dusky. Closely allied to *Paraliparis (Hilgendorfia) membranaceus* (Günther), but with very heavy lower jaw, which projects beyond the upper. The premaxillaries are high, on the level of the lower rim of the orbit, and the form is much more elongate. The pectorals are inserted much lower, and the vent is farther forward. Length 3¼ inches. Off the coast of Oregon in deep water. (Gilbert.) (*mento*, having a projecting chin.)

Paraliparis mento, GILBERT, Proc. U. S. Nat. Mus. 1891, 562, off Coast of Oregon at Albatross Station 3071, in 685 fathoms.

2476. PARALIPARIS COPEI, Goode & Bean.

Head 6¼; depth about 6; snout 3½; eye 3¼. D. 60; A. 55; pectoral 17 + 3; caudal 10. Snout broad, obtuse; interorbital width equaling length of posterior part of head; nostril about midway between eye and tip of snout, in a very short tube. Five large mucous pores on each side of snout, 3 smaller ones on cheek under eye, and 6 on mandible and edge of opercular bones. Teeth minute, villiform, and in 2 series. Gill opening a very small slit, edged with black at upper angle of pectoral. Origin of dorsal distant from snout 1½ times length of head; tip of pectoral when extended about under second or third ray of dorsal; anal origin under ninth or tenth dorsal ray; length of middle caudal ray 10½ times in standard length; longest pectoral ray equaling ½ distance from snout to dorsal; longest ray of the isolated portion of the fin equaling that of middle caudal ray; distance of vent from tip of snout nearly equal to length of head; vent a narrow slit placed horizontally, directly under base of pectoral. Pyloric cæca 4 to 6. Color, anterior half of body very pale, whitish; posterior half very light brown; snout, chin, and edge of gill opening and region about the vent black. This species has nearly the same number of rays in the dorsal and anal as are recorded for *P. bathybius*, but the pectoral has 17 rays in the upper portion and 3 in the lower. The jaws are shorter than in *P. bathybius;* the form is more elongate and the coloration is strikingly different, anterior half of the body pale, almost whitish, while the snout, chin, and anal origin are black. Length about 8 inches. The eggs of an example a little smaller than the type have a diameter of about 3 mm. At the same time the ovaries contain some eggs, which have not reached maturity, whose diameter is only ½ mm. The stomach is a stout subspherical receptacle, its greatest width ⅔ of its length. Gulf Stream, in deep water. (Named for Prof. Edward Drinker Cope.)

Paraliparis copei, GOODE & BEAN, Oceanic Ichthyology, 279, fig. 253, 1896, Gulf Stream, at Albatross Station 2232, in 39° 12′ 17″ N., 72° 9′ 30″ W., in 520 fathoms. (Type, No. 35637.)

2477. PARALIPARIS DACTYLOSUS, Gilbert.

Head 5; depth 6¼. D. about 56; A. about 46. Shape of head much the same as in *P. ulochir*, the snout broadly rounder, slightly, if at all, overlapping the horizontal mouth, and the cheeks vertical. Occiput and nape not conspicuously swollen. Lower jaw included; maxillary reaching vertical behind middle of pupil, 2⅔ in head. Eye large, 3 in head; bony interorbital space 5; snout 4½. Gill opening a narrow slit extending to opposite third or fourth pectoral ray, its length equaling diameter of pupil. Teeth acute, in narrow bands in each jaw, each band made up of oblique rows. Opercle produced into a pointed lobe, which is separated from upper pectoral ray by a distance equaling diameter of pupil. Dorsal beginning slightly behind upper base of pectorals; pectorals inserted lower than in *P. ulochir*, the upper ray on a level with or below margin of pupil; as in *P. ulochir*, the 2 lobes joined by a series of about 8 more widely spaced rays, none of the rays is free; anterior ends of the fins meeting under throat at a point vertically below the pupil; pectoral rays 30, of which about 10 constitute the lower lobe; upper rays extending beyond front of anal, but the lower lobe apparently much shorter. Vent anterior in position, its distance from pectoral symphysis ½ its distance from front of anal. Each of the types has lost the epidermis. In this condition the head and body are light or slightly dusky, except the eyes, opercles, gill membranes, and abdomen, which are black; gill cavity and mouth black; probably black everywhere in life. Very close to *P. ulochir*, but with the upper insertion of pectoral much lower, opposite the lower margin of pupil, and the head and body much more slender and elongate. Length about 3¼ inches. Off Santa Cruz, California, in deep water. (*dactylosus*, δάκτυλος, fingered.)

Paraliparis dactylosus, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 469, pl. 34, fig. 2, off Santa Cruz, California, at Albatross Station 3112, in 296 fathoms.

Subgenus HILGENDORFIA,* Goode & Bean.

2478. PARALIPARIS ULOCHIR, Gilbert.

Head about 5 in length, equaling the depth. D. about 65; A. about 60. Snout broad and short, very slightly projecting beyond the mouth, the

* The subgenus *Hilgendorfia*, Goode & Bean, is thus defined: "Liparids with large, high, compressed head, whose bones are exceedingly thin, cavernous jaws, even in front. Vertical fins represented by broad median folds, confluent at the posterior end of the body, where the caudal is represented by 2 or 3 exceedingly fine, slender, terminal filaments. Pectoral very large, broad based, with membranes fringed posteriorly and having several of the lower rays free, though not forming a distinct portion of the fin. No ventral disk. Vent far advanced, opposite hind margin of orbit. Gill opening as in *Amitra*. The type described by Günther is only 60 mm. in length, and he remarks concerning it that it is uncertain whether it represents a form in which embryonic characters are persistent, or merely an early stage of development. Its peculiarities seem sufficient to warrant its being set apart for further study before it is merged with *Paraliparis*. It is named for Dr. Franz Hilgendorf, of the Royal Zoological Museum, Berlin." (Oceanic Ichthyology, 280, 1896.)

distance from tip of snout to front of eye $4\frac{1}{2}$ in length of head. Eye large, 3 to $3\frac{1}{4}$ in head, nearly twice the bony interorbital width. Differing from other Pacific species in the high insertion of the pectorals and their short horizontal limb, and from *P. holomelas* in having the fins not divided into 2 separate lobes. In general appearance greatly resembling *P. holomelas*, being also uniformly black in color, having the same broadly rounded snout, horizontal mouth with included lower jaw, and prominent occipital and nuchal region. The maxillary reaches vertical just behind pupil and is contained $2\frac{1}{10}$ times in head. Teeth acute, in rather broad bands in both jaws. Gill opening a narrow slit, restricted to area above base of pectoral, not longer than diameter of pupil. Opercle forming a short, pointed lobe posteriorly, which touches base of upper pectoral ray. The head is denuded of skin, so the nature of the mucous pores can not be made out. Dorsal beginning above upper base of pectorals; pectoral placed higher than in any other species known, its base describing a gentle, even curve, convex backward, horizontal for a very short distance anteriorly, its upper end above level of pupil, and its lower anterior end vertically below posterior margin of orbit; upper and lower rays of the fin fine and crowded, the middle third of the base being occupied by 4 or 5 more widely spaced rays; the fin with 25 rays, of which 9 belong to the lower lobe; upper lobe extending beyond origin of anal fin; none of the rays free; lower anterior ends of pectorals closely approximated, without perceptible interspace. Vent anterior in position, a distinct, though short, anal papilla. Distance from vent to anterior end of pectoral base $\frac{1}{3}$ its distance from front of anal. Longest specimen 85 mm. Uniformly black, including mouth and branchial cavity. Gulf of California and Bering Sea north of Unalaska Island, in deep water. (*οὐλός*, complete; *χείρ*, hand.)

Paraliparis ulochir, Gilbert, Rept. U. S. Fish. Comm. 1893(1896), 441, Gulf of California, at Albatross Station 3010, in 1,005 fathoms.

791. RHINOLIPARIS, Gilbert.

Rhinoliparis, Gilbert, Rept. U. S. Fish Comm. 1893 (1896), 445 (*barbulifer*).

Allied to *Paraliparis*, from which it differs in the gently produced snout, which much overlaps the mouth, and bears at its tip a pair of barbels. No sucking disk. Vent anterior, between the pectoral fins; pectorals deeply notched, continuous; gill openings narrow, mostly above the pectorals. Teeth acute, in a broad band in each jaw, arranged in oblique series within the band. Deep sea. (*ῥίν*, snout; *Liparis*.)

2479. RHINOLIPARIS BARBULIFER, Gilbert.

Head $5\frac{3}{4}$; depth 7; eye nearly 3. Slender, compressed, the greatest depth just in front of dorsal, the nuchal region not greatly swollen. Body tapering into an extremely slender, almost filamentous, tail. Mouth small, horizontal, inferior, overpassed by the broadly rounded, very soft snout for a distance equaling diameter of pupil. At the tip of the snout,

separated by a space ½ as wide as pupil, are 2 barbels directed forward, each as long as the interspace. Maxillary reaching vertical from posterior border of orbit, 2¼ in head; bony portion of interorbital width narrow, ¾ diameter of pupil; gill slit narrow, beginning opposite upper pectoral rays, ⅔ diameter of orbit; no pseudobranchiæ; opercle prolonged posteriorly into a narrow pointed flap; round mucous pores along under side of snout and suborbital, and on under side of mandible. Pectorals of 2 lobes, the lower narrow, containing but 4 or 5 rays, the upper with about 15, 1 or 2 rays widely spaced connecting the 2, none of the rays free, the fin inserted high, the upper end of base on a level with upper edge of pupil; below, the fins are not approximated as closely as usual, the lowermost rays of the 2 fins separated by an interspace as wide as pupil, inserted on a vertical through middle of cheek; dorsal originating slightly in front of gill slit. Vent anterior in position, its distance from front of pectorals ⅔ its distance from front of anal fin. In spirits, light gray, dusky along bases of dorsal and anal fins and on the nape; the black lining of abdominal cavity, gill cavity, and mouth can be seen through the transparent integuments; eye also black; barbels transparent, hence very inconspicuous. Length about 3½ inches. Bering Sea, north of Unalaska Island, in deep water. (*barbula*, a small barbel; *fero*, I bear.)

Rhinoliparis barbulifer, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 445, Bering Sea, north of Unalaska Island, at Albatross Stations 3227, 3325, 3326, and others, in 225 to 576 fathoms.

Suborder CRANIOMI.

Scapular arch abnormal, the post-temporal forming an integral part of the cranium and the postero-temporal crowded out of place by the side of the proscapula above or at the edge of the post-temporal. In other respects essentially as in the *Loricati*, from which the *Craniomi* are derived. The suborbital stay characteristic of both groups, in the *Craniomi* is highly developed, its surface forming part of the bony armature of the head. Fishes mostly of the warm seas, often singular in form, the head always with a coat of mail. (κράνιον, skull; ὦμος, shoulder.)

a. Myodome* developed and cranial cavity open in front; prosethmoid and anteal normally connected by suture. Infraorbital chain with its anterior bones excluded from the orbit and functional as rostrolateral, the series covering the cheeks, the third a large buccal bone articulating with the anterior wall of the preoperculum; post-temporal suturally connected with the opiotic and pterotic by inferior processes, and with the upper surface forming an integral part of the cranium; intermaxillaries with the ascending pedicles atrophied and connected with the knob of the anteal by ligament. Postero-temporal contiguous to the proscapula; ventral rays I, 5; the fins inserted wide apart; gill membranes free from the isthmus.

b. Pectoral fin with its 3 lowermost rays detached as feelers; teeth present.
TRIGLIDÆ, CLXXXIV.

bb. Pectoral fin with its 2 lowermost rays detached as feelers; mouth toothless.
PERISTEDIIDÆ, CLXXXV.

* The skeletal characters here given are taken from Dr. Gill. (Proc. U. S. Nat. Mus. 1888, 567–592.)

aa. Myodome undeveloped, the cranial cavity mostly closed in front by expansions from the subtectals suturally connected with corresponding expansions of the prootics and the parasphenoid; prosethmoid and anteal entirely disconnected, leaving a capacious rostral chamber opening backward mesially into the interorbital region. Infraorbital chain with its second and third bones crowded out of the orbital margin by junction of the first and fourth, and leaving a wide interval between the suborbitals and preoperculum; the first very long and extending backward, the second under the fourth, and the third developed as a small special bone (pontinal) bridging the interval between the second suborbital and the antero-inferior angle of the preoperculum; post-temporal suturally connected with the posterior bones of the cranium, and with the upper surface forming a large part of the roof of the head; intermaxillaries with well-developed ascending pedicles gliding into the cavity between the anteal and prosethmoid. Postero-temporal distant from the proscapula, and manifest as an ossicle on the edge of the post-temporal.

c. Pectoral fin divided to the base into 2 unequal parts; no free feelers; gill membranes broadly joined to the isthmus; ventral rays I, 4, the fins contiguous. CEPHALACANTHIDÆ, CLXXXVI.

Family CLXXXIV. TRIGLIDÆ.

(THE GURNARDS.)

Body elongate, usually more or less fusiform, covered with scales or bony plates. Head externally bony, entirely cuirassed with rough, bony plates, some of which are armed with spines; eyes high; mouth terminal or subinferior; premaxillaries protractile; maxillary without supplemental bone, slipping under the preorbital; teeth very small, in bands in the jaws, and usually on vomer and palatines; gills 4, a large slit behind the fourth; pseudobranchiæ present; gill rakers various; gill membranes free from the isthmus. Ventral fins thoracic, wide apart, separated by a flat area, their rays I, 5. Spinous dorsal present, short; soft dorsal similar to the anal, which is without spines; caudal narrow, few-rayed; pectoral large, with broad base, with 3 lower rays detached, forming feelers. These free rays are used chiefly in search for food, turning over stones, exploring shells, etc.* Air bladder present; pyloric cæca usually present, few in number. Singular looking fishes, found in all warm seas. Genera 5; species about 40, some of them in rather deep water, these red in color, the others living about rocks. (*Triglidæ*, pt., Günther, Cat., II, 191–210.)

a. Palatines with teeth.

b. Dorsal spines low, the longest usually much shorter than head; scales moderate, 50 to 80 pores. PRIONOTUS, 792.

bb. One or 2 of the dorsal spines greatly elevated, about as long as body; scales large, rough, the pores 40. BELLATOR, 793.

aa. Palatines toothless; scales small.

c. Lateral line without enlarged bony plates. CHELIDONICHTHYS, 794.

cc. Lateral line armed with a series of transverse bony plates. TRIGLA, 795

* See note by Albro D. Morrill, Journ. Morphology, XI, 1895, 177.

792. PRIONOTUS,* Lacépède.

(GURNARDS.)

Prionotus, LACÉPÈDE, Hist. Nat. Poiss., III, 37, 1802 (*evolans*).
Ornichthys, SWAINSON, Nat. Hist. Class'n Fishes, II, 262, 1839 (*punctatus*).
Chriolax, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1878, 374 (*evolans*).
Gurnardus, JORDAN & EVERMANN, new subgenus (*gymnostethus*).
Merulinus, JORDAN & EVERMANN, new subgenus (*carolinus*).

Body subfusiform; profile of head descending to the broad depressed snout, which is much longer than the small eye; eyes close together, high up; surface of head entirely bony, the bones rough with ridges and granulations; scales on head few or none; preopercle with 1 or 2 sharp spines at its angle; opercle with a sharp spine; nape with 2 strong spines; a spine on shoulder girdle. Mouth rather broad; bands of small, almost granular, teeth on jaws, vomer and palatines; gill membranes nearly separate, free from isthmus; gill rakers rather long. Body covered with small, rough scales, which are not keeled; lateral line continuous; scales on breast very small. Dorsal fins distinct, the first of 8 to 10 rather stout spines, the third usually highest, but mostly shorter than head; anal fin similar to soft dorsal; pectoral fin with the 3 lower anterior rays thickened, entirely free from each other and from the fin; ventrals I, 5, wide apart, with a flat space between them, the inner rays longest. Pyloric cæca in moderate number; air bladder generally with lateral muscles and divided into 2 lateral parts; vertebræ 10 or 11+15. Species numerous, all but 1 being American.† Representing in America the Old World genus *Trigla*. Some of them in deep water. They are well defined and easily recognized, but vary considerably with age, and are not easily thrown into subordinate groups. Most of the characters in the following analysis have been taken from adult individuals. Young examples in most cases differ from the adults in the following respects, in addition to those characters which usually distinguish young fishes: The spines on the head are sharper, more conspicuous, and more compressed in the young, and some spines, especially those on the side of the head, disappear entirely with age. The interorbital space is more concave in the young. The pectoral fins are also much shorter. The gill rakers are longer in the young, and proportionately more slender, and some of the color markings—especially the darker cross shades—are more conspicuous, while the spots on body and fins are less so. (*πρίων*, saw; *νῶτος*, back; 3 free saw-like spines being said to intervene between the two dorsals.)

a. Mouth comparatively small, the maxillary less than ½ the length of the head, the mandible usually not extending backward as far as the vertical from the front of the eye; generally a more or less distinct cross groove on top of head behind eye; black spot on spinous dorsal usually more or less distinct.

GURNARDUS (from Gurnard):

b. Snout distinctly birostrate, its tip deeply emarginate; anterior profile of head strongly concave; pectoral fin very short, little if any longer than head; interorbital space deeply concave.

* For a detailed account of the species of *Prionotus* see Jordan & Hughes, Proc. U. S. Nat. Mus. 1886, 327, *et seq.*

† The single species found outside our limits is *Prionotus japonicus*, Bleeker.

c. Preopercular spine with a smaller one at base in front.

d. Center of radiation of cheek with a strong spine, there being 4 spines in a right line from rostral plate to preopercular spine; rostral projections very narrow; body slender; scales small; 3 dorsal spines granulate.

BIROSTRATUS, 2480.

dd. Center of radiation of cheek without spine; 1 or 2 spinules below preopercular spine.

e. Breast and belly wholly naked; dorsal spines 10 or 11. GYMNOSTETHUS, 2481.

ee. Breast and belly with small scales; dorsal spines 8 or 9. XENISMA, 2482.

cc. Preopercular spine with no smaller one at base in front; dorsal spines 10; pores 50; body with olive vermiculations and oblique brown bars. LOXIAS, 2483.

MERULINUS (*Merula*, a robin):

bb. Snout not distinctly birostrate, the anterior profile usually not strongly concave.

f. Pectoral fin long, reaching past front of anal.

g. Pectoral fin not reaching base of caudal; gill rakers moderate, 8 or 10 developed; snout not strongly emarginate; no spine on cheek bone or edge of snout; dorsal spines 10.

h. Body not very slender, the depth 5 in length; head not very small, its length 3 in body; groove across top of head behind eye, very conspicuous; interorbital area moderately concave, rather broad, about equal to diameter of eye; bones of head comparatively smooth, the preocular, postocular, occipital, and nuchal spines low, depressed; temporal ridge conspicuous, without spines. Dorsal spines low, the second $2\frac{1}{3}$ in head, the first moderately serrate; base of soft dorsal equal to distance from tip of snout to tip of humeral spine; caudal fin lunate, its outer rays $\frac{1}{3}$ to $\frac{1}{4}$ longer than inner; pectoral fin somewhat rounded, the longest ray about the fifth; free rays of pectoral expanded toward tip, with decurrent membrane; scales rather large; about 58 pores. Body and fins nearly plain, mottled with darker, but without well-defined spots except the dorsal ocellus; back with 4 obscure cross blotches; 2 or 3 oblique pale streaks across spinous dorsal. Gill membranes dusky. Young with head rougher, pectoral fins shorter, dark spots on body more distinct. CAROLINUS, 2484.

hh. Body very slender, the depth about 6 in length; groove across top of head behind eye, conspicuous; interorbital area narrow, deeply concave, its width about $\frac{2}{3}$ the diameter of the eye; bones of head very smooth, the striations very weak; spines on top of head (preocular, supraocular, occipital, and nuchal) short and sharp, not depressed; temporal ridge blunt, without spine.

i. Pectoral short, reaching little past front of anal, not $\frac{1}{5}$ length of body. Dorsal spines very high, the second $1\frac{3}{4}$ in head, the first moderately serrate; soft dorsal high, its base about $\frac{1}{4}$ longer than head; caudal truncate; free rays of pectoral a little expanded at tip; 52 pores. Body covered with roundish bronze spots of various sizes; smaller bronze spots on the head; both dorsals, caudal, and pectoral fins with similar bronze spots, these especially numerous and distinct on soft dorsal. SCITULUS, 2485.

ii. Pectoral longer, reaching past middle of anal, more than $\frac{1}{4}$ body; pores 62; color rose-red, not spotted; pectorals dusky. ROSEUS, 2486.

gg. Pectoral fins very long, reaching base of caudal, the rays graduated; 50 pores in the lateral line; gill rakers shortish, 1+6 in number; body rather stout, the depth 4 in length; palatine teeth few, feeble; caudal subtruncate; second dorsal spine longest, $\frac{1}{3}$ the length of head; first spine strongly serrated in front; preopercular spine with a smaller one at its base; head $2\frac{1}{3}$ in length. D. X-12; A. 11. Body with 4 faint cross bands; caudal with black tip and 2 paler cross shades; spinous dorsal with small dark spots besides the large one; soft dorsal plain; pectorals clouded. ALATUS, 2487.

PRIONOTUS:

aa. Mouth comparatively large, the maxillary 2 to 2¾ in the length of head, the mandible extending backward to opposite eye, or nearly so; usually no distinct cross groove on top of head; free rays of pectoral tapering, not expanded at tip; black blotch on spinous dorsal diffuse, not ocellated, involving the membranes of more than two spines.

j Preopercular spine without a distinct smaller spine at its base in front.

k. Pectoral fins very long, reaching at least to beyond the second third of the soft dorsal.

l. Scales rather small, about 75 pores in lateral line; head large, comparatively smooth, the interorbital area narrow, concave, not ⅔ diameter of eye; orbital rim low, scarcely serrate; preorbital serrulate; pectoral fin long, reaching middle of anal; coloration plain, slightly clouded; form rather slender; head 2⅔; depth 4⅓. D. X–12; A. 11. MILES, 2488.

ll. Scales moderate, 50 to 55 pores in lateral line.

m. Head large, nearly plane above, the interorbital space scarcely concave, its width equal to eye; snout very short, 2⅓ in head; maxillary 2⅛ in head, reaching past front of orbit; bands of palatine teeth very narrow; bones of head smoothish, little striate; gill rakers rather long, about 10 below angle; first dorsal spine slightly rough in front, the third spine nearly ½ the length of the head; caudal slightly concave; pectoral reaching ninth anal ray, its outline rounded; scales about 55. Color pale olivaceous, back and sides with a few dark spots; second dorsal with 3 rows of black spots; caudal with 3 rows of dark blotches; pectorals with round dark blotches. D. X–12; A. 11. STEPHANOPHRYS, 2489.

mm. Head not plane above, the interorbital space more or less concave; dorsal spines 10.

n. Supraorbital cirrus wanting.

o. Pectoral fins moderate, not reaching in adult beyond dorsal and anal.

p. Interorbital space moderately concave, its width nearly equal to eye; orbital rim little elevated; preorbital with 15 to 20 fine serræ, the last one directed backward; preopercular spine long. QUIESCENS, 2490.

pp. Interorbital space very narrow, deeply concave, its width about ⅔ length of eye; rim elevated and serrate; preorbital with about 20 small serræ, blunt and even; head well armed; snout and jaws white. ALBIROSTRIS, 2491.

oo. Pectorals very long, reaching in adult beyond base of dorsal and anal; interorbital space moderately concave, its width about ⅓ length of eye; no cirrus above the eye; distance from supraocular spine to nuchal scales about equal to eye; supraocular and nuchal spines low; occipital spines wanting; temporal ridge sharp, ending in a blunt spine; preorbital projecting, strongly serrate; a blunt spine on each side of snout, behind serræ of preorbital; a blunt spine behind this above angle of mouth; no spine on cheek bone in adult; upper opercular spine almost obsolete; bones of head rather strongly striate, but not granulate; gill rakers short; mouth moderate; scales rather large; about 52 pores. D. X–11; A. 10. First dorsal spine not much shorter than second, which is 2⅛ in head; caudal very slightly concave; ventrals reaching a little past vent; head 3; depth 5. Color nearly plain brownish, with darker clouds; no distinct spots anywhere on body or fins; the pectorals marbled with paler. RUBIO, 2492.

nn. Supraorbital cirrus present, fringed. Interorbital space very deeply concave, its width about ⅔ length of eye; a fringed cirrus above the eye; distance from supraocular spine to nuchal scales about ½ the eye; occipital as well as nuchal spines distinct; temporal ridge with a small spine; no spine on cheek bone; bones of the head with fine, sharply defined striæ, but no granulations; upper opercular spine well developed;

scales moderate (about 50 pores). D. VIII–13; A. 11. First dorsal spine longest, 1⅓ in head; caudal subtruncate; pectorals nearly twice as long as head, reaching nearly to last rays of dorsal; ventrals about reaching to vent; head 3 in length; depth 4⅓. Color crimson red, nearly plain; caudal with 2 dark cross shades. OPHEVAS, 2493.

kk. Pectoral fins short, not reaching beyond middle of dorsal; head much smoother than in any other species, the bones of the head faintly striate, with small granulations; the cranial spines little developed; the supraocular, occipital, and temporal spines wholly wanting, there being only 3 pairs of spines on the head; mouth large, the maxillary 2 in head. Gill rakers short and thick in adult, slender in young, about 10 developed; interorbital space concave, rather broad, its width, in adult, rather more than length of eye; first dorsal spine granulated; caudal slightly lunate; pectoral subtruncate, the second ray the longest, as long as head in adult; scales large, 48 pores in the lateral line. Head large, 2⅔ in length; depth 3⅔. D. X–12; A. 11. Color crimson, with darker clouds and small spots; both dorsals with dark cross streaks; head and pectoral fins conspicuously reticulated with blackish (in adult); anal plain, whitish; free rays of pectoral unspotted. STEARNSI, 2494.

jj. Preopercular spine with a distinct smaller one at the base; gill rakers slender.

q. Cheek bone without distinct spine at the center of radiation; edge of preorbital granular-serrate, without distinct spine, the serræ about 12 in number on each side; temporal ridges roughish but without spines; bones of the head with the striæ coarsely granular; mouth moderate, the maxillary about 2⅔ in head; head not very broad, the spines above, except the nuchal spines, not conspicuous; gill rakers long and slender, 15 to 20 developed; head 2⅔ in length; depth about 4. D. X–12; A. 11. Coloration brownish; side with a very distinct dusky bronze band below the lateral line and parallel with it, this becoming broken posteriorly into a series of roundish dark spots; some dark streaks and clouds below this stripe; fins with dark clouds, the soft dorsal with 2 dark blotches, which extend as bars on the back; head with scattered dark spots; dusky area below eye.

r. Pectoral with its rays each crossed by fine black bars, these especially distinct toward the base of the fin; free rays spotted; scales comparatively small, 10+1+23 in a vertical line from last dorsal spine to vent; interorbital area broad and almost flat, its width a little more than length of eye; first dorsal spine granulated; second spine 2¾ in head; pectorals about ⅓ the length of the body. STRIGATUS, 2495.

rr. Pectoral fin with its rays all plain blackish; free rays plain dusky; scales larger, 8+1+21 in a vertical line from last dorsal spine to vent; interorbital space more deeply concave, its width in adult not quite length of eye; first dorsal spine nearly smooth; second spine 3 in head; pectorals a little more than ⅓ of the body. EVOLANS, 2496.

qq. Cheek bone with a spine (small in the adult, larger in the young) at the center of radiation, this rarely obsolete in old examples.

s. Spines on bones of head moderate, not knife-like; preorbital with a series of serræ and 1 or more bluntish spines.

t. First 3 dorsal spines little if at all serrate; pectorals reaching past middle of anal, their length not quite ⅓ the body; gill rakers rather long and slender, about 10 developed; maxillary 2⅔ in head; a bluntish spine on edge of snout behind the serræ; behind this, 1 or 2 smaller ones, at least in the young; usually a small spine on cheek bone; a shallow groove behind the eye evident; interorbital area rather narrow, concave; preocular, supraocular, occipital and nuchal spines rather prominent. Dorsal spines high, the third 2⅓ in head;

first spine not serrate; caudal truncate. Head 2⅘ in length; depth 2⅔. D. X-12; A. 12. Pores about 50. Back obscurely spotted; dorsal and caudal fins spotted with brown, the first dorsal with a black blotch besides; the pectoral with obscure dark spots, and margined with blue. PUNCTATUS, 2497.

tt. First 3 dorsal spines more or less serrate; gill rakers rather long, about 8 developed; maxillary about 2⅔ in head; a groove behind eye; interorbital space narrow, ⅔ eye; third dorsal spine high, ⅓ head. D. X-12; A. 11. Pores 50. Color brownish yellow; spinous dorsal with a black blotch; pectoral with 2 longitudinal broad dark areas separated and surrounded by paler. BEANII, 2498.

ss. Spines on bones of head elevated, knife-like; head very large, more than ⅔ length; temporal ridge with 2 bluntish spines; bones of the head very sharply striate; young with 4 sharp, knife-like spines on side of cheek and snout, in a line before the preocular spine, these nearly disappearing with age; maxillary about 2⅓ in head; sides without dark longitudinal stripe.

u. Pectoral fin moderate, about ⅓ body in adult, 2⅓ in young; gill rakers slender in the young, becoming shorter and thicker with age, about 10 developed on lower part of arch; head broad, the spines on its upper surface very prominent, all of them more or less compressed and knife-like, especially in the young. Second dorsal spine 2⅓ in head; head 2⅓; depth 4⅓. D. X-12; A. 11. Body brownish, much mottled with grayish and dusky, and with 3 or 4 obscure dark cross bands; head and dorsal fins with many dark spots; caudal with 2 dusky shades; free rays of pectoral spotted. TRIBULUS, 2499.

uu. Pectoral fin short, about 3 in body; gill rakers long and slender, 5 developed; scales small, 100; spines as in *P. tribulus*, but still larger and more knife-like, much larger than in any other species; pectoral fins reaching third anal ray; interorbital space broad, little concave, about as wide as eye. Body brownish; no black blotch on spinous dorsal; soft dorsal and caudal barred, pectoral with a broad black band. HORRENS, 2500.

Subgenus GURNARDUS, Jordan & Evermann.

2480. PRIONOTUS BIROSTRATUS, Richardson.

Head 3; depth 5. D. IX-12; A. I, 10; C. 10. Body slender and elongate; width at nape between costal spines 5½ in head; head long, profile before eyes concave, the snout with a projecting lobe on each side, the lobe longer and narrower than in any other species, its length ⅔ eye; a backward projecting spine at base, besides numerous retrorse serræ along its sides; surface of bones very rough; mouth small; maxillary barely ⅓ head, not reaching front of eye; eye 4½ in head; interorbital space very narrow, deeply concave, granular, striate, its length ⅔ eye, 7 in head; orbital rim elevated, with a stout spine anteriorly, before which are 3 or 4 others forming a raised ridge; a stout spine above posterior part of orbit, another behind orbit; still another on the occipital beyond ridge; 2 or 3 blunt spines behind eye; a stout spine on opercle which reaches slightly beyond base of the still larger humeral spine; a stout spine on preopercle

with another strong spine at its base; another before it on radiation center of cheek; another on the preorbital, these 4 forming a straight line with retrorsal projection and all of them hooked backward; no curve across top of head; membranaceous flap of opercle not scaly; gill rakers unknown; scales rather small, the number not to be counted, apparently none on breast, the naked skin of breast forming an acute angle behind; 7 rows of scales between occiput and dorsal; spinous dorsal rather low, its first 3 spines serrate in front, the second the highest, $\frac{1}{2}$ head; soft dorsal with first row serrated, its length $2\frac{1}{6}$ in head; anal lower, its longest ray $2\frac{2}{3}$ in head; caudal truncate, $1\frac{2}{3}$ in head; pectoral short, its middle rays longest, reaching fifth ray of soft dorsal, $2\frac{2}{3}$ in body; ventrals $1\frac{2}{3}$ in head; free rays tapering, $1\frac{2}{3}$ in head. Coloration not described; body and fins in figure unmarked, except pectoral, which is blackish at tip. Gulf of Fonseca; known from a single specimen 6.7 inches long. Here described from Richardson's figure, which gives details not mentioned in his description. Apparently a well-marked species with the spines of *Prionotus horrens*, but with a small mouth and rostral beak different from that of any other species. (*birostratus*, two-beaked.)

Prionotus birostratus, RICHARDSON, Voyage of the *Sulphur*, Ichthyology, pt. 2, 81, April, 1845, Gulf of Fonseca, west Coast of Central America. (Coll. Edward Belcher.)

2481. PRIONOTUS GYMNOSTETHUS, Gilbert.

Head $2\frac{1}{3}$; depth $3\frac{1}{3}$; eye 4 in head; snout $2\frac{7}{8}$; interorbital width 10; pectoral $1\frac{1}{2}$; ventrals $1\frac{1}{3}$; second dorsal spine 2. D. X or XI–10 or 11; A. 11. Body heavy anteriorly, tapering rapidly to the slender tail. Head large; striæ fine, numerous, scarcely to be distinguished on top of head, which is finely granular; snout long; profile concave, the orbital region abruptly elevated; preorbital on each side projecting beyond the mouth in a narrow lobe, which is usually sharply triangular, the 2 lobes strongly divergent; free edge of preorbital with a row of very fine serræ; anterior nostril with a long flap; mouth small, the maxillary $\frac{1}{3}$ length of head, a naked area behind it. Eyes very prominent, the preocular ridge high, bearing 1 strong spine; interorbital space very narrow, concave, $\frac{1}{10}$ length of head; a deep transverse groove on top of head behind eyes. Supraocular spine short and blunt; occipital and nuchal spines flat, bluntish; temporal ridge with 2 spinous projections; opercular, preopercular, and humeral spines, long and sharp; preopercular spine with a very strong accessory spine at base, from which a low ridge extends forwards to middle of cheek, where it ends in a weak spine. Gill rakers 1+7, slender, of moderate length. Second dorsal spine the highest, the first 2 spines minutely serrulated in front; pectorals very short, $\frac{1}{3}$ the length, reaching but little past front of anal; caudal slightly lunate. Scales very small, none on breast or belly, the naked area extending upwards behind pectoral fin to humeral spine, and narrowing backward to front of anal; region in front of spinous dorsal naked. Color in spirits, brownish above, light below, the dorsal region and top of head often with small dark spots; 1, or rarely 2, small black ocel-

lated spots between fourth and fifth dorsal spines; pectoral mostly dusky, light at base below, and edged with white; a dusky blotch on outer portion of lower caudal lobe. Gulf of California; several specimens taken in shallow water. Length 3½ inches. Closely related to *Prionotus xenisma*, differing conspicuously in the wholly naked breast and belly. (Gilbert.) (γυμνός, naked; στῆθος, breast.)

Prionotus gymnostethus, GILBERT, Proc. U. S. Nat. Mus. 1891, 559, Gulf of California. (Coll. *Albatross.*)

2482. PRIONOTUS XENISMA, Jordan & Bollman.

Head 2½ to 2⅝ (3 to 3¼ in total); depth 3⅛ to 3½ (4 to 4¼); eye 4 in head. D. VIII–11; A. 10; scales in a longitudinal series 60 to 70, about 43 oblique series between gill opening and tail; 35 to 40 pores in lateral line. Body short and robust, little compressed, the width of the nape, between occipital spines, 6 to 6½ in head; head short and high; eyes prominent; the profile angulated, concave before eye, convex above it; snout short and broad, its breadth greater than its length, which is 2⅝ in head, its anterior margin deeply emarginate, the preorbital produced on each side into a broad, triangular, spiniferous lobe, which is nearly equal to ½ diameter of eye. Surface of bones of head finely and densely granular striate. Mouth small, maxillary 3 to 3⅛ in head, not reaching front of eye. Band of palatine teeth rather broad. Interorbital space narrow, deeply concave and granular striate, its least width 6 to 6½ in head; orbital rim elevated, its edges granular serrate; no distinct preorbital spine, whole edge of preorbital finely serrulate; a groove across top of head behind the blunt supraorbital spine. Occipital ridges present, the outer granular and not spinous, the inner rather sharply elevated into a short spine, the outer rather wide, extending to second dorsal spine; temporal region with an elevated roughish ridge, on which are 2 bluntish prominences; preopercular spine large and strong, with a smaller one at its base; opercular spine long and sharp; humeral spine usually larger than opercular; no spines on suborbital. Breast scaly; membranaceous flap of opercles not scaly. Gill rakers short, slender, longer than interspaces, and about equal to ⅔ of pupil; naked skin of throat with numerous papillæ; scales rather small, of about equal size on breast and belly, extending beyond base of pectorals, anterior margin forming an obtuse angle; 3 or 4 rows of rudimentary scales between occiput and dorsal. Spinous dorsal high, the first spine longest, 1⅔ to 1⅞ in head; anterior margin of first 3 spines with numerous granulæ; second spine slightly more than 2 in head, all the spines stout; longest ray of second dorsal shorter than snout, 3⅛ in head; longest anal ray 3⅜ in head; caudal slightly lunate, 1¾ in head; pectorals quite short, reaching fifth ray of second dorsal, 3 in body; ventrals reaching anal, 1⅔ in head. Three short granular interspinal bones projecting through the skin between first and second dorsal, the anterior less robust than others. Coloration in spirits, grayish (probably red in life); dusky above and irregularly mottled; a distinct dark spot before base of caudal; dorsal dusky, with a large black, ocellated spot between fourth and fifth spines; dorsal spines with a row of

dark spots on their anterior margin; soft dorsal with 3 rows of dark spots; pectorals blackish, a few of the upper rays pale on the inner side; ventrals and anal pale; caudal barred with dusky. Length 4 inches. Pacific coast of Colombia. Very many specimens dredged at a depth of 33 fathoms. (ξένισμα, a surprise.)

Prionotus xenisma, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 169, Pacific Ocean off coast of Colombia, at Albatross Station 2795, 7° 57′ N., 78° 55′ W. (Type, No. 41151.)

2483. PRIONOTUS LOXIAS, Jordan.

Head 2½; depth 3½. D. X–10 or 11; A. 10; scales about 50. Body stout; head large, rough; mouth moderate, maxillary not reaching front of orbit, 2⅔ in head; eye large, 3¾ to 4 in head; snout 2½ in head; opercular spine strong, nearly as large as preopercular spine; humeral spine small, not ½ as large as either of the others; interorbital area narrow, its ridge evenly concave, its width 2⅔ in eye; preopercular spine without smaller one in front; no spine at center of radiation of cheek; preorbital edge prominent, finely denticulated; supraorbital ridge prominent, with a bluntish spine before and behind; a transverse groove on head behind eye; snout broad, slightly emarginate at tip, the rostral plates not much projecting, their edges sharply and finely serrate, with 10 to 12 serræ; occipital ridges a short distance behind supraorbital ones, ending in a bluntish spine, as also the nuchal ridges. Teeth on jaws, vomer, and palatines in bands; lower jaw included; base of mandible below front of orbit; bones of cheeks and opercles with strong striæ, the rest of the bones of the head roughish. Gill rakers short, about 10 below the angle, the anterior ones tubercle-like; breast closely scaled. Pectorals short, 3¼ to 3½ in the length of the body, scarcely longer than longest detached ray, their tips reaching about third ray of anal fin, length 3½ in the body; ventrals long, their tips almost reaching tips of pectorals, 1⅜ in head; first dorsal spine the longest, its length 1⅘ in head, serrate in front; first dorsal ray slightly serrulate at base, its length 3 in head; longest anal ray 4 in head; caudal fin lunate, with pointed lobes, 1¾ in head. Color, brownish above, grayish below; head and anterior parts more or less distinctly vermiculated with dark olivaceous, these markings especially distinct on bones of head; sides with 6 to 15 narrow brown oblique bands extending downward and backward from the lateral line about halfway to anal fin, these obsolete or less conspicuous on anterior portion of the body; both dorsals mottled with olive; caudal with 3 broad blackish bars which do not cross the upper and lower ray, the last bar broad and very conspicuous; upper ray of caudal dark olive; no black spot at base of caudal; anal and ventrals white; pectorals blackish, faintly barred with darker and margined with white. Pacific coast of Central America. Here described from many specimens, 3 to 6 inches long, from *Albatross* Station 2805, where it occurs with *Prionotus xenisma*, but more abundantly than the latter. It has not yet been seen elsewhere. This species resembles *Prionotus xenisma* very closely. It is, however, a little more elongate, with rather smaller scales, the snout longer, the caudal fin barred, the

body more vermiculate and with oblique bars, and no distinct dark spot at base of caudal. There is no trace of a second spine at the base of the large preopercular spine, and the humeral spine is much smaller than the opercular. Both have the snout somewhat birostrate, the forehead and eyes very prominent, and the pectorals very short. The projection of the lobes of the snout is, however, much less than in *Prionotus birostratus*, with which species the present one was at first identified. (λοξίας, λοξός, oblique, from the cross bars.)

Prionotus birostratus, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 170; not of RICHARDSON.

Prionotus loxias, JORDAN, in GILBERT, Proc. U. S. Nat. Mus. 1896, 452, at Albatross Station 2805, south of Panama, in 51½ fathoms. (Type, No. 47580, U. S. Nat. Mus.; cotype, No. 165, L. S. Jr. Univ. Mus.)

Subgenus **MERULINUS**, Jordan & Evermann.

2484. PRIONOTUS CAROLINUS (Linnæus).

(COMMON GURNARD; RED-WINGED SEA-ROBIN.)

Head 3; depth 5. D. X–13; A. 12; scales 58. Body not very slender; head moderate; mouth comparatively small, the maxillary about 3 in head, the mandible not reaching vertical from front of eye; groove across top of head behind eye, very conspicuous; interorbital area moderately concave, rather broad, about equal to diameter of eye; bones of head comparatively smooth, the preocular, postocular, occipital and nuchal spines low, depressed; temporal ridge conspicuous, without spines; 1 or 2 small spinules on lower edge of preopercle, below the preopercular spine; preopercular spine with no smaller one at its base in front; pectoral fin short, reaching little past front of anal, its length less than ½ the body; gill rakers of moderate length, about 10 developed; no spine on cheek bone or on sides of snout. Dorsal spines low, the second 2⅓ in head, the first moderately serrate; base of soft dorsal equal to distance from tip of snout to tip of humeral spine; caudal fin lunate, its outer rays ⅓ to ¼ longer than inner; pectoral fin somewhat rounded, rather short, not reaching last ray of anal and not more than ½ the length of the body; the longest ray about the fifth; free rays of pectoral expanded toward tip, with decurrent membrane; scales rather large, about 58 pores. Body and fins nearly plain, mottled with darker, but without well-defined spots except the dorsal ocellus; back with 4 obscure cross blotches; 2 or 3 oblique pale streaks across spinous dorsal; gill membranes dusky. Young with head rougher, pectoral fins shorter, dark spots on body more distinct. Coast of Maine to South Carolina, chiefly northward; very abundant on the coasts of southern New England and New York, but rarely taken as far south as Charleston. Our specimens are from Menemsha Bight, Marthas Vineyard. (*carolinus*, from Carolina.)

Trigla carolina, LINNÆUS, Mantissa, 176, 528, Carolina.

Trigla palmipes, MITCHILL, Trans. Lit. and Phil. Soc. New York 1814, 431, pl. 4, fig. 5, New York Harbor.

Prionotus pilatus, STORER, Proc. Bost. Soc. Nat. Hist., II, 1845, 77, Massachusetts Bay, adult specimens; STORER, Hist. Fish. Mass., 68, pl. 6, fig. 1, 1867; GOODE & BEAN, Bull. Essex Inst., XI, 12, 1879.

Prionotus carolinus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 90, 1829; STORER, Report Fishes Mass., 14, 1839; DE KAY, New York Fauna: Fishes, 46, pl. 5, fig. 15, 1842; AYRES, Bost. Journ. Nat. Hist., IV, 1842, 258; GÜNTHER, Cat., II, 192, 1860; GILL, Cat. Fish. East Coast N. Am., 21, 1873; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1878, 373; BEAN, Proc. U. S. Nat. Mus. 1880, 84; GOODE & BEAN, Bull. Essex Inst., XI, 12, 1879; JORDAN & HUGHES, Proc. U. S. Nat. Mus. 1886, 333.

Prionotus palmipes, STORER, Hist. Fish. Mass., 66, pl. 5, fig. 1, 1867; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1883, 614; JORDAN & GILBERT, Synopsis, 734, 1883; GOODE, Nat. Hist. Aquatic Animals, 255, pl. 71, 1884; GOODE & BEAN, Oceanic Ichthyology, 468, 1896.

2485. PRIONOTUS SCITULUS, Jordan.

Head 2¾ to 3⅜; depth 5⅛ to 6⅓. D. X–13; A. 12; scales rather small, about 60; 52 pores; body much slenderer than in any other species; head small, low, rather pointed. Snout rather long, a little shorter than rest of head, its width between angles of mouth about 2½ in head. Maxillary not reaching front of eye, 2⅔ in head. Sides of snout finely and evenly serrate; mouth comparatively small, the maxillary less than ⅔ the length of the head; groove across top of head behind eye, little conspicuous; interorbital area narrow, deeply concave, its width about ⅔ the diameter of the eye; bones of the head very smooth, the striations very weak; spines on top of head preocular, supraocular, occipital, and nuchal) short and sharp, not depressed; temporal ridge blunt, without spine; 1 or 2 small spinules on lower edge of preopercle, below the preopercular spine; dorsal spines very high, the second 1¾ in head, the first moderately serrate; soft dorsal high, its base about ¼ longer than head; caudal truncate, its third ray longest, the others, to the tenth, little shorter; free rays of pectoral a little expanded at tip; scales rather small. Bands of palatine teeth narrow. Gill rakers long and slender. Pectoral in male 2⅛ to 2⅖ in body, in female scarcely more than ⅛ length of body, reaching to base of fifth or sixth dorsal ray. Coloration of female in life, dark olive above; back and sides covered with numerous round spots of different sizes, and not arranged in series, these spots bronze color in life, becoming brownish after death; spinous dorsal dusky, with lighter streaks; a distinct black spot on upper half of spinous dorsal, between the fourth and fifth spine, this spot being ocellated below and behind; a second black blotch on upper half of first spine and membrane, also ocellated behind; second dorsal and caudal spotted and finally blotched with black; anal largely black, with a pinkish border; pectorals blackish; ventrals pale; branchiostegals pinkish. Male in life, light olive brown, with 4 saddle-like dark blotches on back, 1 downward and forward from middle of spinous dorsal to humeral spine; a second from front of soft dorsal; a third from end of dorsal downward and forward to below lateral line, thence continued forward as a narrow horizontal streak; a fourth on caudal peduncle; sides everywhere with reddish brown spots, as in the female; opercle reddish brown; branchiostegal membrane and palatine region largely jet-black; spinous dorsal olive brown, with 2 irregular, lengthwise, translucent streaks and an intense well-defined black spot on membrane above, between fourth and fifth spines; second dorsal olive brown, vermiculated with whitish translucent, and without round spots; caudal reddish brown, blackish

toward tip, with a conspicuous, white, longitudinal streak on upper lobe; anal blackish, with white base and margin; pectoral dark brown, irregularly barred and blotched with greenish and light brown; free rays of pectorals and inner face of ventrals dusky, tinged with orange. South Atlantic coast of United States, Beaufort to St. Augustine; a very well-marked species, rather common within the region from which it is known. Length 5 to 6 inches. (*scitulus*, slender.)

Prionotus punctatus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1878, 373; GOODE, Proc. U. S. Nat. Mus. 1879, 111; GOODE & BEAN, Proc. U. S. Nat. Mus. 1879, 338; JORDAN & GILBERT, Synopsis, 734, 1883; not *Trigla punctata*, BLOCH.

Prionotus scitulus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 288, Beaufort (Coll. Jordan, Brayton & Gilbert); JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 614; JORDAN, Cat. Fish. N. A., 114, 1885; JORDAN & HUGHES, Proc. U. S. Nat. Mus. 1886, 333.

2486. PRIONOTUS ROSEUS, Jordan & Evermann.

Head $3\frac{1}{6}$ in length (4 with caudal); depth 6 ($7\frac{1}{2}$); eye moderate, $5\frac{1}{5}$ in head. D. X–12; A. 12; scales (transverse series) 65 to 70; pores in lateral line 60 to 65. Allied to *Prionotus scitulus*. Body quite slender, little compressed, narrowed above, width of nape between occipital spines being about $\frac{1}{2}$ of head; head quite short and rather high; eyes prominent, so that the anterior profile forms a sharp angle at front of eye, and is somewhat concave; snout rather narrow, $2\frac{1}{3}$ in head, its tip somewhat emarginate; edges of snout finely serrulate and without spine; surface of bones of head comparatively smooth, the small granulations arranged in distinct, fine, radiating striæ. Mouth small, the maxillary $3\frac{1}{4}$ in head, the mandible not quite reaching front of orbit; band of palatine teeth narrow. Apparently a slight cirrus above eye, on one side, in typical example, possibly a result of mutilation. Interorbital space narrow, deeply concave, its least width $7\frac{1}{4}$ in head; orbital rim considerably elevated, both in front and behind, its edge granular serrate; a shallow groove across top of head behind orbital rim, which does not end in a distinct spine; occipital ridges weak, the inner pair without spines, the outer with short ones which reach somewhat beyond front of dorsal; temporal region with an elevated roughish ridge, but without distinct spine; preopercle with a single moderate spine, which has no smaller one at its base, either in front or below; opercular spines small and sharp; humeral spine moderate; no trace of spines on suborbital or preorbital, the head being provided with but 5 pairs of spines, including the humeral spine; membranaceous flap of opercle scaly. Gill rakers short and thickish, about 8 developed, these little longer than the interspaces, and not $\frac{1}{2}$ length of pupil and nearly $\frac{1}{2}$ as broad as high. Scales small, those on breast much reduced in size, about 12 between occiput and dorsal. Spinous dorsal high, the first spine serrulate in front, shorter than the second, which is $1\frac{3}{4}$ in head. (Second dorsal and anal mutilated.) Caudal slightly and unequally lunate, the lower lobe the longer, $1\frac{1}{6}$ in head; pectoral reaching nearly to last rays of dorsal, a little more than $\frac{1}{2}$ length of body; ventrals as long as head. Coloration in spirits, grayish, unspotted, more dusky above; dorsal dusky,

with no very distinct markings; caudal fin yellowish, marked at base with dusky, its tip black; pectorals mostly black; lower fins pale. In life the type was chiefly pinkish red, which color still persists on the inside of opercles. Deep waters of Gulf of Mexico, known only from the "spewings" of Red Groupers, on the Snapper Banks off Tampa Bay and Pensacola. Length of type 6½ inches. (*roseus*, rose red.)

Prionotus scitulus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 288, mutilated specimen from Pensacola; not type.

Prionotus roseus, JORDAN & EVERMANN, Proc. U. S. Nat. Mus. 1886, 470, off Tampa Bay, Florida. (Type, No. 37989; Coll. Chas. H. Bollman.)

2487. PRIONOTUS ALATUS, Goode & Bean.

Head 2⅓; depth 4. D. X–12; A. 11; caudal 3–7 + 5–4; pectoral 13 + 3; ventrals I, 5; lateral line about 100; tube-bearing scales about 50. Body normal in form, rather robust; its greatest width ⅕ its greatest length without caudal; least height of tail contained 12 times, or nearly so, in the standard body length, and 3 times in height of body. The number of rows of scales, counting diagonally around the body from origin of anal, is 21 below and 7 above the lateral line. Width of head equal to its height; upper limb of orbit encroaching upon the upper profile of head, the center of the pupil equidistant from tip of snout and tip of the prolonged preopercular spine; length of snout measured obliquely from anterior margin of orbit equal to that of the postorbital portion of head to end of opercular spine; a robust spine at lower angle of preoperculum, curving slightly upward, the length equal to that of first dorsal ray; this spine serrated upon its outer edge, and with a small spine at its base which is also serrated; the tip of the spine extending to perpendicular from center of the interspace between third and fourth dorsal spines, while that of the humeral spine extends to the perpendicular from the interspace between the fourth and fifth, and that of the opercular to the perpendicular from the center of base of third; a strong scapular spine extends back to the posterior edge of the second dorsal spine. Length of upper jaw equal to ⅓ that of head. Palatine teeth in short feeble bands, hardly perceptible, even with a strong magnifying glass. Gill rakers 6, besides several rudimentary ones, 5 being below the angle, and the longest equal in length to ⅓ diameter of eye. First dorsal fin inserted above tip of upper opercular spine and at a distance from snout equal to twice length of the fourth dorsal spine, height of first dorsal spine, which is equal to that of the third and slightly less than that of the second, equal to ½ length of head, its anterior margin strongly serrated, while those of the second and third spines are less markedly so; length of base of first dorsal equal to greatest height of body; the distance between its insertion and that of second dorsal fin equal to the length of the longest and superior detached pectoral ray; second dorsal fin inserted in the perpendicular over the interspace between the second and third anal rays, the length of its longest ray equaling twice the least height of tail, and the length of its base equaling the greatest

length of the ventral rays, its first ray conspicuously serrated on its anterior edge. The insertion of the anal fin is in the perpendicular below the end of the first dorsal fin; the length of its longest rays is equal to ½ that of the middle caudal rays; caudal truncated, very slightly emarginate; pectoral very peculiar in structure, its longest ray, the ninth, reaching to base of the caudal rays and equal in length to 4 times that of the fourth dorsal spine; the tenth ray a trifle shorter, extending nearly to the end of the soft dorsal; the eleventh, twelfth, and thirteenth rays graduated, decreasing in regular proportion, the thirteenth being less than ¼ as long as the tenth; the eighth about midway between the tenth and the eleventh; the first slightly longer than the twelfth, and those intermediate between the first and the eighth are graduated in length, so as to form a rounded outline for the anterior or upper portion of the fin; the pectoral appendages slender, the third being slightly greater in length than the thirteenth ray, being ⅞ as long as the first, while the second is intermediate between the other two; the ventral inserted directly under the base of the pectoral appendages, its first spine about equal in length to the preopercular spine from the base of the supplemental spines, its longest, the third and fourth, exactly equal in length to the base of the second dorsal. Color brownish above, with about 4 indistinct transverse band-like blotches, 1 of which is on the base of the caudal, whitish beneath; vertical fins uniform, the tips of the caudal rays blackish, with 2 indistinct cloud-like bands in advance of the terminal bands thus formed; a black blotch, with whitish anterior margin on the membrane between the fourth and fifth dorsal spines; a very inconspicuous blackish spot on the membrane between the fifth and sixth; others still less conspicuous on the succeeding interspaces; the pectoral blotched and clouded with blackish brown and white. Off Charleston, South Carolina; 1 specimen obtained in the same haul with *Notosema dilectum*. (Goode & Bean.) It is a very well-marked species, distinguished especially by its very long pectoral fin. (*alatus*, winged.)

Prionotus alatus, Goode & Bean, Bull. Mus. Comp. Zool., xix, 210, 1883, deep sea off Charleston, South Carolina; Jordan, Cat. Fish. N. Am., 114, 1885; Jordan & Hughes, Proc. U. S. Nat. Mus. 1886, 332; Goode & Bean, Ocean. Ichth., 467, 1896.

Subgenus PRIONOTUS.

2488. PRIONOTUS MILES, Jenyns.

Head 2⅓; depth 4½; eye 4⅓ in head. D. X–12; A. 10 or 11; scales about 78. Body elongate, rather slender. Head moderate, not very rough; mouth moderate, maxillary scarcely reaching front of orbit, 2½ in head; interorbital area concave, narrow, 1⅔ in eye; no transverse postorbital groove; no spine on center of radiation of cheek, opercular spine rather weak, less prominent than the humeral spine, preopercular spine slightly larger than the humeral spine, no smaller one before it; all of the spines simple; preorbital ridge serrulate, no conspicuous postorbital ridge; temporal ridges present, all of the bones of the head with radiating striæ; snout emarginate, about 10 prominent spinules on each lobe, teeth on

vomer in 2 patches, on jaws and palatines in bands. Gill rakers rather long, about 10 below the angle, 3 or 4 tubercles on anterior part of arch. Pectoral fins long, reaching to middle of the anal fin; first dorsal spine weaker and shorter than the second, its length 2⅞ in the head, longest dorsal spine 2⅛ in head; ventrals reaching front of anal, 1⅛ in the head; first dorsal spine finely serrulate, other spines not serrated; caudal emarginate, 4½ in the length of the body. Color uniform brownish above, distal ¾ of spinous dorsal black; soft dorsal and caudal faintly barred; ventrals and anal white; pectorals blackish with white margins. Galapagos Islands; not rare. Here described from numerous fine specimens, 8 to 10 inches in length, from the Galapagos. (*miles*, a soldier.)

Prionotus miles, JENYNS, Zool. Beagle, Fishes, 29, pl. 6, 1842, Chatham Island, Galapagos Archipelago (Coll. Charles Darwin); GÜNTHER, Cat., II, 196, 1860.

2489. PRIONOTUS STEPHANOPHRYS, Lockington.

Head 2⅔; depth 4; eye 1⅛ in snout. D. X–12 or 13; A. 11 or 12; P. 13+3; scales 55. Body rather stout; head large, compressed, broad, and very nearly plane above, the interorbital region not concave; no transverse furrow behind orbits; orbital rim with a slightly raised, serrated crest; snout very short, ⅔ length of head; mouth large, the broad maxillary reaching beyond front of orbit; bands of palatine teeth very narrow; gill rakers rather long and slender; head less rough than usual, the bones little striate; occipital process not reaching first dorsal spine; preorbital little projecting; preopercle with a strong smooth spine, reaching slightly beyond membrane of opercle; opercle ending in 2 points, the lower a long spine, the membrane connecting the 2 scaly; scales thin, ciliate, not closely imbricate; first dorsal spine granulate in front; pectorals reaching beyond middle of anal, about to base of ninth ray; free rays very slender, the uppermost more than ½ length of fin; ventrals not reaching vent; caudal slightly emarginate. Pale olivaceous, abruptly white at level of pectorals; back and sides with a few scattered dark spots; branchiostegals mostly saffron yellow; spinous dorsal dusky, with diffuse dark blotches most distinct between fourth, fifth, and sixth spines; second dorsal spotted; caudal with dark blotches; pectorals dark, with large, round, black spots. Deep water off San Francisco, Point Reyes, Monterey, and Lower California; 4 specimens known. Here described from Mr. Lockington's type, and from 2 others collected by the *Albatross* at Station 3041, coast of Lower California. (στέφανος, crown; ὀφρύς, eyebrow.)

Prionotus stephanophrys, LOCKINGTON, Proc. U. S. Nat. Mus. 1880, 529, Point Reyes, near San Francisco; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 454; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 62; JORDAN & GILBERT, Synopsis, 736, 1883; JORDAN & HUGHES, Proc. U. S. Nat. Mus. 1886, 334.

2490. PRIONOTUS QUIESCENS, Jordan & Bollman.

Head 2½ to 2¾ (3⅛ to 3⅜ with caudal); depth 4 to 4⅓ (5 to 6); eye moderate, 4⅛ to 5 in head. D. X–12; A. 11; scales, in lateral line, 50 to 55; in a longitudinal series 60 to 70. Body rather slender, compressed, not much

narrowed above, the width of the nape between the occipital spines 4 to 4⅓ in head; head moderately elongate, not elevated; eyes not prominent, the profile from snout to nape almost straight or slightly convex; snout broad, of about equal length and breadth, 2½ to 2⅔ in head; anterior margin emarginate; preorbital little projecting, its edge with about 15 to 20 fine serræ, the anterior strongest with the exception of the last one, which is directed backward. Surface of bones of head smoother than usual in this genus and with fine radiating granular striæ, those on opercles, cheeks, and top of head most strongly developed. Mouth rather large, maxillary 2½ in head, reaching slightly beyond anterior orbital rim. Band of palatine teeth narrow. Orbital rim not especially elevated, its edges granulated, especially anteriorly, preorbital and postorbital spines small and blunt; interorbital not deeply concave, rather wide, its least width 4⅔ to 5 in head; no groove across top of head behind orbital rim; occipital ridges present, the inner very low, ending in a very small spine; outer large, ending in a moderate spine, the pair diverging, their inner edges serrulate, the spines extending to nearly opposite first dorsal spine; temporal region with a slight elevated roughish ridge, but no spine; preopercular spine long and sharp, its anterior edge somewhat serrulate, no smaller spine below it and none on suborbital stay; opercular and humeral spines well developed, sharp; no trace of spines on suborbital or preorbital. There are but 3 distinct spines on each side of the head, occipital, opercular, and preopercular. Membranaceous flap of opercle with a few scales. Gill rakers long and slender, about equal to ½ diameter of eye, 8 to 10 well developed. Scales quite small, those on breast (between ventrals) larger than those on belly or throat; scales extending beyond base of pectorals to isthmus; about 12 to 15 rows between occiput and front of dorsal. Spinous dorsal rather low; first spine not strongly serrulate, ⅔ length of second, which is 2½ in head; first ray of second dorsal weakly serrulate at base; longest ray shorter than snout and slightly less than 3 in head; longest anal ray 3½ in head; caudal lunate, 1½ to 1⅔ in head; pectoral generally reaching last dorsal ray, a little more than ½ body; ventrals reaching vent, 1⅔ to 1½ in head. Coloration in spirits, grayish, unspotted, more dusky above; spinous dorsal dusky, a distinct black spot between fifth and sixth spines; soft dorsal with 3 rows of diffuse spots; caudal dusky on the outer ⅔ and base; pectorals mostly black, with faint pale cloudings; ventrals and anal pale; body largely red in life. Related to *Prionotus stephanophrys*, Lockington, but the interorbital area concave; the bones of head much striate and granulated, and the caudal differently colored. It also bears some resemblance to the Atlantic species *Prionotus stearnsi*. Pacific Ocean, the types taken off the coast of Colombia; abundant in various places in 7 to 60 fathoms; also found at *Albatross* Station 3039, in the Gulf of California. (*quiescens*, resting quietly.)

Prionotus quiescens, Jordan & Bollman, Proc. U. S. Nat. Mus. 1889, 166, off Pacific coast of Colombia, at Albatross Stations 2800, 2801, 2802, and 2805. (Type, No. 41153, U. S. Nat. Mus. Coll. *Albatross*.)

2491. PRIONOTUS ALBIROSTRIS, Jordan & Bollman.

Head 3 in length ($3\frac{3}{4}$ with caudal); depth $4\frac{1}{2}$ ($5\frac{3}{4}$); eye rather large, $4\frac{1}{3}$ in head. D. X–12; A. 11; pores along lateral line 50 to 55. Body moderately elongate, little compressed, narrowed above, the width of the nape between the occipital spines being $4\frac{1}{3}$ in head; head rather short and high; eyes prominent, the anterior profile regularly concave, the eyes and forehead less prominent than in *P. rubio.* Snout broad, its breadth at angle of mouth almost equal to its length, $2\frac{1}{3}$ in head, its anterior margin not produced, but slightly emarginate; serræ short and even, bluntish, about 20 well developed; whole edge of preorbital with fine serræ. Anterior nostril with a large flap. Surface of bones of head with strong radiating striæ, those in front of eyes most broken up into granulations. Mouth moderate, maxillary $2\frac{3}{4}$ in head, not reaching front of eye; band of palatine teeth rather broad. Interorbital space narrow, deeply concave, smoother than rest of head, its least width $6\frac{1}{2}$ in head; orbital rim elevated, with coarse spine-like striæ in front, inner largest, forming the preocular spine; upper margin with moderate, strong serræ ending behind in a large, supraocular spine; no groove across top of head behind orbital rim; occipital ridges strong, the inner pair with a few asperities at base, ending in a compressed spine, the outer with stronger serræ at base and extending to opposite first dorsal spine; temporal ridge slightly crenulate, with 2 blunt spines; preopercular spine without a smaller one at base, but with a high sharp ridge before it, serrulate at base, this ridge not ending in a distinct spine; edge serrulate; suborbital stay with an elevated serrulate ridge, but no spine; opercular spine small and blunt, smaller than the strong humeral spine; no spines on suborbital or preorbital; membranaceous flap of opercle scaly. Gill rakers rather short, longer than interspaces, the longest rather less than $\frac{1}{2}$ eye, 5 most strongly developed. Scales small, those on belly smaller than those on breast, not extending before a line drawn between base of pectorals and ventrals; about 7 scales between occiput and dorsal; spinous dorsal moderately high, the first spine very strongly serrulate in front, shorter than second, which is $\frac{1}{2}$ head; first ray of second dorsal serrulate at base, the longest ray very slightly longer than snout; longest anal ray 3 in head; caudal subtruncate, $1\frac{1}{3}$ in head; pectorals long, reaching to the last dorsal ray or even farther in young specimens, $1\frac{1}{10}$ in body; ventrals reaching third anal ray, $1\frac{1}{4}$ in head. Coloration in spirits, grayish, unspotted, darkest above, and with darker cross shades; snout and jaws white; the tip of each jaw, a bar across each jaw, and 1 behind angle of mouth, black; a black bar on anterior and another on posterior part of interorbital, the latter extending across the cheek; first dorsal dusky, vaguely clouded with darker; second irregularly spotted, its posterior half dusky; caudal broadly black at base and tip, its middle part yellowish; pectorals dark, with a slight violet shade and traces of darker mottlings; middle of anal dusky; ventrals dusky on upper surface. Pacific coast of tropical America; taken at *Albatross* Station 3014, in Gulf of California; also known from several specimens, the largest $5\frac{1}{2}$ inches long, dredged at *Albatross* Station 2795, at a depth of 33 fathoms. It approaches *Prionotus quiescens* in technical

characters, but is very different in appearance; easily distinguished by the form, armature, and coloration of the head. (*albus*, white; *rostrum*, snout.)

Prionotus albirostris, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 168, Pacific Ocean off the coast of Colombia, at Albatross Station 2795, 7° 57′ N., 78° 55′ W. (Type, No. 41162, U. S. Nat. Mus.)

2492. PRIONOTUS RUBIO, Jordan.

(RUBIO VOLADOR.)

Head 3; depth 5. D. X–11; A. 10; scales about 57, pores 52. Body not very slender; gill rakers very short, tubercle-like, 9 or 10 developed, little if any longer than the interspaces; first dorsal spine nearly smooth; mouth not very large, the maxillary 2½ to 2¾ in head; interorbital space moderately concave, its width about ⅓ length of eye; no cirrus above the eye; distance from supraocular spine to nuchal scales about equal to eye; supraocular and nuchal spines low; occipital spines wanting; temporal ridge sharp, ending in a blunt spine; preorbital projecting, strongly serrate; a blunt spine on each side of snout, behind serræ of preorbital; a blunt spine behiud this above angle of mouth; no spine on cheek bone in adult; upper opercular spine almost obsolete; bones of head rather strongly striate, but not granulate; scales rather large, about 52 pores; first dorsal spine not much shorter than second, which is 2¼ in head; caudal very slightly concave; pectorals longer than in any other species (except *alatus*), reaching entirely past bases of dorsal and anal, its tip subtruncate, the longest ray about the ninth; ventrals reaching a little past vent. Color in life, dark olive, with rivulations of light green; sides shaded with pale salmon color; edge of pectoral light blue, ventrals reddish; upper fins marked with different shades of brown. From related species, *P. rubio* is well distinguished by its long pectorals, and by its very short gill rakers, much shorter than in any other species, *P. ophryas* coming nearest it in this respect. West Indies; not rare; our specimens from Cuba and Jamaica. (*rubio*, robin, the Spanish name.)

Rubio volador, PARRA, Descr. Dif. Piezas de Hist. Nat., 1787, lam. 38, Havana.
Prionotus punctatus, POEY, Synopsis, 304, 1868; POEY, Ennumeratio, 41, 1875; JORDAN & GILBERT, Synopsis, 956; not *Trigla punctata* of BLOCH.
Prionotus rubio, JORDAN, Proc. U. S. Nat. Mus. 1886, 50, Havana (Coll. Jordan); JORDAN & HUGHES, Proc. U. S. Nat. Mus. 1886, 334.

2493. PRIONOTUS OPHRYAS, Jordan & Swain.

Head 3 in length (3⅞ with caudal); depth 4½ (5⅝); eye 4½ in head. D. VIII–13; A. 11; scales (transverse series) about 75; pores in lateral line about 50. Body rather slender, narrower anteriorly and more compressed above than in other species, the width of the nape between the outer pair of occipital spines being not quite ¼ the length of the head; upper profile of head peculiar, being nearly straight from above front of eye backward, and steep and strongly concave from front of eye to tip of snout, the snout, therefore, steeper, more depressed, and rather shorter than in related species, its length being 2¼ in head; snout not very broad, its front

broadly rounded, its tip, as usual, emarginate, its edge with fine serræ directed backward, but no spines; surfaces of bones of the head covered with fine, sharply defined striæ, but with none of the small granulations which are found in *P. strigatus* and other species. Mouth rather wide, the maxillary reaching nearly to front of eye, the mandible quite to front of eye; maxillary 2½ in head; band of palatine teeth of moderate length, as long as eye. Eye large, placed high; upper part of eye with a fleshy cirrus rather enlarged toward the tip and fringed, this resembling the cirri in *Scorpæna*, its length about ½ that of the eye; interorbital area very narrow and very deeply concave, its least width little more than ⅛ the length of the head and not ⅔ the length of the eye; depth of interorbital area nearly ⅛ length of eye; bone forming anterior portion of orbital rim very prominent, forming a strongly striated crest, each of the striæ ending in a projecting point or spinule; upper portion of orbital rim prominent, even, ending behind the eye in a sharp backward-directed spine, behind which is a short cross groove, which does not extend across the top of the head; distance from the base of this spine to the scales on nape very short, not more than ½ the length of the eye; both pairs of occipital spines distinct, the outer and larger ones extending to opposite front of dorsal; a small spine on temporal region in front of outer pair of spines; preopercle with a single moderate spine at the end of a long ridge; no smaller spine at its base; opercle strongly striate, with 2 strong spines, of which the upper one is proportionately larger than usual; a single, rather strong humeral spine; membranaceous flap of opercle scaly. Gill rakers very short and thick, about 9 developed, these not ½ longer than the interspaces, and not ¼ length of eye; they are about ½ as broad as high, thus having a form very different from that seen in *P. evolans*, *P. strigatus*, *P. tribulus*, etc. Scales rather large, the scales on the back little reduced in size, about 10 between occiput and dorsal fin (17 in *P. strigatus*). Dorsal spines high and rather slender, the first rather the highest, its length 1½ in head, its anterior margin not granulated; soft dorsal rather high, its longest ray 2 in head; caudal 1⅓ in head; longest anal ray 2⅓ in head; pectorals rather long, extending nearly to last rays of dorsal, their length almost twice head; detached rays moderate, the uppermost or longest 1⅓ in head; ventrals 1⅓ in head. Coloration largely faded in the typical example; deep crimson in life; pale below; caudal with 2 dark cross bands; pectorals dusky, the free rays with dusky spots; ventrals pale, with some dusky bands; head nearly plain, the cirri dark. A second specimen, in very bad condition, has since been obtained by us from the same source. In this the undigested parts of the head and body are of a deep crimson. Probably all the deep-water species of this genus will be found to be red in life. Gulf of Mexico, in deep water; known only from the Snapper Banks, near Pensacola. Described from the original type, 7¼ inches long, taken from the stomach of a Red Snapper (*Neomænis aya*), from the Pensacola Snapper Banks, by Mr. Silas Stearns. (ὄφρυας, having projecting eyebrows.)

Prionotus ophryas, Jordan & Swain, Proc. U. S. Nat. Mus. 1884, 542, Snapper Banks off Pensacola (Type, No. 36944. Coll. Silas Stearns); Jordan, Cat. Fish. N. Am., 115, 1885; Jordan & Hughes, Proc. U. S. Nat. Mus. 1886, 334.

2494. PRIONOTUS STEARNSI, Jordan & Swain.

Adult example: Head 2⅘ in length (3⅛ with caudal); depth 4⅕ (5⅓); eye 5½ in head. D. VIII or IX–12; A. 12; scales in transverse series about 50; pores in lateral line about 52; length of specimen, 14½ inches. Body rather robust, formed much as in *P. tribulus;* width of nape between occipital spines not quite ⅓ length of head; head very large, broad, and unusually smooth, the profile, except for the prominence of the orbital region, forming a very gentle and somewhat regular arch; snout very broad, truncate at tip, its breadth at tip 3⅕ in head, its length 2⅔; edge of snout granular, without spines. Surface of all bones of head very finely, evenly, and regularly striated, the striæ much finer than in *P. tribulus,* their granulations all minute. Mouth wide, the maxillary reaching front of eye, its length about 2⅓ in head. Band of palatine teeth well developed. Eye large, placed high; no cirri; interorbital space about as in *P. tribulus,* rather broad and moderately concave, wider than the eye, and about 5 in head. Bone at anterior portion of orbital rim very prominent, serrulate, its principal ridge ending in a stout, blunt spine; supraorbital rim little prominent and without spine; no cross groove on top of head; a slight, bluntish spine behind eye; no spines on temporal ridge; outer pair of occipital spines short, strong, compressed, not quite reaching front of dorsal; inner pair and their ridges obsolete; opercular and preopercular spines short, the latter with no smaller one at its base; upper opercular spine very weak; humeral spine moderate; membranaceous flap of opercle with about 5 rows of scales; no spine on cheek bone. Gill rakers short and few, little longer than in *P. ophryas,* the longest about ⅓ interorbital width, about ½ longer than the interspaces, and perhaps 5 times as high as broad; about 9 gill rakers developed. Scales comparatively large, those on the back little reduced in size, about 10 before dorsal fin; 7 scales in a vertical row from first ray of soft dorsal to lateral line. Dorsal fins rather low and strong (the first injured), with its anterior margin not granulated; the third 2⅔ in head; soft dorsal moderate, its longest ray 3¼ in head; caudal very slightly concave, its longest rays 1⅓ in head. Pectoral fins rather short, reaching third ray of anal, 2½ in body, their length little more than length of head; detached rays tapering, a little more than ½ head; ventrals about reaching vent, 1¼ in head. Color in alcohol, nearly plain brownish olive, with dark shades at the bases of many of the scales, giving a mottled appearance; head everywhere conspicuously reticulate with blackish, in fine pattern; pectoral fin dusky, with a network of fine black cross streaks; dorsals similarly marked, the spinous dorsal with a diffuse black blotch between the fourth and sixth spines; caudal plain, slightly dusky; anal and ventrals pale; pectoral filaments nearly so. In the young the body is more slender, the snout less broad, with a slightly emarginate tip, eye larger, 4⅓ in head; no spine or groove behind the eye; gill rakers more slender, about 13 in number. Gulf of Mexico; in deep water. Known from 2 specimens, both taken on the Snapper Banks, off Pensacola, by Mr. Silas Stearns. The original type is a small specimen, not 4 inches long. The other is very large, about 13 inches long, larger than any other specimen of *Prionotus* which we have

ever seen. In spite of the remarkable differences in appearance of the two specimens, there is little reason to doubt their specific identity, as very similar differences distinguish the young and old of *P. tribulus*. According to Mr. Stearns, the large specimen was in life of a bright crimson red. Of all the species of the genus the present one has the spines of the head least developed, its upper surface being almost smooth. (Named for the late Silas Stearns, of Pensacola, long a volunteer assistant to the U. S. Fish Commission, and a most intelligent student of economic questions in marine ichthyology.)

Prionotus stearnsi, JORDAN & SWAIN, Proc. U. S. Nat. Mus. 1884, 541, Pensacola, young (Type, No. 36943. Coll. Silas Stearns); JORDAN, Cat. Fish. N. Am., 115, 1885; JORDAN, Proc. U. S. Nat. Mus. 1886, 228, adult; JORDAN & HUGHES, Proc. U. S. Nat. Mus. 1886, 335; GOODE & BEAN, Oceanic Ichthyology, 466, 1896.

2495. PRIONOTUS STRIGATUS (Cuvier & Valenciennes).

(NORTHERN STRIPED GURNARD; BROWN-WINGED SEA-ROBIN.)

Head $2\frac{2}{7}$; depth about 4. D. X-12; A. 11; scales 50 to 60 pores. Body rather robust; head moderately large; preopercular spine with a distinct smaller one at the base. Caudal fin very slightly lunate; pectoral fin subtruncate. Cheek bone without distinct spine at the center of radiation; edge of preorbital granular-serrate, without distinct spine, the serræ about 12 in number on each side; temporal ridges roughish, but without spines; bones of the head with the striæ coarsely granular; mouth moderate, the maxillary about $2\frac{3}{5}$ in head; head not very broad, the spines on its upper surface, except the nuchal spine, inconspicuous. Gill rakers longer and slenderer than in other species, 15 to 20 developed. Coloration brownish; side with a very distinct dusky bronze band below the lateral line and parallel with it, this becoming broken posteriorly into a series of roundish dark spots; some dark streaks and clouds below this stripe; fins with dark clouds, the soft dorsal with 2 dark blotches, which extend as bars on the back; head with scattered dark spots; dusky area below eye; pectoral with its rays each crossed by fine black bars, these especially distinct toward the base of the fin; free rays spotted; interorbital area broad and almost flat, its width a little more than length of eye; first dorsal spine granulated; second spine $2\frac{3}{4}$ in head; pectorals about $\frac{1}{2}$ the length of the body. Atlantic coast of the Northern States, Cape Cod to Virginia; very common in shallow water. The specimens here described from Marthas Vineyard. It is extremely close to *Prionotus evolans*, of which it may be a geographical variety. We have, however, as yet seen no intermediate examples, which should be looked for off the coast of Virginia. (*strigatus*, striped.)

Trigla lineata, MITCHILL, Trans. Lit. and Phil. Soc. New York, I, 1814, 430, pl. 4, fig. 4; not *Trigla lineata*, BLOCH.

Trigla strigata, CUVIER, Règne Animal, Ed. II, vol. 2, 161, 1829, New York; after *lineata*, MITCHILL.

Prionotus lineatus, DE KAY, New York Fauna: Fishes, 45, pl. 4, fig. 12, 1842; STORER, Synopsis, 50, 1846; GÜNTHER, Cat., II, 192, 1860.

Prionotus evolans var. *lineatus*, JORDAN & GILBERT, Synopsis, 736, 1883.

Prionotus strigatus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., 86, 1829; STORER Report Fish. Mass., 12, 1839; AYRES, Bost. Journ. Nat. Hist., iv, 1842, 258; JORDAN & GILBERT, Synopsis, 974, 1883, note; JORDAN, Cat. Fish. N. Am., 115, 1885; JORDAN & HUGHES, Proc. U. S. Nat. Mus. 1886, 335.

Prionotus evolans, GOODE & BEAN, Bull. Essex Inst., XI, 12, 1879; BEAN, Proc. U. S. Nat. Mus. 1880, 84; GOODE, Nat. Hist. Aquatic Animals, 255, pl. 71, 1884, and of writers; not *Trigla evolans*, LINNÆUS.

2496. PRIONOTUS EVOLANS (Linnæus).

(SOUTHERN STRIPED GURNARD.)

Head $2\frac{3}{5}$; depth $4\frac{3}{4}$. D. X–12; A. 11; lateral line with 53 pores; soft dorsal high, longest ray equaling longest spine, $2\frac{1}{2}$ in head; caudal $3\frac{1}{2}$ in length. Body and head stouter than in *Prionotus strigatus;* edge of preorbital granular-serrate, without distinct spine, the serræ about 12 in number on each side; mouth moderate; preopercular spine with a smaller one at its base; scales larger than in *Prionotus strigatus*, 8+1+21 in a vertical line from last dorsal spine to vent; interorbital space more deeply concave, its width in adult not quite length of eye; first dorsal spine nearly smooth; second spine 3 in head; pectorals a little more than $\frac{1}{3}$ of the body. Color in life, olive brown above, becoming light olive on sides, white below; back with 3 brown cross bars, the first under spinous dorsal, the second under first third of second dorsal, the third under its end, all of these bars extending downward and forward to lateral line, the posterior forming a brown blotch on base of last dorsal rays; back and sides with numerous small white spots, irregular in shape and size; these often wanting; a lateral line running in a narrow brown streak; distinct broad reddish-brown streak from humeral spine backward to opposite end of anal; traces of a narrow streak above this; branchiostegal membrane yellowish above; a dark-brown streak from angle of mouth to base of preopercular spine; opercle dusky brown without, deep reddish brown within; caudal with a light-brown bar at base, then a broad translucent bar, the terminal $\frac{1}{3}$ orange yellow, narrowly margined behind with white; spinous dorsal dusky, with a diffuse black blotch between fourth and sixth rays above; soft dorsal translucent brownish, without streaks of any kind; anal wine color, translucent at base and tip; ventrals light reddish; pectorals glaucous green within, the lower rays reddish, the upper white; the outer side dark greenish brown, unbarred, with a very narrow blue margin behind. This form is in some respects intermediate between *Prionotus strigatus* and *P. tribulus*. The color is in most particulars like that of *P. tribulus*, but the white spots on back and sides are much less numerous, or wholly wanting, and the brown bar backward from humeral spine is present, as in *P. strigatus*, and the dorsal fin is not barred; the gill rakers are, as in *P. strigatus*, slender and fine, 18 to 20 developed on lower limb; the spines on the head are not strong as in *P. tribulus*, that above orbit behind not conspicuously raised above surface of head. In 2 specimens from Beaufort, North Carolina, the pectorals are much lengthened, reaching nearly to base of caudal, but this seems to be here, as in *P. tribulus*, a very variable feature, as specimens from

Charleston have the pectorals but ⅘ length of body. The description of *Trigla evolans* given by Linnæus is of very little value, but the redescription of the type given by Dr. Bean leaves little doubt that it is this species. In this species the gill rakers are longer than in any other except its analogue, *P. strigatus*. Here described from specimens from Charleston and Beaufort. South Atlantic coast of United States; known only from North and South Carolina, where it is locally abundant. (*evolans*, flying out.)

Trigla evolans, Linnæus, Syst. Nat., Ed. xii, 493. 1766, Carolina (Coll. Dr. Alex. Garden); Bean, Proc. U. S. Nat. Mus. 1885, 204; description of Linnæan type.

Prionotus carolinus, Jordan & Gilbert, Proc. U. S. Nat. Mus. 1882, 615. Charleston; Beaufort (Coll. Jordan & Gilbert); Jordan & Gilbert, Synopsis, 974, 1883; Jordan & Swain, Proc. U. S. Nat. Mus. 1884, 541.

Prionotus evolans, Gill, Cat. Fish. East Coast N. Am., 21. 1873; Jordan & Gilbert, Proc. U. S. Nat. Mus. 1878, 374. Jordan & Gilbert, Synopsis, 735, 1883. Jordan, Cat. Fish. N. Am., 115, 1885; Jordan, Proc. U. S. Nat. Mus. 1884, 541; Jordan & Hughes, Proc. U. S. Nat. Mus. 1886, 336.

2497. PRIONOTUS PUNCTATUS (Bloch).

Head 2¼; depth 2¼; eye 6 in head. D. X-12; A. 12; about 50 pores in lateral line. Body stout; head large; preopercular spine with a smaller one at its base; pectorals reaching past middle of anal, their length not quite ½ the body; gill rakers rather long and slender, about 10 developed; maxillary 2½ in head; a bluntish spine on edge of snout behind the serræ; behind this 1 or 2 smaller ones, at least in the young; no spine on cheek bone; groove behind the eye evident; interorbital area rather narrow, concave; preocular, supraocular, occipital and nuchal spines rather prominent; dorsal spines high, the third 2⅔ in head; first spine not serrate; mouth large, maxillary 2⅔ to 2¾ in head, and reaching nearly to the eye; a small spine on center of radiation of cheek and 1 before it. Color nearly plain; spinous dorsal with dark clouds and without black ocelli; pectoral dark, with some round brown spots above; caudal dark barred; a whitish area on back between dorsals. Our description is taken from 2 small specimens collected (probably at Tuxpan) on the east coast of Mexico, by Mr. T. Salt; from the specimens in the museum at Paris, the types of Cuvier and Valenciennes, and from a specimen taken by the *Albatross* at Bahia. This species is certainly the *Prionotus punctatus* of Cuvier and Valenciennes, but it may not be the species figured by Plumier, to which Bloch has given the name of *Trigla punctata*. The figure of Plumier shows a bright-red body, with many small spots of a darker red, while red spots are scattered over all of the fins, except the spinous dorsal and the ventrals. In general form and in the armature of the head, so far as this is shown in the plate, Plumier's figure most resembles the present species, but the red color suggests a possibility that some of the deep-water species may have been intended. The present species corresponds better to the figure than any other yet known. Bloch's figure of *Trigla carolina*, which has been identified with *P. punctatus*, is almost certainly *P. tribulus*. West Indies and coast of South America; not known from the coasts of the United States. (*punctatus*, spotted.)

Trigla punctata, BLOCH, Ichthyol., pl. 353, 1793, Martinique; on a drawing by PLUMIER; CUVIER, Règne Animal, Ed. 2, vol. II, 161, 1829.

Prionotus punctatus, CUVIER & VALENCIENNES, Hist. Nat. Poiss.,* IV, 93, 1829; GÜNTHER, Cat., II, 193, 1860 (in part); JORDAN & HUGHES, Proc. U. S. Nat. Mus. 1886, 332; JORDAN, Proc. U. S. Nat. Mus. 1886, 544; JORDAN, Proc. U. S. Nat. Mus. 1890, 328.

2498. PRIONOTUS BEANII, Goode.

Head 3; depth nearly 4; greatest width 4; eye nearly 4 in head or 2 in snout. D. X–12; A. I, 10; P. 13+3; V. I, 5; snout 2 in head, equal to mandible. Least height of tail nearly $\frac{1}{5}$ length of head, and equal to short diameter of eye; width of interorbital space on the bone about $\frac{1}{3}$ length of snout; a furrow behind eyes which is interrupted on nape; snout produced into 2 short, obtuse, serrated points, flanked behind on each side by a short but stout compressed spine; behind this on snout another short spine; a short spine on cheek bone; anterior nostril in a tube which is produced posteriorly into a flap; posterior nostril in a large, shorter tube; preopercular spine, also, with a short spine at its base; preocular and supraocular spines present; a pair of postocular spines on each side, one in front of the other; a pair of blunt spines on occiput and another on nape; length of opercular spine, measured back to the edge of preopercle, equal to length of postorbital part of head; humeral spine well developed, its length nearly equal to that of preocular; maxillary not reaching front of eye, its length equal to postorbital part of head; mandible reaching about to vertical front of eye, its length nearly $\frac{1}{2}$ length of head. Teeth in narrow, villiform bands in jaws and on vomer and palate; 8 developed gill rakers on the anterior arch, the longest about $\frac{1}{2}$ as long as eye; 5 rudiments below and 2 above the developed rakers of the anterior arch, these mere tubercles scarcely raised above the general surface. Distance of dorsal from tip of snout a little greater than length of head; length of base of spinous dorsal nearly 3 times length of eye; first spine serrate on its anterior margin for the greater portion of its height, nearly as long as the second, its length $\frac{2}{3}$ length of head; third spine longest, its length $\frac{1}{2}$ length of head; last 2 spines very small; third spine also serrated along its anterior margin for the greater portion of its height; a very slight interspace between the 2 dorsals; length of first ray of soft dorsal equal to that of second spine of dorsal; length of last ray equal to $\frac{1}{2}$ length of spinous dorsal base; length of middle caudal rays 3 times length of eye; caudal slightly emarginate; origin of anal immediately under origin of soft dorsal; length of anal base $\frac{1}{4}$ distance from anal to tip of snout, the spine only about $\frac{2}{3}$ as long as first ray, its length $\frac{1}{2}$ length of maxilla; length of longest anal ray $\frac{1}{2}$ length of spinous dorsal base; ventral extending to origin of anal, its length equal to anal base; pectoral when extended reaching to the line connecting the fifth ray of dorsal with sixth ray of anal, this fin emarginate behind, its longest ray slightly longer than

* The types of Cuvier & Valenciennes in Paris seem to be the species usually called *Prionotus punctatus*. There is also a specimen in the museum, labeled, apparently in the handwriting of Valenciennes: "*Trigla punctata* nobis, Bl., 253. *Tr. carolina*, Bl., 252. *Prionotus evolans*, Lacèp. *Rubio volador*, Parra, tab. 38, du Brésil. Quoy et Gaimard, exp'n Freycinet." This specimen, 0.25 m. long, in good condition, is of the same species as the one described above. Longest dorsal spine $2\frac{5}{6}$ in head. Pores in lateral line 85 to 90.

head and more than twice length of longest dorsal spine, 7 rows of scales between lateral line and origin of second dorsal, 19 rows between lateral line and origin of anal, 50 pores in lateral line, and about 93 oblique rows of scales can be counted; scales on breast conspicuously smaller than the rest. Color in alcohol, light brownish yellow above, lighter below; spinous dorsal with a black blotch between the fourth and fifth spines, its length about ⅔ that of eye; the membrane connecting the spines of dorsal with faint dusky shades in several places; membrane connecting the last 3 or 4 rays of second dorsal slightly dusky; pectoral with 2 broad dark areas, separated and surrounded by lighter, the dark markings on the pectorals not taking the form of bands, but having their greatest length nearly parallel with the axis of the fish. Length 4½ inches. Trinidad. (Goode & Bean.) (Named for Dr. Tarleton H. Bean.)

Prionotus beanii. Goode, in Goode & Bean, Oceanic Ichthyology, 468, pl. cxii, fig. 383, 1896, off Trinidad, in 73 fathoms. (Type, No. 39318. Coll. *Albatross.*)

2499. PRIONOTUS TRIBULUS (Cuvier).

(Big-headed Gurnard.)

Head 2¼; depth 5 (3 to 4 in young). D. IX or X–12 or 13; A. 11; P. 13+3; scales 49 (tubes). Body robust; head shorter and broader, snout shorter, and bones more strongly striate than in *P. evolans;* interorbital space deeply concave; occipital and supraorbital spines very strong and much compressed; band of palatine teeth as long as eye; gill rakers shortish, 9 below angle; membranaceous edge of opercle scaly; preopercular spine with a smaller one at its base, which is high and sharp; cheek bone with a spine (small in the adult, larger in the young) at the center of radiation; temporal ridge with 2 bluntish spines; bones of the head very sharply striate; young with 4 sharp, knife-like spines on side of cheek and snout, in a line before the preocular spine, these nearly disappearing with age; maxillary about 2⅓ in head; sides without dark longitudinal stripe. Gill rakers slender in the young, becoming shorter and thicker with age, about 10 developed on lower part of arch; head broad, the spines on its upper surface very prominent, all of them more or less compressed and knife-like, especially in the young. Second dorsal spine 2⅓ in head; pectorals moderate, 2 in body in the adult, 2½ in the young. Coloration in life, light olive green, the head and body everywhere reticulated with dark olive green, in definite patterns, the dark lines on the head conspicuous, arranged in a series of curves and concentric circles; the dark streaks on the body mostly undulating and ascending backward; a diffuse band along side of bright orange; belly white; 2 faint diffuse dark bands downward and forward from soft dorsal, the hindmost ascending on the fin; a fainter band on spinous dorsal; spinous dorsal reddish, clouded with darker; a large dark blotch, not ocellated, between fifth and sixth spines; second dorsal translucent reddish, with darker spots; anal similar, paler, the spots almost obsolete; caudal reddish, with 3 darker bands; ventrals plain light reddish; pectorals light, clear green on the front side, grayish behind; with about 5 somewhat irregular dark cross

bands, the 3 median broadest and forked or Y-shaped above; upper edge of pectoral pale; pectoral appendages reddish, barred with darker. Young with soft dorsal, caudal, anal and ventral fins plain; spines proportionally longer and fins shorter. South Atlantic coast, from Long Island to Brazos Santiago; very common southward; our description chiefly from Galveston specimens, verified on others from Pensacola, Cedar Keys, Charleston, and Beaufort. A very abundant species, well distinguished from the others of the Atlantic by the greater development of the spines of the head. The young have these spines much larger and more compressed than the adult, and in the very young 3 or 4 strong knife-like spines are developed on each side of the snout, as in *P. horrens*. In very young examples the spine at the base of the preopercular spine is much larger than the latter. (*tribulus*, scraping, from the thorny head.)

Trigla carolina, BLOCH, Ichthyologia, 352, 1793, Carolina; not of LINNÆUS.
Trigla tribulus, CUVIER, Règne Animal, Ed. 2, vol. 2, 161, 1829, America.
Prionotus tribulus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., IV, 98, pl. 74, 1829, New York; Carolina; DE KAY, New York Fauna: Fishes, 48, 1842; GÜNTHER, Cat., II, 195, 1860; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1878, 373 and 374; GOODE, Proc. U. S. Nat. Mus. 1879, 111; GOODE & BEAN, Proc. U. S. Nat. Mus. 1879, 128; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 288; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 615; JORDAN & GILBERT, Synopsis, 735, 1883; BEAN, Cat. Fishes, London Intern. Exhibit., 49, 1883; JORDAN & SWAIN, Proc. U. S. Nat. Mus. 1884, 233; JORDAN, Cat. Fish. N. A., 115, 1885; JORDAN & HUGHES, Proc. U. S. Nat. Mus. 1886, 366.

2500. PRIONOTUS HORRENS, Richardson.

Head $2\frac{2}{3}$; depth $4\frac{1}{2}$; eye $5\frac{1}{2}$ in head. D. X–12; A. 10; scales about 100; gill rakers rather long, about 5 developed. Body rather stout; head large; interorbital area concave, $1\frac{1}{3}$ times diameter of the eye; bones of the head with strong, radiating striæ; a preocular and 2 postocular spines; occipital and temporal ridges not very prominent, each ending in a spine; opercular and humeral spines simple; preopercular spine with a smaller one at its base; nuchal spine present; snout emarginate, its lobes dentate, its length $2\frac{1}{4}$ in head; cheek with spine at point of radiation of striæ; a similar spine about $\frac{1}{3}$ distance to tip of snout; first dorsal spine nearly as long as second, its edge nearly smooth; the second the longest, its length 2 in head; edge of first dorsal ray smooth; longest dorsal ray $2\frac{1}{2}$ in head; longest anal ray $3\frac{1}{4}$ in head; caudal fin emarginate, $1\frac{3}{4}$ in head; pectorals rather short, their tips reaching about third anal ray, 3 in length of the body; ventrals not reaching anal by a distance equal to the diameter of the eye, $3\frac{3}{4}$ in length of body. Color, uniform brownish above, lighter below; spinous dorsal dusky; no distinct black blotch; soft dorsal and caudal fins irregularly barred; ventrals and anal white; pectoral dusky on basal third, the rest of the fin lighter, with an interrupted broad black transverse band across middle, a narrower one across the tip. Pacific coast of tropical America; not rare in shallow water; known from Mazatlan to the Galapagos. The specimens here described from Mazatlan and *Albatross* Station 3041, off Lower California. Young examples in the British Museum are almost exactly like the young of *P. tribulus*, differing chiefly in the still larger proportionate size of the knife-like spines on the head.

The following notes on the types of *Prionotus horrens* were taken by us in London: Three young specimen, allied to *P. tribulus*, but the spines still larger and more knife-like; first spine on edge of snout broad and serrate, 3 behind this progressively larger, then 2 large spines on preopercle, the posterior one the largest; 2 smaller ones on opercle, and 1 very large on the scapula; 2 sharp ones over each eye; 1 behind the eye; 2 on top of head and 2 on occiput. Mouth large, maxillary reaching front of eye, $2\frac{1}{3}$ in head; gill rakers long and slender, 5; scales small; pectorals short, 3 in body, reaching somewhat past second dorsal front; pectorals and tip of caudal dusky; no groove behind the eye; belt of palatine teeth narrow. (*horrens*, bristling (creating horror), from the large head spines.)

Prionotus horrens, RICHARDSON, Voy. Sulph., Ichth., 79, pl. 42, figs. 1–3, 1843, Gulf of Fonseca; GÜNTHER, Cat., II, 195, 1860.

793. BELLATOR, Jordan & Evermann.

Bellator, JORDAN & EVERMANN, Check-List Fishes, 488, 1896 (*militaris*).

This genus is closely allied to *Prionotus*, differing chiefly in the great development of the first and second dorsal spines, which are about as long as body. Scales large and very rough. Snout short, abruptly descending. West Indies. (*bellator*, warrior.)

a. Body robust; scales large, 40 tubes; snout with serrated process. MILITARIS, 2501.
aa. Body slender; scales moderate, 60 tubes; snout without processes. EGRETTA, 2502.

2501. BELLATOR MILITARIS (Goode & Bean).

Head 3; depth $3\frac{1}{5}$; eye $10\frac{1}{2}$, = interorbital width; snout $2\frac{5}{14}$. D. X–11; A. I, 9; P. 12 + 3; V. I, 5. Body short, stout, its greatest width at base of pectorals nearly $\frac{1}{4}$ of length. Head short, snout abruptly descending and with 2 rather long diverging spinous processes at its tip. Orbits much elevated, spines large, and jaws small; distance measured obliquely from tip of rostral spine to edge of opercular flap $2\frac{1}{2}$ in length; nearly all of the spines of head and exposed edges of preorbital, mandible, and opercles minutely serrate; the diverging spines upon snout themselves armed along margin by numerous spinules; a strong spine on preopercle, with a secondary spine at its base; the spine on preopercle as long as snout; a stout spine on operculum, another in humeral region, another on nape extending backward to base of fourth dorsal spine. Teeth in jaws, and on vomer and palate, very small, in villiform bands. Length of maxillary a little greater than that of eye; length of mandible a little less than that of snout, reaching about to the vertical from front of eye; a furrow across nape immediately behind eyes; 9 developed gill rakers on the anterior arch, besides several rudiments; all of the gill rakers very short. Pseudobranchiæ present. Branchiostegals 7. Distance of dorsal from tip of snout $2\frac{1}{2}$ in body; first 2 dorsal spines much produced; length of first almost equal to standard length; length of second slightly greater than that of first; the short spine about twice as long as mandible; when the dorsal

spines are fully extended they reach nearly to tip of caudal; anterior margins of first 3 spines minutely serrated, the serrations being in several rows; longest ray of soft dorsal about ½ as long as head; length of middle caudal rays equal to length of anal base; caudal slightly emarginate; length of pectoral a little less than 2 in body; longest separate ray of pectoral about 1½ times as long as shortest, its length equal to that of middle caudal rays; length of ventral spine ⅓ of length of longest ventral ray; ventral when extended reaching to third ray of anal; longest anal ray about ½ as long as head; scales very rough, in about 7 rows between origin of second dorsal and lateral line, and 19 rows below lateral line; lateral line with about 40 tubes, the number of rows of scales counted obliquely about 55. Color in life, rosy; head and pectoral speckled with dark brown; 6 or 7 small dark blotches on upper edge of pectoral. Some specimens have the inner surface of pectoral dark on its lower half. In 1 specimen the dark blotches on pectoral are grouped into 4 half bands, of which the middle 2 are very small. Off Cape Catoche, Yucatan, Gulf of Mexico. (Goode & Bean.) (*militaris*, like a soldier, from the high spines.)

Prionotus militaris, GOODE & BEAN, Oceanic Ichthyology, 464, pl. CXI, fig. 380, and pl. CXII, fig. 384, 1896, off Cape Catoche, Yucatan, in 25 fathoms. (Coll. *Albatross*.)

2502. BELLATOR EGRETTA (Goode & Bean).

Head 2⅜; depth 4⅓; eye 3½ in head; snout 2⅓; interorbital width 2 in eye. D. XI–11; A. I, 10; P. 12+3; V. I, 5; gill rakers *x*+9; maxillary nearly 3; mandible 2⅓; scales 9–100–32, 60 tubes. Body rather slender, tapering rapidly posteriorly; head moderately long; snout not descending so abruptly as in *P. militaris*, without projections in front. Spines of head moderately strong, without subsidiary basal spines; opercular and preopercular spines about equal in length, 2 in snout; exposed edges of bones of head minutely serrated, teeth at end of snout slightly enlarged; humeral spine small; nuchal spines not well developed; 1 or 2 spines at front of supraorbital, and 2 or 3 at its posterior portion. Teeth in villiform bands on jaws, and on vomer and palatines, the vomerine band very narrow. A slight groove across the nape immediately behind eyes, continued downward by an interspace between the preopercle and opercle. Distance from tip of snout to origin of dorsal equal to length of head; base of spinous dorsal equal to length of head without snout; first dorsal spine coarsely serrated for first sixth of its length, produced into a filament extending beyond end of caudal, the spine thus exceeding length of fish; second spine nearly as long as base of fin, the last 3 small; several succeeding spines also weakly serrated on their anterior margins; first ray of second dorsal serrated on anterior margin, the rays increasing in size to the ninth which is nearly ½ length of head; caudal somewhat emarginate, length of middle rays a trifle longer than base of spinous dorsal; origin of anal almost directly opposite that of second dorsal; length of anal spine about ⅔ that of first ray, the rays increasing in length posteriorly, the ninth being 2 in base of second dorsal; longest separate pectoral ray as long as base of second dorsal, the shortest 2 in head; ventrals reaching origin of anal. Color light brownish yellow on back, paler

beneath; fins, except pectoral and ventral, pale; pectoral with 5 dark bands, the third extending entirely across the fin. Known only from the type taken off Barbados between 100 and 200 fathoms. (*egretta*, an egret, in allusion to the elongate dorsal ray, resembling the plume of an egret.)

Prionotus egretta, GOODE & BEAN, Oceanic Ichth., 465, fig. 381, 1896, off Barbados, at Blake Station 64. (Type in Mus. Comp. Zool.)

794. CHELIDONICHTHYS, Kaup.

(SMALL-SCALED GURNARDS.)

Chelidonichthys, KAUP, Archiv f. Naturgeschichte 1873, 87 (*hirundo*).

This genus differs from *Prionotus* chiefly in the absence of palatine teeth. The scales are much smaller, and the pectoral fins less developed; a series of bony, spinous plates extends along the base of the dorsal fin, a pair of them to each ray, the fin thus running in a shallow groove; there are no plates along the lateral line; caudal fin usually emarginate. Lateral line usually forking at base of caudal, the branches running to tip of fin. The numerous species abound on the coasts of Europe, Africa, and India; ranging north to Japan. (*χελιδών*, swallow; *ἰχθύς*, fish.)

2503. CHELIDONICHTHYS PICTIPINNIS (Kaup).

The following description* is taken from Dr. Kaup's original type, kindly sent us from the museum in Hamburg by Dr. Georg Pfister, Curator:

Head $3\frac{1}{8}$ (to tip of rostral lobe); depth $5\frac{1}{6}$. D. IX–16; A. 15; pectoral 10 + 3 detached rays; scales about 68. Width of head $1\frac{5}{6}$ in its length; width of nape between occipital spines $3\frac{1}{2}$ in length of head; upper profile of snout straight. Snout long, a little more than twice in head; emarginate at tip, the preorbitals extending beyond it on either side; each preorbital with about 5 blunt spines. Mouth very wide, its greatest width (measured inside) $2\frac{1}{5}$ in head; maxillary reaching front of orbit, $2\frac{2}{3}$ in head. Vomerine teeth present in a small crescent-shaped patch, villiform; no teeth on palatines; a wide band of well-developed villiform teeth present on each jaw. Eye large, $4\frac{3}{5}$ in head. Interorbital space deeply concave, $1\frac{3}{5}$ in eye. Depth of caudal peduncle $1\frac{1}{2}$ in eye. Gill rakers 9, about $2\frac{1}{2}$ in eye. Bones of the head covered with radiating striæ which are comparatively fine, smooth and regular; preopercle with 2 spines at the angle, the upper much the larger; opercle ending in 2 spines; humeral spine long and strong; supraorbital rim anteriorly with 2 spines; a bony groove extending the entire length of the dorsals, formed by 24 plates, each of which ends on the side in a spine; a short, somewhat obscure transverse groove behind each eye, the two not continued across the occiput. Scales small, cycloid, present on back,

* Dr. Pfister sends the following note under date of January 2, 1896:
"Das Kaup'sche Original-Stück von *Trigla pictipinnis* war in unseren Katalogen nicht enthalten; erst durch längere Studien im Archiv ergab sich, dass das Stück, welches ich Ihnen nunmehr übersandt habe, thatsächlich das Original-Stück von Kaup ist; es stand im Museum als *Trigla hirundo*, Barbados, Ug. Ehrhardt."

sides, and belly; none on fins excepting on caudal, on which they extend nearly to tip; those in lateral line elongate, enlarged. Pelvic bone diamond-shaped, not scaled, 1⅔ in head, its width twice in its length; region of axil of ventrals and pectorals naked. Base of spinous dorsal 1½ in head, third spine 1¾ in head, all the spines smooth; base of soft dorsal 1⅛ times as long as head, longest soft ray 2½ in head; base of anal slightly greater than head; pectoral reaching about to seventh anal ray, 2⅔ in body; the upper detached pectoral ray the longest, 1⅛ in head, middle detached ray 1⅛ in upper, lower 1¼ in middle; ventrals reaching origin of anal, 1⅛ in head. Color (in spirits): back and upper half of sides reddish brown; lower portion of sides golden silvery; belly, anal, base of pectorals, and base of ventrals white; head light chocolate brown; dorsals, caudal, ventrals, and detached rays of pectoral translucent, the membrane of pectorals satin-like and very dark brown, the rays white; 6 or 7 conspicuous, small white spots on the inside of pectoral on its lower half. Type 13 inches long, in the museum at Hamburg, said to have been sent from Barbados. This specimen agrees perfectly with descriptions of *Chelidonichthys kumu* (Lesson & Garnot) from New Zealand. It probably came from the South Seas, not from Barbados, and should not be admitted in this work. *Chelidonichthys spinosus* McClelland, of which we have specimens from Japan, although closely related, is a different species.

The following is the substance of Dr. Kaup's account:

Pectoral reaching seventh or eighth anal ray; dorsal spines smooth, reddish brown above, yellow on sides; belly white; pectoral colored on the inner side as in *Trigla peronii* and in *Trigla kumu* ("a large black blotch and white spots in variable number"). First dorsal uniformly colored; 4 black spots along median line of back, and 2 spots on side, parallel with the first 2 dorsal spots; a black spot before the first dorsal spine; a similar spot below the hinder half of the eye. One specimen a foot long in the Hamburg Museum from Barbados. (Kaup.)

The subgenus *Chelidonichthys* to which it is referred is thus described:

Preorbital plate of the steeply truncate snout somewhat prominent, with blunt teeth; spines of preopercle and opercle short; first dorsal with slender spines, the anterior entirely smooth or somewhat rough; first dorsal with 9 or 10; second with 1 spine and 16 soft rays; anal with 15 or 16 soft rays; scales very small; lateral line somewhat prominent, without spines. The species reach a length of 8 inches to a foot or more. (*pictus*, painted; *pinna* fin.)

Trigla pictipinnis, Kaup, Arćhiv für Naturgeschichte 1873, 87, Barbados? (Coll. Ehrhardt. Type, Mus. Hamburg.)

795. TRIGLA (Artedi) Linnæus.

(Mailed Gurnards.)

Trigla (Artedi) Linnæus, Syst. Nat., Ed. x, 300, 1758 (*cuculus*).

This genus differs from *Chelidonichthys*, with which it agrees in the absence of palatine teeth, in having the sides of the body armed with transverse bony plates, crossing the lateral line. Species numerous; very

abundant in the Mediterranean. (*τρίγλα, Trigla,* classical name of *Mullus barbatus,* still used by the fishermen of the Adriatic, transferred to this group by Artedi for no evident reason.)

2504. TRIGLA CUCULUS, Linnæus.

(Red Gurnard.)

Head $3\frac{1}{4}$; depth $5\frac{1}{5}$. D. IX–17; A. 16; plates along lateral line 65; eye $5\frac{3}{5}$ in head; second dorsal spine $1\frac{1}{3}$; pectoral $1\frac{1}{10}$; anal $1\frac{1}{3}$; caudal $1\frac{1}{2}$. Body not anywhere compressed; head everywhere covered with rough, bony plates; mouth moderate, with bands of villiform teeth on jaws and vomer; maxillary scarcely reaching to front of eye; snout truncate, with 2 or 3 spines on each side; eyes placed high; a couple of spines on supraorbital rim above anterior edge of eye; interorbital deeply concave; posttemporal, with its upper surface rugose, ending as a spine behind; a couple of spines on opercle in front of the flap; a spine on clavicle just above pectoral fin; a series of long, narrow plates along lateral line; back and sides covered with small scales, belly naked; a ridge of about 26 spines extending along base of dorsals, ending at posterior end of soft dorsal; spinous dorsal triangular in outline, the second spine the longest, spines rapidly decreasing in length behind it, second spine reaching to soft dorsal when depressed; pectoral long and narrow, reaching to front of anal; ventrals inserted a distance of $\frac{1}{2}$ eye behind lower end of base of pectoral, reaching nearly to tip of pectorals; caudal long and emarginate. Color rose-red on back, white below; inner face of pectoral blackish, the outer face slightly dusky; other fins dusky. Southern Europe; said by Cuvier to have been once brought from New York by Milbert; a very doubtful record. No recent collector has found any species of *Trigla* in American waters. The specimen here described is from the Paris market. (*cuculus,* cuckoo.)

Trigla tota rubens, Artedi, Genera Piscium, 45, 1738.

*Trigla cuculus,** Linnæus, Syst. Nat., Ed. x, 301, 1758, Mediterranean, open sea; after Artedi.

Trigla pini, Bloch, Ichthyol., pl. 355, 1793; Günther, Cat., II, 199; Cuvier & Valenciennes, Hist. Nat. Poiss., IV, 26; De Kay, New York Fauna: Fishes, 43, pl. 70, 1842; Day, Fish. Gt. Brit., 58, 1880.

Family CLXXXV. PERISTEDIIDÆ.

(The Deep-water Gurnards.)

Body elongate, fusiform, covered with bony plates, each of which is armed with a strong spine; head bony; each preorbital produced into a long, flat process, which projects more or less beyond the mouth; mouth small, inferior, like that of a sturgeon; teeth none; lower jaw provided with barbels; gill membranes separate, narrowly joined to the isthmus

* *Polynemus tridigitatus,* Mitchill, Trans. Lit. and Phil. Soc., I, 449, 1815, New York (no description), referred by Dr. Günther to the synonymy of *Trigla cuculus,* is apparently based on a recollection of some species of *Prionotus.* It is certainly not a *Trigla.*

anteriorly; gill rakers slender. Dorsal fin continuous or divided. Pectoral fin short, with the 2 lowermost rays detached. Ventrals I, 5, separated by a broad, flat area. Air bladder simple. Pyloric cæca about 10. Color generally red. Deep-sea fishes; 2 or 3 genera and about 13 species known, bearing some resemblance to young sturgeons.

a. Barbels at angle of mouth in large tufts of fringes; dorsal fins 2. PERISTEDION, 796.
aa. Barbels at angle of mouth minute, simple or nearly so; dorsal fins 2. VULSICULUS, 797.

796. PERISTEDION, Lacépède.

Peristedion, LACÉPÈDE, Hist. Nat. Poiss., III, 368, 1802 (*malarmat=cataphractum*).
Peristethus, KAUP, Proc. Zool. Soc. Lond. 1859, 103 (*cataphractus*); amended spelling.

Barbels large, forming large fringed tufts at angles of mouth and on lower jaw. Dorsal fins 2; characters otherwise included above. (*περί*, around; *στηθίον*, diminutive of *στῆθος*, breast, the breast being mailed.)

a. Body rather robust, the depth 4½ to 5 in length to base of caudal; head about 2⅓ in length; barbels long.
b. Length of preorbital extension about 3½ in snout; color uniform crimson. MINIATUM, 2505.
bb. Length of preorbital extension 2 in snout; color red, with a black blotch near tip of pectoral, and black on caudal. LONGISPATHUM, 2506.
aa. Body slender, the depth 6 to 6½ in length to base of caudal; head 3 to 3½; barbels moderate; fins mottled or blotched.
c. Body very slender; preorbital process 2½ in snout; color yellowish; a pearly lateral band; back dotted; fins mottled. GRACILE, 2507.
cc. Body much depressed; preorbital process 3 in snout; color red, mottled and blotched. PLATYCEPHALUM, 2508.

2505. PERISTEDION MINIATUM, Goode.

Head 2½; depth 5. D. VII–18; A. 17; C. 16; P. 10+2; V. I, 5; plates 27 to 29. Armature of body essentially as in *Peristedion brevirostre*;* spines of abdominal plates very weak posteriorly; length of preorbital process 3½ in distance from its extremity to front of orbit; interorbital space deeply concave; protuberance on the forehead very slight; a pair of spines on upper surface of snout behind base of preorbital processes; a larger pair on preorbital processes; ridge of preopercle ending in a depressed, short, sharp-pointed spine; about 10 small tentacles on each side of lower jaw, those near the symphysis smallest. Long tentacles at angle of mouth fringed, extending to bases of pectorals. Length 12 inches. Gulf Stream, in various localities, at a depth of about 150 fathoms. (Goode.) (*miniatus*, scarlet.)

Peristedion miniatum, GOODE, Proc. U. S. Nat. Mus. 1880, 349, Gulf Stream, off Rhode Island; JORDAN & GILBERT, Synopsis, 733, 1883; GOODE & BEAN, Oceanic Ichthyology, 470, pl. CXIII, figs. 385, 385A, B, 1896.

2506. PERISTEDION LONGISPATHUM, Goode & Bean.

D. VIII–19; A. 19. Body high anteriorly, its greatest height 4½ in total length; length of head without prolongations 2½ in total length, with prolongations 2. Crown of head flat, separated from nuchal plate by a deep

* See Günther, Cat., II, 218, 1860.

furrow, which is convex forward. Interorbital space deeply concave, the supraorbital margins being swollen, its width equaling long diameter of orbit. No protuberance on forehead, which is much depressed, its outline descending abruptly and rapidly in front of eyes; a ridge but no spine beneath eye. Length of snout, including the preorbital extension, more than ½ length of head; preorbital extension equaling ½ length of snout; the processes flat, rounded anteriorly, and covered with minute granulations; they diverge considerably, the distance of the tips apart being nearly twice that at their bases; a ridge arises at base of preorbital process and extends to angle of preoperculum, and its width at the angle is contained twice in diameter of orbit; a narrow, inconspicuous, and interrupted ridge below; a ridge on operculum, ending in a sharp spine at the angle, its length equal to diameter of eye. Jaws feeble, toothless; lower jaw with 2 long, much fringed barbels and 14 shorter ones; length of the long barbels twice diameter of the eye; maxillary not reaching vertical through anterior margin of eye; diameter of eye 4 in length of head without its prolongations; greatest width of head over preopercular ridge, 3 in total length. Dorsal origin directly in a line with upper angle of gill opening; longest spine slightly longer than width of interorbital space; anal origin under that of second dorsal; caudal small, slightly emarginate, length of its middle rays 1½ times diameter of eye; ventrals slightly in advance of pectorals, and extending farther back, reaching slightly beyond vent and to vertical through seventh row of scales; pectoral short, extending to vertical from fifth scale of the lateral line, the longest detached ray to the sixth; 29 rows of scales. Color in life bright roseate; a black blotch near tip of pectoral; dorsal with narrow dark margin; tip of caudal black. The elongation of the preorbital extension is noticeable in the smallest examples. (Goode & Bean.) West Indies, off Cuba and Barbados. (*longus*, long; *spatha*, sheath.)

Peristedium longispatha, Goode & Bean, Bull. Mus. Comp. Zool., xii, No. 5, 166, 1886, off Santa Cruz, in 314 fathoms (Coll. Blake); off Barbados, 209 fathoms (Coll. Blake), and at other stations in the Gulf of Mexico (Coll. *Albatross*); Goode & Bean, Oceanic Ichthyology, 472, pl. cxiv, fig. 386, 1896.

2507. PERISTEDION GRACILE, Goode & Bean.

B. 7 or 8. D. VIII–20; A. I, 18; P. 11 + 2; V. I, 5. Body very slender, its greatest height nearly ⅙ of length. Length of head to tip of the prolonged rostral spine 48 mm.; without the prolongations its length is 36 mm. or twice the greatest height of the body. Interorbital space deeply concave; interorbital width ⅓ length of snout, including rostral spine. Forehead descending gently toward snout, the supraorbital crests very little elevated. The rostral spine on each side continued backward by a slight bony ridge ending in a blunt spine at the angle of preoperculum. Opercular spine small, the length of operculum with the spine being about equal to length of eye. Length of snout with preorbital extension a little less than 4 in body; length of spine alone nearly equal to length of mandible. Jaws feeble and toothless; length of maxillary ⅓ length of head; length of mandible ½ length of snout with preorbital extension;

mandible with 2 groups of barbels, 1 on each side of its lower edge; length of longest barbel equaling that of eye; barbels subdivided into several fringes, the number of barbels on each side of the mandible may be divided up into 8 or 9 clusters; mandible extending to vertical through front of eye; 26 gill rakers on first arch, the longest half as long as eye. Longitudinal diameter of eye nearly equal to width of interorbital space; greatest width of head over preorbital ridge $\frac{1}{2}$ length of snout including the preorbital extension. Spinous dorsal originating directly above end of the opercular flap; the first spine slightly the longest, its length equaling length of snout and its projecting spine, also short diameter of eye; longest ray of second dorsal nearly $\frac{1}{4}$ length of head; least height of tail $\frac{1}{3}$ length of eye. Anal origin under second ray of soft dorsal, the fin extending as far back as the dorsal, the length of its base being 4 times the width of interorbital area, the rays about as long as those of second dorsal; length of middle caudal rays $\frac{2}{3}$ length of eye, the fin being emarginate; ventral base under pectoral base, the fin reaching to vent, its length twice that of eye; pectoral short, reaching to below eighth scute of the lateral line, the longest detached ray reaching to below the ninth scute of the lateral line, its length equaling $\frac{1}{2}$ length of head; 30 scutes in the lateral line. Color (of alcoholic specimen) very light yellow, a broad pearly band along the sides; back stippled with light brownish; pectorals with dark mottlings. Gulf of Mexico. Known from 1 specimen, 5 inches long. (Goode & Bean.) (*gracilis*, slender.)

Peristedion gracile, Goode & Bean, Ocean Ichthyology, 473, pl. cxiv, fig. 387, 1896, Gulf of Mexico, in 142 fathoms, 28° 28′ 30″ N., 85° 52′ 30″ W., at Albatross Station 249.

2508. PERISTEDION PLATYCEPHALUM, Goode & Bean.

D. VIII-17; A. 17. Body much depressed, its greatest height $6\frac{1}{3}$ in body length, $6\frac{3}{4}$ in total. Length of head without prolongations, twice the height of body, $3\frac{1}{2}$ in its length, with prolongations $2\frac{1}{2}$ in body length. Interorbital space deeply concave, the supraorbital margin being swollen, its width equal to the long diameter of the eye. No protuberance on the forehead, which is much depresssd, its outline descending abruptly and rapidly in front of the eyes. A ridge below the eye, not armed; a small vertical spine behind each nostril. Stout spines upon the operculum and several upon the vertex. The length of the snout, with its extensions, is $\frac{1}{2}$ the length of the head, its processes about 3 in its own length. The processes are flat, triangular, diverging slightly, the distance apart of their tips 2 to $2\frac{1}{2}$ that at their bases. A ridge extending backward from base of each process along the lower edge of the preoperculum, ending behind in a sharp, flat spine; the greatest width of the expanded portion, on the preoperculum, only $\frac{1}{4}$ as wide as the eye; beneath this another less conspicuous ridge with minutely serrated edge, which is double in front and single behind, the 2 portions separated by a slight notch. Jaws normal, the 2 tentacles much fringed, their length not much exceeding the diameter of the eye; between them, and placed about equidistant from each, are 2 bunches of short tentacles, about 4 in each. Chin with numer-

ous short tentacles, some of them as long as the eye, arranged for the most part in bunches of 4. Maxillary not reaching anterior margin of orbit. Diameter of eye nearly 4 in greatest length of head, and exactly ⅓ of total length of snout. Greatest width of head, over the preopercular ridges, nearly equal to its own length without the processes. Dorsal origin over the upper angle of gill opening; longest spine equal to that of postorbital portion of head; anal origin about under origin of second dorsal, a trifle farther back, and in the vertical through the space between the seventh and eighth lateral scutes; the fin about as high as the dorsal; caudal small, slightly emarginate, with tips slightly produced, length of middle rays equal to that of dorsal; ventral origin in advance of the axil of the pectorals, the fin extending slightly beyond vent, but not quite to origin of anal, its length about twice length of dorsal. Pectoral rather long, extending to ninth scute of lateral line and past vertical through origin of anal; 29 rows of scutes. Color red; body and fins mottled and blotched with darker. Known from 2 specimens taken off Barbados, West Indies. Type from *Blake* Station LX, in 123 fathoms; the other from 288 fathoms. (Goode & Bean.) (*πλατύς*, broad; *κεφαλή*, head.)

Peristedion platycephalum, Goode & Bean, Bull. Mus. Comp. Zool., xii, No. 5, 167, 1886, off Barbados in 123 fathoms (Coll. the *Blake*); Goode & Bean, Oceanic Ichthyology, 474, pl. cxiv, fig. 388, A, B, 1896.

797. VULSICULUS, Jordan & Evermann.

Vulsiculus, Jordan & Evermann, Check-List Fishes, 489, 1896 (*imberbe*).

This genus differs from *Peristedion* chiefly in the reduction of the barbels on the lower jaw and the angle of the mouth. These are minute and simple, or nearly so, and scarcely appreciable. (*vulsus*, plucked; hairless.)

2509. VULSICULUS IMBERBIS (Poey).

Body somewhat slender, its greatest height 4½ times in distance between tip of snout and base of caudal. Length of head without prolongations 2½ in total length; with prolongations, 2. Crown of head flat; interorbital space concave, with a depressed groove in its middle, branching posteriorly along the base of the supraorbital crests, its width somewhat greater than the horizontal diameter of orbit. No protuberances on forehead or on the snout above, and no ridges or spines beneath eye. Length of snout, including the preorbital extension, equal to length of head; preorbital extension about ⅔ of length of snout. Preorbital processes flat, unarmed, and somewhat divergent; a ridge arising at base of preorbital process and extending to angle of preoperculum, where it terminates in a blunt spine; a low ridge on operculum, ending in a rather inconspicuous spine; another above and close to it, pointing upward. Length of opercular ridge of spines nearly equal to horizontal diameter of eye. Jaws feeble and apparently toothless; barbels on lower jaw so inconspicuous as to be scarcely visible; maxillary extending almost to vertical through anterior margin of eye; diameter of eye 4 times in length

of head without its prolongations. Dorsal origin directly in line with upper angle of gill opening; the origin of anal under that of second dorsal; ventrals slightly in advance of pectorals, and not so far back, though their tips reach somewhat beyond vent; pectorals rather long; about 25 rows of scales. Gulf of Mexico. The type, about 2 inches long, is in the Museum of Comparative Zoology at Cambridge, and has been examined in the preparation of this description, but the example being unique and small, we have not ventured to examine it so closely as would have been practicable with more material. It is, however, apparently a well-marked species, resembling in a general way *P. longispathum*. It was obtained by Poey from the stomach of *Polymixia nobilis*, taken near Cuba in deep water. (Goode & Bean). Near Cuba, in deep water, from stomachs of *Polymixia* (Coll. Poey). A few specimens in bad condition were taken by Jordan & Stearns from stomachs of Groupers and Snappers on the Snapper Banks off Pensacola. In these specimens, which seem to represent Poey's *imberbis*, the dorsal rays are about V–19, scutes 30; each side of body with 4 rows of spinous plates, the ventral series with hooked spines. (*imberbis*, beardless.)

Peristedion imberbe, Poey, Mem. Hist. Nat. Cuba, II, 367, 389, 1861, Cuba; Poey, Rep. Fish. Nat. Cuba, II, 158, 1866; Jordan, Proc. U. S. Nat. Mus. 1884, 38; Goode & Bean, Oceanic Ichthyology, 472, 1896.

Peristedion micronemus, Poey, Ann. Lyc. Nat. Hist. N. Y., IX, 1870, 321, Cuba; substitute name on finding the existence of barbels.

Peristethus micronema, Günther, Challenger Report, Deep Sea Fishes, XXII, 65, 1887.

Family CLXXXVI. CEPHALACANTHIDÆ.

(The Flying Gurnards.)

Body elongate, subquadrangular, tapering behind; head very blunt, quadrangular, its surface almost entirely bony; nasals, preorbitals, suborbitals, and bones of top of head united into a shield; nuchal part of shield on each side produced backward in a bony ridge, ending in a strong spine, which reaches past front of dorsal; interocular space deeply concave; preorbitals forming a projecting roof above the jaws; preopercle produced in a very long rough spine; cheeks and opercles with small scales; opercle smaller than eye; gill openings narrow, vertical, separated by a very broad, scaly isthmus; pseudobranchiæ large; gill rakers minute; mouth small, lower jaw included; jaws with granular teeth; no teeth on vomer or palatines; scales bony, strongly keeled; 2 serrated, knife-like appendages at base of tail; first dorsal of 4 or 5 rather high flexible spines, the first 1 or 2 spines nearly free from the others; an immovable spine between the dorsals; anal and second dorsal short, of slender rays; caudal small, lunate; pectoral fins divided to the base into 2 parts, the anterior portion about as long as the head, of about 6 rays, closely connected; the posterior and larger portion more than twice length of head, reaching nearly to caudal in the adult (*Dactylopterus*); much shorter in the young (*Cephalacanthus*); these rays very slender, simple, wide apart at tip; ventral rays I, 4, the long fins pointed, their bases close together,

the inner rays shortest; air bladder with 2 lateral parts, each with a large muscle; pyloric cæca numerous; vertebræ 9+*13=22. Warm seas; the adult able to move in the air like the true flying-fish, but for shorter distances. One genera and 2 to 4 species.

798. CEPHALACANTHUS, Lacépède.

(Flying Gurnard.)

Cephalacanthus, Lacépède, Hist. Nat. Poiss., III, 323, 1802 (*spinarella*); young examples of the East Indian species.

Dactylopterus, Lacépède, Hist. Nat. Poiss., III, 325, 1802 (*pirapeda*=*volitans*); adults of the American species.

Gonocephalus, Gronow, Cat. Fish., Ed. Gray, 106, 1854 (*macrocephalus*=*volitans*).

Character of the genus included above. Two species known, the following and the East Indian, *Cephalacanthus spinarella*. (κεφαλή, head; ἄκανθα, spine.)

2510. CEPHALACANTHUS VOLITANS (Linnæus).

(Flying Robin; Bat-fish; Volador; Murciélago.)

Head 4⅓; depth 5½. D. II–IV, 8; A. 6; P. 28+6. First 2 dorsal spines free, slightly connected by membrane at base; preopercular spine reaching beyond base of pectorals, not to end of occipital spine; pectorals reaching nearly to base of caudal in adult, very much shorter in young; in the young the spines of the head are much longer. Greenish olive and brown above, of varying shades; below pale, marked irregularly with dusky and bright brick red, varying to salmon yellow; pectoral fins mottled with bright-blue streaks near the base and blue spots and bars toward the tip; their under sides glaucous blue, edged with darker; caudal fin with about 3 brownish-red bars; coloration extremely variable. Length 12 inches. Atlantic Ocean, on both coasts; very abundant on South Atlantic and Gulf coasts; a handsome and singular fish. (Eu.) (*volitans*, flying.)

Pirabebé, Marcgrave, Hist. Brasil., IV, 162, 1648, Brazil.

Milvus cirratus, Sloane, Hist. Jamaica, II, 288, Jamaica.

Trigla digitis vicenis palmatis, Artedi, Genera, 44, 1738, Mediterranean, etc.

Hirundo, Catesby, Nat. Hist. Carolina, II, tab. 8, Bahamas.

Trigla volitans, Linnæus, Syst. Nat., Ed. X, I, 302, 1758; after Artedi; "Mari Mediterraneo Oceanio, Pelago inter tropicos, in Asia ad Cap. b. Spei. Sæpe agitata evolans ex aqua."

Dactylopterus volitans, Cuvier & Valenciennes, Hist. Nat. Poiss., IV, 117, 1829; Günther, Cat. Fishes, II, 221, 1860; Lütken, Spolia Atlantica, 417, 1880.

Trigla tentabunda, Walbaum, Artedi, Piscium, III, 362, 1792; after *Cataphractus*, Klein, Missus, which is after Catesby, Fishes of Carolina, IV, 44, taf. 14, f. 1.

Trigla fasciata, Bloch & Schneider, Syst. Ichth., 16, tab. 3, f. 1, 1801; after *Corystion*, Klein, Missus, IV, 45, taf. 14, f. 2, locality not stated.

Dactylopterus pirapeda, Lacépède, Hist. Nat. Poiss., III, 326, 1802, Mediterranean and almost all warm seas.

Polynemus sexradiatus, Mitchell, Trans. Lit. and Phil. Soc., I, 1815, pl. 4, f. 10, New York.

Callionymus pelagicus, Rafinesque, Amer. Monthly Mag., Jan., 1818, 205, Atlantic Ocean.

Dactylopterus communis, Owen, Osteol. Cat., I, 56, 1851.

Gonocephalus macrocephalus, Gronow, Cat. Fish., Ed. Gray, 106, 1854, pelagic.

* The osteological characters of this family are given (after Gill) on page 2147.

END OF PART II.

SMITHSONIAN INSTITUTION.
UNITED STATES NATIONAL MUSEUM.

BULLETIN

OF THE

UNITED STATES NATIONAL MUSEUM.

No. 47.

THE FISHES

OF

NORTH AND MIDDLE AMERICA:

A DESCRIPTIVE CATALOGUE OF THE SPECIES OF FISH-LIKE VERTEBRATES FOUND IN THE WATERS OF NORTH AMERICA, NORTH OF THE ISTHMUS OF PANAMA.

BY

DAVID STARR JORDAN, Ph. D.,
PRESIDENT OF THE LELAND STANFORD JUNIOR UNIVERSITY AND OF THE CALIFORNIA ACADEMY OF SCIENCES,

AND

BARTON WARREN EVERMANN, Ph. D.,
ICHTHYOLOGIST OF THE UNITED STATES FISH COMMISSION.

PART III.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1898.

SMITHSONIAN INSTITUTION.

UNITED STATES NATIONAL MUSEUM.

THE FISHES

OF

NORTH AND MIDDLE AMERICA:

A DESCRIPTIVE CATALOGUE

OF THE

SPECIES OF FISH-LIKE VERTEBRATES FOUND IN THE WATERS OF NORTH AMERICA, NORTH OF THE ISTHMUS OF PANAMA.

BY

DAVID STARR JORDAN, PH. D.,

PRESIDENT OF THE LELAND STANFORD JUNIOR UNIVERSITY AND OF THE CALIFORNIA ACADEMY OF SCIENCES,

AND

BARTON WARREN EVERMANN, PH. D.,

ICHTHYOLOGIST OF THE UNITED STATES FISH COMMISSION.

PART III.

WASHINGTON:

GOVERNMENT PRINTING OFFICE.

1898.

TABLE OF CONTENTS, PART III.

CLASS III. PISCES—Continued.

ORDER BB. ACANTHOPTERI—Continued.

Family CLXXXIV. Triglidæ—Continued. Page.

Group Gobioidea ... 2184

Family CLXXXVII. Callionymidæ ... 2184

Genus 799. Callionymus, Linnæus ... 2185

2511. bairdi, Jordan ... 2185

2512. himantophorus, Goode & Bean ... 2186

2513. calliurus, Eigenmann & Eigenmann ... 2187

2514. pauciradiatus, Gill ... 2188

Family CLXXXVIII. Gobiidæ ... 2188

Genus 800. Ioglossus, Bean ... 2192

2515. calliurus, Bean ... 2193

Genus 801. Philypnus, Cuvier & Valenciennes ... 2194

2516. dormitor (Lacépède) ... 2194

2517. lateralis, Gill ... 2195

Genus 802. Dormitator, Gill ... 2195

2518. maculatus (Bloch) ... 2196

Genus 803. Guavina, Bleeker ... 2198

2519. guavina, Cuvier & Valenciennes ... 2198

Genus 804. Eleotris (Gronow) Bloch & Schneider ... 2199

2520. amblyopsis (Cope) ... 2199

2521. abacurus, Jordan & Gilbert ... 2200

2522. pisonis (Gmelin) ... 2200

2523. perniger (Cope) ... 2201

2524. pictus (Kner & Steindachner) ... 2201

Genus 805. Alexurus, Jordan ... 2202

2525. armiger, Jordan & Richardson ... 2203

Genus 806. Erotelis, Poey ... 2203

2526. smaragdus (Cuvier & Valenciennes) ... 2204

Genus 807. Gymneleotris, Bleeker ... 2204

2527. seminudus (Günther) ... 2204

Genus 808. Chriolepis, Gilbert ... 2205

2528. minutillus, Gilbert ... 2205

Genus 809. Sicydium, Cuvier & Valenciennes ... 2205

2529. plumieri (Bloch) ... 2206

2530. antillarum, Ogilvie-Grant ... 2206

2531. vincente, Jordan & Evermann ... 2207

2531 (a). punctatum, Perugia ... 2867

Genus 810. Cotylopus, Guichenot ... 2207

Subgenus Sicya, Jordan & Evermann ... 2207

2532. gymnogaster (Ogilvie-Grant) ... 2207

2533. salvini (Ogilvie-Grant) ... 2208

Genus 811. Evorthodus, Gill ... 2208

2534. breviceps, Gill ... 2208

Genus 812. Lophogobius, Gill ... 2209

2535. cyprinoides (Pallas) ... 2209

CLASS III. PISCES—Continued.
ORDER BB. ACANTHOPTERI—Continued.
Family CLXXXVIII. Gobiidæ—Continued. Page.
Genus 813. Gobius (Artedi) Linnæus ... 2210
Subgenus Gobius ... 2216
2536. soporator, Cuvier & Valenciennes ... 2216
Subgenus Ctenogobius, Gill ... 2218
2537. nicholsii, Bean ... 2218
2538. eigenmanni, Garman ... 2218
2539. glaucofrænum (Gill) ... 2219
2540. manglicola, Jordan & Starks ... 2220
2541. stigmaturus, Goode & Bean ... 2220
2542. quadriporus, Cuvier & Valenciennes ... 2221
2543. shufeldti, Jordan & Eigenmann ... 2221
2544. boleosoma, Jordan & Gilbert ... 2221
2545. fasciatus (Gill) ... 2222
2546. encæomus, Jordan & Gilbert ... 2223
2547. stigmaticus (Poey) ... 2224
2548. lyricus, Girard ... 2224
2549. garmani, Eigenmann & Eigenmann ... 2225
2550. zebra, Gilbert ... 2226; 2867
Subgenus Euctenogobius, Gill ... 2226
2551. poeyi, Steindachner ... 2226
2552. badius (Gill) ... 2227
Subgenus Gobionellus, Girard ... 2227
2553. microdon, Gilbert ... 2227
2554. smaragdus, Cuvier & Valenciennes ... 2227
2555. strigatus, O'Shaughnessy ... 2228
2556. sagittula (Günther) ... 2228
2557. hastatus, Girard ... 2229
2558. oceanicus, Pallas ... 2230
Subgenus Lythrypnus, Jordan & Evermann ... 2230
2559. dalli, Gilbert ... 2230
Genus 814. Garmannia, Jordan & Evermann ... 2231
Subgenus Garmannia ... 2232
2560. paradoxa (Günther) ... 2232
2561. hemigymna (Eigenmann & Eigenmann) ... 2233
Subgenus Enypnias, Jordan & Evermann ... 2233
2562. seminuda (Günther) ... 2233
Genus 815. Awaous, Steindachner ... 2234
2563. flavus (Cuvier & Valenciennes) ... 2235
2564. nelsoni, Evermann ... 2235
2565. taiasica (Lichtenstein) ... 2236
2566. mexicanus (Günther) ... 2237
Genus 816. Bollmannia, Jordan ... 2237
2567. ocellata, Gilbert ... 2238
2568. chlamydes, Jordan ... 2238
2569. macropoma, Gilbert ... 2239
2570. stigmatura, Gilbert ... 2239
Genus 817. Aboma, Jordan & Starks ... 2240
2571. etheostoma, Jordan & Starks ... 2240
2572. lucretiæ (Eigenmann & Eigenmann) ... 2241
2573. chiquita (Jenkins & Evermann) ... 2241
Genus 818. Microgobius, Poey ... 2242
2574. gulosus (Girard) ... 2243
2575. eulepis, Eigenmann & Eigenmann ... 2244
2576. thalassinus, Jordan & Gilbert ... 2245
2577. signatus, Poey ... 2246
Genus 819. Zalypnus, Jordan & Evermann ... 2246

CLASS III. PISCES—Continued.
ORDER BB. ACANTHOPTERI—Continued.
Family CLXXXVIII. Gobiidæ—Continued. Page.
2578. cyclolepis (Gilbert) 2246
2579. emblematicus (Jordan & Gilbert) 2247
Genus 820. Eucyclogobius, Gill 2248
2580. newberryi (Girard) 2248
Genus 821. Lepidogobius, Gill 2249
2581. lepidus (Girard) 2249
Genus 822. Gillichthys, Cooper 2249
2582. mirabilis, Cooper 2250
2583. detrusus, Gilbert & Scofield 2251
Genus 823. Quietula, Jordan & Evermann 2251
2584. y-cauda (Jenkins & Evermann) 2251
Genus 824. Ilypnus, Jordan & Evermann 2253
2585. gilberti (Eigenmann & Eigenmann) 2253
Genus 825. Clevelandia, Eigenmann & Eigenmann 2254
2586. ios (Jordan & Gilbert) 2254
2587. rosæ, Jordan & Evermann 2255
Genus 826. Evermannia, Jordan 2256
2588. longipinnis (Steindachner) 2256
2589. zosterura (Jordan & Gilbert) 2256
Genus 827. Gobiosoma, Girard 2257
2590. histrio, Jordan 2258
2591. molestum, Girard 2258
2592. bosci (Lacépède) 2259
2593. crescentale, Gilbert 2259
2594. multifasciatum, Steindachner 2260
Genus 828. Barbulifer, Eigenmann & Eigenmann 2260
2595. ceuthœcus (Jordan & Gilbert) 2260
Genus 829. Typhlogobius, Steindachner 2261
2596. californiensis, Steindachner 2262
Genus 830. Tyntlastes, Günther 2262
2597. brevis (Günther) 2262
2598. sagitta (Günther) 2263
Genus 831. Gobioides, Lacépède 2263; 2868
2599. broussonnetii, Lacépède 2263
2600. peruanus (Steindachner) 2264
Genus 832. Cayennia, Sauvage 2265
2601. guichenoti, Sauvage 2265
SUBORDER DISCOCEPHALI 2265
Family CLXXXIX. Echeneididæ 2265
Genus 833. Phtheirichthys, Gill 2268
2602. lineatus (Menzies) 2268
Genus 834. Echeneis (Artedi) Linnæus 2268
2603. naucrates, Linnæus 2269
2604. naucrateoides, Zuieuw 2270
Genus 835. Remilegia, Gill 2270
2605. australis (Bennett) 2270
Genus 836. Remora, Gill 2271
Subgenus Remora 2271
2606. remora (Linnæus) 2271
Subgenus Remorina, Jordan & Evermann 2272
2607. albescens (Temminck & Schlegel) 2272
Subgenus Remoropsis, Gill 2272
2608. brachyptera (Lowe) 2272
Genus 837. Rhombochirus, Gill 2273
2609. osteochir (Cuvier) 2273
Group Trachinoidea 2273

CLASS III. PISCES—Continued.
ORDER BB. ACANTHOPTERI—Continued. Page.
Family CXC. Malacanthidæ 2274
Genus 838. Malacanthus, Cuvier 2275
2610. plumieri (Bloch) 2275
Genus 839. Caulolatilus, Gill 2276
2611. princeps (Jenyns) 2276
2612. microps, Goode & Bean 2277
2613. cyanops, Poey 2278
Genus 840. Lopholatilus, Goode & Bean 2278
2614. chamæleonticeps, Goode & Bean 2278
Family CXCI. Opisthognathidæ 2279
Genus 841. Opisthognathus Cuvier 2280
2615. lonchurum, Jordan & Gilbert 2281
2616 punctatum, Peters 2281
2617. macrognathum, Poey 2281
2618. ommatum, Jenkins & Evermann 2282
Genus 842. Gnathypops, Gill 2283
2619. scops, Jenkins & Evermann 2283
2620. maxillosa (Poey) 2284
2621. macrops (Poey) 2284
2622. rhomalea (Jordan & Gilbert) 2285
2623. snyderi, Jordan & Evermann 2285
2624. mystacina, Jordan 2286
Genus 843. Lonchopisthus, Gill 2286
2625. micrognathus (Poey) 2287
Family CXCII. Bathymasteridæ 2287
Genus 844. Bathymaster 2288
2626. signatus, Cope 2288
Genus 845. Ronquilus, Jordan & Starks 2289
2627. jordâni (Gilbert) 2289
Genus 846. Rathbunella, Jordan & Evermann 2289
2628. hypoplecta (Gilbert) 2290
Family CXCIII. Chiasmodontidæ 2291
Genus 847. Chiasmodon, Johnson 2291
2629. niger, Johnson 2291
Genus 848. Pseudoscopelus, Lütken 2292
2630. scriptus, Lütken 2292
Family CXCIV. Chœnichthyidæ 2293
Genus 849. Hypsicometes, Goode 2293
2631. gobioides, Goode 2294
Family CXCV. Trichodontidæ 2295
Genus 850. Trichodon (Steller) Cuvier 2295
2632. trichodon (Tilesius) 2295
Genus 851. Arctoscopus, Jordan & Evermann 2297
2633. japonicus (Steindachner) 2297
Family CXCVI. Dactyloscopidæ 2297
Genus 852. Gillellus, Gilbert 2298
2634. semicinctus, Gilbert 2298
2635. arenicola, Gilbert 2299
2636. ornatus, Gilbert 2299
Genus 853. Dactyloscopus, Gill 2300
Subgenus Dactyloscopus 2301
2637. pectoralis, Gill 2301
2638. tridigitatus, Gill 2301
2639. poeyi, Gill 2302
2640. lunaticus, Gilbert 2302
Subgenus Esloscopus, Jordan & Evermann 2303
2641. zelotes, Jordan & Evermann 2303

CLASS III. PISCES—Continued.
ORDER BB. ACANTHOPTERI—Continued.
Family CXCVI. Dactyloscopidæ—Continued. Page.
Genus 854. Dactylagnus, Gill ... 2304
2642. mundus, Gill ... 2304
Genus 855. Myxodagnus, Gill ... 2305
2643. opercularis, Gill ... 2305
Family CXCVII. Uranoscopidæ ... 2305
Genus 856. Astroscopus, Brevoort ... 2306
2644. y-græcum (Cuvier & Valenciennes) ... 2307
2645. zephyreus, Gilbert & Starks ... 2309
2646. guttatus (Abbott) ... 2310
Genus 857. Kathetostoma, Günther ... 2311
2647. averruncus, Jordan & Bollman ... 2311
2648. albiguttum, Bean ... 2312
SUBORDER HAPLODOCI ... 2313
Family CXCVIII. Batrachoididæ ... 2313
Genus 858. Batrachoides, Lacépède ... 2314; 2868
2649. surinamensis (Bloch & Schneider) ... 2314
2650. pacifici (Günther) ... 2314
Genus 859. Opsanus, Rafinesque ... 2315
2651. tau (Linnæus) ... 2315
2652. pardus (Goode & Bean) ... 2316
Genus 860. Porichthys, Girard ... 2317
2653. porosissimus (Cuvier & Valenciennes) ... 2319
2654. notatus, Girard ... 2321
2655. margaritatus (Richardson) ... 2322
Genus 861. Thalassophryne, Günther ... 2323
2656. maculosa, Günther ... 2324
2657. reticulata, Günther ... 2325
Genus 862. Dæctor, Jordan & Evermann ... 2325
2658. dowi (Jordan & Gilbert) ... 2325
SUBORDER XENOPTERYGII ... 2326
Family CXCIX. Gobiesocidæ ... 2326
Genus 863. Caularchus, Gill ... 2327
2659. mæandricus (Girard) ... 2328
Genus 864. Bryssetæres, Jordan & Evermann ... 2328
2660. pinniger (Gilbert) ... 2328
Genus 865. Gobiesox, Lacépède ... 2329
Subgenus Bryssophilus, Jordan & Evermann ... 2330
2661. papillifer, Gilbert ... 2330
Subgenus Gobiesox ... 2331
2662. gyrinus, Jordan & Evermann ... 2331
2663. nigripinnis (Peters) ... 2331
2664. cephalus, Lacépède ... 2332
2665. tudes, Richardson ... 2333
2666. strumosus, Cope ... 2333
2667. virgatulus, Jordan & Gilbert ... 2333
2668. adustus, Jordan & Gilbert ... 2334
2669. funebris, Gilbert ... 2334
2670. pœcilophthalmus, Jenyns ... 2335
2671. rhodospilus, Günther ... 2335
2672. macrophthalmus, Günther ... 2335
2673. cerasinus, Cope ... 2336
Subgenus Sicyases, Müller & Troschel ... 2336
2674. erythrops, Jordan & Gilbert ... 2336
2675. rubiginosus (Poey) ... 2337
2676. carneus (Poey) ... 2337
2677. hæres, Jordan & Bollman ... 2337

CLASS III. PISCES—Continued.
ORDER BB. ACANTHOPTERI—Continued.
Family CXCIX. Gobiesocidæ—Continued. Page.
2678. punctulatus (Poey) ... 2338
2679. fasciatus (Peters) ... 2338
Genus 866. Rimicola, Jordan & Evermann ... 2338
2680. muscarum (Meek & Pierson) ... 2338
2681. eigenmanni (Gilbert) ... 2339
Genus 867. Arbaciosa, Jordan & Evermann ... 2340
2682. rhessodon (Rosa Smith) ... 2340
2683. humeralis (Gilbert) ... 2341
2684. rupestris (Poey) ... 2341
2685. zebra (Jordan & Gilbert) ... 2341
2686. eos (Jordan & Gilbert) ... 2343
Group Blennioidea ... 2343
Family CC. Blenniidæ ... 2344
Genus 868. Enneanectes, Jordan & Evermann ... 2349
2687. carminalis (Jordan & Gilbert) ... 2350
Genus 868(a). Dialommus, Gilbert ... 2868
2687(a). fuscus, Gilbert ... 2868
Genus 869. Heterostichus, Girard ... 2350
2688. rostratus, Girard ... 2351
Genus 870. Gibbonsia, Cooper ... 2351
2689. evides (Jordan & Gilbert) ... 2352; 2869
2690. elegans (Cooper) ... 2353
Genus 871. Neoclinus, Girard ... 2354
Subgenus Neoclinus ... 2354
2691. blanchardi, Girard ... 2354
Subgenus Pterognathus, Girard ... 2355
2692. satiricus, Girard ... 2355
Genus 872. Malacoctenus, Gill ... 2356
2693. ocellatus (Steindachner) ... 2356; 2869
2694. varius (Poey) ... 2357
2695. macropus (Poey) ... 2357
2696. lugubris (Poey) ... 2357
2697. gillii (Steindachner) ... 2358
2698. bimaculatus (Steindachner) ... 2358
2699. delalandi (Cuvier & Valenciennes) ... 2358
2700. versicolor (Poey) ... 2359
2701. biguttatus (Cope) ... 2360
Genus 873. Labrisomus, Swainson ... 2360
2702. herminier (Le Sueur) ... 2361
2703. nuchipinnis (Quoy & Gaimard) ... 2362
2704. xanti, Gill ... 2362
2705. bucciferus, Poey ... 2363
2706. microlepidotus, Poey ... 2363
Genus 874. Mnierpes, Jordan & Evermann ... 2364
2707. macrocephalus (Günther) ... 2364
Genus 875. Gobioclinus, Gill ... 2364
2708. gobio (Cuvier & Valenciennes) ... 2365
Genus 876. Starksia, Jordan & Evermann ... 2365
2709. cremnobates (Gilbert) ... 2365
Genus 877. Cryptotrema, Gilbert ... 2366
2710. corallinum, Gilbert ... 2366
Genus 878. Exerpes, Jordan & Evermann ... 2367
2711. asper (Jenkins & Evermann) ... 2367
Genus 879. Auchenopterus, Günther ... 2369
Subgenus Corallicola, Jordan & Evermann ... 2369
2712. nigripinnis (Steindachner) ... 2369

CLASS III. PISCES—Continued.
ORDER BB. ACANTHOPTERI—Continued.
Family CC. Blenniidæ—Continued. Page.
2713. altivelis (Lockington) ... 2370
2714. marmoratus (Steindachner) ... 2371
Subgenus Auchenopterus ... 2371
2715. affinis (Steindachner) ... 2371
2716. monophthalmus, Günther ... 2372
2717. integripinnis (Rosa Smith) ... 2372
2718. fasciatus (Steindachner) ... 2373
2719. nox (Jordan & Gilbert) ... 2373
Genus 880. Paraclinus, Mocquard ... 2374
2720. chaperi, Mocquard ... 2374
Genus 881. Emmnion, Jordan ... 2375
2721. bristolæ, Jordan ... 2375
Genus 882. Atopoclinus, Vaillant ... 2376
2722. ringens, Vaillant ... 2376
Genus 883. Runula, Jordan & Bollman ... 2377
2723. azalea, Jordan & Bollman ... 2377
Genus 884. Blennius (Artedi) Linnæus ... 2377
Subgenus Lipophrys, Gill ... 2378
2724. carolinus (Cuvier & Valenciennes) ... 2378
2725. fucorum, Cuvier & Valenciennes ... 2379
2726. stearnsi, Jordan & Gilbert ... 2379
2727. favosus, Goode & Bean ... 2380
2728. pilicornis, Cuvier & Valenciennes ... 2380
2729. marmoreus, Poey ... 2381
2730. truncatus (Poey) ... 2381
2731. vinctus, Poey ... 2382
2732. cristatus, Linnæus ... 2382
Genus 885. Scartella, Jordan ... 2384
2733. microstoma (Poey) ... 2384
Genus 886. Hypleurochilus, Gill ... 2385
2734. geminatus (Wood) ... 2385
Genus 887. Hypsoblennius, Gill ... 2386
Subgenus Hypsoblennius ... 2386
2735. gilberti (Jordan) ... 2386
2736. gentilis (Girard) ... 2387
2737. striatus (Steindachner) ... 2388
2738. ionthas (Jordan & Gilbert) ... 2388
2739. hentz (Le Sueur) ... 2390
Subgenus Blenniolus, Jordan & Evermann ... 2390
2740. brevipinnis (Günther) ... 2390
Genus 888. Chasmodes, Cuvier & Valenciennes ... 2391
2741. jenkinsi (Jordan & Evermann) ... 2391
2742. quadrifasciatus (Wood) ... 2392
2743. saburræ, Jordan & Gilbert ... 2392
2744. novemlineatus (Wood) ... 2393
2745. bosquianus (Lacépède) ... 2394
Genus 889. Homesthes, Gilbert ... 2394
2746. caulopus, Gilbert ... 2394
Genus 890. Scartichthys, Jordan & Evermann ... 2395
2747. rubropunctatus (Cuvier & Valenciennes) ... 2396
Genus 891. Rupiscartes, Swainson ... 2396
2748. atlanticus (Cuvier & Valenciennes) ... 2397
Genus 892. Entomacrodus, Gill ... 2397
2749. chiostictus (Jordan & Gilbert) ... 2398
2750. margaritaceus (Poey) ... 2398
2751. decoratus, Poey ... 2399

CLASS III. PISCES—Continued.
ORDER BB. ACANTHOPTERI—Continued.
Family CC. Blenniidæ—Continued. Page.
2752. nigricans, Gill ... 2399
Genus 893. Salariichthys, Guichenot ... 2400
2753. textilis (Quoy & Gaimard) ... 2400
Genus 894. Ophioblennius, Gill ... 2400
2754. webbii (Valenciennes) ... 2401
2755. steindachneri, Jordan & Evermann ... 2401
Genus 895. Emblemaria, Jordan & Gilbert ... 2401
2756. atlantica, Jordan & Evermann ... 2402
2757. nivipes, Jordan & Gilbert ... 2402
2758. oculocirris, Jordan ... 2403
Genus 896. Chænopsis, Gill ... 2403
2759. ocellatus, Poey ... 2403
Genus 897. Lucioblennius, Gilbert ... 2404
2760. alepidotus, Gilbert ... 2404
Genus 898. Pholidichthys, Bleeker ... 2405
2761. anguilliformis, Lockington ... 2405
Genus 899. Psednoblennius, Jenkins & Evermann ... 2406
2762. hypacanthus, Jenkins & Evermann ... 2406
Genus 900. Stathmonotus, Bean ... 2407
2763. hemphillii, Bean ... 2407
Genus 901. Bryostemma, Jordan & Starks ... 2408
2764. polyactocephalum (Pallas) ... 2408
2765. nugator, Jordan & Williams ... 2410
Genus 902. Apodichthys, Girard ... 2411
2766. flavidus, Girard ... 2411
2767. univittatus, Lockington ... 2412
Genus 903. Xererpes, Jordan & Gilbert ... 2413
2768. fucorum (Jordan & Gilbert) ... 2413
Genus 904. Ulvicola, Gilbert ... 2413
2769. sanctæ-rosæ, Gilbert & Starks ... 2413
Genus 905. Pholis (Gronow) Scopoli ... 2414
Subgenus Urocentrus, Kner ... 2415
2770. pictus (Kner) ... 2415
Subgenus Rhodymenichthys, Jordan & Evermann ... 2416
2771. dolichogaster (Pallas) ... 2416
Subgenus Pholis ... 2417
2772. fasciatus (Bloch & Schneider) ... 2417
2773. gunnellus (Linnæus) ... 2419
2774. ornatus (Girard) ... 2419
Genus 906. Gunnellops, Bleeker ... 2420
2775. roseus (Pallas) ... 2420
Genus 907. Asternopteryx, Rüppell ... 2420
2776. gunnelliformis, Rüppell ... 2420
Genus 908. Anoplarchus, Gill ... 2421
2777. atropurpureus (Kittlitz) ... 2422; 2869
Genus 908(a). Alectrias, Jordan & Evermann ... 2869
2778. alectrolophus (Pallas) ... 2421; 2869
Genus 909. Xiphister, Jordan & Starks ... 2423
2779. ulvæ, Jordan & Starks ... 2423
2780. chirus (Jordan & Gilbert) ... 2424
Genus 910. Xiphidion, Girard ... 2424
2781. mucosum, Girard ... 2425
2782. rupestre (Jordan & Gilbert) ... 2426
Genus 911. Cebedichthys, Ayres ... 2426
2783. violaceus (Ayres) ... 2427
Genus 912. Plagiogrammus, Bean ... 2427

CLASS III. PISCES—Continued.
ORDER BB. ACANTHOPTERI—Continued.
Family CC. Blenniidæ—Continued. Page.
2784. hopkinsi, Bean 2428
Genus 913. Opisthocentrus, Kner 2428
2785. ocellatus (Tilesius) 2429
Genus 914. Pholidapus, Bean & Bean 2430
2786. dybowskii (Steindachner) 2430
Genus 915. Plectobranchus, Gilbert 2431
2787. evides, Gilbert 2432
Genus 916. Leptoclinus, Gill 2432
2788. maculatus (Fries) 2433
Genus 917. Poroclinus, Bean 2433
2789. rothrocki, Bean 2434
Genus 918. Lumpenus, Reinhardt 2435
Subgenus Anisarchus, Gill 2435
2790. medius (Reinhardt) 2435
Subgenus Lumpenus 2436
2791. anguillaris (Pallas) 2436
2792. mackayi (Gilbert) 2436
2793. fabricii (Cuvier & Valenciennes) 2437
2794. lampetræformis (Walbaum) 2438
Genus 919. Stichæus, Reinhardt 2439
2795. punctatus (Fabricius) 2439
Genus 920. Ulvaria, Jordan & Evermann 2440
2796. subbifurcata (Storer) 2440
Genus 921. Eumesogrammus, Gill 2441
2797. præcisus (Kröyer) 2441
Family CCI. Cryptacanthodidæ 2442
Genus 922. Delolepis, Bean 2442
2798. virgatus, Bean 2442
Genus 923. Cryptacanthodes, Storer 2443
2799. maculatus, Storer 2443
Genus 924. Lyconectes, Gilbert 2444
2800. aleutensis, Gilbert 2444
Family CCII. Anarhichadidæ 2445
Genus 925. Anarhichas (Artedi) Linnæus 2445
2801. latifrons, Steenstrup & Hallgrimsson 2446
2802. minor, Olafsen 2446
2803. lupus, Linnæus 2446
2804. lepturus, Bean 2447
2805. orientalis, Pallas 2447
Genus 926. Anarrhichthys, Ayres 2447
2806. ocellatus, Ayres 2448
Family CCIII. Cerdalidæ 2448
Genus 927. Cerdale, Jordan & Gilbert 2448
2807. ionthas, Jordan & Gilbert 2449
Genus 928. Microdesmus, Günther 2450
2808. dipus, Günther 2450
2809. retropinnis, Jordan & Gilbert 2450
Family CCIV. Ptilichthyidæ 2451
Genus 929. Ptilichthys, Bean 2452
2810. goodei, Bean 2452
Group Ophidioidea 2453
Family CCV. Scytalinidæ 2453
Genus 930. Scytalina, Jordan & Gilbert 2454
2811. cerdale, Jordan & Gilbert 2454
Family CCVI. Zoarcidæ 2455
Genus 931. Zoarces, Gill 2456

CLASS III. PISCES—Continued.
ORDER BB. ACANTHOPTERI—Continued.
Family CCVI. Zoarcidæ—Continued. Page.
Subgenus Macrozoarces, Gill ... 2457
2812. anguillaris (Peck) ... 2457
Genus 932. Embryx, Jordan & Evermann ... 2458
2813. crassilabris (Gilbert) ... 2458
2814. crotalinus (Gilbert) ... 2458
Genus 933. Lycodopsis, Collett ... 2460
2815. pacificus (Collett) ... 2460
Genus 934. Aprodon, Gilbert ... 2460
2816. cortezianus, Gilbert ... 2461
Genus 935. Lycodes, Reinhardt ... 2461
Subgenus Lycodes ... 2463
2817. esmarkii, Collett ... 2463
2818. vahlii, Reinhardt ... 2463
2819. concolor, Gill & Townsend ... 2463
2820. zoarchus, Goode & Bean ... 2464
2821. reticulatus, Reinhardt ... 2465
2822. perspicillum, Kröyer ... 2465
2823. frigidus, Collett ... 2465
2824. terræ-novæ, Collett ... 2466
2825. digitatus, Gill & Townsend ... 2466
2826. palearis, Gilbert ... 2466
2827. brevipes, Bean ... 2467
Subgenus Lycias, Jordan & Evermann ... 2468
2828. nebulosus, Kröyer ... 2468
2829. seminudus, Reinhardt ... 2468
Genus 936. Lycodalepis, Bleeker ... 2468
2830. polaris (Sabine) ... 2468
2831. mucosus (Richardson) ... 2470
Genus 937. Lycenchelys, Gill ... 2470
2832. verrillii (Goode & Bean) ... 2470
2833. paxillus (Goode & Bean) ... 2471
2834. porifer (Gilbert) ... 2471; 2869
Genus 938. Furcella, Jordan & Evermann ... 2472
2835. diaptera (Gilbert) ... 2472
Genus 939. Lycodonus, Goode & Bean ... 2473
2836. mirabilis, Goode & Bean ... 2474
Genus 940. Lyconema, Gilbert ... 2474
2837. barbatum, Gilbert ... 2474
Genus 941. Bothrocara, Bean ... 2475
2838. pusilla (Bean) ... 2476
2839. mollis, Bean ... 2476
Genus 942. Gymnelis, Reinhardt ... 2477
2840. viridis (Fabricius) ... 2477
2841. stigma (Lay & Bennett) ... 2477
Genus 943. Lycocara, Gill ... 2478
2842. parrii (Ross) ... 2478
Genus 944. Melanostigma, Günther ... 2478
2843. gelatinosum, Günther ... 2479
2844. pammelas, Gilbert ... 2479
Family CCVII. Derepodichthyidæ ... 2480
Genus 945. Derepodichthys, Gilbert ... 2480
2845. alepidotus, Gilbert ... 2480
Family CCVIII. Ophidiidæ ... 2481
Genus 946. Lepophidium, Gill ... 2482
2846. marmoratum (Goode & Bean) ... 2482
2847. emmelas (Gilbert) ... 2483

CLASS III. PISCES—Continued.
ORDER BB. ACANTHOPTERI—Continued.
Family CCVIII. Ophidiidæ—Continued. Page.
2848. stigmatistium (Gilbert) ... 2483
2849. profundorum (Gill) ... 2484
2850. cervinum (Goode & Bean) ... 2484
2851. prorates (Jordan & Bollman) ... 2485
2852. brevibarbe (Cuvier) ... 2485
2853. pardale (Gilbert) ... 2486
2854. microlepis (Gilbert) ... 2486
Genus 947. Ophidion (Artedi) Linnæus ... 2487
2855. beani, Jordan & Gilbert ... 2487
2856. holbrooki (Putnam) ... 2487
2857. graellsi, Poey ... 2488
Genus 948. Chilara, Jordan & Evermann ... 2488
2858. taylori (Girard) ... 2489
Genus 949. Rissola, Jordan & Evermann ... 2489
2859. marginata (De Kay) ... 2489
Genus 950. Otophidium, Gill ... 2490
2860. omostigma (Jordan & Gilbert) ... 2490
2861. indefatigabile, Jordan & Bollman ... 2490
2862. galeoides (Gilbert) ... 2491
Family CCIX. Lycodapodidæ ... 2491
Genus 951. Lycodapus, Gilbert ... 2492
2863. dermatinus, Gilbert ... 2492
2864. fierasfer, Gilbert ... 2493
2865. parviceps, Gilbert ... 2493
2866. extensus (Gilbert) ... 2494
Family CCX. Fierasferidæ ... 2494
Genus 952. Fierasfer, Cuvier ... 2495
2867. affinis (Günther) ... 2495
2868. arenicola, Jordan & Gilbert ... 2496
2869. bermudensis (Jones) ... 2497
Family CCXI. Brotulidæ ... 2498
Genus 953. Brotula, Cuvier ... 2500
2870. barbata (Bloch & Schneider) ... 2500
Genus 954. Stygicola, Gill ... 2500
2871. dentatus (Poey) ... 2500
Genus 955. Lucifuga, Poey ... 2501
2872. subterraneus, Poey ... 2501
Genus 956. Brosmophycis, Gill ... 2502
2873. marginatus (Ayres) ... 2502
Genus 957. Ogilbia, Jordan & Evermann ... 2502
2874. ventralis (Gill) ... 2503
2875. cayorum, Evermann & Kendall ... 2503
Genus 958. Bythites, Reinhardt ... 2504
2876. fuscus, Reinhardt ... 2504
Genus 959. Catætyx, Günther ... 2504
2877. rubirostris, Gilbert ... 2505
Genus 960. Dicromita, Goode & Bean ... 2506
2878. agassizii, Goode & Bean ... 2506
Genus 961. Bassozetus, Gill ... 2507
2879. normalis, Gill ... 2507
2880. compressus (Günther) ... 2508
2881. catena, Goode & Bean ... 2509
2882. tænia (Günther) ... 2510
Genus 962. Mœbia, Goode & Bean ... 2510
2883. promelas (Gilbert) ... 2511
Genus 963. Neobythites, Goode & Bean ... 2512

CLASS III. PISCES—Continued.
ORDER BB. ACANTHOPTERI—Continued.
Family COXI. Brotulidæ—Continued. Page.
2884. gillii, Goode & Bean ... 2512
2885. marginatus, Goode & Bean ... 2513
Genus 964. Benthocometes, Goode & Bean ... 2513
2886. robustus, Goode & Bean ... 2514
Genus 965. Bassogigas, Gill ... 2515
2887. gillii, Goode & Bean ... 2515
2888. stelliferoides (Gilbert) ... 2516
Genus 966. Barathrodemus, Goode & Bean ... 2517
2889. manatinus, Goode & Bean ... 2517
Genus 967. Nematonus, Günther ... 2518
2890. pectoralis, Goode & Bean ... 2518
Genus 968. Porogadus, Goode & Bean ... 2519
2891. miles, Goode & Bean ... 2520
Genus 969. Penopus, Goode & Bean ... 2520
2892. macdonaldi, Goode & Bean ... 2521
Genus 970. Dicrolene, Goode & Bean ... 2522
2893. intronigra, Goode & Bean ... 2522
Genus 971. Mixonus, Günther ... 2523
2894. laticeps (Günther) ... 2523
Genus 972. Barathronus, Goode & Bean ... 2524
2895. bicolor, Goode & Bean ... 2524
Genus 973. Aphyonus, Günther ... 2525
2896. mollis, Goode & Bean ... 2525
Family CCXII. Bregmacerotidæ ... 2525
Genus 974. Bregmaceros, Thompson ... 2526
2897. macclellandii, Thompson ... 2526
2898. atlanticus, Goode & Bean ... 2527
SUBORDER ANACANTHINI ... 2528
Family CCXIII. Merlucciidæ ... 2529
Genus 975. Merluccius, Rafinesque ... 2529
2899. merluccius (Linnæus) ... 2530
2900. bilinearis (Mitchill) ... 2530
2901. productus (Ayres) ... 2531
Family CCXIV. Gadidæ ... 2531
Genus 976. Boreogadus, Günther ... 2533
2902. saida (Lepechin) ... 2533
Genus 977. Pollachius, Nilsson ... 2534
2903. virens (Linnæus) ... 2534
Genus 978. Theragra, Lucas ... 2535
2904. chalcogramma (Pallas) ... 2535
2905. fucensis (Jordan & Gilbert) ... 2536
Genus 979. Eleginus, Fischer ... 2537
2906. navaga (Kölreuter) ... 2537
Genus 980. Microgadus, Gill ... 2538
2907. proximus (Girard) ... 2539
2908. tomcod (Walbaum) ... 2540
Genus 981. Gadus (Artedi) Linnæus ... 2540
2909. callarias, Linnæus ... 2541
2910. macrocephalus, Tilesius ... 2541
2911. ogac, Richardson ... 2542
Genus 982. Melanogrammus, Gill ... 2542
2912. æglefinus, Linnæus ... 2542
Genus 983. Lepedion, Swainson ... 2543
2913. verecundum, Jordan & Cramer ... 2543
Genus 984. Antimora, Günther ... 2544
2914. viola (Goode & Bean) ... 2544

CLASS III. PISCES—Continued.
ORDER BB. ACANTHOPTERI—Continued.
Family CCXIV. Gadidæ—Continued. Page.
2915. microlepis, Bean ... 2545
Genus 985 Uraleptus, Costa ... 2545
2916. maraldi (Risso) ... 2545
Genus 986. Lotella, Kaup ... 2546
2917. maxillaris, Bean ... 2546
Genus 987. Physiculus, Kaup ... 2547
2918. fulvus, Bean ... 2547
2919. nematopus, Gilbert ... 2548
2920. kaupi, Poey ... 2548
2921. rastrelliger, Gilbert ... 2549
Genus 988. Lota (Cuvier) Oken ... 2550
2922. maculosa (Le Sueur) ... 2550
Genus 989. Molva, Fleming ... 2551
2923. molva (Linnæus) ... 2551
Genus 990. Urophycis, Gill ... 2552
Subgenus Urophycis ... 2553
2924. regius (Walbaum) ... 2553
2925. cirratus (Goode & Bean) ... 2553
2926. floridanus (Bean & Dresel) ... 2554
Subgenus Emphycus, Jordan & Evermann ... 2554
2927. carlli (Bean) ... 2554
2928. tenuis (Mitchill) ... 2555
2929. chuss (Walbaum) ... 2555
2930. chesteri (Goode & Bean) ... 2556
Genus 991. Laemonema, Günther ... 2556
2931. barbatulum, Goode & Bean ... 2556
2932. melanurum, Goode & Bean ... 2557
Genus 992. Gaidropsarus, Rafinesque ... 2557
2933. ensis (Reinhardt) ... 2558
2934. argentatus (Reinhardt) ... 2559
2935. septentrionalis (Collett) ... 2559
Genus 993. Enchelyopus, Bloch & Schneider ... 2560
2936. cimbrius (Linnæus) ... 2560
Genus 994. Brosme (Cuvier) Oken ... 2561
2937. brosme (Müller) ... 2561
Family CCXV. Macrouridæ ... 2561
Genus 995. Bathygadus, Günther ... 2563
2938. arcuatus, Goode & Bean ... 2564
2939. favosus, Goode & Bean ... 2565
2940. macrops, Goode & Bean ... 2566
2941. longifilis, Goode & Bean ... 2566
Genus 996. Steindachneria, Goode & Bean ... 2567
2942. argentea, Goode & Bean ... 2568
Genus 997. Trachyrinchus, Giorna ... 2568
2943. helolepis, Gilbert ... 2569
Genus 998. Malacocephalus, Günther ... 2569
2944. occidentalis, Goode & Bean ... 2570
Genus 999. Moseleya, Goode & Bean ... 2570
2945. cyclolepis (Gilbert) ... 2570
Genus 1000. Nematonurus, Günther ... 2571
2946. goodei (Günther) ... 2571
2947. suborbitalis (Gill & Townsend) ... 2572
Genus 1001. Albatrossia, Jordan & Evermann ... 2573
2948. pectoralis (Gilbert) ... 2573
Genus 1002. Bogoslovius, Jordan & Evermann ... 2574
2949. clarki, Jordan & Gilbert ... 2575

CLASS III. PISCES—Continued.
ORDER BB. ACANTHOPTERI—Continued.
Family CCXV. Macrouridæ—Continued. Page.
2950. firmisquamis (Gill & Townsend) ... 2575
Genus 1003. Chalinura, Goode & Bean ... 2576
2951. serrula, Bean ... 2576
2952. filifera, Gilbert ... 2577
2953. simula, Goode & Bean ... 2578
Genus 1004. Coryhænoides, Gunner ... 2578
2954. rupestris, Gunner ... 2579
2955. carapinus, Goode & Bean ... 2579
Genus 1005. Hymenocephalus, Giglioli ... 2580
2956. cavernosus (Goode & Bean) ... 2580
Genus 1006. Macrourus, Bloch ... 2581
2957. berglax, Lacépède ... 2582
2958. holotrachys, Günther ... 2582
2959. bairdii, Goode & Bean ... 2583
2960. lepturus, Gill & Townsend ... 2584
2961. acrolepis, Bean ... 2585
2962. stelgidolepis, Gilbert ... 2585
2963. cinereus, Gilbert ... 2586
Genus 1007. Cœlorhynchus, Giorna ... 2587
2964. occa (Goode & Bean) ... 2588
2965. carminatus (Goode) ... 2588
2966. caribbæus (Goode & Bean) ... 2589
2967. scaphopsis (Gilbert) ... 2590
Genus 1008. Trachonurus, Günther ... 2591
2968. sulcatus (Goode & Bean) ... 2591
Genus 1009. Lionurus, Günther ... 2592
2969. filicauda (Günther) ... 2592
2970. liolepis, Gilbert ... 2593
SUBORDER TÆNIOSOMI ... 2594
Family CCXVI. Regalecidæ ... 2595
Genus 1010. Regalecus, Brünnich ... 2595
2971. glesne (Ascanius) ... 2596
Family CCXVII. Trachypteridæ ... 2597
Genus 1011. Trachypterus, Gouan ... 2599
2972. rex-salmonorum, Jordan & Gilbert ... 2599
2973. trachyurus, Poey ... 2600
Family CCXVIII. Stylephoridæ ... 2601
Genus 1012. Stylephorus, Shaw ... 2601
2974. chordatus, Shaw ... 2601
SUBORDER HETEROSOMATA ... 2602
Family CCXIX. Pleuronectidæ ... 2602
Genus 1013. Atheresthes, Jordan & Gilbert ... 2609
2975. stomias (Jordan & Gilbert) ... 2609
Genus 1014. Reinhardtius, Gill ... 2610
2976. hippoglossoides (Walbaum) ... 2611
Genus 1015. Hippoglossus, Cuvier ... 2611
2977. hippoglossus (Linnæus) ... 2611
Genus 1016. Lyopsetta, Jordan & Goss ... 2612
2978. exilis (Jordan & Gilbert) ... 2612
Genus 1017. Eopsetta, Jordan & Goss ... 2613
2979. jordani (Lockington) ... 2613
Genus 1018. Hippoglossoides, Gottsche ... 2614
2980. platessoides (Fabricius) ... 2614
2981. elassodon, Jordan & Gilbert ... 2615
2982. robustus, Gill & Townsend ... 2616
2983. hamiltoni, Jordan & Gilbert ... 2616

CLASS III. PISCES—Continued.
ORDER BB. ACANTHOPTERI—Continued.
Family CCXIX. Pleuronectidæ—Continued. Page.
Genus 1019. Psettichthys, Girard 2617
2984. melanostictus, Girard 2618
Genus 1020. Verasper, Jordan & Gilbert 2618
2985. moseri. Jordan & Gilbert 2619
Genus 1021. Hippoglossina, Steindachner 2620
2986. stomata, Eigenmann & Eigenmann 2620
2987. macrops, Steindachner 2621
2988. bollmani, Gilbert 2621
Genus 1022. Lioglossina, Gilbert 2622
2989. tetrophthalma, Gilbert 2622
Genus 1023. Xystreurys, Jordan & Gilbert 2623
2990. liolepis, Jordan & Gilbert 2623
Genus 1024. Paralichthys, Girard 2624
2991. californicus (Ayres) 2625
2991(a). Paralichthys magdalenæ, Abbott 2872
2992. æstuarius, Gilbert & Scofield 2626
2993. brasiliensis (Ranzani) 2626
2994. sinaloæ, Jordan & Abbott 2627, 2872
2995. woolmani, Jordan & Williams 2628
2996. dentatus (Linnæus) 2629
2997. lethostigmus, Jordan & Gilbert 2630
2998. squamilentus, Jordan & Gilbert 2631
2999. albiguttus, Jordan & Gilbert 2631
3000. oblongus (Mitchill) 2632
Genus 1025. Ramularia, Jordan & Evermann 2633
3001. dendritica (Gilbert) 2633
Genus 1026. Ancylopsetta, Gill 2634
3002. quadrocellata, Gill 2634
Genus 1027. Notosema, Goode & Bean 2635
3003. dilectum, Goode & Bean 2635
Genus 1028. Gastropsetta, B. A. Bean 2636
3004. frontalis, B. A. Bean 2636
Genus 1029. Pleuronichthys, Girard 2637
3005. decurrens, Jordan & Gilbert 2637
3006. verticalis, Jordan & Gilbert 2638
3007. cœnosus, Girard 2638
Genus 1030. Hypsopsetta, Gill 2639
3008. guttulata (Girard) 2639
Genus 1031. Parophrys, Girard 2640
3009. vetulus, Girard 2640
Genus 1032. Inopsetta, Jordan & Goss 2641
3010. ischyra (Jordan & Gilbert) 2641
Genus 1033. Isopsetta, Lockington 2642
3011. isolepis (Lockington) 2642
Genus 1034. Lepidopsetta, Gill 2642
3012. bilineata (Ayres) 2643
Genus 1035. Limanda, Gottsche 2644
3013. ferruginea (Storer) 2644
3014. aspera (Pallas) 2645
3015. proboscidea, Gilbert 2645
3016. beanii, Goode 2646
Genus 1036. Pseudopleuronectes, Bleeker 2646
3017. americanus (Walbaum) 2647
3018. pinnifasciatus (Kner) 2647
Genus 1037. Pleuronectes (Artedi) Linnæus 2648
3019. quadrituberculatus, Pallas 2648

CLASS III. PISCES—Continued.
ORDER BB. ACANTHOPTERI—Continued.
Family CCXIX. Pleuronectidæ—Continued. Page.
Genus 1038. Liopsetta, Gill 2649
3020. glacialis (Pallas) 2649
3021. putnami (Gill) 2650
3022. obscura (Herzenstein) 2651
Genus 1039. Platichthys, Girard 2651
3023. stellatus (Pallas) 2652
Genus 1040. Microstomus, Gottsche 2653
3024. kitt (Walbaum) 2654
3025. pacificus (Lockington) 2655
Genus 1041. Embassichthys, Jordan & Evermann 2655
3026. bathybius (Gilbert) 2655
Genus 1042. Glyptocephalus, Gottsche 2656
3027. cynoglossus (Linnæus) 2657
3028. zachirus, Lockington 2658
Genus 1043. Lophopsetta, Gill 2659
3029. maculata (Mitchill) 2660
Genus 1044. Platophrys, Swainson 2660
3030. spinosus (Poey) 2662
3031. constellatus, Jordan 2663
3032. ocellatus (Agassiz) 2663
3033. maculifer (Poey) 2664
3034. ellipticus (Poey) 2665
3035. lunatus (Linnæus) 2665
3036. leopardinus (Günther) 2666
Genus 1045. Perissias, Jordan & Evermann 2667
3037. taeniopterus (Gilbert) 2667
Genus 1046. Engyophrys, Jordan & Bollman 2668
3038. sancti-laurentii, Jordan & Bollman 2668
Genus 1047. Trichopsetta, Gill 2669
3039. ventralis (Goode & Bean) 2669
Genus 1048. Syacium, Ranzani 2670
3040. papillosum (Linnæus) 2671
3041. micrurum, Ranzani 2672
3042. latifrons (Jordan & Gilbert) 2673
3043. ovale (Günther) 2674
Genus 1049. Cyclopsetta, Gill 2675
3044. querna (Jordan & Bollman) 2675
3045. chittendeni, B. A. Bean 2676
3046. fimbriata (Goode & Bean) 2676
Genus 1050. Azevia, Jordan 2677
3047. panamensis (Steindachner) 2677
Genus 1051. Citharichthys, Bleeker 2678
Subgenus Orthopsetta, Gill 2679
3048. sordidus (Girard) 2679
3049. fragilis, Gilbert 2680
3050. xanthostigmus, Gilbert 2680
3051. stigmæus, Jordan & Gilbert 2681
Subgenus Citharichthys 2682
3052. dinoceros, Goode & Bean 2682
3053. platophrys, Gilbert 2683
3054. arctifrons, Goode 2683
3055. unicornis, Goode 2683
3056. uhleri, Jordan 2684
3057. macrops, Dressel 2684
3058. spilopterus, Günther 2685
3059. gilberti, Jenkins & Evermann 2686

CLASS III. PISCES—Continued.
ORDER BB. ACANTHOPTERI—Continued.
Family CCXIX. Pleuronectidæ—Continued. Page.
Genus 1052. Etropus, Jordan & Gilbert ... 2687
3060. microstomus (Gill) ... 2687
3061. rimosus, Goode & Bean ... 2688
3062. crossotus, Jordan & Gilbert ... 2689
Genus 1053. Monolene, Goode ... 2690
3063. sessilicauda, Goode ... 2691
3064. atrimana, Goode & Bean ... 2692
Family CCXX. Soleidæ ... 2692
Genus 1054. Achirus, Lacépède ... 2693
Subgenus Baiostoma, Bean ... 2695
3065. achirus (Linnæus) ... 2695
3066. inscriptus, Gosse ... 2696
3067. klunzingeri (Steindachner) ... 2697
3068. lineatus (Linnæus) ... 2697
3069. mazatlanus (Steindachner) ... 2698
3070. fonsecensis (Günther) ... 2699
3071. fischeri (Steindachner) ... 2699
3072. scutum (Günther) ... 2700
Subgenus Achirus ... 2700
3073. fimbriatus (Günther) ... 2700
3074. fasciatus, Lacépède ... 2700
3075. panamensis (Steindachner) ... 2702
Genus 1055. Apionichthys, Kaup ... 2702
3076. unicolor (Günther) ... 2702
Genus 1056. Gymnachirus, Kaup ... 2703
3077. fasciatus, Günther ... 2703
Genus 1057. Symphurus, Rafinesque ... 2704
Subgenus Symphurus ... 2705
3078. piger (Goode & Bean) ... 2705
3079. marginatus (Goode & Bean) ... 2706
3080. atramentatus, Jordan & Bollman ... 2706
3081. fasciolaris, Gilbert ... 2707
3082. elongatus (Günther) ... 2707
3083. atricaudus (Jordan & Gilbert) ... 2707
3084. leei, Jordan & Bollman ... 2708
3085. plagusia (Bloch & Schneider) ... 2709
3086. plagiusa (Linnæus) ... 2710
3087. pusillus (Goode & Bean) ... 2710
3088. diomedeanus (Goode & Bean) ... 2711
3089. williamsi, Jordan & Culver ... 2711
Subgenus Acedia, Jordan ... 2712
3090. nebulosus (Goode & Bean) ... 2712
ORDER CC. PEDICULATI ... 2712
Family CCXXI. Lophiidæ ... 2713
Genus 1058. Lophius (Artedi) Linnæus ... 2713
3091. piscatorius, Linnæus ... 2713
Genus 1059. Lophiomus, Gill ... 2714
3092. setigerus (Vahl) ... 2714
Family CCXXII. Antennariidæ ... 2715
Genus 1060. Pterophryne, Gill ... 2715
3093. histrio (Linnæus) ... 2716
3094. gibba (Mitchill) ... 2717
Genus 1061. Antennarius, Lacépède ... 2717
3095. inops, Poey ... 2718
3096. principis (Cuvier & Valenciennes) ... 2719
3097. tenebrosus (Poey) ... 2719

CLASS III. PISCES—Continued.
ORDER CC. PEDICULATI—Continued.
Family COXXII. Antennariidæ—Continued. Page.
3098. reticularis, Gilbert ... 2719
3099. strigatus, Gill ... 2720
3100. sanguineus, Gill ... 2721
3101. ocellatus (Bloch & Schneider) ... 2721
3102. scaber (Cuvier) ... 2722
3103. tigris, Poey ... 2723
3104. nuttingii, Garman ... 2723
3105. multiocellatus (Cuvier & Valenciennes) ... 2724
3106. radiosus, Garman ... 2725
Genus 1062. Chaunax, Lowe ... 2726
3107. pictus, Lowe ... 2726
3108. nuttingii, Garman ... 2727
Family COXXIII. Ceratiidæ ... 2727
Genus 1063. Ceratias, Kröyer ... 2729
3109. holbolli, Kröyer ... 2729
Genus 1064. Mancalias, Gill ... 2729
3110. uranoscopus (Murray) ... 2729
3111. shufeldti (Gill) ... 2730
Genus 1065. Cryptopsaras, Gill ... 2731
3112. couesii, Gill ... 2731
Genus 1066. Oneirodes, Lütken ... 2732
3113. escrichtii, Lütken ... 2732
Genus 1067. Himantolophus, Reinhardt ... 2732
3114. grœnlandicus, Reinhardt ... 2733
Genus 1068. Corynolophus, Gill ... 2733
3115. reinhardti (Lütken) ... 2733
Genus 1069. Liocetus, Günther ... 2733
3116. murrayi (Günther) ... 2733
Genus 1070. Linophryne, Collett ... 2734
3117. lucifer, Collett ... 2734
Genus 1071. Caulophryne, Goode & Bean ... 2734
3118. jordani, Goode & Bean ... 2735
Family COXXIV. Ogcocephalidæ ... 2735
Genus 1072. Ogcocephalus, Fischer ... 2736
3119. vespertilio (Linnæus) ... 2737
3120. nasutus (Cuvier & Valenciennes) ... 2737
3121. radiatus (Mitchill) ... 2738
Genus 1073. Zalieutes, Jordan & Evermann ... 2738
3122. elater (Jordan & Gilbert) ... 2738
Genus 1074. Halieutichthys, Poey ... 2739
3123. aculeatus (Mitchill) ... 2739
3124. caribbæus, Garman ... 2741
Genus 1075. Halieutæa, Cuvier & Valenciennes ... 2741
3125. spongiosa, Gilbert ... 2741
Genus 1076. Halieutella, Goode & Bean ... 2742
3126. lappa, Goode & Bean ... 2742
Genus 1077. Dibranchus, Peters ... 2743
3127. atlanticus, Peters ... 2743

LIST OF NEW NAMES.

The following is a list of the new generic, subgeneric, specific, and subspecific names which appear as new in Part III of the present work:

	Page.
Sicydium vincente, Jordan & Evermann	2207
Enypnias, Jordan & Evermann	2231
Gnathypops snyderi, Jordan & Evermann	2285
Dactyloscopus zelotes, Jordan & Gilbert	2303
Dæctor, Jordan & Evermann	2325
Bryssophilus, Jordan & Evermann	2329
Gobiesox gyrinus, Jordan & Evermann	2331
Corallicola, Jordan & Evermann	2369
Blenniolus, Jordan & Evermann	2386
Homesthes, Gilbert	2394
Homesthes caulopus, Gilbert	2394
Scarticnthys, Jordan & Evermann	2395
Ophioblennius steindachneri, Jordan & Evermann	2401
Emblemaria atlantica, Jordan & Evermann	2402
Enedrias, Jordan & Gilbert	2414
Embryx, Jordan & Evermann	2458
Lycias. Jordan & Evermann	2461
Emphycus, Jordan & Evermann	2552
Albatrossia, Jordan & Evermann	2573
Bogoslovius, Jordan & Evermann	2574
Hippoglossoides hamiltoni, Jordan & Gilbert	2616
Verasper, Jordan & Gilbert	2618
Verasper moseri, Jordan & Gilbert	2619
Ramularia, Jordan & Evermann	2633
Perissias, Jordan & Evermann	2667
Carcharhinus cerdale, Gilbert	2746
Carcharhinus velox, Gilbert	2747
Myliobatis asperrimus, Gilbert	2754
Aspistor, Jordan & Evermann	2763
Galeichthys xenauchen, Gilbert	2777
Tachysurus emmelane, Gilbert	2785
Aztecula, Jordan & Evermann	2799
Notropis chamberlaini, Evermann	2800
Notropis louisianæ, Evermann	2801
Pisoodonophis daspilotus, Gilbert	2803
Muræna clepsydra, Gilbert	2805
Stolephorus rastralis, Gilbert & Pierson	2811
Stolephorus mundeolus, Gilbert & Pierson	2812
Stolephorus naso, Gilbert & Pierson	2813
Stolephorus starksi, Gilbert & Pierson	2813
Cetengraulis engymen, Gilbert & Pierson	2815
Argyrosomus alascanus, Scofield	2817
Osmerus albatrossis, Jordan & Gilbert	2823
Bathylagus milleri, Jordan & Gilbert	2825

Page.
Zaphotias, Goode & Bean 2826
Characodon garmani, Jordan & Evermann 2831
Siphostoma sinaloæ, Jordan & Starks 2838
Rhynchias, Gill 2841
Oligoplites mundus, Jordan & Starks 2844
Hemicaranx zelotes, Gilbert 2845
Ulocentra meadiæ, Jordan & Evermann 2852
Lobotes pacificus, Gilbert 2857
Porocottus bradfordi, Rutter 2862
Sigmistes Rutter 2863
Sigmistes caulias, Rutter 2863
Crystallichthys, Jordan & Gilbert 2864
Crystallichthys mirabilis, Jordan & Gilbert 2865
Allinectes, Jordan & Evermann 2866
Prognurus, Jordan & Evermann 2866
Prognurus cypselurus, Jordan & Gilbert 2866
Sicyosus, Jordan & Evermann 2867
Alectrias, Jordan & Evermann 2869
Furcimanus, Jordan & Evermann 2869
Salmo clarkii tahoensis, Jordan & Evermann 2870
Oligocottus snyderi, Greeley 2871
Flammeo, Jordan & Evermann 2871
Paralichthys magdalenæ, Abbott 2872
Paralichthys sinaloæ, Jordan & Abbott 2872

THE FISHES

OF

NORTH AND MIDDLE AMERICA.

BY DAVID STARR JORDAN AND BARTON WARREN EVERMANN.

PART III.

PREFATORY NOTE.

This volume is the third of a descriptive catalogue of the fishes and fish-like vertebrates of North and Middle America. For the sake of greater completeness the marine fishes of the Galapagos Islands and the South American coast north of the equator have been included, as all of these are sure, sooner or later, to be found within our limits. For the same reason the few species known from Kamchatka and the Kuril Islands are included as a part of the fauna of the Alaskan Sea.

The pagination and the numbering of the species, genera, and higher groups are continuous throughout the three parts.

Part I, *Branchiostomatidæ* to *Priacanthidæ* inclusive (pages 1 to 1240), was published October 3, 1896; Part II, *Lutianidæ* to *Cephalacanthidæ* inclusive (pages 1241 to 2183), was published October 3, 1898; and Part III, *Callionymidæ* to *Ogcocephalidæ* appears on November 26, 1898. Parts I, II, and III have each their own table of contents, while in Part IV (the Atlas) is given a table of contents complete for the entire work and corrected to include the Addenda.

The present part includes also an artificial key to the families of true fishes, an addendum containing species overlooked or described subsequently to the publication or casting of the part to which they belong, a glossary of scientific terms, and a general index complete for the entire work.

A fourth volume, or Atlas of plates, containing illustrations of one or more species of each of the more important genera, will follow within the year.

The preparation of the manuscript for this work was begun by the senior author in 1891. In 1893 the junior author became associated with him, and since then both have given to it such of their time and energy as could be spared from engrossing official duties to which systematic ichthyology bears no relation.

The insertion of the comma between generic and specific names and the authorities for them, as practiced in this publication, is in accordance with the views held by the authorities of the United States National Museum, and does not express the views of the authors of this work.

Class PISCES—Concluded.

Subclass TELEOSTOMI—Concluded.

Order BB. ACANTHOPTERI—Concluded.

Group GOBIOIDEA.

(The Gobies.)

Body elongate, variously scaled or naked; head usually large, armed or not, the suborbital ring without a bony stay for the preopercle; gill openings reduced, the membranes attached to the isthmus. Gills 4, a slit behind the last; pseudobranchiæ present. Ventral rays I, 4 or I, 5, inserted below pectoral, the fins close together or united or widely separated or otherwise peculiar; dorsal fins separate or united, the first of a few weak spines, sometimes wanting; anal rather long, usually with a single weak spine, similar to soft dorsal; caudal rounded. Usually no air bladder nor pyloric cæca. Vertebræ 24 to 35. Carnivorous bottom fishes, mostly of small size in warm regions, some marine, others of the fresh waters. Two families.

a. Ventral fins widely separated; preopercle strongly armed; lateral line present. Callionymidæ, clxxxvii.

aa. Ventral fins close together, usually united; preopercle with a weak spine or none; no lateral line. Gobiidæ, clxxxviii.

Family CLXXXVII. CALLIONYMIDÆ.

(The Dragonets.)

Body elongate, naked; head usually broad and depressed; the mouth narrow, the upper jaw very protractile; teeth very small, in jaws only; preopercle armed with a strong spine, which is usually branched. Eyes moderate, usually directed upward. Lateral line present, often duplicated. Dorsal fins 2, the anterior with 3 or 4 flexible spines; soft dorsal and anal short, the latter without distinct spine; ventrals I, 5, widely separated from each other; pectoral fins large. Gill openings small, the membranes broadly attached to the isthmus; gills 4, a slit behind the fourth; pseudobranchiæ present; no air bladder. Vertebræ usually $8+13=21$. Small fishes of the shores of warm seas, chiefly of the old world. Allied to the Gobies, but often resembling the *Cottidæ* in form. Genera 4, species about 30. (*Gobiidæ Callionymina*, Günther, Cat. Fishes, III, 138–152.)

a. Ventrals entire, the outer ray not detached; head depressed; gill opening reduced to a very small foramen on upper surface of head; lateral line single. Callionymus, 799.

799. CALLIONYMUS, Linnæus.

Callionymus, Linnæus, Syst. Nat., Ed. x, 249, 1758 (*lyra*).

This genus includes Dragonets with the ventral fins entire, without detached ray, the gill opening reduced to a small foramen opening upward, and the lateral line single; head triangular, depressed; eyes directed upward; preopercular spine very large; sexual differences strongly marked. Species numerous, living on sea bottoms at some depth. (κάλλις, beauty; ὄνομα, name.)

a. Dorsal rays IV, 8 or 9; anal rays 8; some of the dorsal spines filamentous.
  b. Preopercular spine very long, armed with about 9 hooks or spinules; caudal not filamentous. BAIRDI, 2511.
  bb. Preopercular spine strong, bifurcate; caudal fin more or less produced or filamentous. HIMANTOPHORUS, 2512.
aa. Dorsal rays III, 6 or IV, 6; anal rays 4.
  c. Preopercular spine with 2 barbs, the anterior turned forward; body with white spots. CALLIURUS, 2513.
  cc. Preopercular spine with 3 teeth above, ending in an acute point. PAUCIRADIATUS, 2514.

2511. CALLIONYMUS BAIRDI, Jordan.

Head $3\frac{1}{5}$; depth $9\frac{1}{3}$. D. IV, 9; A. 8. Body long and low, very slender, the head much depressed, the least depth of the caudal peduncle about equal to the diameter of the eye. Head triangular as viewed from above, its breadth $\frac{3}{5}$ its length, exclusive of the preopercular spine. Snout bluntish as seen from above, sharp in profile, its outline straight and moderately steep until above the eyes; profile behind the eyes considerably depressed. Snout $2\frac{3}{5}$ in head to gill opening; eye 4; mouth small, inferior, the maxillary reaching front of eye, as long as snout; lower lip conspicuous. Teeth slender, in villiform bands in both jaws, none on vomer. Interorbital area a simple narrow ridge. Bones of head behind eyes rugose; a low rough tubercle of bare bone above the temporal region on each side, somewhat behind each eye. Preopercular spine very long, as long as eye, its exterior ridge with a single antrorse spinule at its base, its posterior edge with 8 conspicuous hooks turned forward and inward, these growing progressively smaller from the second. Gill opening reduced to a pore at upper posterior angle of opercle, its width rather less than that of pupil. Dorsal spines strong, the first ending in a slender filament, the whole as long as head; second and third spines broken (probably each with a short filament in life, as a short filament is still present on the fourth spine); fourth spine well behind third (leaving room for another spine, although no trace of such spine is present); soft dorsal high, most of its rays slightly filamentous at tip, the longest about $\frac{3}{5}$ head; caudal subtruncate, not filamentous, about as long as head to base of preopercular spine; anal fin rather high, the length of its base 3 in body; pectorals about as long as ventrals, each as long as head without preopercular spine. Lateral line single. Color light grayish, mottled or spotted with yellowish and dark brown; cheeks with steel-bluish spots; first dorsal with dusky reticulations around pale gray spots; second dorsal and caudal with nar-

row dusky cross streaks; anal with its posterior half chiefly black, the anterior pale; ventrals black; pectorals pale. Type, a specimen 4½ inches long, in good condition, from the "spewings" of a Snapper or a Grouper (*Neomænis aya* or *Epinephelus morio*), taken on the Snapper Banks, between Pensacola and Tampa; 1 other specimen known. ("I have named this species for Prof. Spencer F. Baird, to whom I have been indebted for aids of many kinds in connection with my studies of American fishes." Jordan.)

Callionymus bairdi, JORDAN, Proc. U. S. Nat. Mus. 1887, 501, Snapper Banks off Pensacola. (Type, No. 39300. Coll. Silas Stearns.)

2512. CALLIONYMUS HIMANTOPHORUS, Goode & Bean.

Head 3⅓; depth of head equal to length of its postorbital portion or to greatest depth of body. Greatest depth of body at the head and the anterior portion of the trunk. D. IV, 8; A. 8; P. 19; V. I, 5. Body slender, moderately elongate, fins all well developed, the tail tapering and with some of its rays produced into a filament. Caudal peduncle very slender, the least height of tail scarcely more than ¼ greatest height of body. Profile descending very rapidly at snout. Mouth small and the intermaxillary very protractile, but may be almost entirely concealed under the preorbitals. Intermaxillary reaching to front of orbit. Maxillary a roundish, slender bone, extending backward to end of intermaxillary. Mandible about as long as eye, extending to vertical through front of pupil. Teeth in villiform bands on intermaxillary and mandible. Interorbital space very narrow, less than ¼ length of eye, which is 1⅓ times as long as snout and nearly ⅓ of total without caudal. A strong bifurcated spine at angle of the preoperculum extending backward slightly beyond the gill opening; length of this spine at its upper articulation ⅔ length of eye. Gill opening reduced to a small slit, placed at a distance behind eye about equaling length of eye and above median line of body. Skin naked. Lateral line abruptly arched over gill opening and connected across nape with its fellow of the opposite side. Spinous dorsal somewhat elevated in front, the first spine nearly twice as long as last, its length about ⅛ total length of caudal; sixth and seventh rays longest, their length nearly equaling that of base of fin; caudal consisting of 4 simple and 8 divided rays; of the divided rays the fifth and sixth are the longest, the lower portion of the fifth and the upper portion of the sixth being produced into a filament, making these rays as long as the distance from the tip of the intermaxillary to the fourth anal ray. It is worthy of remark that in another example of the same species and of about the same size as the type, the sixth of the divided rays alone contributes to form the filament; and in a young example, about ⅓ as large as the type, the first dorsal spine when laid back reaches to the end of soft dorsal. Some of the numerous examples of this species have none of the caudal rays much produced, even in large individuals. Anal fin beginning directly under third ray of soft dorsal, its rays increasing in length to the sixth, which is the longest and twice as long as the first, its length 5⅔

in total without caudal. All the rays simple except the last, which is divided. The pectoral beginning under middle of spinous dorsal and extending to below the fifth ray of the soft dorsal, its rays all simple. The ventral base overlapping lower extremity of pectoral base, its origin under the gill opening. The fourth and longest ray equaling ⅓ of total length without caudal. A small but distinct anal papilla. Color generally light brown, the back with numerous narrow streaks and blotches of slightly darker brown; a dark blotch on membrane between the third and fourth dorsal spines, in some cases occupying nearly all the membrane, in other cases more limited and nearly elliptical in shape; anal with a broad subvertical dark band, the tips of rays and a small area of the membrane behind each ray pale; the lower caudal lobe with a narrow submarginal dark band; ventral with 2 indistinct narrow dark bands on its outer half. From Blake Station XXX, off Barbados, in 209 fathoms; Station CLXXX, at 137 fathoms; Station XXXIII, off Santa Cruz, at 115 fathoms; Station 2 CCXVI, at 119 fathoms; Station CCXXX, at 84 fathoms. (*ἵμας*, whip; *φορέω*, bear.)

Callionymus himantophorus,* GOODE & BEAN, Ocean. Ichth., 296, pl. LXXVI, figs. 268, 268a, b, 1896, off Barbados.

2513. CALLIONYMUS CALLIURUS, Eigenmann & Eigenmann.

Head 3½ to tip of opercular spine (5 in total); depth 7 (9). D. IV, 6; A. 4. Body flat below, the ventral surface bordered on each side with a fold of skin which is wider than the pupil; a single lateral line; diameter of eye equaling length of snout, 3½ in head; maxillary not extending to eye; preopercular spine with 2 barbs above, the anterior one larger and turned forward; gill opening a minute foramen opening upward. The last dorsal ray equaling length of head, and the first dorsal spine reaching its tip when the fin is depressed; ventral fins connected by a broad membrane to the middle of the outer pectoral region; pectoral fins as long as the head. Cheeks, opercles, connecting membrane of ventral fins and antepectoral region with milk-white spots; lower jaw black near the rictus; a series of black dots on branchiostegal membranes, 1 or 2 similar dots in front of pectorals, 2 on the cheek forming a series with the second branchiostegal spot; 4 black spots on the marginal membrane of the belly, other black spots above it; lower half of body with numerous dirty white spots; pectorals transparent, ventrals dusky; membrane of anal sprinkled with minute black points aggregated into black spots in places, and with opaque white spots; caudal transparent, having minute points, its upper half with opaque milk-white bars running obliquely downward and backward from ray to ray; lower half with interrupted longitudinal lines of opaque white, alternating with black spots; dorsal transparent, with white and dark dots most conspicuous between last rays; body marbled with light and darker. Key West, Florida; 1 specimen dredged in 5 fathoms. (*κάλλος*, beauty; *οὐρά*, tail.)

Callionymus calliurus, EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 76, South Beach, Key West. (Type, No. 26265. M. C. Z.)

* The species was listed by Eigenmann, Proc. Cal. Ac. Sci., 2d ser. 1, 78, as "*Callionymus agassizii*, Goode & Bean," a name only, accompanied by no description.

2514. CALLIONYMUS PAUCIRADIATUS, Gill.

"D. III, 6; A. 4. The preopercular spine is armed with three teeth above and terminates in an acute point." (Gill.) Matanzas, Cuba; an imperfectly described species, known only from the above note. (*pauci,* few; *radiatus,* rayed.)

Callionymus pauciradiatus, Gill, Ann. Lyc. Nat. Hist. N. Y., VIII, 1865, 143, Matanzas, Cuba.

Family CLXXXVIII. GOBIIDÆ.

(The Gobies.)

Body oblong or elongate, naked or covered with ctenoid or cycloid scales. Dentition various, the teeth generally small; premaxillaries protractile; suborbital without bony stay. Skin of head continuous with covering of eyes. Opercle unarmed; preopercle unarmed or with a short spine; pseudobranchiæ present. Gills 4, a slit behind the fourth; gill membranes united to the isthmus, the gill openings thus restricted to the sides. No lateral line. Dorsal fins separate or connected, the spinous dorsal least developed, of 2 to 8 flexible spines, rarely wanting; anal usually with a single weak spine, similar to the soft dorsal; ventral fins close together, separate or fully united, each composed of a short spine and 5 (rarely 4) soft rays, the inner rays longest; the ventral fins, when united, form a sucking disk, a cross fold of skin at their base completing the cup; caudal fin convex; anal papilla prominent. No pyloric cæca; usually no air bladder. Carnivorous fishes, mostly of small size, living on the bottoms near the shores in warm regions. Some inhabit fresh waters, and others live indiscriminately in either fresh or salt water. Many of them bury in the mud of estuaries. Few of them are large enough to be of much value as food. Genera about 80; species nearly 600. The species are for the most part easily recognized, but their arrangement in genera is a matter of extreme difficulty. Until the multitude of Asiatic forms are critically studied, any definition of the American genera must be tentative only. (*Gobiidæ,* part; groups *Gobiina, Amblyopina,* and *Trypauchenina,* Günther, Cat. Fishes, III, 1–138.)

Analysis of Genera of North American Gobiidæ.

a. Ventral fins separate; body scaly.

Oxymetopontinæ:

b. Ventral rays I, 4.

c. Forehead bluntly rounded, without sharp keel; tongue very slender, sharp; body elongate, compressed, covered with very small scales; head short, compressed, rather broad above; mouth oblique, the lower jaw projecting; teeth in few series, some of them canine-like; isthmus narrow. Dorsals separate, the first of 6 slender spines; soft dorsal and anal elongate; caudal lanceolate. Ioglossus, 800.

Eleotridinæ:

bb. Ventral rays I, 5.

d. Vomer with a broad patch of villiform teeth; gill openings extending forward to below posterior angle of mouth, the isthmus thus very narrow;

teeth villiform, the outer scarcely enlarged; vertebræ 12 + 13 (*dormitor*); skull above with conspicuous elevated ridges, one of these bounding the orbit above, the orbital ridges connected posteriorly above by a strong cross ridge; a sharp longitudinal ridge on each side of the occipital, the two nearly parallel, the post-temporals being attached to the posterior ends. Insertions of post-temporals widely separated, the distance between them greater than the rather narrow interorbital width; the post-temporal bones little divergent; top of head depressed, both before and behind the cross ridge between eyes; a flattish triangular area between this and the little elevated supraoccipital region; preopercle without spines; lower pharyngeals with slender, depressible teeth, and without lamelliform appendages; scales of moderate size, ctenoid. PHILYPNUS, 801.

dd. Vomer without teeth; isthmus broad; gill openings scarcely extending forward below to posterior angle of preopercle; skull without crests.

e. Body scaly, both anteriorly and posteriorly.

f. Lower pharyngeal teeth stiff and blunt; the bones with an outer series of broad flexible lamelliform appendages, which are rudimentary gill filaments; body short and elevated; teeth slender, those in the outer row scarcely larger, and movable; top of head without raised crests, flattish, its surface uneven; post-temporal bones rather strongly diverging, the distance between their insertions about ⅓ the broad flattish interorbital space; no spine on preopercle or branchiostegals; scales large, ctenoid. Species herbivorous. DORMITATOR, 802.

ff. Lower pharyngeals normal, subtriangular, the teeth stiff, villiform, no lamelliform appendages; scales of moderate or small size; body oblong or elongate.

g. Body moderately robust, the depth 4 to 5½ times in the length to base of caudal; scales ctenoid; cranium without distinct median keel; a small supraoccipital crest.

h. Post-temporal bones little divergent, not inserted close together, the distance between their insertions greater than the moderate interorbital space, or 3⅔ in length of head; top of skull little gibbous; lower pharyngeals narrower than in *Eleotris;* preopercle without spine; scales very small, about 110 in a longitudinal series. Vertebræ 11 + 13; teeth moderate, the outer series on lower jaw enlarged. GUAVINA, 803.

hh. Post-temporal bones very strongly divergent, their insertions close together, the distance between them about ⅔ the narrow interorbital space, and less than ⅐ length of head; top of skull somewhat elevated and declivous; interorbital area somewhat convex transversely; lower pharyngeals rather broad, the teeth bluntish; preopercle with partly concealed spine directed downwards and forward at its angle; scales moderate, 45 to 60 in a longitudinal series; vertebræ (*pisonis*) 11 + 15; teeth small. ELEOTRIS, 804.

gg. Body very slender, elongate, the depth 8 to 9 times in length to base of caudal; scales very small, cycloid.

i. Preopercle with a partly concealed antrorse hook at its angle; caudal with numerous accessory rays at base. ALEXURUS, 805.

ii. Preopercle without spine; caudal without many accessory rays at base; post-temporal bones short, strongly divergent, the distance between their insertions about equal to the narrow interorbital space, or about ⅓ length of head; top of head with a strong median keel, which is highest on the occipital region; no supraoccipital crest; mouth very oblique; the teeth small. EROTELIS, 806.

ee. Body naked on the anterior part; head naked; lower jaw with 4 larger recurved teeth. GYMNELEOTRIS, 807.

eee. Body entirely naked. CHRIOLEPIS, 808.

aa. Ventral fins united.

j. Dorsal fins separate, free from caudal.

SICYDIINÆ:

k. Ventral disk short, adnate to belly; body subcylindrical, covered with ctenoid scales; lips very thick; upper teeth mostly small and movable, lower fixed; dorsal spines 6.

l. Teeth simple; no canines in front of lower jaw. SICYDIUM, 809.

ll. Teeth trifid (or bifid); no canines in front of lower jaw. COTYLOPUS, 810.

GOBIINÆ:

kk. Ventral disk free from the belly.

m. Dorsal spines 4 to 8; eyes well developed.

n. Teeth emarginate, uniserial, those of the lower jaw nearly horizontal; dorsal spines 6; scales large, ctenoid; gill openings moderate. EVORTHODUS, 811.

nn. Teeth simple.

o. Body scaly, more or less.

p. Maxillary normal, not prolonged behind the rictus; skull of the usual gobioid form, comparatively short and abruptly broadened behind the orbits; occiput depressed; supraoccipital and temporal ridges continuous.

q. Dorsal spines 6; scales evidently ctenoid; head naked (the nape scaly as usual.)

r. Interorbital area anteriorly elevated, with a large foramen-like depression in front of eye; body short, compressed, formed much as in *Dormitator;* nape with a fleshy crest; scales large. Vertebræ 11 + 15. LOPHOGOBIUS, 812.

rr. Interorbital area not elevated in front; body more elongate; no fleshy nuchal crest; isthmus broad.

s. Inner edge of shoulder girdle without fleshy cirri or papillæ; cranium anteriorly short; interorbital space narrower, grooved, with a low median ridge or none; median crest on cranium low.

t. Body scaly anteriorly and posteriorly (sometimes a naked strip on back or belly). Vertebræ 12 + 16 to 10 + 15. GOBIUS, 813.

tt. Body entirely naked anteriorly, the posterior half scaled; scales moderate or small. GARMANNIA, 814.

ss. Inner edge of shoulder girdle with 2 or 3 conspicuous dermal flaps; preorbital region very long; premaxillary and maxillary strong; interorbital groove with a conspicuous median crest; scales rather small (45 to 70.) AWAOUS, 815.

qq. Dorsal spines 7 or 8 (very rarely 6, especially in *Eucyclogobius*.)

u. Scales large, ctenoid; shoulder girdle without dermal flaps.

v. Sides of head scaled; soft dorsal and anal rather short, of 11 to 14 rays each; deep-water species. BOLLMANNIA, 816.

vv. Sides of head naked; soft dorsal and anal short, of 10 to 12 rays each; shore species. ABOMA, 817.

uu. Scales very small, cycloid or nearly so.

w. Inner edge of shoulder girdle without fleshy processes; head naked; body more or less compressed; mouth very oblique; teeth strong; interorbital groove with or without a median ridge. Vertebræ 11 + 15 or 16; soft dorsal and anal long, of 15 to 17 rays each.

x. Body chiefly scaly, anteriorly as well as posteriorly. MICROGOBIUS, 818.

xx. Body naked anteriorly, scaled posteriorly. ZALYPNUS, 819.

ww. Inner edge of shoulder girdle with 2 or 3 dermal flaps, or processes, as in *Awaous*.

y. Head naked, the interorbital groove with the median ridge high, not extending forward to orbit; body rather robust; soft dorsal and anal short; fresh-water species. EUCYCLOGOBIUS, 820.

yy. Head scaled like the body; the interorbital groove with the median ridge little developed; soft dorsal and anal long; body elongate; marine species. LEPIDOGOBIUS, 821.

pp. Maxillary much produced backward, extending beyond the gill opening in the adult; skull comparatively long, gradually (not abruptly) broadened behind orbits; median crest of cranium well developed; scales small, cycloid; head naked, occipital region narrowed forward; supraorbital and temporal crests not continuous.

z. Occiput depressed, with a blunt median keel.

a'. Shoulder girdle without dermal flaps; dorsal spines 6; soft dorsal and anal short; mouth very large; isthmus broad; vertebræ 14 + 16 (*mirabilis*).
GILLICHTHYS, 822.

aa'. Shoulder girdle with 1 to 3 small dermal flaps on the inner edge; dorsal spines 5; soft dorsal and anal long.
QUIETULA, 823.

zz. Occiput transversely rounded without median keel.

b'. Shoulder girdle with 1 to 3 small dermal flaps on its inner edge; dorsal spines 5; soft dorsal and anal long. ILYPNUS, 824.

bb'. Shoulder girdle without dermal flaps; dorsal spines 4 or 5; soft dorsal and anal long. CLEVELANDIA, 825.

oo. Body and head entirely naked.

c'. Dorsal spines 4; body long and slender; mouth large, the lower jaw projecting; no barbels; soft dorsal and anal long; male with ornate colors.
EVERMANNIA, 826.

cc'. Dorsal spines 7 (rarely 6).

d'. Chin without barbels; mouth small, little oblique; body robust, soft dorsal and anal short. GOBIOSOMA, 827.

dd'. Chin with a fringe of short barbels; mouth terminal, oblique; soft dorsal and anal very short. BARBULIFER, 828.

CRYSTALLOGOBIINÆ:

mm. Dorsal spines 2 (or 1); body wholly naked.

e'. Eyes reduced to small rudiments; interorbital area forming a sharp median range; skull rather abruptly widened behind orbits; anterior portion of skull unusally long; no flaps on shoulder girdle; skull highest at nape, depressed above the eyes; soft dorsal and anal short.
TYPHLOGOBIUS, 829.

GOBIOIDINÆ:

jj. Dorsal fin continuous, the soft part and the anal joined to base of caudal; eyes minute; body elongate; scales minute or wanting; mouth very oblique, the lower jaw projecting; gill openings moderate.

f'. Dorsal rays VI, 16 to 23; anal rays 17 to 23.

g'. Teeth small, in a single series; scales present. TYNTLASTES, 830.

gg'. Teeth in a band, those of the outer series being very strong; scales present.

h'. Body entirely scaled. GOBIOIDES, 831.

hh'. Anterior part of body naked. CAYENNIA, 832.

800. IOGLOSSUS, Bean.

Ioglossus, BEAN, in Jordan & Gilbert, Proc. U. S. Nat. Mus. 1882, 297 (*calliurus*).

Body elongate, strongly compressed, of equal depth throughout, covered with very small, mostly cycloid, scales. Head short, compressed, not keeled above; mouth large, oblique, the lower jaw projecting; teeth in narrow bands or single series, some of them canine; no teeth on vomer or palatines; tongue very slender, sharp; opercles unarmed. Gill openings

very wide, the membranes narrowly joined to isthmus on median line. No lateral line. Branchiostegals 5. Dorsals separate, the first of 6 very slender, flexible spines; the second elongate, similar to the anal; caudal long and pointed, free from dorsal and anal; ventrals close together, separate, each of 1 spine and 4 rays, their insertion below or behind pectorals; anal papilla present. A remarkable type, belonging to the *Oxymetopontinæ*, differing widely from our other Gobioid fishes. Gulf of Mexico, in rather deep water. (ὄϊος, arrow; γλῶσσα, tongue.)

2515. IOGLOSSUS CALLIURUS, Bean.

Head 5; depth 7 to 7⅓. D. VI–22 to 24; A. I, 21 to 23. Body very elongate, slender, much compressed, of equal depth throughout; head compressed, without osseous crest; mouth very oblique, the lower jaw strongly projecting; premaxillaries in front on the level with pupil; maxillary extending to opposite front of pupil, its length 2¾ in head; upper jaw with a narrow band of about 2 series of conical cardiform teeth, those of the outer row much larger than the others, behind these 2 small conical curved canines; lower jaw with a single row of smaller teeth, behind which are about 4 short canines directed somewhat backward; the posterior pair strongly curved; no teeth on vomer or palatines. Tongue narrow, pointed. Eye large, nearly twice length of snout, 3½ in head, its diameter considerably more than depth of cheek, about ⅓ more than interorbital width; opercles unarmed. Pseudobranchiæ present. Gill openings wide, extending forward below, the membranes attached mesially to the very narrow isthmus, across which they do not form a fold. Gill rakers long and slender. Dorsal fins separated by a short interval, the first of very slender somewhat filamentous spines, the longest about as long as head; second dorsal little more than ½ as high as first, apparently nearly uniform, separated from the caudal by an interval nearly ½ length of head; caudal lanceolate, its middle rays filamentous, about ½ the length of rest of body; anal rather high, similar to soft dorsal; ventrals I, 4, inserted very slightly in advance of base of pectorals, the 2 fins very close together, but apparently quite separate and without basal fold of skin, the fin little longer than head, the inner rays filamentous; pectoral with broad base, about 1¼ in head. Anal papilla very short, midway between tip of snout and base of caudal. Body with very small, nonimbricate, embedded scales, these a little larger and imbricate on the tail; cheeks with embedded cycloid scales; scales very weakly ctenoid, most of them appearing cycloid; no lateral line. Color light olive, everywhere densely punctate; dorsals edged with black; middle of caudal reddish, with paler bluish edgings. Length 4½ inches. Here described from specimens from off Pensacola. Gulf of Mexico; known only from the Snapper Banks off Pensacola, in rather deep water. (κάλλος, beauty; οὐρά, tail.)

Ioglossus calliurus (Bean MS.), in Jordan & Gilbert, Proc. U. S. Nat. Mus. 1882, 297, Pensacola, Florida; Bean, Proc. U. S. Nat. Mus. 1882, 419; Jordan & Gilbert, Synopsis, 949, 1883; Jordan, Proc. U. S. Nat. Mus. 1884, 437; Jordan & Eigenmann, Proc. U. S. Nat. Mus. 1886, 481.

801. PHILYPNUS,* Cuvier & Valenciennes.

(GUAVINAS.)

Gobiomorus,† LACÉPÈDE, Hist. Nat. Poiss., II, 699, 1798, in part (*dormitor*, etc.); restricted to *dormitor* by JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 571; restricted to *Gobiomorus taiboa*, Lacépède (Valenciennes' *strigata*), by GILL, Proc. U. S. Nat. Mus. 1888, 79, in accordance with the law of exclusion.

Philypnus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 255, 1837 (*dormitator*).

Lembus, GÜNTHER, Cat. Fishes, I, 505, 1859 (*maculatus*).

Body elongate, terete anteriorly, compressed behind. Head elongate, depressed above. Mouth large; lower jaw considerably projecting; teeth in jaw rather small, slender, recurved, the outer scarcely enlarged; teeth on vomer villiform, in a broad, crescent-shaped patch; gill openings extending forward to below posterior angle of mouth, the isthmus very narrow. Scales moderate, ctenoid, covering most of the head, 55 to 66 in a longitudinal series. Dorsal with 6 spines and 9 or 10 rays; anal rays I, 9 or 10; ventrals separate. No preopercular spine; insertion of post-temporals almost midway between occipital crest and edge of skull; parietals with a crest running from insertion of post-temporal forward to just behind eye, where they are connected by a thin, high, transverse crest; supraocular with a short, high crest, extending from above front of eye back to posterior edge of orbit, thence extending outward parallel with the transverse crest, leaving a deep groove between them; bony projections before and behind eye prominent. Vertebræ 12 + 13 = 25; lower pharyngeals triangular, with slender teeth. Largest of the Gobies, some of the species reaching a length of 2 or 3 feet and valued as food. Tropical rivers. (*φίλυπνος*, slumber-loving; *φίλος*, loving; *ὕπνος*, sleep.)

a. Coloration rather obscure, the dark lateral band indistinct or wanting; scales 55 to 57. DORMITOR, 2516.

aa. Coloration bright, the black lateral band distinct; scales 52 to 55. LATERALIS, 2517.

2516. PHILYPNUS DORMITOR (Lacépède).

(SLEEPER; GUAVINA.)

Head $2\frac{3}{5}$ to $2\frac{9}{10}$; depth 5 to $5\frac{3}{4}$. D. VI–10; A. I, 9; scales 55 to 57; eye $6\frac{1}{8}$ to $7\frac{1}{2}$ in head; snout $3\frac{2}{5}$; maxillary $2\frac{3}{5}$. Body elongate, terete anteriorly, compressed behind. Head elongate, depressed above. Mouth large; maxillary reaching to middle of pupil. Lower jaw considerably projecting. Teeth on jaws slender, depressible. Interspace between dorsals slightly greater than interorbital width; dorsal spines slender, the second the longest, $2\frac{1}{4}$ in head; length of base of anal about $2\frac{1}{3}$ in head; ventrals reaching $\frac{2}{3}$ of the distance to vent; tips of pectorals reaching ventral.

* The *Eleotrinæ* have been made the subject of a special paper (A Review of the American Eleotridinæ, in Proc. Ac. Nat. Sci. Phila. 1885, 66–80) by Eigenmann & Fordice. The *Gobiidæ* of America have been discussed in detail by Jordan & Eigenmann (Proc. U. S. Nat. Mus. 1886, 477–518) and later by Eigenmann & Eigenmann (Proc. Cal. Ac. Sci., 2d ser., vol. 1, 1888, 51–78). In this paper are valuable notes on the specimens in the Museum of Comparative Zoology.

† For the reasons in favor of the use of the name *Gobiomorus* for *Valenciennea*, Bleeker, instead of using it for the present genus, see GILL, Proc. U. S. Nat. Mus. 1888, 69.

Dark brownish or olive, lighter below; an interrupted dark lateral band extending from base of pectoral to base of caudal (not always present); fins dusky, and with the exception of the anal and ventrals, all distinctly mottled; spinous dorsal margined with blackish; head often with dark spots. Streams of the West Indies and Atlantic shores of Central America, Mexico, and Surinam; everywhere common, reaching a length of 2 feet or more. Here described from Cuban specimens. (*dormitor*, sleeper.)

Guavina, PARRA, Descr. Dif. Piezas Hist. Nat. Cuba, tab. 39, fig. 1, 1787, Havana.
Gobiomorus dormitor, LACÉPÈDE, Hist. Nat. Poiss., II, 599, 1798, Martinique; from a drawing by PLUMIER; EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 52.
Platycephalus dormitator, BLOCH, Ichth., 1801, Martinique; after LACÉPÈDE.
Batrachus guavina, BLOCH & SCHNEIDER, Syst. Ichth., 44, 1801; based on *Guavina* of PARRA.
Eleotris longiceps, GÜNTHER, Proc. Zool. Soc. Lond. 1864, 151, Nicaragua; GÜNTHER, Fish. Centr. Amer., 440, 1869.
Electris dormitatrix, CUVIER, Règne Animal, Ed. II, vol. 2, 246, 1829, Antilles; GÜNTHER, Cat. Fish., III, 119, 1861.
Gobiomorus dormitator, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 572.
Philypnus dormitator, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 255, 1837; POEY, Mem. de Cuba, II, 381, 1860; GIRARD, U. S. and Mexican Boundary Survey, Zool., 27, pl. 12, fig. 13, 1859; JORDAN & GILBERT, Synopsis, 631, 1883.

2517. PHILYPNUS LATERALIS, Gill.

(ABOMA DE MAR.)

Head $2\frac{3}{10}$; depth $5\frac{1}{3}$. D. VI–10; A. 1, 10; scales 52 to 55; eye 6 to $6\frac{1}{2}$ in head; snout $3\frac{1}{2}$ to $3\frac{3}{4}$; maxiliary $2\frac{1}{2}$ to $2\frac{3}{4}$. Brownish, lighter or white below; a distinct dark brown or blackish band extending from base of pectoral to base of caudal; dorsals, pectoral and caudal dusky; ventrals and anal lighter; dorsals, caudal, and in some specimens the anal, distinctly blotched. The only constant difference between this species and *Philypnus dormitor* seems to be the brighter coloration of *lateralis*. Streams of Pacific Coast of Mexico and Central America, from Sonora to Panama, entering the sea; common, reaching a much larger size than any other of our Gobies. Here described from specimens from Rio Presidio, Mazatlan. (*lateralis*, pertaining to the side.)

Philypnus lateralis, GILL, Proc. Ac. Nat. Sci. Phila. 1860, 123, Cape San Lucas (Coll. Xantus); JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 377.
Eleotris lateralis, GUNTHER, Cat., III, 122, 1861.

802. DORMITATOR, Gill.

(PUÑECAS.)

Prochilus, CUVIER, Règne Animal, Ed. 1, vol. II, 294, 1817 (*nacrolepidota* = *maculatus*); name preoccupied.
Dormitator, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 240 (*gundlachi*).

Body short, robust; head broad and flat above; mouth little oblique; maxillary reaching to anterior margin of orbit; lower jaw little projecting; no teeth on vomer; lower pharyngeal teeth stiff and blunt, the bones with an external series broad, flexible, lamelliform, these being rudimentary gill filaments; scales large, ctenoid, 30 to 33 in a longitudinal

series; skull much as in *Eleotris;* D. VII–I, 8; A. I, 9 or 10; no spine on preopercle; post-temporals inserted midway between occipital crest and edge of skull; supraoccipital crest low. (*dormitator,* one who sleeps.)

2518. DORMITATOR MACULATUS (Bloch).

(Guavina Mapo; Pañeca.)

Head 3⅕; depth about 3 in adult. D. VII–I, 8 or 9; A. I, 9 or 10; lateral line 33. Body short, robust; head broad and flat above; eye small, less than snout; caudal a little shorter than head; mouth little oblique; maxillary reaching to anterior margin of orbit; lower jaw little projecting; no teeth on vomer; interspace between dorsals equaling orbit; highest anal ray 1¾ in head; highest dorsal ray 1⅓ in head; skull much as in *Eleotris,* but everywhere broader; no spine on preopercle; post-temporal inserted midway between occipital crest and edge of skull; supraoccipital crest low; scales large, becoming much smaller on belly, 25 series on median line from base of ventrals to vent; 18 series across breast from pectoral to pectoral; 18 on a median line from posterior border of orbit to dorsal. Dark brown, with lighter bluish spots; a faint dark stripe along sides; a conspicuous large dark blue spot edged with black above base of pectorals; a dark streak from eye to angle of mouth; 2 dark streaks on side of head; branchiostegal membrane blackish; dorsals barred with spots; anal dusky, barred with bluish, and with white margin; a dark bar on base of pectoral. Length 1 to 2 feet. Both coasts of America, ranging from South Carolina through the West Indies to Pará, Cape San Lucas, and Panama, in fresh or brackish water; everywhere abundant and used as food. Dr. Eigenmann observes:

There seem to be 2 forms of the adult—one with the profile gibbous, the dorsal outline forming a regular curve; the other having the profile depressed over the eyes, the anterior portion being subhorizontal. The specimens from Gurupa and the Rio Grande have the profile depressed; all the other specimens have a gibbous profile. A comparison in detail of the two forms is appended. Only extreme differences are given.

West Indian specimens 5 to 7⅓ inches.	Rio Grande specimens 5, 6½, and 7¼ inches.
Profile regularly curved from first dorsal spine to snout.	Profile depressed over eye, becoming horizontal anteriorly.
Head 3⅓ to 4; depth 3 to 3¼; depth always greater than length of head.	Head 3; depth 3 to 3⅓; depth usually less than length of head.
Highest anal ray 1⅓ to 1⅔ in head.	Highest anal ray 1⅔ to 2 in head.
Distance from first dorsal spine to snout greater than distance from first dorsal spine to first anal ray.	Distance from first dorsal spine to snout equals distance from first dorsal spine to base of last anal ray.
Scales in median series 29 to 32.	Scales in median series 30 to 34.
Color usually dark brown, a black spot above base of pectoral, a short bar on base of pectoral.	Color gray, a jet-black spot above base of pectoral; a black bar at base of pectoral; a black line from eye to mouth; longitudinal black lines on cheeks and opercles; dark spots on back; some silvery scales on sides.

Among our specimens from Mazatlan are 3 markedly different forms which seem like distinct species. In view of the great variations to which this species is subject we do not, however, regard them as such, especially as none of the three corresponds exactly to the account above given of the 2 Atlantic forms.

I. Deep-bodied Specimens (*Dormitator latifrons*, Richardson).

Head 3; depth 3. D. VII–I, 8; A. I, 8 or 9; scales 30 to 33; eye $4\frac{3}{4}$ to $5\frac{1}{4}$ in head; snout $3\frac{1}{4}$ to $3\frac{1}{2}$ in head; interorbital width $2\frac{1}{5}$ in head; ventrals reaching $\frac{2}{3}$ the distance to vent, $1\frac{1}{2}$ to $1\frac{2}{3}$ in head; highest anal ray $1\frac{3}{4}$ to 2 in head. Body short, robust, the back elevated; head broad and flat above, the anterior profile from first dorsal spine to tip of snout oblique, descending abruptly; mouth oblique, maxillary reaching anterior margin of orbit; lower jaw little projecting. Color greenish, lighter below; body with cross bars of dark brown; fins dusky, the dorsals distinctly blotched with darker; a dark cross bar at base of pectorals; a dark-blue humeral blotch, becoming blackish in spirits; 3 or 4 dark cross bands extending from eye and below eye to posterior margin of preopercle; a dark band extending from below eye to below tip of maxillary. Two specimens from Rio Presidio, Mazatlan.

II. Common Form, at Mazatlan.

Head $3\frac{1}{6}$; depth $3\frac{2}{3}$ to $3\frac{3}{4}$. D. VII–I, 7; A. I, 8; scales 33 or 34; eye $4\frac{1}{4}$ to $4\frac{2}{5}$ in head; snout $3\frac{1}{2}$ to 4 in head; interorbital width 3 to $3\frac{1}{6}$ in head; ventrals reaching about $\frac{2}{3}$ the distance to vent, $1\frac{1}{2}$ in head; highest anal ray $1\frac{5}{6}$ to 2 in head. Body short, compressed, the back little elevated; head rather broad and slightly convex above, the anterior profile from first dorsal spine to tip of snout slightly convex; mouth oblique, maxillary reaching anterior margin of orbit; lower jaw little projecting. Color olive brown, with cross bars of darker brown, lighter below; fins dusky, the dorsals with about 3 darker cross bars; pectorals with a darker cross bar at base; a distinct dark-brown humeral spot slightly larger than eye; 3 or 4 dark cross bands extending from eye and below eye to posterior margin of preopercle; a distinct dark-brown bar extending from below eye to below tip of maxillary; a dark lateral band extending from base of pectoral to base of caudal. Many specimens from Mazatlan.

III. Large-headed Form.

Head $3\frac{1}{6}$; depth $3\frac{1}{4}$. D. VI–I, 8; A. I, 9; scales 32 or 33; eye $4\frac{1}{3}$ in head; snout $3\frac{1}{2}$ in head; interorbital width $2\frac{1}{2}$ in head; ventrals reaching $\frac{4}{7}$ the distance to vent, $1\frac{1}{2}$ in head; highest anal ray 2 in head. Body moderately compressed, the back little elevated; head very broad above, convex; the anterior profile from first dorsal spine to tip of snout oblique, gently descending; mouth oblique, maxillary reaching anterior margin of orbit; lower jaw little projecting. Color brownish, middle of back darker, lighter below; body with darker cross bands; ventrals yellowish; other fins dusky; dorsals with darker blotches; a dark crossbar at base of pectoral; a dark humeral spot; four cross bands extending from eye and below eye to posterior margin of preopercle; a dark band extending from below eye to below top of maxillary; a dark lateral band extending from base of pectoral to base of caudal. One specimen, from near Mazatlan. (*maculatus*, spotted.)

Sciæna maculata, BLOCH, Ichth., pl. 299, fig. 2, 1790, West Indies.
Eleotris mugiloides, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 226, 1837, Martinique; Surinam.
Eleotris sima,* CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 232, 1837, Vera Cruz.
Eleotris latifrons, RICHARDSON, Voy. Sulphur, Fishes, 57, pl. 35, figs. 4 and 5, 1837, locality unknown, supposed to be from Pacific coast, Central America.
? Eleotris grandisquama,† CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 229, 1837, America; locality unknown.
Eleotris somnolentus, GIRARD, Proc. Ac. Nat. Sci. Phila. 1858, 169, near mouth of Rio Grande.
Eleotris omocyaneus, POEY, Memorias, II, 269, 1860, Havana.
Dormitator microphthalmus, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 170, Panama. (Coll. Capt. John M. Dow.)
Dormitator gundlachi, POEY, Synopsis, 396, 1868, Cuba.
Dormitator lineatus, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 271, Savannah.
Dormitator maculatus, JORDAN & GILBERT, Synopsis, 632, 1883; JORDAN & EIGENMANN, *l. c.*, 482; EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci., 2d series, vol. 1, 1888, 52.

803. GUAVINA, Bleeker.

Guavina, BLEEKER, Esquisse d'un Syst. Nat. Gobioid., 302, 1874 (*guavina*).

This genus is allied to *Eleotris*, differing in having the post-temporal bones little divergent, not inserted close together, the distance between their insertions greater than the moderate interorbital space, or $3\frac{4}{5}$ in length of head; top of skull little gibbous; lower pharyngeals narrower than in *Eleotris;* preopercle without spine; scales very small, ctenoid, about 110 in a longitudinal series. Vertebræ 11 + 13; teeth moderate, the outer series on lower jaw enlarged. Fresh waters of the West Indies and Brazil. Two species known; *Guavina brasiliensis* (Sauvage) from Bahia, and the following. (*Guavina*, the Spanish name.)

2519. GUAVINA GUAVINA (Cuvier & Valenciennes).

(GUARUBACO; GUAVINA.)

Head $3\frac{1}{3}$; depth $4\frac{1}{3}$ to $5\frac{1}{4}$. D. VI, or VII–I. 10; A. I, 9 or 10. Body stoutish, oblong; mouth oblique; maxillary reaching opposite middle of eye, its length $2\frac{1}{2}$ to $3\frac{1}{2}$ in head. Lower jaw little projecting; teeth in broad bands, the outer ones on lower jaw enlarged. Scales on head embedded; those on body very small, ctenoid on sides, cycloid on back and belly, 100 to 110 in a longitudinal series. Isthmus very broad. Pectorals reaching to middle of spinous dorsal. Highest anal ray $1\frac{4}{5}$ in head. Post-temporals inserted twice as far from occipital crest as in *Eleotris pisonis*. Parietals ending

* Types, 2 specimens in poor order, from Vera Cruz, 0.09 mm. long. Snout a little more steep and convex than usual in *Dormitator maculatus*. Head $3\frac{1}{3}$ in length; depth $3\frac{1}{4}$. Eye $4\frac{1}{2}$ in head. D VII, 9; A. 11; scales 31–11. Soft dorsal very high, with round black spots. Caudal and anal plain. This seems to be inseparable from *Dormitator maculatus*.

† We have the following note on the type of *Eleotris grandisquama:* Type specimen in fair condition, 0.14 mm. long, from "Amérique Méridionale?" Head slenderer than in *D. maculatus*, and much depressed, its depth at the eyes less than its width, which is less than that of body. Anterior profile almost concave. Caudal fin large; other fins moderate. Dorsal VI, 9; anal I, 9; scales about 29 11. A few dusky spots on dorsal and anal. According to Dr. Eigenmann, specimens of *Dormitator maculatus* from the Rio Grande agree fairly with this type, and it is not likely that it is different.

in a sharp point behind. Preopercular spine none, a broad, thin extension on the lower limb of preopercle taking its place. Lower pharyngeals triangular, normal, rather narrow; the teeth small. Vomer without teeth. Length 1 foot. East coast of tropical America, Cuba to Rio Janeiro, in fresh and brackish waters; very common. (*guavina*, Spanish name.)

Eleotris guavina, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 223, 1837, Martinique.
Guavina guavina, JORDAN & EIGENMANN, *l. c.*, 483.

804. ELEOTRIS (Gronow) Bloch & Schneider.

Eleotris, GRONOW, Zooph., 83, 1763 (nonbinomial).
Eleotris, BLOCH & SCHNEIDER, Syst. Ichth., 65, 1801 (*pisonis*).
Culius, BLEEKER, Esquisse d'un Syst. Nat. des Gobioid., 303, 1874 (*fuscus*).

Body long and low, compressed behind. Head long, low, flattened above, without spines or crests, almost everywhere scaly. Mouth large, oblique, lower jaw projecting. Lower pharyngeals rather broad, the teeth small,* bluntish. Preopercle with a small concealed spine below, its tip hooked forward. Branchiostegals unarmed. Eyes small, high, anterior; isthmus broad. Post-temporal bones very strongly divergent, their insertions close together, the distance between them about $\frac{2}{3}$ the narrow interorbital space, and less than $\frac{1}{7}$ length of head; top of skull somewhat elevated and declivous; interorbital area slightly convex transversely; dorsal fins well apart, the first of 6 or 7 flexible spines; ventrals separate. Scales moderate, ctenoid, 45 to 62 in a longitudinal series; vertebræ (*pisonis*) 11 + 15. Tropical seas, entering fresh waters. (ἠλεός, bewildered.)

a. Teeth subequal, those of inner or outer series enlarged.
b. Cheek entirely scaled.
c. Teeth of inner series of each jaw enlarged.
d. Scales in a median series 40 to 51, in a cross series 12 to 20.
e. Eye large, 5 to 6 in head; scales 40 to 44—12 to 14. AMBLYOPSIS, 2520.
ee. Eye small, 8 in head; scales 51–20. ABACURUS, 2521.
dd. Scales in a median series 57 to 66; in a cross series 18 to 24. PISONIS, 2522.
bb. Lower half of cheek naked; scales 61. PERNIGER, 2523.
aa. Teeth all equal; scales 60. PICTUS, 2524.

2520. ELEOTRIS AMBLYOPSIS (Cope).

Head $3\frac{2}{3}$; depth $4\frac{1}{4}$. D. VI–9; A. I, 8; scales 46 (40 to 44–12 to 14 according to Eigenmann); eye $5\frac{1}{2}$ in head, 2 in interorbital width; preopercular spine strong, decurved; width of head $\frac{3}{5}$ in its length; chin prominent; premaxillary spines forming a projection in profile. Brown, a black spot above at base of pectoral; first dorsal and anal dusky; second dorsal and caudal delicately cross-barred with blackish; 3 black lines from orbit behind and below. Surinam. Described from 3 specimens each 3 inches long. (Cope.) Dr. Eigenmann mentions 15 other examples,

* The characters of the skeleton are taken from *Eleotris pisonis* and have not been verified on other species. The hooked preopercular spine supposed to characterize *Culius* is found on the typical species of *Eleotris*, as well as in *Alexurus*.

the longest 2½ inches long, from Surinam, in the Museum of Comparative Zoology. (ἀμβλύς, blunt; ὄψις, face.)

Electris amblyopsis, COPE, Trans. Amer. Philos. Soc. 1870, 473, Surinam (Coll. Dr. Charles Hering); JORDAN & EIGENMANN, *l. c.*, 483, 1886; EIGENMANN & EIGENMANN, *l. c.*, 55.

2521. ELEOTRIS ABACURUS, Jordan & Gilbert.

Head 3; depth 4⅓. D. VI–9; A. I, 8; scales 51–20; eye 8 in head, 2⅓ in interorbital width; pectoral 1⅓; ventral 1½; highest dorsal ray 2; highest anal ray 2; caudal 1¼. Body slender, compressed, the head depressed, becoming very narrow anteriorly, its width ⅜ its length; a notable depression above orbits, the premaxillary processes protruding before it; lower jaw the longer; maxillary reaching vertical behind pupil, 2⅜ in head. Teeth in jaws in narrow villiform bands, becoming a single series on sides of lower jaw, those of the outer and inner series in each jaw somewhat enlarged, the largest being a single series in sides of lower jaw. Preopercular spine as usual in the genus. Scales smooth above and below, ctenoid on sides. Color in spirits, brown, lighter above and below; each scale on middle of sides with a dusky streak, these forming obscure lengthwise lines; back anteriorly with a few small black spots; under parts, including sides of head, very thickly punctulate with black; no dark stripes from orbit; lips black; a dark streak from snout through eye to upper angle of preopercle; 2 dusky streaks from eye downward and backward across cheek; a very conspicuous black blotch as large as eye in front of upper pectoral rays; pectorals and ventrals transparent, dusky; vertical fins all barred with light and dark in fine pattern. Coast of South Carolina. Known from a single specimen, 4 inches long, taken in the harbor of Charleston. This species agrees very well with Cope's account of *Culius amblyopsis*, but the scales are larger, the eye is smaller, and there is some difference in color, besides the remote habitat. (ἄβακος, checker; οὐρά, tail.)

Culius amblyopsis, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 610; not of Cope.
Eleotris abacurus, JORDAN & GILBERT, Proc. Cal. Ac. Sci. 1896, 228, Charleston. (Coll. Dr. C. H. Gilbert. Type, No. 2009, L. S. Jr. Univ. Mus.)

2522. ELEOTRIS PISONIS (Gmelin).

(GUAVINA TÉTARD; SLEEPER.)

Head 3 to 3⅓ in body; depth 4⅓ to 5. D. VI–9; A. I, 8; scales 62; eye 5¼ to 8 in head; maxillary 2⅔; pectoral 1½; ventral 2; caudal 1½. Body not much compressed; head somewhat depressed; mouth rather large, the maxillary reaching to below posterior margin of pupil; lower jaw much projecting, a knob at symphysis; wide bands of villiform teeth in jaws, none on vomer or palatines; interorbital region nearly twice as wide as the horizontal diameter of eye; top of head, cheeks, and opercles covered with small scales; a stout, concealed spine projecting downward on edge of preopercle. Origin of dorsal about midway between tip of snout and end of last dorsal rays; tips of first dorsal spines not reaching front of

second dorsal when fin is depressed; origin of anal a little behind that of soft dorsal; pectorals reaching to posterior spine of first dorsal; ventrals inserted very slightly behind base of pectorals; caudal peduncle as wide as length of maxillary. Color brownish; fins with dark spots and wavy lines; ventrals dusky; 2 dark stripes behind the orbit. Here described from specimens, 6 or 7 inches long, collected in the Rio Almendares, Cuba, by Dr. Jordan. Streams of the West Indies, generally common from southern Florida to Rio Janeiro. Dr. Eigenmann enumerates many specimens from various localities in Brazil. (Named for Dr. William Piso, of the University of Leyden, associate of George Marcgraf and Prince Maurice of Nassau, in 1648, in the study of the natural history of Brazil.)

Amore pixuma, MARCGRAVE & PISO, Hist. Brasil., IV, 166, 1648, Brazil.
Eleotris capite plagioplateo, GRONOW, Mus. Ichth., II, 168, 1757; after MARCGRAVE.
Gobius pisonis, GMELIN, Syst. Nat., 1206, 1788; based on *Eleotris* of GRONOW.
Gobius amorea, WALBAUM, Artedi Piscium, III, 205, 1792; based on *Eleotris* of GRONOW.
Eleotris gyrinus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 220, pl. 356, 1837, Martinique; San Domingo; Surinam.
Eleotris (*Culius*) *belizianus*,* SAUVAGE, Bull. Soc. Philom. Paris 1879 (1880), 55, Belize (Coll. Morelet), Cayenne (Coll. Mélinon); EIGENMANN & FORDICE, Proc. Ac. Nat. Sci. Phila. 1885, 75; EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 55.
Eleotris pisonis, JORDAN & EIGENMANN, *l. c.*, 483; EIGENMANN & EIGENMANN, *l. c.*, 55.

2523. ELEOTRIS PERNIGER (Cope).

Head $4\frac{1}{4}$; depth $4\frac{2}{3}$. D. VI–I, 9; A. I, 9; scales 61; eye 3 in interorbital width; no vomerine teeth. A strong spine at posterior angle of preoperculum, directed downward. Premaxillary spines not prominent in profile; scaling of vertex extending to their extremities. Longitudinal diameter of orbit $\frac{1}{6}$ length of head. Color black, abdomen brown, fins dusky; first dorsal with white extremity and 2 longitudinal black bars, 1 along the base; other fins with small black bars; [no] maxillary or caudal spot or ocellus. Length 5 inches. West Indies, south to Rio Janeiro. (Cope.) A specimen in our collection from Jamaica. It is close to *E. pisonis*, but the cheeks are not fully scaled. (*perniger*, very black.)

Culius perniger, COPE, Trans. Am. Philos. Soc. 1870, 473, St. Martins. (Coll. Dr. R. E. van Rijgersma.)
Eleotris perniger, EIGENMANN & EIGENMANN, *l. c.*, 55.

2524. ELEOTRIS PICTUS, Kner & Steindachner.

(GUAVINA.)

Head 3 to $3\frac{1}{3}$; depth 6. D. VI–I, 7 or 8; A. I, 7 or 8; lateral line 60; 24 scales in an oblique series from front of soft dorsal downward and back-

* *Eleotris belizianus* is described as follows: Head 4 in total; depth 5. D. VI–I, 9; A. I, 8; scales 60; eye 5 in head. Preopercle with a spine turned downward; 16 rows of scales between soft dorsal and anal; scales of top of head a little smaller than those of body, extending forward nearly to front of eyes; cheeks scaly; scales ciliate. Interocular space flattened, $\frac{1}{2}$ broader than eye; snout depressed a little longer than eye; lower jaw prominent; outer teeth enlarged; maxillary reaching front of eye. Dorsals contiguous. Color brownish, faint dark streaks on the fins. Belize; Cayenne. (Sauvage.) Length 100 mm. Evidently not different from *E. pisonis*.

ward to anal; about 20 in a vertical series. Body elongate, depressed anteriorly; head especially very broad and flat; mouth large, broad, very oblique, the maxillary reaching nearly or quite to opposite posterior margin of eye, its length 2⅛ to 2⅜ in head; lower jaw considerably projecting. Teeth in jaws all equal, in broad bands, the outer not at all enlarged. Eye small, anterior, its length in adult 2 in interorbital width, which width is about 3 in head; a conspicuous knob at upper anterior and posterior angles of orbit; preopercular spine well developed, strong, compressed, directed downward and forward. Scales on head very small, mostly cycloid, covering cheeks and opercles and upper part of head to the eyes; scales on body smaller and smoother than in most other species, those on belly much smaller than those on sides; scales on back and belly cycloid, only those on sides distinctly ctenoid. Pectoral fins moderate, reaching to near end of base of first dorsal, 1⅜ in head; ventrals inserted just behind axil, reaching halfway to vent, about 2 in head. Interspace between dorsals about equal to diameter of eye. Soft dorsal and anal short and high, very similar, coterminous; last ray of anal a little longer than ½ length of head; caudal peduncle long, a little shorter than head. Caudal fin rounded, 1⅛ in head. Color* dark, dull olivaceous brown, paler below; younger individuals mottled below with bluish and speckled with dark brown; sides without longitudinal stripes; fins dusky, all of them finely mottled and speckled with darker, the dark markings on dorsal and anal forming undulated longitudinal stripes; on pectorals and ventrals forming dark bars. Distinguished from related species by the larger mouth with small, equal teeth, and the small, smoothish scales. Length about 18 inches. Streams of the Pacific Coast, from Sonora, south to Panama; abundant in Rio Presidio, at Mazatlan, where the types of *E. æquidens* were taken; not rare about Panama. (*pictus*, painted.)

Eleotris pictus, KNER & STEINDACHNER, Abh. Ak. Wiss. Wien 1864, 18, pl. 3, f. 1, Rio Bayano, near Panama; depth 6 to 7 in total length; scales 60.

Culius æquidens, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 461, Rio Presidio, near Mazatlan. (Types, Nos. 28268 and 29240. Coll. Gilbert.)

Eleotris æquidens JORDAN & EIGENMANN, *l. c.*, 483.

805. ALEXURUS, Jordan.

Alexurus, JORDAN, Proc. Cal. Ac. Sci. 1895, 512 (*armiger*).

Body elongate, covered with small cycloid scales; preopercle with a small, concealed, hooked spine at its angle, as in *Eleotris;* caudal fin broad, its base with many procurrent rays. In other respects similar to *Eleotris.* One species known; marine. (ἀλέξω, to defend; οὐρά, tail, from the caudal fulcra.)

* A young example shows the following details of coloration in life: Blackish everywhere, sides with faint whitish streaks, along rows of scales a broad, blackish lateral band occupying whole of side, back and belly paler, traces of faint dark cross bands; caudal black, with a pale margin and some dark cross shades; pectorals, dorsals, and ventrals more or less barred with black; preopercular spine well developed; a whitish bar at base of caudal with a darker one before it.

2525. ALEXURUS ARMIGER, Jordan & Richardson.

Head $4\frac{2}{3}$; depth 8. D. VI–13; A. 11; V. I, 5; scales about 102–30; eye 8 in head; maxillary $2\frac{3}{8}$; mandible $2\frac{1}{2}$; snout $5\frac{3}{8}$; interorbital $4\frac{1}{8}$; pectoral $1\frac{4}{5}$; caudal equals head; ventral 2; last dorsal ray $1\frac{3}{5}$. Body long and low, compressed posteriorly, depressed in front. Head flattish and broad above, the cheeks moderately tumid. Eyes small, high up, separated by a broad, flattish, interorbital space; snout short; mouth moderate, very oblique, the maxillary ceasing below the center of pupil; lower jaw very heavy, oblique, projecting beyond upper, its outline horseshoe-shaped, obtuse in front. Teeth in rather broad bands, the outer enlarged below, but scarcely so above; none of them canine-like. Top of head with very small scales; cheeks and opercles with rudimentary scales above; preopercle with a concealed antrorse hook below, as in *Eleotris;* scales on body very small, perfectly smooth, partially embedded; scales on nape and throat minute. Gill membranes extending a little forward below, so that the branchiostegals are free from the isthmus. Insertion of dorsal twice as far from middle of base of caudal as from tip of snout; the fin low, its slender rays slightly filamentous; soft dorsal low, its last ray highest; anal similar, beginning under second dorsal ray; caudal long, bluntly pointed behind, with strongly procurrent base above and below, the base above $\frac{2}{5}$ length of head, formed of 14 short rays, that below a little shorter, of 12 rays, this procurrent portion forming an angle with the caudal proper where it joins it; pectorals and ventrals short, the ventrals inserted under pectorals. Color olive green, dusky above, paler below, but everywhere covered with fine black dots; both dorsals with the membranes pale, the rays each barred with black; caudal mesially blackish, all the rays barred or checkered in fine pattern; pectoral and anal pale, similarly speckled, base of pectoral dusky; ventral finely speckled. La Paz, Lower California; 1 specimen, $6\frac{1}{2}$ inches long, taken by Mr. James A. Richardson. (*armiger,* bearing arms, from the concealed spine.)

Alexurus armiger, Jordan & Richardson, Proc. Cal. Ac. Sci. 1895, 511, pl. 48, La Paz. (Type in L. S. Jr. Univ. Mus. Coll. James A. Richardson.)

806. EROTELIS, Poey.

(Esmeraldas de Mar.)

Erotelis, Poey, Memorias, II, 273, 1861 (*valenciennesi = smaragdus*).

Body very slender, elongate, covered with minute cycloid scales. Ventrals separate, the rays I, 5. No teeth on vomer. Lower pharyngeals subtriangular, the teeth stiff, villiform, none of them lamelliform. Posttemporal bones short, strongly divergent, the distance between their insertions about equal to the narrow interorbital space; top of head with a strong median keel, highest on the occipital region; no supraoccipital crest; no preopercular spine. Mouth very oblique. One species known; strictly marine. (Name an anagram of *Eleotris.*)

2526. EROTELIS SMARAGDUS (Cuvier & Valenciennes).

(Esmeralda Negra; Esmeralda de Mar.)

Head $4\frac{1}{3}$ to $5\frac{1}{2}$; depth 8 to 12. D. VI–I, 10; A. I, 9; V. I, 5; scales 100. Body very long and slender, compressed behind, the form much as in *Gobius oceanicus*. Head depressed, flattish above, the eyes mostly superior, not $\frac{1}{2}$ the width of the interorbital area, which has a knob near its middle. Mouth very oblique, the lower jaw much projecting, the maxillary about reaching front of eyes; teeth rather small, in bands. Fins rather high; dorsal spines slender, lower than the highest soft rays, which are $1\frac{1}{2}$ in head; caudal lanceolate, $\frac{1}{5}$ longer than head; ventrals moderate, 2 in head. Scales very small, cycloid. Color very dark green, almost black; the fins mostly bluish, the dorsal with brown lines; some dark markings about eye and on base of pectoral above. Length 8 inches. Coral shores among green algæ; known from Key West and Cuba; not common; not entering rivers. Here described from Key West specimens. (σμαραγδός, emerald.)

Eleotris smaragdus, Cuvier & Valenciennes, Hist. Nat. Poiss., xii, 231, 1837, Cuba; Jordan & Gilbert, Proc. U. S. Nat. Mus. 1884, 143.
Erotelis valenciennesi, Poey, Memorias, ii, 273, 1861, Cuba.
Erotelis smaragdus, Jordan & Eigenmann, *l. c.*, 484.

807. GYMNELEOTRIS, Bleeker.

Gymneleotris, Bleeker, Esquisse d'un Syst. Nat. des Gobioid., 304, 1874 (*seminuda*).

Body scaled only posteriorly, the anterior half and the head naked. Ventrals separate, I, 5. Vomer without teeth. Isthmus broad; skull without crests. Lower jaw with 4 large recurved teeth. Otherwise essentially as in *Eleotris*, the preopercle probably without spine. (γυμνός, naked; *Eleotris*.)

2527. GYMNELEOTRIS SEMINUDUS (Günther).

Head $3\frac{1}{3}$. D. VII–11; A. 9. Head depressed, broader than high, flat above. Snout rather obtuse, longer than eye, lower jaw somewhat prominent; cleft of mouth extending to below anterior margin of orbit. Teeth in upper jaw in a narrow band, the lower having 4 somewhat larger and recurved teeth in front, appearing to form a single series; palate toothless. None of the fin rays prolonged; pectoral not quite extending to origin of second dorsal; ventral much shorter than pectoral, its inner ray the longest, the others gradually decreasing in length outward; caudal fin rounded. Head and trunk naked; tail covered with small scales. Brown, with numerous well-defined white cross stripes on head as well as on body; vertical fins black. Panama. (Günther); known from the type only, a young example, $1\frac{3}{4}$ inches long; not seen by us. (*seminudus*, half-naked.)

Eleotris seminuda, Günther, Proc. Zool. Soc. London 1864, 24, pl. 4, figs. 2, 2a, Panama; Günther, Fish. Centr. Amer., 441, 1869.
Gymneleotris seminuda, Jordan & Eigenmann, *l. c.*, 484.

808. CHRIOLEPIS, Gilbert.

Chriolepis, GILBERT, Proc. U. S. Nat. Mus. 1891, 557 (*minutillus*).

This genus differs from *Gymneleotris*, Bleeker, in the total absence of scales, and the absence of enlarged canines in the front of the mandible. Head and body compressed, the former as deep as wide. Ventrals separate, near together, the inner rays longest, each with 1 spine and 5 soft rays. Teeth in a rather wide band in upper jaw, the outer series somewhat enlarged. Teeth in mandible in a single series, similar to outer row in upper jaw, none of them canine-like. Gill slits narrow; no dermal flaps on inner edge of shoulder girdle. Size small. (*χρεία*, want; *λεπίς*, scale.)

2528. CHRIOLEPIS MINUTILLUS, Gilbert.

Head 3⅓; depth 4⅓ in length. D. VII–12; A. 11. Mouth oblique, the maxillary reaching to below middle of orbit, 2½ in head; eyes high up, but with lateral range, separated by a narrow interorbital space less than diameter of pupil; diameter of orbit nearly twice length of snout, 3⅓ in head; dorsal spines high and slender, but not filamentous, the longest ½ length of head; soft dorsal rays higher, nearly ⅔ length of head; the anal lower; caudal short, broadly rounded, the depth of peduncle ½ length of head; length of pectoral equaling that of head without snout. Color uniform light brown on head and body, above and below; fins dusky, the anal blackish. A single specimen, 1 inch long, from *Albatross* Station 2825, off the east coast of Lower California. (*minutillus*, very small.)

Chriolepis minutillus, GILBERT, Proc. U. S. Nat. Mus. 1891, 558, Albatross Station 2825, Gulf of California, in 79 fathoms.

809. SICYDIUM, Cuvier & Valenciennes.

Sicydium, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 168, 1837 (*plumieri*).

Body subcylindrical, covered with rather small ctenoid scales; head oblong and broad, with cleft of mouth nearly horizontal; upper jaw prominent; snout obtusely rounded; lips very thick, the lower with a series of numerous slender horizontal teeth, of which sometimes only the extremities are visible; upper jaw with a single uniform series of numerous movable small teeth attached by ligament to edge of maxillary; behind this outer visible series lie numerous other parallel series of young teeth hidden in the gum, which succeed the former as they become worn out or broken; lower jaw with a series of widely set conical teeth; teeth all simple, slender, the distal half bent inward nearly at a right angle; eyes of moderate size; 2 dorsal fins, the anterior with 6 (5 or 7) flexible spines; caudal quite free; ventrals united into a short cup-shaped disk; gill openings of moderate width; 4 branchiostegals. Species few in the streams of the West Indies. (*σικύδιον*, diminutive of *σικύα*, a gourd, or gourd-shaped cupping glass, from the ventral disk.)

a. Body covered with small scales.
  b. Scales very small, about, 84. PLUMIERI. 2529.
  bb. Scales moderate, about 68. ANTILLARUM, 2530.
aa. Body nearly naked. VINCENTE, 2531.

2529. SICYDIUM PLUMIERI (Bloch).

(SIRAJO.)

Head 4 to $4\frac{2}{3}$; depth $4\frac{1}{2}$; eye 6 to 7 in head, 2 to 3 in interorbital width. D. VI–I, 10; A. I, 10; scales 84. Teeth in upper jaw long, slender, bent inward at right angles, only the lips protruding from the gums. Front teeth of lower jaw not larger than those behind; a single row of inconspicuous papillæ on the gum beneath the upper lip, a large median papilla above the maxillary suture; a median cleft in the upper lip. Pectorals longer than head; third, fourth, and fifth dorsal spines produced into long ribbons, the fourth, which is the longest, being 2 to 3 times height of body. Body usually covered with small scales, reduced in size on neck and belly; frequently almost naked, the scales present only on posterior part of body. Caudal deeply emarginate. Color olive or violet brown, with about 7 more or less distinct dark vertical bars; a dark bar at base of pectoral; dorsal with irregular dark markings; anal fin with a dark marginal band, sometimes edged with white; an H-shaped figure on base of caudal fin, and a black bar on its posterior half. Fresh waters of the West Indies. (Named for Père Charles Plumier, who discovered the species at Martinique.)

Gobins plumieri, BLOCH, Ichth., 125, pl. 178, fig. 3, 1786, Martinique; on a drawing by PLUMIER.

Sicydium siragus, POEY, Memorias, II, 278, 1861, Santiago de Cuba.

Sicydium plumierii, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 168, 1837; GILL, Proc. Ac. Nat. Sci. Phila. 1860, 101; GÜNTHER, Cat., III, 92, 1861; OGILVIE-GRANT, Proc. Zool. Soc. Lond. 1884, 156, pl. 11, fig. 1; JORDAN & EIGENMANN, *l. c.*, 484; EIGENMANN & EIGENMANN, *l. c.*, 56.

2530. SICYDIUM ANTILLARUM, Ogilvie-Grant.

Head $4\frac{2}{5}$; depth 6; width of head $\frac{2}{3}$ length. D. VI–I, 10; A. I, 10; scales 68. Teeth in upper jaw long, slender, and bent inward over the gum at right angles. A row of small lamelliform transverse papillæ on the gum beneath upper lip, with a larger median lamelliform papilla above maxillary suture; a slight median cleft in upper lip; maxillæ at right angles to one another; horizontal teeth conspicuous. Scales on body and tail subequal and larger than those on neck and belly. Maxilla not extending to vertical from posterior margin of eye, the diameter of which is contained $6\frac{1}{2}$ times in length of head and twice in interorbital space. Length of pectoral greater than that of head. The third, fourth, and fifth dorsal spines produced into long narrow ribands; the fourth, which is longest, nearly 3 times height of body; second dorsal considerably higher than body. Color uniform violet brown; dorsal fins with irregular wavy dark markings; anal with a black and white marginal band; caudal with a dark band on upper margin. One specimen, $4\frac{3}{4}$ inches long, from Barbados (Ogilvie-Grant); not seen by us. (*antillarum* of the Antilles.)

Sicydium antillarum, OGILVIE-GRANT, Proc. Zool. Soc. Lond. 1884, 157, Barbados.

2531. SICYDIUM VINCENTE, Jordan & Evermann, new species.

Another species of *Sicydium* or of some related genus is thus mentioned by Dr. Eigenmann: "Mr. Samuel Garman collected several hundred specimens of this species at Kingston, St. Vincent. Most of these specimens are less than an inch in length, the longest 1¼ inches; they differ considerably in coloration from the adult; most are entirely naked, a few of those examined having scales only on the posterior part of the body. Caudal deeply emarginate. There are traces of about 7 dark vertical bars; a black bar at base of pectoral; dorsals with several series of black spots; an H-shaped figure on base of caudal, a black bar on the posterior half of caudal fin; belly and lower part of body plain; everywhere else with black points. The specimens collected by Mr. Garman may be the types of a new species. No large specimens were collected at the Island of St. Vincent. Specimens 1½ inches in length from Hayti have the fins plain and a series of blotches along the middle of the posterior part of the body; the body, except the belly, is entirely covered with scales which are plainly ctenoid." (Eigenmann.) (Named for St. Vincent.)

Sicydium vincente, JORDAN & EVERMANN, Check-List Fishes, 456, 1896, St. Vincent Island; name only.

810. COTYLOPUS, Guichenot.

Cotylopus, GUICHENOT, in Maillard, Notes sur l' Isle de la Réunion, II, Addendum 9, 1864 (*acutipinnis*).

Sicya, JORDAN & EVERMANN, Check-List Fishes, 456, 1896 (*gymnogaster*).

This genus is closely allied to *Sicydium*, agreeing closely with the latter in external characters and in the absence of larger teeth in front of lower jaw; it differs chiefly in the form of the upper teeth which are curved, tricuspid, and trident-shaped, the middle cusp either permanent (*Cotylopus*) or else worn away leaving the tooth apparently bicuspid (*Sicya*). (κοτύλη cup; πούς, foot.)

SICYA (σικύα, a gourd, or gourd-shaped cup):

a. Teeth in upper jaw curved, tricuspid, trident-shaped, the lateral lobes long, the middle short and suspended between the outer lobes, and soon wearing away leaving the tooth apparently bicuspid.

b. Neck and belly naked; a double or triple row of small papillæ on the gum beneath the upper lip. GYMNOGASTER, 2532.

bb. Neck and belly covered with small scales; gum beneath the upper lip smooth. SALVINI, 2533.

Subgenus SICYA, Jordan & Evermann.

2532. COTYLOPUS GYMNOGASTER (Ogilvie-Grant).

Head 4½ to 5; depth 5½ to 6. D. VI–I, 10; A. I, 10; scales 60 to 64; eye 6 in head, twice in interorbital space. Teeth in the upper jaw tricuspid, the middle cusp, which is situated at the anterior end of tooth, is very soft and soon becomes worn away. A double or treble row of small papillæ on the gum beneath the upper lip, without a larger median papilla; upper lip with a very slight median notch; maxillæ containing an angle of about 75°; horizontal teeth more or less inconspicuous. Scales strongly ctenoid; neck and belly naked. Length of pectoral greater than

that of head. Second, third, and fourth dorsal spines produced into filaments; the third, which is the longest, twice height of body; second dorsal higher than body. Color violet brown, yellowish in young specimens, shaded with indistinct transverse bands of darker; irregular brown spots on axis of pectoral, and a broad dark band from base of pectoral to root of caudal, both more or less indistinct in adult specimens; fins violet, clouded with darker. Length 4⅝ inches. Streams about Mazatlan (Ogilvie-Grant); not seen by us. (*γυμνός*, naked; *γαστήρ*, belly.)

Sicydium gymnogaster, Ogilvie-Grant, Proc. Zool. Soc. Lond. 1884, 158, pl. 11, fig. 2, and pl. 12, fig. 6, Mazatlan.
Sicyopterus gymnogaster, Jordan & Eigenmann, *l. c.*, 485.

2533. COTYLOPUS SALVINI (Ogilvie-Grant).

Head 4⅝; depth 6¼. D. VI-9 or 10; A. I, 10; scales 78; eye 5½ in head, twice in interorbital space. Teeth in upper jaw tricuspid; the middle cusp, which is situated at anterior end of tooth, very soft and soon becomes worn away. Gum beneath upper lip smooth; a median papillose tubercle above maxillary suture; upper lip with a small median notch; maxillæ containing an angle of about 75°; horizontal teeth conspicuous. Scales ctenoid, those on neck and belly smaller than those on body and tail. Length of pectoral rather greater than that of head. Second and third dorsal spines subequal and produced into short filaments, 1½ times height of body; second dorsal not so high as body. Color olive brown; anal yellow, with a black and white band along margin; membrane of second dorsal clear, spotted with brown; caudal with a dark and yellow band round the extremity. Length 4⅜ inches. Streams near Panama; 1 specimen known. (Ogilvie-Grant.) (Named for Osbert Salvin, who collected largely in Central America for the British Museum.)

Sicydium salvini, Ogilvie-Grant, Proc. Zool. Soc. Lond. 1884, 159, pl. 12, fig. 2, Panama.
Sicyopterus salvini, Jordan & Eigenmann, *l. c.*, 485.

811. EVORTHODUS, Gill.

Evorthodus, Gill, Proc. Ac. Nat. Sci. Phila. 1859, 195 (*breviceps*).

Body elongate, covered with ctenoid scales of moderate size. Head thick, short, naked. Isthmus moderate. Teeth in a single series, with the crown emarginate, those of the lower jaw horizontal; no canines. First dorsal of 6 spines; ventral fins united, not adherent to the belly, otherwise as in *Gobius*, so far as known. (*εὖ*, well; *ὀρθός*, straight; *ὀδους*, tooth.)

2534. EVORTHODUS BREVICEPS, Gill.

Head 4½, about as deep as wide; depth 4½. D. VI-I, 10; A. I, 11; eye 3. Teeth emarginate, uniserial, those of lower jaw nearly horizontal. Snout blunt, profile evenly decurved; caudal rounded, 3 in length of body; some of the dorsal rays filamentous. Color light brown, with irregular blackish blotches along sides; 2 black spots at base of caudal fin, 1 above the other, alternating with 1 more anterior on the peduncle; first dorsal

with 2 bands parallel with its upper margin; second dorsal with 3 narrow longitudinal bands. (Gill.) Fresh waters of Trinidad and Surinam; not seen by us. (*brevis*, short; *-ceps*, head.)

Evorthodus breviceps, GILL, Proc. Ac. Nat. Sci. Phila. 1859, 195, Trinidad; JORDAN & EIGENMANN, *l. c.*, 486.

812. LOPHOGOBIUS, Gill.

(CRESTED GOBIES.)

Lophogobius, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 240 (*cristagalli=cyprinoides*).

Dorsal spines 6; scales evidently ctenoid. Body short, compressed, form much as in *Dormitator*; nape with fleshy crest; scales large. Vertebræ 11 + 15. Interorbital area of cranium anteriorly elevated, with a large foramen-like depression in front of eye. One species, differing considerably in form from the other Gobies. The study of its skeleton shows no distinction of much importance unless the peculiar form of its interorbital area be regarded as such. (λόφος, crest; *Gobius*.)

2535. LOPHOGOBIUS CYPRINOIDES (Pallas).

Head 3⅔; depth 3⅔; greatest width 5½ to 6½. D. VI or VII–10 or 11; A. 9 or 10; scales 26 to 30; vertebræ 11 + 15; eye 3½ to 4. Body short and deep, little compressed, formed much as in *Cyprinodon;* head naked, a prominent naked dermal crest extending from above middle of eye to near front of spinous dorsal; interorbital width slightly less than diameter of eye; profile convex; snout short, bluntish, about as long as eye; mouth very oblique, the gape slightly curved; front of upper lip on level of lower border of eye; lower jaw somewhat projecting; teeth in both jaws in bands, the outer series erect and somewhat enlarged, those of the inner series small; scales large, reduced on breast and nape; a few scales on upper part of opercle; median line before dorsal naked; dorsal spines produced in short filaments; last rays of soft dorsal reaching caudal; caudal rounded; pectorals lanceolate, reaching beyond insertion of anal, the upper rays not silk-like; skull very broad and short, with low, median crest, highest behind; double crests of temporal region joining at the upper posterior angles of the eyes and forming a bridge over the interorbital area, the crests ending abruptly above the anterior part of the orbit, forming a decided angle, the bridged interorbital leaving a large foramen in front of this angle. Color blackish green in life; spinous dorsal black; soft dorsal, ventrals, and anal dark, plain; pectorals lightish, plain; caudal finely mottled. Length 2 inches. West Indies, north to southern Florida; generally common in the streams and brackish waters of Cuba and other islands. Recently taken by Dr. Evermann in brackish water at Biscayne Bay, Florida. (κυπρῖνος, carp; εἶδος, resemblance.)

Gobius cyprinoides, PALLAS, Spicilegia, Zool., VIII, 17, pl. 1. fig. 5, 1770, Amboina; CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 129, 1837; GÜNTHER, Cat. Fish., III, 8, 1861.

Gobius cristagalli, VALENCIENNES, in CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 130, 1837, Havana; GUICHENOT, in Ramon de la Sagra, Hist. Cuba, 128, pl. 3, fig. 3, 1850.

Lophogobius cyprinoides, POEY, Repertorio, I, 335, 1867; POEY, Synopsis, 393, 1868; POEY, Enumeratio, 125, 1876; JORDAN & EIGENMANN, Proc. U. S. Nat. Mus. 1886, 487; EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897, 131, plate 9, fig. 13.

813. GOBIUS (Artedi) Linnæus.

(Gobies.)

Gobius, Artedi, Genera, 28, 1738 (*Gobius ex nigricante varius*, etc., = *niger*).
Gobius, Linnæus, Syst. Nat., Ed. x, 262, 1758 (*niger*, etc.), and of authors generally.
Gobionellus, Girard, Proc. Ac. Nat. Sci. Phila. 1858, 168 (*hastatus* = *oceanicus*).
Ctenogobius, Gill, Fish. Trinidad, 374, 1858 (*fasciatus*).
Euctenogobius, Gill, Annals Lyc. Nat. Hist. New York 1859, 45 (*badius*).
Smaragdus, Poey, Memorias, II, 279, 1861 (*smaragdus*).
?Pomatoschistus, Gill, Proc. Ac. Nat. Sci. Phila. 1863, 263, footnote (*minutus*).
Coryphopterus, Gill, Proc. Ac. Nat. Sci. Phila. 1863, 263 (*glaucofrænum*).
?Deltentosteus, Gill, Proc. Ac. Nat. Sci. Phila. 1863, 263, footnote (*quadrimaculatus*).
?Gobiichthys, Klunzinger, Fisch. Rothen Meeres, 479, 1871 (*petersii*).
?Mesogobius, Bleeker, Esquisse d'un Syst. Nat. Gobioid., 317, 1874 (*guavina*).
?Stenogobius, Bleeker, *l. c.*, 317 (*gymnopomus*).
?Oligolepis, Bleeker, *l. c.*, 318 (*melanostigma*).
?Gnatholepis, Bleeker, *l. c.*, 318 (*anjerensis*).
?Callogobius, Bleeker, *l. c.*, 318 (*hasselti*).
?Hypogymnogobius, Bleeker, *l. c.*, 318 (*xanthozona*).
?Hemigobius, Bleeker, *l. c.*, 318 (*melanurus*).
?Cephalogobius, Bleeker, *l. c.*, 320 (*sublitus*).
?Acentrogobius, Bleeker, *l. c.*, 321 (*chlorostigma*).
?Porogobius, Bleeker, *l. c.*, 321 (*schlegeli*).
?Amblygobius, Bleeker, *l. c.*, 322 (*sphinx*).
Zonogobius, Bleeker, *l. c.*, 323 (*semifasciatus*).
?Odontogobius, Bleeker, *l. c.*, 323 (*bynoënsis*).
?Stigmatogobius, Bleeker, *l. c.*, 323 (*pleurostigma*).
?Oxyurichthys, Bleeker, *l. c.*, 324 (*belosso*).
Lythrypnus, Jordan & Evermann, Check-List Fishes, 458, 1896 (*dallii*).

Body oblong or elongate, compressed behind. Head oblong, more or less depressed. Eyes high, anterior, close together; opercles unarmed. Mouth moderate. Teeth on jaws only, conical, in several series, those in the outer row enlarged; no canines. Isthmus broad. Shoulder girdle without fleshy flaps or papillæ. Skull depressed, abruptly widened behind the eyes and without distinct median keel. Scales moderate, ctenoid, permanently covering the body; cheeks usually naked; belly generally scaly. Dorsal with 6 rather weak spines; pectorals well developed, the upper rays sometimes very slender and silky; ventrals completely united, not adnate to the belly; caudal fin usually obtuse. Species very numerous. The genus *Gobius*, as here understood, comprises a very large number of species more or less closely related to the European type of the genus, *Gobius niger*, and its American relative, *Gobius soporator*. An examination of skulls or skeletons of numerous European and American species shows a remarkable uniformity in most respects. The general form and structure of the cranium is the same in all, the only differences being very minor ones in the height of certain crests. *Gobius oceanicus* seems the most aberrant, but seems to be inseparable generically on account of intermediate forms. Probably several of the many genera indicated by Bleeker will prove valid, but only a thorough study of skeletons can establish them. It is not unlikely that *Ctenogobius*, to which group most of our species belong, may be separable from *Gobius*. (κωβιός; Latin, *Gobius* or *Gobio*, a name applied to the gudgeon (*Gobio gobio*) and

to other small fishes; allied to *Cobitis*, chub, etc. According to Apostolides κωβιός and γωβιός are common names in modern Greek for all species of the genus *Gobius*. Aristotle κωβιός, 610*b*, 4, 598*a* 11 16, 508*b* 16, 569*b* 23, 621*b* 13 19, 567*b* 11, 591*b* 13, 601*b* 22, 835*b* 14. The κωβιός has many pyloric appendages above the stomach, spawns near the land on the rocks, the bunches of eggs are flat and crumbling; it feeds on mud, seaweed, sea moss, etc.; lives near the land, gets fat in the rivers, and is found in schools. The white κωβιός, found in the Euripus of Lesbos, never leaves that lagoon for the open sea as the other fishes found there do. Latin *Gobio* and *Cobio*, Plin. *Gobius*, Ovid., Hal. 12, 8. Martial 13, 88. Horace A. Hoffman.)

GOBIUS:

a. Upper rays of pectoral fin silk-like; i. e., short and very slender and flexible, free for nearly their whole length.

b. Body robust, compressed posteriorly; depth 4⅔ to 5¼ in length; head broad, low, rounded in profile, its length 3 1/10 to 3⅔ in body; eye 4 to 5 in head; mouth large, little oblique; lips thick; teeth in both jaws in bands, the outer series enlarged; those on lower jaw subequal; scales large, strongly ctenoid, smaller on nape and belly; dorsal spines short, none filamentous. Color olivaceous, light or dark, varying from sand color to greenish black, everywhere mottled and marbled with dark and paler; a faint dusky spot behind eye. D. VI–9 or 10; A. I, 7 to 9. Scales 36 to 41. SOPORATOR, 2536.

CTENOGOBIUS (κτεις, comb; *Gobius*):

aa. Upper rays of pectoral normal, not silk-like, similar to the others.

c. Scales large, 25 to 33.

d. Color in life olivaceous, more or less spotted, never red.

e. Dorsal soft rays 12 to 14; vertex and nape with a slight median fold of skin.

f. Body compressed, its depth 5 in length; head 3⅓ to 3⅔; eye 3 to 3¼ in head; vertex and nape with a slight median fold of skin; maxillary reaching about to front of pupil; lower jaw very slightly produced; teeth in bands, the outer slightly enlarged. Olivaceous; spinous dorsal black at tip; second dorsal finely checkered in adult. D. VI–14; A. I, 11. Scales 25 or 26–10. NICHOLSII, 2537.

ff. Body long, not much compressed; head 3½; eye 3 in head; no median fold on vertex and nape; a dark spot on first dorsal. EIGENMANNI, 2538.

ee. Dorsal soft rays 10 to 12; no median fold of skin on vertex and nape.

g. Caudal with 2 spots at its base; jaws unequal, the lower slightly produced; body robust, compressed behind, the depth 5 in total length; head 4⅓; eye longer than snout, 3⅓ in head; maxillary reaching pupil; teeth in a band, the outer enlarged and distant, the inner enlarged and bent backward. Brownish; a faint blue spot on each scale; six spots along middle of back; similar spots on scapular region and middle of sides; 2 spots on base of caudal; a dark spot above opercle; blue dots on head; a straight blue line crossing cheek above and continued on opercle; dorsals faintly spotted. D. VI–10; A. 10. Scales 25–7. (Gill.) GLAUCOFRÆNUM, 2539.

gg. Caudal plain or with but a single spot at its base.

h. Dorsal spines low, the highest little longer than head.

i. Region from nape to dorsal entirely scaled.

j. Pores on preopercle not very conspicuous; no canine teeth.

k. Body very slender, compressed, the depth 5⅔ in length; caudal much longer than head; mouth rather large, the lower jaw projecting; teeth unequal, rather strong; yellowish, much spotted with darker. D. VI-12; A. 12. Scales 35.

MANGLICOLA, 2540.

kk. Body subfusiform, little compressed; depth 4⅓ in length; head blunt, 4 in length, rounded in profile; eye equal to snout, 4 in head. Mouth small, horizontal, the lower jaw included; maxillary 3 in head, reaching to below eye. Teeth small, in bands in both jaws, the outer enlarged, those of the upper jaw very slender. Scales large, ctenoid, those of nape and belly little reduced. Longest dorsal spine shorter than head. Caudal scarcely pointed, about as long as head. Color whitish gray, middle of sides with 4 or 5 dark blotches, from each of which a narrow dark bar extends downward and forward; a large black blotch above pectorals, obsolete in female; a small black spot at base of caudal; a dark mark below eye; vertical fins barred. D. VI-12; A. 11 or 12. Scales 33.

STIGMATURUS, 2541.

jj. Pores on preopercle very conspicuous; lower jaw with small canies. D. VI-I, 9; A. I, 9.

QUADRIPORUS, 2542.

ii. Region between nape and dorsal with a narrow naked median strip. Body moderately elongate, subfusiform, the depth 5⅓ in length. Head large, not so blunt as in *G. boleosoma*, its length 3⅞ to 3⅔ in length; anterior profile gently decurved; snout 3⅓ to 3⅓ in head; eye 4; mouth large, slightly oblique; maxillary entending to front of pupil, 2⅓ in head. Teeth small, slender and curved, in moderate bands. Scales moderate, ctenoid, those in front much reduced in size; breast naked. Longest dorsal spine 1⅓ in head. Caudal as long as head, somewhat pointed. Olivaceous, mottled with gray; about 5 rounded dark blotches along middle of sides, the last forming a spot at base of caudal; no dark spot on side of nape; some dark marks on head; vertical fins barred. D. VI-12; A. 13. Scales 33 to 35. SHUFELDTI, 2543.

iii. Region between nape and dorsal entirely naked.

l. Highest rays of second dorsal little more than ⅓ head, none of them reaching base of caudal.

m. Profile much decurved, skull rounded behind, without distinct median ridge; mouth horizontal. Body elongate, deepest below front of dorsal, tapering regularly backward, the greatest depth 5⅓ in length. Head short, blunt, pro-

file anteriorly abruptly decurved, cheek somewhat swollen. Length of head 3½ in body. Snout about equal eye, 3⅔ in head. Mouth horizontal, maxillary reaching to below pupil (in male); lower jaw included. Teeth in each jaw in a band, the outer row of the upper jaw large, recurved. Scales large, ctenoid, somewhat reduced anteriorly. Nape, breast, and belly naked. Dorsal spines about ⅔ of head. Caudal pointed, 2⅔ to 3½ in body. Color olivaceous, with numerous dark reticulations on the back; 5 black spots along the sides, the last forming a spot on base of caudal, sometimes with V-shaped dark bars extending from them to dorsal; breast and sides of belly with numerous dark specks in male; a dark line between eyes; a dark line from eye to middle of premaxillary, some dark spots below eye, sometimes forming bars, sometimes a stripe; a large oblique spot above pectorals, continued on opercle; a black spot at base of pectoral; dorsals and caudal barred, anal uniform dusky, ventrals and pectorals black in male, white in female. D. VI–11; A. 10 to 12. Scales 25 to 30.

BOLEOSOMA, 2544.

mm. Profile moderately decurved; eye longer than snout, 3¾ in head. Color yellowish, oblong dark blotches on middle of sides; dorsal and caudal barred. Head 4; depth 6. D. VI–12; A. 10.

FASCIATUS, 2545.

ll. Highest rays of second dorsal as long as head, the last reaching base of caudal. Body elongate, the back not arched; depth 6 in length; head 4, not compressed, the cheeks tumid. Profile abruptly decurved, the snout 3½ in head. Mouth large, nearly horizontal, the maxillary reaching posterior edge of eye in males, middle of eye in females. Teeth in narrow bands in each jaw, the outer somewhat enlarged, the outer in some (males?) much enlarged above and recurved, the enlarged teeth fixed, the others movable. Scales large, ctenoid, reduced anteriorly; belly naked. Dorsal spines little filamentous, the longest about equal to head; caudal 2¼ to 3 in body. Males dark olive, with 4 oblong dark blotches along middle of sides; a dark caudal spot; a black blotch larger than eye on each side of shoulder; dorsal spotted;

caudal reddish above, dusky below; females with 5 oblong dark blotches on sides, the last on base of caudal; from each of the middle blotches a V-shaped bar runs to the back; a black shoulder blotch; a dark bar from eye to mouth; ventrals pale, with 2 dark streaks. D. VI–11; A. 12. Scales 30 (27 to 33) at least in males.

ENCÆOMUS, 2546.

kk. Dorsal spines high, the highest reaching past middle of second dorsal. Nape scaly. Body elongate, moderately compressed.

n. Depth 5 to 6 in length. Profile little decurved, skull flattish behind, much broader than in *Gobius boleosoma*, with an evident median ridge; mouth very oblique, much larger than in *G. boleosoma;* lower jaw thin and flat. Back slightly arched. Body a little deeper and rather less compressed than in *G. encæomus*, the depth 5 to 6 in length. Head 4. Anterior profile moderately decurved. Eye $3\frac{1}{3}$ in head. Mouth large, oblique; maxillary reaching to below pupil in both sexes. Teeth above uniserial, some of them enlarged and recurved; lower teeth in a narrow band; males sometimes with the hindermost of the outer series a strong, exserted, recurved canine; belly naked. Longest dorsal spine $\frac{2}{3}$ head in females, elevated in males; soft dorsal elevated in males; caudal $3\frac{1}{2}$ in body. Color light greenish; sides of male with 5 or 6 narrow, straight, rather sharply defined whitish or yellowish cross bars, regularly placed; 4 dark bars, 3 below eye and 1 on opercle; a small dark spot behind and above opercle; vertical fins barred; female with a row of irregular dark spots connected by a dusky streak, and with the pale cross bars obsolete. D. VI–12; A. 13. Scales 27.

STIGMATICUS, 2547.

nn. Depth $4\frac{2}{3}$ in length; the profile very obtuse anteriorly; mouth nearly horizontal, the maxillary extending beyond pupil, $2\frac{2}{3}$ in head. Teeth strong, uniserial, 4 shortish canines in lower jaw behind the other teeth; upper teeth largest. Some of the dorsal spines filamentous, reaching (in male) past middle of second dorsal; caudal $\frac{1}{4}$ longer than head; scales large, ctenoid, those on nape and belly much reduced in size. Dark olive, with 4 or 5 irregular, confluent, blackish cross bands, besides irregular, dark blotches; head marked with darker; fins mostly dusky; caudal dark blue with 2 red longitudinal stripes. D. VI–11; A. 10. Scales 27. LYRICUS, 2548.

nnn. Depth 4 in length; mouth nearly horizontal; teeth short and thick, uniserial; yellowish, much mottled and blotched. D. VI–11; A. 11. Scales 30. GARMANI, 2549.

dd. Color in life, cherry red, with many bluish cross bars; body stout; depth 4⅓ in length. D. VI–11; A. 9. Scales large. ZEBRA, 2550.

cc. Scales moderate or small, 40 to 90.

o. Soft dorsal and anal short, each of 10 to 14 rays; body more or less elongate. EUCTENOGOBIUS (εὖ, well; κτείς, comb; *Gobius*):

p. Caudal rounded, not much longer than head.

q. Scales 40; dorsal with 9 soft rays only; anal with 9; depth 6⅓ in total length; head broad, flattish; snout short, decurved; eye 4¼ in head, 1⅓ in interorbital area, longer than snout; maxillary extending to below middle of eye. Some of the dorsal spines produced in filaments, the third 1⅓ times depth of body; caudal short, rounded. Two rows of ill-defined blotches on upper half of body; 2 rows of brownish spots on second dorsal, the upper strongly marked POEYI, 2551.

qq. Scales 50; dorsal and anal with 10 soft rays each; profile very oblique. Color dark brown. BADIUS, 2552.

GOBIONELLUS* (diminutive of *Gobius*):

pp. Caudal lanceolate, much longer than head; lower jaw thin; usually a green spot on roof of mouth in life.

r. Body rather deep, the depth about 5 in length.

s. Teeth minute, seen with a lens only. Dorsal spines filamentous; scales much reduced below. D. VI–13; A. 14. Scales 62. MICRODON, 2553.

ss. Teeth well developed.

t. Scales rather large, 39 to 42; body moderately elongate, compressed; depth 5¼; head 4. Head not compressed, the cheeks tumid, the snout short, abruptly decurved; mouth large, little oblique, the jaws equal, the maxillary 2⅓ in head, reaching to below pupil; eye 5 in head; teeth above large, unequal, uniserial, some of them fixed, those below small, in a band. Scales anteriorly cycloid, becoming larger posteriorly, and ctenoid; dorsal spines scarcely filamentous, none of them as high as body; caudal 2¼ in body. Light olive, with dark olive blotches; body and head with many conspicuous round spots of cream color, each surrounded by a dusky ring, these most distinct on the head, all smaller than pupil; snout with dusky streaks; dorsals and caudal sharply barred; anal and ventrals dusky (in male); a small round spot at base of caudal. D. VI–11; A. 11. SMARAGDUS, 2554.

tt. Scales comparatively small (53). Body elongate, compressed behind; head a little compressed, 3¾ in length; depth 5; eye 3⅓ in head, shorter than the rounded snout; maxillary reaching to below middle of eye; teeth small, the outer a little enlarged; dorsal spines all shorter than head, not filamentous. Nape scaly, its scales much reduced in size; scales ctenoid. Two violet stripes from eye to mouth; 8 or 9 violet bars on sides; 3 or 4 bars on caudal; second dorsal spotted. D. VI–12; A. 11 or 12. STRIGATUS, 2555.

rr. Body elongate, the depth 6⅓ to 9 in length; head 4⅓; teeth well developed; caudal very long.

* *Gobionellus* is probably generically distinct from *Gobius* and *Ctenogobius*, but at present we do not know how to limit it, and therefore we are unable to define it.

u. Scales 55 to 60; eye longer than snout, 4 in head; mouth slightly oblique, the jaws equal, the maxillary not reaching center of eye; teeth in a narrow band, the outer much enlarged and separated from the others by a narrow interspace. Second dorsal spine not equal to depth of body. Caudal 3⅓ in body. Scales on nape and axil very small, those on posterior part of body much larger. Light olive green; a series of brown spots along middle of tail; sides of head with dusky blotches, vertical fins dotted with black. D. VI–13; A. 14. Scales 58–20. SAGITTULA, 2556.

uu. Scales very small (60 to 90); caudal more than twice as long as head in adult. Body compressed, extremely elongate, the depth 6 to 9 in length; head higher than wide, short, compressed, 4½ to 5 in length; mouth wide, oblique; maxillary in adult reaching to below posterior border of eye. Lower jaw very thin and flat; teeth in both jaws small, subequal, those in the upper jaw in a single series, those of the lower in a narrow band; outer teeth somewhat movable. Scales anteriorly small, cycloid, embedded, those behind larger and ctenoid; a few scales on upper anterior corner of opercle; dorsal fins high, some of the spines filamentous, longer than head. Caudal very long filamentous, 2 to 2⅔ in body. Light olive; fins dusky in male; a round, black spot on side, a little larger than eye, below spinous dorsal; first dorsal spine with 2 or 3 black spots; a small dusky spot at base of caudal; emerald spot on tongue conspicuous, fading in spirits. D. VI–14; A. 14 or 15.

v. Head 5½ to 6 in length; scales 60 to 70; patch of scales on opercle obsolete. HASTATUS, 2557.

vv. Head 7 to 8 in length; scales about 90; patch of scales on opercle well developed. OCEANICUS, 2558.

LYTHRYPNUS (λύθρον, gore; ὑπνός, slumberer; a red sleeper):

oo. Soft dorsal and anal very long; D. VI–17; A. 14. Body short, compressed; mouth very oblique; jaws with distant canine-like teeth. Coral red, with bluish crossbands and markings. Scales 40. DALLII, 2559.

Subgenus GOBIUS.

2536. GOBIUS SOPORATOR,* Cuvier & Valenciennes.

(SLEEPER; MAPO; CAIMAN.)

Head 3$\frac{1}{10}$ to 3$\frac{3}{8}$; depth 4$\frac{3}{5}$ to 5$\frac{1}{2}$; eye 4 to 5. D. VI–I, 9 or 10; A. I, 7 to 9; scales 35 to 41—13 to 15. Vert. 11 + 16. Body robust, compressed pos-

*The specimens examined are from Panama, Barbados, Pará, Itapuana, Cuba, Galapagos, Sambara, Bahia, Orange Key, Bahamas, Pernambuco, St. Thomas, Tortugas, Florida Keys, Martinique, Sao Matheas, Curuca, Rio de Janeiro, Rio Doce. "The color variations among examples of this species are very great, specimens from one locality varying from plain sand color, or gray, to greenish black; some dark brown specimens have light bars across the back; in others the scales have light centers forming horizontal series of light lines; sometimes there are light spots on sides of head and cheek; some specimens are conspicuously marbled with light and dark brown, and white spots occur in the centers of some of the scales on specimens of any ground color, these white spots being brighter on some of the scales than on the others, forming interrupted longitudinal lines. If any value could be placed upon the coloration, almost every specimen would be a distinct species. The color variation is irrespective of locality, some localities having all the above-described variations. The types of Poey's *mapo*, *lacertus*, and *brunneus* prove to be color varieties of *Gobius soporator*." (Eigenmann.)

teriorly; head broad, low, rounded in profile; mouth large, little oblique; lips thick; teeth on upper jaw in a broad band, those of outer series enlarged, the inner ones minute; teeth on lower jaw in a broad band, the outer row enlarged, but not quite as large as the outer series on upper jaw. Anterior half of trunk scaled, head naked; scales large, strongly ctenoid, smaller on nape and belly. Dorsal spines short, not filamentous; upper rays of pectoral fin silk-like, short, and very slender and flexible, free for nearly their whole length; caudal short. Skull posteriorly much as in *Lophogobius cyprinoides*, but the median crest reduced to a slight ridge. Lateral crests very high and closely approximated, rising obliquely outward; the inner crests meeting behind eye, the outer ones forming a very high border about the orbit. Interorbital very narrow and deep, with a median ridge. Coloration that of the rocks, usually granite gray or olivaceous, light or dark, varying from sand color to greenish black, everywhere mottled and marbled with darker and paler, often with brassy or greenish; a faint dusky spot behind eye; coloration varying indefinitely with the surroundings; pectorals, dorsals, and caudal generally mottled; anal and ventrals usually plain. Length 3 to 6 inches. Specimens from Pensacola show the following characters: Head $3\frac{1}{3}$ (4 in total); depth 4 (5). D. VI-10; A. I, 9; scales 30 to 38; 12 rows of scales from first dorsal downward and backward to anal. Scales on nape extremely small, those on sides firm, ctenoid; first dorsal with an oblique median shade of blackish, the base in front and the distal part light orange; second dorsal dusky at the base, with some spots, its margin light orange; caudal reddish, with dusky cross lines or spots; anal and ventral dusky, yellowish at base in the female; pectoral olivaceous, yellowish at base, reddish at tip, 2 dark spots on base of pectorals. Form robust. Head rather blunt and heavy, the snout less abruptly decurved than in *G. lyricus*. Mouth moderate, the jaws equal, the maxillary reaching about to front of pupil, $2\frac{2}{3}$ in head. Teeth in moderate bands, the outer series somewhat enlarged. Cheeks full, tumid. Eyes moderate, placed rather high, much broader than the interorbital space. Dorsal spines slender, the first longer than the other, but not filamentous, $1\frac{2}{3}$ in head; caudal rounded, $1\frac{1}{3}$ in head; upper rays of pectorals silk-like, the fin somewhat longer than the ventral, $1\frac{1}{4}$ in head. Color in life, very deep olive green, the back and sides obscurely barred and much marbled with different shades of olive green; cheeks with dark markings, forming reticulations around pale spots; whole under part of head blackish in the males, yellowish in the females. Tropical seas; universally distributed and almost everywhere common, lurking among stones or on sand in shallow water, or in rock pools, moving very quickly when disturbed; north on our coast to Carolina and Gulf of California. The commonest of all shore fishes in tropical America. Among our species it seems to be the one most nearly related to the European *Gobius niger*, and it may, therefore, be held to represent the subgenus *Gobius*, if our other species be placed in different subgenera. Perhaps all the others will ultimately be removed from *Gobius*.

Gobius soporator, Cuvier & Valenciennes, Hist. Nat. Poiss., xii, 56, 1837, Martinique; Günther, Cat. Fish., iii, 26, 549, 1861; Poey, Enumeratio, 124, 1876; Jordan & Gilbert, Synopsis, 634, 1883.

Gobius lineatus, JENYNS, Zool. Voy. Beagle, 95, pl. 19. fig. 2, 1842, Galapagos Archipelago. (Coll. Charles Darwin.)

Gobius catulus, GIRARD, Proc. Ac. Nat. Sci. Phila. 1858, 169, St. Joseph Island, Texas; GIRARD, U. S. and Mex. Bound. Survey, Zool., 26, pl. 12, figs. 9 and 10, 1859; JORDAN & EIGENMANN, *l.c.*, 493.

Gobius mapo, POEY, Memorias, II, 277, 1861, Cuba; POEY, Synopsis, 392, 1868.

Gobius lacertus, POEY, Memorias, II, 278, 1861, Cuba; POEY, Synopsis, 392, 1868; POEY, Enumeratio, 125, 1876.

Gobius andrei, SAUVAGE, Bull. Soc. Philom., Ser. 7, IV, 44, 1880, Rio Guayas, Ecuador. (Coll. André.)

Gobius carolinensis, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 268, Charleston, South Carolina; JORDAN & GILBERT, Synopsis, 634, 1883.

Gobius brunneus, POEY, Synopsis, 393, 1868, Havana; name preoccupied.

Evorthodus catulus, JORDAN & GILBERT, Synopsis, 632, 1883.

According to Dr. Eigenmann, *Gobius albopunctatus* of the Western Pacific can not be separated from *Gobius soporator*. In this case several other synonyms should be added.

Subgenus CTENOGOBIUS, Gill.

2537. GOBIUS NICHOLSII, Bean.

Head $3\frac{1}{8}$ to $3\frac{3}{5}$; depth 5 to $5\frac{1}{4}$. D. VI–I, 12 to 14; A. I, 11; scales 25 or 26–10. Body compressed; width of head about twice in its length. Mouth oblique, the maxillary reaching to front of pupil, $2\frac{1}{2}$ to 3 in head; lower jaw very slightly produced. Teeth present on both jaws; the outer series of long, conical teeth, placed at a considerable distance apart; the enlarged teeth on lower jaw not extending on the sides; the inner series of a band of small teeth. Interorbital space very narrow, equaling pupil. Snout 4 to $4\frac{1}{8}$ in head. Eyes large, placed high, 3 to $3\frac{1}{4}$ in head. Caudal peduncle $2\frac{3}{4}$ to $3\frac{1}{6}$ in head. Scales large, caducous, ctenoid; lacking on head, nape, and fins. Dorsal spines slender, flexible; base of first dorsal $1\frac{3}{8}$ to 2 in head; soft dorsal and anal similar; base of anal $1\frac{1}{5}$ to $1\frac{1}{3}$ in base of soft dorsal, and $1\frac{1}{4}$ to $1\frac{1}{3}$ in head; ventrals $1\frac{1}{4}$ to $1\frac{2}{3}$ in head, inserted below or slightly behind origin of pectorals; pectorals reaching a considerable distance beyond ventrals, $1\frac{1}{12}$ to $1\frac{1}{4}$ in head; caudal rounded, not equaling head. Color in spirits, light yellowish brown, with traces of darker, lighter below; ventrals usually dusky; spinous dorsal narrowly margined with black; second dorsal finely checkered in adult; other fins yellowish, not distinctly marked. Length 2 to $3\frac{1}{2}$ inches. Coast of British Columbia; not rare. Here described from 5 specimens from *Albatross* Station 2944, numbered 66 in the L. S. Jr. Univ. Museum. (Named for Capt. Henry E. Nichols, U. S. N., its discoverer.)

Gobius nicholsii, BEAN, Proc. U. S. Nat. Mus. 1881, 469, Departure Bay, British Columbia; JORDAN & GILBERT, Synopsis, 946, 1883; JORDAN & EIGENMANN, *l. c.*, 494.

2538. GOBIUS EIGENMANNI, Garman.

D. VII–12; A. 13; P. 19; scales 27–7. Body rather stout, body cavity more than $\frac{1}{2}$ the length from snout to base of caudal. Head $\frac{2}{9}$ of the total length or $\frac{2}{7}$ of the distance to the caudal base, blunt and rounded anteriorly, very narrow between the eyes, slightly compressed. Eyes large, $\frac{1}{3}$ of the

head, very close together. Snout short, little more than ½ as long as the eye. Mouth wide; maxillary reaching a vertical from the middle of the eye, moderately oblique. First dorsal higher, anterior 3 rays prolonged in the filaments, third ray longest and reaching to the eighth ray of the second dorsal; origin of anal fin midway from edge of preopercle to base of caudal; pectorals nearly as long as the head; caudal as long as head, pointed. Scales large, thin, deciduous, 27 in a longitudinal series, 2 rows above the lateral line. Yellowish, with a few punctulations of black near the bases of the caudal rays, with a light-edged black spot on the outer halves of the fourth to the sixth rays of the first dorsal and with a black streak around the mouth immediately above the maxillary. The long body, the large eye, the dorsal spot, and the streak above the mouth serve to distinguish this species from its nearest allies of the same locality. Off Key West, in 60 fathoms. ("The specific name is given in honor of the distinguished ichthyologists who have added so much to our knowledge of the American Gobiidæ, C. H. and R. S. Eigenmann.")

Gobius eigenmanni, GARMAN, Bull. Lab. Nat. Sci. State Univ. Iowa, vol. IV, No. 1, 88, 1896; off Key West in 60 fathoms. (Coll. Iowa Univ. Bahama Expedition.)

2539. GOBIUS GLAUCOFRÆNUM (Gill).

Head 3⅓; depth 4⅓. D. VI–10; A. I, 9; P. 18; lateral transverse 7. Body robust, compressed; head naked; mouth oblique, the lower jaw slightly projecting, the maxillaries extending to below pupil; teeth long, in many series, the outer curved; scales ctenoid, large. Pectoral fin with the upper rays little branched, not silk-like; cheeks scarcely tumid; caudal and pectoral longer than ventrals, about as long as head. Tawny, with a faint blue spot in the center of each scale, and with 6 spots, each formed by aggregation of dark dots, on the ridge of the back between the second dorsal spine and the axil of the soft dorsal fin; another row of similar but fainter spots runs from the scapular region, and a third row along the middle of the sides; head tawny, with dark spots and blue dots; a straight blue line across the cheek; dorsal fins with faint blue spots. Length 1½ inches. Florida Keys; said to have come from the coast of Washington, but this is probably an error, as the species has not since been taken there, while 1 apparently identical has been taken at Tortugas.* (*glaucus*, glaucous; *frænum*, bridle.)

* Dr. Eigenmann thus describes the specimens from the Tortugas examined by him: *Gobius glaucofrænum* (Gill). Head 3⅔ (4⅜ in total); depth 4⅓ (5⅓). D. VI–10; A. 10; scales in a median series, 23, in a transverse series, 8; eye as long as snout, 3⅓ in head, jaws equal, maxillary barely reaching pupil. Teeth in bands in both jaws, those of the outer row of lower jaw enlarged. Dorsal spines scarcely filamentous, the third highest and equaling depth of body. Posterior dorsal rays highest, as high as spines; anal similar to soft dorsal; pectoral long and narrow, longer than head, 3⅓ in body; ventral reaching past vent; scales large, thin, finely toothed, reduced on breast; nape naked. Color in spirits, light yellowish brown; a light spot on each scale, the spots especially conspicuous near shoulder; 6 dark spots on middle of back; fainter but similar spots along middle of sides; a conspicuous dark spot above opercle; a wavy light line extending forward from it through lower rim of eye to snout; a straight pale-blue bar extending parallel to it across preopercle and cheek to corner of mouth; a narrow faint bar below it; a triangular dark spot at corner of mouth; cheeks and preopercle purplish chocolate; opercle and snout plain yellowish; 2 brown spots at base of caudal; the smaller specimens differing from this in having the markings more distinct. Length of 4 specimens examined, 1⅞, 1 7/12, 1⅓, 1⅛ inches. (Eigenmann.)

Coryphopterus glaucofrænum, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 263, Coast of Washington (evidently an error).

Gobius glaucofrænum, JORDAN & GILBERT, Synopsis, 635, 1883; JORDAN & EIGENMANN, *l. c.*, 494; EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 59.

2540. GOBIUS MANGLICOLA, Jordan & Starks.

Head 4¼; depth 5⅔. D. VI–12; A. 12; scales about 35, not to be exactly counted; caudal lanceolate, 2⅔ in body; pectoral about equal to head; dorsal spine slender, not filamentous, 1⅔ in head; eyes large, close together, the range partly vertical, the narrow interorbital deeply furrowed; no flaps on shoulder girdle; scales moderate, ctenoid anteriorly, becoming smooth behind; median keel on head slight; head naked. Body long, compressed, the head depressed, the cheeks tumid; snout bluntly truncate; mouth large, the maxillary reaching the middle of eye, not produced backward, truncated behind, somewhat oblique, the lower jaw a little the longer; lower jaw flat; teeth strong, the outer in both jaws enlarged; cranium without median crest, abruptly widened behind eyes. Color light olive, mottled with darker; 6 oblong blotches of blackish on sides as in *Gobius boleosoma*, the last at base of caudal; dorsals and caudal finely checkered and barred with dark brownish orange and blackish; anal mottled; a dark shoulder spot; a dark bar before eye and 1 below eye; ventrals dusky, the edge pale. One specimen, 1¼ inches long. Mazatlan; found in the mud of the Astillero among the roots of mangrove bushes (*Rhizophora mangle*), (whence the name *mangle; colo*, I inhabit).

Gobius manglicola, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 496, Mazatlan. (Coll. Hopkins Expedition to Mazatlan. Type, 3095, L. S. Jr. Univ. Mus.)

2541. GOBIUS STIGMATURUS, Goode & Bean.

Head 4; depth 4½; eye 4, about equal to snout. D. VI–12; A. 11 or 12; scales 33. Body subfusiform, little compressed; head blunt, the profile rounded. Mouth small, horizontal, the lower jaw included; maxillary 3 in head, reaching eye. Teeth small, in bands in both jaws, the outer enlarged, those of the upper jaw very slender. Region from nape to dorsal entirely scaled, the scales large, ctenoid, those on nape and belly little reduced. Dorsal spines short, the longest shorter than the head; caudal fin scarcely pointed, about as long as head. Grayish white, middle of sides with 4 or 5 dark blotches, from each of which a dark bar extends downward and forward; a large black blotch above pectoral, obsolete in the female; a small black spot at base of caudal, and a dark mark below the eye; vertical fins barred. Two specimens taken in a shallow bay at Key West are thus described: Very pale olive, everywhere freckled and spotted; lower part of sides silvery, crossed by faint and narrow cross streaks of light brown; sides with about 5 faint dark blotches; a dark blotch below eye and 1 on opercle; a round black spot at base of caudal; bars on verticle fins light olive. Numerous other specimens are less freckled in coloration, and have a more diffuse caudal spot as well as a vague dark spot at the shoulder. The dusky marks on the sides are larger.

We find no other differences, and refer all of them to *G. stigmaturus*. The relations of *G. boleosoma*, *G. stigmaturus*, and *G. encaemus* are certainly very intimate. Florida Keys, not very common, our specimens from Key West. (*στίγμα*, spot; *οὐρά*, tail.)

Gobius stigmaturus, GOODE & BEAN, Proc. U. S. Nat. Mus. 1882, 418, no type locality given, but specimens probably from Florida Keys; JORDAN & GILBERT, Synopsis, 946, 1883; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1884, 140; JORDAN & EIGENMANN, *l. c.*, 495.

2542. GOBIUS QUADRIPORUS, Cuvier & Valenciennes.

D. VI–I, 9; A. I, 9; scales as in *Gobius caninus*. The 2 pores on the vertical arm of preopercle very open; 2 smaller ones above them; teeth of outer series small; 2 small canines on each side of lower jaw; dorsal spines not prolonged as filaments. Color yellowish, with lighter lines which follow the rows of scales; brown spots on dorsal; 2 lines on cheek. Surinam. (Cuvier & Valenciennes.) Not seen by us. (*quatuor*, four; *porus*, pore.)

Gobius quadriporus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 87, 1837, Surinam; EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 61.

2543. GOBIUS SHUFELDTI, Jordan & Eigenmann.

Head $3\frac{2}{3}$ to $3\frac{3}{5}$; depth $5\frac{1}{3}$; eye 4; snout $3\frac{1}{3}$ to $3\frac{1}{2}$. D. VI–12; A. 13; scales 33 to 35. Body moderately elongate, subfusiform; head less blunt than in *Gobius boleosoma*, the anterior profile gently decurved; mouth large, slightly oblique; maxillary extending to front of pupil, $2\frac{1}{3}$ in head. Teeth small, slender, and curved, in moderate bands; scales covering anterior half of trunk; head and breast naked; scales moderate, ctenoid, those in front much reduced. Longest dorsal spine $1\frac{1}{2}$ in head; caudal fin as long as head, somewhat pointed. Olivaceous, mottled with gray; about 5 round dark blotches along middle of side, the last at base of caudal; no dark spot on side of nape; some dark marks on head; vertical fins barred. Gulf coast of the United States, known as yet only from fresh waters about New Orleans. (Named for Dr. Robert Wilson Shufeldt, U. S. A., who collected the types.)

Gobius shufeldti, JORDAN & EIGENMANN, Proc. U. S. Nat. Mus. 1886, 495, New Orleans. (Type, No. 35202.)

2544. GOBIUS BOLEOSOMA, Jordan & Gilbert.

Head 4 (5 in total); depth $4\frac{1}{2}$ to $5\frac{1}{2}$. D. VI–12; A. I, 10 to 12; scales 25 to 30. Body slender, subfusiform, little compressed; head moderate, not very blunt, the anterior profile somewhat evenly decurved, the snout not very short, scarcely shorter than the large eye; mouth not very large, horizontal, the lower jaw included, the maxillary extending slightly beyond front of pupil, its length about 3 in head; teeth small, slender, in narrow bands, those of the outer series longer than the others; eyes placed high, about 4 in head; interorbital space not wider than pupil; scales moderate, ctenoid, those on nape and belly not much reduced in size; gill opening not continued forward above opercle; first dorsal with

the spines slender but rather firm, none of them filamentous, the longest about $\frac{3}{5}$ head; second dorsal and anal rather large; caudal long, pointed, slightly longer than head; pectorals large, slightly longer than head, none of the upper rays silk-like; ventrals slightly shorter than head, inserted below axil of pectorals; skull rounded behind, no ridges nor crests; crests at side minute; interorbital very narrow. Color in life: Male, deep olive green, mottled with darker; middle of side with 4 or 5 vague darker blotches; a jet-black spot above gill opening, on side of back; head mottled, dusky below; usually a dark bar below eye; dorsals tipped with bright yellowish, each crossed by numerous narrow, somewhat oblique, interrupted bars or series of spots, these being of a rich reddish brown color; caudal barred with black, its upper edge tinged with orange; anal nearly plain, with a slight orange tinge; ventrals bluish black, their edges whitish. Female, paler and duller in color, more mottled, the black spot above gill opening obsolete or nearly so; a dark spot at base of caudal; upper fins barred, as in the male; lower fins mostly pale, tinged with orange. Many specimens of this species, the largest about 2 inches in length, were obtained in the Laguna Grande at Pensacola. It lurks in sea wrack on muddy bottoms in very shallow water (6 to 12 inches). In form, size, coloration, and movements this little fish bears a remarkable resemblance to the percoid, *Boleosoma olmstedi*. Gulf of Mexico, Pensacola to Key West; common in shallow sandy bays, lurking in sea wrack at the depth of a foot (whence the name *βολίς*, dart; *σῶμα*, body).

Gobius boleosoma, Jordan & Gilbert, Proc. U. S. Nat. Mus. 1882, 295, Laguna Grande, Pensacola; ibid, Synopsis, 946, 1883; Jordan & Eigenmann, *l. c.*, 495.

2545. GOBIUS FASCIATUS (Gill).

Head 4 in length ($4\frac{3}{8}$ in total); depth 6 (7). D. VI–12; A. 10; scales 30–7. Body slender, elongate; head somewhat pointed; profile rounded, not as much as in *Gobius garmani* and *Gobius boleosoma*; eye large, slightly longer than snout, $3\frac{1}{4}$ in head; interorbital area scarcely wider than pupil; mouth slightly oblique, maxillary extending to below anterior margin of pupil, $3\frac{1}{2}$ in head; lower jaw thin and flat; teeth strong, recurved, in a band in each jaw, the teeth of the outer series of the upper jaw enlarged, several times as large as those of the inner series. Scales finely ctenoid (fallen off anteriorly in specimen examined); antedorsal region and breast naked. Dorsal spines slender, filamentous near tip, not reaching second dorsal, $1\frac{1}{2}$ in length of head; second dorsal of moderate height; caudal (tips broken) about 5 in length, $1\frac{2}{3}$ in length of head; ventral not reaching vent, $1\frac{1}{3}$ in head; pectorals pointed, equaling the head in length. Color yellowish, marbled with darker above; 4 oblong dark blotches along middle of sides; a darker spot at base of caudal; narrow dark stripes across nape; a faint dark stripe along upper margin of opercle, through lower margin of eye to snout; another extending from angle of mouth to edge of preopercle, then extending down along the margin of the preopercle and ending in a dark blotch on the lower part of the cheek; a dark spot on opercle; first dorsal with 2 curved bars; caudal with 3 rather broad dark bars; anal

dusky; connecting membrane of ventral white, its first rays blackish, outer rays yellowish; lower parts yellowish. West Indies; not seen by us. This description by Eigenmann, from a specimen 1¾ inches long, No. 13231, M. C. Z., collected in Hayti by Dr. Weinland. (*fasciatus*, branded.)

Ctenogobius fasciatus, GILL, Fishes Trinidad, 378, 1858, Trinidad.
Gobius fasciatus, GÜNTHER, Cat., III, 34, 1861; JORDAN & EIGENMANN, Proc. U. S. Nat. Mus. 1886, 495; EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 62.

2546. GOBIUS ENCÆOMUS, Jordan & Gilbert.

Head 4; depth 6; snout 3⅓. D. VI–11; A. 12; scales 27 to 33. Body very elongate, much tapering backward; head compressed, the cheeks high and vertical; snout very short, compressed, obtusely rounded vertically. Mouth nearly horizontal, low, large, the maxillary 2 in head, nearly reaching vertical from posterior margin of orbit. Teeth in very narrow bands in both jaws, those of the outer series in the upper jaw much enlarged and recurved in some specimens; eyes inserted high, the interorbital space very narrow, about as wide as pupil; diameter of orbit much greater than snout, nearly ⅓ of head. Gill opening 2⅓ in head, the isthmus wide. Dorsals contiguous, the membrane of spinous dorsal reaching nearly to base of soft dorsal; dorsal spines high, of nearly uniform length, the last reaching well beyond origin of soft dorsal when depressed; the longest spine about ½ length of head; soft dorsal and anal long and high, the posterior rays of both fins reaching at least to base of caudal when depressed; caudal lanceolate, the middle rays produced, 2⅔ in body; ventrals reaching vent, somewhat longer than pectorals, which about equal length of head; ventral sheath well developed, its length ⅔ that of fin. Body wholly covered with large, strongly ctenoid scales, which are much reduced in size anteriorly; head, antedorsal region, and breast naked. In female specimens the mouth is evidently smaller, and the caudal less elongate. Colors in life: Male, light olivaceous, mottled above with darker olive brown; a series of about 4 obscure oblong dark blotches along middle of sides; a dark spot at base of caudal; each side of nape with an intense blue-black spot larger than eye; an obscure dusky streak from eye forward to mouth; a small dusky spot sometimes present on upper portion of base of pectorals; both dorsals translucent, with a series of bright reddish-brown spots as large as pupil; upper lobe of caudal light reddish, the lower lobe blue black; anal and ventrals dusky bluish, pectorals slightly dusky, with a narrow, bright pink border behind. Female, without bright markings; body light olive, with 5 oblong dark blotches on sides, the last on base of caudal; from each of the 3 middle blotches a V-shaped bar runs to the back (these visible also in males); back somewhat mottled with dusky; a black blotch on scapula; a small one on opercle; a dark bar from eye forward to mouth. Vertical fins with dusky streaks, these appearing on caudal in the form of cross bars; ventrals light, with 2 lengthwise dark streaks; pectorals plain. South Carolina to Key West, in sandy bays; scarce. Length 2 inches. (*ἐγκαίω*, brand; *ὦμος*, shoulder.)

Gobius encæomus,* JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 611, Charleston, South Carolina (Type, No. 29673, 3 specimens. Coll. C. H. Gilbert); JORDAN & GILBERT, Synopsis, 945, 1883; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1884, 142; JORDAN & EIGENMANN, *l. c.*, 496.

2547. GOBIUS STIGMATICUS (Poey).

Head 4; depth 5 to 6; eye $3\frac{1}{3}$. D. VI–12; A. 12 or 13; scales 27. Body a little deeper and less compressed than in *Gobius encæomus*. Anterior profile moderately decurved; back slightly arched; skull flattish behind, much broader than in *G. boleosoma*, with an evident median ridge; mouth oblique, large, lower jaw thin and flat, maxillary reaching to below pupil. Teeth above uniserial, some of them enlarged and recurved; lower teeth in a narrow band, males sometimes with the hindmost of the outer series a strong, exserted, recurved canine (present in Poey's type). Anterior half of body scaled except region between nape and dorsal, which is naked; breast naked. Longest dorsal spine $\frac{2}{3}$ head, sometimes elongate; caudal $3\frac{1}{2}$ in body. Light greenish, sides of male with 5 or 6 narrow, straight, whitish or yellowish cross bars, regularly placed; 4 dark bars on head, 3 below the eye and 1 on opercle; a small dark spot behind and above opercle; ventral fins barred; female with a row of irregular dark spots connected by a dusky streak, the pale cross bars obsolete. Coast of North Carolina, Florida Keys, the West Indies, southward to Rio Janeiro; common at Havana. Subject to considerable variation. Brazilian specimens said by Eigenmann to be darker, the bars on cheek conspicuous; third dorsal spine often much elongate, reaching fifth dorsal ray, last soft ray sometimes reaching caudal. (*stigmaticus*, spotty.)

Smaragdus stigmaticus, POEY, Memorias, II, 281, 1861, Cuba.

Gobionellus stigmaticus, POEY, Synopsis, 394, 1868; POEY, Enumeratio, 126, 1876; JORDAN & GILBERT, Synopsis, 947, 1883.

Gobius stigmaticus, JORDAN, Proc. U. S. Nat. Mus. 1886, 49; JORDAN & EIGENMANN, *l. c.*, 496.

2548. GOBIUS LYRICUS, Girard.

Head $4\frac{1}{3}$; depth $4\frac{2}{3}$. D. VI–11; A. I, 10; scales 27. Body rather elongate, moderately compressed; head rather short, the profile very obtuse, descending abruptly from before the front of the eye to the snout; eyes small, placed high, about as long as snout, and about $4\frac{1}{2}$ in head; mouth nearly horizontal, much below level of eye, the maxillary extending to beyond pupil, $2\frac{2}{3}$ in head; jaws subequal; teeth strong, in 1 series in each jaw; in the lower jaw about 4 shortish, canine-like teeth behind the other teeth; anterior teeth of lower jaw small, of upper jaw rather large; gill

* One small specimen, taken with the seine in a shallow bay, at Key West, is described as follows:

Light green, with 5 diffuse spots of darker green on sides, the posterior one most conspicuous; pectorals, both dorsals, and caudal edged above with pale orange; ventrals mostly black, edged with paler; anal dark; a conspicuous dusky shoulder spot; maxillary reaching to below middle of eye; caudal about $\frac{1}{3}$ longer than head. Lateral line about 30. This little specimen appears to be identical with that described by us from Charleston under the name *Gobius encæomus*. The species is allied to *G. stigmaturus*, but has a much slenderer body. The number of scales in a lateral series is less than 37, the number originally stated by us. There are about 33 in this specimen. (Jordan & Gilbert.)

opening not continued forward above opercle; first dorsal with 2 or 3 spines filamentous, the longest reaching past the middle of the second dorsal, which is of moderate height and similar to the anal; caudal long and pointed, ¼ longer than the head; pectoral as long as head, about reaching front of anal; upper rays of pectorals not silk-like; ventrals somewhat shorter than head, their insertion below front of pectorals; scales large, rough, those on nape, pectoral region, and belly reduced in size; head naked. Color in life, dark olive, with 4 or 5 irregular confluent blackish cross bands, besides dark blotches and irregular markings; head marbled with darker, the jaws, opercles, and branchiostegals blackish; first dorsal mostly dusky translucent, somewhat barred; second dorsal and anal plain dusky; caudal dark blue, with 2 longitudinal stripes of bright red; pectoral finely barred or reticulated with blackish and pale; head and belly yellowish. Female specimens duller and paler. Gulf of Mexico, from Galveston to Cuba and the Lesser * Antilles; rather common. (*lyricus*, pertaining to a lyre, apparently an allusion to the dorsal spines.)

Gobius lyricus, GIRARD, Proc. Ac. Nat. Sci. Phila. 1858, 169, Brazos Santiago, Texas; GIRARD, U. S. and Mex. Bound. Surv., 25, pl. 12, figs. 4 and 5, 1859; GÜNTHER, Cat., III, 550, 1861; JORDAN & EIGENMANN, *l. c.*, 496; EIGENMANN & EIGENMANN, *l. c.*, 63.

Smaragdus costalesi, POEY, Memorias, II, 280, 1861, Havana. (Type, No. 13109, M. C. Z. Coll. Felipe Poey.)

Gobius wurdemanni, GIRARD, Proc. Ac. Nat. Sci. Phila. 1858, 169, Brazos Santiago; probably the female; JORDAN & GILBERT, Synopsis, 634.†

Euctengobius lyricus, JORDAN & GILBERT, Synopsis, 633, 1883.

2549. GOBIUS GARMANI, Eigenmann & Eigenmann.

Head 4 in length (5½ in total); depth 4 (5½). D. VI–11; A. 11; scales 30–7. Body robust, head short and blunt; profile in front of eye abruptly decurved, rounded much as in *Gobius boleosoma;* mouth inferior, horizontal; lower jaw included; maxillary extending to below pupil, 2½ in head; lips thin; teeth short and thick, in a single series in each jaw. Dorsals contiguous; dorsal spines filamentous, the second and third longer than the rest, reaching past first third of second dorsal; last dorsal rays reaching base of caudal; pectorals equaling head in length; ventral short and broad, 5 in body; caudal rather long and pointed, 3 in body. Scales large, slightly reduced and cycloid on nape. Color yellowish, marbled with brown; a series of irregular blotches along the sides; a light spot at base

*A specimen from St. Kitts is thus described by Eigenmann: "Depth 5 in length; head 4. The second and third dorsal spines extend to base of caudal; dorsal scarcely less than length of head, the last rays reaching past base of caudal; the caudal fin is ⅔ longer than the head, 2¼ in body. Color light brown, faintly marked with darker; the first dorsal with minute dark points, the lower fourth of the spines with simple dark spots, above which are jet-black spots ocellated with white; the second dorsal fin dusky, darker posteriorly, the basal portion of the last half of the fin evenly black, the anterior 4 rays marked with dark points similar to the spots on the lower parts of the spines of the first dorsal; caudal dusky, with 2 light bars; anal plain, darker than body; ventral fins blackish, edged with white; pectorals blackish, with many series of white spots on the membrane, and short, white bars at base; branchiostegal membrane black, with a light margin.

†*Gobius wurdemanni*, Girard. Appearance of *Gobius lyricus*. Reddish brown, obscurely barred with dusky. Head larger; caudal shorter; ventrals shorter; anal lower; scales smaller than in *G. lyricus*; teeth very slender, much smaller than in *G. lyricus*; third dorsal spine filamentous. D. VI-11; A. 12. Brazos Santiago, Texas. (Girard.) (Named for Dr. Gustav Würdemann, its collector.)

of caudal, partly or wholly surrounded by a broad ring of dark brown; head slate color, white below; 3 dark bars extending forward and downward from eye to mouth; a triangular spot on opercle; dorsals, caudal, and pectorals finely barred with black; a chocolate bar on base of ventral; anal margined with white; an irregular black bar on shoulder and upper half of pectoral; everywhere more or less blotched with darker, the blotches at times forming numerous bars across the back. Dominica, Fort de France, Martinique, St. Kitts. (Eigenmann & Eigenmann.) Not seen by us. Apparently very close to *Gobius lyricus*, if not the same. (Named for its discoverer, Prof. Samuel Garman.)

Gobius garmani, EIGENMANN & EIGEMANN, Bull. Cal. Ac. Sci. 1888, 61, Dominica, Fort de France, Martinique, St. Kitts. (Coll. Samuel Garman.)

2550. GOBIUS ZEBRA, Gilbert.

Head 3; depth $4\frac{1}{3}$; eye $3\frac{1}{4}$ in head. D. VI–11 or 12; A. 9. Body not elongate, the snout short, the mouth oblique, with maxillary reaching below middle of orbit. Mouth small, the maxillary $2\frac{1}{4}$ in head. Interorbital space very narrow. Teeth in upper jaw in a narrow band or double series, the outer row enlarged and spaced; lower jaw apparently with a single series, similar to the outer row in the upper jaw. Scales cycloid, large, wanting on nape and a narow strip along base of spinous dorsal. Color cherry red, head and sides with 15 blue cross bars, a little narrower than interspaces, encircling the body posteriorly, lacking for a short distance on belly and under side of head; on upper side of head and nape these bars run obliquely forward and downward, but elsewhere vertical; on middle of each interspace a very narrow blue line, becoming indistinct on lower part of sides; on cheeks the blue bars are connected by narrow cross lines, forming blue reticulations surrounding round spots of the ground color. Length of types $\frac{1}{2}$ inch. Two specimens from *Albatross* Station 2989, west coast of Mexico, in 36 fathoms. (*zebra*, zebra, from the stripes.)

Gobius zebra, GILBERT, Proc. U. S. Nat. Mus. 1890, 73, Albatross Station 2989, west Coast of Mexico.

Subgenus EUCTENOGOBIUS, Gill.

2551. GOBIUS POEYI, Steindachner.

Head broad and flattish; depth $6\frac{1}{2}$ in total length; eye $4\frac{1}{4}$, $1\frac{1}{5}$ in interorbital width, longer than snout; snout short and decurved. D. VI–9; A. 9; scales 40. Maxillary extending to below middle of eye. Some of the dorsal spines produced and filamentous, the third $1\frac{1}{2}$ times depth of body; caudal short, rounded. Two rows of ill-defined blotches on upper half of body; dorsals and caudal sharply barred, anal and ventrals dusky (male). A small round dark spot at base of caudal. (Steindachner.) Barbados; not seen by us. (Named for Prof. Felipe Poey.)

Gobius poeyi, STEINDACHNER, Ichthyol. Notizen, VI, 44, 1867, Barbados; JORDAN & EIGENMANN, *l. c.*, 497.

2552. GOBIUS BADIUS (Gill).

Head 6 in total; depth 7. D. VI–I, 10; A. I, 10; scales 50–18. Anterior profile very oblique; a line of pores above each eye; 2 on upper ascending margin of preopercle; eye 4 in head; interorbital space $3\frac{1}{8}$ in eye; caudal 5 in total length; pectoral 6. Color dark bay with a posteriorly straight heavy dot in the center of each scale on back and sides above; head plumbeous, with 2 livid blue bands from eye to upper jaw. (Gill). About mouth of Amazon; not seen by us. (*badius*, bay color, dark red.)

Euctenogobius badius, Gill, Ann. Lyc. Nat. Hist. N. Y., vii, 1857, 47, Amazon.
Gobius bosci, Sauvage, Bull. Soc. Philom. Paris, iv, 44, 1880.
Gobius badius, Eigenmann & Eigenmann, *l. c.*, 65.

Subgenus **GOBIONELLUS**, Girard.

2553. GOBIUS MICRODON, Gilbert.

Head $4\frac{1}{6}$; depth 5. D. VI–13; A. 14; scales 62. Head and body compressed, everywhere deeper than wide. Mouth at lower profile of snout, nearly horizontal, the lower jaw extremely weak, broadly rounded anteriorly; maxillary reaching vertical from hinder margin of pupil, nearly $\frac{1}{2}$ length of head. Teeth minute scarcely perceptible without the use of a lens, those in upper jaw in a single series. Mandible with a close set outer series of teeth, separated by an interval from an inner narrow band of still smaller teeth. Interorbital space narrow, less than diameter of pupil. Isthmus wide, the gill slits extending little below base of pectorals. Scales minute and cycloid anteriorly and on belly, becoming larger posteriorly; on sides they are everywhere ctenoid behind the middle of spinous dorsal; belly wholly scaled; nape scaled forward nearly to orbits, but with a narrow median naked streak running back to front of dorsal; breast and sides of head naked. Dorsal fins not connected. First 4 spines filamentous, the longest longer than head, reaching when depressed to base of third ray of soft dorsal. Soft dorsal and anal similar, not high, the last rays not extending beyond the base of caudal; caudal lanceolate, much longer than head; pectorals and ventrals about equal, reaching vent. Color nearly uniform light olive, with minute darker punctulations which sometimes form darker margins to the scales; an oblique dusky streak on opercle; 3 or 4 oblique obscure dark cross bars on spinous dorsal, and 4 or 5 on tail; ventrals with white pigment. Length 2 inches. San Juan Lagoon, west coast of Mexico. (Gilbert.) (*μικρός*, small; *ὀδούς*, tooth.)

Gobius microdon, Gilbert, Proc. U. S. Nat. Mus. 1891, 554, San Juan Lagoon, north of Rio Ahomè, Mexico. (Coll. Gilbert.)

2554. GOBIUS SMARAGDUS, Cuvier & Valenciennes.

(Esmeralda.)

Head 4; depth $5\frac{1}{4}$ to $5\frac{7}{8}$; eye 4 to 5. D. VI–11 or 12; A. 11 or 12; scales 39 to 42. Body moderately elongate, compressed; head not compressed; the cheeks tumid; the snout short and abruptly decurved; mouth large, little oblique; lower jaw slightly inferior; maxillary reaching to below

pupil or to posterior margin of orbit, $2\frac{1}{8}$ to $2\frac{1}{3}$ in head; outer row of teeth on upper jaw enlarged; the narrow band of teeth back of this row separated from it by a space; teeth on lower jaw in a band, subequal. Scales cycloid anteriorly, becoming larger and ctenoid posteriorly. Caudal $2\frac{1}{4}$ to $2\frac{1}{2}$ in body. Male, light olive, with dark-olive blotches; body and head with many conspicuous round cream-colored spots, each surrounded by a dusky ring, these smaller than pupil and most distinct on head; snout with dusky streaks; dorsal and caudal plainly barred; pectoral crossed with dark wavy lines, dusky at base; anal and ventrals dusky; a small dark spot at base of caudal; a shining deep-green spot inside the mouth in life. Female, plain olivaceous, nearly or quite immaculate. West Indies, south to Rio Janeiro, north to St. Augustine, Florida (Dr. Oliver P. Hay), and to Charleston (C. H. Gilbert); specimens before us from Marco Island, Florida (J. A. Henshall). (*σμάραγδος*, emerald, from the bright-green spot on the tongue.)

Gobius smaragdus, Cuvier & Valenciennes, Hist. Nat. Poiss., xii, 120, 1837, Cuba; Jordan & Eigenmann, *l. c.*, 497.
Smaragdus valenciennei, Poey, Memorias, ii, 280, 1861, Cuba.
Gobionellus smaragdus, Poey, Synopsis, 394, 1868; Poey, Enumeratio, 126, 1876.

2555. GOBIUS STRIGATUS, O'Shaughnessy.

Head $3\frac{3}{4}$; depth 5; eye $3\frac{1}{2}$, shorter than the rounded snout. D. VI–12; A. 11 or 12; scales 53–13. Body elongate, compressed posteriorly; head little compressed; maxillary reaching to below middle of eye; teeth small, the outer a little enlarged; dorsal spines all shorter than head, not filamentous. Head naked; anterior half of body covered with ctenoid scales, those on nape much reduced in size. Two violet stripes from mouth to eye, 8 or 9 violet bars on side, 3 or 4 on caudal; second dorsal spotted. (Steindachner.) Coast of Surinam. (*strigatus*, striped.)

Gobius strigatus, O'Shaughnessy, Ann. Mag. Nat. Hist., series 4, xv, 1875, 145, Surinam.
Gobius kraussi, Steindachner, Ichth. Beiträge, viii, 16, 1879, Surinam; Jordan & Eigenmann, *l. c.*, 497.

2556. GOBIUS SAGITTULA (Günther).

Head $4\frac{1}{2}$ to 5 in length to base of caudal; depth 6 to 8; eye $4\frac{3}{4}$ to $5\frac{1}{4}$. D. VI–13 or 14; A. 13; scales about 66 in longitudinal, 15 in transverse series, counted just below space between the two dorsals. Body slender, tapering pretty regularly from middle of first dorsal to caudal, most compressed posteriorly, depth about uniform from head to origin of second dorsal. Head short, depressed, and broad; mouth large, nearly horizontal, the maxillary in adults $2\frac{3}{5}$ in head, reaching beyond middle of eye; distance between maxillaries at their posterior ends greater than their length; eye about $\frac{2}{3}$ the bony interorbital space. Teeth in a narrow band in each jaw, those in lower jaw uniform, the outer series in upper jaw considerably enlarged and separated by an interspace from the inner band. Pseudobranchiæ well developed. Gill rakers short and flexible. Longest dorsal spine about $\frac{3}{5}$ head; distance between dorsals less than diameter of eye; pectorals $1\frac{1}{3}$ in head, their tips reaching past middle of spinous dorsal; ventrals about equaling pectorals, reaching more than halfway to origin of anal; anal equal and opposite to the second dorsal, but slightly lower; caudal fin greatly elongate, more than $\frac{1}{4}$ head and

body in largest specimens, 2½ in smaller ones, its relative length increasing with age. Head scaleless, predorsal region with small scales; body covered with close-set ctenoid scales, small and greatly crowded anteriorly, toward the caudal fin growing gradually larger and more strongly ctenoid. General color light yellowish, palest below, upper parts darker; middle of sides with 5 elongate black blotches, most distinct in the young; the first under first dorsal, second under origin of second dorsal, the third, which is sometimes almost double, at about middle of second dorsal, the fourth near its posterior end, and the last at base of caudal; a large black spot upon each shoulder just above origin of pectoral fin; head plain; lips and maxillary dark; opercle with a dark blotch; basal portion of dorsal fins with dark lines formed of spots; anal unmarked; pectorals with cross lines formed of dots; ventrals plain; caudal crossed by numerous narrow dark bars. Reaching a length of 8 inches. Gulf of California and neighboring waters south to Panama; very common in lagoons and mouths of rivers. The types of *sagittula* are evidently the young, those of *longicauda* the adults of the same species. (diminutive of *sagitta*, arrow.)

Euctenogobius sagittula, GÜNTHER, Proc. Zool. Soc. London 1861, 3, West coast Central America, young individuals; GÜNTHER, Cat. Fishes, III, 555, 1861; GÜNTHER, Fishes of Centr. Amer., 389, 1869.

Gobius longicauda, JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 146, adult examples, Guaymas. (Coll. Evermann & Jenkins. Type, No. 39636.)

Gobius sagittula, JORDAN & EIGENMANN, *l. c.*, 497.

2557. GOBIUS HASTATUS, Girard.

(EMERALD FISH; SHARP-TAILED GOBY.)

Head 4⅓ to 5; depth 6 to 7⅔. D. VI–14; A. 14 or 15; scales 60; vertebræ 11+15. Body compressed, extremely elongate; depth nearly equal throughout; head short, compressed, deeper than wide; mouth wide, oblique, the jaws equal; maxillary in adult reaching to below posterior border of eye; lower jaw very thin and flat; teeth in each jaw small, subequal, those in the upper jaw in a single series, those in the lower jaw in a narrow band; outer teeth somewhat movable; scales anteriorly small, cycloid, and embedded, those behind larger and ctenoid; the scales larger than in *Gobius oceanicus;* a few scales on upper anterior corner of opercle, but without the large patch seen in *G. oceanicus;* dorsal fins high, some of the spines filamentous and longer than the head; caudal very long and filamentous, 2 to 2⅔ in body; pectoral slightly longer than head or than ventrals, none of its rays silk-like. A single specimen from Ceylon belongs to this species, which appears to be characterized by a longer head (5 in length, 7 in total), by the much larger scales (60 in a lateral line), by the obsolescence of the patch of scales on opercles, and by slightly different coloration. This may be really only the extreme of variation of *G. oceanicus*, with which species most authors have hitherto confounded it. The two need detailed comparison. Coast of Texas. (*hastatus*, spear-like.)

Gobionellus hastatus, GIRARD, Proc. Ac. Nat. Sci. Phila. 1858, 168, St. Josephs Island, Texas; GIRARD, U. S. and Mex. Bound. Surv., 25, pl. 12, figs. 7 and 8, 1859.

Gobius lanceolatus, GÜNTHER, Cat., III, 50, 1861, and of authors; not of BLOCH.

2558. GOBIUS OCEANICUS, Pallas.

(Esmeralda; Endormi Émeraude; Bacalhao Sabara.)

Head 4⅓ to 6; depth 6⅛ to 8½; eye 4 to 5 in head; ventral 6 to 6½; pectorals 5⅞ to 6½. D. VI–14. A. I, 14 or 15; scales about 65. Body extremely elongate; head very short; upper part of opercle scaled, head otherwise naked. Scales on body very small, becoming much larger behind. All the dorsal spines more or less filamentous; caudal fin nearly half length of rest of body. Skull behind eye broad and short, its length 1½ in width, no decided ridges nor crests; lateral crests large and stout behind, minute forward; interorbital area narrow, deeply grooved, with a median ridge. Color in spirits, reddish olive; a distinct, round, blackish blotch below spinous dorsal, twice as large as orbit; an indistinct dusky shade along middle of sides, terminating in a dusky blotch on base of caudal; middle of sides with a series of marks, formed by very veiny lines widely diverging backward; a similar narrow line from eye to maxillary, and 1 from eye backward to upper angle of preopercle; evident traces of the emerald spot at base of tongue; 2 small dark spots on first dorsal spine; spinous dorsal dusky, with a light and dusky streak at base; soft dorsal dusky, a light (bluish in life) area behind each ray; anterior rays barred with light and dark; anal and ventrals whitish (probably blue in life), the ventrals without dark markings; pectorals dusky, the base lighter, and with some indistinct dusky bars; a dusky half bar on the upper part of the axil; base of tongue tuberculate, and shining with bright blue and green reflections like a precious stone (hence the names *smaragdus, esmeralda,* etc.), this color fading in spirits. Vertebræ elongate, 11 + 15 = 26. Length a foot. South Atlantic and Gulf coasts of the United States and southward through the West Indies; not rare, perhaps intergrading with the preceding. Here described from a specimen 11 inches long, taken by Dr. Gilbert in Charleston Harbor. (*oceanicus,* ocean.)

Gobius cauda longissima acuminata, Gronow, Zooph., 82, No. 277, pl. 4, fig. 4, 1763, locality unknown.

Gobius oceanicus, Pallas, Spicilegia, viii, 4, 1769, locality unknown; after Gronow; Jordan & Eigenmann, *l. c,* 497.

Gobius lanceolatus, Bloch, Fische Deutschlands, ii, 8, pl. 38, fig. 1, 1783, Martinique, figure probably from Plumier; Cuvier & Valenciennes, Hist. Nat. Poiss., xii, 114, 1837; Poey, Synopsis, 393, 1868.

Gobius bacalaus, Cuvier & Valenciennes, Hist. Nat. Poiss., xii, 119, 1837, Surinam (Coll. Le Valliant); Cayenne (Coll. Richard); Cuba (Coll. Poey).

Gobionellus occanicus, Jordan & Gilbert, Proc. U. S. Nat. Mus. 1882, 613; Jordan & Gilbert, Synopsis, 636, 1883.

Subgenus LYTHRYPNUS, Jordan & Evermann.

2559. GOBIUS DALLI, Gilbert.

Head 3⅜; depth 4⅛. D. VI–17; A. 14; scales 40. Body short, compressed, resembling *Microgobius.* Head high, mouth moderate, very oblique; upper pectoral rays normal; scales ctenoid, of moderate size; anterior dorsal spines much produced. Mouth very oblique, the maxillary

reaching vertical from front of pupil, 2½ in length of head. Snout short, ⅔ diameter of orbit, which is 3 in head. Jaws with an outer series of long, distant, canine-like teeth, and an inner series or a narrow band of minute teeth. Dorsal spines 6, the 2 anterior greatly elongate, not free, in our largest specimen extending beyond middle of soft dorsal; membrane from last dorsal spine reaching to, or nearly to, base of first soft ray; soft dorsal rather high, the fin long; caudal rounded, less than length of head; ventrals free from belly, fully united; pectorals short, the upper rays not free nor silk-like. Scales of moderate size, ctenoid, covering entire trunk, with possible exception of the nape; the scales are readily caducous, and are lacking on nape and frequently on anterior third of body in our specimens. Color light coral red, anteriorly with 4 to 6 narrow blue bands not reaching ventral outline, the posterior ones growing narrower and fainter; a blue streak upward and backward from each orbit, the 2 uniting on occiput; a transverse interorbital bar, a continuation of which encircles the orbit anteriorly; below orbit, a blue bar consisting of 2 portions, 1 running downward and obliquely backward, the other upward and backward; in the largest specimen a blue streak runs from occiput along profile to front of dorsal; the first blue bar runs from nape obliquely downward and forward, ending on opercle; the second vertically downward from front of spinous dorsal, the third under middle of spinous dorsal, the remaining bars under soft dorsal; fins unmarked. Several small specimens, the largest 1 inch long, from *Albatross* Station 3001, in 33 fathoms. A single slightly larger example dredged by Dr. W. H. Dall, in about 35 fathoms, off Catalina Harbor, California. Probably the type of a distinct genus distinguished by the many-rayed fins and the form of the body and head. (Named for its discoverer, William Healey Dall.)

Gobius dalli, GILBERT, Proc. U. S. Nat. Mus. 1890, 73, Albatross Station 3001, Lower California (Coll. *Albatross*); Catalina Harbor (Coll. W. H. Dall).

814. GARMANNIA, Jordan & Evermann.

(HALF-NAKED GOBIES.)

Garmannia, JORDAN & EVERMANN, Proc. Cal. Ac. Sci. 1895, 495, pl. 49 (*paradoxus*).
Enypnias, JORDAN & EVERMANN, new subgenus (*seminudus*).

Anterior half of body naked; posterior half covered with moderate or small scales; teeth rather strong, unequal, usually 2 small curved canines in front. Very small gobies. Otherwise essentially as in *Gobius*. ("Named for Mr. Samuel Garman, the accomplished ichthyologist of the Museum of Comparative Zoology at Cambridge, Mass., in recognition of his important contributions to ichthyology.")

GARMANNIA:

a. Scales moderate.

b. Scales ctenoid, 13 or 14 series developed; first dorsal spine filamentous; D. VI–11; A. 9. Body rather robust, the depth about 4⅔ in length; the head 3⅓; lower jaw with 2 curved canines. PARADOXA, 2560.

bb. Scales smaller, 17 series developed; depth 4⅔ in length. D. VI–10; A. 8; first dorsal spine not filamentous; lower jaw with small canines.
HEMIGYMNA, 2561.

ENYPNIAS (ἐνύπνιος, in one's sleep):

aa. Scales excessively minute; body slender, the depth 6 in length. D. VI–15; A. 10; dorsal spines not filamentous; lower jaw with 2 small curved canines in front. SEMINUDA, 2562.

Subgenus GARMANNIA.

2560. GARMANNIA PARADOXA (Günther).

Head about $3\frac{1}{2}$ ($4\frac{1}{4}$ in total); depth about $4\frac{2}{3}$ ($5\frac{2}{3}$ with caudal). D. VI–11; A. 9; scales 14. Head nearly as broad as high, its width being rather more than $\frac{1}{2}$ of its length. Eyes rather close together, of moderate size. Snout obtuse, rounded, as long as the eye; cleft of the mouth slightly oblique, with the jaws equal in length, and with maxillary extending to below middle of the eye. Teeth in villiform bands; 2 curved canine teeth on each side of the lower jaw. Head and trunk entirely naked to between second dorsal and anal, the remainder covered with ctenoid scales of moderate size, 9 or 10 of them in 1 of the anterior transverse series. First dorsal spine elongate, filiform, sometimes extending to the base of the caudal; caudal rounded, shorter than head; none of pectoral rays silk-like; ventral terminating at a great distance from vent. Blackish in spirits; caudal and ventral fins black, dorsal filament whitish. (Günther.) Panama to Mazatlan; scarce. Our single specimen from the estuary at Mazatlan differs somewhat from Dr. Günther's account. It is thus described: Head $3\frac{1}{2}$; depth $4\frac{1}{3}$. D. VI–11; A. 9; scales 12; eye 4 in head; snout $4\frac{1}{4}$; pectoral $1\frac{1}{6}$ in head; dorsal spine $1\frac{1}{3}$. Form of *Gobiosoma bosci*. Body compressed; head broad and depressed, with tumid cheeks; snout not very blunt, short, oblique-truncate; eyes rather large, high, the maxillary not produced, extending to their posterior margin; mouth large, oblique; lower jaw heavy, slightly projecting; teeth strong; gill openings narrow, not wider than base of pectoral. First dorsal rather high, the first spine filamentous, reaching past soft dorsal; other fins low. Head and anterior half of body to front of soft dorsal naked; scattering scales coming in above, 12 rows of imbricated slightly ctenoid scales along median line of caudal peduncle and forward to middle of soft dorsal, the scaled area about as long as head, the upper parts better scaled than lower. No flaps on shoulder girdle. Olivaceous, with 7 or 8 dark cross shades, 2 on head, 1 across gill openings, 1 behind pectoral, and a broad 1 below soft dorsal; dorsals dusky, the filamentous ray pink; lower half of soft dorsal yellowish, upper dusky; lower fins black; caudal dusky; a dark speck at angle of opercle; skin everywhere punctate with black; a pale olive bar at base of caudal. Skull without median crest; interorbital space not concave; head not very abruptly widened behind eyes. Pacific coast of Mexico and Central America. One specimen, $1\frac{1}{2}$ inches long, recently obtained on muddy bottoms among the mangroves lining the estuary at Mazatlan. (*paradoxus*, paradox.)

Gobius paradoxus, GÜNTHER, Proc. Zool. Soc. Lond. 1861, 3, west coast Central America; GÜNTHER, Cat., III, 549, 1861; JORDAN & EIGENMANN, Proc. U. S. Nat. Mus. 1886, 498.
Garmannia paradoxa, JORDAN, Proc. Cal. Ac. Sci. 1895, 497, pl. 59.

2561. GARMANNIA HEMIGYMNA (Eigenmann & Eigenmann).

Head $3\frac{3}{8}$ ($4\frac{1}{2}$ in total); depth $4\frac{3}{8}$. D. VI–10; A. 8; scales smaller than in *Garmannia paradoxa*, 17–7. Body compressed, depressed anteriorly, the greatest depth in this specimen being at origin of anal and second dorsal fins. Head wider than deep, rounded; profile much decurved from eye to mouth as in *paradoxa;* eye perfectly round, smaller than in *paradoxa*, $1\frac{1}{3}$ in rounded snout, 5 in head; interorbital space scarcely wider than orbit; mouth somewhat oblique, larger than in *paradoxa*; maxillary reaching beyond posterior rim of orbit; lower jaw slightly shorter than upper; teeth in upper jaw in a band, the outer series remote, and the teeth several times as large as in the inner row, all more or less movable; teeth in lower jaw similar, a recurved canine on each side near the front. Scales very weakly ctenoid, covering only the sides of the posterior half of body, not extending quite to base of dorsal or anal fins even at their posterior insertion; the upper and lower edges of the caudal peduncle likewise free from scales, the scaly region, however, widest on peduncle and tapering forward to the central point opposite beginning of anal, where the scales are smallest. First spine of the dorsal not elongate as in *G. paradoxa*, $1\frac{1}{2}$ in head, the third, fourth, and fifth spines slightly exceeding the first in height, equaling the posterior rays of soft dorsal, which are little higher than the anterior rays of the soft dorsal; caudal rounded, about 4 in length of body, $1\frac{1}{4}$ in head; ventral not reaching vent, $1\frac{1}{3}$ in head; pectorals rounded, rather short and broad, $1\frac{1}{4}$ in head. Color light olivaceous, without distinct markings, everywhere with minute dark punctulations; 8 faint cross bars from dorsal to middle of sides, which, close under dorsal fins, are formed of 2 blackish dots; 8 black dots along lateral line, the last at base of caudal; fins all smutty, the pectoral lightest, white on its anterior half, 2 dusky spots at its base; opercle ashy; a light bar at base of caudal; iris blackish blue, a short straight streak of same color from eye to upper lip; an irregular bluish mark on cheeks formed of punctulations closely crowded. West Indies, exact locality unknown; taken with the dredge. (*ἥμι*, half; *γυμνός*, naked.)

Gobius hemigymnus, Eigenmann & Eigenmann, Proc. Cal. Ac. Sci. 1888, 66, dredged in the West Indies.

Subgenus **ENYPNIAS**, Jordan & Evermann.

2562. GARMANNIA SEMINUDA (Günther).

Head 4; depth 6. D. VI–15; A. 10. Head and anterior portion of trunk naked; sides with exceedingly small scales, becoming somewhat larger posteriorly. Head with the cheeks swollen, depressed, broader than high, its width $\frac{2}{3}$ length. Eyes close together, directed upward, of moderate size; snout obtuse, as long as the eye; cleft of the mouth slightly oblique, with the jaws equal anteriorly, and with the maxillary extending to below the middle of the eye. Teeth in villiform bands, the anterior of the lower jaw slightly enlarged; 2 small curved canine teeth on each side of lower jaw. Dorsal fins rather low, the hind part of the spinous dorsal scarcely lower than anterior; caudal rounded, as long as pectoral; none of pectoral

rays silk-like; ventral rather short, terminating at a great distance from vent. Blackish; fins and sides of head dotted with black; ventrals black. (Günther.) Panama; not seen by us; probably the type of a distinct genus. (*seminudus*, half-naked.)

Gobius seminudus, Günther, Proc. Zool. Soc. London 1861, 3, west coast Central America; Günther, Cat., III, 554, 1861; Jordan & Eigenmann, Proc. U. S. Nat. Mus. 1886, 498.

815. AWAOUS, Steindachner.

Awaous,* Steindachner, Verh. Mat. Phys. Naturw. 1860, 289; after *les Awaous* of Cuvier & Valenciennes (*ocellaris*, etc).

Chonophorus, Poey, Memorias, II, 274, 1861 (*buccelentus*=*taiasica*).

Awaous, Bleeker, Esquisse d'un Syst. Nat. Gobioides, 320, 1874 (*ocellaris*); after *les Awaous* of Cuvier & Valenciennes.

Inner edge of shoulder girdle with 2 or more conspicuous dermal flaps; preorbital region very long; premaxillary and maxillary strong; lips thick; scales rather small, ctenoid, 40 to 80 in a longitudinal series; interorbital groove with a conspicuous median crest; otherwise essentially as in *Gobius*. The species reach a large size and are confined to the fresh waters of the tropics of America and the Hawaiian Islands. The Asiatic species of similar habit have much larger scales and seem to form a distinct genus. *Rhinogobius*, Gill. The physiognomy in each is peculiar, the snout being long and convex. (*Awaou*, a Hawaiian name.)

a. Scales about 53, little crowded anteriorly, 21 before dorsal on nape; depth 5⅔ in length; head 4; eyes placed high, interorbital area equal to diameter of eye; mouth horizontal; maxillary extending to middle of eye, 2⅓ in head, lower jaw more flat than in *A. taiasica*; teeth small, in narrow bands, those of the outer row above enlarged, some large teeth in band of lower jaw. D. VI–I, 12; A. I, 10. Uniform yellowish in spirits. FLAVUS, 2563.

aa. Scales 60 to 70, crowded anteriorly, about 30 scales before the dorsal on nape; body compressed posteriorly, rather depressed anteriorly; greatest depth 5¼ in length; head 3⅓ in length. Olivaceous, a series of irregular, roundish blotches along middle of sides; narrow dark streaks radiating from eye; a blackish streak running across upper margin of opercle and extending obliquely across base of upper pectoral rays; belly white; dorsal and caudal more or less distinctly barred with wavy blackish lines.

b. About 15 scales between second dorsal and base of anal. NELSONI, 2564.

bb. About 21 scales between second dorsal and base of anal. TAIASICA, 2565.

aaa. Scales 76 to 82; 24 scales between second dorsal and anal; head as broad as high; depth of body 6⅔ in length; head 4; head flat above, snout elongate, upper profile oblique; eye ⅓ of head, equals interorbital area (in adult); mouth horizontal; lower jaw included; maxillary reaching to below anterior margin of eye; teeth of the outer series enlarged; canine teeth none; scales ctenoid, those on nape and anterior part of body very small; head naked; dorsal fins lower than body, none of the spines produced; caudal rounded, 7 in length of body. Yellowish olive; back and sides reticulated with blackish; head, dorsal, caudal, and pectoral fins dotted with blackish, the spots forming streaks on second dorsal; 6 cross series of dots on the caudal; an irregular small blackish spot on the upper part of the root of pectoral. D. VI–11; A. 11; scales about 80. MEXICANUS, 2566.

* The name "Les Awaous," given to this group by Valenciennes, was a French plural, not a generic appellation, and if used as the name of a genus must be dated from its use in that sense by Steindachner or Bleeker. The Hawaiian type of "*Awaous*" agrees with the American species (*Chonophorus*) in the character of the flaps on the shoulder girdle, as well as in general appearance. The Asiatic genus *Rhinogobius*, Gill (*similis*), seems to be very close to *Chonophorus*, but the scales are larger, 28 in the lateral series.

2563. AWAOUS FLAVUS (Cuvier & Valenciennes).

Head 4; depth 5⅔ to 6½; eye equal to the interorbital width, placed high. D. VI–I, 12; A. I, 10; scales about 53 to 55, little crowded anteriorly, 21 before the dorsal. Mouth horizontal, maxillary extending to middle of eye, 2⅓ in head; lower jaw flatter than in *Awaous taiasica;* teeth small, in narrow bands, those of the outer row enlarged; some large teeth in band of lower jaw. Yellowish, with a row of faint ocellated spots along middle of sides; dorsal and caudal faintly barred; lines radiating from eye, a line along opercle halfway to pectoral, sometimes uniform blue-black. Rivers of Surinam and Brazil, south to Bahia. (*flavus,* yellow.)

Gobius flavus, Cuvier & Valenciennes, Hist. Nat. Poiss., xii, 60, 1837, Surinam; Günther, Cat. Fish., iii, 13, 1861.
Chonophorus flavus, Jordan & Eigenmann, *l. c.*, 500; Eigenmann & Eigenmann, *l. c.*, 67.

2564. AWAOUS NELSONI, Evermann.

Head 3½; depth 6; eye 5¾ in head; snout 3; maxillary 2⅞. D. VI–11; A. 11; scales about 63. Body long, compressed and tapering posteriorly; head large, quadrate, mouth nearly horizontal, lower jaw included; snout abruptly decurved; top of head flat, the interorbital with a slight median groove with a thin, raised edge on each side; maxillary reaching about to vertical of anterior edge of pupil; teeth in bands on jaws very small, the outer somewhat enlarged; pectoral rays normal, the longest 1⅓ in head; ventrals completely united, the disk free from belly, 1¾ in head. Dorsal fins separated by a space about ⅔ diameter of eye; dorsal spines slender, weak, about 1⅔ in head; soft dorsal and anal similar, each free from caudal; caudal fin rather short and rounded, its middle rays about 1¼ in head. Gill membranes broadly united to the isthmus; eyes moderate, high up, the interorbital width equal to the eye's diameter. Scales ctenoid, very small and irregularly crowded anteriorly, much larger posteriorly, about 15 rows counting from origin of soft dorsal downward and backward to the anal fin; head naked, but with slight indication of a few minute embedded scales on opercles. Color grayish; head mottled and blotched with dark; side with 7 or 8 black blotches, the largest under middle of pectoral fin; dorsals pale, crossed by several lines of black spots; caudal pale, with about 6 or 7 dark cross bars; ventrals and anal pale; pectorals pale, dusted with dark specks and with a small dark blotch at base of upper rays. Close to *A. taiasica,* but with broader interorbital, longer snout and larger scales on posterior part of body. Length 4 inches. Known only from fresh water at Rosario, Sinaloa, where 8 specimens were obtained July 27, 1897, by Mr. E. W. Nelson. (Named for Mr. Edward William Nelson, the well-known ornithologist, in recognition of his work upon the fishes of Illinois in 1876.)

Awaous nelsoni, Evermann, Proc. Biol. Soc. Washington, vol. xii, 1898, 3, fresh-water pools at Rosario, Sinaloa, Mexico. (Type, No. 48836, U. S. Nat. Mus.; cotypes, No. 533 U. S. Fish Comm., 5793 L. S. Jr. Univ. Mus., and 48837 U. S. Nat. Mus.)

2565. AWAOUS TAIASICA (Lichtenstein).

(Guavina Hoyera; Aboma de Rio.)

Head 3¼; depth 5¼; eye small, less than interorbital width (in adult), 3 in snout (twice in young), and about 7 in head. D. VI–11; A. 11; scales 60 to 70, crowded anteriorly, about 30 before dorsal fin, 21 between second dorsal and anal. Body compressed posteriorly, rather depressed anteriorly; head broader than deep. Distance from eye to mouth 3½ in head, the preorbital being much enlarged; mouth large, horizontal, maxillary extending to below anterior part of orbit in adult male, shorter in young; lower jaw included. Teeth of the upper jaw in 2 series, those in anterior series much enlarged and recurved; teeth of lower jaw in a narrow band, the outer series scarcely enlarged. Inner edge of the shoulder girdle with 2 or 3 rather long papillæ. Body covered with ctenoid scales, much reduced in size anteriorly; nape closely scaled, breast scaly, head naked. Dorsal fins less than depth of body, the spines scarcely filamentous, not as long as the soft rays; caudal rounded, shorter than the head; ventrals very broad and short, 1½ to 1⅔ in head, the rays very much branched. Skull rounded behind, with a very short crest in its middle; lateral crests high and thin, converging into 1 opposite the insertion of suprascapula, inner crests not meeting behind eye, the outer ones extending around orbit. A low, blunt ridge between the posterior corners of orbit, becoming much higher forward, continued as the ethmoid and ending abruptly some distance in advance of orbit. Teeth in upper jaw in a few series, those of outer series many times larger than the others, which are minute; those of lower jaw all alike small, in a band. Olivaceous, with a series of irregular, roundish blotches along middle of side, and narrow dark streaks radiating from eye; a blackish streak running across upper margin of opercle and extending obliquely across base of upper pectoral ray; belly white; dorsal and caudal more or less distinctly barred with wavy blackish lines. Length a foot or more. Extremely variable in form and coloration, as is the case with most widely distributed fresh-water fishes. Fresh waters of the West Indies and both coasts of Mexico, south to Brazil; common in Cuba, in Sinaloa, and about La Paz in Lower California, thence southward to Panama. (*taiasica*, Brazilian name of some other goby.)

Amore guacu, Marcgrave, Hist. Brasil., 166, 1648, Brazil.

Gobius taiasica, Lichtenstein, Berl. Abhandl., 273, 1822, Brazil; not *Tajasica* Marcgrave.

Gobius banana, Cuvier & Valenciennes, Hist. Nat. Poiss., xii, 103, 1837, San Domingo; Günther, Cat., iii, 59, 1861.

Gobius martinicus, Cuvier & Valenciennes, Hist. Nat. Poiss., xii, 105, 1837, Martinique.

Chonophorus bucculentus, Poey, Memorias, ii, 275, 1861, Cuba.

Rhinogobius contractus, * Poey, Memorias, ii, 424, 1861, Cuba; Poey, Enumeratio, 125, 1875

* The following are the characters assigned to *Awaous contractus* (Poey): Head 4; depth 5½; D. VI–11; A. 11; eye 7 in head; maxillary ceasing ½ an eye's diameter before eye. Head smaller than in *A. taiasica*. Greenish brown; the cheeks with brown lines; body with brown points; dorsals brownish, with brown longitudinal bands more numerous on the second; caudal with 7 brown bands, made of lanceolate spots on the rays; pectorals speckled; ventrals and anal rose color. Cuba (Poey); probably not different from *A taiasica;* said to differ in the small mouth, which probably varies with age and sex.

Gobius dolichocephalus, COPE, Trans. Amer. Philos. Soc. Phila. 1869, 403, near Orizaba, Mexico.
Euctenogobius latus, O'SHAUGHNESSY, Ann. Mag. Nat. Hist., Series 4, XV, 1875, 146, Bahia. (Coll. Dr. Wucherer.)
Chonophorus taiasica, JORDAN & EIGENMANN, *l. c.*, 500.

2566. AWAOUS MEXICANUS (Günther).

Head 4; depth $6\frac{3}{5}$; eye 8. D. VI–11; A. 11; scales 76 to 82, 24 between second dorsal and anal. Head as broad as deep, flat above, snout elongate, upper profile oblique; mouth horizontal, lower jaw included, maxillary reaching to below anterior margin of eye. Teeth of the outer series enlarged; no canine teeth. Scales ctenoid, those on nape and anterior part of body very small; head naked. Dorsal fins lower than depth of body, none of the spines produced; caudal rounded, 7 in length of body. Yellowish olive; back and sides reticulated with blackish; head, dorsal, caudal, and pectoral fins dotted with blackish, the spots forming streaks on second dorsal; 6 cross series of dots on caudal; an irregular, small blackish spot on the upper part of the base of the pectoral. (Günther.) Fresh-water streams of the eastern slope of Mexico; known to us only from Dr. Günther's description.

Gobius mexicanus, GÜNTHER, Cat., III, 61, 1861, Mexico.
Chronophorus mexicanus, JORDAN & EIGENMANN, *l. c.*, 501.

816. BOLLMANNIA, Jordan.

Bollmannia, JORDAN, Proc. U. S. Nat. Mus. 1889, 164 (*chlamydes*).

This genius differs from *Lepidogobius* by having no fleshy processes on inner edge of shoulder girdle, the interorbital area of skull narrower and without trace of median keel, and by very large ctenoid scales. From *Gobius* proper it is distinguished by the presence of 7 dorsal spines and by the presence of large scales on the cheeks. Species inhabiting the depths of the Pacific; not found in shoal waters as is the case with most other gobies. ("I have named this species in honor of my late colleague, Mr. Charles Harvey Bollman, whose untimely death, while engaged in the exploration of the rivers of Georgia, took place while this paper was passing through the press."—Jordan.)

a. * A conspicuous black spot on posterior portion of spinous dorsal. Body deep, the least depth of caudal peduncle greater than diameter of orbit.
b. Filamentous dorsal spines very long, reaching beyond middle of soft dorsal when depressed. Lower caudal rays black; dorsal spot conspicuously ocellated. Eye large, 3 to $3\frac{1}{4}$ in head. OCELLATA, 2567.
bb. Filamentous dorsal spines shorter. Lower caudal rays not black, and dorsal spot not ocellated. Eye smaller, $3\frac{3}{4}$ to 4 in head. CHLAMYDES, 2568.
aa. No black spot on spinous dorsal. Body slender, the depth $\frac{1}{5}$ the length. Least depth of caudal peduncle not greater than diameter of orbit.
c. Head large, 3 to $3\frac{1}{4}$ in length. No black spot at base of caudal. Fins low. MACROPOMA, 2569.
cc. Head smaller, $3\frac{2}{3}$ in length. A black spot at base of caudal. Fins higher. STIGMATURA, 2570.

* This analysis of species is taken from Gilbert, Proc. U. S. Nat. Mus. 1891, 555.

2567. BOLLMANNIA OCELLATA, Gilbert.

Head 3½ to 3⅔ in length; depth 4½. D. VII–14 or 15; A. 14; scales 27. Very close to *Bollmannia chlamydes,* differing from the latter constantly in the following respects: The eye is larger, 3 to 3⅕ in head (3¾ to 4 in *chlamydes*); the filamentous rays of spinous dorsal are much longer, reaching in adults, when laid back, to or nearly to end of base of soft dorsal, 1½ to 1⅞ times length of head; rarely the filamentous dorsal rays are little more elongate than in *chlamydes.* Teeth in a narrow band in each jaw, the outer series in upper jaw, and both outer and inner series in lower jaw enlarged, but not canine-like; maxillary not reaching vertical from middle of pupil, ½ length of head; interorbital width less than ½ diameter of pupil; opercle short, its length being less than the diameter of the eye; pectorals nearly as long as head, a trifle more than length of ventrals, which scarcely reach vent; caudal much longer than head in adults, 7 or 8 scales before dorsal. Black spot on posterior part of spinous dorsal jet black, conspicuously ocellated with white; a black streak along lower margin of caudal, including several of the lower rays, and running from base to tip of fin; no dusky bars visible on sides in any of the types; fins dusky; membranes uniting outer rays of ventrals white instead of black, as in *chlamydes;* no black spot at base of caudal; branchiostegal membrane with a medial black streak; anal blackish. Numerous specimens from the northern part of the Gulf of California, at *Albatross* Stations 3031 and 3035, in 30 and 33 fathoms. This species may vary into the typical *chlamydes,* but the material before us does not justify us in so identifying it. (Gilbert.) (*ocellatus,* ocellated.)

Bollmannia ocellata, Gilbert, Proc. U. S. Nat. Mus. 1891, 555, Gulf of California.

2568. BOLLMANNIA CHLAMYDES, Jordan.

Head 3½ (5 to 5½ in total); depth 4½ (6⅔ to 7). D. VII–15; A. 15; scales in a longitudinal series about 28, 8 or 9 in a cross series at vent. Body rather robust, compressed; head large and heavy, its profile evenly curved; mouth very large, oblique, the lower jaw projecting; maxillary reaching to opposite pupil, 2½ to 2⅔ in head; teeth small, sharp, in several series, the outer, especially in lower jaw, somewhat enlarged; eye longer than snout, 3¾ to 4 in head; interorbital area very narrow, concave, its least width about ⅓ of eye or almost equal to pupil; scales very large, ctenoid; little reduced on breast and nape; about 8 before dorsal, where they are little smaller than on body; top and sides of head with large scales; scales on cheek in 4 rows; 2 rows on upper part of opercle; the scales on head lost in some of the specimens; dorsal spines slender, filamentous, fifth longest, 1⅕ in head; first 2 in head, last 3½ to 4; first soft dorsal ray 2⅔ in head, the antepenultimate longest and about equal to head; first anal ray equal to snout, the antepenultimate 1¼ in head; middle caudal rays very long, somewhat more than ½ length of body; pectorals 1⅕ in head; ventrals 1⅓. Color olivaceous, darkest above; scales with a few black dots, some of the posterior occasionally dark edged; sides with 8 or 10 obscure dusky vertical bars, which are narrower than the inter-

spaces, and in some specimens wholly obsolete; snout bluish; opercles with a dark shade; lips, gular region, and anterior branchiostegals very dark in males; upper part of spinous dorsal darkest, with a few lighter dark-edged oval spots, a well-marked black blotch between last 2 spines; soft dorsal dusky, usually with about 3 well-developed rows of lighter, dark-edged oval spots; anal dusky, crossed by 2 narrow bluish streaks; some of the last rays occasionally with a few spots similar to those on dorsal; caudal, pectorals, and ventrals dusky, tinged with blue; ventrals edged with pale. Length 4¾ inches. West coast of Colombia. Many specimens of this abundant species were dredged at *Albatross* Stations 2800 in 7 fathoms and 2805 in 51½ fathoms. (*χλαμύδης*, cloaked.)

Bollmannia chlamydes, JORDAN, Proc. U. S. Nat. Mus. 1889, 164, Pacific Ocean, off coast of Colombia, Station 2800, 8° 51′ N., 79° 41′ 30″ W., and Station 2805, 7° 56′ N., 79° 41′ 30″ W. (Type, No. 41158, U. S. Nat. Mus. Coll. *Albatross.*)

2569. BOLLMANNIA MACROPOMA, Gilbert.

Head 3 to 3½; depth 5. D. VII–14; A. 14; scales 28. Characterized by its slender form, low fins, large opercle, and comparatively plain coloration. Caudal peduncle correspondingly slender, its least height equaling diameter of eye. Head very large and heavy; opercle conspicuously larger than in *B. ocellata*, agreeing in this respect more nearly with *B. chlamydes*; dentition as in other species of the genus; eye large, 3¼ to 3⅓ in the head. Dorsal spines slender, comparatively little produced, the longest usually not reaching the base of the first ray of second dorsal, and never beyond the base of the second or third ray; soft dorsal and anal low, the posterior rays usually not reaching the rudimentary caudal rays when depressed, about ½ length of head; pectoral long, extending beyond front of anal; the ventrals to or nearly to vent; middle caudal rays produced as usual, varying in length; scales 8 to 10 in front of dorsal. Color in spirits, light brownish, the sides with 3 vertical dusky bars; spinous dorsal dusky, but without distinct black spot; caudal slightly dusky, with rather large elliptical light spots, as in *B. chlamydes*, the lower rays not black and no black spot at its base; ventrals blackish, including anterior membrane; second dorsal and anal dusky, without evident light spots; branchiostegal membranes sometimes slightly dusky, but not black. Many specimens from the Gulf of California just north of La Paz Bay, at *Albatross* Station 2996, in 112 fathoms. (Gilbert.) (*μακρός*, large; *πῶμα*, opercle.)

Bollmannia macropoma, GILBERT, Proc. U. S. Nat. Mus. 1891, 556, Albatross Station 2996, near La Paz, Lower California.

2570. BOLLMANNIA STIGMATURA, Gilbert.

Head short, 3⅜ in length; depth 5; least depth of caudal peduncle slightly less than diameter of eye; eye large, 2⅓ in head. D. VII–15; A. 14; lateral line 28. Dorsal spines filamentous, longer than in *B. macropoma*, the longest reaching base of fifth to seventh ray of second dorsal; posterior rays of second dorsal and anal often reaching base of median rays when depressed; pectorals not reaching beyond front of anal. Color

almost uniform light brownish; lips black, the fins only slightly dusky, the caudal with elliptical light spots; a roundish dusky spot at base of caudal; branchiostegal membranes not black. Many specimens from the northern part of the Gulf of California, at *Albatross* Stations 3016 and 3017, in 76 and 58 fathoms. This species agrees with *Bollmannia macropoma* in its elongate form, comparatively low fins, and in the absence of a black spot on the spinous dorsal. It differs conspicuously in the very short head and narrow opercle, and in the presence of a black spot at base of tail. The eye is also larger and the fins higher. None of the specimens shows dusky bars on the sides, a conspicuous feature in *B. macropoma*. (Gilbert.) (*στίγμα*, spot; *οὐρά*, tail.)

Bollmannia stigmatura, GILBERT, Proc. U. S. Nat. Mus. 1891, 556, Gulf of California, Albatross Stations 3016, 3017.

817. ABOMA, Jordan & Starks.

Aboma, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 497 (*etheostoma*).

This genus, allied to *Microgobius*, is distinguished by the large, ctenoid scales, which cover the body; head naked, rather long, pointed in profile; the mouth moderate, not very oblique; teeth rather strong. Dorsal spines more than 6, none of them filamentous; soft dorsal and anal short; no flaps on shoulder girdle. Cranium with a slight median crest. (The name *Aboma* is used by the Mexicans in Sinaloa as synonymous with goby.)

a. Scales very large, 26 or 27; profile not very steep, the snout rather pointed.
b. Sides with a jet-black lateral band; caudal with dark cross bars; maxillary 3 in head. ETHEOSTOMA, 2571.
bb. Sides with 4 oblique dark cross bars; a large dark spot at base of caudal; mouth larger, the maxillary 2 in head. LUCRETIÆ, 2572.
aa. Scales smaller, about 37; profile very steep, the snout rounded; sides with numerous pale cross bands with darker spots. CHIQUITA, 2573.

2571. ABOMA ETHEOSTOMA, Jordan & Starks.

Head 3⅓; depth 5. D. VIII–11; A. 10; scales 26; longest dorsal spine 1¾ in head; eye 3; snout 4; maxillary 3. Body long and low, moderately depressed and pointed forward. Scales large, ctenoid behind, none on head, those on nape and belly much reduced. Mouth moderate, terminal, moderately oblique; the maxillary reaching middle of pupil; jaws subequal, or the lower a little the longer; teeth rather strong. No flaps on shoulder girdle. Cranium with a slight median crest. Interorbital ridge not hollowed out; skull not abruptly widened behind. Color olivaceous, side with a very broad jet-black lateral band, 3 times interrupted by silvery; caudal white, with 4 < shaped bands, growing progressively fainter behind; pectoral mottled gray, with a jet-black oblique crescent toward its base surrounding a large yellow spot; side of head with 4 round gray spots separated by black, the largest below eye, with a black streak before it; first dorsal jet-black, second mottled, the produced spine with yellowish; ventral and anal pale. A single small specimen, 1⅓ inches long,

found in the mud on a shallow bottom in the Astillero at Mazatlan. (*Etheostoma*, a darter, which this species strongly resembles.)

Aboma etheostoma, Jordan & Starks, Proc. Cal. Ac. Sci. 1895, 498, pl. 50, Mazatlan. (Coll. Hopkins Expedition to Mazatlan.)

2572. ABOMA LUCRETIÆ (Eigenmann & Eigenmann).

Head 3½ in length; depth 5⅓. D. VII–10; A. 12; scales 28–8. Body slightly compressed posteriorly; head little wider than high; eye placed high, its diameter equaling length of snout, 4½ in head; profile little decurved; mouth large, oblique; maxillary extending below posterior margin of orbit, 2 in head; intermaxillary anteriorly on a level with center of pupil; teeth all recurved, large, those of upper jaw in a narrow band; teeth of outer and inner series enlarged, those of lower jaw similar, largest in front. No dermal flaps on shoulder girdle. Scales large, very weakly ctenoid, becoming cycloid and very much crowded above and below pectoral; head, breast, and anterior part of nape naked. As seen through a lens, these regions seem to be covered with minute embedded scales; this effect is, no doubt, due to light reticulations on a darker ground. Dorsal spines slender, not filamentous; caudal pointed, 3 in length of body; ventrals 1¼ in head; pectorals longer than head. Color light brownish, with 4 oblique dark cross bars as wide as interspaces; 4 narrower transverse bars on nape and back; a large dusky spot at base of caudal; upper half of base of pectoral black; a black spot on opercle, margined below and behind with silvery; fins dusky. Pearl Island, Gulf of Panama; only 1 specimen known. (Named for Mrs. Lucretia M. Smith of San Diego, mother of Mrs. Eigenmann.)

2573. ABOMA CHIQUITA (Jenkins & Evermann).

Head 3⅓ to 3½; depth 4⅓ to 4⅜. D. VII–11; A. 10; eye 4⅜ in head in adult, 4 in young; scales 37–17. Body rather stout, compressed; head short, somewhat depressed, widened behind orbits; snout short and narrowly rounded; profile in front of eye very steep, less so to occiput, and nearly straight from there to caudal fin; eyes moderate, well up; interorbital space very narrow, less than eye; greatest width of head equaling greatest depth of body. Top of head, opercles, and space in front of dorsal naked, rest of body covered with small, strongly ctenoid scales, which increase in size upon the caudal peduncle. Spinous dorsal with its first spine filamentous in adult, much longer than head and reaching middle of soft dorsal, this filament wanting in young; distance from snout to origin of spinous dorsal a little more than ⅓ distance to base of caudal; second dorsal but slightly separated from spinous, its origin about midway of total length of fish; anal of about the same shape and size as soft dorsal, but beginning a little behind it; pectorals tapering, about equaling head in length, their tips not reaching origin of anal, but to origin of soft dorsal; ventrals united, free from belly, inserted behind pectorals, but their tips not reaching tips of pectorals. Teeth apparently in a single series, small and weak. Ground color pale yellowish, thickly mottled with fine punc-

tulations of dark; about 7 pretty well-defined larger spots of dark brown along middle of side; 8 or 9 faint cross bars of lighter, a number of small light spots scattered irregularly over the sides; head dark; dorsal, anal, and ventral fins covered with fine black points; in some specimens the dorsals and anal quite dark; pectorals plainer; caudal similar to ventrals; "the cranium is depressed and flattish behind the orbits, without distinct median keel on occiput or on interorbital area. The form of the head is as in typical *Gobius*, the occiput abruptly widened behind the eyes; the ridges also similar, the orbital ridge bounding the orbit behind as well as above the eye and joining the temporal ridge laterally." (Gilbert MS.) Length 1 to 2 inches. Gulf of California; abundant. The original description from young examples, here corrected in accordance with Dr. Gilbert's notes on many adults taken by him at La Paz. (Spanish, *chiquito;* a diminutive of *chico,* a little one.)

Gobius chiquita, JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 146, Guaymas, Sonora. (Type, No. 39634. Coll. Jenkins & Evermann.)

818. MICROGOBIUS, Poey.

Microgobius, POEY, Enumeratio, 127, 1875 (*signatus*).

Dorsal spines 7 or 8; scales very small, cycloid or weakly ctenoid, the body scaled anteriorly as well as posteriorly, the head naked, the nape, belly, and breast usually so. Inner edge of shoulder girdle without fleshy processes; body more or less compressed; mouth large, very oblique; the lower jaw conspicuous, teeth strong; interorbital groove with or without a median ridge. Vertebræ 11+15 or 16. (*μικρός*, small; *Gobius.*)

a. Scales about 42. Body elongate, moderately compressed, the depth 4 to 5 in length; head long and large, rather sharp in profile, 3 to 3½ in body; eye longer than snout, 4 in head; mouth large, very oblique, the lower jaw strongly projecting; maxillary 1½ to 2½ in head, extending to opposite middle of eye, or much beyond front of orbit; teeth in few series, the outer very long and slender, curved, the lower longest, none canine-like; scales small, some of them with short, thick teeth, those of anterior part of body not well developed; dorsal spines more or less filamentous, the third and fourth or fourth and fifth sometimes with long filaments; caudal pointed, about as long as head. Grayish olive, with rather sharply-defined markings of darker brown overlaid with orange in life; head with a pale bluish or gilt stripe from maxillary backward across suborbital region to upper edge of gill opening; another pale gilt streak from snout along lower part of eye, another from angle of mouth upward and backward; rest of head dark; opercle with an oblique blackish bar; top of head and nape with dark marbling surrounded by paler reticulations; back with a series of black cross blotches mostly separated on the median line; 2 narrower dark vertical bars behind pectoral; middle line of side posteriorly with longitudinally oblong black blotches; besides these, numerous other blotches not regularly arranged; first dorsal with 2 or 3 oblique black bands; second dorsal pale, with about 4 series of black dots; caudal spotted with black; pectoral yellowish; ventral black, its center yellowish (male); anal pale. D. VII–15; A. 16 or 17. GULOSUS, 2574.

aa. Scales about 50; snout not pointed; depth 5½ in length; mouth large, the maxillary 2¼ in head; teeth strong. Color yellowish, much dotted, but without bars. EULEPIS, 2575.

aaa. Scales 65 or more.

b. Caudal fin more than ⅛ (⅞) length of body. Scales very small, cycloid, deciduous. Body elongate, much compressed, highest in front of ventrals, tapering regularly to the very narrow, short caudal peduncle; greatest depth 4⅜ in length; head 3½. Head compressed, much higher than wide; snout very short, acute, preorbital not as wide as pupil; mouth terminal, very wide and oblique; jaws equal; maxillary reaching vertical from middle of orbit, 2 in head. Outer series of teeth enlarged. Eye 3 in head. Dorsals closely contiguous; spines very slender, the fifth slightly produced and filamentous; pectorals as long as head. Head and body translucent, overlaid by brilliant green luster, formed by minute, close-set green points; 3 conspicuous translucent bars wider than the interspaces, crossing body close behind head; head with 2 brilliant narrow blue and green lines running obliquely across cheek below eye; dorsal whitish, with 2 or 3 lengthwise series of large reddish-brown spots; spinous dorsal blackish at base, upper caudal rays marked with red, the lower portion of caudal and most of the anal fin blackish, anal whitish at base, the anterior rays tipped with white. In spirits, body dusted with dark points; 2 light cross bars toward head; lower part of caudal and anal black. D. VII–16; A. 15. THALASSINUS, 2576.

bb. Caudal fin less than ⅛ length of body. Scales small, cycloid, embedded. Body very much compressed, more or less elongate, greatest depth at ventrals 4 (female) to 6⅔ (male) in length; head 3⅓ to 4. Head much compressed, much deeper than wide. Snout very short, acute, the anterior profile not decurved, not steep; preorbital not as wide as pupil; mouth very large, very oblique or almost vertical; maxillary extending to below pupil, 2 in head (in male, 2¼ in female). Lower jaw projecting, the teeth of the outer series enlarged, recurved. Eye 3¼ to 4 in head. Dorsals contiguous, spines very fine, produced in filaments, the third highest, a little longer than head; second dorsal and anal high. Head and nape naked. In the female the depth is greater, mouth less oblique, smaller; profile from spinous dorsal oblique. First dorsal spine highest, 3⅓ in length. Ventrals much shorter than in males. Dark gray; female with a short bright blue bar bordered by blackish above pectorals; a blotch of sky blue and orange below eye; fins dusky, the ventrals pale in female, dusky in males. Males with the body plain bluish gray. D. VII–17 to 20; A. 18 to 21; scales 68 to 70. SIGNATUS, 2577.

2574. MICROGOBIUS GULOSUS (Girard).

Head 3 to 3½; depth 4 to 5; eye 4 in head, longer than snout. D. VII–15; A. 16 or 17; scales about 42; vertebræ 11 + 15. Body elongate, moderately compressed; head long and large, rather sharp in profile; mouth large, very oblique, the lower jaw strongly projecting; maxillary 1½ to 2½ in head, extending to opposite middle of eye. Teeth in few series, the outer very long and slender, curved, the lower longest, none canine-like. Body entirely scaled, except the nape, belly, breast, and head, which are naked; scales small, some of them with short thick teeth, those on anterior part of body not well developed. Dorsal spines more or less filamentous, the third to fifth sometimes with long filaments; caudal pointed, as long as head. Ventrals as long as pectorals, which are 1¼ in head. Skull flattened behind, with a median ridge extending from eyes back to end of skull. Double crests bordering skull in front and on sides, the inner ones meeting in front of median crest. Interorbital very narrow and deeply grooved, with a median ridge. Frontal bones very thin and fragile.

Teeth on each jaw in narrow bands, all alike. Coloration in life, light grayish olive, with rather sharply defined markings of darker brown; head with a pale bluish stripe from behind the angle of the mouth upward and forward parallel with the gape to below front of eye, then turning abruptly backward across suborbital region to upper edge of gill opening; another pale streak from snout along lower part of eye; between this and the first streak a dusky area; below the first-mentioned streak a dusky region on cheek; opercle with an oblique blackish bar; top of head with dark marblings surrounded by paler reticulations; back with a series of black cross blotches, mostly separated on the median line; 2 narrow vertical dark bars behind pectoral; middle line of side posteriorly with longitudinally oblong black blotches; besides these numerous other blotches not regularly arranged; first dorsal with 2 or 3 oblique black bands; second dorsal pale, with about 4 series of black dots; caudal spotted with black, pectoral yellowish, ventral black, its center yellowish; anal pale; lower side of head pale; jaws dusky. Coast of Florida to Texas, in sandy or weedy bays, common north to Indian River. A strongly marked species with no near relative among our other gobies. The specimens here described from Pensacola. (*gulosus*, large-mouthed.)

Gobius gulosus, Girard, Proc. Ac. Nat. Sci. Phila. 1858, 169, Indianola, Texas; Girard, U. S. and Mex. Bound. Surv., Zool., 26, 1859; Jordan & Gilbert, Synopsis, 634, 1883.
Lepidogobius gulosus, Jordan & Gilbert, Proc. U. S. Nat. Mus. 1882, 294; Jordan & Gilbert, Synopsis, 945, 1883.
Microgobius gulosus, Jordan & Eigenmann, *l. c.*, 505.

2575. MICROGOBIUS EULEPIS, Eigenmann & Eigenmann.

Head 4 in length (5⅛ in total); depth 5½ (7). D. VII–15; A. 16; scales 50–14. Body elongate, scarcely compressed; head slightly higher than wide, the depth 1⅛ in its length; eye large, longer than snout, 3⅛ in head; snout 5 in head, rather broad, not pointed as in *M. thalassinus;* preorbital narrower than pupil; mouth very oblique, maxillary not extending beyond anterior margin of pupil, 2¼ in head; teeth in upper jaw in a very narrow band, slightly enlarged in outer series, largest toward angle of mouth; teeth of lower jaw in a similar band, some of outer ones in front long and slender. Scales cycloid, rather large, crowded anteriorly, regularly arranged, not embedded as in *M. signatus*, not deciduous as in *M. thalassinus;* breast, nape, and region along spinous dorsal naked. First dorsal spine equidistant from tip of snout and first anal ray; longest dorsal spine 1½ in head; caudal fin about 4 in body; ventral not reaching vent, equaling length of head, the basal membrane ¼ of its actual length; pectoral equaling length of head. Color yellow or very light brown, dotted with minute dark points above; scales along back with a dark margin; head and nape with minute points; spinous dorsal transparent, a marked black spot on upper part of membrane between fourth and fifth dorsal spines; other fins plain; a light vertical bar on posterior margin of preopercle;

no other bars or stripes anywhere. Fortress Monroe, Virginia; known from a specimen 1⅞ inches long. (*εὖ*, well; *λεπίς*, scaled.)

Microgobius eulepis, EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 69, Fortress Monroe, Virginia. (Type, No. 27123, M. C. Z. Coll. Mrs. C. N. Willard.)

2576. MICROGOBIUS THALASSINUS, Jordan & Gilbert.

Head 3½ in length; depth 4¾. D. VII–16; A. 15; eye 3 in head. Body elongate, much compressed, highest in front of ventrals, thence tapering regularly to a very narrow, short caudal peduncle; the body with a peculiar, translucent, fragile appearance, common also to *Z. emblematicus*. Head compressed, much higher than wide; snout very short, acute, the preorbital not as wide as pupil; mouth terminal, very wide and oblique, the jaws equal; maxillary reaching vertical from middle of orbit, ½ length of head; teeth in a narrow band in each jaw, the outer series enlarged, canine-like (under a microscope the band of small teeth behind the outer series seems evident, but the size of our specimens does not enable us to verify it with certainty); eyes placed high, separated by a narrow ridge, the diameter about ⅓ length of head. Dorsals very closely contiguous; spines very slender, the fifth slightly produced and filamentous, reaching (in our specimens) to base of third soft ray when depressed; caudal lanceolate, very long and pointed, the middle rays produced, 2⅔ in body; pectorals as long as head; the upper rays not silk-like; ventrals with basal membranes well developed; the fin long, reaching to or slightly beyond front of anal, somewhat longer than head. Body covered with rather small cycloid scales; head naked; the scales very readily deciduous; as they have in our specimens mostly fallen off, the count can not be given. Head and body translucent, overlaid by brilliant green luster, which is formed by exceedingly minute close-set green points; the luster is intense toward the head, where it assumes a blue tint, and becomes hardly noticeable on caudal peduncle; 3 conspicuous translucent bars, wider than the interspaces, crossing body immediately behind head; head with 2 brilliant narrow blue or green lines running obliquely across cheek below eye; opercle with greenish luster; branchiostegal membrane white; dorsals whitish, with 2 or 3 lengthwise series of large reddish-brown spots; spinous dorsal blackish at base; upper caudal rays marked with red, the lower portion of caudal and the most of the anal fin blackish, anal whitish at base, the anterior rays tipped with brilliant white; ventrals light buff; pectorals translucent. In spirits, the body appears dusted with dark points; 2 light cross bars toward head; lower part of caudal and anal black. Coast of South Carolina; two specimens, the largest 1½ inches long (No. 29674, U. S. Nat. Mus.), were taken in muddy tide pools in Charleston Harbor. (*θαλασσίνός*, thalassinus, sea-green; *θάλασσα*, the sea.)

Gobius thalassinus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 612, Charleston Harbor, South Carolina. (Coll. C. H. Gilbert.)
Lepidogobius thalassinus, JORDAN & GILBERT, Synopsis, 947, 1883.
Microgobius thalassinus, JORDAN & EIGENMANN, *l. c.*, 505.

2577. MICROGOBIUS SIGNATUS, Poey.

Head 3½ to 4; depth 4 (female) to 6½ (male); eye 3¼ to 4. D. VII–17 to 20; A. 18 to 21; scales 68 to 70; vertebræ 14 + 15. Body very much compressed, more or less elongate; head much compressed, deeper than wide; snout very short, acute, the anterior profile not decurved, not steep; preorbital not as wide as pupil; mouth very large, almost vertical; maxillary extending to below pupil, 2 in head in male, 2¼ in female; lower jaw projecting; teeth of the outer series enlarged and recurved. Dorsals contiguous, spines very fine, produced in filaments, the third longest, a little longer than head; second dorsal and anal high. Scales as in *M. gulosus*. Skull rounded, very fragile; a median crest which is highest between eyes; lateral crests developed, the inner ones meeting above posterior part of eye; interorbital comparatively broad, the median crest ending above anterior part of the orbit. Teeth in each jaw in 2 or 3 series; outer series of the upper jaw enlarged and recurved, the inner ones minute; outer series of lower jaw smaller than those of upper jaw, the one nearest angle of mouth an enlarged canine. Dark gray; female with a short bright blue bar, bordered by blackish above pectoral; a blotch of sky blue and orange below eye; fins dusky, the ventrals pale in female; males with the body plain bluish gray. The sexual differences in this species are very strongly marked. West Indies, in salt water; common in Cuba; one of the smallest gobies, barely 2 inches long. Here described from Havana examples collected by Dr. Jordan. (*signatus*, marked.)

Microgobius signatus, Poey, Enumeratio, 127, pl. 5, fig. 3, 1875, Cuba (Type in M. C. Z. Coll. Poey); Jordan, Proc. U. S. Nat. Mus. 1886, 49; Jordan & Eigenmann, *l. c.*, 505.

819. ZALYPNUS, Jordan & Evermann.

Zalypnus, Jordan & Evermann, Check-List Fishes, 459, 1896 (*emblematicus*).

This genus differs from *Microgobius* in having the anterior half of the body naked. Soft dorsal and anal long, of 16 or 17 rays. Two species known. (ξάλη, surf; ὕπνος, slumber.)

a. Scales 48; shoulder with a round black spot; none of the dorsal spines elongate. CYCLOLEPIS, 2578.

aa. Scales 65; a silvery cross bar behind pectorals; some of the dorsal spines usually elongate. EMBLEMATICUS, 2579.

2578. ZALYPNUS CYCLOLEPIS (Gilbert).

D. VII–16; A. 17; scales 48. Body somewhat elongate, compressed, the mouth very large, narrow, and oblique; maxillary produced beyond the rictus for a distance equaling ⅔ diameter of orbit, reaching vertical from posterior margin of pupil, 1⅔ in head; snout short, 5 in head; eye larger, 3¾ in head; interorbital width ½ orbit; teeth in upper jaw in 2 series, the outer enlarged and distant; in lower jaw apparently in a single series, similar to outer series of upper jaw, with 2 stronger canines anteriorly. Inner edge of shoulder girdle without fleshy prominences. Dorsal spines

7, none of them elongate, the membrane of last spine reaching base of first soft ray; soft anal rays of moderate height, 1½ in head, the tips of last rays reaching base of caudal, the fin similar to soft dorsal but lower; caudal long, apparently rounded posteriorly, longer than head (mutilated in our specimen); ventrals and pectorals reaching vent. Scales cycloid, small, absent on belly, nape, and on sides in front of fourth dorsal spine. Color in spirits, light olive, the fins dusky; a conspicuous round black spot on shoulder, ½ size of eye, its posterior margin denser black. Resembling *Zalypnus emblematicus*, differing in its larger scales and different coloration. A single specimen, about 2 inches long, from Lower California, in 7 fathoms. (Gilbert.) (*κύκλος*, circle, cycloid; *λεπίς*, scale.)

Microgobius cyclolepis, Gilbert, Proc. U. S. Nat. Mus. 1891, 74, Albatross Station 3020, Lower California.

2579. ZALYPNUS EMBLEMATICUS (Jordan & Gilbert).

Head 3⅜; depth 5. D. VII–16; A. 17; scales about 65. Anterior part of body naked; teeth of upper jaw in one series; body elongate, compressed, heaviest forward; depth 5 in length; head 3⅜; snout short, rather broad, acute in profile; mouth terminal, very oblique; gape wide, its length nearly ⅓ head; maxillary reaching to opposite middle of pupil; lower jaw projecting. Teeth in lower jaw partly in 2 series in front, forming a single row laterally; anterior teeth in both jaws strong, incurved. Eyes very large, about ⅓ of head; snout less than orbit. Scales extremely small, cycloid, scarcely increasing in size toward caudal peduncle; head and anterior part of body to front of dorsal fin naked; a narrow naked strip along base of anterior ⅓ of spinous dorsal. Dorsal spines very slender and weak, some of the middle ones usually prolonged, sometimes reaching nearly to the base of caudal, sometimes little elevated; second dorsal and anal similar to each other, the rays high, the last when depressed nearly reaching to the base of caudal; caudal pointed, a little longer than head. Light olivaceous; above thickly punctate with pale dots; sides very thickly covered with golden-green specks; back with 6 pairs of golden-green spots on each side of the dorsal fin, each nearly as large as pupil; sides of head and anterior half of body with wide streaks and bars alternately of purplish blue and golden bronze; those on cheek longitudinal; those on opercle extending obliquely upward and backward, those on body vertical; first dorsal dusky, second dorsal with about 3 series of light-blue spots; anal pale; caudal yellowish green below, dusky above, a very conspicuous narrow bright-red streak from the lower end of the base to the tip of the fifth or sixth ray from the bottom, thus crossing the rays obliquely; ventrals bluish. In spirits, plain light olive, with a silvery cross bar behind pectorals. Length 3½ inches. Panama; known only from the original types. (*ἔμβλημα*, a banner, from the high dorsal.)

Gobius emblematicus, Jordan & Gilbert, Bull. U. S. Fish Comm. 1881, 330, Bay of Panama.
Lepidogobius emblematicus, Jordan & Eigenmann, *l. c.*, 505.

820. EUCYCLOGOBIUS, Gill.

Eucyclogobius, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 279 (*newberryi*).

This genus is allied to *Lepidogobius*, differing chiefly in the naked head and short, chubby body; shoulder girdle with a few dermal flaps; opercle adnate to shoulder girdle from the angle upward; dorsal spines 6 or 7; soft dorsal short; scales all cycloid; cranium depressed behind the parietal region, somewhat excavated, the supraoccipital crest rather high, not extending so far forward as the orbit. Species small, in fresh or brackish waters of California. (*εὖ*, well; *κύκλος*, circle (cycloid); *Gobius*.)

2580. EUCYCLOGOBIUS NEWBERRYI (Girard).

Head 3⅜ to 3¾; depth 4¼ to 5½. D. VI or VII*–11; A. 10 or 11 (8 in one specimen, perhaps abnormal); scales about 60 to 70, too irregular for exact counting. Body moderately elongate, somewhat compressed, tapering posteriorly; head rounded above, its width 2½ in its length; mouth large, oblique, the maxillary reaching to or beyond posterior margin of orbit, 2 to 2⅙ in head; interorbital space wide, 4 to 4½ in head; snout bluntish, broad, a little longer than interorbital width; eye small, 5 in head; teeth present on both jaws, slender, canine-like, arranged in series, the outer row enlarged; caudal peduncle 3 to 3⅓ in head; gill slit about 2½ in head, its upper edge opposite or slightly above uppermost ray of pectoral; scales minute, cycloid, inconspicuous, wanting on head, nape, and fins; shoulder girdle with 2 or 3 small dermal flaps; dorsals separated by a narrow space; dorsal spines very slender; base of spinous dorsal 2⅛ to 2⅜ in head; anal similar to soft dorsal, its base about 1⅓ in head; caudal subtruncate, 1¼ to 1½ in head; ventrals inserted under or slightly behind lower edge of base of pectorals, 1⅞ to 2 in head; pectorals 1½ to 1⅞ in head. Dark olivaceous, mottled with darker; head with some dusky markings; the sides and back with irregular dark markings as in species of *Ethcostominæ;* dorsals distinctly mottled; the first 3 or 4 dorsal spines margined with paler; caudal with faint, broad, wavy cross bars, a faint spot at its base; anal dusky; ventrals yellowish, dusky in males; pectorals plain. Length about 2 inches. Streams of California, in small clear brooks near the sea; locally common in San Luis Obispo Creek, where the specimens here described were taken; probably confined to fresh waters. (Named for Dr. John Strong Newberry of Columbia College, then also on the U. S. Geological Survey.)

Gobius newberryi, GIRARD, Proc. Ac. Nat. Sci. Phila. 1856, 136, Tomales Bay (Coll. E. Samuels); GIRARD, Jour. Bost. Soc. Nat. Hist. 1857, 530, pl. 25, figs. 5 to 8; GIRARD, Pac. R. R. Surv., x, 128, 1858.

Lepidogobius newberryi, JORDAN & GILBERT, Synopsis, 637, 1883; JORDAN & EIGENMANN, *l. c.*, 503.

* Of the nine specimens examined from San Luis Obispo Creek, five have 7 dorsal spines and the other four 6. Girard gives the fin rays as D. VIII–13; A. 12; but we have seen no specimens either with 8 spines or 13 rays. Six specimens from Wadell Creek, Santa Cruz County, California, show the following fin variation: D. VI in 4; D. VII in 1; D. V (?) in 1; D. rays 10 in 4; D. rays 9 in 2; A. 10 in 2; A. 8 in 1; A. 9 in 3.

821. LEPIDOGOBIUS, Gill.

Lepidogobius, GILL, Ann. Lyc. Nat. Hist. N. Y. 1859, 14 (*lepidus*).
Cyclogobius, STEINDACHNER, S. B. K. Ak. Wiss. Wien, XLII, 1860, 284 (*lepidus*).

This genus contains small gobies with the head and body covered with small cycloid scales; dorsal spines 7; inner edge of shoulder girdle with 2 or 3 dermal flaps; interorbital groove with the median ridge of skull little developed; body elongate, subterete; otherwise essentially as in *Gobius*, the skull nearly as in *Gillichthys*, with a median keel and not abruptly widened behind the eye. Pacific Ocean; not entering rivers. (λεπίς, scale; *Gobius*.)

2581. LEPIDOGOBIUS LEPIDUS (Girard).

Head $4\frac{1}{6}$, regularly conical; depth 7; eye 4, equal to snout, twice as long as deep. D. VII–16 to 18; A. 15; scales about 86. Body elongate, subfusiform, little compressed. Snout not obtuse in profile; interorbital space narrow, about equal to diameter of pupil. Mouth rather large, maxillary reaching to below posterior edge of pupil, $2\frac{1}{3}$ in head; teeth small, all similar, those of upper jaw in 2 or 3 series, those of lower jaw close set, in a broad band. Body covered with small cycloid scales which are very much reduced anteriorly, especially on the nape; cheeks, sides of head, and upper posterior part of opercles covered with small scales; top of head scaly to eye; breast scaled. Dorsal spines weak, the longest 2 in head; soft dorsal low, none of the rays reaching caudal; caudal long, somewhat pointed. Color very pale olive, with roundish blotches of rusty red on back and sides; vertical fins mottled with reddish; distal half of all fins and under side of head blackish, especially in the males. "This species is remarkable for numerous lines of papillæ on mandible, snout, and sides of head. The occipital region of the skull is somewhat more depressed than in *Gobius soporator*, and has much lower ridges. A low median carina is present and the low supraorbital ridges are continuous behind the eyes with the temporal crests." (Gilbert MS.) Pacific coast of North America, from Vancouver Island to Lower California; in rather deep water off San Francisco Bay; often seined in great numbers and sold in restaurants as "whitebait." (*lepidus*, pretty.)

Gobius gracilis, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 134, San Francisco; preoccupied by *Gobius gracilis*, JENYNS.
Gobius lepidus, GIRARD, Pac. R. R. Surv., X, 127, pl. 25a, figs. 5 and 6, 1858; substitute for *gracilis;* GÜNTHER, Cat., III, 78, 1861.
Lepidogobius gracilis, GILL, Ann. Lyc. Nat. Hist. N. Y. 1859, 14; JORDAN & GILBERT, Synopsis, 637, 1883; JORDAN & EIGENMANN, *l. c.*, 502.

822. GILLICHTHYS, Cooper.

Gillichthys, COOPER, Proc. Cal. Ac. Sci. 1863, 109 (*mirabilis*).
Gillia, GÜNTHER, Zool. Record 1864, 157 (*mirabilis*); name preoccupied.
Saccostoma (GUICHENOT MS.) SUAVAGE, Bull. Soc. Philom. Paris 1882, 171 (*gulosum*); name preoccupied.

Body moderately elongate, compressed, covered with small, cycloid, embedded scales; belly and head naked. Scales of the young more or

less ciliated. Eyes small, almost superior. Gape wide, the maxillary in the adult inordinately developed, prolonged backward to the base of the pectorals, its posterior part a cartilaginous expansion, connected to an expansion of the skin of the lower jaw, thus forming a channel backward from the mouth, almost exactly as in *Neoclinus* and *Opisthognathus*, genera otherwise very different. Teeth small, even, in broad bands. Skull in adult with a strong median keel, not abruptly widened behind the eye, triangular behind; young with the keel obsolete. Dorsal fins 2, the second high, the first of 6 very weak spines, none of which is exserted; soft dorsal and anal short; caudal less rounded; pectorals large; isthmus broad. Singular little fishes, in brackish waters, burrowing in the mud; confined to the Pacific. (Named for Theodore Gill.)

a. Head moderately depressed; dorsal fins close together. MIRABILIS, 2582.
aa. Head very broad and depressed; distance between dorsals ½ length of first dorsal. DETRUSUS, 2583.

2582. GILLICHTHYS MIRABILIS, Cooper.

(Long-jawed Goby.)

Head 3⅓; depth 5; eye 6 to 7; snout longer than eye, low, little decurved. D. VI–12; A. 10; vertebræ 15+17. Body stout, somewhat compressed behind, broad and depressed anteriorly; head broader than deep, its width 1½, its depth 2 or more in its length; interorbital space greater than eye. Mouth very large; maxillary variable, extending to base of pectoral in adult, broadened behind; fold of lower lip extending its full length. Teeth all alike, small, fixed, and in bands, the band of the lower jaw broader than that of the upper. Scales small, cycloid, irregularly placed, largest from front of dorsal backward, decreasing in size anteriorly; head, breast, belly, and ½ of nape naked. Dorsal spines not filamentous, not as long as the soft rays which are little more than ½ depth of body; caudal broad, short, rounded; pectorals broad and rounded, longer than ventrals, 2 in head. Skull not abruptly widened behind eye, as in *Gobius*, being triangular posteriorly. No lateral ridges; a strong median keel; a short transverse crest behind orbit. Interorbital not deeply grooved, with a blunt median ridge. Orbit not bordered by any prominent ridges. Teeth in both jaws, close set, in bands, all alike. Dull olive, very finely marbled with darker; sides of head and maxillary finely punctuate; fins olive; belly yellowish. Length 8 inches. Pacific coast of North America, from San Francisco to Cerros Island; a most remarkable little fish; very abundant in the mud flats in shallow water along the California coast, burrowing in holes in the mud like a crawfish, and readily taking the hook baited with flesh or worm when dropped into the mouth of the burrow. (*mirabilis*, wonderful.)

Gillichthys mirabilis, Cooper, Proc. Cal. Ac. Sci. 1863, 109, San Diego Bay; Lockington, Amer. Nat. 1879; Jordan & Gilbert, Synopsis, 636, 1883; Jordan & Eigenmann, *l. c.*, 510; Evermann & Jenkins, Proc. U. S. Nat. Mus. 1891, 162.

Gobius townsendi, Eigenmann & Eigenmann, Proc. U. S. Nat. Mus. 1888, 463, San Diego; young.

2583. GILLICHTHYS DETRUSUS, Gilbert & Scofield.

Allied to *Gillichthys mirabilis,* Cooper, differing in the broader and more depressed head, the larger anal fin, and greater distance between the 2 dorsals. Head 3½; depth 5; eye 7 in head; snout 4; interorbital 5½. D. VI–13; A. 11 developed rays (10 in *G. mirabilis*); scales very fine anteriorly but becoming much larger posteriorly; about 75 scales from base of pectoral to caudal, and about 25 longitudinal rows between front of anal and front of second dorsal. The head is depressed, the frontals broad, the shortest distance across being contained in the head 8 times (11 times in *G. mirabilis.*) The postfrontals are small and project but very little, differing from *G. mirabilis,* where the postfrontals project into an elevated wing-like process. The width of the isthmus contained 3 times in the head; maxillary 1⅓ and mandible 1⅔ in head. Least depth of caudal peduncle 2⅔ in head. Distance between dorsals equal to ½ length of first dorsal; length of first dorsal 2¼ in head; second dorsal 1½; anal 2 in head; length of longest pectoral ray 1⅔ in head. Color a very pale olive, some with dark punctulations about the head and fins. The pale coloration is probably due to their life in shallow water on bottom of pale sand. Gulf of California. The types and numerous other specimens, the longest about 5 inches long, were taken by Dr. C. H. Gilbert at Horseshoe Bend, near the mouth of the Colorado River, in Mexico, where they are quite abundant. These are numbered 3836 in L. S. Jr. Univ. Mus. (*detrusus,* depressed.)

Gillichthys detrusus, Gilbert & Scofield, Proc. U. S. Nat. Mus., xx, 1897, 498, pl. 38, Horseshoe Bend, mouth of Colorado River. (Type, No. 48127. Coll. Gilbert & Alexander.)

823. QUIETULA, Jordan & Evermann.

Quietula, Jordan & Evermann, Proc. Cal. Ac. Sci. 1895, 839 (*y-cauda*).

This genus is closely related to *Gillichthys,* from which it differs in the presence of 2 or 3 cutaneous flaps on the inner edge of the shoulder girdle. Maxillary elongate, as in *Gillichthys*; scales rather small, cycloid; cranium essentially as in *Gillichthys.* Small gobies living in the mud of lagoons and river mouths. (A diminutive, from *quies,* quiet.)

2584. QUIETULA Y-CAUDA (Jenkins & Evermann).

Head 3½ (4); depth 7 (8); eye 3½. D. V–14 or 15; A. 15; scales about 50–18; B. 5. Body moderately elongate, compressed, narrowing regularly from shoulder girdle to caudal fin; head not greatly depressed, broader than body, its length 4 in body; snout rounded, short, about equal to diameter of eye; interorbital space narrow, not greater than ½ diameter of eye; mouth rather large, its gape extending nearly to vertical of posterior margin of orbit; maxillary somewhat variable in length, but usually prolonged behind eye for a distance nearly equal to diameter of eye. Scales small, cycloid, about 50 in longitudinal series, 18 in transverse. Teeth in a narrow band on premaxillaries and mandible, short, blunt, and curved slightly backward, most closely set and most numerous

on premaxillaries. Shoulder girdle with 2 or 3 small cutaneous flaps on its inner edge. Fins moderate; dorsal of 5 spines and 16 soft rays, the spines unconnected with the rayed portion, the space between them about equal to ½ diameter of eye; the spines weak and flexible, their length ½ that of head; soft dorsal beginning at a point a little nearer end of snout than tip of caudal and extending nearly to caudal, its height about equal to that of spinous portion, the first few rays slightly graduated; anal having 15 rays and beginning a little behind origin of soft dorsal, the rays about equaling those of dorsal in length; pectorals moderate, inserted a little below axis of the body, their length greater than depth of body, their tips reaching a vertical from posterior part of spinous dorsal; ventrals united, but not adnate to belly, inserted slightly in front of pectorals and their tips not quite reaching those of pectorals. Ground color light; head and body pretty uniformly covered with dark punctulations; an irregular dark bar across occiput; breast and belly pale; a row of 9 or 10 small dark blotches along middle of side, the one at base of caudal plainest and having a shape something like the Greek letter Υ; about 6 dark blotches along median line of back; peritoneum dark. Length about 1⅜ inches. Pacific coast of North America, from Guaymas to Vancouver Island; excessively abundant from San Diego southward in mud flats; specimens recorded from Saanich Arm, Vancouver Island, San Diego, mouth of Colorado River, San Luis Gonzales Bay, St. Georges Bay, Concepcion Bay, Guaymas, and La Paz. It was at first confounded with the young of *Gillichthys mirabilis*, from which genus it differs in the presence of dermal flaps on the shoulder girdle.* (*cauda*, tail, which has a Y-like mark.)

Gillichthys y-cauda, JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 147, Guaymas, Sonora. (Type, No. 39637. Coll. Jenkins & Evermann.)

Quietula y-cauda, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 839.

Gillichthys guaymasiæ,† JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 148, Guaymas, Sonora; young specimens 2⅝ inches long. (Type, No. 39637. Coll. Jenkins & Evermann.)

* "The cranium is similar to that of *Gillichthys mirabilis*, the occiput being depressed, wedge-shaped, narrowed anteriorly with a blunt median carina, the supraorbital and temporal ridges not continuous behind the eye. As in *Gillichthys mirabilis*, the supraorbital ridges end in wing-like expansions immediately behind the interorbital space." (Gilbert MS.)

† *Gillichthys guaymasiæ* is thus described: Head 3 (3⅜ in total); depth 6 (7). D. V–14; A. 13; eye 5. Body quite slender, elongate, but little compressed; head long, narrow, not much widened behind the eyes, not depressed, forming ⅓ the length to base of caudal. Profile gently arched from snout to ½ the distance to dorsal fin, from there nearly straight to dorsal, and then gently curved to caudal peduncle; ventral outline nearly straight; a considerable prominence on the snout made by the enlarged end of the turbinal bone. Eye somewhat above the median line, not quite equaling the snout in length; interorbital space narrow, 1½ times in the eye. The maxillaries are much produced, in some specimens nearly reaching the gill openings, broadest at the middle, and tapering to a blunt point posteriorly; premaxillaries not protractile, but little movable at the symphysis, more than ½ as long as the maxillaries. Gill rakers 2 above the angle, 10 below, short and blunt, the first 4 the largest, those on the second arch but little developed. Teeth well developed, in a single series, on mandible and premaxillaries, all slightly curved backward. Tongue not so broad as in *Gillichthys mirabilis*, Cooper; it is gently rounded at the tip, which is free for a much greater length than in *Gillichthys mirabilis*. Peritoneum black or blackish, and the intestine short, but little longer than the head, and not at all convoluted. Scales small, embedded, and scarcely perceptible except on sides; no pores appear to be developed. First dorsal of fine flexible spines, distance of origin from snout 2⅔ length of body, and separated from the second dorsal by a distance but little greater than length of snout; second dorsal of 14 rays of nearly equal length, which equals the

824. ILYPNUS, Jordan & Evermann.

Ilypnus, JORDAN & EVERMANN, Check-List Fishes, 460, 1896 (*gilberti*).

This genus is allied to *Clevelandia*, from which it differs chiefly in the presence of dermal flaps on the inner edge of the shoulder girdle; scales minute, embedded, cycloid; dorsal with 5 spines; occiput transversely rounded, without median keel; maxillary moderate. Small gobies, inhabiting mud flats. (ἰλύς, mud; ὕπνος, slumber.)

2585. ILYPNUS GILBERTI (Eigenmann & Eigenmann.)

Head 3 to $3\frac{1}{3}$ ($3\frac{2}{3}$ to 4 in total); depth 5 to $5\frac{1}{2}$ (6 to 7). D. V–15 to 17; A. 14 to 16; B. 5; vertebræ 15 + 19. Form elongate, compressed. Head long, subconical, about as high as wide, its width $2\frac{1}{4}$ in its length. Profile nearly straight from eyes to spinous dorsal, decidedly decurved in front of eyes. Eye entirely above the premaxillary level, 1 in snout, $4\frac{1}{2}$ in head, $\frac{1}{2}$ in interorbital. Mouth slightly oblique; maxillary extending to below middle of eye, lower jaw slightly included. Teeth villiform, in a broad band in each jaw, the outer series of lower jaw somewhat enlarged. One, rarely 2, dermal flaps on inner edge of shoulder girdle. Scales cycloid, embedded, very small; head, nape, and breast naked. Distance from tip of snout to insertion of spinous dorsal $2\frac{3}{5}$ in length; highest dorsal spine about $\frac{2}{3}$ length of head; soft dorsal rays lower; interdorsal area about $\frac{1}{2}$ orbital diameter; tip of last dorsal ray not reaching base of caudal; caudal broad and rounded when expanded; anal similar to soft dorsal fin; ventral fins large, nearly reaching vent in specimens $1\frac{3}{4}$ inches long. Pectorals usually shorter than ventrals. Color in life, sand color; head and body with small rust-colored spots, which are dotted with black, the punctulations forming a more or less regular network; dorsal fins hyaline at base, bright rust-colored above, and rather broadly margined with white, everywhere black punctate except on margins; about 3 groups of black dots on each ray, giving a barred appearance to these fins; caudal margined with white, upper and lower parts of fin rust colored, median portion dark gray; about 5 wavy, rustlike, vertical bars; entire fin dotted with black except its margin; anal fin hyaline at base, sparsely dotted, its middle third jet-black, margined with white; pectorals and ventrals milky white, yellowish, sparingly black dotted and white edged; a large, conspicuous, metallic blue-black spot on opercle; top of head blackish; belly white or yellowish; chin and throat white, sometimes

---

distance from end of snout to middle of pupil; length of base of soft dorsal not quite equaling length of head; distance of posterior end from caudal fin equaling distance between the 2 dorsal fins. Origin of anal behind that of soft dorsal and a little posterior to middle of total length of fish; its base $1\frac{1}{4}$ times in base of soft dorsal, or about 4 in length of fish to base of caudal fin; pectorals moderate, a little more than $\frac{1}{2}$ length of head; ventrals inserted slightly behind the pectorals and about equaling them in length. Color in life whitish beneath, grayish or mottled above; 6 double white spots along the back, alternating with fine blackish areas; a white spot behind each eye on top of head; cheek with 2 dark bands extending obliquely backward and downward from eye; a number of dark splotches on opercles; about 7 dusky areas along the side, the last and most marked being upon the base of the caudal fin; dorsal fins finely marked lengthwise by about 4 series of small dark spots; caudal crossed by 5 or 6 wavy vertical bars of very fine dark spots or points; anal, pectorals, and ventrals plain. In alcohol these markings are less plain, especially the white and black areas upon the back. Length $2\frac{3}{4}$ inches.

punctate. Young lighter, showing the reticulations, but the other markings faint or undeveloped. Length about 2¼ inches. (Eigenmann.) "This species agrees with *Lepidogobius* in the presence of papillæ on the inner edge of shoulder girdle. It differs decidedly in the shape of the occipital region of the cranium, which is transversely evenly convex as in *Clevelandia;* not abruptly widened behind the orbits, not continuous laterally with the temporal ridge as in *Gobius, Lepidogobius,* etc. From *Clevelandia* and *Gillichthys, Lepidogobius gilberti* differs in the presence of papillæ on the shoulder girdle, and from *Gillichthys y-cauda* in the shape of the cranium." (Gilbert MS.) San Diego Bay and southward; found by Dr. Gilbert abundant at Magdalena Bay, at Concepcion Bay, and St. Georges Bay, in the Gulf of California. (Named for Charles Henry Gilbert, professor of Zoology in the Leland Stanford Junior University.)

Lepidogobius gilberti, EIGENMANN & EIGENMANN, Proc. U. S. Nat. Mus. 1888, 464, San Diego Bay. (Type, No. 40128, U. S. Nat. Mus. Coll. C. H. Eigenmann.)

825. CLEVELANDIA, Eigenmann & Eigenmann.

Clevelandia, EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 73 (*longipinnis*, EIGENMANN & EIGENMANN, = *rosæ*).

This genus is closely allied to *Gillichthys*, differing chiefly in the form of the skull, which is rounded above, strongly convex in transverse profile, perfectly smooth, without ridges or crests. Body long and slender; maxillary much produced, but not extending to the gill opening; mouth horizontal; dorsal spines 4 or 5, very weak; body covered with minute cycloid embedded scales; soft dorsal and anal long, each of 14 to 17 rays. (Named for Daniel Cleveland, esq., president of the San Diego Society of Natural History, a gentleman deeply interested in scientific matters.)

a. Caudal short, rounded; dorsal spines 5. IOS, 2586.
aa. Caudal pointed, scarcely shorter than head; dorsal spines 4. ROSÆ, 2587.

2586. CLEVELANDIA IOS (Jordan & Gilbert).

Head 3¼ in length of body; depth 6. D. V–16; A. 14; eye 6½ in head; maxillary 1⅓; pectoral 1⅜; ventrals 1⅗; caudal 1⅛; base of soft dorsal 3 in length of body; base of anal 3½. Body long and slender, compressed, the back not elevated; caudal peduncle moderately wide; head long, profile steep to within a short distance of the front of the eye, thence horizontal; mouth very large, not very oblique, the maxillary projecting to opposite the middle of the cheek; jaws subequal; teeth in narrow villiform bands; eye small, longer than wide, set high in head; interorbital space narrow, about as wide as eye. Body covered with very small cycloid scales, too small to count; spinous dorsal well separated from soft dorsal, the spines slender; soft dorsal the higher, its origin a little nearer base of caudal fin than tip of snout; anal about equal to soft dorsal in height, its origin a little behind first dorsal ray, ending at about the same comparative place as soft dorsal; ventrals inserted slightly behind pectorals, reaching midway between their base and front of anal; caudal short, its end rounded.

Color light olivaceous, the cheeks and sides with many dark points which form mottlings; snout dark; a dark spot on upper part of opercle; top of head black; dorsals light, with 3 or 4 dark lines running across the rays; some dark spots on base of anal; pectorals crossed with dark wavy lines; caudal with about 5 irregular cross bars. Puget Sound and neighboring waters. Here described from 2 specimens, each 2 inches in length, dredged off Port Orchard by Mr. Edwin C. Starks. The original description is imperfect and partly incorrect, the single type, from the stomach of *Hexagrammos asper*, being in bad condition. (*Ἰός*, arrow.)

Gobiosoma ios, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 437, Saanich Arm, Vancouver Island (Coll. Jordan & Gilbert); JORDAN & GILBERT, Synopsis, 948, 1883; JORDAN & EIGENMANN, *l. c.*, 509.

Clevelandia ios, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 839, pl. 100.

2587. CLEVELANDIA ROSÆ, Jordan & Evermann.

Head 4 ($4\frac{3}{4}$ in total); depth $6\frac{3}{8}$ (7). D. IV–16; A. 17; scales 70–18. Body very much elongate, slender; head long and slender, depressed anteriorly much as in *Lucius;* profile straight; eye moderate, slightly shorter than snout, $4\frac{1}{3}$ in length of head; interorbital area about as wide as pupil; anteorbital area scarcely $\frac{1}{2}$ diameter of eye; mouth large, maxillary extending much beyond orbit; lower jaw flat, slightly curved upward anteriorly; mouth very much as in *Lucius;* teeth all small, in narrow bands in each jaw; the outer ones of the upper jaw slightly larger than the others. Scales minute, slightly enlarged posteriorly; the margins plain, anterior part of the exposed area lengthwise striated; breast and antedorsal area naked. Distance from snout to insertion of first dorsal spine $2\frac{3}{5}$ in body; the spines slender and short, 3 in head; interdorsal area equals snout and eye; dorsal rays slightly longer than spines, the last ray not extending halfway to caudal; caudal pointed, scarcely shorter than head; ventrals not reaching halfway to vent, $1\frac{2}{3}$ in head; pectoral $1\frac{1}{2}$ in head; vent slightly behind middle of body. Color light brownish; numerous darker spots of aggregated points along nape and upper half of body; belly white; head slightly darker than body; posterior edge of opercle white; an oblique silvery bar on the lower half of opercle, and a light blotch at the upper corner of opercle; cheek with black points; some light areas below eye; lower surface of head and posterior part of maxillaries plain; 2 dark bars on spinous dorsal; second dorsal with 3 or 4 dark bars; a curved black bar at base of caudal; remainder of caudal irregularly barred with dark; other fins plain. Length $1\frac{7}{8}$ inches. San Diego Bay (Eigenmann & Eigenmann); at first incorrectly identified by Mr. and Mrs. Eigenmann with *Evermannia longipinne* (Steindachner), a species similar in habit but wholly scaleless. (Named for Mrs. Rosa Smith Eigenmann.)

Clevelandia longipinnis, EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 73; not *Gobiosoma longipinne*, STEINDACHNER.

Clevelandia rosæ, JORDAN & EVERMANN, Proc. Cal. Ac. Sci. 1896, 229, San Diego. (Coll. R. S. Eigenmann.)

826. EVERMANNIA, Jordan.

Evermannia, JORDAN, Proc. Cal. Ac. Sci., IV, 1895, series 2, 592 (*zosterura*).

Body slender, compressed behind, entirely naked. Head long, slender. Snout rather pointed; mouth moderate, terminal, the maxillary more or less produced backward; teeth small and slender, the outer above slightly enlarged. Skull with a small median crest, not much widened behind. Interorbital space very narrow, channeled; no dermal flaps on shoulder girdle; first dorsal of 4 to 6 spines; second dorsal and anal long, of 14 or 15 rays. Caudal lanceolate. Ventrals formed as in *Gobius* and *Gobiosoma*. Size small, the sexes not colored alike. Species living in holes in sand and mud between tide marks. (Named for "my former student and later scientific associate, Dr. Barton Warren Evermann, now ichthyologist of the United States Fish Commission, in recognition of his work on the fishes of the Gulf of California."—Jordan.)

a. Head $3\frac{2}{3}$; depth $5\frac{1}{4}$; body and fins dotted. LONGIPINNIS, 2588.
aa. Head $3\frac{1}{3}$; depth $6\frac{2}{3}$; vertical fins in males banded with black and with white edgings. ZOSTERURA, 2589.

2588. EVERMANNIA LONGIPINNIS (Steindachner).

Head $3\frac{4}{5}$; depth 5 to $5\frac{1}{3}$. D. IV to VI–16 or 17; A. 16 or 17; snout slightly decurved in profile, $3\frac{1}{2}$ in head; eye 6, greater than interorbital width. Body very slender. Mouth somewhat oblique, the jaws equal; maxillary extending beyond middle of head to a distance behind eye equal to diameter of eye. Teeth in each jaw in 2 series laterally and 3 in front, those of the outer series somewhat enlarged. Fins low, the longest dorsal spine 2 in head; pectoral a little shorter than caudal, scarcely longer than ventrals. Caudal rounded, shorter than head, probably $4\frac{5}{8}$ in body. Body and head completely naked.* Brownish yellow; upper parts of head and body with small, irregularly placed brown spots and streaks; dorsals and caudal finely barred with dark specks. (Steindachner). Gulf of California; not seen by us; known from 3 specimens $37\frac{1}{2}$ mm. long. We refer this species provisionally to *Evermannia*, with which genus it agrees in external respects, although the mouth is much larger. It may be the type of a distinct genus. It differs from *Clevelandia* in the entire absence of scales. (*longus*, long; *pinna*, fin.)

Gobiosoma longipinne, STEINDACHNER, Ich. Beitr., VIII, 27, 1879, Las Animas Bay, Gulf of California; JORDAN & EIGENMANN, *l. c.*, 509.
Evermannia longipinnis, JORDAN, Proc. Cal. Ac. Sci. 1896, 229.

2589. EVERMANNIA ZOSTERURA (Jordan & Gilbert).

Head $3\frac{1}{4}$; depth 6. D. IV–15; A. 14; eye equals snout, 5 in head; P. $1\frac{2}{3}$; C. $1\frac{1}{3}$. Body compressed, profile convex; snout short, not very blunt; eyes high, the maxillary reaching to their posterior margin; mouth oblique, jaws equal; first spine of dorsal filamentous, reaching to middle

* At our request Dr. Steindachner has reexamined the types of this species. He still finds them "vollkommen schuppenlos."

of soft dorsal (male); body entirely naked. Body everywhere speckled with dots of dark brown. Male sometimes with traces of 8 olive cross bands; fins very ornate, the dorsal and anal yellowish at base, then a broad median band of jet black, then a broad white margin; middle of caudal yellow to the tip, with a black band above and below, and a white edge above and below this as in dorsal and anal; no bands on tail. Female with dorsal filament short, reaching about to first soft ray; dorsals and anal checkered with blackish; caudal faintly barred; all vertical fins with pale edgings, but without the black stripe of the males. Length 2 inches. Very common on sandy bottoms, everywhere about the estuary of Mazatlan, the numerous specimens here described being dug out of the sand. It is seldom found much, if any, below the mark of low tide. It is a very handsomely colored species, the male being more strikingly marked than any other of our gobies. (*ζωστήρ*, band; *οὐρά*, tail.)

Gobiosoma zosterurum, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 361, Mazatlan (fin rays incorrect), (Type, No. 29245, U. S. Nat. Mus. Coll. C. H. Gilbert); JORDAN & EIGENMANN, *l. c.*, 509.

Evermannia zosterura, JORDAN, Proc. Cal. Ac. Sci., 2d ser., vol. IV, 1895, 498, pl. 51.

827. GOBIOSOMA, Girard.

(NAKED GOBIES.)

Gobiosoma, GIRARD, Proc. Ac. Nat. Sci. Phila. 1858, 169 (*alepidotus*).

Body entirely naked; mouth moderate, horizontal; snout blunt; teeth in several series, the outer row enlarged; no canines; dorsal spines normally 7, rarely 5 or 6; second dorsal and anal short; no barbels about head; shoulder girdle without flaps. Species chiefly American. (*Gobius: σῶμα*, body.)

a. Coloration olivaceous, mottled with darker; no red nor blue.

b. Maxillary extending to beyond pupil, 4¼ in head; color blackish, with sharply defined cross bars of whitish. Body rather short, the depth 5⅔ in length; head 3¼; snout low, little obtuse; mouth large, rather oblique, the maxillary 2⅔ in head; teeth small, in few series above, in a band below, the outer enlarged; fins low; caudal 1⅔ in head. Cross bands on body as wide as eye, not quite meeting below; a dark blotch on base of pectoral, a fainter one on base of caudal; fins nearly plain. D. VII–13; A. 12. HISTRIO, 2590.

bb. Maxillary extending to below posterior part of orbit; coloration not sharply defined, the body usually with dark cross streaks.

c. Body rather short, chubby, the depth about 4 in length; head about 3⅔; head rounded above; teeth in several series, slender, the outer ones somewhat elongate, none of the inner ones specially enlarged. Color olivaceous, with dark points; sides with narrow, alternating light and dark bars; a row of small linear dark spots along middle of sides; first dorsal with 3 oblique dark bars; second dorsal, caudal, and pectorals finely barred, base and edge of anal light, middle dark; breast with many well-defined spots; a dark line running forward and downward from eye to angle of mouth, another extending straight down; a black bar on edge of preopercle, a black spot on upper edge of opercle. D. VII–13; A. 10. MOLESTUM, 2591.

cc. Body more elongate, depth 5 to 6 in body; head very broad, flattish above, with tumid cheeks, its length 3⅓ in body; eye small, longer than snout, 5 in head; mouth large, little oblique, the jaws subequal.

d. Soft dorsal with 14 rays; no crescent at base of caudal; maxillary extending to below posterior part of orbit (at least in male), 2⅓ in head; teeth in few series, the outer considerably enlarged; 2 teeth on each side of inner series of lower jaw especially large canines; dorsal spines slender, none filamentous; caudal rounded. Olivaceous, with darker cross shades of rounded spots; vertical fins dusky, faintly barred. Teeth of the female similar to those of the male but smaller; head narrower; more slender. D. VII–14; A. 10. BOSCI, 2592.

dd. Soft dorsal with 12 rays; a brown crescent at base of caudal. CRESCENTALE, 2593.

aa. Coloration not plain olivaceous; head with a red bar; anterior dorsal rays not produced in filaments; head and body compressed; greatest depth 5⅔ in total length, head about 4; angle of mouth little behind center of eye; eye 4 in head; teeth pointed, in several series, those of the outer series a little enlarged; caudal rounded. Head light yellow; a carmine-red bar extending along upper edge of head, from upper corner of gill opening to snout, where it joins its fellow, ending behind over the pectoral in a small indigo-blue spot; body with 16 or 17 light green, well-defined cross bars, separated by narrow white stripes; fins chiefly greenish. D. VII (VI)–11 or 12; A. 10. MULTIFASCIATUM, 2594.

2590. GOBIOSOMA HISTRIO, Jordan.

Head 3¼; depth 5⅓. D. VII–12 or 13; A. 11 or 12; maxillary 2¼ in head; caudal 1¼. Body rather short; snout depressed, little obtuse; mouth large, rather oblique, maxillary reaching to below posterior part of orbit; chin without barbels; many series of minute papillæ along mucous pores of head. Teeth small, in few series above, in a band below, the outer enlarged. Fins low. Cross bands on body whitish, as wide as eye, not quite meeting below; a dark blotch on base of pectoral, a fainter one on base of caudal; fins nearly plain. Length 2 inches. Gulf of California; known only from the Gulf of California at Guaymas (Emeric; Evermann & Jenkins) and La Paz (Gilbert). (*histrio*, a harlequin.)

Gobiosoma histrio, JORDAN, Proc. U. S. Nat. Mus. 1884, 260, Guaymas, Mexico (Coll. H. F. Emeric); JORDAN & EIGENMANN, *l. c.*, 508; EVERMANN & JENKINS, Proc. U. S. Nat. Mus. 1891, 162.

2591. GOBIOSOMA MOLESTUM, Girard.

Head about 3⅔; depth 4. D. VII–13; A. 10; vertebræ 12 + 15. Body rather short, maxillary extending to below posterior part of orbit. Teeth in several series, slender, the outer ones somewhat elongate, none of the inner ones specially enlarged. Skull flattish, with a slight median keel; lateral crests developed, lower and stronger than in *Gobius;* interorbital very narrow, bounded by 2 minute crests; bones of the skull very weak and fragile. Teeth in both jaws recurved, in 2 or 3 series. Olivaceous, with dark points; sides with narrow, alternating light and dark bars; a row of small dark spots along middle of side; first dorsal with 3 oblique dark bars; second dorsal, caudal, and pectorals finely barred; base and

edge of anal light, middle dark; breast with many well-defined spots; a dark line running forward and downward from eye to angle of mouth, another extending straight downward from eye; a black bar on edge of preopercle, and a black spot on upper edge of opercle. A specimen taken at Key West is thus described: Pale olive, with darker cross bands formed of dark dots; a row of dark dots along middle of side; vertical fins all mottled and faintly barred with dark olive; pectorals and ventrals nearly plain. Length 2½ inches. Gulf coast of the United States; common in shallow waters along the coast from Key West to Texas and south to Bahia. (*molestus*, disturbed.)

Gobiosoma molestum, GIRARD, Proc. Ac. Nat. Sci. Phila. 1858, 169, Indianola, Texas; GIRARD, U. S. and Mex. Bound. Surv., 27, pl. 12, fig. 14. 1859; GÜNTHER, Cat., III, 556, 1861; JORDAN & GILBERT, Synopsis, 638, 1883; JORDAN & EIGENMANN, *l. c.*, 508.

Gobiosoma alepidotum, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 297, Laguna Grande, Pensacola. (Coll. Dr. Jordan.)

2592. GOBIOSOMA BOSCI (Lacépède).

(NAKED GOBY.)

Head 3⅓; depth 5 to 6. D. VII–14; A. 10; eye 5, longer than snout. Body more elongate; head very broad, flattish above, with tumid cheeks. Eye small. Mouth large, little oblique, jaws subequal, the maxillary extending to below posterior part of orbit (at least in male), 2⅓ in head. Teeth in few series, the outer considerably enlarged; 2 teeth on each side of inner series of lower jaw especially large canines. Dorsal spines slender, not filamentous; caudal rounded. Olivaceous, with darker cross shades of rounded spots; vertical fins dusky, faintly barred. Atlantic coast of the United States, Cape Cod to Florida; generally common, especially southward in shallow grassy bays. (Named for M. Bosc, French consul at Charleston in the last century; an ardent naturalist.)

Gobius bosci, LACÉPÈDE, Hist. Nat. Poiss., II, 555, pl. 16, fig. 1, 1798, Charleston, South Carolina. (Coll. M. Bosc.)

Gobius alepidotus, BLOCH & SCHNEIDER, Syst. Ichthyol., 547, 1801, after LACÉPÈDE; DE KAY, N. Y. Fauna: Fishes, 160, pl. 23, fig. 70, 1842.

Gobius viridipallidus, MITCHILL, Trans. Lit. and Philos. Soc. N. Y., I, 1814, 379, pl. 1, fig. 8, New York.

Gobiosoma bosci, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 613; JORDAN & EIGENMANN, *l. c.*, 508.

Gobiosoma alepidotum, GÜNTHER, Cat., III, 85, 1861; JORDAN & GILBERT, Synopsis, 638, 1883.

2593. GOBIOSOMA CRESCENTALE, Gilbert.

Head 3¾ in length; depth 6¼; eye 5⅓ in head; snout 5⅓. D. VII–12; A. 11. Body very slender, the head depressed, broad and flattened above, the head and body of nearly equal depth throughout. Mouth small, oblique, the maxillary not extending beyond the vertical from posterior border of orbit, 2½ in head; eyes small, 1⅔ in the rather broad interorbital space. Teeth in bands in both jaws, the outer series enlarged, canine-like, and distant. Fins all small, the caudal short and rounded from a broad base, pectoral as long as head without snout; ventrals short, not

reaching ⅔ the distance from their base to vent; dorsal spines not filamentous; skin wholly naked. Color in spirits, lower half of head and body uniform warm brown, the back much lighter, the two areas separated by a well-defined line along middle of sides; this line passing through orbit and through the middle of the base of the pectoral fin; back light grayish, with brownish reticulations, which tend to form 5 or 6 indistinct darker bars uniting with the darker area below the lateral line; a conspicuous brown crescent at base of caudal and pectorals, broad below, narrowing above, margined in front with whitish; anal brown at base; dorsal and caudal with small brown spots forming faint cross series. A single specimen known. Off coast of Lower California. (Gilbert.) (*crescentalis*, pertaining to a crescent.)

Gobiosoma crescentalis, Gilbert, Proc. U. S. Nat. Mus. 1891, 557, off coast of Lower California, at Albatross Station 2825, 24° 22′ N., 110° 19′ 15″ W., in 79 fathoms.

2594. GOBIOSOMA MULTIFASCIATUM, Steindachner.

Head about 4; depth 5⅔. D. VII–12 (Poey), VI–11 (Steindachner); eye 4 in head. Body and head compressed. Angle of mouth little behind center of eye. Teeth pointed, in several series, those of outer series somewhat enlarged. Dorsal rays not filamentous; caudal fin rounded. Head light yellow; a carmine-red bar extending along upper edge of head, from upper corner of gill opening to snout, where it joins its fellow, ending behind over the pectoral in a small indigo-blue spot; body with 16 or 17 light-green, well-defined cross bars, separated by narrow white stripes. (Steindachner.) West Indies; known from Cuba, St. Thomas, and the Lesser Antilles; not seen by us. Its coloration is very different from that of *Gobiosoma*, and it may belong to a distinct genus. (*multus*, many; *fasciatus*, banded.)

Gobius lineatus,* Poey, Memorias, II, 424, 1861, Cuba; name preoccupied by *Gobius lineatus*, Jenyns.

Gobiosoma multifasciatum, Steindachner, Ichth. Beiträge, V, 183, 1870, Lesser Antilles; Jordan & Eigenmann, *l. c.*, 509; Eigenmann & Eigenmann, *l. c.*, 73.

828. BARBULIFER, Eigenmann & Eigenmann.

Barbulifer, Eigenmann & Eigenmann, Proc. Cal. Ac. Sci. 1888, 70 (*papillosus*).

A series of numerous minute barbels around the mouth and chin; otherwise as in *Gobiosoma*; body naked, the dorsal spines 7; second dorsal and anal very short. (*barbula*, a small barbel; *fero*, I bear.)

2595. BARBULIFER CEUTHŒCUS (Jordan & Gilbert).

Head 3⅔; depth 7. D. VII–10; eye 4; A. 10. Body slender; head narrow and slender, depressed; snout not blunt; mouth terminal, oblique, the maxillary reaching to below eye, 3 in head; eyes close together; chin

* *Gobius lineatus* is thus described: Head 3⅓; depth of body 6 in length. D. VII–12; eye 6 in head. Body elongate, subcylindrical, maxillary extending almost to below middle of eye; pectorals rounded; dorsals high. Yellowish green; the body with 20 vertical yellow bands; a red band extending from snout to point of opercle; fins yellowish. Cuba. (Poey.) Type .43 mm. in length.

with a fringe of short barbles; vertical fins high, rays not filamentous. Upper half of head and body brown, finely speckled; 4 oblong, colorless areas along base of dorsals and a smaller one on back of caudal peduncle; lower parts abruptly pale; back with 5 or 6 blackish cross bars reaching to middle of sides, below which they extend as 5 or 6 short V-shaped projections; a brownish streak below eye; a small brown bar on base of pectoral, and a jet black bar at base of caudal. About Key West; scarce. (κεῦθος, a cavity; οἰκέω, to inhabit; the type specimen taken from the cavity of a sponge.)

Gobiosoma ceuthœcum, JORDAN & GILBERT, Proc. U.S.Nat.Mus. 1884, 29, Key West; young (Type in U. S. N. M.); JORDAN & EIGENMANN, *l. c.*, 508.

Barbulifer papillosus,* EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 70, Key West, Florida; adult.

829. TYPHLOGOBIUS, Steindachner.

(BLIND GOBIES.)

Typhlogobius, STEINDACHNER, Ichth. Beiträge, VIII, 24, 1879 (*californiensis*).
Othonops, ROSA SMITH, Proc. U. S. Nat. Mus. 1881, 19 (*eos* = *californiensis*).

Body moderately elongate, compressed, covered with loose, smooth, naked skin. Head large, depressed, with tumid cheeks. Mouth large, the maxillary reaching to beyond the orbit; jaws equal, each with a narrow band of villiform teeth, the outer teeth slightly enlarged; lower jaw capable of little motion; snout rounded; no cirri. Eyes very small, reduced to mere vestages, covered by skin, and functional only in the young. Skull greatly modified, the brain case quadrate. Fins low; first dorsal of 2 flexible spines; second dorsal moderate; anal very short; caudal rounded; ventral disk as in *Gobius*. Gill openings rather narrow. One species known; singular blind gobies, living like slugs under rocks between tide marks. (τυφλός, blind; *Gobius*.)

*This species, which we suppose to be the adult of *Barbulifer ceuthœcus*, is thus described by Dr. Eigenmann: Head 3½ (4⅔ in total); depth 4⅓ (5⅔). D. VII–9; A. 9. Body short and robust, deepest below first dorsal spine; head blunt, profile straight from first dorsal spine to eye, much curved in front of eye; eye longer than snout, 3½ in head; interorbital area ⅔ diameter of eye; snout blunt; mouth small, oblique; maxillary 3 in head, reaching to below anterior margin of pupil; lips thick. About 21 barbels, in length ½ orbital diameter or longer, arranged as follows: A series of 7 cross the snout from one angle of the mouth to the opposite angle, the anterior 3 on the snout rather thick and colored (2 of them nasal), all the others yellowish, the barbel nearest each angle of the mouth longer than any of the others; on the lower jaw a barbel near each rictus, 2 on the chin, behind which are 2 pairs of barbels; posterior to these and below the rictus are 2 barbels on each side; 1 slender barbel on each side of preopercle below the posterior margin of the eye. Numerous rows of pores or papillæ on the head; 1 series extending straight downward on the anterior part of the opercle, from the upper end of which another series extends perpendicularly backward; other pores irregularly scattered on the opercle; a double series extending along edge of preopercle, the pores becoming larger and especially conspicuous below, meeting on the chin; 6 or 7 series radiating from eye, extending to snout, maxillary, and opercular series below; a row of pores nearly surrounding mouth, curving backward, encircling the nasal opening; 1 series about the eye posteriorly, otherwise none on top of head or nape; fins high and rounded; second dorsal higher than first, 1⅓ in head, caudal very broad and rounded, equal to the head in length; anal lower than soft dorsal; ventral reaching ⅔ to vent, 1½ in head; pectoral 1½ in head. Color yellow; upper half of body with a broad band of purplish spots; 6 diamond-shaped spots of darker cross the band, extending above and below it; nape, top of head, and upper part of cheek covered with dark points; opercle light yellow, cheeks darker; an oblique bar of black points on upper half of pectoral base, a curved bar of fainter spots on base of caudal; fins otherwise colorless and transparent. Length 1⅛ inches. (Eigenmann.)

2596. TYPHLOGOBIUS CALIFORNIENSIS, Steindachner.

(Blind Goby of Point Loma; Pink-fish.)

Head 3½; depth 5; eye 6; eye concealed, very small; D. II-12; A. 12. Vertebræ 17 + 13. Body subcylindrical, the males more compressed behind; head very broad behind, its greatest width ⅔ its length. Interorbital space a mere ridge; skin about mouth and eye very loose; a small papilla in front of nasal opening. Lower lip developed as a fold; another fold of skin behind it, bordered with fine cilia; behind this fold is a row of short, thick papillæ; edge of jaw rounded. Spinous dorsal remote from the soft dorsal in the male, but connected with it by a low membrane, this membrane absent in the female; soft dorsal much higher than the spinous; caudal broad, rounded; anal very short, inserted under sixth dorsal ray, and coterminous with dorsal; pectorals little longer than ventrals, 2 in head. Body naked; males with small tubercular plates irregularly placed. Skull highest at its posterior part, depressed forward; the bones all thick and strong. No lateral crests; a median keel which is lowest behind. Orbit not bounded by any ridges. Two keels diverge from the posterior end of the median keel to the insertion of the suprascapula. Premaxillaries and mandible very long. Teeth of the upper jaw all alike, long, close-set, in a broad band, those of the lower jaw in a narrow band, the inner ones apparently larger. Color uniform light pink. Length 2 inches. Coast of Lower California, from San Diego southward to Cerros Island; an extraordinary fish, found attached to the lower side of rocks in shallow water or surf; especially common at Point Loma.

Typhlogobius californiensis, Steindachner, Ichth. Beiträge, viii, 24, 1879, False Bay, San Diego, California (Coll. Prof. Essmark); Jordan & Gilbert, Synopsis, 639, 1883; Jordan & Eigenmann, *l. c.*, 511.

Othonops eos, Rosa Smith, Proc. U. S. Nat. Mus. 1881, 53, Point Loma, California.

830. TYNTLASTES, Günther.

Tyntlastes, Günther, Proc. Zool. Soc. London 1862, 193 (*sagitta*).

Body elongate, compressed, covered with small, imbricate, cycloid scales. Head elongate, quadrangular. Mouth wide, oblique, the lower jaw projecting; teeth small, in single series, none on vomer or palatines. Eyes very small, or rudimentary. Dorsal fin single, continuous, about 6 of its anterior rays simple; caudal fin pointed, more or less joined to the dorsal and anal; ventral fins united. Air bladder very small or absent. No pseudobranchiæ. Vertebræ 11 + 20. Pacific Ocean. (τυντλαίστης, a mud-dabbler.)

a. Dorsal and anal each with 15 soft rays; head 4½ in length. BREVIS, 2597.
aa. Soft dorsal and anal each with 21 unbranched or soft rays; head 5½ in length. SAGITTA, 2598.

2597. TYNTLASTES BREVIS (Günther).

Head 4½; depth 8. D. VI, 15;* A. 15. Eyes minute. Jaws each with a

* The dorsal formula is apparently VIII, 14 in 2 half-digested specimens taken from the stomach of a *Centropomus* at Panama. (Gilbert.)

series of wide-set teeth. Caudal fin black. (Günther.) Panama; not seen by us. (*brevis*, short.)

Amblyopus brevis, GÜNTHER, Proc. Zool. Soc. London 1864, 151, Panama; GÜNTHER, Fishes Centr. Amer., 441, 1869.
Tyntlastes brevis JORDAN & EIGENMANN, *l. c.*, 512.

2598. TYNTLASTES SAGITTA (Günther).

Head $5\frac{2}{3}$; depth $9\frac{2}{3}$. D. VI, 21; A. 21. Body and head elongate, compressed. Maxillary reaching to behind eye; teeth subhorizontal, very small. Scales becoming larger posteriorly. Caudal arrow-shaped, about 4 in body; pectorals as long as ventrals, 2 in head. Grayish, sides and under parts silvery; an ovate gray spot before each dorsal ray; caudal grayish. (Günther.) Length $9\frac{1}{2}$ inches. Coast of Lower California; exact locality unknown. (*sagitta*, arrow.)

Amblyopus sagitta, GÜNTHER, Proc. Zool. Soc. London 1862, 193, "California," probably from Lower California.
Tyntlastes sagitta, JORDAN & GILBERT, Synopsis, 639, 1883; JORDAN & EIGENMANN, *l. c.*, 512.

831. GOBIOIDES, Lacépède.

(BARRETOS.)

Gobioides, LACÉPÈDE, Hist. Nat. Poiss., II, 580, 1798 (*broussonnetii*).
Plecopodus, RAFINESQUE, Analyse de la Nature, 87, 1815 (*broussonnetii*); substitute for *Gobioides*, regarded as objectionable.
Ognichodes, SWAINSON, Nat. Hist. Class'n Animals, II, 183 and 278, 1839 (*broussonnetii*).

Body greatly elongate, compressed behind, the scales very minute; head small; eyes very small; mouth large, oblique, the lower jaw projecting; gill openings moderate. Teeth in a band, those in the outer series being very strong. Dorsal rays V to VII, 15 to 23; anal rays 16 to 23. Dorsal fin low, continuous, the spines similar to the soft rays, but more widely separated; the soft dorsal and the anal are joined to base of caudal; ventrals 45, united in a disk which is formed much as in *Gobius*. No air bladder; no pseudobranchiæ. From *Tæniοides* (=*Amblyopus*) the genus *Gobioides* is distinguished by the absence of barbels, the presence of scales, and by the much smaller number of rays in its vertical fins. Brackish waters of the Tropics, reaching a considerable size. (*Gobius*; εἶδος, resemblance.)

a. Eye small, but evident; scales evident, larger behind. BROUSSONNETII, 2599.
aa. Eye minute, not evident; scales minute. PERUANUS, 2600.

2599. GOBIOIDES BROUSSONNETII, Lacépède.

Head $5\frac{1}{4}$ (young) to 7 (adult); caudal $3\frac{1}{2}$ to 5; eye small but evident, 7 to 10 in head; interorbital space 1 to $1\frac{2}{3}$ diameter of eye. D. VII, 16; A. I, 16. Body elongate, mouth oblique, maxillary extending beyond eye; teeth in bands, the outer series enlarged, shorter, and closer set than in *Gobioides peruanus;* scales twice as large as in *peruanus*, those on anterior part of body not imbricated, much smaller than those on posterior part,

which are elongate oval in form. Violet bars extending downward and forward on the upper part of body; sometimes a violet spot with a lighter or darker dot at end of the bars; head marbled or spotted with dark violet or brown. (Steindachner.) Length 20 inches or more. West Indies to Brazil; common southward, ascending rivers; once taken near New Orleans (Bean & Bean). (Named for Dr. Augustin Broussonnet, professor in the University of Montpelier.)

Gobioides broussonnetii, LACÉPÈDE, Hist. Nat. Poiss., II, 580, 1798, probably from Surinam, "given by Holland to France."

Amblyopus brasiliensis, BLOCH & SCHNEIDER, Syst. Ichth., 69, 1801, Brazil; on drawing made by Prince Maurice; CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 121, 1837.

Gobious oblongus, BLOCH & SCHNEIDER, Syst. Ichth., 548, 1801; based on LACÉPÈDE.

Gobioides barreto, POEY, Memorias, II, 282, 1861, Cuba; POEY, Synopsis, 394, 1868; POEY, Enumeratio, 125, 1876.

Amblyopus mexicanus, O'SHAUGHNESSY, Ann. Mag. Nat. Hist., series IV, vol. XV, 1875, 147, Mexico.*

Gobioides broussoneti, JORDAN & EIGENMANN, *l. c.*, 512; BEAN & BEAN, † Proc. U. S. Nat. Mus. 1895, 631.

2600. GOBIOIDES PERUANUS (Steindachner).

Head 5; depth 11. D. VII, 17; A. I, 16. Eye scarcely visible, much smaller than in *G. broussoneti;* scales very minute; snout $2\frac{1}{2}$ in postorbital part of head; interorbital 5 in head; lower jaw slightly projecting; maxillary $2\frac{3}{5}$ in head; a series of large slender teeth in each jaw, behind which, in each jaw, is a narrow band of fine teeth; caudal $4\frac{1}{2}$ in

* The following is Mr. O'Shaughnessy's description of *Amblyopus mexicanus:* D. VII, 15; A. I, 15. Depth 13 in total length. Body covered all over with scale-shaped crypts. Head naked. Dorsal $\frac{3}{4}$ height of body. Eye small, but distinct. Snout obtuse; lower jaw extending a little beyond upper. Teeth small, close set, the outer series much smaller and more closely set than in *G. broussonnetii*. Dorsal and anal connected with the caudal. Upper parts dark brown, with a series of white spots along the whole length of the side; lower parts of sides and belly white. One specimen in the British Museum, from Mexico, purchased. Length $20\frac{1}{4}$ inches. (O'Shaughnessy.)

This seems to differ from *G. broussonnetii* in color only.

† The following description is given by Bean & Bean of *Gobioides broussonnetii* (Lacépède): Head 7; depth 14. D. VI, 17; A. I, 16. The greatest depth of the head equals the length of the upper jaw, or about $\frac{1}{3}$ the length of head without snout. The body is compressed. Its greatest thickness is contained $1\frac{2}{3}$ times in its greatest depth. The teeth are in narrow bands in each jaw, some of those in the outer row enlarged, canine-like, and curved inward. All of the teeth are more or less curved inward and depressible. The vomer and palate are toothless. The mouth is oblique, the lower jaw projecting slightly beyond the upper. The maxilla extends well behind the eye, its length is slightly more than $\frac{1}{3}$ that of head without the snout. It is not much expanded posteriorly. Eyes very small, their diameter equaling $\frac{1}{2}$ length of snout, about equal to width of interorbital space. The snout scarcely equals more than $\frac{1}{5}$ of the head's length. Gill openings wide, the membranes wholly joined to the isthmus. Branchiostegals much curved, 4 in number. The dorsal begins at a distance from the nape equal to the postorbital part of the head, the origin being about over the end of the extended pectoral. The ventral reaches farther back than the pectoral, and is longer than that fin, its length equaling postorbital part of head. The distance of the vent from the tip of the snout equals somewhat more than 3 times the length of the head; it is under the interspace between the last spine and first ray of the dorsal, with a small genital papilla behind it. The caudal is very long and tapering, $1\frac{2}{3}$ times as long as the head. The dorsal spines are long and slender, the fifth nearly as long as the postorbital part of the head. The second dorsal ray is slightly longer. The anal rays are about as long as those of the dorsal. The scales are thin, not imbricated, except on the posterior part of the head, where they are long and elliptical in shape. The head and breast are naked. The colors have faded out in alcohol; the ground color appears to have been light brown, with darker blotches on the median line of the body under the spinous portion of the dorsal and the anterior part of the soft dorsal. (Bean & Bean.) Here described from a specimen obtained in the Gulf of Mexico by Mr. Robert S. Day, of New Orleans, Louisiana, and is No. 38220, U. S. Nat. Mus.

body, connected by membrane to dorsal and anal; sides with regular cross series of pores. Body with narrow angular cross bars; dorsal rays violet, the membrane yellowish. (Steindachner.) Shores of Ecuador and Peru, ascending rivers.

Amblyopus peruanus, STEINDACHNER, Fisch-Fauna des Cauca und Flüsse bei Guayaquil, 42, 1880, Guayaquil.
Gobioides peruanus, EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci., 2d ser., I, 1888, 75.

832. CAYENNIA, Sauvage.

Cayennia, SAUVAGE, Bull. Sci. Philom., ser. 7, IV, 1880, 57 (*guichenoti*).

Body much elongate; dorsals united, caudal free from dorsal and anal; ventrals united, not adhering to belly; teeth small, the outer enlarged; anterior part of body naked, posterior part covered with cycloid scales. Otherwise as in *Gobioides*, from which the genus may not be separable. (Name from Cayenne.)

2601. CAYENNIA GUICHENOTI, Sauvage.

Head 9; depth 17. D. VI, 17; A. I, 16; vertebræ about 36. Head deeper than wide; eye small, placed well forward; maxillary reaching to below posterior margin of eye; a low membrane connecting dorsal and caudal; caudal 7 in length; ventrals 1⅓ in head. Color brownish, marbled with black anteriorly. Cayenne (Sauvage); not seen by us. (Named for A. Guichenot, formerly ichthyologist of the Muséum d'Histoire Naturelle at Paris.)

Cayennia guichenoti, SAUVAGE, Bull. Soc. Philom., ser. 7, IV, 1880, 57 Cayenne; EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1888, 76.

Suborder DISCOCEPHALI.

Bony fishes "with a suctorial transversely laminated oval disk on the upper surface of the head (homologous with a flat dorsal fin), thoracic ventral fins with external spines, a simple basis cranii, intermaxillary bones flattened, with the ascending processes deflected sideways, and with the supramaxillary bones attenuated backward, flattened, and appressed to the dorsal surface of the intermaxillaries; hypercoracoid (or scapula) perforated nearly in the center, and with 4 short actinosts (carpals)." (Gill.)

This remarkable group consists of a single family, *Echeneididæ*. (δίσκος, disk; κεφαλή, head).

Family CLXXXIX. ECHENEIDIDÆ.

(THE REMORAS.)

Body fusiform, elongate, covered with minute, cycloid scales. Mouth wide, with villiform teeth on jaws, vomer, palatines, and usually on tongue. Premaxillaries not protractile. Lower jaw projecting beyond upper. Spinous dorsal modified into a sucking disk, which is placed on

the top of the head and neck, and is composed of a double series of transverse, movable, cartilaginous plates, serrated on their posterior or free edges. By means of this disk these fishes attach themselves to other fishes or to floating objects, and are carried for great distances in the sea. Opercles unarmed. Pectoral fins placed high; ventral fins present, thoracic and close together, I, 5; dorsal and anal fins long, without spines, opposite each other; caudal fin emarginate or rounded. Branchiostegals 7. Gills 4, a slit behind the fourth; gill rakers short; gill membranes not united, free from the isthmus. Pseudobranchiæ obsolete. Several pyloric appendages. No air bladder. No finlets. No caudal keel. Vertebræ more than 10 + 14. Genera 4; species about 10, found in all seas, all having a very wide range. The species of this group are apparently descended from a fossil genus, *Opisthomyzon*,* Cope (*glaronensis*), characterized by the small posterior disk and slender body.

The following description of this family is given by Dr. Gill: Body elongated, subcylindrical, diminishing backward gradually from the head and into the slender caudal peduncle. Anus subcentral. Scales cycloid, very small, and not, or scarcely, imbricated. Lateral line nearly straight and very faint. Head above oblong and with a flattened straight upper surface, furnished with an adhesive oblong or elongated, laminated disk. The eyes are rather small, submedian, and overhung by the disk. Suborbital bones forming a slender infraorbital chain; the first or preorbital triangular and thick. Opercular apparatus normally developed and unarmed. Nostrils double, close together. Mouth terminal or, rather, superior, the lower jaw projecting, but with the cleft nearly horizontal and not extending laterally to the eyes. Teeth present on the jaws and palate. Branchial apertures ample and fissured forward. Branchiostegal rays 7 (or 8) on each side. The adhesive disk on the upper surface of the head is a modified first dorsal fin, and from the snout generally extends more or less posteriorly on the nape and back; it is oblong or elongated and of an oval or elliptical form, divided into equal halves by a longitudinal septum, and with more or less numerous transverse laminæ in each division, the laminæ being slightly erectile and depressible. Dorsal fin oblong or elongated on the posterior half of the body (including head), ending some distance from the caudal. Anal fin opposite and similar to the dorsal. Caudal fin rather small, variable in outline, but never deeply forked. Pectoral fins moderate, inserted high on the sides. Ventral fins thoracic, each with a spine and 5 branched rays. The vertebral column has vertebræ in slightly increased numbers, the abdominal vertebræ being about 12 to 14 and the caudal 15 or 16. The stomach is cæcal and the pyloric cæca are present in moderate numbers. The air bladder is obsolete.

* "A careful comparison of the proportions of all the parts of the skeleton of the fossil *Echeneis* with those of the living forms, such as *Echeneis naucrates* or *Echeneis remora*, shows that the fossil differs nearly equally from both, and that it was a more normally shaped fish than either of these forms. The head was narrower and less flattened, the preoperculum wider, but its two jaws had nearly the same length. The ribs, as also the neural and hæmal spines, were longer, the tail more forked, and the soft dorsal fin much longer. In fact, it was a more compressed type, probably a far better swimmer than its living congeners, as might be expected, if the smallness of the adhesive disk is taken into account." (Storms.) This form (*Echeneis glaronensis*, Wellstein) is made the type of the genus *Opisthomyzon*, Cope, the name referring to the posterior portion of the small disk. The vertebræ in *Opisthomyzon* are 10 + 13 = 23.

Concerning the relations of this family, Dr. Gill has the following pertinent remarks:

"The family of *Scombéroïdes* was constituted by Cuvier for certain forms of known organization, among which were fishes evidently related to *Caranx*, but which had free dorsal spines. In the absence of knowledge of its structure, the genus *Elacate* was approximated to such because it also had free dorsal spines. Dr. Günther conceived the idea of disintegrating this family, because, inter alias, the typical *Scomberoides* (family *Scombridæ*) had more than 24 vertebræ and others (family *Carangidæ*) had just 24. The assumption of Cuvier as to the relationship of *Elacate* was repeated, but inasmuch as it has 'more than 24 vertebræ' (it has $25 = 12 + 13$) it was severed from the free-spined *Carangidæ* and associated with the *Scombridæ*. *Elacate* has an elongated body, flattish head, and a colored longitudinal lateral band; *Echeneis* has also an elongated body, flattened head, and a longitudinal lateral band; therefore *Echeneis* was considered to be next allied to *Elacate* and to belong to the same family. The very numerous differences in structure between the two were entirely ignored, and the reference of the *Echeneis* to the *Scombridæ* is simply due to assumption piled on assumption. The collocation need not, therefore, longer detain us. The possession by *Echeneis* of the anterior oval cephalic disk in place of a spinous dorsal fin would alone necessitate the isolation of the genus as a peculiar family. But that difference is associated with almost innumerable other peculiarities of the skeleton and other parts, and in a logical system it must be removed far from the *Scombridæ*, and probably be endowed with subordinal distinction. In all essential respects it departs greatly from the type of structure manifested in the *Scombridæ* and rather approximates—but very distantly—the *Gobioidea* and *Blennioidea*. In those types we have in some a tendency to flattening of the head, of anterior development of the dorsal fin, a simple basis cranii, etc. Nevertheless, there is no close affinity nor even any tendency to the extreme modification of the spinous dorsal exhibited by *Echeneis*. In view of all these facts *Echeneis*, with its subdivisions, may be regarded as constituting not only a family but a suborder. * * * Who can consistently object to the proposition to segregate the *Echeneididæ* as a suborder of teleocephaleous fishes? Not those who consider that the development of 3 or 4 inarticulate rays (or even less) in the front of the dorsal fin is sufficient to ordinarily differentiate a given form from another with only 1 or 2 such. Certainly the difference between the constituents of a disk and any rays or spines is much greater than the mere development or atrophy of articulations. Not those who consider that the manner of depression of spines, whether directly over the following, or to the right or left alternately, are of ordinal importance; for such differences again are manifestly of less morphological significance than the factors of a suctorial disk. Nevertheless, there are doubtless many who will passively resist the proposition because of a conservative spirit, and who will vaguely recur to the development of the disk as being a 'teleological modification,' and as if it were not an actual fact and a development correlated with radical modifications of all parts of the skeleton at least. But whatever may be the closest relations of *Echeneis*, or the systematic

value of its peculiarities, it is certain that it is not allied to *Elacate* any more than to others of the hosts of Scombroid, Percoid, and kindred fishes, and that it differs in toto from it, notwithstanding the claims that have been made otherwise. It is true that there is a striking resemblance, especially between the young—almost as great, for example, as that between the placental mouse and the marsupial *Antechinomys*—but the like is entirely superficial, and the scientific ichthyologist should be no more misled in the case than would the scientific therologist by the likeness of the marsupial and placental mammals."

a. Body very slender, the vertebræ 14 + 16 = 30; ventrals narrowly adnate to abdomen; lower jaw produced in a flap; pectorals acute, with flexible rays.
  b. Laminæ 10 only. PHTHEIRICHTHYS, 833.
  bb. Laminæ 20 to 28. ECHENEIS, 834.
aa. Body rather robust, the vertebræ 12 + 15=27; ventrals broadly adnate to abdomen; lower jaw not produced; pectorals rounded.
  c. Laminæ 24 to 27. REMILEGIA, 835.
  cc. Laminæ 16 to 20.
    d. Pectoral rays soft and flexible. REMORA, 836.
    dd. Pectoral rays stiff and ossified. RHOMBOCHIRUS, 837.

833. PHTHEIRICHTHYS, Gill.

Phtheirichthys, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 239 (*lineata*).

Disk with 10 laminæ; palatines with sharp teeth; teeth in pairs, uniform in all ages; otherwise as in *Echeneis*. A single species, found attached to spearfishes and Barracudas. (*ψθείρ*, a louse; *ἰχθύς*, fish.)

2602. PHTHEIRICHTHYS LINEATUS (Menzies).

Head 5; disk twice as long as broad, its length 4⅓ in body. D. X–33; A. 33. Lower jaw very narrow, much projecting. Body blackish, with 2 whitish lateral bands; all the fins white-margined. Tropical seas, ranging north to South Carolina and Pensacola; rather rare. (*lineatus*, striped.)

Echeneis lineata, MENZIES, Trans. Linn. Soc. London, I, 1791, 187, pl. 17, fig. 1, Pacific Ocean between the tropics; GÜNTHER, Cat., II, 382, 1860.
Echeneis tropica, EUPHRASEN, Nya Handl., XII, 317, 1791, Atlantic between the Tropics.
Echeneis apicalis, POEY, Memorias, II, 254, 1861, Cuba. (Coll. Poey.)
Echeneis sphyrænarum, POEY, Memorias, II, 255, 1861, Cuba, on Barracudas. (Coll. Poey.)
Phtheirichthys lineatus, JORDAN & GILBERT, Synopsis, 969, 1883.

834. ECHENEIS (Artedi) Linnæus.

Echeneis (ARTEDI) LINNÆUS, Syst. Nat., Ed. X, 260, 1758 (*naucrates*).
Leptecheneis, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 60 (*naucrates*); the name *Echeneis* being transferred to *E. remora*, the only species known to Artedi.

Body comparatively elongate, the vertebræ 14 + 16 = 30; disk long, of 20 to 28 laminæ; pectoral pointed, its rays soft and flexible; soft dorsal and anal long, of 30 to 41 rays each; caudal lunate in the adult, convex in the young. Species of wide distribution, attaching themselves mainly to sea turtles and large fishes. (*ἐχενηΐς*, an ancient name, from *ἔχω*, to hold back; *ναύς*, a ship.)

a. Disk of 22 to 26 laminæ (rarely 21 or 28), its length less than ¼ body. NAUCRATES, 2603.

aa. Disk of 20 or 21 laminæ, its length more than ¼ body. NAUCRATEOIDES, 2604.

2603. ECHENEIS NAUCRATES, Linnæus.

(Shark-sucker; Pega; Pegador; Sucking-fish.)

Head 5¼; depth 11 to 12. D. XXII to XXVIII (rarely XXI)–32 to 41; A. 31 to 38. Breadth between pectorals 7½; disk 4 to 5 in body; eye 5 in head; snout 2⅕; maxillary 3; from angle of mouth to tip of lower jaw 2⅜; pectoral 1⅖; ventrals 1½; middle caudal rays 1⅘; highest anal ray 2; highest dorsal ray 2⅕; width of disk 2½ in its length; base of dorsal 2½, anal 2½, in body. Body elongate, subterete, slender. Lower jaw strongly projecting, the tip flexible; maxillary reaching nostril; teeth uniform in the adult, the young with series of small slender teeth in advance of the others; gill rakers short and slender, about equal to pupil; vertical fins low. Anal rays higher than dorsal anteriorly; pectorals reaching very slightly past tips of ventrals; origin of ventral spine under middle of pectoral base; inner rays of ventral fins narrowly adnate to the abdomen; dorsal and anal commencing and ending opposite each other; caudal with the middle rays produced in the young, the fin becoming emarginate or lunate with age. Color brownish; belly dark, like the back, as usual in this family; sides with a broad stripe of darker edged with whitish extending through eye to snout; caudal black, its outer angles whitish; pectorals and ventrals black, sometimes bordered with pale; dorsal and anal broadly edged with white anteriorly; adult nearly uniform dark brown, not paler below. Warm seas, universally distributed; common north to Cape Cod and occasionally to San Francisco, attaching itself to turtles and to large fishes. This species is very common in the tropics, being found attached to sharks, groupers, or any other large fish, without regard to species. Few large sharks at Key West are without them. They are often caught with hook and line from the wharf, where they frequently forsake their host to take the bait. Lütken's remark that only *Remora remora* has been recorded from sharks is no longer true. Several writers have recognized 2 species of *Echeneis* proper—*naucrates*, with 22 to 26 laminæ, the disk 4 to 5 in body, and *naucrateoides* (= *albicauda* = *holbrooki* = *lineatus*), in which the disk is longer, 3⅗ to 4 in body, but composed of fewer, 20 or 21, laminæ. The latter form is rather common on our coast, the specimens from Key West above mentioned having 21. We doubt the existence of any permanent difference between the two, but provisionally retain *Echeneis naucrateoides* as a species distinct from *Echeneis naucrates* until more complete comparison can be made. (*naucrates*, a pilot; *ναῦς*, ship; *κρατέω*, to govern, guide.)

Echeneis neucrates (misprint for *naucrates*), Linnæus, Syst. Nat., Ed. x, 261, 1758, "in Pelago Indico;" Günther, Cat., II, 384, 1860; Jordan & Gilbert, Synopsis, 416, 1883.
Echeneis albicauda, Mitchill, Amer. Monthly Mag., II, 1817, 244, New York.
Echeneis lunata, Bancroft, Proc. Comm. Zool. Soc., I, 1830, 134, Kingston, Jamaica.
? *Echeneis vittata*, Lowe, Proc. Zool. Soc. Lond. 1839, 89, Madeira.
Echeneis fasciata, Gronow, Ed. Gray, 92, 1854, Mediterranean Sea.
Leptecheneis naucrates, Gill, Proc. Ac. Nat. Sci. Phila. 1864, 60.

Echeneis vittata, RÜPPELL, Neue Wirb. Fische, 82, 1835, Red Sea.
Echeneis guaiacan, POEY, Memorias, II, 248, 1861, Cuba; young. (Coll. Poey.)
Echeneis verticalis, POEY, Memorias, II, 253, 1861, Cuba; young.
Echeneis metallica, POEY, Memorias, II, 252, 1861, Cuba; D. XXIII, 40; A. 37; large specimen, metallic green, the bands faint. (Coll. Poey.)
Echeneis fusca, GRONOW, Cat. Fish., 92, 1854; after *E. naucrates*, L.

2604. ECHENEIS NAUCRATEOIDES, Zuiew.

Head 5; depth 11. D. XX or XXI–32 to 35; A. 33 to 35. Disk $3\frac{3}{8}$ to $3\frac{4}{5}$ in total, twice width of body between pectorals. In all other respects essentially as in *Echeneis naucrates*, the disk longer, but composed of fewer laminæ, the laminæ being farther apart. Color of *Echeneis naucrates*. Cape Cod to West Indies, common on our south Atlantic coast; specimens before us from Key West. (*naucrates*, *ναυκράτης*, a pilot; *εἶδος*, resemblance.)

Echeneis neucratoides, ZUIEW, Nova Acta Acad. Sci. Imp. Petropol., IV, 1789, 279, no locality.
Echeneis lineata, HOLBROOK, Ichth. S. C., 102, 1860, Charleston, South Carolina; not of MENZIES.
Echeneis holbrooki, GÜNTHER, Cat., II, 382, 1860, Jamaica; D. XIII, 35; A. 33.
Leptecheneis naucrateoides, GILL, *l. c.*, 61.

835. REMILEGIA, Gill.

Remilegia, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 61 (*australis*).

This genus differs from *Remora* chiefly in the length of the sucking disk, which has 24 to 27 laminæ; the soft dorsal and anal are proportionately short. (A metathesis for *remeligo*, the delayer or hinderer.)

2605. REMILEGIA AUSTRALIS (Bennett).

D. XXVII–22; A. 21 to 23. The length of the disk is $2\frac{1}{3}$ in the total, the width of the body between the pectorals $5\frac{3}{5}$. Caudal truncated; dorsal and anal fins not continued to the caudal. Color brown. This species has the general habit of *E. remora*, but may be readily distinguished from all the others by the extraordinary size of the disk, which is elongate, subelliptical, obtusely rounded anteriorly and posteriorly, and formed by 27 pairs of laminæ; it extends backward beyond the vertical from the tip of the ventrals, and its length is $2\frac{1}{8}$ in the total. The spines with which the single lamina are armed are less conspicuous than in the other species, and do not offer the same resistance to the touch. There is a large posterior portion of the disk which is not provided with laminæ, but quite smooth. The width of the disk, taken between the extremities of the bony laminæ, is $\frac{1}{3}$ of its length; the membranaceous margin is bent upward. The head and the body below the disk are depressed, and their height is $9\frac{1}{2}$ in the total length, whilst the width between the pectorals is $5\frac{3}{4}$ in it. The body between the disk and the vertical fins is quadrangular, tapering posteriorly. The upper jaw is subtruncated, and overreached by the lower, which is much narrower; both are armed with a broad band of villiform teeth, and with an outer series of larger ones on

the sides; the vomerine and palatine bones have a continuous band of teeth, narrowest on the vomer; the tongue is hard, cartilaginous, and destitute of teeth. The cleft of the mouth reaches only to the vertical from the nostril; the eye is small. The pectoral is rounded and small, its length being $\frac{1}{9}$ of the total; the ventrals are slightly pointed, and, as in all the species of the genus, composed of 1 spine, hidden in the skin, and 4 soft rays; they are inserted immediately behind the vertical from the pectoral, which they equal in length; they can be received in a shallow groove on the abdomen. The distance between the dorsal and the disk is $3\frac{1}{2}$ in the length of the latter; the dorsal is low, and enveloped in a thick membrane. The caudal is truncated when stretched out. The anal is very similar to the dorsal, and its origin and termination fall vertically below those of the latter. The scales are minute, and can be perceived only by the aid of a magnifier; they are embedded in pore-like cavities. (Günther: description of type of *Echeneis scutata*.) Tropical seas; rare; recorded by Dr. Lütken from 10° N., 39° W. (Coll. Capt. V. Hygom) from a dolphin; not seen by us. (*australis*, southern.)

Echeneis australis, BENNETT, Narr. Whaling Voyage, II, 273, pls. 24–26, 1840.
Echeneis scutata, GÜNTHER, Ann. Mag. Nat. Hist. 1860, 401, pl. 10, f. B, Ceylon (Coll. Dr. Sibbald); GÜNTHER, Cat. Fish., II, 381, 1860; LÜTKEN, Vid. Medd. Kjöbenh. 1875, 42.

836. REMORA, Gill.

(REMORAS.)

Remora, GILL, Proc. Ac. Nat. Sci. Phila. 1862. 239 (*remora*).
Echeneis, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 60 (*remora*); not *Echeneis*, GILL, 1862, restricted to *naucrates*.
Remoropsis, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 60 (*brachypterus*).
Remorina, JORDAN & EVERMANN, Check-List Fishes, 490, 1896 (*albesens*).

Body rather robust, the vertebræ $12 + 15 = 27$; disk shortish, of 13 to 18 laminæ; pectoral rounded, its rays soft and flexible; soft dorsal and anal moderate, of 20 to 30 rays; caudal subtruncate. Species attaching themselves to large fishes, especially to sharks. (*Remora*, an ancient name, "holding back.")

REMORA:
a. Laminæ about 18; soft dorsal with 23 rays. — REMORA, 2606.
aa. Laminæ 13 to 16.
REMORINA:
b. Dorsal rays XIII, 22. — ALBESCENS, 2607.
REMOROPSIS (*Remora*; ὄψις, appearance):
bb. Dorsal rays XIV, XVI, 29 to 32. — BRACHYPTERA, 2608.

Subgenus REMORA.

2606. REMORA REMORA (Linnæus).

(REMORA.)

Head 4; disk $2\frac{3}{4}$; width between pectorals $5\frac{1}{4}$. D. XVIII–23; A. 25; vertebræ $12 + 15$. Body comparatively robust, compressed behind. Pectoral fins rounded, short, and broad, their rays short and flexible; ventral fins

adnate to the abdomen for more than ½ the length of their inner edge. Tip of lower jaw not produced into a flap; head broad, depressed; disk longer than the dorsal or the anal fin; maxillary scarcely reaching front of orbit. Caudal lunate; vertical fins rather high; pectoral ⅔ length of head. Color blackish, nearly uniform above and below. Length 15 inches. Warm seas, north to New York and San Francisco, where it is not rare; usually found attached to large sharks; very common in the West Indies; more robust than *Echeneis naucrates*, and reaching a smaller size.

Echeneis remora, LINNÆUS, Syst. Nat., Ed. x, 260, 1758, "in Pelago Indico;" GÜNTHER, Cat., II, 378, 1860; LÜTKEN, Vid. Medd. Kjöbenh. 1875, 38; JORDAN & GILBERT, Synopsis, 417, 1883.

Echeneis squalipeta, DALDORF, Skrivt. Naturh. Selsk., II, 1797, 157, Atlantic Ocean between the tropics; GÜNTHER, Cat., II, 377, 1860.

Echeneis jacobæa, LOWE, Proc. Zool. Soc. London 1839, 89, Madeira.

Echeneis remoroides, BLEEKER, Batoë, II, 70, Batoe.

Echeneis parva, GRONOW, Cat. Fish., Ed. Gray, 92, 1854, no locality; after *E. remora*, L.

Echeneis postica, POEY, Memorias, II, 255, 1861, Havana. (Coll. Poey.)

Remora jacobæa, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 239.

Subgenus REMORINA, Jordan & Evermann.

2607. REMORA ALBESCENS (Temminck & Schlegel).

Length of disk 3⅕ to 3¼ in total length; width between pectorals 5 to 5⅓; number of laminæ on disk 13 or 14. D. XIII–22; A. 22. Angle of mouth in the vertical from the third lamina of the disk. Length of ventral fins equal to the distance between root of pectoral and posterior margin of eye. Color uniform grayish brown. (Günther.) Tropical Pacific, straying to America; a specimen taken at La Paz, Gulf of California (Streets), and 1 in the Gulf of Mexico (Bean). (*albescens*, whitish.)

Echeneis albescens, TEMMINCK & SCHLEGEL, Fauna Japonica, Poiss., 272, pl. 120, fig. 3, 1842, Japan; GÜNTHER, Cat., II, 377, 1860; STREETS, Bull. U. S. Nat. Mus., VII, 54, 1877.

Remora albescens, JORDAN, Cat. Fishes, 66, 1885.

Subgenus REMOROPSIS, Gill.

2608. REMORA BRACHYPTERA (Lowe).

Head nearly 4; width between pectorals 6½. D. XIV to XVI–29 to 32; A. 25 to 30. Body robust, the greatest depth nearly twice the length of the short pectoral fins; disk shorter than base of dorsal, rather broad; upper jaw angular. Caudal nearly truncate. Light brown, darker below, fins paler. Warm seas, occasionally north to Cape Cod. (*βραχύς*, short; *πτερόν*, fin.)

Echeneis brachyptera, LOWE, Proc. Zool. Soc. London 1839, 89, Madeira; GÜNTHER, Cat., II, 378, 1860; JORDAN & GILBERT, Synopsis, 417, 1883.

Echeneis sexdecimlamellata, EYDOUX & GERVAIS, Voy. Favorite, V, 77, pl. 31, 1839, Indian Ocean?

Echeneis quatuordecimlaminatus, STORER, Rept. Fishes Mass., 155, 1839, Holmes Hole.

Echeneis pallida, TEMMINCK & SCHLEGEL, Fauna Japonica, Poiss., 271, pl. 120, fig. 2, 3, 1842, Japan.

Echeneis nicuhofii, BLEEKER, Sumatra, II, 279, Sumatra.

Remoropsis brachypterus, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 60.

837. RHOMBOCHIRUS, Gill.

Rhombochirus, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 88 (*osteochir*).

This genus agrees with *Remora* in every respect excepting the structure of the pectoral fins. These are short and broad, rhombic in outline, the rays all flat, broad and stiff, being partially ossified, although showing the usual articulation; upper rays of pectoral broader than the others. One species known. (*ῥόμβος*, rhomb; *χείρ*, hand.)

2609. RHOMBOCHIRUS OSTEOCHIR (Cuvier).

Head $4\frac{3}{4}$ in length; disk $2\frac{1}{4}$; width between pectorals 5. D. XVIII–21 to 23; A. 20 or 21; P. 20. Mouth very small, maxillary not nearly reaching to the line of the orbit; outer series of teeth longer than the others. Disk very large, broader and rougher than in *Remora remora*, extending forward beyond the tip of the snout. Caudal fin emarginate, with rounded angles. Light brown; underside of head, ventral line, part of ventrals and a spot on pectorals pale. West Indies north to Cape Cod; parasitic on species of *Tetrapturus;* rather rare. (*ὀστέον*, bone; *χείρ*, hand.)

Echeneis osteochir, CUVIER, Règne Animal, Ed. 2, vol. II, 348, 1829, no locality given; GÜNTHER, Cat., II, 381, 1860; JORDAN & GILBERT, Synopsis, 418, 1883.
Echeneis tetrapturorum, POEY, Memorias, II, 256, 1858, Cuba. (Coll. Poey.)

Group TRACHINOIDEA.

(THE TRACHINOID FISHES.)

A large group of transitional forms, some of them of doubtful relationships, showing affinities with the *Percoidea* on the one hand and with the *Batrachoididæ* and *Blennoidea* on the other. In general, the spinous dorsal is short or weak, the soft dorsal long and similar to the anal, and the squamation is less complete and less ctenoid than in the *Percoidea*. The skull is, in general, depressed, with the supraocular crest low, and the suborbital stay is wanting, although in some genera the suborbital bones are enlarged. The bones of the skull are not strongly armed, and the ventral fins are often inserted well forward, and they are sometimes reduced in size. The group is divided by Dr. Gill into *Percophidoidea*, *Trachinoidea*, and *Uranoscopoidea*. The two latter groups are natural and related, but, as Dr. Gill observes, "the *Percophidoidea* are undoubtedly a heterogeneous group and need a thorough revision." The relations of *Bathymaster*, *Trichodon*, and *Latilus* especially are uncertain. Several of the leading families of this group are confined to the South Temperate Zone, and none of the *Trachinidæ* occurs within our limits.

a. Mouth horizontal or moderately oblique, the lips not fringed; eyes lateral; ventral rays I, 5, their insertion more or less before the pectorals; suborbitals moderate; gills 4, a slit behind the fourth.
b. Snout subconic, not prolonged and spatulate; ventrals not widely separated.
c. Body covered with scales; dorsal spines flexible.
d. Lateral line complete; caudal fin forked; vertebræ 24 to 27.

dd. Lateral line incomplete, running close to the back; caudal rounded or lanceolate; dorsal fin continuous.

e. Vertebræ about 27; scales cycloid; maxillary more or less dilated behind, with a supplemental bone; middle rays of ventrals longest. OPISTHOGNATHIDÆ, CXCI.

ee. Vertebræ about 50; scales ctenoid; maxillary not dilated, without supplement bone; inner rays of ventrals longest. BATHYMASTERIDÆ, CXCII.

cc. Body naked; snout short; mouth very large, the maxillary much produced behind; jaws with sharp canines; lateral line well developed; dorsals 2; caudal forked. CHIASMODONTIDÆ, CXCIII.

bb. Snout much prolonged and spatulate; ventrals widely separated; body scaly or naked; lateral line near the back; dorsal usually divided. CHÆNICHTHYIDÆ, CXCIV.

aa. Mouth vertical, the lips fringed.

f. Eyes lateral; gills 4, a slit behind the last; preopercle armed; body naked, compressed; caudal lunate, on a slender peduncle; vertebræ about 48. TRICHODONTIDÆ, CXCV.

ff. Eyes superior; gills more or less reduced, usually $3\frac{1}{2}$, the last slit small or wanting; suborbitals more or less dilated; body scaly or naked.

g. Lateral line well developed, concurrent with the back anteriorly; dorsal spines slender, not pungent; vertebræ about 25 to 30.

h. Ventral rays I, 3. DACTYLOSCOPIDÆ, CXCVI.

gg. Lateral line obscure; dorsal spines few, more or less pungent, sometimes obsolete. URANOSCOPIDÆ, CXCVII.

Family CXC. MALACANTHIDÆ.

(THE BLANQUILLOS.)

Body more or less elongate, fusiform or compressed. Head subconical, the anterior profile usually convex; suborbital without bony stay; the bones not greatly developed; cranial bones not cavernous; opercular bones mostly unarmed. Mouth rather terminal, little oblique; teeth rather strong; no teeth on vomer or palatines; the premaxillary usually with a blunt posterior canine, somewhat as in the *Labridæ;* premaxillaries protractile; maxillary without supplemental bone, not slipping under the edge of the preorbital. Gills 4, a long slit behind the fourth; pseudobranchiæ well developed; gill membranes separate, or more or less united, often adherent to the isthmus; lower pharyngeals separate. Scales small, ctenoid; lateral line present, complete, more or less concurrent with the back; dorsal fin long and low, usually continuous, the spinous portion always much less developed than the soft portion, but never obsolete; anal fin very long, its spines feeble and few; caudal fin forked; tail diphycercal; ventrals thoracic or subjugular, I, 5, close together; pectoral fins not very broad, the rays all branched; vertebræ in normal or slightly increased number (24 to 30). Pyloric cœca few or none. Fishes of the temperate and tropical seas, some of them reaching a large size. Genera about 6; species about 8 to 10, mostly American. The relationships of the family are obscure, and it may be that the genera here associated are not really closely allied. (*Malacanthidæ*, Günther, Cat., III, 359, 1861; *Trachinidæ*, part, Günther, Cat., II, 225–264, 1860.)

MALACANTHINÆ:

a. Vertebræ 24; preopercle entire.

b. Soft dorsal and anal extremely long, each with more than 40 rays; preopercle entire; form slender; scales very small. MALACANTHUS, 838.

aa. Vertebræ more than 24; preopercle more or less serrate.

CAULOLATILINÆ:

c. Soft dorsal and anal moderate, each with 22 to 27 soft rays; preopercle serrate; scales rather small; form robust.

d. Upper jaw with posterior canines; dorsal spines graduated. CAULOLATILUS, 839.

LATILINÆ:

cc. Soft dorsal and anal short, each of 13 to 15 soft rays; preopercle denticulate; scales small; form robust.

e. Nape with a large adipose appendage; a fleshy prolongation on each side of the labial fold, extending forward behind angle of mouth. LOPHOLATILUS, 840.

838. MALACANTHUS, Cuvier.

(MATAJUELO BLANCO.)

Malacanthus, CUVIER, Règne Animal, Ed. 2, vol. II, 205, 1829 (*plumieri*).

Body elongate, slightly compressed; cleft of mouth horizontal, with the jaws equal; eyes lateral; scales very small, minutely ciliated; one continuous dorsal, with the first 4 to 6 rays not articulated; dorsal and anal very long; pectoral rays all branched; jaws with villiform teeth; an outer series of stronger teeth, some of them canine-like, and with a canine at the posterior extremity of the intermaxillary; no teeth on the palate; preopercle entire; opercle with a spine; gill rakers little developed; vertebræ in small number, $10+14=24$. One species, a shore fish of tropical America. (*μαλακός*, soft; *ἄκανθα*, spine.)

2610. MALACANTHUS PLUMIERI (Bloch).

(MATAJUELO BLANCO.)

Head $3\frac{2}{3}$; depth $6\frac{1}{2}$. D. VI, 49; A. 48; scales 14–130–30; eye $5\frac{1}{2}$ in head; maxillary $2\frac{1}{3}$; snout $2\frac{1}{2}$; P. 2; longest dorsal rays 3, equal to anal ray; upper caudal lobe $1\frac{2}{3}$. Body elongate, little compressed. Head moderately long and pointed; eye placed high; interorbital flat, as wide as eye; profile of head obliquely straight from tip of snout to above nostril, where there is a slight angle formed, thence nearly horizontally straight to dorsal. Mouth large, maxillary reaching slightly past the vertical from posterior nostril; jaws equal; a band of villiform teeth in upper jaw growing broader anteriorly, and another row of small, even, conical teeth at the sides, and 6 well-developed canines in front, the 2 outer ones the largest; a canine on premaxillary at angle of mouth; villiform teeth in lower jaw not extending very far back; large recurved canines on side of jaw anteriorly, small conical teeth in front and on sides posteriorly, with a single large canine at angle of mouth; gill rakers rudimentary, about $5+7$. Top of head forward from above middle of eye, preorbital, and lower jaw, naked; fins without scales. Dorsal and anal similar, long and low, continuous; pectoral reaching past tips of ventrals to front of

anal; ventrals not reaching to vent, origin of ventral spine slightly behind base of pectoral; caudal forked, the lobes elongate, sometimes produced into a filament. Color in spirits, uniform, pale olive brown above, white below; fins light brownish; no distinct markings. Length 15 inches. West Indies, rather common; used as food. Here described from specimens from Havana. (Named for Père Plumier, of Martinique.)

Mateguelo blanco, PARRA, Dif. Piezas Hist. Nat. Cuba, 22, tab. 13, f. r. 1787, Cuba.
Coryphæna plumieri, BLOCH, Ichthyol., v, 119, pl. 175, 1787, Martinique; from a drawing by PLUMIER.
Malacanthus trachinus, VALENCIENNES, in CUVIER, Règne Animal, pl. 90, fig. 3.
Sparus oblongus, BLOCH & SCHNEIDER, Syst. Ichth., 283, 1801; after PARRA.
Malacanthus plumieri, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XIII, 319, 380, 1839, specimens from San Domingo; GÜNTHER, Cat., III, 359, 1861.

839. CAULOLATILUS, Gill.

(BLANQUILLOS.)

Caulolatilus, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 240 (no diagnosis), and GILL, Proc. Ac Nat. Sci. Phila. 1865, 66 (*chrysops*).
Dekaya, COOPER, Proc. Cal. Ac. Sci. 1864, 70 (*princeps*), not *Dekayia*, MILNE-EDWARDS & HAIME, 1851, a genus of corals.

Body elongate, subfusiform, not strongly compressed, heavy forward, tapering to a rather slender caudal peduncle; profile of head strongly arched; mouth moderate, little oblique, the jaws nearly equal; lips thick; maxillary narrow, not slipping under the preorbital; teeth in villiform bands, preceded by a row of stronger acute teeth; posterior teeth in each jaw canine-like, directed forward; posterior canines of upper jaw largest; no teeth on vomer or palatines; preopercle pectinate, the teeth nearly even; opercle with a blunt, flat spine; eyes large, lateral; gill membranes slightly connected, forming a fold across the isthmus, with which they are narrowly joined; branchiostegals 6; gill rakers short and stout; nostrils double, round, close together; scales small, firm, ctenoid; lateral line continuous, concurrent with the back; dorsal with 7 to 9 slender, pointed, graduated spines and 22 to 27 soft rays; anal similar to soft dorsal, with 1 or 2 small spines and more than 20 soft rays; caudal fin forked; ventral fins thoracic; no adipose appendage at the nape; vertebræ 12+15=27. Large fishes of the warm seas of America; valued as food. (*καυλός*, stem; *Latilus;* being distinguished from *Latilus* by the many rays.)

a. Scales small, about 125 in the lateral line, about 50 in a transverse series.
b. Eye large, 4½ in the head; depth 4 in length; scales 16-125-40. PRINCEPS, 2611.
bb. Eye small, 6 in head; depth 3½ in body; scales 13-120-35. MICROPS, 2612.
aa. Scales larger, about 108 in the lateral line, about 25 in a transverse series; scales 12—108—25. CYANOPS, 2613.

2611. CAULOLATILUS PRINCEPS (Jenyns).

(BLANQUILLO; WHITE-FISH.)

Head 3⅜; depth 4. D. IX, 24; A. II, 23; scales 16-125-40. Flesh of the occiput becoming thick with age, as in *Harpe*. Eye large, about ⅓ the convex interorbital space, 4½ in head; maxillaries reaching front of eye;

teeth rather strong; preopercle finely, evenly, and acutely serrate behind, nearly entire below; preopercle, interopercle, and preorbital naked; cheeks and opercles scaly; top of head scaled on the median line to between the eyes; dorsal spines flexible; ventrals slightly behind the pectorals, the outer rays longest; caudal moderately forked, the upper lobe the longer; caudal peduncle short and slender, abruptly contracted; pectorals falcate, longer than caudal, $\frac{5}{6}$ length of the head. Olivaceous, with bluish reflections; brownish above, greenish below; fins light greenish olive, tinged with bluish and orange, the colors always pale; dorsal and anal greenish, with a bluish band near the tip; axil dusky. Rocky islands of the Pacific coast from Monterey southward to the Galapagos; abundant about the Santa Barbara Islands; a food fish of considerable importance. Length 40 inches. We are unable to detect any differences by which the Californian form, *Caulolatilus anomalus*, can be separated from *Caulolatilus princeps*. (*princeps*, a leader.)

Latilus princeps, JENYNS, Zool. Beagle, Fishes, 52, pl. 11, 1840, Chatham Island, Galapagos Archipelago (Coll. Charles Darwin); GÜNTHER, Cat., II, 253, 1860.

Dekaya anomala, COOPER, Proc. Cal. Ac. Sci. 1864, 70, coast of Southern California.

Caulolatilus affinis, GILL, Proc. Ac. Nat. Sci. Phila. 1865, 68, Cape St. Lucas. (Coll. John Xantus.)

Caulolatilus princeps, GILL, *l. c.* 68.

Caulolatilus anomalus, GILL, *l. c.* 68; STREETS, Bull. U. S. Nat. Mus., VII, 48, 1877; JORDAN & GILBERT, Synopsis, 625, 1883.

2612. CAULOLATILUS MICROPS, Goode & Bean.

Head $3\frac{1}{4}$; depth $3\frac{1}{2}$. D. VII, 25; A. I, 23; scales 105 counting the oblique series, 120 counting the row above lateral line; transverse rows 12 + 30; eye $5\frac{3}{4}$ in head; snout 2; maxillary $2\frac{1}{2}$; pectoral $1\frac{1}{3}$; ventral $2\frac{1}{10}$; highest dorsal spine $3\frac{1}{3}$; highest anal rays $3\frac{1}{2}$; upper caudal lobe $1\frac{1}{3}$. Body rather robust; upper profile of head rather steep, evenly rounded from tip of snout to dorsal; nostrils small, midway between eye and tip of snout, separated by a distance equal to $\frac{1}{2}$ diameter of pupil; mouth large, maxillary scarcely reaching to anterior margin of eye; lips thick; lower jaw included; jaws with small conical teeth, the outer row enlarged, canine-like, a large tooth on posterior end of maxillary at angle of mouth; preopercle finely and evenly serrate on its vertical limb; a broad flat spine on opercle; snout, preorbital and lower jaw naked; fins scaleless; dorsal and anal similar, long and low; pectorals reaching far past tips of ventrals to vent; origin of ventral spine about the length of 2 scales behind the vertical from pectoral base; caudal fin lunate when spread, its upper lobe slightly the longer. Color reddish, marked with yellow; a yellow band below the eye; a dark blotch in and above axil of pectoral; dorsal light at base, darker above, with many indistinct brownish spots. Gulf of Mexico, in rather deep water; not rare. Here described from a specimen from the Pensacola Snapper Banks, 26 inches in length. A rather doubtful species, perhaps not distinct from *C. cyanops* or *C. chrysops*. (*μικρός*, small; *ὤψ*, eye.)

Caulolatilus microps, GOODE & BEAN, Proc. U. S. Nat. Mus. 1878, 43, off Pensacola, Florida. (Coll. Silas Stearns.)

Caulolatilus chrysops, JORDAN & GILBERT, Synopsis, 626, 1883; not *Latilus chrysops*, CUVIER & VALENCIENNES.

2613. CAULOLATILUS CYANOPS,* Poey.

(Blanquillo.)

Head 4 in total length. D. VII, 24; A. I, 22 (scales 10–108–25. Bean). Profile convex before the eye, not ascending to the nape; no scales on the fins; soft rays little divided; caudal slightly lunate; first caudal vertebra spoon-like, its cavity receiving the air bladder; vertebræ 12 + 15; no pyloric cæca, stomach short, air bladder large. Color greenish above, a faint, broad, interrupted brown band above the lateral line; some small brown spots above and below it; region below the eye clear blue, not very different from the color of the belly; soft dorsal brown, paler at its base, edged with orange; spinous dorsal orange. (Poey.) Coast of Cuba; not seen by us. Both this and the preceding species may be identical with *Caulolatilus chrysops*, a species described from the coast of Brazil. (κυάνεος, blue; ὤψ, eye.)

? *Latilus chrysops*, Cuvier & Valenciennes, Hist. Nat. Poiss., ix, 496, 1833, Brazil (Coll. M. Gay); Günther, Cat., II, 253. 1860.
Caulolatilus cyanops, Poey, Repertorio, I, 312, 1867, Cuba. (Coll. Poey.)

840. LOPHOLATILUS, Goode & Bean.

(Tile-Fishes.)

Lopholatilus, Goode & Bean, Proc. U. S. Nat. Mus. 1879, 205 (*chamæleonticeps*).

Body stout, somewhat compressed; mouth moderate, maxillary reaching anterior margin of the orbit; opercle and preopercle scaly, the latter finely denticulate; upper jaw with outer series of stronger teeth, behind which is a band of villiform teeth; lower jaw with a few large canines, and an inner series of small conical teeth; vomer and palatines toothless; nape with a large adipose appendage; a fleshy prolongation upon each side of the labial fold, extending backward beyond the angle of the mouth; stomach small, siphonal, barely more than a loop in the very large intestine; alimentary canal short, less than total length of the body; air bladder simple, with thick muscular walls, strongly attached to the roof of the abdominal cavity by numerous root-like appendages, resembling somewhat that of *Pogonias*. Deep-sea fishes. (λόφος, crest; *Latilus*.)

2614. LOPHOLATILUS CHAMÆLEONTICEPS, Goode & Bean.

(Tile-Fish.)

Head 3; depth 3½. D. VII, 15; A. II, 13; scales 8–93–30. Body stout, somewhat compressed, its greatest width equaling length of caudal peduncle; intermaxillaries supplied with a series of from 19 to 23 canine teeth, behind which is a band of villiform teeth, widest at the symphysis; mandible with about 12 large canines; eye rather small, its diameter 6½ in

* The characters distinguishing *Caulolatilus chrysops* are thus given by Poey: Head 4½ in total length. D. VIII, 24; A. II, 22. Profile most gibbous behind the eye; a very bright gilded band below the eye, broader anteriorly; dorsal fin brown with irregular blue spots; axillary spot green. Coast of Brazil.

head, and about twice length of labial appendages; distance between posterior nostril and eye equal first anal spine, and ½ distance from tip of snout to anterior nostril. Caudal fin emarginate, middle rays 1½ in outer rays; vent under interval between fourth and fifth dorsal rays. Back bluish, with a green tinge, iridescent, changing through purplish blue and bluish gray to rosy white below, and milky white toward median line of belly; head rosy, iridescent, with red tints most abundant on forehead, blue under the eyes, cheeks fawn-colored; throat and under side of head pearly white, with an occasional tint of lemon yellow, most pronounced in front of ventrals and on anterior portion of ventral fins; back with numerous maculations of bright yellow or golden; anal purplish, with blue and rose tints, iridescent; margin of anal rich purplish blue, iridescent, like the most beautiful mother-of-pearl, this color pervading more or less the whole fin, which has large yellow maculations, the lower border rose-colored, like the belly, base of the fin also partaking of this general hue; dashes of milk white on base of anal between the rays; dorsal gray; in front of the seventh dorsal the upper third posterior to the upper two-thirds dark brown; spots of yellow, large, elongate, on or near the rays; adipose fin whitish brown or yellow, a large group of bright yellow confluent spots at the base; pectorals sepia-colored, with rosy and purplish iridescence. (Goode & Bean.) Deep waters of the western Atlantic, at times very abundant; now rare or almost extinct. "The tilefish was first observed in 1879 by fishermen fishing for cod on Nantucket Shoals. From its abundance it was thought to become of some economic importance. In March and April, 1882, vessels arriving at New York, Philadelphia, and Boston reported having passed large numbers of dead and dying fish, the majority of which were tilefish. Captain Collins estimated the area covered by dead and dying fish to be from 5,000 to 7,500 square statute miles, the number of fish to be 1,000,000,000. Several visits were made by the Fish Commission vessels to the grounds where these fishes were formerly abundant, but no specimen was obtained, and it was thought to have become extinct. In 1892 several specimens were taken by the *Grampus* in latitude 38° to 40° N., and longitude 71° to 73° W. The wholesale destruction of the tilefish in 1882 is thought by Colonel McDonald to be due to climatic causes." (Goode & Bean.) (*chamæleon, χαμαιλέων*; *-ceps*, head.)

Lopholatilus chamæleonticeps, Goode & Bean, Proc. U. S. Nat. Mus. 1879, 205, Nantucket Shoals; Jordan & Gilbert, Synopsis, 624, 1883; Collins, Rept. U. S. Fish Comm. (1882) 1884, 237; Lucas, Rept. U. S. Nat. Mus. (Smithsonian Report) 1889, 647, with plate; Goode & Bean, Oceanic Ichthyology, 284, 1896.

Family CXCI. OPISTHOGNATHIDÆ.

(The Jaw-Fishes.)

Body oblong or elongate, low, moderately compressed, covered with small cycloid scales; lateral line present, straight, running close to the dorsal fin, not extending much behind middle of body. Head large, naked, the anterior profile decurved, no ridges, spines, or crests above. Mouth

terminal, horizontal, its cleft usually very wide, the maxillary sometimes greatly dilated; supplemental maxillary present; premaxillaries protractile; jaws subequal, with conical or cardiform teeth; vomer usually with a few teeth; palatines toothless; opercles unarmed; no suborbital stay. Pseudobranchiæ present. Gill rakers rather long; gills 4, a slit behind the fourth; gill membranes somewhat united, free from the isthmus. Branchiostegals 6. Air bladder present. No pyloric cæca. Vertebræ large, about 27 in number. Dorsal fin long, continuous, its anterior half composed of slender, flexible spines, which pass gradually into soft rays; caudal distinct, rounded or lanceolate; tail not isocercal, the last vertebra expanded (27 to 34); anal long and low, without distinct spines; ventrals separate, jugular, I, 5, the middle rays longest; pectorals fan-shaped. Three genera, of about 15 species; small fishes inhabiting rocky bottoms in tropical seas, many of them with bright markings. The American species are all rarities, living about rocks in deep or shallow water; nowhere abundant and none of the species well represented in collections. (*Trachinidæ*, genus *Opisthognathus*, Günther, Cat., II, 254–256.)

a. Maxillary of great length, nearly as long as head, produced behind in a flexible lamina. OPISTHOGNATHUS, 841.

aa. Maxillary normal, truncate behind, much shorter than head.

b. Caudal moderate, rounded behind; body oblong, moderately compressed. GNATHYPOPS, 842.

bb. Caudal lanceolate, long and pointed; body elongate. LONCHOPISTHUS, 843.

841. OPISTHOGNATHUS, Cuvier.

Opisthognathus, CUVIER, Règne Anim., Ed. 2, vol. II, 240, 1829 (*sonnerati*).

Maxillary prolonged backward in a long flexible lamina, which reaches about to base of pectoral. Characters of the genus otherwise included above. It has been suggested that the species of *Gnathypops* are females of analogous species of *Opisthognathus*, the long maxillary being a character of the male. This seems impossible, but deserves an investigation. The fact that *Gnathypops maxillosa* has but 27 vertebræ, while its long-jawed cognate, *Opisthognathus macrognathum*, is said by Poey to have 34 vertebræ, is opposed to this view, as is also the fact that the analogous species do not in other respects exactly correspond, as in *Gnathypops mystacina*, the scales are smaller than in *Opisthognathus lonchurum; Gnathypops rhomalea* has fewer fin rays than *Opisthognathus punctatum*, etc. But the parallelism of species in the two genera living in the same waters is remarkable. (ὄπισθε, behind; γνάθος, jaw.)

a. Scales moderate, about 67; D. X, 15; A. II, 13; body nearly plain olivaceous, the maxillary not distinctly striped within. LONCHURUM, 2615.

aa. Scales very small, 100 to 150 in longitudinal series; dorsal rays about XI, 17; A. II, 16; body and fins much variegated, the maxillary within with 2 ink-black stripes on a milk-white ground.

b. Dorsal without large black spot in front; scales 120. PUNCTATUM, 2616.

bb. Dorsal with a large black spot more or less ocellated.

c. Scales about 100. MACROGNATHUM, 2617.

cc. Scales about 140. OMMATUM, 2618.

2615. OPISTHOGNATHUS LONCHURUM, Jordan & Gilbert.

Head $3\frac{1}{3}$; depth $4\frac{3}{5}$. D. 25; A. 15; scales 67. Head moderate; snout very short, shorter than pupil; eye $3\frac{1}{3}$ in head; maxillary $1\frac{1}{3}$ in head, rather narrow; lower jaw included; vomer with 5 rather large teeth. Longest dorsal spine about as long as head, slightly higher than soft rays; caudal long, the middle rays scarcely shorter than head; longest anal rays $1\frac{1}{5}$ in head; pectoral little more than $\frac{1}{2}$ head. Scales moderate. Olivaceous; margin of upper lip with a narrow black stripe; caudal with 3 dusky bars; color of rest of body uniform. Gulf of Mexico, in deep water. Two specimens known, taken from the stomach of a Red Snapper at Pensacola, Florida. The species resembles *Gnathypops mystacinus*, found in the same waters, but the latter species has smaller scales. (λόγχη, lance; οὐρά, tail.)

Opisthognathus lonchurus, Jordan & Gilbert, Proc. U. S. Nat. Mus. 1882, 290, Snapper Banks, off Pensacola, Florida (Type, No. 29671. Coll. Jordan & Stearns); Jordan & Gilbert, Synopsis, 943, 1883.

2616. OPISTHOGNATHUS PUNCTATUM, Peters.

D. 28; A. 18. Body moderately elongate; scales very small, about 125 in lateral line. Dorsal spines continuous with the soft rays. No vomerine teeth. Maxillary very long, extending slightly beyond head. Head everywhere finely speckled with black, the body more coarsely and irregularly spotted; pectoral finely and closely speckled, its edge plain; ventral fin dusky, similarly marked; dorsal without large black blotch, finely spotted, the spots behind gradually forming the boundaries of white ocelli, the base of the fins having rings of white around black spots, the upper part with dark rings around pale spots; caudal with pale spots, its edge, like that of the dorsal, somewhat dusky, not black; anal with a broad, blackish edge, and with dark spots, those near the base of the fin largest; lining membrane of maxillary with the usual bands of white and inky black. Mazatlan. Only the type of this species is yet known, this description having been taken by us from the original specimen. It bears considerable resemblance to *Gnathypops rhomalea*, which is found in the same waters, differing in the generic character of the dilated maxillary. (*punctatus*, spotted.)

Opisthognathus punctatus, Peters, Berliner Monatsberichte 1869, 708, Mazatlan; Jordan, Proc. Ac. Nat. Sci. Phila. 1883, 290; Jordan, Cat. Fish. N. A., 118, 1885.

2617. OPISTHOGNATHUS MACROGNATHUM, Poey.

Head $3\frac{2}{3}$; depth 5. D. XI, 16; A. II, 16 or 17; P. 17; scales 100. Body moderately elongate, somewhat compressed. Head blunt anteriorly; snout very short, about as long as pupil; eye large, 4 in head; maxillary reaching slightly past edge of preopercle, but not to end of head, its length contained $3\frac{3}{4}$ times in length of body. Teeth rather strong, wide set, forming 2 distinct series, directed backward, especially in the upper jaw; lateral teeth of lower jaw largest; a single vomerine tooth. Gill rakers long and slender, nearly 20 below angle. Scales very small.

Dorsal fin low, continuous, the soft rays but little higher than the spines, which are slender and flexible, the longest $3\frac{1}{3}$ in head; caudal short, rounded, its length $5\frac{3}{5}$ in body; anal similar to soft dorsal; pectoral $\frac{1}{2}$ as long as head. Grayish olive, much variegated with yellowish and dark olive; about 6 irregular dusky bands on the body, which extend on the dorsal fin; whitish markings on body forming roundish spots, surrounded by reticulations of grayish olive; head marbled, its posterior part, as well as the sides of the back and pectoral base, with small blackish dots; membrane lining inside of maxillary with 2 curved inky-black bands on a white ground; angle of mouth with a black spot; lining of opercle black; fins all variegated like the body. Florida Keys to Cuba. Here described from the type of *O. scaphiurum*, from Garden Key, but *O. macrognathum* seems to be the same. (*μακρός*, long; *γνάθος*, jaw.)

Opisthognathus macrognathus,* Poey, Memorias, II, 284, July, 1860, Cuba. (Coll. Poey.)

Opisthognathus megastoma, Günther, Cat., II, 255, September, 1860, Gulf of Mexico. (Haslar Collection.)

Opisthognathus scaphiurus, Goode & Bean, Proc. U. S. Nat. Mus. 1882, 417, Garden Key, Florida (Type, No. 5936, U. S. Nat. Mus. Coll. Dr. Whitehurst); Jordan & Gilbert, Synopsis, 943, 1883.

2618. OPISTHOGNATHUS OMMATUM, Jenkins & Evermann.

Head 3; width of head 5; depth 5; eye 3 in head. D. 28; A. 18; scales about 140. Body moderate, compressed, depth $4\frac{1}{7}$; width behind the head $8\frac{3}{10}$ in length of body. Head large, its breadth equaling its depth, being 5 in length of body. Scales small, embedded; head naked, lateral line extending past middle of dorsal fin. Mouth large. Maxillary long, $1\frac{1}{2}$ in head; postorbital portion $2\frac{3}{10}$ in head, not extending beyond head; snout short, its length less than $\frac{1}{2}$ diameter of eye; distance from tip of snout to end of maxillary $3\frac{3}{10}$ in length of body and $1\frac{1}{8}$ in head. Teeth in front part of each jaw in several series, on sides of jaws reduced to a single series, the outer series strong; a tooth on the vomer; gill membranes connected; the interorbital space very narrow, $11\frac{3}{5}$ in head. Distance from snout to origin of dorsal but little greater than length of head; space between dorsal and caudal fins $\frac{1}{2}$ greater than length of snout; no depression between spinous and soft rays of dorsal fin, the dorsal equaling the anal in height, its longest ray $1\frac{3}{5}$ times the eye; pectorals slightly longer than ventrals, being 2 in head; breadth of pectorals 3 in head; ventrals inserted slightly in front of pectorals; caudal rounded and narrow. Coloration: Body irregularly mottled with dark, head evenly blackish; dorsal fin blackish on the posterior portion, with 2 rows of 4 or 5 pale spots well separated; a large ocellated spot from the third to the sixth spines, including them, greater than diameter of eye; anal fin black, with a series of pale spots on the rays, the base pale; caudal black, with 2 pale

*Poey thus describes his specimens of *Opisthognathus macrognathum:* "Head $3\frac{1}{4}$ in total; depth $5\frac{1}{4}$; eye nearly 4, twice length of snout, 3 times interorbital width. D. XI, 16; A. II, 16. Vomer with 2 teeth; spines not pungent. Body covered with large yellowish points on a brown ground; 7 broad brown bands on sides, not reaching belly, but extending to middle of dorsal, which, like the anal, has yellow points; a large black ocellus between sixth and ninth spines of dorsal; maxillary with 2 ink-black bands on a milk-white ground; pectorals, ventrals, and caudal yellowish with black points. No pyloric cæca; vertebræ $10 + 24 = 34$, the first 5 strong." (Poey.)

spots at the base and a row of spots across the middle; lining of maxillary with bands of black and white. Bay of Guaymas; 3 specimens known. (*ὀμματός*, eyed, from its ocellate dorsal.)

Opisthognathus ommata, JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 153, Guaymas. (Type, No. 39640. Coll. Jenkins & Evermann.)

842. GNATHYPOPS, Gill.

Gnathypops, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 241 (*maxillosus*).

This genus differs from *Opisthognathus* in having the maxillary of medium length and truncate behind, not extending to edge of opercle; caudal moderate, rounded behind. Species in form and habit agreeing closely with those of *Opisthognathus*. (*γνάθος*, jaw; *ὕπο*, below; *ὤψ*, eye.)

a. Body and fins spotted with black and often with pale.
  b. Dorsal fin with a conspicuous dusky blotch in front.
    c. Scales very small, about 120; dorsal rays 26. SCOPS, 2619.
    cc. Scales moderate, about 65; dorsal rays 15. MAXILLOSA, 2620.
  bb. Dorsal fin without distinct blotch in front; scales small, about 100.
    d. Dorsal rays about XI, 16; dark spots on head and body few. MACROPS, 2621.
    dd. Dorsal rays about XI, 13; dark spots on head and body numerous.
      e. Dorsal fin distinctly notched; lateral line not reaching middle of dorsal. RHOMALEA, 2622.
      ee. Dorsal fin not notched; lateral line reaching middle of dorsal. SNYDERI, 2623.
aa. Body and fins nearly uniform olive; the spots few and spare; dorsal rays 24; scales 100; no black on membrane of maxillary. MYSTACINA, 2624.

2619. GNATHYPOPS SCOPS, Jenkins & Evermann.

Head 3½; width of head 5¾, its depth 4½. D. 26 (X, 16); A. 19 (II, 17); scales 3–122–40. Scales small, none on head; lateral line extending to about middle of dorsal fin; mouth large; maxillary extending beyond eye a distance 4$\frac{1}{16}$ in head; snout 6½ in head; teeth in bands, outer series on upper jaw rather strong; a single tooth on vomer; gill membranes connected. Opercle ending in a long flap, which extends upward and backward, nearly meeting over the back in front of the dorsal fin. Eye large, 2½ in head; interorbital space narrow, 11 in head. No depression between the dorsal spines and the soft rays, which are scarcely distinguishable; height of dorsal equal to that of anal; ventrals inserted in front of pectorals; pectorals equal to ventrals in length, 7 in body; caudal rounded. Coloration, in alcohol: Body pale, covered with many dark spots about the size of 3 to 6 scales; top of head with smaller dark spots; sides of head with whitish spots; dorsal fin with a black ocellated spot equal to eye on the space between second and fifth spines; remainder of fin dark, with many white spots running into each other on some portions, so as to form irregular lines; base of anal pale, the outer edge black; caudal dark with 2 whitish spots at the base, and a row of 6 white spots across the middle on alternate rays; pectorals lighter, with small whitish specks; ventrals dusky; belly pale. Guaymas; 3 specimens known, respectively 115 cm., 10 cm., and 7 cm. in length to base of caudal. This species is the

analogue of *Opisthognathus ommatum*, also from Guaymas. (*scops*, the screech owl; *σκώψ*, from *σκοπέω*, to look, in allusion to the large eyes.)

Gnathypops scops, JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 152, Guaymas. (Type, No. 39641, U. S. Nat. Mus. Coll. Jenkins & Evermann.)

2620. GNATHYPOPS MAXILLOSA (Poey).

Head $3\frac{2}{5}$; depth $4\frac{1}{3}$. D. VIII, 17; A. II, 13; scales 65. Body moderately compressed; head not very large; maxillary truncate behind, extending behind eye for a distance for about $\frac{2}{3}$ diameter of eye, its length $1\frac{2}{3}$ in head; eye $3\frac{2}{3}$ in head. Teeth conical, curved, well separated, mostly in a single series; no teeth on vomer. Fins moderate; dorsal continuous, its spines slender; caudal short, its length $\frac{2}{3}$ head. Color grayish olive, with 7 irregular Λ-shaped bars of darker, everywhere much marbled and variegated; fins all similarly marked, the ventrals dusky, the dorsal with a dusky blotch in front. Cuba, north to Florida, from which locality the specimen here described was taken. Evidently very close to the Brazilian species *G. cuvieri*, the eye perhaps smaller. According to Poey, *Gnathypops maxillosa* has the eye 4 in head; D. VIII, 18; A. II, 15; 2 teeth on vomer; spinous dorsal lower than soft dorsal; body covered with large yellow spots on a ground color of clear brown; 6 brown cross bands reaching middle of dorsal, which is variegated with yellow and reddish, as is the anal; maxillary yellowish on its posterior border, the middle blackish; other spots on the jaws; ventrals, pectoral, and caudal yellowish, the ventrals finely spotted with brown, the caudal with 5 brown bands; base of pectorals with dark spots. Vertebræ $10 + 17 = 27$. (*maxillosus*, pertaining to the jaw.)

Opisthognathus maxillosus, POEY, Memorias, II, 286, 1860, Cuba. (Coll. Poey.)

Gnathypops maxillosus, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 241; POEY, Synopsis, 400; JORDAN & GILBERT, Synopsis, 942, 1883.

2621. GNATHYPOPS MACROPS (Poey).

Head $3\frac{1}{3}$ in total (with caudal?); depth about $4\frac{1}{2}$. D. XI, 16; A. II, 15 or 16; scales 100. Eye $3\frac{1}{3}$ in head; maxillary extending beyond eye $\frac{2}{3}$ of a diameter, $5\frac{1}{5}$ in head; vomer with 6 conical teeth. Color (faded in the type) reddish olive, with round, yellowish spots and vestiges of vertical bands; dorsal and anal plain, pectorals with brown bands; jaws not spotted with brown and white. Coast of Cuba (Poey); known from 1 specimen 132 mm. long; not seen by us. According to Poey, it may not be distinct from *G. maxillosa*, which in turn may possibly be the female of *Opisthognathus macrognathum*. This species may also be identical with the Brazilian species *Gnathypops cuvieri*,* but the latter has a dorsal ocellus and apparently larger scales. (*μακρός*, large; *ὤψ*, eye.)

Opisthognathus macrops, POEY, Memorias, II, 287, 1860, Cuba. (Coll. Poey.)

* *Gnathypops cuvieri* (Valenciennes). Head $3\frac{1}{8}$; depth $4\frac{1}{2}$; eye $3\frac{1}{8}$ in head. D. X, 18; A. II, 16; scales 70. Maxillary reaching beyond the vertical from posterior margin of orbit, $1\frac{1}{3}$ in head; eye $3\frac{1}{3}$ in head, dorsal fin not notched. Olivaceous; a large dark-blue ovate ocellate spot between the fourth and eighth dorsal spines; dorsal and anal mottled, the edge dusky posteriorly; caudal with 3 dark-bluish bands. Bahia (Valenciennes). (Named for Georges Dagobert Cuvier.)

Opisthognathus cuvieri, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 504, 1836, Bahia (Coll. Blanchet); GÜNTHER, Cat., II, 256, 1860.

2622. GNATHYPOPS RHOMALEA (Jordan & Gilbert).

Head $2\frac{7}{8}$; depth 4. D. XI, 13; A. II, 13; scales 103 (pores fewer). Body rather robust, compressed; head very large, ovoid, thicker and deeper than body, with swollen cheeks, the occipital region high, the snout somewhat truncate, the intermediate profile forming a nearly even curve; greatest depth of head equal to its thickness and $\frac{2}{3}$ its length. Eye not very large, 6 in head, longer than snout, about equal to the width of the flattish interorbital space. Mouth large, the maxillary extending well beyond the eye, but not to the margin of the preopercle nor to the mandibulary joint, its posterior margin truncate; supplemental bone small, but distinct; length of maxillary from end of snout $1\frac{3}{4}$ in head. Teeth moderate, in both jaws, in broad bands which become narrow on the sides; outer series of teeth somewhat enlarged, especially in upper jaw; 1 rather small, blunt tooth on middle of vomer. Gill membranes scarcely connected; gill rakers long and slender, about $\frac{3}{4}$ diameter of eye, 9 + 19; pseudobranchiæ situated in a cavity above the gill arches. Head naked; scales on body small, smooth, somewhat embedded; breast naked; lateral line ceasing opposite anterior third of second dorsal; 103 scales in a longitudinal series from head to caudal. Dorsal fin high; a rather deep notch separating the spines from the soft rays; the longest spines 3 in length of head, more than $\frac{1}{2}$ longer than the last spine, and scarcely lower than the soft rays. Insertion of dorsal opposite tip of the bony opercle, the opercular flap extending to opposite the third spine, last rays of dorsal and anal reaching past the base of caudal rays; caudal fin rounded, about $\frac{1}{2}$ length of head; anal higher than soft dorsal, its longest rays $2\frac{1}{4}$ in head; ventrals large, close together, inserted in front of pectoral, $1\frac{1}{3}$ in head; pectorals short and broad, $1\frac{1}{3}$ in head. Color in spirits, olivaceous, slightly brownish above, scarcely paler below, everywhere more or less tinged and mottled with greenish; head everywhere thickly and closely covered with small rounded dark-brown spots, largest above and on cheeks, where they are about as large as pin heads; smaller on lips and opercles, most thickly set on the anterior part of the head; eye thickly spotted; spots similar to those on the head extending along upper part of back, forming a vague band, which grows narrower backward and disappears opposite front of second dorsal; front side of pectoral and first 3 or 4 dorsal spines with dark spots; dorsal dusky olive, with darker clouds, and with some dark spots, especially on the spinous part; caudal and anal plain dusky or faintly marbled with paler; ventrals blackish, greenish at base; pectorals dusky green. Gulf of California, in shallow water; 1 specimen known, the type (above described) 16 inches long. This is perhaps the largest species of the genus. (*ῥομαλέος*, robust.)

Opisthognathus rhomaleus, Jordan & Gilbert, Proc. U. S. Nat. Mus. 1881, 276, Santa Maria Cove, Lower California. (Type, No. 29382. Coll. Lieut. Henry L. Nichols.)

2623. GNATHYPOPS SNYDERI, Jordan & Evermann, new species.

Head $3\frac{1}{3}$; depth $4\frac{1}{3}$. D. X, 14; A. II, 13; scales 93; 60 pores; 3 or 4 teeth on vomer; lateral line very distinct, extending to fourteenth ray of dorsal; no notch separating the dorsals, the spines and soft rays not sepa-

rable, the last spine not much shorter than the longest, which is $2\frac{1}{2}$ head; longest anal ray 2, pectoral $2\frac{1}{3}$. Vertebræ $10+17=27$. Body olivaceous, with 5 broad faint dusky cross shades; head with many round black spots of varying sizes, some as large as pin heads, the largest below and between eyes, covering both jaws and the membrane of the maxillary, few on cheeks, most numerous on forehead; similar spots extending along side of back to end of lateral line; dorsal dusky, with 6 round dusky blotches at its base, corresponding to the dark shades on body; caudal, anal, and pectorals plain dusky olive; ventrals blackish. Gulf of California; known from 1 specimen collected by Dr. Gilbert in San Luis Gonzales Bay. The species is close to *G. rhomalea*, but has a different dorsal fin and lateral line. Type, No. 2014 L. S. Jr. Univ. Mus., about 8 inches long. (Named for John O. Snyder, curator of fishes in Leland Stanford Junior University.)

2624. GNATHYPOPS MYSTACINA, Jordan.

Head $3\frac{1}{12}$ in length ($3\frac{5}{6}$ to tip of caudal); depth $4\frac{4}{5}$ ($5\frac{5}{6}$). D. 23 or 24 (X, 14); A. II, 11; lateral line with about 54 tubes; 100 scales between gill opening and caudal. Head rather elongate, very blunt in profile; snout very short, not longer than pupil; eye large, about $3\frac{1}{3}$ in length; maxillary $1\frac{2}{3}$ in length of head, 5 in length to base of caudal, $6\frac{1}{3}$ in total length to tip of caudal; end of maxillary abruptly truncate, not ending in a flexible lamina, the supplemental bone well developed; lower jaw slightly included. Teeth in each jaw in a narrow band, the outer slender, enlarged; vomer with about 4 slender teeth; palatines toothless. Gill rakers long and slender. Gill membranes nearly separate, free from the isthmus. Scales very small; lateral line extending to below anterior part of soft dorsal, its length $\frac{5}{4}$ that of head. Dorsal spines not distinguishable from the soft rays, the rays apparently fewer than usual, none of them very high, the last ray $2\frac{1}{4}$ in head; caudal short, apparently truncate, $1\frac{1}{4}$ in head; anal rather low; pectorals 2 in head; ventrals $1\frac{2}{3}$. Color nearly plain olive green, without bands or spots on body or fins; vertical fins tipped with blackish; maxillary with a faint median blackish stripe; pectoral with 2 dusky cross shades; no black or white on lining membrane of jaws. Length $3\frac{1}{2}$ inches. Deep waters of Gulf of Mexico; the few specimens known from the stomachs of Red Snappers (*Neomænis aya*) from the Pensacola Snapper Banks. It resembles *Opisthognathus lonchurum*, but the scales are smaller. (μύστα̃ξ, mustache, from the maxillary stripe).

Gnathypops mystacinus, Jordan, Proc. U. S. Nat. Mus. 1884, 37, Snapper Banks off Pensacola. (Coll. Jordan & Stearns. Type, 34976, U. S. Nat. Mus.)

843. LONCHOPISTHUS, Gill.

Lonchopisthus, Gill, Proc. Ac. Nat. Sci. Phila. 1862, 241 (*micrognathus*).

This genus differs from *Gnathypops* in the slender, compressed body, the still smaller maxillary and the lanceolate caudal fin. The single species is very rare. (λόγχη, lance; ὄπισθε, behind, from the form of the caudal.)

2625. LONCHOPISTHUS MIGROGNATHUS (Poey).

Head 5 in total; depth 6. D. X, 17; A. II, 16; scales 80; eye 3 in head. Body elongate, compressed, snout short; maxillary reaching ⅓ an eye's diameter behind the eye; no teeth on vomer; no second row of teeth in jaws; no scales on head except on cheek; lateral line almost touching profile of back. Caudal long and pointed as in *Gobius oceanicus*. Dark brown, paler below; 20 narrow vertical whitish bands from back to belly, the first 2 on cheek, the third on opercle; fins colored like body, except the pectorals, which are yellow, the edge orange. Vertebræ 10 + 18 = 28. Length 4 inches. Cuba; rare (Poey); only the types known, examined by us in the National Museum. (*μικρός*, small; *γναθος*, jaw.)

Opisthognathus micrognathus, Poey, Memorias, II, 287, 1860, Cuba. (Coll. Poey.)
Lonchopisthus micrognathus, Gill, Proc. Ac. Nat. Sci. Phila. 1862, 241.

Family CXCII. BATHYMASTERIDÆ.

(The Ronquils.)

Body rather elongate, moderately compressed, covered with small, ctenoid scales. Head rather large, subconic. Eyes large. Mouth moderate, nearly horizontal, the lower jaw slightly projecting; lips full; premaxillaries protractile, not extending to angle of the mouth; maxillary without supplemental bone, not slipping under the narrow preorbital. Teeth moderate, in a cardiform band in each jaw, the outer somewhat enlarged; bands of teeth on vomer and palatines. No barbels; no crests or spines on head. Branchiostegals 6. Gill membranes scarcely or broadly connected, free from the isthmus; gill rakers few, very short. Pseudobranchiæ large. Opercular bones unarmed. Mucous pores numerous on top and sides of head, sometimes provided with fringed flaps. Lateral line conspicuous, placed high, not quite reaching the caudal fin, its scales sometimes enlarged. Dorsal fin long, continuous, moderately high, a few of the foremost rays inarticulate, none of them pungent or spine-like; the posterior rays branched; anal fin long, similar to the dorsal; caudal convex; pectorals rather broad, their bases extending obliquely downward and backward, their rays all branched; ventrals slightly in front of pectorals, I, 5, close together, the inner rays longest. Skeleton well ossified. Pyloric cæca few (2 or 3). No anal papilla. Vertebræ in large numbers, about 14 + 35. Three species known, from the Northern Pacific; here referred to 3 genera. The relations of the group are uncertain; externally they resemble the *Opisthognathidæ*, but the relation can not be close, and the number of vertebræ is greatly increased.

a. Gill membranes not connected below.
  b. Head naked; scales in lateral line not enlarged; only first 3 or 4 rays in dorsal fin unbranched; pores of head with small flaps. Bathymaster, 844.
  bb. Head scaly on cheeks; scales in lateral line enlarged; anterior 20 to 30 rays of dorsal fin unbranched; pores of head mostly without flaps. Ronquilus, 845.
aa. Gill membranes broadly connected; cheeks scaly; scales in lateral line enlarged; about 15 of anterior rays of dorsal simple. Rathbunella, 846.

844. BATHYMASTER, Cope.

Bathymaster, COPE, Proc. Amer. Phil. Soc. 1873, 31 (*signatus*).

Head naked; pores of head large, many of them with dermal flaps; gill membranes scarcely connected; scales of lateral line similar to the others; dorsal fin with but 3 or 4 of its anterior rays unbranched; characters otherwise included above. (*βαθύς*, deep; *μαστήρ*, searcher.)

2626. BATHYMASTER SIGNATUS, Cope.

Head $3\frac{1}{2}$; depth 5. D. 47; A. 34; scales 6–95–19; eye $4\frac{1}{3}$ in head; maxillary $2\frac{1}{3}$; snout $4\frac{1}{3}$; pectoral $1\frac{4}{5}$; ventral $2\frac{1}{2}$; highest dorsal ray $2\frac{3}{4}$; highest anal ray $3\frac{1}{2}$; caudal $2\frac{2}{3}$. Body compressed, elongate, anterior profile convex from tip of snout to dorsal; mouth not very oblique, the maxillary reaching the vertical from posterior edge of orbit; snout about equal to eye; jaws equal, with bands of small conical teeth, outer row enlarged; lower jaw with a single row at the sides; well developed conical teeth on vomer and palatines. Branchiostegal membranes not united; margin of preopercle free, furnished with 5 or 6 conspicuous mucous pores; large pores on top and sides of head, each with a small flap; opercle ending in a flap behind; gill rakers moderately long and slender, $\frac{2}{3}$ eye, about 7 + 18; many mucous pores on top of head and under eye; head entirely naked; dorsal and pectoral with fine scales running about halfway up the fin; anal naked; a naked strip from nape to dorsal; pectoral broad and fan-shaped, its lower rays smaller, reaching to front of anal; origin of ventral spine about the diameter in front of the lower end of pectoral base; dorsal about uniform in height for nearly its entire length, higher than anal; dorsal and anal rays about reaching to base of caudal rays; first 3 or 4 rays of dorsal simple, the others branched; caudal truncate or slightly rounded. Color almost uniform warm brown with darker shades, the fins somewhat mottled with yellowish, the anal and ventrals blackish, other fins dusky; a conspicuous black ocellated blotch on front of dorsal, covering tips of 4 or 5 spines.* Shores of southern Alaska, from Unimak

* Concerning this species, Dr. Gilbert has the following note: "*Bathymaster signatus* is taken very abundantly in our series of shallow-water dredgings along the southern shore of the Alaskan Peninsula, and northward through Unimak Pass. The stations at which it was obtained are numbered 3211, 3212, 3213, 3214, 3215, 3217, 3220, 3222, and 3223, and the depth range from 34 to 56 fathoms. In addition, a very few small specimens were secured at Stations 3262, 3309, 3221 and 3333, north of the Aleutian Islands, in depths of 19 to 71 fathoms, but the species is evidently not abundant in Bering Sea. No examples were taken in any of the very numerous dredgings made in Bristol Bay. In life the sides are olive brown, and the upper parts show faint traces of 6 or 7 broad dusky cross bars, which correspond to or alternate with an equal number below the lateral line; the anal and ventral fins, the branchiostegal and gular membranes, the lower pectoral rays, and the snout blue black; anterior edge of orbit and front edge of preorbital light yellow; the pores on edge of preopercle, 2 pores above and behind maxillary, and 3 at upper edge of opercle, bright scarlet; a large black blotch on anterior dorsal rays; distal half of anterior portion of dorsal fin and the upper pectoral rays yellow. Outer ventral ray simple and inarticulate, followed by 5 branched rays. Only the first 2 dorsal rays spinous, being soft and flexible, but unjointed. The third and all following rays jointed and forked. All of the anal rays jointed. A specimen from *Albatross* Station 3211, 35 mm. in length to base of caudal, shows that the ventrals occupy very different positions in adults and in young. In the latter they are truly thoracic in position, and are inserted as much behind base of pectorals as they are located in advance of this point in adults. A specimen 65 mm. long is entirely similar to adults in this respect."

Pass to Sitka; not uncommon in water of moderate depths. Here described from a specimen collected by the *Albatross* (No. 2143, L. S. Jr. Univ. Mus.), Station 3214, 11 inches in length. Other specimens taken in rock pools at Sitka are dark green, almost black. (*signatus*, marked.)

Bathymaster signatus, COPE, Proc. Amer. Philos. Soc. 1873, 31, Sitka (Coll. Prof. George Davidson); GILBERT, Proc. U. S. Nat. Mus. 1888, 554.

845. RONQUILUS, Jordan & Starks.

Ronquilus, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 838 (*jordani*).

Cheeks scaly; scales of lateral line enlarged; anterior half of dorsal fin more or less composed of unbranched rays; mucous pores on head without conspicuous flaps; gill membranes separate. One species. (*Ronquil*, a Spanish name of the typical species, possibly from ῥόγχος, one who grunts.)

2627. RONQUILUS JORDANI (Gilbert).

(RONQUIL.)

Head $4\frac{1}{6}$; depth $6\frac{2}{3}$. D. 41; A. 33; V. I, 5; P. 18; scales 92+6 (tubes), about 200 transverse. Body rather elongate, moderately compressed. Eye large, about as long as snout, 4 in head, its diameter much more than the interocular space; maxillary extending to below front of pupil; cheeks closely scaly; rest of head entirely naked; a narrow, naked area in front of dorsal, bounded by rows of mucous pores; skull with large mucous cavities behind the eyes, which are translucent in life; scales of lateral line enlarged, twice as far apart as the others. Dorsal fin inserted at a distance behind the occiput, less than the diameter of the eye; pectorals $\frac{2}{3}$ the length of the head; fourth ray of ventrals longest; vent much nearer snout than root of caudal; vertebræ 14+35=49; olivaceous, tinged with brown; about 8 round, faint-bluish blotches along the sides, each surrounded by rings of yellow spots; a yellow ring around the eye and a yellow band along the cheek; fins translucent, the anal with a yellowish strip and a deep-bluish or black edging; dorsal reddish or yellow, with a dusky blotch in front; ventrals dusky; pectorals with the lower rays blackish or dark blue, larger specimens nearly uniformly dark, the color varying with the surroundings. Bristol Bay to Puget Sound, about rocks, in water of moderate depth. Length 6 to 10 inches. Known from Seattle and from Wrangel and Bristol Bay, the latter specimen in 32 fathoms. (Named for its discoverer, David Starr Jordan.)

Bathymaster signatus, JORDAN & GILBERT, Synopsis, 623, 1883; not *B. signatus*, COPE.
Bathymaster jordani, GILBERT, Proc. U. S. Nat. Mus. 1888, 554, Elliott Bay at Seattle (Coll. Jordan) and Fort Wrangel, Alaska. (Coll. *Albatross*.)
Ronquilus jordani, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 838, pl. 99.

846. RATHBUNELLA, Jordan & Evermann.

Rathbunella, JORDAN & EVERMANN, Check-List Fishes, 463, 1896 (*hypoplectus*).

This genus differs from *Ronquilus* in having the gill membranes broadly united across the isthmus. The unbranched anterior rays form about $\frac{1}{3}$

of the dorsal fin. (Named for Mr. Richard Rathbun, then chief of the Division of Scientific Inquiry in the U. S. Fish Commission, in recognition of his many services to science.)

2628. RATHBUNELLA HYPOPLECTA (Gilbert).

Head $4\frac{2}{3}$ in length; depth 7. D. 46; A. 33. Head and body compressed, elongate, the anterior profile of head compressed, declivous; mouth somewhat oblique, at lower side of snout, small, the maxillary reaching vertical from middle of pupil, $3\frac{1}{6}$ in head; snout very slightly shorter than orbit, $4\frac{1}{3}$ in head; diameter of orbit 4 in head; teeth well developed, in broad bands on jaws, vomer and palatines, the vomer and palatine patches nearly continuous; branchiostegal membranes broadly united, free from isthmus, forming a fold whose depth exceeds $\frac{1}{2}$ diameter of orbit. Margin of preopercle adnate behind, slightly free below, furnished with a series of 6 conspicuous mucous pores; head without spines, ridges or filaments; inner margin of shoulder girdle conspicuously notched above and below, but without hook; gill rakers tubercular, few in number; a well-marked slit behind last gill. Distance from nape to front of dorsal fin equals its distance from posterior border of eye; anterior 10 or 12 dorsal rays simple and apparently not articulate, but flexible and not spine-like; distance from front of anal to base of ventrals $2\frac{1}{3}$ in its distance from base of caudal, all but first $\frac{1}{3}$ of dorsal rays, and all of anal rays forked at tip; dorsal not high, the longest rays $\frac{1}{2}$ head; highest anal ray equals snout and $\frac{1}{2}$ eye; last dorsal and anal rays entirely disconnected from caudal, leaving a free space on caudal peduncle $\frac{1}{3}$ diameter of orbit; ventrals I, 5, in advance of base of pectorals, narrowly triangular, the inner rays longest; pectorals with curved base running backward and downward, the rays all branched, 18 in number, the width of base of fin $3\frac{1}{4}$ in head, the longest ray $1\frac{1}{3}$ in head; caudal rounded, $\frac{3}{5}$ length of head. Body covered with small, partially embedded, cycloid scales, including antedorsal region, belly, breast, and area in front of base of pectorals; cheeks covered with similar but smaller scales, the opercles and rest of head naked. Lateral line running high, parallel with back, on a series of enlarged scales, which are also partly embedded in the thick skin; the lateral line fails to reach base of caudal by a distance equaling $\frac{1}{3}$ of head, and is present on 82 scales. Color, dark olive-brown above, lighter below; a series of about 12 quadrate dark blotches below lateral line, connected more or less by dusky streaks with an alternating series along base of dorsal; no bright colors; dorsal, pectorals, ventrals and branchiostegal membranes dusky straw color; anal black, the rays white tipped; caudal blackish; peritoneum white. A single specimen, 8 inches long, from *Albatross* Station 2944, off Santa Barbara Islands, in 30 fathoms.

Bathymaster hypoplectus, GILBERT, Proc. U. S. Nat. Mus. 1890, 97, off Santa Barbara Islands, California, at Albatross Station 2944. (Coll. *Albatross*.)

Family CXCIII. CHIASMODONTIDÆ.

(The Black Swallowers.)

Body elongate, subcylindrical, or slightly tapering; head subconic. Skin naked; lateral line continuous, placed low; 2 dorsal fins, the first rather short, of slender spines, the second dorsal and anal long; ventrals normal, thoracic, inserted before pectorals, the rays I, 5; pectorals long and narrow; mouth very deeply cleft, reaching beyond the eyes, with numerous long, sharp, movable teeth, the anterior canines movable; teeth on palatines; upper jaw not protractile, the maxillary produced backward. Opercular apparatus very oblique and reduced; no spines or cirri on head; caudal fin forked. Genera 2; species 2; deep-sea fishes, notable for the sharp teeth and for the extensible stomach. (*Chiasmodontidæ*, Gill, in Jordan & Gilbert, Synopsis, 964, 1883.)

a. Jaws with some of the anterior canines extremely long and movable, the 2 anterior crossing each other; lower jaw projecting. Chiasmodon, 847.

aa. Jaws with slender, close-set teeth, none of them greatly produced; lower jaw not prominent. Pseudoscopelus, 848.

847. CHIASMODON, Johnson.

(Black Swallowers.)

Chiasmodon, Johnson, Proc. Zool. Soc. London 1863, 408 (*niger*).
Chiasmodus, Günther, change of spelling.

Body elongate, compressed, and tapering posteriorly, naked; belly pendent, its walls membranaceous, capable of great dilation. Mouth very large; lower jaw longer than upper; each jaw with 2 series of large, pointed teeth, some of the anterior being very large and movable; vomerine teeth none; palatines with teeth similar to those in the jaws. Gills 4. No pseudobranchiæ. Gill openings very wide, the membranes joined to the isthmus for a short distance. Dorsal fins 2; anal single; ventrals inserted below pectorals, each of 5 soft rays. Tail truncate at base of caudal. Caudal forked, free from dorsal and anal. Singular fishes of the deep sea, remarkable for their ability to swallow fishes of many times their own size by means of the great distensibility of the walls of the body. (*χίασμα*, a mark of the form of the letter *χ*; *ὀδούς*, tooth; the 2 anterior canines crossing each other when depressed.)

2629. CHIASMODON NIGER, Johnson.

Head 3½. D. XI-28; A. 27; P. 13; V. 5. Head compressed, elongate, the crown flat, its depth less than ½ its length; maxillary reaching angle of preopercle; both jaws armed with long, pointed, wide-set teeth, nearly all of which are movable; 2 anterior teeth of upper jaw very long, crossing each other when depressed; 3 anterior pairs of teeth in lower jaw

likewise prolonged, the third pair the longest; palatines with a longer, fixed tooth in front. Eye moderate, above the anterior part of maxillary, 4½ in head, shorter than snout, as wide as interorbital space. Lateral line in a longitudinal groove. First dorsal of slender rays, its base 2⅓ in in that of second dorsal; anal commencing behind second dorsal, its anterior rays without connection with vertebral column; posterior rays of anal and dorsal very feeble; pectoral as long as head without snout; ventral ½ as long as pectoral. Color entirely black. Length 12 inches. (Günther.) Deep waters of the Atlantic; a remarkable fish, the walls of the body inordinately extensible; taken at Madeira, in the mid-Atlantic, near the island of Dominica, and off the coast of Massachusetts.* (*niger*, black.)

Chiasmodon niger, JOHNSON, Proc. Zool. Soc. London 1863, 408, Madeira; JORDAN & GILBERT, Synopsis, 964; GOODE & BEAN, Oceanic Ichthyology, 292, 1896.
Chiasmodus niger, GÜNTHER, Cat., v, 435, 1864; CARTER, Proc. Zool. Soc. 1866, 38; GÜNTHER, Challenger Report, Deep Sea Fishes, XXII, 99, 1887.

848. PSEUDOSCOPELUS, Lütken.

Pseudoscopelus, LÜTKEN, Spolia Atlantica, Scopelini, 64, 1892 (*scriptus*).

Body perciform, scaleless, naked; mouth very large; eyes moderate; the slender maxillary reaching far beyond eye; jaws and palate with slender, close-set teeth; ventral fins short, subthoracic, of 1 spine and 5 rays; first dorsal short, of about 8 slender spines; posterior dorsal long, similar to the anal. Each jaw with a distinct line of pores, a median line of pores before ventrals, a cross line connecting ventrals, a series of pores from the vent passing around anal on each side. Lateral line well developed, running high. Head without spines. Gill openings very broad. Pectorals long; caudal short, forked. One species known, in deep water. (ψευδής, false; *Scopelus*.)

2630. PSEUDOSCOPELUS SCRIPTUS, Lütken.

Head 3½; depth 4½. D. VIII–22; A. 22; V. I, 5. Body subfusiform, somewhat compressed. Head large, the snout short and pointed, 4½ in head, the small eye, about 5. Jaws subequal, maxillary 1½ in head; cheek V-shaped, very oblique; bones of head not serrate. Form of head and mouth much as in *Engraulis* or *Scopelus*. Pectoral nearly as long as head, reaching past front of anal; soft dorsal higher than spinous, the anterior rays of soft dorsal and anal elevated. Pores as above described.

* The first specimen of this remarkable fish was obtained at Magdalena (Madeira), at a depth of 312 fathoms, in 1850, by Lowe, who, however, omitted to give a description of it. The species was rediscovered at the same locality by Johnson twelve years later. A third specimen was picked up from the surface, near the island of Dominica. A fourth example was obtained by the *Challenger* in mid-Atlantic, at Station 107, in 1,500 fathoms, on August 26, 1873. A fifth was obtained by the U. S. National Museum from Capt. Thomas F. Hodgdon of the Gloucester schooner *Bessie W. Somers*. It was found on Le Have Bank, floating on the surface, in June, 1880. (Goode & Bean.)

One specimen from Old Bahama Straits. (Lütken.) A singular fish of uncertain relationships, remarkable for the development of mucous pores. (*scriptus*, written.)

Pseudoscopelus scriptus, LÜTKEN, Spolia Atlantica, Scopelini, 64, 1892, Old Bahama Straits.

Family CXCIV. CHÆNICHTHYIDÆ.

Body rather elongated, gradually and regularly declining from the nape to the caudal fin; anteriorly subcylindrical or scarcely compressed. Skin naked or covered with small scales. Lateral line high on the sides and near the dorsal fin. Head moderate or large, with the snout prolonged, depressed, and spatuliform. Crown depressed, not relieved by crests or ridges. Preorbital bones large; suborbital chain very narrow, not articulated with the preopercle. Opercular bones all present, the interopercle and subopercle moderately developed. Mouth terminal, with the cleft lateral and large, extending to the vertical of the eye; upper jaw with its border formed almost entirely by the premaxillaries, whose posterior processes are very short; maxillaries with their articulations entirely posterior to the premaxillaries, slender and gradually enlarged toward their extremities. Teeth on the jaws; palate unarmed. Gill openings wide; gill membranes inferiorly deeply emarginated behind. Branchiostegals 6. Pseudobranchiæ developed. Dorsal fin with its spinous portion short, and usually distinct from the soft, the rays of the latter often simply articulated and not branched; anal fin a little shorter than the dorsal, its rays divided, the membrane notched behind each; caudal fin not forked; pectoral fins well developed, with their inferior rays divided; ventral fins jugular or subjugular, separated by a rhomboid area, each with a spine and 5 rays, the first of which is frequently thickened and entire. Cranium flattened behind, the crests little developed or obsolete. The spatuliform snout is principally formed by the elongated frontal bones. Stomach of moderate size and cæcal. Pyloric cæca in very small number. The chief distinctive characteristic of this family is doubtless the spatuliform extensions of the snout. This, combined with the extent of the fins, structure of the head, and general form, distinguish the group from all others. It appears to be most closely allied to the *Harpagiferidæ* and *Nototheniidæ*. From the former it is separated by the form of the head, as well as by that of the body. From the latter, by the same features, and also by the naked skin. (Gill.) Genera 3 or 4, with about 6 species; inhabiting rather deep waters, mostly in the Tropics. (*Chænichthyoidæ*, Gill, Proc. Ac. Nat. Sci. Phila. 1861, 507.)

a. Body covered with cycloid, deciduous scales; maxillary with a flap; opercle with a dermal flap. HYPSICOMETES, 849.

849. HYPSICOMETES, Goode.

Hypsicometes, GOODE, Proc. U. S. Nat. Mus. 1880, 347 (*goboides*).

Body elongate, subcylindrical, tapering posteriorly. Head very large, much depressed, with snout elongate, spatulate; cleft of mouth very wide,

horizontal, with lower jaw much the longer; the posterior margin of the maxillary wide, free, and with a long cutaneous flap. Eyes very large, close together, subvertical. Scales large, cycloid, deciduous; lateral line conspicuous and continuous, descending abruptly behind pectorals, its scales smaller than those of the body adjoining. Teeth acicular, in bands on the jaws, vomer, and palatines, the largest being upon the palatines, the vomer, and upon 2 pads on either side of the symphysis of the maxillaries. A sharp, short, strong scapular spine. Opercle with 3 feeble, sharp spines, each at the end of a strong feeble ridge; a long, skinny opercular flap extending far beyond the bony portion, and covered with scales. Branchiostegals 6. Gill membranes free from the isthmus, except far in front, where they are united to it, the left-hand flap overlapping the right at the point of junction. Pseudobranchiæ present. Gill rakers short. (*ὕψι*, below, i. e., in deep water; *κωμήτης*, dweller.)

2631. HYPSICOMETES GOBOIDES, Goode.

Head about $2\frac{1}{3}$; depth $7\frac{1}{2}$; orbit $4\frac{1}{2}$ in head, or $1\frac{1}{2}$ in snout. D. VI–15 to 17; A. 16 to 18; V. I, 5; P. 26; scales 65. Mouth very wide, horizontal, the maxillary, which is expanded spoon-like posteriorly, reaching considerably beyond vertical from anterior margin of orbit; eye considerably nearer tip of snout than end of flap, and equidistant between tip of snout and tip of uppermost spine of operculum; entire upper surface of head, cheeks, and opercula covered with scales, except upon bony portion of snout; first dorsal fin placed far forward, not far behind vertical from axil of pectoral; interspace between termination of first dorsal and beginning of second equal to diameter of the orbit, this fin composed of 6 spines, the first and second of which are longest, equal to distance from anterior margin of orbit to tip of lower jaw, and triangular in form; origin of second dorsal almost vertical from that of anal, and terminating a little in advance of the latter; second dorsal fin highest in front and low behind; length of caudal peduncle a little less than length of snout; caudal rounded; pectoral very broad at base, rounded, extending beyond vent and nearly to vertical from origin of anal; lower rays branched; ventrals far apart, horizontal, *Trigla*-like, composed of 1 flexible spine and 5 branched rays, their insertion far forward and far in advance of base of pectorals. Color grayish brown; lighter and yellowish below. Known only from a very small specimen, in which many of the important characters were not discernible. This specimen (No. 26007, U. S. Nat. Mus.) was taken by the *Fish Hawk* from Station 871, in 40° 02′ 54″ N. lat., 70° 23′ 40″ W. lon., at a depth of 115 fathoms, and is much contracted and distorted from immersion in strong alcohol. (Goode.) (*Gobius*; *εἶδος*, resemblance.)

Hypsicometes goboides, Goode, Proc. U. S. Nat. Mus. 1880, 348, lat. 40°, 02′, 54″ N., lon. 70° 23′ 40″ W., in 115 fathoms (Coll. *Fish Hawk*); Jordan & Gilbert, Synopsis, 808, 1883; Goode & Bean, Oceanic Ichthyology, 290, fig. 263, 1896.

Family CXCV. TRICHODONTIDÆ.

(The Sand-fishes.)

Body rather elongate, compressed, naked. Head short, flat on top, the sides vertical. Eyes large, high up, but not superior. Mouth large, almost vertical; lower jaw projecting, its tip entering the profile; lips fringed; premaxillaries protractile; maxillary very broad, without supplemental bone, not slipping under the very narrow preorbital. Teeth moderate, slender and sharp, but not setiform, in bands on jaws and vomer; palatines toothless; inner teeth of jaws depressible. Gill rakers short, slender; gill membranes narrowly united, free from the isthmus. Branchiostegals 5. Gills 4, a slit behind the fourth. Pseudobranchiæ large. Preopercle with 5 prominent spines, the 2 upper directed strongly upward, the 2 lower downward, the middle 1 downward and backward; no barbels; opercle small, strongly striate, unarmed; preorbital with spines; no suborbital stay. Lateral line obsolete. Dorsal fins separate, the first the larger, of numerous slender spines; anal fin elongate, without distinct spines, the rays of anterior third of the fin much shorter than the others, the beginning of the fin below middle of spinous dorsal; pectorals with a very broad, curved, procurrent base; a broad lunate area between pectoral and gill opening, nearly covered by the opercle; soft rays of dorsal, anal, and pectoral fins all simple; ventrals I, 5, close together, thoracic, but behind the pectorals, the middle rays longest; caudal lunate, with many accessory rays, on a slender peduncle. Vertebræ numerous, 48 in typical species. Two genera and 2 species known; from the North Pacific; living in sand near the shore. The fringed lips and other characters indicate the relationship of these fishes with the *Uranoscopidæ*. (*Trachinidæ*, genus *Trichodon*, Günther, Cat., II, 250.)

a. First dorsal long and rather low, of 14 or 15 spines. Trichodon, 850.
aa. First dorsal short and high, of 10 spines. Arctoscopus, 851.

850. TRICHODON (Steller) Cuvier.

(Sand-fishes.)

Trichodon, Steller, in Tilesius, Mem. Acad. St. Petersburg, IV, 1811, 468 (*trichodon*).
Trichodon, Cuvier, Règne Animal, Ed. II, vol. 2, 149, 1829 (*trichodon*).

Characters of the genus included above, the first dorsal long and rather low, of 15 spines. One species. (*θρίξ*, hair; *ὀδούς*, tooth.)

2632. TRICHODON TRICHODON (Tilesius).

(Sand-fish.)

Head from tip of upper jaw, 3¼; depth 3¼. D. XIII-I, 18; A. 28; P. 22; eye 4⅓ in head, snout 4⅓; maxillary 2; interorbital 3; pectoral 1⅛; ventral 1⅔; height of spinous dorsal 3⅛. Body moderately elongate, compressed; dorsal outline slightly concave and sloping gently upward from snout to dorsal, thence turning at a very slight angle nearly straight to caudal;

ventral outline well rounded from chin to caudal peduncle, the curve much more gradual posteriorly; head and body everywhere covered with thin naked skin. Mouth large, superior, nearly vertical, the lower jaw projecting, its tips entering the profile; lips fringed; maxillary reaching to middle of pupil; teeth in 2 or 3 rows, small, sharp and recurved; teeth on vomer; palatines toothless. Eyes placed high, their diameter equal to length of snout; interorbital wide and flat, a third wider than eye; top of head smooth, sometimes rugose in younger individuals, covered with thin smooth skin; anterior nostril ending in a tube; preopercle with 5 spines, the 1 at angle largest, the 2 upper ones pointing upward and backward, the middle one pointing downward and backward, the 2 lower ones pointing downward and forward; opercle with radiating ridges; gill rakers short and slender, numerous. Origin of spinous dorsal behind base of pectoral, its distance from snout 3 in body, the spines not varying greatly in length, the last one connected by a membrane to the back; soft dorsal well separated from spinous, its rays about equal to spines in length, highest in front; anal long, its origin nearer to the snout than base of caudal by a distance equal to the length of the eye. Pectoral, when spread, broadly rounded behind, its lower rays rapidly decreasing in size below, reaching well past front of anal; ventrals inserted behind base of pectorals a distance equal to ⅔ eye, their tips reaching to vent. Lateral line running high. Vertebræ $17 + 30 = 47$. Color silvery, light brown above; a dark brown streak following the lateral line, broken up into spots anteriorly; quadrangular, dark brown marks along the back at base of dorsals, chain-like markings in front of dorsal on nape; snout and tip of lower jaw dark; a dark line at lower part of eye; dorsals light, a dark streak along upper part of spinous dorsal; pectorals dusky; ventrals and anal colorless. Length 8 to 10 inches. North Pacific, on sandy shores, from Bering Sea to Monterey; very abundant northward; burying in the sand. Here described from a specimen, 8½ inches in length, from Herendeen Bay, Alaska (*Albatross* collection). Possibly detailed comparison may show a difference between California specimens and those from Bering Sea.

Trachinus trichodon, Tilesius,* Mem. Acad. St. Petersburg, iv, 1811, pl. 15, fig. 8, 473, Kamchatka; Pallas, Zoographia Rosso-Asiatica, iii, 235, 1811.†

* The specific name *trichodon* should apparently date from Tilesius, 1811. Although Vol. iv, of the Mem. Acad. St. Petersburg bears the date 1813 it was for the year 1811, and it is evident that the plate containing the figure of this species was accessible to Pallas as early as 1811, for, in his "Zoographia," printed in 1811, though not published until 1831, Pallas refers to the plate of Tilesius in very definite terms. The fact that Pallas was, in 1811, thus able to refer definitely to Tilesius's plate of *Trachinus trichodon*, fixes the date of publication of that plate at least as early as 1811. That this plate appeared in the volume of Memoirs for 1811 (though the volume was not published until 1813), fixes 1811 as the date for the name. Though the "Zoographia" of Pallas was not formally published until 1831, it was printed in 1811, and Cuvier & Valenciennes evidently had a copy in 1829, as they refer to it.

† Tilesius confused matters greatly by using, in one and the same article, three different names or combinations of names for this fish. At the beginning of this article (p. 406) in a bald list of the species discussed in the paper, he has "*Drachinus trichodon*." On page 466 he has "*Trachinus gasteropelecus*," accompanied by a full description of the species. In a footnote on page 473, he has "*Trachinus trichodon*" together with a description which he says applies to the young, and finally his pl. 15, fig. 8, is marked "*Trachinus trichodon*."

Drachinus trichodon, TILESIUS, Mem. Acad. St. Petersburg, IV, 1811, 406; name only.
Trachinus gasteropelecus, TILESIUS, *l. c.*, 466, 1811, Kamchatka.
Trichodon stelleri, CUVIER & VALENCIENNES, Hist. Nat. Poiss., III, 154, pl. 57, 1829; based on *Trachinus trichodon* PALLAS; GÜNTHER, Cat., II, 251, 1860; JORDAN & GILBERT, Synopsis, 627, 1883.
Trichodon lineatus, AYERS, Proc. Ac. Nat. Sci. Phila. 1860, 60, San Francisco; D. XV–18; A. 28; P. 23.

851. ARCTOSCOPUS, Jordan & Evermann.

Arctoscopus, JORDAN & EVERMANN, Check-List Fishes, 464, 1896 (*japonicus*).

This genus differs from *Trichodon* in the short, high, triangular spinous dorsal which is composed of 10 spines. (ἄρκτος, northern; σκοπός, gazer; for *Uranoscopus*.)

2633. ARCTOSCOPUS JAPONICUS (Steindachner).

Head $3\frac{3}{4}$; depth $3\frac{3}{4}$. D. X or XI–13; A. 30 or 31; P. 25. Form of body and coloration of *Trichodon trichodon*. First dorsal high, triangular, the spines slender, separated by a long interval from the second dorsal. Preopercle with 5 sharp spines; the 2 spines on the preorbital very small. Pectoral well developed, all its rays simple, the lower a little thickened, the fin considerably longer than the head and reaching past the last spine of the dorsal; anal fin with its rays gradually longer posteriorly. Dentition as in *Trichodon trichodon*, but the mouth rather more oblique. Length $4\frac{1}{2}$ inches. North Pacific; scarce. Recorded from Strietok, in the Sea of Japan, and Sitka, Alaska, by Steindachner, and by Jordan & Gilbert from Iturup Island (Kurils). (*japonicus*, from Japan.)

Trichodon japonicus, STEINDACHNER, Ichth. Beitr., X, 4, 1881, Strietok; Sitka; JORDAN, Cat. Fishes N. A., 117, 1885.
Arctoscopus japonicus, JORDAN & GILBERT, Rept. Fur Seal Investig., 1898.

Family CXCVI. DACTYLOSCOPIDÆ.

(THE SAND STAR-GAZERS.)

Body oblong, low, compressed posteriorly, covered with moderate, cycloid, imbricated scales; lateral line complete, anteriorly running along side of back, posteriorly median; head oblong, nearly plane above; eyes small, superior, well forward; suborbital bones enlarged, but without bony stay connecting with the preopercle; nostrils double; opercles fringed; mouth nearly vertical; premaxillaries protractile, not forming the entire edge of the upper jaw; lips fringed as in *Uranoscopidæ;* gill openings very broad, the membranes separated and free from the isthmus, pseudobranchiæ present or obsolete. Dorsal fin very long, continuous or divided, several of the anterior rays spinous; anal very long, commencing close behind the vent, which is near the breast; caudal diphycercal, free from dorsal and anal; pectorals variable, the base broad and procurrent; ventrals jugular, I, 3; vertebræ more than 10 + 14; pyloric cæca none. Genera 4; species about 10; small fishes living on sandy shores of tropical

America. This family is nearly related to *Uranoscopidæ*, of which group it seems to be a reduced or degenerate branch. Its relations with the Asiatic family *Leptoscopidæ* are most intimate, the incomplete ventrals and simple pectoral rays of *Dactyloscopidæ* being the chief distinctive features. (*Dactyloscopidæ*, Gill, Arrangm. Families Fishes, 1872.)

a. Dorsal fin divided, the first dorsal composed of 3 spines inserted on the nape; head not cuboid; chin without flap; fringes of lips small. GILLELLUS, 852.

aa. Dorsal fin continuous.

b. Dorsal fin commencing at the nape; pseudobranchiæ very small or obsolete; head cuboid. DACTYLOSCOPUS, 853.

bb. Dorsal fin commencing far behind the nape; pseudobranchiæ well developed.

c. Head cuboid, formed as in *Dactyloscopus;* the mouth vertical. DACTYLAGNUS, 854.

cc. Head elongate-conoid, the lower jaw projecting, with a fleshy flap at tip. MYXODAGNUS, 855.

852. GILLELLUS, Gilbert.

Gillellus, GILBERT, Proc. U. S. Nat. Mus. 1890, 98 (*semicinctus*).

A separate dorsal fin on the nape composed of 3 spines. Lateral line descending posteriorly, its dorsal and median portions about equal. Fringes of upper lip obsolete, those of lower lip little evident. Head not cuboid, the mouth moderately oblique, the lower jaw rounded in front and without symphyseal flap. The physiognomy is intermediate between *Dactyloscopus* and *Myxodagnus*, from each of which the genus is well separated by the characters of the dorsal fin and the lateral line. ("Named in honor of Dr. Theodore Gill, to whom we owe our knowledge of the previously described members of this most interesting group." Gilbert.)

a. Tip of lower jaw projecting.

b. Anterior portion of lateral line longer than posterior portion; the scales 25 to 28 + 3 + 15 to 18 = 43 to 49. D. III-IX, 28; A. II, 30 or 31. SEMICINCTUS, 2634.

bb. Anterior portion of lateral line much shorter than posterior portion, 2⅓ times in the latter; scales 18 + 3 + 27 = 48. D. II-IX, 31; A. II, 35. ARENICOLA, 2635.

aa. Tip of lower jaw scarcely projecting; anterior portion of lateral line 1¼ times in posterior. D. I-IX, 31; A. II, 34. ORNATUS, 2636.

2634. GILLELLUS SEMICINCTUS, Gilbert.

Head 3⅔; depth 5⅓. D. III-IX or X, 28; A. II, 30 or 31; scales 25 to 28–3–15 to 18 (43 to 49 scales in all). Body deep, tapering rapidly either way from front of dorsal. Mouth moderately oblique, the maxillary extending beyond orbit, 3 in head; tip of lower jaw projecting; teeth in a narrow band in front of jaws, becoming a single series laterally; none of the teeth enlarged. Opercular fringes well developed, 8 or 9 in number; fold of membrane between rami of lower jaw well developed; pseudobranchiæ apparently not developed; gill rakers obsolete. Dorsal fin beginning at a distance from occiput less than diameter of eye, the first 3 rays entirely detached from the rest of the fin, the first ray the highest, the second and third shortened; of the remaining part of the fin the first 9 or

10 rays are unarticulated and spinous; first 2 anal rays not articulated; caudal about $1\frac{4}{5}$ in head; pectorals $1\frac{1}{3}$. Lateral line running anteriorly along the very base of spinous dorsal, no scales intervening between it and base of fin; it descends to middle of sides posteriorly, the median portion of its length shorter than the dorsal portion. Color light olivaceous, the back with 6 broad cross bars of pink, narrowly margined behind and in front with blackish, terminating below on middle of sides; the lower of these bars frequently black; a black bar across caudal peduncle, and sometimes a black line at base of caudal; along median line of sides frequently a series of small black spots alternating with the cross bars; a similar series along median dorsal line; a large pink blotch covering occiput; a dusky bar across interorbital space, running downward and backward across cheek; silvery spots and blotches on cheeks and anterior portions of opercles; fins unmarked. Specimens have been obtained in the Gulf of California by the *Albatross*, at Stations 2827 and 2829, and by the *Grampus* in the Atlantic, at Stations 5108 and 5112, off the coast of Florida; no specific difference among them noticed, but the Atlantic form needs further study. (Gilbert.) (*semi*, half; *cinctus*, belted.)

Gillellus semicinctus, GILBERT, Proc. U. S. Nat. Mus. 1890, 98, Albatross Stations 2827, 2829, Gulf of California (Coll. *Albatross*); JORDAN, Proc. Cal. Ac. Sci. 1896, 229, pl. 32.

2635. GILLELLUS ARENICOLA, Gilbert.

Head $4\frac{2}{3}$ in length; depth $8\frac{3}{4}$. D. II–IX, 31; A. II, 35; scales 18–3–27. Body very slender and elongate, much as in *Myxodagnus*, the snout sharp, the mandible produced at symphysis and conspicuously projecting; labial fringes apparently obsolete; maxillary reaching vertical from middle of orbit; eye small, about equaling length of snout, 6 in head; opercular fringes nearly obsolete, 3 or 4 small ones at upper edge of opercle. Anterior dorsal inserted close behind occiput, composed of 3 rays, and separated by a short interspace from rest of fin; pectorals longer than head. Lateral line anteriorly running along base of dorsal, from which it is not separated by intervening scales, the anterior portion contained $2\frac{1}{3}$ times in the posterior median portion. Color light olivaceous, the head with grayish blotches and small pearly spots; 11 dark bars downward from back, the alternate ones narrower and fainter and not extending to middle of sides, as do the others; the margins of the larger bars darker than the median portion, the bars not continued onto dorsal fin; all the fins translucent. A single specimen $1\frac{1}{2}$ inches long, from Cape San Lucas. (Gilbert.) (*arena*, sand; *colo*, I inhabit.)

Gillellus arenicola, GILBERT, Proc. U. S. Nat. Mus. 1890, 99, Cape San Lucas. (Coll. Gilbert.

2636. GILLELLUS ORNATUS, Gilbert.

Head $4\frac{1}{3}$ in length; depth 8. D. III–IX, 31; A. II, 34; scales not counted. With the elongate form and general appearance of *Gillellus arenicola*, but differing in the subequal jaws and in the long anterior portion of the lateral line. Head conical, acute, very small; jaws nearly equal, the lower slightly longer than the upper, but not noticeably protruding. In this

respect the species resembles most strongly *G. semicinctus*, from which it varies widely in the general form and proportions. Snout extremely short, scarcely equaling diameter of the minute eye; diameter of orbit about 7 in head. Mouth oblique, the maxillary 4 in head, reaching nearly to vertical from posterior margin of orbit. Lips without fringes. Eyes separated by a narrow septum, the interorbital width being less than the diameter of the pupil. Opercular fringes few and small, flat, and not terminating evident ridges as in *Dactyloscopus*. Dorsal beginning well forward, its origin less than diameter of orbit behind the posterior line of occiput; anterior detached part of fin consisting apparently of 3 rays, the first of which is the longest, the second and third equal and short; fourth spine again longer; spines as usual slenderer than the rays, and showing no articulations, but with some difficulty discriminated from them; pectoral as long as head. Anterior part of lateral line running immediately along base of dorsal, without intervening scales, as in other members of this genus. It is much longer than in *G. arenicola* and is contained 1¼ times in the posterior median portion. There are 3 scales between the posterior part of the lateral line and the base of the dorsal. Color similar to that of *G. arenicola* and *G. semicinctus*, light olivaceous, unmarked below the middle of the sides, the back and upper half of sides with 8 brown bars which extend downward to lateral line; the upper part of each bar with a lighter central area, the light areas between the bars marked more or less with brown, which sometimes forms indistinct secondary bars; a blackish bar at base of caudal, and a faint streak below eye; a large pearly blotch on opercle. A single specimen, about 2 inches long, from *Albatross* Station 2828 in the Gulf of California. (Gilbert.) (*ornatus*, adorned.)

Gillellus ornatus, GILBERT, Proc. U. S. Nat. Mus. 1891, 558, Gulf of California. (Coll. Gilbert.)

853. DACTYLOSCOPUS,* Gill.

Dactyloscopus, GILL, Proc. Ac. Nat. Sci. Phila. 1859, 132 (*tridigitatus*).
Esloscopus, JORDAN & EVERMANN, Check-List Fishes, 465, 1896 (*zelotes*).

Body moderately elongate, covered with rather large, cycloid scales; head cuboid, oblong, and nearly flat above; eyes small; interorbital space broad; mouth nearly vertical; lower jaw not dilated beneath nor emarginate in front, without barbels; no intralabial filament; teeth villiform, on jaws only; pseudobranchiæ very small or obsolete. Dorsal commencing at the nape, with 6 or 12 slender spines, the soft rays numerous; anal

*This genus is thus defined by Dr. Gill: "Body elongate with the dorsal and abdominal outlines slowly converging to the caudal fin. Scales large, regularly imbricated. Lateral line straight, and running along the middle of the side. Head oblong, subcubical, and smooth. Preopercle entire, opercle radiately fringed behind. Mouth nearly vertical. Tongue thick, narrowed anteriorly, attached to the floor of the mouth. Labial velum without a barbel. Anus a short distance behind the base of the pectoral fins. Dorsal fin subequal, single, and very long, commencing above or before the anus and continued almost to the base of the caudal. Anal fin commencing behind the anus, and with the same form and termination as the dorsal. Caudal fin small and narrow, posteriorly subtruncated. Pectoral fins subangular. Ventral fins jugular, closely approximated, and each with 3 stout simple and articulated rays."

inserted behind dorsal; ventral rays I, 3. (δάκτυλος, finger; σκοπός, gazer, short for *Uranoscopus*.)

DACTYLOSCOPUS:
a. Dorsal rays X to XII, 22 to 31; anal rays less than 35.
b. Soft dorsal with 22 soft rays; anal with 26. PECTORALIS, 2637.
bb. Soft dorsal with 28 to 31 rays; anal with 32 or 33; scales about 45.
c. Body rather slender, the depth about 6 in length (7 with caudal); opercular fringe of 15 filaments. TRIDIGITATUS, 2638.
cc. Body rather stout, the depth 5¼ in length (6 in total with caudal); opercular fringe of 18 filaments.
d. Back not barred; head blotched and dotted. POEYI, 2639.
dd. Back with about 10 pale cross bars; head marked with whitish; a dark bar at base of caudal. LUNATICUS, 2640.
ESLOSCOPUS (ἐσλός, good; σκοπός for *Uranoscopus*):
aa. Dorsal rays VI, 38; anal rays II, 37; scales 6–51–5. ZELOTES, 2641.

Subgenus DACTYLOSCOPUS.

2637. DACTYLOSCOPUS PECTORALIS, Gill.

Head about 5 in total length with caudal; depth about 7 (in total). D. XII, 22; A. II, 26; P. 12; V. I, 3. Width of head behind operculum 7 in total length with caudal; eye small, 10 in head; interorbital space equals diameter of eye; preoperculum broader at the angle than in *Dactyloscopus tridigitatus;* pores well developed; opercular fringe of 11 or 12 free filaments; origin of dorsal between ⅓ and ⅙ length of fish from tip of snout; origin of anal under sixth or seventh dorsal ray, the first 12 dorsal and 2 anal rays simple. Pseudobranchiæ obsolete. Color light brownish yellow, with dark spots on the back, arranged in lines forming the outlines of about 6 quadrangular areas, from the angles of which irregular lines proceed downward, converging toward those departing from the angles of adjoining areas; more scattered and irregular spots and dots often present below the lateral line; head lighter, diffused with pink above. Each orbit with 4 diverging bands, 1 in front, a bifurcated one from the antero-inferior angle, and 2 from posterior border, a transverse sinuated nuchal line; upper angle of operculum whitish, bounded in front by a dark line or spot. (Gill.) Cape San Lucas; not seen by us. (*pectoralis*, pertaining to the breast.)

Dactyloscopus pectoralis. GILL, Proc. Ac. Nat. Sci. Phila. 1861, 267, Cape San Lucas. (Coll. John Xantus.)

2638. DACTYLOSCOPUS TRIDIGITATUS, Gill

Head 5 (in total) with caudal; depth 7. D. XII, 28; A. II, 32; P. 13; V. I, 3; scales 11 + 4 + 30 = 45. Body slender, much compressed posteriorly; opercular fringe of 15 separate filaments. Origin of dorsal fin over the lower angle of the base of the pectorals, or immediately before the margin of the operculum, its distance from snout to dorsal 5 in total length of body. Pseudobranchiæ very small (overlooked by Dr. Gill, but evident in living specimens). In life, pale sand color above, the lower part whitish; above 12 narrow cross bands of whitish on the back, not

extending down far on the sides; head mottled above; fins all pale. West Indies, north to Key West; rather common in coral sand in shallow water about Key West. (*tres*, three; *digitus*, finger, from the 3 ventral rays.)

Dactyloscopus tridigitatus, GILL, Proc. Ac. Nat. Sci. Phila. 1859, 132, Barbados (Coll. Dr. Gill); GILL, *l. c.*, 1861, 264; GILL, *l. c.*, 1862, 505; GÜNTHER, Cat., III, 279, 1861; JORDAN & GILBERT, Synopsis, 753, 1883; JORDAN, Proc. U. S. Nat. Mus. 1884, 140.

2639. DACTYLOSCOPUS POEYI, Gill.

Head 5 in total length; depth $6\frac{1}{4}$ in total. D. XI, 31; A. II, 32. Body more robust than in *D. tridigitatus;* head plane above and obtusely angulated at the sides of the plane; thickness of the head behind the preoperculum exceeding $\frac{1}{2}$ of its length; interorbital space $\frac{3}{8}$ diameter of eye. Eye about 7 in head; preopercle as in *D. tridigitatus*, pores indistinct or obsolete; opercular fringe of about 18 filaments, the lowest of which are scarcely extended beyond the margin; origin of dorsal fin $\frac{1}{6}$ distance from tip of snout; origin of anal fin under sixth dorsal ray; scales of moderate size and regularly imbricated. Color reddish brown, dotted with darker above the lateral line; head blotched and dotted with darker; opercles variegated; opercular bones nearly immaculate. (Gill.) Cuba; not seen by us. (Named for Prof. Felipe Poey.)

Dactyloscopus poeyi, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 266, Cuba. (Coll. Felipe Poey.)

2640. DACTYLOSCOPUS LUNATICUS, Gilbert.

Head (to end of opercular fringes) $3\frac{3}{8}$, from tip of lower jaw to base of fringes 4; depth greater than in related species, $5\frac{1}{4}$ in length. D. X or XI, 29 or 30; A. II, 32 or 33; scales about 11 + 4 + 30 = about 45. Head cuboid, narrowed forward, the vertex gently convex; width at occiput $\frac{1}{2}$ length of head (to base of fringes on opercle). Mouth nearly vertical, maxillary $2\frac{3}{8}$ in head. Labial fringes short but evident. A short nasal filament. Teeth in a rather broad cardiform band on front of upper jaw, becoming narrow laterally; in lower jaw a single series, or an irregular double series anteriorly; vomer and palatines toothless. Eyes small, very close together, the interorbital width about $\frac{1}{2}$ their diameter, which equals length of snout, or about $\frac{1}{3}$ head. Gill laminæ much reduced in size; a small round pore behind inner arch. Gill rakers obsolete; pseudobranchiæ small but evident. Opercular fringes composed of 18 filaments. Dorsal beginning at a distance behind occiput equaling diameter of orbit, its anterior rays but partly joined by membrane, the first 10 or 11 slender and not articulated, the last ray distant from base of caudal about a diameter of orbit; origin of anal under sixth dorsal spine, the 2 anterior rays not articulated; pectorals short, $1\frac{3}{8}$ in head, containing 14 or 15 rays; caudal very small, with 10 developed rays, its length $2\frac{3}{4}$ in head. Lateral line running high in its anterior portion, declining on 3 or 4 scales, the posterior portion on middle of sides with 29 or 30 tubes; 4 scales between median portions of lateral line and base of dorsal. Color light olivaceous, a dark streak along back, 1

along middle of sides, and a fainter one along base of anal, formed by darker margins to the scales; median dorsal line with 10 or 11 more or less evident narrow pearly white cross bars; top of head and front of mandible colored like the back, the pearly blotches varying in size and shape, but symmetrically arranged, many of them narrowly edged with black; nasal tentacle white; white streaks on preopercle; caudal with a narrow black bar at base. Gulf of California. Three specimens, the longest 3 inches, from *Albatross* Stations 2797 and 3012, the latter in 22 fathoms. (Gilbert.) (*lunaticus*, moon-struck.)

Dactyloscopus lunaticus, GILBERT, Proc. U. S. Nat. Mus. 1890, 99. Gulf of California. (Coll. *Albatross.*)

Subgenus **ESLOSCOPUS**, Jordan & Evermann.

2641. DACTYLOSCOPUS ZELOTES, Jordan & Gilbert, new species.

Head 4¼ in length; depth 6⅔. D. VI, 38; A. II, 37; V. 3; scales 6–51–5; B. 6. Head and body slender, compressed, the greatest width at occiput, ⅔ length of head; the greatest depth immediately behind insertion of anal fin, thence tapering to a very narrow tail. Head narrow, cuboid, compressed, the upper surface nearly plane, the cheeks vertical. Eyes very small, superior, with little lateral range; diameter of orbit about ⅓ length of head; snout very short, about equaling orbit; anterior nostril in a short tube; gape subvertical, the lower jaw very heavy, projecting, as in *Uranoscopus;* premaxillaries protractile, the processes reaching far behind orbits; lips fringed; both jaws with bands of villiform teeth; no teeth on tongue, vomer, or palatines. Subopercle and interopercle very wide, flexible, striate, the latter overlapping throat and base of ventral fins, the former wholly covering base of pectoral fins; the striations of opercle terminate posteriorly in a wide, coarse, membranaceous fringe; branchiostegal membranes not united, free from the isthmus; pubic bones forming a sharp projection at throat; no pseudobranchiæ; gills small, a round pore behind the fourth. Dorsal beginning on the nape, its distance from snout about equaling depth of body, the first 6 rays shorter than those following and not connected by membrane; as no traces of articulation can be found, they are probably flexible spines, but are not clearly differentiated from those immediately following; origin of anal under fourth dorsal spine; caudal distinct, narrow, short; ventrals inserted under anterior margin of preopercle; ventrals 2 in head; pectorals 1¼. Scales large, with entire edges, wanting on head, breast, and region behind pectoral fins. Lateral line beginning at upper posterior angle of opercle, running parallel with the back on about 12 scales, then obliquely downward to middle of body. Color in spirits, light olivaceous, the edgings of the scales, some vermiculations on top of head, and the labial fringes clear brown; fins translucent, the caudal with a brown bar at base; eyes dark. Length 3¼ inches. Panama; 1 specimen known. The present description copied from the original in Proc. Nat. Mus. 1882, 628. (ζηλώτης, an imitator, from its resemblance to *Dactylagnus mundus.*)

Dactyloscopus, sp. nov., JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 628, Panama.
Dactyloscopus zelotes, JORDAN & GILBERT, new species (MS. 1882), Panama (Coll. Capt. Dow).

854. DACTYLAGNUS, Gill.

Dactylagnus, Gill, Proc. Ac. Nat. Sci. Phila. 1862, 505 (*mundus*).

Body moderately elongated, covered with rather large and uniform scales. Head cuboid, oblong, scarcely convex transversely above. Eyes small, directed obliquely upward, and situated near the snout on the upper surface of the head. Interorbital area moderate and channeled. Mouth very oblique or subvertical, the snout truncated in front; lower jaw transversely convex in front and with no barbel; teeth acute, in a narrow band along each jaw; palate smooth. Dorsal fin perfectly entire, commencing rather farther behind than the anal, and with its anterior portion armed with about 10 slender spines; anal fin longer than the dorsal. This genus closely resembles *Dactyloscopus* externally. It differs from the latter genus chiefly in the structure of the dorsal fin and the well-developed pseudobranchiæ. (δάκτυλος, finger; ἄγνος, *Agnus*, an old name of *Uranoscopus scaber*.)

2642. DACTYLAGNUS MUNDUS, Gill.

Head 4⅓; depth 6⅓. D. X, 31; A. II, 38; scales 2–48–10; eye 6 in head; maxillary 2⅔; snout equals eye; highest dorsal spine 3; highest anal ray 2½; pectoral equals head; caudal 1⅚. Body elongate, compressed, tapering posteriorly; upper profile of head nearly horizontal, slightly convex; eyes superior, looking upward; interorbital narrow, concave; lower jaw strongly projecting, mouth nearly vertical; teeth small and conical, in narrow bands, widest in front; vomer and palatines toothless; lips furnished with labial fringes about as long as diameter of eye; nostril ending in a tube; preopercle entire; opercle fringed on its upper edge, a flap of skin downward from opercle covers the branchiostegals; pseudobranchiæ present; gill rakers not developed; head and belly naked; fins naked. Lateral line running near the back through 14 scales, deflected on 4, and thence continued along the middle through 36. Dorsal low, long, and continuous, distance from its origin to tip of snout 3¾ in body; anal similar, slightly higher and longer; posterior rays of dorsal and anal reaching to base of caudal rays; upper rays of pectoral the longest, reaching to the vertical from tenth anal ray, the lower rays short, graduated, tip of fin slightly curved up; origin of ventrals in front of pectorals, the inner rays the longest, reaching about to vent; caudal truncate, or very slightly rounded. Color in spirits, light brown above, white below, each scale on back with a dark brown spot; top of head with a few brown spots; fins colorless. Length 4½ inches. Gulf of California. Here described from specimens collected by the *Albatross* at Carmen Island, Gulf of California; the type from Cape San Lucas. (*mundus*, neat.)

Dactylagnus mundus, Gill, Proc. Ac. Nat. Sci. Phila. 1862, 505, Cape San Lucas. (Coll. Xantus.)

855. MYXODAGNUS,* Gill.

Myxodagnus, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 269 (*opercularis*).

This genus differs from *Dactyloscopus* in the form of the head, which is elongate-conoid, the lower jaw obtusely pointed and provided with a short flap in front. The pseudobranchiæ are well developed and the dorsal fin commences far behind the nape. One species known. (*Myxodes*, a genus of blennies, which this fish resembles in form; *Agnus*, ἄγνος, an old name of *Uranoscopus scaber*.)

2643. MYXODAGNUS OPERCULARIS, Gill.

Head 5 without lower jaw; depth 7. D. 36; A. II, 36; scales 2–44–9; pectorals equal head; ventrals 1¾; caudal 1½. The body is deepest at front of dorsal fin, tapering regularly to the caudal fin. Head elongated, acutely conical; profile nearly straight, but slightly concave in front of the eyes; the crown is transversely arched and smooth; the frontal bones between the eyes are exceedingly narrow, so that the orbits appear separated by little more than a mere septum; eyes large, longitudinally elliptical; opercular pores obsolete; the postorbital or temporal ridge is nearly as long as the diameter of the orbit; the opercular fringe is composed of 6 or 7 short filaments; origin of dorsal above vent, the fin very low and continuous, its last rays not reaching to base of caudal rays; anal commencing slightly in front of dorsal, similar to it but higher, its last ray reaching to base of caudal rays; pectoral large and pointed, reaching to curve in lateral line; rays of ventral subequal, reaching about to vent; caudal truncate; scales moderate, finely striated concentrically and arranged in 11 rows on each side; the lateral line runs through 12 scales on the sides of the back, is then deflected through 3, and thence runs along the fifth row from the back through 36. Color light yellowish brown, rendered darker on the back by congregations of dark spots on the scales; there is a pearly patch behind and beneath the eye, and the operculum is also colored in the same manner. (Gill.) Cape San Lucas. Described from a specimen 2½ inches in length. Not obtained by recent collectors. (*opercularis*, pertaining to the gill cover.)

Myxodagnus opercularis, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 270, Cape San Lucas. (Coll. Xantus.)

Family CXCVII. URANOSCOPIDÆ.

(THE STAR-GAZERS.)

Head large, broad, partly covered with bony plates. Body elongate, conic, subcompressed, widest and usually deepest at the occiput. Body either naked or covered with very small, smooth, adherent scales, which

* This genus is thus defined by Dr. Gill: Body quite slender, the greatest height contained about 10 times in length. Head rather elongated and acutely conical, about twice as long as high; eyes large and elliptical, and very closely approximated; frontal bones extremely narrow. Mouth oblique; lower jaw projecting much beyond the upper and furnished with a short, compressed, and wide flap or barbel in front of the symphysis; villiform teeth present only on the jaws. Dorsal fin inserted behind the vertical of the anus, and furnished with simple and articulated rays; anal fin as long as or longer than the dorsal.

are arranged in very oblique series running downward and backward; the scales on the belly inconspicuous or obsolete. Lateral line little developed, running high. Eyes small, on anterior and upper portion of head, with vertical rings. Mouth vertical, with strong and prominent mandible; teeth moderate, on jaws, vomer, and palatines. Premaxillaries freely protractile; maxillary broad, without supplemental bones, not slipping under the preorbital. Gill openings wide, continued forward; gill membranes nearly separate, free from isthmus. Pseudobranchiæ present; 6 branchiostegals; 3½ gills, a slit behind the last; no anal papilla. Spinous dorsal very short or wanting; second dorsal long. Anal and pectorals large, the latter with broad oblique bases, the lower rays rapidly shortened, most of them branched; ventrals jugular, close together, I, 5, the spine very short, innermost ray longest; caudal not forked. Air bladder generally absent; pyloric cæca in moderate number. Vertebræ 24 to 26. Carnivorous fishes, living on the bottom of the shores of most warm regions. Genera 8; species 25.

URANOSCOPINÆ:

a. Spinous dorsal of 4 or 5 well-developed spines; scales present.

b. Head above not entirely covered with bone, the occipital plate ceasing much behind the orbits; from the middle line anteriorly a Y-shaped bony process extends forward, the tips of the fork between the eyes; a trapezoidal space on either side of the Y, covered by naked skin, bounded by the Y, the eyes, the suborbitals, and the occipital plate. A covered furrow behind and on the inner side of each eye terminating near front of orbits, its edges fringed. Head without spines; humeral spine obsolete; lips and nostrils fringed; no retractile tentacle in mouth. ASTROSCOPUS, 856.

KATHETOSTOMATINÆ:

aa. Spinous dorsal obsolete; no scales; head above covered with bone except the groove of the premaxillary spine; the bony occipital plate coalescing with the orbital rims; humeral spine well developed; no distinct protuberances on top of head; no spine in front of humeral spine; 2 small forward-directed spines in front of eye; 3 small spines on lower margin of preopercle; upper lip scarcely fringed; no retractile tentacle in mouth; ventral fin not largely adnate to abdomen. KATHETOSTOMA, 857.

856. ASTROSCOPUS, Brevoort.

(ELECTRIC STAR-GAZERS.)

Astroscopus (BREVOORT) GILL, Proc. Ac. Nat. Sci. Phila. 1860, 20 (*anoplos;* young).

Agnus, GÜNTHER, Cat. Fishes, II, 229, 1860 (*anoplos*).

Upselonphorus, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 113 (misprint for *Upsilonphorus*) (*y-græcum;* adult).

Body robust. Head above not entirely covered with bone, the occipital plate ceasing much behind the orbits; from the middle line anteriorly a Y-shaped bony process extends forward, the tips of the fork between the eyes; a trapezoidal space on either side of the Y, covered by naked skin, bounded by the Y, the eyes, the suborbitals, and the occipital plate. A covered furrow behind and on the inner side of each eye terminating near front of orbits, its edges fringed. Head without spines; humeral spine obsolete; lips and nostrils fringed; no retractile tentacle in mouth. Young individuals with top of head largely covered by bone. Head scaleless; back and sides covered with close-set scales; belly mostly naked. Humeral

spine obsolete; no spine before the ventrals. First dorsal small, of 4 or 5 low, stout, pungent spines, connected by membrane to the second dorsal, which is rather high and long; pectorals and ventrals large. Species American, distinguished from the Old World genus, *Uranoscopus,** chiefly by the unarmed head. (*ἀάστρον*, star; *σκοπέω*, to look.)

a. Naked space between forks of the Y on top of head long and narrow, but shorter than the vertical limb of the Y; no distinct spines before eye; sides with round pale spots, each with a dark ring.

b. Dorsal spines 4, rather high; scales normal. Y-GRÆCUM, 2644.

bb. Dorsal spines 5, lower than in *y-græcum;* scales of sides cohering in oblique series. ZEPHYREUS, 2645.

aa. Naked space between the forks of the Y short and broad, but longer than the very short vertical limb of the Y; 2 distinct spines directed forward before the eye; sides with small pale spots, not dark-edged. GUTTATUS, 2646.

2644. ASTROSCOPUS Y-GRÆCUM (Cuvier & Valenciennes).

Head, without lower jaw, $2\frac{5}{6}$; depth $3\frac{1}{3}$. D. IV–I, 12; A. 13; scales 80; eye $12\frac{1}{2}$ in head; maxillary 2; pectoral $1\frac{1}{10}$; second dorsal spine 4; highest dorsal ray 2; highest anal ray 3; caudal $1\frac{1}{3}$. Body moderately elongate, very robust forward, greatest depth at occiput; anteriorly subcylindrical, posteriorly somewhat compressed. Head large and broad; mouth large, vertical, a fringe of barbels on each jaw, slightly longer than the diameter of the eye; tongue extremely large and fleshy, forming a pad under membrane of lower jaw which projects forward somewhat. Teeth conical, small and movable, in many bands in upper jaw, in lower jaw the teeth are larger and in fewer bands; teeth on vomer and palatines. Eyes very small but prominent, set on top of head; interorbital very wide, $3\frac{1}{2}$ times wider than the eye; bones on top of head coarsely granular; Y-shaped ridge on top of head conspicuous, on each side of which is a broad naked area; naked space between forks of Y on top of head long and narrow, but shorter than vertical limb of the Y which is very long; edges of nostrils fringed, anterior nostril round, separated from the eye by a high granular ridge; posterior nostril ending in a long curved furrow, which runs obliquely across the naked area behind eye, its posterior end not curved forward, its length $2\frac{1}{2}$ times the diameter of the eye; 2 or 3 small blunt spines in front of the eye; surface of the bones of opercle, preopercle, and humeral process coarsely granular; gill rakers not developed; pseudobranchiæ small. Head entirely scaleless; belly naked below a line drawn from fifth anal ray to upper end of pectoral base; fins without scales; scales very small and somewhat embedded. Width of pectoral at base less than $\frac{1}{2}$ length of the head, the upper rays longest, the lower rays very short, graduated from the lower side to the upper; fin somewhat pointed behind and curved up, its tip reaching to the vertical from base of sixth dorsal ray; the rays of ventrals very thick and swollen, the inner

* The following are the characters of *Uranoscopus:* Head with spines; humeral spine well developed; 1 strong spine on subopercle, 4 smaller ones on preopercle, all directed downward; 1 small spine directly above and in front of humeral spine; 4 low, stout protuberances on top of head pointing backward; naked space between eyes extending back to posterior part of orbits; upper lip and nostrils not fringed; retractile tentacle in mouth more or less developed. First dorsal with about 4 pungent spines; scales well developed.

rays the longest, reaching midway from their base to end of pectorals; origin of fin a distance of the width of pectoral in front of the lower edge of pectoral base; soft dorsal much higher than anal; posterior rays reaching slightly past the vertical from base of the last anal ray; end of the last anal ray about reaching to base of caudal rays; caudal truncate or slightly rounded; a ridge of skin along middle of belly from the ventrals to vent. Dark brown above, paler below; upper parts densely covered with small rounded white spots, each surrounded by a black ring; lower jaw and labial fringes similarly spotted; spinous dorsal black, white posteriorly; soft dorsal brown anteriorly with a horizontal white and black band, then tipped with white; posteriorly with 2 vertical black stripes and a white one between them; caudal black, tipped with white, with 2 to 4 white longitudinal stripes, its upper and lower edges narrowly white; the anal white at base and tip, with a black median band, $\frac{1}{3}$ depth of fin, darkest posteriorly; pectorals brown, with a black band below, the lower edge white, the upper ray spotted; ventrals white with a black lengthwise streak. Old examples lose the black ring around the spots, and the edges of the spots are blended into the dark brown of the back; a dark stripe running from the upper angle of gill opening to caudal. South Atlantic coast from Cape Hatteras to the Caribbean Sea, in sandy bays, rather common in shallow water, varying much with age. Here described from a specimen, 15 inches in length, from Charleston, South Carolina. It is recorded from Charleston, Beaufort, Matanzas River, St. Johns River, Pensacola, Key West, and "the Caribbean Sea." According to Dr. James A. Henshall, the naked area on top of the head in *Astroscopus* is the seat of electric power. This interesting statement needs verification. (Named from the armature of the head, in the form of the Greek *Y*.)

Uranoscopus y-græcum, CUVIER & VALENCIENNES, Hist. Nat. Poiss., III, 308, 1829, origin unknown; GÜNTHER, Cat., II, 229, 1860.

Uranoscopus anoplos,* CUVIER & VALENCIENNES, Hist. Nat. Poiss., VIII, 493, 1831 (young examples), Charleston, South Carolina.

Upsilonphorus y-græcum, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 113; KIRSCH, *l. c.*, 263, 1889.

Astroscopus y-græcum, BEAN, Proc. U. S. Nat. Mus. 1879, 58; JORDAN & GILBERT, Synopsis, 628, 1883.

Agnus anoplus, GÜNTHER, Cat., II, 229, 1860.

Astroscopus anoplus, JORDAN & GILBERT, Synopsis, 629, 1883.

Astroscopus anoplos, KIRSCH, Proc. Ac. Nat. Sci. Phila. 1889, 262.

* The genus *Astroscopus* was based on small specimens which, in our present opinion are simply immature examples of the species *y-græcum*. The supposed genus is thus described in distinction from *Upsilonphorus*, which seems to us the adult of the same type: Head covered above with bone except a small region between and in front of the eyes, the bony occipital plate coalescing with the orbital rims; no spines on head; humeral spines obsolete; occipital region with bluntish projections; naked space between eyes extending back to near middle of orbits; lips and nostrils fringed; no retractile tentacle in mouth.

The following characters are assigned by Dr. Kirsch to *Astroscopus anoplos:* Head $2\frac{2}{3}$; depth $3\frac{1}{4}$. D. IV-I, 13; A. 13. Pectorals rather large, their longest ray equaling in length base of second dorsal and extending to front of that fin; ventrals equaling pectorals in length, and extending to front of that fin; the second dorsal equaling anal but its anterior insertion slightly posterior to that; anal rays reaching base of caudal; vent much nearer base of caudal than to tip of snout. Color dark brown above, yellowish below; lighter portions of body covered with small white specks; chin jet-black; all the fins whitish. Length 2 inches. (Specimen from Key West). Small individuals are found along the coast from Cape Hatteras to Florida wherever *A. y-græcum* is found. The adult differs mainly in the armature of the top of the head, a characteristic which is developed at different ages in different individuals.

2645. ASTROSCOPUS ZEPHYREUS, Gilbert & Starks.

Head, without lower jaw, $2\frac{2}{3}$; depth $3\frac{2}{5}$. D. V, 13; A. 14; scales 84; eye 12 in head; maxillary $2\frac{1}{3}$; pectoral $1\frac{1}{4}$; second dorsal spine 7; highest dorsal ray $2\frac{1}{3}$; highest anal ray $3\frac{1}{6}$; caudal $1\frac{3}{4}$. Body robust, widest at occiput, slightly compressed posteriorly; anteriorly subcylindrical. Head very large and broad, wider than the body; mouth large, vertical, a fringe of barbels curving over mouth on each jaw; length a little greater than the diameter of the eye; tongue very large and fleshy, forming a pad under the membrane of lower jaw, which projects forward somewhat; teeth conical, small and movable, in many bands in upper jaw; in lower jaw the teeth are larger and in 2 or 3 rows; vomer and palatines with teeth; eyes very small but prominent, set on top of head; interorbital very wide, 4 times as wide as the eye; bones on top of head coarsely granular; Y-shaped ridge on top of head conspicuous, on each side of which is a broad naked area, the form of these and other bones of the head exactly as in *A. y-græcum;* edges of nostrils closely fringed, anterior nostril round, the ridge between it and eye not very high or conspicuous; posterior nostril ending in a long curved furrow which runs obliquely across the naked area behind eyes; at its posterior end it turns sharply forward, its length $2\frac{2}{3}$ times the diameter of the eye; 2 very short blunt spines in front of the eye; surface of the opercle, preopercle, and humeral process granular, not so rough as in *Astroscopus y-græcum;* gill rakers not developed; pseudobranchiæ very small. Head entirely scaleless; belly naked below a line drawn from first anal ray to the middle of the pectoral base; fins without scales; scales small and nearly square, grown together side by side, forming series of oblique plates. Width of pectoral at base slightly less than $\frac{1}{2}$ length of head, the lower rays very short and graduated to the long upper rays, the fin pointed and slightly turned up, its tip reaching to the vertical from base of the third dorsal ray; the ventral rays thick and swollen, the inner rays the longest, its tip reaching about midway between its base and tips of pectorals; origin of fin in front of pectorals a distance equal to the width of pectoral base; soft dorsal somewhat higher than anal, its posterior rays reaching to the vertical from base of last anal ray; tip of last anal ray nearly reaching to the base of caudal rays; caudal truncate or slightly rounded; a fold of skin along middle line of belly from ventrals to vent. Color dark brown above, paler below; upper parts with many round white spots of various sizes, edged with rings of dark brown; spinous dorsal black, light posteriorly; soft dorsal light at base, the ends of the rays with black and white stripes; pectoral and anal dusky with light edge; caudal with longitudinal black and white stripes. Pacific coast of Mexico. One specimen, numbered 333, in the Leland Stanford Junior University Museum, collected by the *Albatross* at Magdalena Bay, Lower California. It is 12 inches in length. A distinct electric shock was given by this fish when alive, the electric organs being in the fleshy areas on top of head behind

eyes. (Gilbert.) A second large specimen was sent from Mazatlan by Dr. George W. Rogers, having been taken by Ygnacio Moreno in January, 1896. (ζεφύριος, western; ζέφυρος, the west wind.)

Astroscopus zephyreus, GILBERT & STARKS, Proc. U. S. Nat. Mus. 1896, 453, pl. 53, fig. 2, and pl. 54, Magdalena Bay, Lower California (Type No. 47743. Coll. *Albatross*).

2646. ASTROSCOPUS GUTTATUS (Abbott).

Depth 4 in length in young and 3¼ in adult. D. IV or V–13 or 14; A. 13; V. I, 5. Eye 5½ in interorbital space. Naked space between forks of Y on top of head short and broad, but longer than the vertical limb of the Y, which is very short; 2 distinct spinules directed forward before eye; white spots on body very small and irregular without dark rings; base of dorsals equaling in length the distance from front of first dorsal to tip of snout; base of first dorsal twice length of its longest spine; first spine equaling second in length, and 3 times length of last; length of middle caudal rays a little less than that of ventrals; pectorals slightly longer than ventrals, 3½ in total length, and extending to fifth anal ray. Color of upper parts of body and lower jaw bright chocolate; belly and throat white; darker portions covered with numerous circular spots much lighter than ground color; membrane of first dorsal black; second dorsal white with 3 irregular bands of dull black obliquely across it; the caudal with 3 parallel bands of blackish brown, the middle of which appears to be the continuation of a variable longitudinal band on the center of each side; the anal having a variable band of dull brown, darker upon the posterior termination. Length 12 inches. Atlantic coast of the United States, from Long Island to Virginia; apparently scarce. Recorded from Cape May; Tompkinsville, New York; Norfolk, Virginia; Somers Point, New Jersey, etc; not known south of Cape Hatteras. In *Astroscopus guttatus* the pale spots are much smaller, less sharply defined, and occupy a smaller area than in *A. y-græcum;* the lower part of the head has 2 black blotches in each species; the second dorsal, anal, and ventrals are nearly or quite plain. The naked area behind each eye is (in *A. guttatus*) lunate, its length barely twice that of the snout; the bony Y-shaped plate is short and broad, concave on the median line, and forked for about ½ its length, the posterior undivided portion broader than long; the bony bridge across the occiput but little shorter than the part of the head which precedes it. In *A. y-græcum* the naked area is trapezoidal, longer than broad, and about 4 times the length of the snout; the Y is forked for more than ½ its length, its undivided part more than twice as long as broad, and not concave; the occipital plate is not ½ as long as the part of the head which precedes it. (*guttatus*, spotted, as with rain drops.)

Astroscopus guttatus, ABBOTT, Proc. Ac. Nat. Sci. Phila. 1860, 365, Cape May, New Jersey. *Upsilonphorus guttatus*, BEAN, Proc. U. S. Nat. Mus. 1879, 60; KIRSCH, *l. c.*, 264, 1889.

857. KATHETOSTOMA, Günther.

Kathetostoma, GÜNTHER, Cat. Fish., II, 231, 1860 (*lævе*).

Body robust, formed as in *Astroscopus* and *Uranoscopus*. Scales none. One continuous dorsal without spines; ventrals jugular not adnate to the abdomen; pectoral rays branched; some bones of the head armed. Cavity of the gills without superior opening; 6 branchiostegals; pseudobranchiæ present. Air bladder none. Three species known, the type, *Kathetostoma lævе*, being from Australia. (*κάθετος*, vertical; *στόμα*, mouth.)

a. Dorsal rays 13; anal 13; body shaded and dotted with blackish. AVERRUNCUS, 2647.
aa. Dorsal rays 10; anal 12; body spotted with white. ALBIGUTTA, 2648.

2647. KATHETOSTOMA AVERRUNCUS, Jordan & Bollman.

Head $2\frac{3}{8}$, $3\frac{1}{3}$ with caudal; depth $3\frac{3}{4}$. D. 13; A. 13. Body short and robust, its width behind base of pectorals equal to length of top of head. Head very large, its width at peropercle less than its length by $\frac{1}{2}$ length of eye. Mouth large, vertical; maxillary 2 in head. Snout $1\frac{2}{3}$ in eye. Eye rather small, 5 in head. Teeth of lower jaw largest, inner row of each jaw enlarged and movable; vomer and palatines with a few large, conical teeth. Lower jaw without tentacle. Interorbital space lightly concave, $1\frac{1}{4}$ times length of eye. Premaxillary groove as broad as long, $1\frac{1}{2}$ in eye, obtuse behind, extending backward just past middle of pupil. Distance between bases of humeral spines $1\frac{1}{4}$ in top of head. Preorbital with 3 spines in front directed forward and downward. Preopercle with 3 spines below angle directed downward and forward. Two antrorse spines on mandible, and 2 on breast before ventrals. Bones of top of head coarsely granular, striate, no naked area above except premaxillary groove; 2 points on occipital region whence granular ridges radiate; opercles and orbital bones coarsely granular, but not striate. No trace of scales or of spinous dorsal. Base of dorsal equal to base of anal, $1\frac{1}{3}$ in head, longest ray equal to depth of cheek; pectorals $\frac{1}{2}$ eye, length greater than that of top of head; ventrals reaching more than halfway to vent, their length equal to that of top of head. A few small depressions resembling embedded scales on region before dorsal and above head. Color blackish brown, mottled with paler; lower parts pale, dusted with brown; lips and gular region black; dorsal dusty, with 5 indistinct, partly confluent, whitish spots along its base; anterior part of anal pale, posterior thickly dusted with blackish, tips of rays pale; pectorals blackish, faintly barred; axil dusted outside, inner part very pale; ventrals pale; caudal with 3 irregular oblique dark bars; floor of mouth pinkish; tongue dusted with dark specks. Length $4\frac{1}{2}$ inches. Pacific Ocean, off coast of Colombia; a single specimen dredged at a depth of 7 fathoms; a most singular fish. (*averruncus*, a deity which wards off; from the mailed head.)

Kathetostoma averruncus, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 163, off coast of Colombia, at Albatross Station 2800, 8° 57′ N., 79° 31′ 30″ W.; KIRSCH, *l. c.*, 259, 1889; JORDAN, Proc. Cal. Ac. Sci. 1896, 229, pl. 31.

2648. KATHETOSTOMA ALBIGUTTA, Bean.

Head 3; greatest width 3; depth 3½. D. 10; A. 12; interorbital space 4 in head, containing a deep groove, the length of which is slightly greater than its width and nearly equaling length of eye. Mouth nearly vertical when closed; intermaxillary slightly protractile, the length of its tooth-bearing surface ⅔ length of head; maxillary very broadly expanded behind, its greatest width about 3 in length, extending almost to vertical from middle of eye; end of mandible not much farther back; length of mandible 4⅜ in length; mandible having 2 blunt prominences at its posterior end; the exposed portion of the maxillary traversed by radiating striæ. The lower limb of preoperculum with 3 stout spines along its lower border; length of humeral spine 3 in head; humerus very strongly rugose on its upper border; 3 short spines on the anterior edge of preorbital. Teeth in villiform bands in the intermaxillary and mandidible, and on vomer; palatines in a very short band; a cavity between head of vomer and the processes of the intermaxillary ending in a semicircular canal behind, which is separated from the anterior cavity by a flap of skin. Gill openings very wide and only narrowly attached to the isthmus, leaving a free posterior border. Pseudobranchiæ present, small; a small, narrow slit behind the last gill, its length about ⅔ that of eye; gill rakers tubercular, none on anterior arch. A pair of short but stout spines in front of ventrals. The origin of dorsal a little nearer to root of caudal than to tip of snout, midway between base of caudal and middle of eye; length of dorsal base about 3 in length, the third ray the longest, its length nearly ½ length of base of fin, the last ray about as long as eye, and the first scarcely longer than this. The anal origin directly under that of dorsal, the base of fin slightly longer than that of dorsal; the seventh, eighth, and ninth anal rays about the longest, their length equaling about ½ that of middle caudal rays; the first ray not much more than ½ as long as the longest and the rays gradually increasing in size to the ninth; length of pectoral 3½ in body; length of lowermost ray less than ¼ length of head; only the first ray simple, the rest divided. Ventral origin under eye; the longest ray of ventral slightly shorter than mandible. Caudal slightly rounded when expanded, the middle rays as long as head without snout. The lateral line beginning near the root of humeral spine, curving upward slightly and running along back to end of dorsal, then curving downward to near the middle of the caudal base; skin naked. Color, upper parts light brown, the upper surface of the head minutely dotted with white; the back with numerous roundish spots and oblong blotches of whitish; lower parts pale; the dorsal with 2 or three dark blotches near the margin, in some cases not much larger than eye, in others fully twice as long; caudal with 9 black blotches, those on outer rays largest, differing in size in different specimens, these blotches distributed over the greater portion of the fin; anal pale, with the exception of a brownish blotch on the membrane of the last 3 rays; pectoral with a brownish submarginal band on its outer half, this band sometimes broken up on the mem-

brane; ventrals pale. Length about 6½ inches. Gulf of Mexico, in 27 to 88 fathoms; 1 specimen known. (*albus*, white; *gutta*, spot.)

Cathetostoma albigutta, BEAN, Proc. U. S. Nat. Mus. 1892, 121, Gulf of Mexico, at Albatross Station 2403, Lat. 28° 42′ 30″ N., Lon. 85° 29′ 00″ W. (Type, No. 39304, U. S. Nat. Mus.)

Suborder HAPLODOCI.

This group is distinguished mainly by the undivided post-temporal, the reduction in the number of gill arches to 3, and by the absence of peculiarities shown by related forms. One family. (*ἁπλόος*, simple; *δοκος*, a shaft or beam, from the form of the post-temporal.)

Family CXCVIII. BATRACHOIDIDÆ.

(THE TOAD-FISHES.)

Body more or less robust, depressed anteriorly, compressed behind; head large, depressed, its muciferous channels well developed; mouth very large, the teeth generally strong; premaxillaries protractile; gills 3, a slit behind the last; pseudobranchiæ none; gill openings restricted to the sides, the membranes broadly united to the isthmus; branchiostegals mostly 6; gill rakers present, moderate; suborbital without bony stay; post-temporal bone simple, undivided; scales small, cycloid, or wanting; dorsal fins 2, the first of 2 or 3 low, stout spines; soft dorsal very long; anal fin similar, but shorter; ventrals rather large, jugular, I, 2 or I, 3; pectorals very broad, the rays branched; pyloric cæca none; tail diphycercal, the caudal fin distinct, rounded; vertebræ in large number, 32 to 45. Carnivorous coast fishes, mostly of the warm seas, some of them ascending rivers; the young of some or all the species fasten themselves to rocks by means of an adhesive ventral disk, which soon disappears. In some species the spines of the head and dorsal fin are provided with poison glands. Genera 7; species about 15. (*Batrachidæ*, Günther, Cat., III, 166–177.)

a. Dorsal spines 3; opercle developed as 2 strong diverging spines; subopercle rather strong, with 2 spines similar to those of opercle; no venom glands.

b. Body scaly; branches of subopercular spine subequal and diverging; frontal region broad, flat, and slightly depressed, its median ridge rather prominent. BATRACHOIDES, 858.

bb. Body scaleless; branches of subopercular spine parallel, the lower branch much the shorter; vertebræ 10+22; frontal region not depressed, its median ridge prominent; axil with a large foramen. OPSANUS, 859.

aa. Dorsal spines 2; opercle very small, its posterior part developed as a single strong spine; subopercle feebly developed, narrowed, and not ending in a spine; body scaleless.

c. Spines solid, without venom glands; several lateral lines on sides of head and body, composed of pores and shining spots, some of these accompanied by cirri; canine teeth present; vertebræ 12+31; frontal region depressed, forming a triangular area below level of temporal region, its median ridge very low. PORICHTHYS, 860.

cc. Spines of dorsal fin and operculum hollow and connected with venom glands; lateral line on sides of body single; no canine teeth.

d. Dorsal and anal free from caudal. THALASSOPHRYNE, 861.

dd. Dorsal and anal fully joined to caudal. DACTOR, 862.

858. BATRACHOIDES, Lacépède.

Batrachoides, LACÉPÈDE, Hist. Nat. Poiss., III, 306, 1798 ("*tau*," Lacépède*=*surinamensis*).
Batrachus (KLEIN), BLOCH & SCHNEIDER Syst. Ichth., 42, 1801 ("*tau*," *didactylus*, *surinamensis*, etc.; substitute for *Batrachoides*).
Batrictius, RAFINESQUE, Anal. Nat. 1815, 82 (substitute for *Batrachoides*).

Body robust, formed as in *Opsanus*. Dorsal spines 3; opercle developed as 2 strong diverging spines; subopercle strongly developed; branches of subopercular spine subequal and diverging; body covered with small ctenoid scales; frontal region broad, flat, and slightly depressed, its median ridge rather prominent. Mucous pores of sides not greatly developed. No poison glands. Shore fishes of warm regions. (βάτραχος, frog; εἶδος, resemblance.)

a. Teeth small, about 14 on the vomer; anterior teeth of lower jaw in a band; lateral teeth of palatine enlarged and canine-like; irregularly arranged.
SURINAMENSIS, 2649.

aa. Teeth larger, about 8 on vomer; anterior teeth of lower jaw in 2 rows; 3 teeth on middle of palatines enlarged and canine-like, the middle one the smallest.
PACIFICI, 2650.

2649. BATRACHOIDES SURINAMENSIS (Bloch & Schneider).

(SAPO.)

Head 3¼ in length of body; depth 6. D. III–29; A. 26. Teeth small, about 14 on vomer; anterior teeth on lower jaw in a band; lateral teeth on palatines enlarged and canine-like, irregularly arranged; pectoral without pores on its inner surface. Color grayish, darker on sides and head; base of soft dorsal pale, with a dark, irregular line above; upper part of fin lighter; caudal nearly black; anal fin light with some dark markings. Coasts of Guiana and Brazil; not rare on sandy shores; our specimen from Curaçao.

Batrachoides tau, LACÉPÈDE, Hist. Nat. Poiss., 306, pl. 12, fig. 1, 1798; not *Gadus tau*, LINNÆUS.
Batrachus surinamensis, BLOCH & SCHNEIDER, Syst. Ichth., 43, 1801, Surinam; from a specimen in the Museum of Vaillant in Paris; GÜNTHER, Cat., III, 173, 1861; MEEK & HALL, Proc. Ac. Nat. Sci. Phila. 1885, 61.

2650. BATRACHOIDES PACIFICI (Günther).

Head 3 in length; depth about 6. D. III–26; A. 22. Teeth rather large, about 8 on vomer; anterior teeth on lower jaw in 2 rows; lateral teeth on lower jaw gradually increasing to middle of jaw, behind which they become abruptly smaller and then gradually increase to end of jaw; 3 teeth on middle of palatines enlarged and canine-like, the middle one the smallest; pectoral with a row of pores on inner surface. Color olivaceous brown; some indistinct dark cross bands on body; dorsal with about 7 very irregular oblique dark bars, anal with about 5; pectorals and caudal

* "Il est revêtu d'écailles molles, petites, minces, rondes, brunes, bordées de blanc, et arrosées par une mucosité très abondante, comme celles de la lote et de la mustelle." (Lacépède.) Lacépède's specimen was therefore one of the scaly species, not an *Opsanus*. No species of the latter group seems to have been known to Lacépède or to Schneider.

dark, with few light cross bands. Panama; locally common, close to the preceding but with smaller teeth and fewer fin rays. The specimen examined by us collected by Dr. Gilbert.

Batrachus pacifici, GÜNTHER, Cat., III, 173, 1861, Panama; GÜNTHER, Fishes Centr. Amer., 435, 1869.

Batrachoides pacifici, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 170; MEEK & HALL, Proc. Ac. Nat. Sci. Phila. 1885, 62.

859. OPSANUS,* Rafinesque.

(TOAD-FISHES.)

Opsanus, RAFINESQUE, Amer. Monthly Mag. 1817, 203 (*cerapalus*).

Batrachus, JORDAN & GILBERT, Synopsis, 751, 1883, and of authors; not of BLOCH & SCHNEIDER.

Body comparatively short and robust, scaleless; head large, depressed; jaws, vomer, and palatines each with a single series of strong blunt teeth; mandible with an additional external series at symphysis; teeth of upper jaw small; dentary bones forming an acute angle at symphysis; lips fleshy; upper angle of opercle with 2 diverging spines, more or less concealed in the skin; no poison glands; spinous dorsal of 3 stout, short spines, the second the longest; axil of pectoral with a large foramen;† lateral line obscure, its pores not conspicuous; young with a series of small, tufted cirri on back and sides; branchiostegals 6; vertebræ 12 + 22. Shore fishes, mostly of temperate regions; voracious creatures, living on the bottoms, feeding on mollusks and crustacea, and having great strength of jaw. (*ὤψ*, eye; *ἄνω*, upward; "the name means looking up." Rafinesque.)

a. Nostrils with fleshy tentacle between them. Color brownish or dusky greenish, mottled with darker and lighter, the dark on sides of body in large irregular blotches extending from base of dorsal to about ⅔ distance to base of anal, and more or less covered with small pale spots; belly and chin plain white or yellowish. TAU, 2651.

aa. Nostrils without fleshy tentacle. Color whitish or gray, everywhere blotched or spotted with brownish yellow and black, the black spots on top of head smaller and more numerous than on rest of body; a large black blotch at base of spinous dorsal, running up on fin; 3 black blotches along base of soft dorsal, which do not extend ⅓ the distance to base of anal; pectoral with black spots which do not form cross bands; ventrals with more dark markings than in *tau*; dorsal, anal, and caudal marked nearly as in *tau*. PARDUS, 2652.

2651. OPSANUS TAU (Linnæus).

(TOADFISH; SAPO; SLIMER; OYSTER-FISH.)

Head 2⅔; depth 4⅓. D. III–26 to 28; A. 24. Body robust, naked, the head broad; mouth large, the very strong jaws closing with great force; teeth blunt, those on mandible small anteriorly, regularly increasing in

* The name *Batrachus* should not be used for this genus, as it was originally given merely as a substitute for *Batrachoides*, having properly the same type, *surinamensis*, wrongly supposed to be *tau* of Linnæus, a species unknown to Lacépède and Bloch & Schneider. No congener of *tau* was placed in *Batrachus* by Bloch & Schneider. Prior to any use of *Batrachus* as the generic name of the naked toadfishes, allied to *tau*, Rafinesque had given to one of the latter the generic name *Opsanus*, which can not be set aside for *Batrachus*, the latter being an unnecessary synonym of *Batrachoides*.

† The Brazilian genus, *Marcgravia* (*cryptocentra*), in which this foramen is wanting, has not been recorded from north of the equator.

size backward, those on vomer prominent; a broad flap above orbit; tip of maxillary and lower side of mandible with conspicuous cirri; a series of smaller cirri along margin of preopercle; subopercle ending in a long, sharp spine; orbit about equaling interorbital width or length of snout; pectoral with a large foramen in the axil. Dusky olive, with black markings confluent on the sides and forming irregular, indistinct bars; belly and under side of head lighter; sides often with many pale yellow or whitish spots; soft dorsal with 6 to 9 oblique light bands; anal with 5 to 9; caudal and pectoral fins with 5 to 7 light cross bands, these formed chiefly from light spots; ventrals with some dark markings. In specimens from shallow water or algæ, the brown becomes nearly black and more extended, the belly and chin spotted with darker, and top of the head has no distinct markings. The deeper-water specimens are lighter in coloration than those from near the surface, and those from the coral reefs (var. *beta*, Goode & Bean) are paler than those from the green algæ and sea wrack; otherwise no differences seem to exist. In young individuals the head is more narrow and rounded, and the lower branch of the subopercular spine proportionally larger than in the adult. Cape Cod to Cuba; very abundant among rocks and weeds close to the shore northward, in deeper water southward; the young clinging to rocks by a ventral sucking disk, which is soon lost. Length 15 inches. Not valued as food. (*tau*, T, the bones on the head when dried showing a T-shaped figure.)

Gadus tau, LINNÆUS, Syst. Nat., Ed. XII, 440, 1766, Carolina. (Coll. Dr. Garden.)
Cottus glaber, SCHÖPF, Schrift. Naturf. Freunde, VIII, 1788, 146, Long Island; D. 25; V. 3; A. 21; short cirri below mouth.
Cottus chætodon, BLOCH & SCHNEIDER, Syst. Ichth., 62, 1801, New York; after SCHÖPF.
Lophius bufo, MITCHILL, Trans. Lit. and Phil. Soc. 1815, 463, New York.
Opsanus cerapalus, RAFINESQUE, Amer. Monthly Mag., Jan., 1817, 204, south coast of Long Island. (Coll. C. S. Rafinesque.)
Batrachoides vernullas, LE SUEUR, Mém. Mus., V, 1819, 157, pl. 17, coast of Rhode Island.
Batrachoides variegatus, LE SUEUR, Jour. Ac. Nat. Sci. Phila., III, 1823, 399 and 401, Egg Harbor, New Jersey.
Batrachus celatus, DE KAY, New York Fauna: Fishes, 170, pl. 50, f. 161, 1842, New York.
Batrachus tau beta, GOODE & BEAN, Proc. U. S. Nat. Mus. 1882, 236, Gulf of Mexico.
Cottus glaber, WALBAUM, Artedi Piscum, III, 392, 1792; after SCHÖPF.
Batrachus tau, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 478, 1837; DE KAY, N. Y. Fauna: Fishes, 168, pl. 28, fig. 26, 1842; GÜNTHER, Cat., III, 167, 1861; JORDAN & GILBERT, Synopsis, 751, 1883; MEEK & HALL, Proc. Ac. Nat. Sci. Phila. 1885, 59.
Batrachus variegatus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 484, 1837.

2652. OPSANUS PARDUS (Goode & Bean).

(SAPO.)

Head to end of opercular spine 3; depth 4. D. III–26; A. 22; maxillary $1\frac{4}{5}$ in head; pectoral $2\frac{1}{6}$; ventral 2; highest dorsal ray $2\frac{5}{6}$; highest anal ray $3\frac{1}{3}$; caudal 2. Body short and robust, compressed posteriorly; head large, somewhat depressed, wider than the body; eyes placed high, not so wide as the slightly concave interorbital space; mouth large, the maxillary reaching far beyond the eye, the lower jaw slightly projecting; a double row of small blunt teeth in upper jaw, not running very far back at the sides; lower jaw with a single row of much larger pebble-like teeth running well back and biting against a single row of similar teeth on pala-

tines; a few teeth in front of jaw which bite against the premaxillary teeth; vomer with 1 or 2 irregular rows of large blunt teeth; head with many fleshy tentacles, 1 over each eye, a row around lower jaw, 1 on end of maxillary, and a row around preopercle; opercle ending in 2 diverging spines, the lower shorter; subopercle ending in a spine, its tip equal with the lower opercular spine, these spines not piercing the skin; gill rakers very short, scarcely developed. Body and fins covered with a soft smooth skin, which is exceedingly loose nearly to the ends of fin rays, and entirely covering the dorsal spines. Dorsal spines very short but stout; soft dorsal longer and higher than anal, but in other ways similar, reaching past base of caudal rays; pectoral short, as wide as long, round and fan-shaped behind, reaching to vertical from base of fourth dorsal ray; origin of ventral far in front of pectorals, the fins reaching to the vertical from the posterior edge of spinous dorsal; caudal well rounded, fan-shaped when spread. Color very pale yellowish brown, thickly covered with round spots of dark brown, those on head smaller; belly with numerous spots, the largest as large as eye; back with many oblong blotches, besides small round spots; fins blotched and banded. Gulf of Mexico, in deep water. This form has a very different coloration from *O. tau* and the texture of its skin and flesh is also less firm, but the technical differences are slight and it is rather a deep-water variety than a species. (*pardus*, leopard.)

Batrachus tau pardus, Goode & Bean, Proc. U. S. Nat. Mus. 1879, 336, Pensacola Snapper Banks; Jordan & Gilbert, Synopsis, 751, 1883; Meek & Hall, Proc. Ac. Nat. Sci. Phila. 1885, 60.

860. PORICHTHYS, Girard.

(Midshipmen.)

Porichthys, Girard, Proc. Ac. Nat. Sci. Phila. 1854, 141 (*notatus*).

Body rather elongate; head not very broad, depressed, the lower jaw projecting. Dorsal spines 2; pectoral broad, without foramen in axil; opercle very small, its posterior part developed as a strong, single spine; suboperculum feebly developed, narrowed and not ending in a spine; no scales on body; spines solid, without venom glands; several lateral lines on sides of head and body, composed of pores and shining spots, some of these accompanied by cirri; canine teeth present; vertebræ 12 + 31; frontal region depressed, forming a triangular area below level of temporal region, its median ridge very low. Branchiostegals 6; interorbital area short, wide, and with shallow grooves. Air bladder more or less deeply divided into 2 lateral parts. Pyloric appendages none. Species American; remarkable for the very great development of mucous pores, some of which simulate the photophores of *Myctophum*, but are different in origin and not at all luminous. (πόρος, pore; ἰχθύς, fish; in allusion to the extraordinary development of the mucous system.)

Note.—The following account of the distribution, structure, and development of the phosphorescent organs of *Porichthys* is furnished us by Prof. Charles Wilson Greene, who has made a careful study of these organs:

"*Porichthys* has numerous lines of conspicuous bright silvery spots distributed in rows over the surface of the body. These spots have been called phosphorescent organs, although no such function has yet been observed, the name arising out of a superficial

resemblance. These so-called phosphorescent organs are arranged in rows over the body, and are definite and characteristic and quite constant in location in different individuals. They are accompanied by rows of epidermal sense organs, the two having an intimate relation in distribution over the surface of the fish. In surface view the shining organs have a bright silvery appearance, are more or less round in outline, size from a mere dot to 0.8 mm. in diameter, and surrounded or bordered on one side by an increased amount of pigment. The end buds present a round, transparent, or pellucid, and usually slightly raised, point. Each end bud is bordered by a pair of papillæ. There are about 20 well-defined lines as follows: The *lateral* row, from posterior upper border of pectoral straight alongside to upper third of base of caudal, 35 pairs with an end bud between each pair, upper series small or rudimentary, segmentally arranged and between myomeres. The *pleural* row, from middle of base of pectoral, curves backward and downward to a point above first anal ray then straight nearly to base of caudal, 43 to 62 organs. End buds below each organ to above middle of caudal, 31 organs. The *caudal* rows, end buds only, 2 longitudinal rows on upper and lower thirds of fin. The *anal* row, on either side base of anal fin from third anal ray to base of caudal. Phosphorescent organs in pairs, a pair for each anal ray, 1 end bud for each pair. The *gastric* row, from front around lower edge of pectoral and along side of belly to opposite anal papilla, 30 phosphorescent organs. The *gular* row, from isthmus along ventral side of ventral fin then outward to join gastric row, spur runs forward along external side of ventral fin, 27 organs. A parallel line of 50 end buds follows the gular row and posterior end of gastric. The *ventral* row with its fellow forms a parenthesis on the stomach from the side of the anus ¾ the distance to the ventral fin, 34 organs; no end buds. The *branchiostegal* row, from the isthmus outward over branchiostegal membrane and between first and second rays, no end buds. The *mandibular* row of phosphorescent organs extends around inner edge of ridge formed by the dentary bones; the row of end buds along the outer rim of the same ridge. The *opercular* rows, upper and lower, extend backward and upward across opercle. The *scapular* row, from above opercular spine straight back above pectoral fin, the curves in toward the base of the dorsal fin opposite the third dorsal ray. The *dorsal* row, along base of dorsal fin to base of caudal. This row and the scapular row consist of well developed end buds and rudimentary phosphorescent organs. The *occipital* and *frontal* rows, along the occipital and frontal regions, short rows of small and poorly developed organs. The *nasal*, from the posterior nasal tube to base of anterior tube. The *suborbital* and *postorbital*, from posterior nasal opening around under eye backward and downward to opercle. A *malar* row, from the suborbital down across the cheek. A *maxillary* across the posterior end of maxillary bone. The rows on the head consist of well-developed end buds with rudimentary and irregularly placed phosphorescent organs. The phosphorescent organs are embedded in the connective tissue dermis of the skin, and in section show a uniform general structure throughout the body. A typical organ from the anal or ventral rows consists of an outer spherical group of cells called a lens, resting in a deeper cup-like structure, the capsule, and this in turn in a cup of fibrillar connective tissue called the reflector. The lens consists of cells, polygonal in the center of the group and flattened or fusiform around the periphery. They have a large conspicuous nucleus and a dense, homogeneous, highly refracting cell body. The outlines of the cells are very distinct. In the cells of the capsule the nuclei stain readily, but the granular protoplasm with difficulty, and the cell boundaries are indistinct and usually obliterated. In some specimens connective tissue septa penetrate the capsule. Blood capillaries are always present. The reflector extends well up around the sides of the lens; it consists of fibrillar connective tissue which strongly reflects light. Much pigment is embedded in its meshes. No nerves have yet been traced to the organ. The developing phosphorescent organs do not appear in the embryo fish until it is 15 to 16 mm. long. Then a bud appears in the lower layer of the epidermis, which soon becomes constricted off as a spherical mass of shells lying in the subepidermal connective tissue. This mass later slightly elongates and gives rise by constriction to the lens and the capsule. The reflector is developed from the surrounding connective tissue, so also the pigment cells. Mature organs are not found until the fish reaches a length of over 20 mm. The end buds appear much earlier, 9 to 10 mm." (Charles Wilson Greene.)

a. Abdomen with 4 longitudinal series of pores, each of which is accompanied by a shining silvery body; 4 rows of shining spots on sides of body; a white blotch below eye, with a black crescent below it.

b. Teeth on palatines few (4 or 5), 1 to 3 of them developed as very strong canines, as large as canines on vomer; dorsal fin with distinct black blotches; back with dark saddles; third lateral line extending nearly to base of caudal. POROSISSIMUS, 2653.

bb. Teeth on palatines numerous, none of them canine, and all much smaller than canines on vomer.

c. Third lateral line ceasing at second third of anal; cross bands on back and dorsal fin very faint or wanting; dorsal fin with a faint dark edge; sides of head and shoulder without distinct spots; body rather elongate. NOTATUS, 2654.

cc. Third lateral line extending nearly to end of anal; cross bands on back and dorsal fin very distinct, appearing as roundish blotches, those on the dorsal fin along the margin; sides of head and humeral region much spotted with brown; body robust. MARGARITATUS, 2655.

2653. PORICHTHYS POROSISSIMUS (Cuvier & Valenciennes).

(BAGRE SAPO.)

Head $3\frac{3}{4}$ ($4\frac{1}{4}$ in total); depth $5\frac{2}{3}$ (6). D. II–37; A. 34. Body rather elongate, tapering and compressed behind. Head depressed, $\frac{2}{3}$ as broad as long and $\frac{1}{2}$ wider than deep; lower jaw considerably projecting, maxillary reaching to well behind eye, its length $1\frac{1}{4}$ in head. Teeth in single series on jaws, vomer, and palatines, those of upper jaw very small, a few of the anterior and 2 or 3 of the lateral teeth somewhat enlarged, the latter strongly hooked forward; teeth in lower jaw strong, rather weaker than in *P. margaritatus*, those in the front of the jaw hooked strongly inward; the lateral teeth, which are larger, hooked backward and inward; 1 or 2 strong canines on each side of vomer, these curved backward and outward; teeth on palatines distant, few in number (usually 4 or 5); among these are 1 to 3 very strong canines (usually, but not always, much larger than canines on vomer), strongly curved forward and inward. In *P. margaritatus* and *P. notatus*, the palatine teeth are not especially enlarged, subequal and more numerous, the canines on the vomer being much larger than any of the other teeth. Gill openings extending from the upper edge of pectoral to just below lower edge. Pectoral without axilliary foramen; height of soft dorsal about 3 in head; length of caudal nearly 2; height of anal $3\frac{1}{3}$; length of pectorals $1\frac{3}{5}$; of ventrals, $2\frac{2}{3}$. Color in life, light brown above, the top of head much darker and clouded with dark brown; a row of about 10 bar-like dark blotches along middle of side, each larger than eye, those anteriorly deeper than long, the others longer than deep; each of these blotches usually more or less confluent with a saddle-like dark blotch across the back; a crescent-shaped pale translucent area below the eye; below this a larger blue-black area, irregularly crescent-shaped, covering the preorbital and suborbital region, bounded below and behind by a row of shining mucous pores; on it are about 4 large pores, and above and behind it, close behind and below eye, is a large shining pore bordered with black; cheek steel bluish; sides of body silvery, becoming golden below; lower

part of head and belly bright golden; a dark stripe along base of dorsal; soft dorsal with 2 or 3 rows of small round dark olive spots, the upper row posteriorly becoming a dark edging to the fin; caudal, dull red, edged with dusky; anal very pale, edged with blackish; pectorals light orange, usually with some small dark spots above; ventrals orange, slightly darker anteriorly. Numerous series of pores on the body, those of the lateral line accompanied by shining golden bodies, as in other species of the genus. According to fishermen, these bodies are phosphorescent, shining at night; a statement which is probably true, although we have been unable to verify it; pores on sides of back not shining. Most of the pores, as in other species, accompanied by numerous small cirri or cilia; the arrangement of the lines of pores and shining bodies not very different from that found in *P. notatus*. It may be thus described in detail: A series of pores beginning at tip of snout, extending down around preorbital region, bounding the dark subocular blotch and joining almost at a right angle with a series of pores which extends downward from lower posterior corner of eye to angle of mouth. Another series diverges from the first in front of eye, passing close below eye, then upward above cheek, ending in a large pore behind preopercle. A curved series of pores extending backward along opercle, and another parallel with it along subopercle. Two obscure series from front of eye along top of head, becoming wide apart at the vertex, converging at the nape, then slightly diverging, converging in front of spinous dorsal, then again diverging to pass around the fin, each at last becoming straight at front of soft dorsal, extending close to its base to its last ray, there being about 2 pores to each ray. Just below this series, at front of soft dorsal on each side, begins a second series, with the pores wider apart and somewhat irregular, ceasing near the middle of the soft dorsal fin. The lateral line proper next begins above upper posterior angle of preopercle, whence a short branch passes directly upward. Opposite front of soft dorsal, the lateral line is interrupted for a distance a little more than diameter of eye. A short branch arises at this interruption and passes upward and backward at an angle from the end of the anterior part; thence the lateral line passes straight to base of caudal. The next series arises just behind axil of pectoral, then curves abruptly downward and backward, becoming straight opposite third ray of anal, thence proceeding to base of caudal, the pores small and close-set, anteriorly bead-like and shining, becoming dull toward the tail. Next comes a double series on each side of base of anal, the 2 series converging behind and finally coalescing. Another series begins at the middle of the base of the pectoral in front, curves downward, around the base of the fin, and, proceeding directly backward, ceases opposite vent. A series begins midway between gill opening and ventral and, extending straight backward, ceases opposite base of pectoral. Another begins, on each side, on lower side of head, directly below angle of mouth, the two diverging slightly between ventrals, then converging a little behind ventrals, then abruptly diverging, joining the series last mentioned, on each side, just in front of base of pectoral. A cross series of pores extends straight across belly, between

vent and anal fin. At each end of this cross series a series of pores turns abruptly forward, the two meeting in an acute angle on the belly just in front of a vertical from base of pectorals. Finally, 3 parallel series on each side of lower parts of head meet in front, the two anterior in obtuse curves, the posterior in an acute angle. The anterior series along the mandible ends at the corner of the mouth. The next just behind the mandible ends just below the corner of the mouth. The next passes along the branchiostegal region, ending at the gill opening. Mandible with 2 large foramina. A series of dark-colored pores along each side of tongue. Length 8 inches. South Carolina to Texas, and southward to Argentina, on sandy shores; not very common, and found in rather deep water. Not rare about Galveston, but unknown to fishermen at Pensacola. Here described from the types of *P. plectrodon*, the North American form. The types of *P. porosissimus* examined by us in Paris agree in dentition and other respects. Except for the remote locality there is no suggestion of differences. We are informed by Dr. Vaillant that the type specimen of *P. porosissimus* from St. Catherine, has 33 anal rays, that from Rio Janeiro 32, and that the number 27, given by Valenciennes for this species, represents an error in counting. According to Valenciennes, *P. porosissimus* has D. II-36; A. 27; each palatine bone with a row of small, pointed, unequal teeth; row of pores above anal reaching base of caudal. Color grayish brown above, silvery white below; dorsal and anal whitish, edged with brown; pectoral with longitudinal lines; ventrals brownish on the outer edge; caudal whitish at base, the rest brownish; some specimens with dark cross bands. (*porosissimus*, most porous.)

Batrachus porosissimus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 501, 1837, Surinam (Coll. Leschenault & Doumerc), Cayenne (Coll. Poiteau), Rio Janeiro (Coll. Delalande), St. Catherine (Coll. Lesson & Garnot).

Porichthys plectrodon, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 291, Galveston, Texas (Type, No. 30894. Coll. D. S. Jordan); JORDAN & GILBERT, Synopsis, 958, 1883.

Porichthys porosissimus, GÜNTHER, Cat., III, 176, 1861; JORDAN, Proc. Ac. Nat. Sci. Phila. 1883, 291; BERG, An. Mus. Nac. Buenos Aires, 1895, 70.

Porichthys porosissimus, JORDAN & GILBERT, Synopsis, 751, 1883; MEEK & HALL, Proc. Ac. Nat. Sci. Phila. 1885, 57.

2654. PORICHTHYS NOTATUS, Girard.

(SINGING FISH; MIDSHIPMAN; CABEZON; SAPO.)

Head $3\frac{2}{3}$; depth $6\frac{1}{2}$. D. II-37; A. 33; V. I, 2; P. 18; eye 8 in head; maxillary 2; pectoral $1\frac{3}{4}$; ventral $2\frac{1}{3}$; caudal $2\frac{2}{3}$. Head narrowed forward; opercle developed as a strong spine; maxillary reaching beyond orbit; lower jaw with a single row of about 10 large, recurved teeth, behind which is a patch of small teeth; sides of jaw with a single series of canines similar to those in front, but larger; upper jaw with an irregular series of small teeth; palatines with a single series of conical teeth; 2 large curved canines on vomer; head with several rows of fringed pores; 1 row along lower line of opercle and subopercle; another along upper edge of cheek, this branching behind and below the orbit, 1 branch running forward

below the orbit and around the snout, the other vertically downward behind the maxillary; a series of fringes behind the lower lip; behind this a series of pores without fringes; a short straight series of pores on each side of vertex; a row of pores along the base of the dorsal fin, curving at front of dorsal, and terminating at upper angle of opercle; a row below this, not reaching base of pectoral; the third row not reaching base of caudal, but ceasing at second third of anal to about its twentieth ray, and is anteriorly strongly curved upward to base of pectoral; 2 concentric series on the abdomen, the outer extending forward between bases of ventrals. The so-called "shining pores" on the sides are not pores, but bright round pieces of shiny membrane, showing through a translucent skin; each of the spots has above it a pair of fringed flaps with a small pore between them; the rows of flaps along dorsal and anal similar, long and low, their last rays reaching base of caudal rays; pectoral broad, somewhat pointed behind, reaching to the vertical from the fifth anal ray; origin of ventrals in advance of pectorals, in distance equal to length of maxillary, their tips not reaching to pectoral base; caudal well rounded. Olive brown above, with coppery reflections, the belly brassy-yellow; sides with irregular broad vertical cross blotches, most distinct in the young; dorsal grayish, with oblique dark bars; vertical fins sometimes margined with black; pores of lateral line bead-like, shining silvery; a white space below eye, with a black crescent below it; head yellowish brown, with no dark spots on opercle and shoulder; peritoneum black. Length 15 inches. Pacific coast; very abundant from Lower California to Puget Sound; living under stones, near the shore northward, in deeper water southward. It makes a peculiar humming noise with its air bladder, hence the name singing fish. (*notatus*, spotted; noted.)

Porichthys notatus, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 141, San Francisco; GIRARD, Pac. R. R. Surv., x, Fishes, 134, 1858.

Porichthys margaritatus, MEEK & HALL, Proc. Ac. Nat. Sci. Phila. 1885, 56; not of RICHARDSON.

Porichthys porosissimus, JORDAN & GILBERT, Synopsis, 751, 1883 (not of CUVIER & VALENCIENNES); GÜNTHER, Cat., III, 176, 1861 (in part).

2655. PORICHTHYS MARGARITATUS (Richardson).

Head $3\frac{1}{3}$ to $3\frac{2}{3}$; depth $4\frac{1}{3}$ to $5\frac{1}{2}$. D. II–37; A. 33. Similar to *Porichthys notatus*, differing chiefly in color. Top and sides of head and space above pectorals with numerous round dark brown spots and freckles, behind pectorals 6 to 8 vertical $\frac{1}{2}$ cross bars; dorsal not margined with black, but with 8 to 10 black submarginal spots; anal, with the exception of a few posterior rays, pale; caudal black at base and tip; pectorals with a few dots at base and on upper rays; a roundish white blotch below eye, below this a jet-black crescent. Palatine teeth small, 1 or 2 slightly enlarged. Series of shining spots arranged as in *P. notatus*, except that the third series extends almost to end of anal, to about its thirtieth ray. Pacific coast of tropical America. This species was obtained by the *Albatross* in large numbers off the west coast of Colombia, at Station 2795 at

a depth of 33 fathoms, and at Station 2802 at a depth of 16 fathoms. The largest specimens are about 4½ inches long. In dentition it agrees with *Porichthys notatus*, but in color and arrangement of spots it resembles *P. porosissimus*. (*margaritatus*, bearing pearls; *μάργαρος*.)

Batrachus margaritatus, RICHARDSON, Voyage Sulphur, Fishes, 67, 1845, Pacific coast of Central America; coloration and arrangement of lines identical with *porosissimus*.

Porichthys nautopœdium,* JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 171, Pacific Ocean, off coast of Colombia, Albatross Station, No. 2802, 8° 38′ N., 78° 31′ 30″ W., in 16 fathoms. (Type, No. 41145, U. S. Nat. Mus. Coll. *Albatross*.)

861. THALASSOPHRYNE, Günther.

(POISON TOAD-FISHES.)

Thalassophryne, GÜNTHER, Cat. Fishes, III, 174, 1861 (*maculosa*.).

Body rather elongate, compressed; head moderate. Dorsal spines 2;† soft dorsal and anal rather short, free from caudal; opercle very small, its posterior part developed as a single strong spine; subopercle feebly developed, narrowed and not ending in a spine; no scales on body. Spines hollow, and connected with venom glands. Lateral line on sides of body single; jaws without canine teeth. Species all South American, some of them ascending rivers; all of them noted for their venomous spines.‡ (*θάλασσα*, the sea; *φρύνη*, toad.)

* *ναυτοπαίδιον*, sailor-boy, from the common name "midshipman," a name given in allusion to the "buttons" on the belly of the fish.

† In *Thalassothia*, Berg, a South American genus, likewise with poison glands, 4 dorsal spines are present.

‡ The poison organs of *Thalassophryne reticulata* are thus described by Dr. Günther: "In this species I first observed and closely examined the poison organ with which the fishes of this genus are provided. Its structure is as follows: (1) The opercular part: The operculum is very narrow, vertically styliform, and very mobile; it is armed behind with a spine, 8 lines long in a specimen of 10½ inches, and of the same form as the venom fang of a snake; it is, however, somewhat less curved, being only slightly bent upward; it has a longish slit at the outer side of its extremity, which leads into a canal perfectly closed, and running along the whole length of its interior; a bristle introduced into the canal reappears through another opening at the base of the spine, entering into a sac situated on the opercle and along the basal half of the spine; the sac is of an oblong-ovate shape, and about double the size of an oat grain. Though the specimen had been preserved in spirits for about 9 months, it still contained a whitish substance of the consistency of thick cream, which on the slightest pressure freely flowed from the opening in the extremity of the spine. On the other hand, the sac could be easily filled with air or fluid from the foramen of the spine. No gland could be discovered in the immediate neighborhood of the sac; but on a more careful inspection I found a minute tube floating free in the sac, whilst on the left-hand side there is only a small opening instead of the tube. The attempts to introduce a bristle into this opening for any distance failed, as it appears to lead into the interior of the basal portion of the operculum, to which the sac firmly adheres at this spot. (2) The dorsal part is composed of the 2 dorsal spines, each of which is 10 lines long. The whole arrangement is the same as in the opercular spines; their slit is at the front side of the point; each has a separate sac, which occupies the front of the basal portion; the contents were the same as in the opercular sacs, but in somewhat greater quantity. A strong branch of the lateral line ascends to the immediate neighborhood of their base. Thus we have 4 poison spines, each with a sac at its base; the walls of the sacs are thin, composed of a fibrous membrane, the interior of which is coated over with mucous. There are no secretory glands embedded between these membranes, and these sacs are probably merely the reservoirs in which the fluid secreted accumulates. The absence of a secretory organ in the immediate neighborhood of the reservoirs (an organ the size of which would be in accordance with the quantity of fluid secreted), the diversity of the osseous spines which have been modified into poison organs, and the actual communication indicated by the foramen in the sac, lead me to the opinion that the organ of secretion is either that system of muciferous channels which is found in nearly the whole class of fishes, and the secretion of which has poisonous qualities in a few of them, or at least an independent portion of it. This description was

a. Dorsal and anal fins not joined to the caudal.

b. Dorsal and anal fins rather short; D. II-19; A. 18; pectoral fins short, their tips reaching to origin of anal. Color brown, marbled with darker; pectoral fins and sides of body with some round black spots; chin and ventrals brownish; belly white. MACULOSA, 2656.

bb. Dorsal and anal fins longer; D. II-24; A. 24; pectoral fins longer, their tips reaching to sixth anal ray. Color of head, body, and fins brown, with a network of yellowish lines; dorsal, anal, caudal, and pectoral fins with white margins. RETICULATA, 2657.

2656. THALASSOPHRYNE MACULOSA, Günther.

D.II-19; A.18; V.I, 2. The head is somewhat longer than broad, its length being contained $3\frac{1}{3}$ in the total; it is moderately depressed. The snout is short, obtuse, with the cleft of the mouth ascending obliquely upward, and with the chin prominent. The maxillary extends to the vertical from the posterior margin of the orbit. The teeth are obtusely conical, standing in single series, except anteriorly in the lower jaw, where they form 2 series, and in the upper, where they are cardiform, in a narrow band. The eyes are directed upward and very small, their width being $\frac{1}{4}$ of that of the bony bridge between the orbits. Gill covers with a single spine; it is long, slender, cylindrical, like one of the dorsal spines, and has the operculum for its base. Gill opening not very narrow; it extends from the upper

made from the first example; through the kindness of Captain Dow I received 2 other specimens, and in the hope of proving the connection of the poison bags with the lateral-line system, I asked Dr. Pettigrew, of the Royal College of Surgeons, a gentleman whose great skill has enriched that collection with a series of the most admirable anatomical preparations, to lend me his assistance in injecting the canals. The injection of the bags through the opening of the spine was easily accomplished; but we failed to drive the fluid beyond the bag, or to fill with it any other part of the system of muciferous channels. This, however, does not disprove the connection of the poison bags with that system, inasmuch as it became apparent that, if there be minute openings they are so contracted by the action of the spirit in which the specimens were preserved, as to be impassable to the fluid of injection. A great part of the lateral-line system consists of open canals; however, on some parts of the body, these canals are entirely covered by the skin; thus, for instance, the open lateral line ceases apparently in the suprascapular region, being continued in the parietal region. We could not discover any trace of an opening by which the open canal leads to below the skin; yet we could distinctly trace the existence of the continuation of the canal by a depressed line, so that it is quite evident that such openings do exist, although they may be passable only in fresh specimens. Thus, likewise, the existence of openings in the bags, as I believed to have found in the first specimen dissected, may be proved by examination of fresh examples. The sacs are without an external muscular layer, and situated immediately below the loose, thick skin which envelops their spines to their extremity; the ejection of the poison into a living animal, therefore, can only be effected by the pressure to which the sac is subjected the moment the spine enters another body. Nobody will suppose that a complicated apparatus like the one described can be intended for conveying an innocuous substance; and therefore I have not hesitated to designate it as poisonous; and, Captain Dow informs me in a letter lately received, 'the natives of Panama seemed quite familiar with the existence of the spines and of the emission from them of a poison which, when introduced into a wound, caused fever, an effect somewhat similar to that produced by the sting of a scorpion; but in no case was a wound caused by one of them known to result seriously. The slightest pressure of the finger at the base of the spine caused the poison to jet a foot or more from the opening of the spine.' The greatest importance must be attached to this fact, inasmuch as it assists us in our inquiries into the nature of the functions of the muciferous system, the idea of its being a secretory organ having lately been superseded by the notion that it serves merely as a stratum for the distribution of peripheric nerves. Also the objection that the Stingrays and many Siluroid fishes are not poisonous, because they have no poison organ, can not be maintained, although the organs conveying their poison are neither so well adapted for this purpose nor in such a perfect connection with the secretory mucous system as in *Thalassophryne*. The poison organ serves merely as a weapon of defense. All the Batrachoids with obtuse teeth on the palate and in the lower jaw feed on Mollusca and Crustaceans." (Günther.)

base of the pectoral obliquely downward and forward to the level of the inferior base of the pectoral. The 2 dorsal spines are slender, pungent, about $\frac{1}{3}$ the length of the head. Dorsal and anal fins terminate immediately before the root of the caudal, the length of which is $\frac{1}{7}$ the total; pectoral obliquely rounded, extending to the origin of the anal; ventral rather short, not quite $\frac{1}{2}$ the length of the head, extending to the base of the pectoral. Skin perfectly smooth, with some very short tentacles at the lower jaw. Two short horizontal muciferous channels on the cheek and the lateral line are very distinct; they are not, as usually, composed of a series of distant pores, but the pores are confluent, forming 1 continuous groove of a white color. Other muciferous channels, as for instance along the base of the anal, are composed of separate indistinct pores. Color brown, marbled with darker; pectoral fins and sides of the body with some round black spots; chin and ventrals brownish; belly white. The general habit is that of a *Batrachus* [*Opsanus*]. One specimen, from Puerto Cabello, Caribbean Sea. (Günther.) (*maculosus*, spotted.)

Thalassophryne maculosa, GÜNTHER, Cat., III, 175, 1861, Puerto Cabello; GÜNTHER, Fishes of Centr. Amer., 436, pl. 68, fig. 1, 1869; MEEK & HALL, Proc. Ac. Nat. Sci. Phila. 1885, 54.

2657. THALASSOPHRYNE RETICULATA, Günther.

D. II-24; A. 24; V. I, 2; P. 16. The length of the head is $\frac{2}{7}$ of the total length (without caudal). The teeth on the palate are in a single series, very short, obtuse, incisor-like. Pectoral very large, extending back to the sixth anal ray. Head, body, and fins brown, with a network of yellowish lines; vertical and pectoral fins with a white margin. In other respects this species agrees with *T. maculosa*. Length 13 inches. Panama; not rare. (*reticulatus*, netted.)

Thalassophryne reticulata, GÜNTHER, Proc. Zool. Soc. London 1864, 150, 155, Panama; GÜNTHER, Fish. Centr. Amer., 437, pl. 68, fig. 2, 1869; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 62; MEEK & HALL, Proc. Ac. Nat. Sci. Phila. 1885?.

862. DÆCTOR, Jordan & Evermann, new genus.

(POISON TOAD-FISHES.)

Dæctor, JORDAN & EVERMANN, new genus (*dowi*).

This genus differs from *Thalassophryne* in the more elongate body and the many-rayed soft dorsal and anal fins, the last rays of which are fully joined to the caudal. (δαίκτωρ, slayer; from δαίζω, to slay.)

2658. DÆCTOR DOWI (Jordan & Gilbert).

Head 4 in length ($4\frac{2}{3}$ with caudal); depth $5\frac{2}{3}$ ($6\frac{2}{3}$). D. II-33; A. 30. Body comparatively elongate, compressed behind. Head low and rather narrow, its width $1\frac{1}{3}$ in its length. Eye very small, the diameter not $\frac{1}{2}$ the interorbital space, and about as long as snout, 8 in head. Interorbital width about $5\frac{1}{2}$ in head. Opercular spine short, nearly 4 in head. Mouth oblique, the lower jaw much projecting. Maxillary 2 in head, extending

to beyond eye. Teeth small, those on the palatine largest; teeth of upper jaw smaller than those of the lower; anterior teeth of the lower jaw in about 2 series. Pectoral fins long, $1\frac{1}{5}$ in head, reaching about to fifth anal ray; last rays of dorsal and anal fully joined to the caudal. Color olivaceous, with darker blotches; first dorsal black; under parts pale; posterior portion of anal edged with dark. Pacific coast of North America, from Punta Arenas to Panama; rare. (Named for Capt. John M. Dow, who obtained a fine specimen (now destroyed) from Panama.)

Thalassophryne dowi, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1887, 388, Punta Arenas (Type, No. 39085, U. S. Nat. Mus. Coll. Cornell University); JORDAN, Proc. Cal. Ac. Sci. 1896, 231, pl. 38.

Suborder XENOPTERYGII.

(THE CLING-FISHES.)

Breast with a broad sucking disk, between the wide-set ventral fins, this formed from the skin of the breast, not from the ventral fins themselves. Ventral rays I, 4 or I, 5; no scales; no spinous dorsal; no suborbital ring; palatine arcade materially modified; no air bladder; vertebræ in increased numbers; gill arches reduced. A well-marked group of small fishes, constituting a single family. (ξένος, strange; πτέρυξ, fin.)

Family CXCIX. GOBIESOCIDÆ.

(CLING-FISHES.)

Body rather elongate, tadpole-shaped, broad and depressed in front, covered by smooth, naked skin; mouth moderate; upper jaw protractile; teeth usually rather strong, the anterior conical or incisor-like; posterior canines sometimes present; suborbital ring wanting; no bony stay from suborbital across cheek; opercle reduced to a spine-like projection concealed in the skin, behind the angle of the large preopercle, this spine sometimes obsolete; pseudobranchiæ small or wanting, gills 3 or $2\frac{1}{2}$; gill membranes broadly united, free or united to the isthmus; dorsal fin on the posterior part of the body, opposite to the anal and similar to it, both fins without spines; ventral fins wide apart, each with 1 concealed spine and 4 or 5 soft rays. Between and behind the ventrals is a large sucking disk, the ventrals usually forming part of it. This sucking disk, which is wholly different in structure from that of *Cyclopterus* and *Liparis*, is thus described by Dr. Günther: "The whole disk is exceedingly large, subcircular, longer than broad, its length being (often) $\frac{1}{4}$ of the whole length of the fish. The central portion is formed merely by skin, which is separated from the pelvic or pubic bones by several layers of muscles. The peripheric portion is divided into an anterior and posterior part by a deep notch behind the ventrals. The anterior peripheric portion is formed by the ventral rays, the membrane between them and a broad fringe, which extends anteriorly from one ventral to the other. This fringe is a fold of the skin containing on one side the rudimentary ventral spine, but

no cartilage. The posterior peripheric portion is suspended on each side on the coracoid, the upper bone of which is exceedingly broad, becoming a free, movable plate behind the pectoral. The lower bone of the coracoid is of a triangular form, and supports a very broad fold of the skin, extending from one side to the other, and containing a cartilage which runs through the whole of that fold. Fine processes of the cartilage are continued into the soft striated margin, in which the disk terminates posteriorly. The face of the disk is coated with a thick epidermis, like the sole of the foot in higher animals. The epidermis is divided into many polygonal plates. There are no such plates between the roots of the ventral fins." (Günther, Cat., III, 495.) No air bladder; intestines short; pyloric cæca few or none; skeleton firm; vertebræ 13 or 14+13 to 22=26 to 36. Carnivorous fishes of small size, chiefly of the warm seas, usually living among loose stones between the tide marks and clinging to them firmly by means of the adhesive disk. Their relations are obscure, but they are probably descended from allies or ancestors of the *Cottidæ* or *Batrachoididæ*. Genera about 15; species 50. The principal genus is *Gobiesox*. (*Gobesocidæ*, Günther, Cat., III, 489–515.)

GOBIESOCINÆ:

a. Gill membranes free from the isthmus; gills 3; posterior part of sucking disk with no free anterior margin.

b. Incisors of lower jaw with entire edges.

c. Vertebræ about 32; anal fin long, nearly as long as dorsal. CAULARCHUS, 863.

cc. Vertebræ about 26; anal fin short.

d. Dorsal fin very long, of about 17 rays, twice as long as the moderate anal, which has 8 or 9 rays; disk broad; upper teeth in several rows. BRYSSETÆRES, 864.

dd. Dorsal fin moderate or short, of 4 to 13 rays.

e. Disk more or less broad, its length 2½ to 3 in body; dorsal and anal not very short, their rays 6 or more; body tapering rapidly backward; opercular spine strong. GOBIESOX, 865.

ee. Disk very narrow, its width 4 to 5 in body; head short, 3½ to 4 in body; dorsal and anal very short and small; a patch of teeth in each jaw behind the large teeth; sucking disk small. RIMICOLA, 866.

bb. Incisors of lower jaw tricuspid or serrate; dorsal and anal fins short; vertebræ about 28. ARBACIOSA, 867.

863. CAULARCHUS, Gill.

Caularchus, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 330 (*mæandricus*).

This genus differs from *Gobiesox* chiefly in the numerous vertebræ, 32 in the only species known. The incisors are entire, the anal fin similar to the dorsal, each having 12 or 13 rays. The single species reaches a large size and is found farther north than any other of the group, a fact in accord with the increased number of vertebræ. (καυλός, stem; ἀρχός, anus; from the many-rayed anal.)

2659. CAULARCHUS MÆANDRICUS (Girard).

(Suck-fish.)

Head $2\frac{3}{4}$; depth $6\frac{1}{3}$. D. 13; A. 12; V. I, 4; vertebræ 13+19=32; eye $7\frac{1}{2}$ in head; distance from vent to caudal $2\frac{3}{4}$ in length of body; sucking disk as broad as long, $3\frac{1}{2}$ in length. Head broad, nearly circular when viewed from above; interorbital width 3 in head; mouth wide, its width more than $\frac{1}{2}$ length of head; maxillary extending to below eye; outer teeth of upper jaw rather strong, close set, vertical, conical, or slightly compressed, a narrow band of small, conical teeth behind them; lower jaw with larger teeth, 6 or 8 of the anterior broad, incisor-like, with entire edges, placed nearly horizontally; lateral and posterior teeth small, as in upper jaw; nostrils ending in tubes; spine on opercle sharp, but not projecting through the skin; origin of dorsal fin a little in advance of vent, the fin much higher than the anal; vent midway between anal and posterior edge of disk; pectorals short and broad, not extending back past the margin of the ventral disk, the 3 lower rays forming part of disk; caudal rounded. Color light olive, everywhere reticulated with brownish orange; middle of upper lip black; a light bar between eyes and 1 across cheek; vertical fins dusky; caudal with 2 faint brownish bars near its base. Specimens from red algæ are light pink, mottled with darker, the pale band between eyes very distinct. Length 6 inches. Pacific coast of United States, from Vancouver Island to Point Concepcion; everywhere very abundant in rock pools; the largest species of *Gobiesocidæ*. (*mæandricus*, meandering, in allusion to the reticulated streaks.)

Lepadogaster reticulatus, Girard, Proc. Ac. Nat. Sci. Phila. 1854, 155, San Luis Obispo, California; name preoccupied.

Lepidogaster mæandricus, Girard, Pacific R. R. Surv., x, Fishes, 130, 1858, San Luis Obispo, California; substitute for *reticulatus*, preoccupied in *Lepadogaster;* Günther, Cat., iii, 505, 1861.

Gobiesox reticulatus, Jordan & Gilbert, Synopsis, 749, 1883.

864. BRYSSETÆRES, Jordan & Evermann.

Bryssetæres, Jordan & Evermann, Proc. Cal. Ac. Sci. 1896, 230 (*pinniger*).

This genus differs from *Gobiesox* solely in the great development of the dorsal fin, which has 17 rays, the moderate anal having but 8 or 9; the vertebræ 10+16, as usual in *Gobiesox*. One species known. (*βρύσσος*, sea-urchin; *ἕταιρος*, comrade, the species living in rock pools with the sea-urchins.)

2660. BRYSSETÆRES PINNIGER (Gilbert).

Head $2\frac{3}{5}$ to $2\frac{4}{5}$; width of body $4\frac{1}{3}$; of head 3 in length. D. 16 or 17; A. 8 or 9. Interorbital width $3\frac{1}{4}$ in head; eye $\frac{1}{3}$ interorbital width; width of mouth $1\frac{3}{5}$ to $1\frac{4}{5}$ in head. Teeth in upper jaw conic, acute, in several series, the anterior row in front enlarged, unequal; in lower jaw the teeth mesially in 2 distinct series, those in middle of anterior row narrow, entire incisors, those laterally conic, canine-like. No evident oper-

cular spine. Disk about as broad as long, its length about that of head. Front of dorsal varying in position, about midway between snout and base of caudal, its length about ½ that of body. Vent nearly equidistant between disk and front of anal, the base of the latter 3½ to 3¾ in body. Caudal 1½ to 1¾ in head. Pectorals about ⅓ of head, with a distinct fold at base. Color variable; anteriorly usually with reticulating dark lines surrounding yellowish spots; a narrow dark streak forward, 1 downward, and 1 backward from orbit; below dorsal fin about 6 dark bars running obliquely downward and backward, these sometimes in greater number, frequently more or less irregular and interconnected, often divided by vertical streaks or series of dots; body sometimes light in spirits, with out distinctive markings; vertical fins usually dusky, narrowly margined with white, sometimes lighter with dark margins. Length 2¼ inches. (Gilbert.) Gulf of California, abundant; specimens known from Puerto Refugio (Angel Island), San Luis Gonzales Bay, and La Paz. Well distinguished by its long dorsal. (*pinniger*, fin-bearing.)

Gobiesox pinniger, GILBERT, Proc. U. S. Nat. Mus. 1890, 94, Puerto Refugio, Gulf of California. (Coll. *Albatross*.)
Brysaetæres pinniger, JORDAN, Proc. Cal. Ac. Sci. 1896, 230, pl. 34.

865. GOBIESOX, Lacépède.

(CLING-FISHES.)

Gobiesox, LACÉPÈDE, Hist. Nat. Poiss., II, 595, 1799 (*cephalus*).
Megaphalus, RAFINESQUE, Analyse de la Nature 1815, 86 (*cephalus*, substitute for *Gobiesox*, regarded as an objectionable compound).
Sicyases, MÜLLER & TROSCHEL, Archiv fur Naturgesch. 1843, 298 (*sanguineus;* small species, with upper teeth uniserial).
Tomicodon, BRISOUT DE BARNEVILLE, Rev. Zool., 144, 1846 (*chilensis* = *Sicyases*).
Sicyogaster, BRISOUT DE BARNEVILLE, Rev. Zool., 144, 1846 (*marmoratus* = *Gobiesox*).
Bryssophilus, JORDAN & EVERMANN, new subgenus (*papillifer*).

Body anteriorly very broad and depressed, posteriorly slender, covered with tough, smooth skin; opercle with a strong spine; head large, rounded in front; mouth terminal, crescent-shaped; lower jaw with a series of strong incisors in front, their edges rounded or truncate; upper jaw with a series of strong teeth, behind which are sometimes smaller teeth; no teeth on vomer or palatines; gills 3; gill membranes broadly united under the throat, not attached to the isthmus; sucking disk large, the posterior portion without anterior free margin. Dorsal and anal moderate, the dorsal rays 6 to 12, the anal rays 6 to 10. Vertebræ about 26, as far as known. Species numerous, all American; mostly tropical, clinging to rocks near the shore. (*Gobius; Esox;* the resemblances either to the goby or the pike being few or remote.)

BRYSSOPHILUS (βρύσσος, sea urchin; φιλέω, to love):

a. Dorsal fin comparatively long, of about 13 rays; anal rays 9; disk broad; upper teeth in several rows; lower incisors narrow; papillæ below chin; color olivaceous. PAPILLIFER, 2661.

aa. Dorsal fin moderate or short, its rays 6 to 11.

GOBIESOX:

c. Upper teeth in more than 1 series (character not verified in a few species); head broad.

d. Coloration in life chiefly olivaceous, without red, sometimes banded with darker or paler.

e. Dorsal rays 12; anal rays 7. GYRINUS, 2662.

ee. Dorsal rays 11; anal rays 6; fins black. NIGRIPINNIS, 2663.

eee. Dorsal rays 9 or 10; anal rays 6. CEPHALUS, 2664.

eeee. Dorsal rays 8; anal rays 6. TUDES, 2665.

eeeee. Dorsal rays 11; anal rays 10. STRUMOSUS, 2666.

eeeeee. Dorsal rays 10; anal rays 8. VIRGATULUS, 2667.

eeeeeee. Dorsal rays 9; anal rays 7.

f. Width of head $3\frac{2}{3}$ in length; color plain brown. ADUSTUS, 2668.

ff. Width of head 5 in length; color blackish, with yellow vermiculations. FUNEBRIS, 2669.

eeeeeeee. Dorsal rays 7; anal rays 7; eyes variegated. PŒCILOPHTHALMUS, 2670.

dd. Coloration in life chiefly bright red, or else with red spots or bands, the color not fading in spirits.

g. Color red, with deep red spots. D. 6; A. 5. RHODOSPILUS, 2671.

gg. Color uniform red, unspotted, the color not fading in spirits; dorsal rays 6 to 8; anal rays 6.

h. Lower jaw with short incisors on each side, followed by canines. MACROPHTHALMUS, 2672.

hh. Lower jaw with 2 horizontal incisors on each side, the third horizontal tooth not incisor-like; no distinct canines. CERASINUS, 2673.

SICYASES (*σικύα*, a sucking cup made of a gourd):

cc. Upper teeth in a single series (character not verified on some species); dorsal and anal short.

i. Color chiefly red.

j. Body with cross bands of deep red; iris red; dorsal rays 6; anal rays 5; head broad, the eyes very large. ERYTHROPS, 2674.

jj. Body with dark cross bands and with spots of clear blue; body rather slender. D. 6 or 7; A. 6. RUBIGINOSUS, 2675.

jjj. Body plain, light red; form rather slender. CARNEUS, 2676.

ii. Color olivaceous or brownish, not red.

k. Dorsal rays 9; anal rays 6.

l. Color olivaceous, without bands. HÆRES, 2677.

ll. Color greenish, with 3 dark cross bands and many dots. PUNCTULATUS, 2678.

kk. Dorsal rays 7; anal rays 7; body with dark cross bands. FASCIATUS, 2679.

Subgenus **BRYSSOPHILUS**, Jordan & Evermann.

2661. GOBIESOX PAPILLIFER, Gilbert.

Head $2\frac{2}{5}$; width of body $3\frac{2}{3}$; width of head $2\frac{1}{5}$. D. 13; A. 9. Width of mouth $1\frac{1}{4}$ in head; interorbital width 3; eye $\frac{1}{5}$ interorbital width; teeth in upper jaw conic, acute, very small, in 2 or more series, 2 of them slightly enlarged, canine-like; teeth in lower jaw in 2 series, the outer anteriorly, narrow entire incisors, with rounded tips, becoming conical laterally; opercular spine sharp, evident, though not projecting through the integument; lips and lower side of head anteriorly with fleshy papillæ; disk about as broad as long, its length $1\frac{1}{5}$ in head; distance from

front of dorsal to base of caudal $1\frac{1}{2}$ in its distance from tip of snout; vent exceptional in position, immediately in front of anal fin; base of anal $1\frac{3}{4}$ in head; caudal rather acute, $1\frac{3}{5}$ in head; pectorals $2\frac{3}{5}$ in head, a distinct fleshy fold at base. In spirits, uniform dark olivaceous, lower side of head and disk light; pectorals dusky; vertical fins with a black bar at base, then a white bar, followed by a wide, dusky area, and narrowly margined with white; caudal with all these marks except the black bar, having the posterior outlines curved, following margin of fin. Length $1\frac{1}{2}$ inches. Magdalena Bay, Lower California. (Gilbert.) Possibly related to the genus *Caularchus*. (*papilla; fero*, I bear.)

Gobiesox papillifer, GILBERT, Proc. U. S. Nat. Mus. 1890, 96, Magdalena Bay, Lower California. (Coll. *Albatross*.)

Subgenus GOBIESOX.

2662. GOBIESOX GYRINUS, Jordan & Evermann, new species.

B. 6; D. 12; A. 7; V. I, 4; P. 20. A vertical fold of skin at base of pectoral; coracoid distinctly below level of upper margin of pectoral; teeth of upper jaw cardiform, lower jaw with very narrow but compressed incisors, which are as short as the other teeth. Lateral profile of head nearly semicircular; head much depressed, as long as broad, its length being $\frac{2}{7}$ of the total; width of interorbital space somewhat less than $\frac{1}{2}$ greatest width of head, or 3 times diameter of eye; cleft of mouth extending beyond anterior margin of eye; distance of origin of dorsal from caudal more than $\frac{1}{2}$ of its distance from snout. Brownish, with scattered dark spots; a black blotch anteriorly on the dorsal fin. Length 3 inches. (Günther.) West Indies; not seen by us. A valid species, according to Dr. Günther, but apparantly as yet without tenable specific name, as the original *Cyclopterus nudus*, Linnæus, must have been some other fish. (*gyrinus; γυρῖνος*, a tadpole.)

Lepadogaster nudus, BLOCH & SCHNEIDER, Syst. Ichth., 2, 1801; in part, description taken from *Cyclopterus nudus*, Linnæus, except the count of fin rays. D. 12; A. 6.

Cotylis nuda, MÜLLER & TROSCHEL, Hor. Ichth., III, 18, pl. 3, f. 2,

Gobiesox nudus, GÜNTHER, Cat. Fish., III, 502, 1861, Island of Cordova. (Coll. G. U. Skinner.)

Gobiesox gyrinus, JORDAN & EVERMANN, Check-List Fishes, 491, 1896, Cordova; after GÜNTHER; name only.

2663. GOBIESOX NIGRIPINNIS (Peters).

D. 11; A. 6; P. 22. "Nostrils, mouth, teeth, opercular spine, and fin rays as in *Cotylis stannii* (*Gobiesox cephalus*), but the dorsal fin longer. Light brown above (minutely dotted with black, if viewed by a magnifier); vertical fins black." Puerto Cabello (Peters); not seen by us; a doubtful species, perhaps identical with *G. cephalus* or *G. nudus*. (*niger*, black; *pinna*, fin.)

Cotylis nigripinnis, PETERS, Berl. Monatsber. 1859, 412, Puerto Cabello.

Gobiesox nigripinnis, GÜNTHER, Cat., III, 502, 1861; after PETERS; GÜNTHER, Fish. Centr. Amer., 390, 1869.

2664. GOBIESOX CEPHALUS, Lacépède.

(TÉTARD; TESTAR.)

D. 9 or 10; A. 6; C. 12; P. 19 or 20. Head and anterior part of body very broad, much depressed; skin tough, naked, and smooth; head nearly as broad as long, with its profile semielliptical, the snout being very obtuse and rounded. The upper surface of the head is quite flat, gently sloping downward in a straight line from the nape to the snout. The greatest width of the interorbital space is ½ of that of the head, or 4 times the diameter of the eye. The cleft of the mouth is horizontal, curved, wide, extending to below the center of the eye; the lips are thick, the lower being divided into 5 portions by 4 vertical grooves, the central portion being the smallest, the lateral ones the largest and hanging downward. The upper jaw is slightly protractile, and there is a broad velum behind the teeth in each jaw. A band of short conical teeth in the upper jaw; a single series in the lower, the anterior ones being slightly compressed incisors, and small like the lateral teeth, which are conical. The eye is small, situated immediately below the upper profile of the head. Two nostrils, close together, opposite the upper angle of the orbit, their margins being slightly raised. The lower angle of the opercular apparatus terminates posteriorly in an obtuse movable point enveloped in skin and directed backward. The gill openings are somewhat narrow in consequence of the small degree of expansibility of the gill covers, but the gill membranes have the margin quite free, being united together under the throat, and not attached to the isthmus. There are only 3 gills; the pseudobranchiæ are quite rudimentary, indicated by 2 or 3 short lamellæ. The distance of the origin of the dorsal fin from the caudal is nearly ⅛ of its distance from the snout, its first ray is much shorter than the others, and apparently without articulations. The caudal rounded and of moderate length; the anal is only ½ as long as the dorsal, commencing below its middle and terminating in the same vertical. The pectoral is broad and short, its lower ½ being longer than the upper; it is slightly connected with the ventral. A vertical fold of the skin at the base of the pectoral; the coracoid is so high as to reach to the upper margin of the pectoral. The adhesive apparatus as broad as long, its length being contained 3½ times in the total. The vent and the porus urogenitalis are close together, situated midway between the margin of the ventral disk and the anal. The anal papilla is small. The color is brown (in spirits), whitish inferiorly. Length of adult, 7 inches. (Günther.) Caribbean Sea, said to be common; not seen by us. The original *G. cephalus* seems nearer the next species, if the 2 are really different. If that be the case the present species may stand as *Gobiesox stannii*. But we have no material adequate to settle this question. (*cephalus*, big-headed; *κεφαλή*, head.)

Gobiesox cephalus, LACÉPÈDE, Hist. Nat. Poiss., II, 595, 1798, Martinique; on a drawing by PLUMIER; D. 8; A. 4 or 5; color plain reddish; anal inserted behind dorsal; head broad; eyes blue; GÜNTHER, Cat., III, 499, 1861.

Lepadogaster testar, BLOCH & SCHNEIDER, Syst. Ichth., 445, 1801, Martinique; after PLUMIER.

Cotylis stannii, MÜLLER & TROSCHEL, Hor. Ichthyol., III, 18, taf. 3, fig. 3, 1845.

2665. GOBIESOX TUDES, Richardson.

Head $2\frac{1}{2}$; depth $4\frac{2}{3}$; width of head $2\frac{1}{4}$. D. 8; A. 6 in plate (5 in the description, the first short ray apparently not counted by Richardson). Head very broad, as broad as long, abruptly truncated anteriorly; mouth large, the maxillary reaching front of eye; lower jaw included; teeth entire; eye large, $4\frac{3}{4}$ in head, a little more than $\frac{1}{2}$ interorbital width, $1\frac{1}{2}$ in snout. Distance from front of dorsal to caudal about equal to length of head; insertion of dorsal before vent; the anal behind dorsal and much shorter than it; pectorals short. Color uniform, probably greenish, without spots or stripes. Length 5 inches. Locality "unknown, but supposed to be from China." (Richardson.) The species is, however, certainly not Chinese and is more likely to be from the West Indies. This species differs from *Gobiesox cephalus*, as described by Günther, in the larger eye and shorter dorsal. It is probably the same species. (*tudes*, hammer.)

Gobiesox tudes, Richardson, Voy. Sulphur, Fish., 103, pl. 46, figs. 1–3, 1845, habitat unknown, erroneously supposed to be China.

2666. GOBIESOX STRUMOSUS, Cope.

D. 11; A. 10; C. 16; P. 21. Head extremely wide, its width $2\frac{5}{6}$ in total length; this width partly produced by a large fleshy mass extending from end of maxillary to end of interopercle; eye small; profile of head descending abruptly from posterior line of orbits. Superior dental series 12 on each side, externally, but the 3 median teeth conceal some series of which the second 3 external teeth are a continuation; inferior teeth 11 on each side; 4 median incisors horizontal and subequal; no marked canine. Bluish plumbeous, fins blackish. (Cope.) Hilton Head, South Carolina, and Indian River, Florida; 4 specimens recently taken at Titusville by Evermann & Bean; apparently distinguished from *G. virgatulus* by its longer anal. (*strumosus*, from struma, a scrofulous tumor, alluding to the swollen cheek.)

Gobiesox strumosus, Cope, Proc. Ac. Nat. Sci. Phila. 1870, 121, Hilton Head, South Carolina; Jordan & Gilbert, Synopsis, 749, 1883; Evermann & Bean, Fishes of Indian River, Florida, in Rept. U. S. Fish Comm. 1896, 248.

2667. GOBIESOX VIRGATULUS, Jordan & Gilbert.

Head $2\frac{3}{4}$ ($3\frac{3}{5}$ with caudal); width of head $3\frac{1}{4}$; depth 6 (7 in total). D. 10; A. 8 or 9; vertebræ $10+16=26$. Body rather slender, the head low and rather broad, broadly rounded anteriorly; eyes very small, about 4 to 6 in head, about $2\frac{1}{2}$ in interorbital width; interorbital space broad, slightly convex. Cheeks prominent; opercle ending in a sharp spine. Cleft of mouth extending to below front of orbit; lower jaw somewhat shorter than upper. Teeth of upper jaw in a narrow band of about 2 series; 4 teeth of outer series a little larger than the rest, somewhat canine-like; middle teeth of lower jaw incisor-like and partly horizontal, their edges entire or somewhat concave. Ventral disk considerably shorter than head. Distance from root of caudal to front of dorsal $2\frac{4}{5}$ in length. Pectoral

short, about $2\frac{2}{3}$ in head. Color in life olivaceous, with numerous paler spots and broad diffuse dark bars; the whole body covered with rather faint, wavy, longitudinal stripes or lines of a light orange-brown color, about as wide as the interspace, much as in some species of *Liparis*, these entirely disappearing in alcohol; skin everywhere with dark punctulations; caudal dusky, slightly barred with paler, its tip abruptly yellowish; dorsal and anal dusky, the darker parts corresponding to dark bars on the body, barred. A rather large species. Length 2 to 4 inches. Common among ballast rocks, from Pensacola Bay north to Charleston. Our specimens from Pensacola and Charleston. (*virgatulus*, narrowly striped.)

Gobiesox virgatulus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 293, Pensacola, Florida (Coll. Jordan & Stearns); JORDAN & GILBERT, Synopsis, 958, 1883; GOODE & BEAN, Proc. U. S. Nat. Mus. 1882, 236; JORDAN, Proc. U. S. Nat. Mus. 1884, 149.

2668. GOBIESOX ADUSTUS, Jordan & Gilbert.

Head 3; depth $5\frac{1}{4}$. D. 9; A. 7. Head and body broad and flat, much depressed; width of head nearly equal to its length, $3\frac{3}{8}$ in body. Incisors in middle of lower jaw entire, broad; those in upper jaw narrow, blunt, little compressed, entire, shorter than the lateral teeth; behind these 2 or 3 series of smaller teeth. Eyes rather large, separated by a broad interorbital space, which is $\frac{1}{3}$ length of head and about $\frac{1}{2}$ greater than diameter of eye. Opercular spine sharp. Pectoral short, about $\frac{1}{2}$ length of head; ventral disk as long as head; distance from base of caudal to front of dorsal equaling $\frac{3}{10}$ of the length; caudal rounded behind. Brown, banded with blackish on body, head marbled with darker brown; front of dorsal black, the fins dusky with darker points. Pacific Coast of Mexico. Three specimens, the largest about 2 inches long, were obtained in a tide pool at Mazatlan. (*adustus*, scorched; brown.)

Gobiesox adustus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 360, Mazatlan, Mexico; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 627; JORDAN & GILBERT, Bull. U. S. Fish Comm. 1882, 108.

2669. GOBIESOX FUNEBRIS, Gilbert.

Body rather slender, its width 5 in length; width of head $3\frac{1}{2}$ to nearly 4; head $2\frac{3}{4}$ to 3 in length; depth $\frac{1}{2}$ head. D. 9; A. 6 or 7. Teeth in upper jaw conical, in several series, unequal but without canines; in lower jaw mesially in 2 series, the outer of narrow, entire incisors, truncate or rounded, without lateral canines. Interorbital space wide, 3 in head, the eye small, $\frac{2}{5}$ interorbital width. Mouth very wide, $\frac{1}{2}$ or more than $\frac{1}{2}$ length of head. Ventral disk wider than long, its length $1\frac{1}{2}$ to $1\frac{3}{4}$ in head. No evident opercular spine. Distance from front of dorsal to base of caudal $2\frac{3}{4}$ to 3 in length anterior to dorsal; distance from vent to front of anal fin $1\frac{1}{2}$ to $1\frac{3}{4}$ in distance from vent to disk; base of dorsal from $1\frac{1}{2}$ to $1\frac{1}{4}$ in head; base of anal about $\frac{1}{2}$ head; caudal rounded, $1\frac{3}{5}$ to $1\frac{3}{4}$ in head; pectorals $3\frac{1}{4}$ in head. Color varying from dark olive brown to black, everywhere covered with fine, yellowish vermiculations, usually arranged to

form narrow lighter bars on the sides; 3 or 4 obscure dark streaks radiating from the eye; blackish below, the fins varying from blackish to straw color. Length 2½ inches. (Gilbert.) Gulf of California; abundant at Puerto Refugio (Angel Island) and La Paz. (*funebris*, funereal, from the dark color.)

Gobiesox funebris, GILBERT, Proc. U. S. Nat. Mus. 1890, 95, Puerto Refugio, Gulf of California. (Coll. *Albatross*.)

2670. GOBIESOX PŒCILOPHTHALMUS, Jenyns.

Head 3, as wide as long. D. 7; A. 7. Opercular spine long and slender; teeth strong, somewhat crowded in front, the anterior in both jaws incisor-like; upper teeth conical, with smaller ones behind; 6 middle teeth of lower jaw incisor-like, projecting forward, their form not described (probably entire). Eyes large, close together, less than a diameter apart. General color olivaceous or brownish white, unmarked; iris golden, with pink and blue. Length 1⅝ inches. (Jenyns.) Chatham Island, Galapagos; only the single type known. (*ποικίλος*, variegated; *ὀφθαλμός*, eye.)

Gobiesox pœcilophthalmus, JENYNS, Voy. Beagle, Fishes, 141, pl. 27, figs. 2, 2a, 2b, 1842, Chatham Island (Coll. Darwin); GÜNTHER, Cat., III, 503, 1861.

2671. GOBIESOX RHODOSPILUS, Günther.

D. 6; A. 5; C. 8 or 9; P. 16. A vertical fold of skin along lower half of base of pectoral. Distance from front of dorsal to caudal 2⅜ in its distance from snout; anal before third dorsal ray. A very narrow band of short conical teeth in upper jaw, 1 lateral tooth larger than the others, recurved, canine-like; lower jaw with 1 series of teeth, the anterior narrow incisors, the outer distinctly canine, like the outer above. Rose-colored with rose-red transverse spots, each with an edge of deep-red dots. Panama. (Günther.) Not seen by us; known from 2 specimens, each 1½ inches long. (*ῥόδον*, rose; *σπίλος*, spot.)

Gobiesox rhodospilus, GÜNTHER, Proc. Zool. Soc. Lond. 1864, 25, Panama (Coll. Captain Dow); GÜNTHER, Fish. Centr. Amer., 445, 1869.

2672. GOBIESOX MACROPHTHALMUS, Günther.

Eye 4½ in head. D. 8; A. 6; C. 12; P. 22. Head and anterior part of body very broad and much depressed, the head as broad as long, its profile semi-elliptical, the snout obtuse and rounded; top of head quite flat; interorbital width equal to eye. Mouth horizontal, curved, moderate, the cleft reaching beyond anterior margin of eye; an acute spine at lower angle of opercle; 4 short incisors on each side in lower jaw, separated from the conical lateral teeth by a larger canine-like tooth. Insertion of dorsal nearer caudal than snout; caudal rounded. Color, uniform reddish. (Günther.) Locality unknown; probably West Indies. (*μακρός*, large; *ὀφθαλμός*, eye.)

Dr. Eigenmann gives the following notes on a small specimen from St.

Thomas, which seems referable to *Gobiesox macrophthalmus*, differing in the slightly shorter dorsal and larger eye:

"Dorsal 6 or 7; anal about 7; head about 3; width of body 3½; width of mouth 2 in head; interorbital width 4½, equal to snout. Eye large, 2⅔ in head. Teeth in the upper jaw conic, in more than 1 series in front, some in the outer row enlarged; teeth on the lower jaw in a single series, about 4 blunt incisors on each side followed by the canines, behind which the teeth are much smaller and conic. Width of disk 1¾ in its length, 2½ in head; opercular spine strong; distance of origin of dorsal from caudal 2½ in its distance from tip of snout. Sides and back uniform bright red; eye black, iris bright red; lower surface yellow, dotted with bright red. One specimen 23 mm. long in the Museum of the University of Indiana from St. Thomas (Coll. Edward W. Brigham), much shrunken and fins hardened by strong alcohol." (Eigenmann, in lit.)

Gobiesox macrophthalmus, GÜNTHER, Cat., III, 502, 1861, locality unknown.

2673. GOBIESOX CERASINUS, Cope.

Head 3 in total with caudal. D. 6; A. 6; C. 12; P. 24; V. 4. Head very wide, ovate, as broad as long to upper base of pectoral. Eye large, 3½ in head, equal to frontal width. Ten teeth on each side of each jaw, none of the upper being incisors, the 2 median on each side larger than the others; 3 teeth on each side in lower jaw horizontal, the others vertical, 2 of the horizontal teeth incisors, the median one on each side of these much the larger; each horizontal tooth with a small one behind it; no canines. Profile regularly descending from supraoccipital; a long subopercular spine. Dorsal beginning with last fourth of distance between tip of snout and base of caudal. Body and fins light crimson lake above, whitish below; no spots. One specimen, 2½ inches long, from St. Martins. (Cope.) (*κεράσινος, cerasinus*, cherry color.)

? *Cyclopterus nudus*, LINNÆUS, Syst. Nat., Ed. X, 260, 1758, "India;" from a specimen in Mus. Adolph Fred. (tab. 27), said to be 2 inches in length; the head broad with a sharp spine behind; dorsal rays 6; not *Lepadogaster nudus*, BLOCH & SCHNEIDER, Syst. Ichth., 2, 1801, who give "D. 12, A. 6,"* the description otherwise that of Linnæus: not *Gobiesox nudus* of recent authors, which is a species (*G. gyrinus*) allied to *G. virgatulus*.

Gobiesox cerasinus, COPE, Trans. Am. Phil. Soc., XIV, 1871, 473, St. Martins, West Indies. (Coll. Dr. R. E. Van Rijgersma. Type in Ac. Sci. Phila.)

Subgenus SICYASES,† Müller & Troschel.

2674. GOBIESOX ERYTHROPS, Jordan & Gilbert.

Head 2⅓; depth 6. D. 6; A. 5. Head scarcely longer than broad, proportionately very broad and depressed, its breadth 3 times in total.

*We do not know by what authority the number of fin rays given by Linnæus (D. 6) was altered to "D. 12, A. 6" by Schneider (Syst. Ichth.). The last-named figures agree with *nudus*, as described by Dr. Günther, that is, with our *G. gyrinus*. If the Linnæan type of *nudus* really had D. 6, it must have been *cerasinus* or *macrophthalmus* or some very similar species. The scanty Linnæan description agrees best with *cerasinus*. The name *nudus*, if used at all, must be taken for a species to which the Linnæan description may be applied. In our judgment the uncertainty is too great to justify the substitution of *nudus* for either *cerasinus* or *macrophthalmus*. It could be no other known species, however.

† This subgenus is composed of small species with the upper teeth in 1 series. This character should be verified on all our species, as perhaps none of them belongs to it.

Incisors in both jaws, entire and rather broad, the lateral teeth, as usual, pointed; no canines. Eyes very large, considerably wider than the narrow interorbital area, 3⅛ in head; interorbital area nearly 5 in head. Ventral disk a little longer than head, 2⅔ in body. Pectoral about ⅓ length of head. Distance from front of dorsal to caudal, 3⅔ in body. Caudal truncate, with rounded edges. Light olivaceous; body with 3 or 4 bars of cherry red; head marbled with red; eyes intensely cherry red, their upper border blackish; fins pale, the upper mottled with reddish; caudal barred with red. Two specimens, 1½ inches long, taken in a rock pool at Mazatlan; also recorded from the Tres Marias Islands. (*ἐρυθρός*, red; *ὤψ*, eye.)

Gobiesox erythrops, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 320, Mazatlan, Mexico (Type, No. 29248. Coll. Gilbert); JORDAN & GILBERT, Bull. U. S. Fish Comm. 1882, 108; JORDAN, Fishes of Sinaloa, in Proc. Cal. Ac. Sci. 1895, 499.

2675. GOBIESOX RUBIGINOSUS (Poey).

D. 6 or 7; A. 6; V. 4; P. 25. Head 3, including caudal, its greatest width twice its height; eye 2 in interorbital width. Body slender; head semioval, obtuse; distance from front of dorsal to caudal 1⅓ in length, including caudal; dorsal opposite anal, beginning at fourth seventh of total length; mouth terminal; teeth not examined by Poey. Color red, with 12 dark bands and with many scattered spots of clear blue on the body; eyes with a red circle. Length 22 mm. Cuba (Poey); not seen by us; locally common at Matanzas; perhaps a species of *Arbaciosa*. (*rubiginosus*, reddish.)

Sicyases rubiginosus, POEY, Synopsis, 391, 1868, wharves of Palmasola, Matanzas, Cuba (Coll. Poey); POEY, Enumeratio, 124, 1875.

2676. GOBIESOX CARNEUS (Poey).

Head rounded; body very slender; eyes large, as wide as interorbital space; mouth inferior; teeth not examined by Poey. Color pale red, with some white specks and bands. Length 22 mm. Otherwise essentially as in *Gobiesox rubiginosus*. (Poey.) Matanzas; a doubtful species. (*carneus*, flesh-colored.)

Sicyases carneus, POEY, Synopsis, 392, 1868, wharves of Palmasola, Matanzas, Cuba (Coll. Poey); POEY, Enumeratio, 124, 1875.

2677. GOBIESOX HÆRES, Jordan & Bollman.

Head 2⅔ (3¼ in total); depth 6 (7½). D. 9; A. 6. Body rather slender; head low and broad, greatest breadth not quite equal to length, its anterior margin not so broadly rounded as in *G. virgatulus*. Eyes very small, 1⅔ in interorbital space, 5 in head; interorbital bone appearing convex, least width 3½ in head and about equal to length of snout; cleft of mouth extending to beyond middle of eye; lower jaw included; teeth uniserial, those of upper jaw all canines, the first 3 on each side small, but becoming larger outward, next 3 or 4 much larger, rest smaller than those in front; anterior teeth of lower jaw entire incisors, which have on each side

about 6 large graduated canines and behind these a few smaller ones; teeth of lower jaw slightly oblique. Distance from front of dorsal to root of caudal about $2\frac{5}{6}$ in body ($3\frac{1}{2}$ in total). Pectorals moderate, 2 in head; ventral disk $1\frac{1}{4}$ in head. Color olivaceous, without any distinct bands; the occipital region and the caudal peduncle darker; body irregularly mottled with groups of darker spots; nape, preopercle, cheeks, and snout with numerous dark points; indistinct dark lines radiating from eye; lips dark; fins dusky; dorsal and anal with the first rays black; a pale spot near base of caudal; axil of pectoral dusky. Green Turtle Cay, Bahamas; a single specimen known, $2\frac{1}{4}$ inches in length. (*hæres*, one who clings.)

Gobiesox hæres, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1888, 552, Green Turtle Cay, Bahamas. (Coll. Dr. Charles L. Edwards.)

2678. GOBIESOX PUNCTULATUS (Poey).

Head very broad, 3 in total length with caudal. D. 9; A. 6. Color brown, covered with black points; 3 dark transverse bands; none on the head. Teeth not described, the incisors probably entire. Length 38 mm. Cuba (Poey); not seen by us. (*punctulatus*, speckled.)

Sicyases punctulatus, POEY, Enumeratio, 124, 1875, Havana.

2679. GOBIESOX FASCIATUS (Peters).

D. 7; A. 7; head and body with alternate dark green and yellowish cross bands. Commencement of dorsal before that of anal, its distance from caudal equal to length of its base. Type, 50 mm. long. (Peters); not seen by us; teeth not described. Puerto Cabello. (*fasciatus*, banded.)

Sicyases fasciatus, PETERS, Monatsber. Berl. Acad. 1859, 412, Puerto Cabello; GÜNTHER, Cat., III, 497, 1861; GÜNTHER, Fishes Centr. Amer., 390, 1869.

866. RIMICOLA, Jordan & Evermann.

Rimicola, JORDAN & EVERMANN, Proc. Cal. Ac. Sci. 1896, 231 (*muscarum*).

This genus differs from *Gobiesox* mainly in the very slender body and head. Head $3\frac{1}{4}$ to 4 in length, its width less than its length; dorsal and anal very short, of 4 to 6 rays each; incisors entire; a crescent-shaped patch of teeth in each jaw behind the large teeth; opercular spine weak or obsolete; sucking disk small. Species of small size; living below tide marks. (*rima*, a crevice; *colo*, I inhabit.)

a. Dorsal rays 6; anal 5; color yellowish, with a brown lateral band and numerous brownish spots. MUSCARUM, 2680.

aa. Dorsal rays 4; anal 5; color uniform light green. EIGENMANNI, 2681.

2680. RIMICOLA MUSCARUM (Meek & Pierson).

Head $3\frac{1}{4}$ in length; depth $8\frac{1}{4}$; D. 6; A. 5. Body elongate, slender, depressed anteriorly, but very narrow, slightly compressed posteriorly, the greatest width of body immediately behind head, 7 in length. Head narrow, much depressed, wider posteriorly. Eye small, its diameter $2\frac{1}{2}$ in interorbital width, 5 in head. Maxillary reaching to the front of the eye,

its length less than 3 in head. Teeth in upper jaw conical, acute, curved, forming a crescent-shaped patch, those of the anterior row enlarged; in the lower jaw an anterior row of about 5 broad, entire incisors placed nearly horizontally; behind these a crescent-shaped patch of teeth, similar to those in the upper jaw, becoming canine-like laterally. No evident opercular spine. Ventral disk longer than broad, its length $1\frac{1}{2}$ in head $6\frac{1}{4}$ in length; distance from vent to front of anal $2\frac{1}{2}$ in the distance from vent to disk; pectoral fin broad, short, $2\frac{1}{2}$ in head; dorsal and anal fins small, the anal slightly in advance; caudal fin rounded. Ground color, in alcohol, light yellowish, paler below; above everywhere sparsely covered with distinct brownish-red spots about as large as pupil; a lateral band of the same color begins on the front of the snout, where it joins the one on the opposite side, extends through the eye across the opercle to the caudal, becoming very indistinct posteriorly; this lateral stripe is in strong contrast with the uniform pale ventral surface. Coast of California. Two specimens were dredged in Monterey Bay at a depth of about 10 fathoms. One of these, the type, is $1\frac{1}{2}$ inches long. The second specimen ($1\frac{1}{10}$ inches long) has the dorsal spots confined to the top of the head and nuchal region and the lateral stripe disappearing slightly behind middle of body, and having the ventral surface marked posteriorly with brownish-red spots like the spots on the dorsal surface. (*muscarum*, of the flies, from the fly-speck markings.)

Gobiesox muscarum, MEEK & PIERSON, Proc. Cal. Ac. Sci. 1895, 571, with colored plate, Monterey Bay. (Coll. S. E. Meek and Charles J. Pierson. Type in L. S. Jr. Univ. Mus.)

2681. RIMICOLA EIGENMANNI (Gilbert).

D. 4; A. 5. Head $3\frac{3}{8}$ in length; depth about $\frac{1}{7}$ head. Body very slender and narrow, the width of head $4\frac{1}{5}$ in length; width of body 6. Mouth wide, the distance between its angles $\frac{1}{2}$ length of head, the maxillary scarcely reaching vertical from front of orbit. Interorbital space wide, about $\frac{1}{2}$ head. Eye very small, about 3 in interorbital width. Teeth in upper jaw conic, acute, in several series, the anterior in upper jaw enlarged; teeth in lower jaw also in several series, those of front row narrow incisors, entire, with rounded or truncate edges; disk very small and narrow, its width about $\frac{3}{8}$ its length, the latter $1\frac{3}{5}$ in length of head. Fins all small, the base of dorsal $\frac{2}{5}$ length of head, less than free portion of caudal peduncle; distance from origin of dorsal to base of caudal $3\frac{1}{8}$ in length before dorsal; distance from vent to front of anal fin $1\frac{3}{5}$ in its distance from disk; caudal broadly rounded, its length $1\frac{1}{4}$ in head; pectoral somewhat pointed, about $\frac{1}{2}$ head; coracoid plate small, about $\frac{1}{2}$ height of pectoral and less than $\frac{1}{2}$ its length. Color uniform light olive green, without distinctive markings. Type, a single specimen, about 1 inch long, taken at Point Loma, near San Diego, California. Other specimens were taken some years since at San Cristobal Bay by Mr. Charles H. Townsend, and were referred to as *Gobiesox rhessodon* by Mrs. Eigenmann, Proc. U. S. Nat. Mus. 1884, page 553. (Named for Dr. Carl H. Eigenmann.)

Gobiesox eigenmanni, GILBERT, Proc. U. S. Nat. Mus. 1890, 96, Point Loma, near San Diego, California. (Coll. Gilbert.)

867. ARBACIOSA, Jordan & Evermann.

Arbaciosa, JORDAN & EVERMANN, Proc. Cal. Ac. Sci. 1896, 290 (*humeralis.*)

This genus differs from *Gobiesox* chiefly in the character of the incisor teeth of the lower jaw; these are strongly serrate, or tricuspid, making a ragged cutting edge. Size small; dorsal and anal comparatively short; head not very broad, the jaws contracted; vertebræ (in *Arbaciosa zebra*) 28. Some species provisionally referred to the section *Sicyases* of *Gobiesox* may prove to belong to *Arbaciosa;* small species, living in rock pools, among the sea urchins, by whose spines they are protected. This relation of *Arbaciosa zebra* with the Echinoid *Arbacia stellata* is especially constant. (*Arbacia,* a sea urchin.)

a. Anal fin long, about 10 rays; dorsal rays 11; teeth above in 1 series; color olivaceous. RHESSODON, 2682.

aa. Anal fin of 5 to 7 rays.

b. Dorsal fin of 8 or 9 rays; teeth in single series; color brownish, with red bars and a large black humeral spot. HUMERALIS, 2683.

bb. Dorsal fin of 7 rays; color greenish, with pale spots and numerous pale cross bands; no red; body slender. RUPESTRIS, 2684.

bbb. Dorsal fin of 6 rays; color chiefly red.

c. General color pinkish olivaceous, with some bright red; back with 5 reddish-brown or blackish bars. Upper teeth in more than 1 series. Body comparatively slender, the depth nearly 8 in length. ZEBRA, 2685.

cc. General color bright rosy red, black, with 1 to 3 faint dark bars. Upper teeth nearly uniserial. Body comparatively stout, the depth 5⅓ in length. EOS, 2686.

2682. ARBACIOSA RHESSODON (Rosa Smith).

Head 3¼; depth 6¼. D. 11; A. 10; eye 4½ in head, ⅔ in interorbital space; ventral disk 1⅓ in head; pectoral 2; caudal 2⅛. Form much as in *Gobiesox mæandricus;* snout bluntly and evenly decurved; the greatest height of the body across the pectoral fins; head broader than body but less deep; maxillary extending to below the eye; incisors of lower jaw not much declined, each of them tricuspid, the central cusp longest; teeth of upper jaw conical, in an irregular series of 7 to 9; teeth in each jaw in single series; opercular spine sharp; distance from vent to caudal 2⅔ in length of body; dorsal a little longer than the anal, having its origin in advance of the anal and terminating opposite it; caudal rounded. Color dark olivaceous, usually with 3 broad yellowish cross bands above, the first across interorbital space and cheek, the second very wide, across back and front of dorsal fin, the third below middle of dorsal, some or all of these sometimes wanting; a dark bar at base of caudal; belly yellowish. Length 2½ inches. San Diego to the northern part of the Gulf of California; locally abundant in rock pools. (*ῥήσσω,* to make ragged; *ὀδούς,* tooth.)

Gobiesox rhessodon, ROSA SMITH, Proc. U. S. Nat. Mus. 1881, 140, San Diego, California; JORDAN & GILBERT, Synopsis, 749, 1883; ROSA SMITH, Proc. U. S. Nat. Mus. 1883, 235.

2683. ARBACIOSA HUMERALIS (Gilbert).

Head 3 to 3½; width of body 4¼; width of head 3⅝; eye very small, 3 in interorbital width. D. 8 or 9; A. 7. Body of moderate width, the head not evenly rounded anteriorly, becoming contracted opposite eyes, the snout forming a quadrate projection beyond the profile, as seen from above. Teeth in a single series in each jaw, the anterior narrow incisors, trilobate at tip, the 2 posterior teeth on each side strong, conical canines, somewhat recurved; about 12 incisors in the upper jaw. Interorbital space very wide, about equaling width of mouth, 2⅛ to 2½ in length of head. Ventral disk about as wide as long, its length 1¼ in head. Opercular spine large and strong, but not exposed. Distance from front of dorsal to base of caudal 3 in length anterior to dorsal; base of dorsal 1⅜ in head; base of anal about equals base of dorsal; distance from vent to front of anal half its distance from disk; caudal broadly rounded, 1¾ in head; pectoral ⅓ head, without distinct fold of skin across it. Ground-color dark olive brownish, crossed by many carmine-red bars, these somewhat broken anteriorly and above, to form reticulating lines, posteriorly and on lower part of sides more regular and running obliquely downward and backward; a conspicuous round humeral spot, larger than eye, in life black with golden-green reflections; numerous streaks from eye backward across cheek and opercles. (Gilbert.) Gulf of California; abundant at Puerto Refugio (Angel Island); also known from La Paz. (*humeralis*, pertaining to the shoulder, *humerus*.)

Gobiesox humeralis, Gilbert, Proc. U. S. Nat. Mus. 1890, 95, Puerto Refugio, Gulf of California. (Coll. *Albatross*.)
Arbaciosa humeralis, Jordan, Proc. Cal. Ac. Sci. 1896, 230, pl. 35.

2684. ARBACIOSA RUPESTRIS (Poey).

Head 4 in total length with caudal; depth 6; eye 4 in head; snout less than eye. D. 7; A. 7. Forehead little decurved; eyes well separated; mouth small, with 1 row of compressed, close-set incisors with denticulated edges, 6 on each side in each jaw; snout truncate, as seen from above. Pectorals short, rounded. Dorsal and anal alike, opposed, highest in front. Caudal rounded. Color greenish ash, each side with 6 large oval spots, those behind touching; sides with about a dozen vertical bands of straw yellow or whitish, these bands sometimes interrupted, forming 2 series of points; 2 small similar bands from the eye, another toward tip of snout; a brown pale-edged band between eyes; some white spots on sides of head. Length 1½ inches. Coral reefs of Cuba; not rare. (Poey): not seen by us; said to be distinguished from other Cuban species by the slender body and narrow head. (*rupestris*, living among rocks.)

Gobiesox rupestris, Poey, Memorias, II, 283, 1861, Cuba.
Sicyases rupestris, Poey, Synopsis, 391, 1868; Poey, Enumeratio, 124, 1875.

2685. ARBACIOSA ZEBRA (Jordan & Gilbert).

Head 3⅜ in length; depth nearly 8. D. 6 or 7; A. 5 or 6; vertebræ 11+17=28. Body comparatively very long and narrow, the greatest width about ⅕ the total length. Head narrow, depressed, its width about 4½

times in length of body; eye small, its diameter about ½ interorbital width; opercular spine well developed; ventral disk nearly as long as head; mouth rather small, anterior, maxillary reaching front of eye; incisors of lower jaw nearly horizontal, rather broad, 3-lobed at tip, the middle cusp the longest; upper teeth much smaller, the median ones compressed, blunt, close set, a little shorter than the lateral teeth and with dentate edges, 1 or 2 series of small teeth close behind them; anal beginning under middle of dorsal; the distance from insertion of dorsal to base of caudal contained 3⅜ in length; pectoral ½ as long as head; caudal truncate, with rounded angles. Back with 5 dark cross bars about as wide as the interspaces, 3 of them in front of dorsal fin, the 2 anterior much broader and more distinct than the others; these bars all distinct on back, fading on sides, which are often vaguely clouded with dark; the color of these dark bars varies from reddish brown to black, and that of the interspaces from olivaceous to light pink and bright rose red; top of head bright red, marbled with light slaty bluish; a black blotch on opercle, and 2 very distinct black cross spots, 1 on each side of median line, forming the front of first dorsal bar; cheek sometimes with 2 or 3 pale bluish streaks; dorsal, pectoral, and caudal more or less shaded with dusky; lower fins pale; usually a dark bar at base of caudal and 1 across middle of fin; shade of ground color extremely variable.* Very abundant in the rocky tide pools around Mazatlan, hiding everywhere under the numerous sea-urchins, especially *Arbacia stellata*, the protective coloration of both being that of the *Corallina*, which lines the rock pools. Length 2 to 3 inches. (*zebra*, from the banded coloration.)

Gobiesox zebra, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 359, Mazatlan, Mexico (Type, No. 29250. Coll. Gilbert); JORDAN & GILBERT, Bull. U. S. Fish Comm. 1882, 108; JORDAN, Proc. Cal. Ac. Sci. 1895, 499.

* The following note on the variations is furnished by Miss Susan B. Bristol:

"I find 4 specimens of this species which differ considerably from the typical form. These may represent a distinct species, but at present we are inclined to think that all these forms are modifications of one species, *Arbaciosa zebra*. The following is a description of a specimen 1½ inches long, taken at Mazatlan (No. 4166 in the L. S. Jr. Univ. Museum): Head 3⅓; depth 9. D. 5 or 6; A. 6. Body slender, much depressed, compressed posteriorly, the greatest width 4⅜ in length. Head depressed, its width 1⅓ in its length. Eye very small, about 1⅓ in interorbital width. Snout rather rounded, 3½ in head. Opercular spine present. Interorbital width 2⅜ in head. Ventral disk 1¼ in head. Mouth small, the lower jaw inferior; outer teeth in both jaws serrate. Anal beginning at end of the first ⅛ of dorsal. Distance from front of dorsal to base of caudal 1⅛ in head. Caudal terminate. Pectoral 2⅜ in head. Color bright red, with very irregular yellow mottlings on back and sides, light yellow below; back with 4 irregular dark-red cross bars, the posterior 3 of which are wider than the interspaces; 3 of the cross bars in front of the dorsal fin, and the fourth on either side of the dorsal; 2 conspicuous black spots about ½ as large as eye, 1 on either side of median line on back above the pectorals a short distance behind their origin; snout plain, dark red; pupil white; 2 yellow parallel stripes extending from eye backward and downward, the second ending at a point about ⅓ the distance from tip of snout to end of opercle; dorsal, caudal, and anal dusky; ventrals and pectorals paler; a large red blotch at base of pectorals extending for a considerable distance on the fin. Another specimen from Mazatlan, bright red in color, about ¾ of an inch long (also in bottle No. 4166, L. S. Jr. Univ. Museum), differs from the preceding form in the following respects: In the greater depth, which is 6¾ in length, in the smaller ventral disk, which is 1⅓ in head; in the more pointed snout; in the absence of the 2 black spots above pectorals; and in having the 4 dark red bands on the back more distinctly marked. Two specimens from Guaymas, Mexico, 1 and 1¼ inches long, No. 92 in the L. S. Jr. Univ. Museum, are chocolate brown in color, the shorter having on its back, including the bar at base of caudal, 7 dark brown cross bars and no dark spots above the pectoral, while in the longer there are no cross bars but a dark brown spot about ½ as large as the eye is present above the pectoral; also, in the longer one, the dorsal begins at the end of first third of anal. The eye in the larger specimens of *zebra* is larger than in these 4 specimens, but some of the smaller specimens seem to be intermediate in this regard between the typical form and these forms."

2686. ARBACIOSA EOS (Jordan & Gilbert).

Head 3; depth $5\frac{1}{2}$; eye moderate, $1\frac{1}{2}$ in interorbital width, which is about $3\frac{1}{2}$ in head. D. 6; A. 6. Body comparatively short, stout, and narrow; the head rather broad, but, like the body, much less depressed than in *G. erythrops;* width of head less than its length, or $3\frac{3}{8}$ in body. Incisors serrate or tricuspid. Pectorals about 4 in head; ventral disk shorter than head. Distance from base of caudal to front of dorsal $3\frac{4}{5}$ in total length; caudal truncate. Bright rosy red, sometimes dusky above with black points; back with 1 to 3 faint dark bars; 3 dark lines downward and backward from orbit, and usually 1 or 2 more on opercle; caudal usually with a reddish bar at base and a dusky one toward tip; fins otherwise nearly plain. Pacific coast of Mexico; abundant in rock pools about Mazatlan in company with *Arbaciosa zebra,* hiding under sea-urchins, especially with *Arbacia stellata.* Length $1\frac{1}{2}$ inches. (ἠώς, sunrise; from the red colors.)

Gobiesox eos, Jordan & Gilbert, Proc. U. S. Nat. Mus. 1881, 360, Mazatlan, Mexico (Coll. Gilbert. Type, No. 29247); Jordan & Gilbert, Bull. U. S. Fish Com. 1882, 108; Jordan, Proc. Cal. Ac. Sci. 1895, 499.

Group BLENNIODEA.

(Blennioid Fishes.)

Body more or less elongate, naked or with scales, large or small; ventral fins small, more or less advanced in position, often wanting, the number of soft rays always less than 5; hypercoracoid perforate, the shoulder girdle normally formed; skull not armed with spines; suborbital not developed as a bony stay articulating with the preopercle; pseudobranchiæ present; dorsal fin long, its anterior half, and sometimes the whole fin, composed of spines; anal long; tail homocercal, the caudal usually rounded, rarely forked; vertebræ numerous, especially in the arctic species. A large group, with ill-defined boundaries, the more primitive forms showing affinities with the *Trachinoidea, Cirrhitidæ,* and other more typical fishes, the extremes very aberrant and passing directly into the *Ophidoidea,* and other forms lacking spines in the fins. We begin the series with the least modified of the type, the *Clininæ,* from ancestors of which group the others have doubtless descended.

a. Caudal fin present, sometimes united to dorsal and anal; dorsal spines connected by membrane.

b. Gill openings not reduced to horizontal slits below the pectoral fins.

c. Teeth not developed as coarse molars.

d. Mouth not vertical. Blenniidæ, cc.

dd. Mouth nearly vertical; scales small or wanting; no lateral line; no ventral fins; dorsal composed entirely of slender spines; gill membranes attached to the isthmus; teeth strong. Cryptacanthodidæ, cci.

cc. Teeth developed as coarse molars on vomer, palatines, and sides of lower jaw; dorsal of flexible spines only; scales minute; gill membranes joined to the isthmus; no ventral fins; air bladder present; no lateral line. Anarhichadidæ, ccii.

bb. Gill openings reduced to separate, narrow, nearly horizontal slits below and in front of the pectoral fin; ventrals small; dorsal fin long and low, anteriorly of slender spines; vertical fins connected. CERDALIDÆ, CCIII.

aa. Caudal fin none, the tail tapering to a point; no ventral fins; no lateral line; scales rudimentary; anterior part of dorsal of low free-hooked spines, the posterior part of many slender soft rays; teeth in jaws only, close set in 1 row. PTILICHTHYIDÆ, CCIV.

Family CC. BLENNIIDÆ.

(THE BLENNIES.)

Body oblong or elongate, naked or covered with moderate or small scales which are ctenoid or cycloid; lateral line variously developed, often wanting, often duplicated; mouth large or small, the teeth various; gill membranes free from isthmus or more or less attached to it; pseudobranchiæ present; ventrals jugular or subthoracic, of 1 spine and 1 to 3 soft rays, often wanting; dorsal fin of spines anteriorly, with or without soft rays; anal fin long, similar to soft dorsal; caudal well developed. Vertebræ in moderate or large number, 30 to 80. Carnivorous fishes of moderate or small size, mostly living near the shore in the tropical and temperate or arctic seas; most of them are carnivorous, the *Clininæ*, so far as known, ovoviviparous, the rest mostly oviparous. Genera, about 80; species, about 400; chiefly of the rock pools and algæ; some species in the lakes of Italy. Dr. Gill divides the tropical Blennies into 3 families, *Clinidæ*, *Blenniidæ*, and *Chænopsidæ*. The first and second of these are fairly well defined. The third is now heterogeneous, and some of its members are intermediate between the other two. The arctic Blennies he again divides into *Xiphidiidæ*, *Cebedichthyidæ*, and *Stichæidæ*, but the first and last of these groups intergrade, the *Xiphidiinæ* are modified *Clininæ*, and there are other forms as well entitled to separate rank as *Cebedichthys*. It seems to us better to treat the group as a single family with many subfamilies. (*Blenniidæ*, Günther, Cat., III, 206–297.)

I. Tropical Blennies, with the vertebræ mostly in moderate number, usually fewer than 45; lateral line usually arched high above the pectoral, if present; dorsal fin with soft rays, at least 1 being present; anal spines little developed; ventrals well developed, usually I, 3.

a. Body scaly.

CLININÆ:

b. Lateral line present, arched anteriorly over the pectoral, becoming posteriorly median in position, or else obsolete; species ovoviviparous.

c. Scales ctenoid, very rough, 35 to 40 in lateral line; dorsal divided into 3 fins; no cirri above eye. ENNEANECTES, 868.

cc. Scales cycloid; dorsal fin not divided into 3 fins.

d. Dorsal with 6 to 20 soft rays.

e. Shoulder girdle with a small upturned hook on its inner edge.

f. Scales along lateral line anteriorly not enlarged; snout sharp; first 5 spines of dorsal more or less modified.

g. Caudal fin forked; air bladder present; scales minute; teeth in jaws in more than 1 series, on vomer and palatines; first 5 dorsal spines lengthened, and partly separated. HETEROSTICHUS, 869.

gg. Caudal fin truncate; air bladder wanting; scales minute; teeth in jaws in more than 1 series; teeth on vomer, none on palatines; first 5 dorsal spines lengthened and partly separated. GIBBONSIA, 870.

ee. Shoulder girdle without upturned hook on its inner edge above.

h. Maxillary greatly developed, reaching much beyond eye; teeth on vomer and palatines; scales minute; soft dorsal long. NEOCLINUS, 871.

hh. Maxillary normal, not greatly expanded.

i. Anterior part of lateral line normally formed; usually a comb of filaments at the nape.

j. Palatines without teeth; scales moderate or small, 38 to 110 in lateral line.

k. Teeth in jaws in 1 row only; teeth usually on vomer, none on palatines; usually a comb of filaments at the nape. MALACOCTENUS, 872.

kk. Teeth in jaws in more than 1 row, a band of villiform teeth behind the others; teeth on vomer, none on palatines.

l. Body oblong, the depth 3½ to 4½ in length; a filament above the eye. LABRISOMUS, 873.

ll. Body elongate, the depth about 6 in length; no filaments above the eye. MNIERPES, 874.

jj. Palatines with teeth, those in jaws in more than 1 series; scales large, 30 to 37 in lateral line; no nuchal filaments.

m. Head very broad, depressed; soft dorsal of about 20 rays. GOBIOCLINUS, 875.

mm. Head moderate, not depressed; soft dorsal of about 8 rays. STARKSIA, 876.

ii. Anterior part of lateral line running on a series of enlarged scales without visible pores; teeth in more than 1 series in jaws; teeth on vomer and front of palatines. CRYPTOTREMA, 877.

dd. Dorsal with 1 short soft ray only; scales large; teeth in jaws in more than 1 series; teeth on vomer, none on palatines.

n. Dorsal fin more or less deeply notched behind the third spine.

o. First 3 spines of dorsal very slender, close set, forming a separate ribbon-shaped fin, which is much higher than any of the spines in the second dorsal; anal spines rather high; body strongly compressed, the snout very sharp. EXERPES, 878.

oo. First 3 dorsal spines stiff, wide set, not remote from rest of fin behind dorsal notch; anal spines short; body more elongate, the snout less acute. AUCHENOPTERUS, 879.

nn. Dorsal fin continuous, not notched. PARACLINUS, 880.

EMMNIINÆ:

bb. Lateral line straight, close to the dorsal fin; scales small, cycloid; dorsal notched, its anterior half of slender spines; no cirri on head; ventrals thickish, inserted slightly before pectorals; teeth in bands, the outer enlarged. EMMNION, 881.

aa. Body scaleless; species oviparous, so far as known.

p. Teeth comb-shaped, in a single row in each jaw, behind which are sometimes long canines; vomer and palatines usually toothless; lateral line usually single, with a strong arch anteriorly; dorsal fin long, continuous, or divided into 2 fins, the anterior portion composed of spines, which are stiff or flexible; anal fin long, usually with 1 or 2 small spines; ventrals well-developed, jugular, of 2 or 3 rays.

q. Teeth all fixed, attached to the bone of the jaws and not movable.

RUNULINÆ:

r. Caudal fin lunate or forked; teeth compressed; spines and soft rays of dorsal indistinguishable.

s. Ventral fins very long, each of a spine and a soft ray.
ATOPOCLINUS, 882.

ss. Ventral fins not ½ length of head, each with about 2 soft rays; gill opening reduced to a small slit above pectoral.
RUNULA, 883.

BLENNIINÆ:

rr. Caudal fin rounded; teeth slender; gill membranes not reduced to a small slit.

t. Teeth all fixed, attached to the bone of the jaws.

v. Gill membranes free from the isthmus, or at least forming a distinct fold across it.

w. Jaws one or both with a posterior fang-like canine, much longer than the anterior teeth.
BLENNIUS, 884.

ww. Jaws without canines, the teeth all equal.
SCARTELLA, 885.

vv. Gill membranes broadly united to the isthmus, the gill openings restricted to the sides.

x. Jaws one or both with posterior fang-like canines.
HYPLEUROCHILUS, 886.

xx. Jaws without posterior canines; the teeth equal.

y. Three articulated ventral rays.

z. Mouth small, the maxillary extending scarcely beyond front of eye; the head decurved in profile.
HYPSOBLENNIUS, 887.

zz. Mouth large, the maxillary extending beyond vertical from middle of eye; the head rather pointed in profile. CHASMODES, 888.

yy. Four articulated ventral rays.
HOMESTHES, 889.

SALARIINÆ:

qq. Teeth of front of jaws all movable, implanted on the skin of the lips.

a'. Vomer toothless.

b'. Jaws without posterior canines; dorsal fin deeply notched.
SCARTICHTHYS, 890.

bb'. Jaws one or both with posterior fang-like canines.

x. Dorsal fin continuous. RUPISCARTES, 891.

xx. Dorsal fin divided. ENTOMACRODUS, 892.

aa'. Vomer with a few teeth; posterior canines small.
SALARIICHTHYS, 893.

pp. Teeth unequal, not comb-like; body oblong or elongate, more or less eel-shaped, naked, or rarely with rudimentary scales; supraocular flap sometimes present. Gill membranes united, free from the isthmus; dorsal fin very long, sometimes divided into 2 fins; formed of flexible spines, which often pass gradually into soft rays; anal fin long; ventral

fins thoracic or subjugular, usually, not much, if any, before the pectorals, composed of 2 soft rays each, the spine rudimentary; caudal well developed, the dorsal and anal usually more or less joined to it at base.

OPHIOBLENNIINÆ:

c'. Jaws each with 4 strong hooked canines in front; a hooked posterior canine below; a cirrus above eye and 1 above nostril; body scaleless; caudal fin forked; dorsal fin notched; body not eel-shaped; dorsal and anal free from caudal; ventrals small.

OPHIOBLENNIUS, 894.

cc'. Jaws with numerous teeth, not as above; caudal fin not forked.

EMBLEMARIINÆ:

d'. Body not eel-shaped; dorsal and anal not joined to caudal; no scales; no cirri; no lateral line; ventrals before pectorals; teeth on palatines; caudal fin rounded.

e'. Dorsal fin very high, not notched, the spines passing gradually into the soft rays; jaws long, sharp at tip.

EMBLEMARIA, 895.

dd'. Body elongate or eel-shaped; the dorsal and anal low, joined to base of caudal.

CHÆNOPSINÆ:

f'. Ventrals subjugular, more or less before pectorals; palatines with teeth; jaws long and sharp.

g'. Jaws with strong teeth, not as above described; dorsal fin with its anterior half of flexible spines, the posterior half of soft rays, the former gradually passing into the latter; jaws long, pike-like; ventrals inserted slightly before pectorals; anal with 2 spines; a villiform band of teeth in each jaw behind anterior teeth.

h'. Vomer toothless. Dorsal rays about XVIII, 38; anal II, 38. CHÆNOPSIS, 896.

hh'. Vomer with a few teeth. Dorsal rays XVIII, 32; anal II, 30. LUCIOBLENNIUS, 897.

PHOLIDICHTHYINÆ:

ff'. Ventrals subthoracic, inserted below pectorals; teeth in jaws uniserial; anal fin without spines.

i'. Dorsal fin continuous, its spines indistinguishable from the soft rays. PHOLIDICHTHYS, 898.

ii'. Dorsal divided into 2 fins, the anterior portion of 3 flexible spines behind the nape.

PSEUDOBLENNIUS, 899.

II. Blennies arctic or subartic; the vertebræ in large number, usually 50 or more; lateral line various, usually median; dorsal fin usually without soft rays; scales small, cycloid, rarely wanting.

j'. Gill openings not continued forward below, the membranes broadly united, sometimes joined to the isthmus; ventral fins small or obsolete; scales small, cycloid.

k'. Pectoral fins short or wanting, never pointed, and never more than ½ head; pyloric cæca usually, but not always, obsolete.

l'. Body not covered with crosswise tubes at right angles to the lateral line.

m'. Dorsal fin composed of spines only.

STATHMONOTINÆ:

n'. Body scaleless; ventrals moderately developed; anal spines 2; no lateral line; no pseudobranchiæ. STATHMONOTUS, 900.

nn'. Body covered with small smooth scales.

CHIROLOPHINÆ:

o'. Ventral fins well developed, of 1 spine and 3 rays; no anal spines; top of head with many cirri; a row of large pores above base of pectorals; gill membranes free from isthmus; no pyloric cæca.

p'. Lateral line obsolete, only the row of pores being present. BRYOSTEMMA, 901.

oo'. Ventral fins rudimentary or wanting, not more than 1 soft ray present; dorsal spines all short and rigid.

PHOLIDINÆ:

q'. Lateral line obsolete.

r'. Gill membranes broadly united, free from the isthmus; no pyloric cæca; carnivorous.

s'. Anal fin with a large sheathed spine; ventrals wanting.

t'. Anal spine very long, pen-shaped, its anterior surface channelled; pectoral fins moderate. APODICHTHYS, 902.

tt'. Anal spine moderate or small, not pen-shaped, its anterior edge convex, not channelled.

u'. Pectoral fins very small; anal spine moderate. XEREPES, 903.

uu'. Pectoral fins wholly wanting; anal spine small. ULVICOLA, 904.

ss'. Anal fin with 2 small spines or with none.

v'. Ventral fins reduced to a short spine, followed by a rudimentary ray.

w'. Caudal fin well developed. PHOLIS, 905.

ww'. Caudal fin very narrow, the dorsal and anal united around the tapering tail. GUNNELLOPS, 906.

vv'. Ventral fins entirely wanting; caudal as in *Pholis*. ASTERNOPTERYX, 907.

rr'. Gill membranes joined to the isthmus, sometimes forming a fold across it; no ventral fins; no anal spines; top of head with fleshy crests; pyloric cæca present; body naked anteriorly, with small scales posteriorly. ANOPLARCHUS, 908.

XIPHIDIINÆ:

qq'. Lateral lines several, each with many short cross branches; pyloric cæca present; gill membranes free from isthmus; ventrals none; anal spines 2 or 3, small; herbivorous.

w'. Pectorals small but well developed, much longer than eye.
XIPHISTES, 909.
ww'. Pectorals minute, not longer than eye. XIPHIDION, 910.

CEBEDICHTHYINÆ:
mm'. Dorsal fin with its posterior half composed of soft rays; gill membranes broadly united, free from isthmus; ventrals wanting; lateral line single, high; pyloric cæca present; herbivorous.
CEBEDICHTHYS, 911.

DICTYOSOMATINÆ:
ll'. Body covered with crosswise tubes at right angles with the lateral line and forming a network with it.
x'. Dorsal fin of spines only; teeth strong; ventral fins present, well developed; gill membranes broadly united, free from the isthmus. PLAGIOGRAMMUS, 912.
kk'. Pectoral fins long and rounded or pointed, nearly as long as head; dorsal fin high; gill membranes broadly united, free from the isthmus; no lateral line; species probably all herbivorous.

OPISTHOCENTRINÆ:
y'. Ventral fins wanting.
z'. Dorsal with its posterior spines rigid and sharp; head scaly.
OPISTHOCENTRUS, 913.
zz'. Dorsal with its spines all flexible; head naked. PHOLIDAPUS, 914.

PLECTOBRANCHINÆ:
yy'. Ventral fins well developed; dorsal spines all pungent; body greatly elongate. PLECTOBRANCHUS, 915.
jj'. Gill openings continued forward below, the membranes separate or nearly so, scarcely joined to the isthmus; pectorals and ventrals well developed; dorsal spines slender, pungent, the fin without soft rays; herbivorous species.

LUMPENINÆ:
a''. Lateral line obsolete or obscure; body greatly elongate.
b''. Pectorals with the upper and middle rays shortened, shorter than lower; teeth on vomer and palatines. LEPTOCLINUS, 916.
bb''. Pectorals with the middle rays longest.
c''. Lateral line not wholly obsolete, a series of distant pores along sides; teeth on vomer and palatines. POROCLINUS, 917.
cc''. Lateral line obsolete, only a few small pores being traceable; no teeth on vomer; palatine teeth small or wanting.
LUMPENUS, 918.

STICHÆINÆ:
aa''. Lateral line present, single, double, or triple; body moderately elongate; teeth on jaws, vomer, and palatines.
d''. Lateral line simple, one on each side of back. STICHÆUS, 919.
dd''. Lateral lines 2, or dividing into 2 on each side. ULVARIUS, 920.
ddd''. Lateral line forking, forming 3 on each side. EUMESOGRAMMUS, 921.

868. ENNEANECTES, Jordan & Evermann.

Enneanectes, JORDAN & EVERMANN, Proc. Cal. Ac. Sci. 1895, 501 (*carminalis*).

Body rather robust, covered with large, rough ctenoid scales; lateral line almost obsolete; mouth moderate, the jaws equal; no tentacle above the eye or on nape; no hook on shoulder girdle; eye large; dorsal fin divided into 3 fins, the first of 3 or 4 slender spines, the second of about 10, the soft dorsal of about 7 rays; caudal rounded; anal fin long; pectoral long, the lower rays simple and thickened. Small fish of the rock

pools, closely allied to the Old World genus, *Tripterygion*, Risso, but distinguished by the chubby body, short fins, and large, rough scales. (ἐννέα, nine; νηκτηρ, swimmer, there being 9 fins.)

2687. ENNEANECTES CARMINALIS (Jordan & Gilbert).

Head 3⅜; depth 4½ to 5⅓. D. III–XII, 9 (IV–X, 8 in the specimen before us); A. II, 11 (misprinted II, 17) scales 33 to 40. Body rather stout, heavy forward, rapidly tapering behind. Head short, the snout low and rather pointed, the profile straight and steep from the snout to opposite the front of the eyes, there forming an angle and extending backward nearly in a straight line; eyes very large, longer than snout, 3 in head, high up, and close together; mouth wide, the jaws subequal, the maxillary extending backward to front of pupil; teeth moderate, essentially as in species of *Labrisomus*, those of the outer series enlarged; no evident cirri on the head; scales on body of moderate size, ctenoid, the edges strongly pectinate; belly naked; lateral line extending to opposite last ray of soft dorsal, ascending anteriorly, but without convex curve; dorsals 3, the first and second contiguous, the second and third well separated; first dorsal of 3 spines, the first of which is the highest and about as long as diameter of eye; the second dorsal of higher and slenderer spines, the anterior the highest, the longest about equaling greatest depth of body; soft dorsal shorter and a little lower than second spinous dorsal; caudal small; anal long, beginning nearly under middle of spinous dorsal; pectoral long, longer than head, reaching much past front of anal; ventrals ⅔ length of head. Color light brownish, with 4 dark-brown cross bars on sides, about as wide as the interspaces, which are marked with more or less reddish and with some lighter spots; belly pale; space behind pectoral dark; a dark bar downward and 1 forward from eye; first dorsal mottled with darker, second and third dorsals nearly plain; a narrow, dark bar at base of caudal and a broader one toward the tip, the fin sometimes entirely black; pectorals somewhat barred; lower fins plain. Mazatlan, in tide pools; the types, 4 specimens, each about 1½ inches long. Another from the same locality, since figured by Dr. Jordan, differs somewhat in the count of the fin rays; but the very small size of the specimen prevents us from being entirely sure of its correctness. (*carmen*, a hetchel, from the rough scales.)

Tripterygium carminale, Jordan & Gilbert, Proc. U. S. Nat. Mus. 1881, 362, Mazatlan. (Type, No. 28118. Coll. Gilbert.)

Enneanectes carminalis, Jordan, Proc. Cal. Ac. Sci. 1895, 510, with plate of young example.

869. HETEROSTICHUS, Girard.

Heterostichus, Girard, Proc. Ac. Nat. Sci. Phila. 1854, 143 (*rostratus*).

Body rather elongate, compressed, covered with very small, smooth scales, those along lateral line not enlarged; head long and low, the snout conic, produced, very acute; premaxillaries protractile; mouth moderate, terminal; each jaw with a row of conical teeth, behind which anteriorly is a broad patch of villiform teeth; vomer and palatines with villiform, teeth; gill rakers feeble; gill membranes broadly united, free from the

isthmus; orbital cirri minute or wanting; cheeks scaly. Dorsal fin very long, the posterior rays soft, the 5 anterior spines wider apart than the rest and separated from them by a notch, the first and second spines longest, rather flexible, the other spines stiff; caudal fin forked; ventrals I, 3; pectorals moderate; lateral line simple, complete, abruptly curved behind pectorals; air bladder present, large. Size large. Close to *Gibbonsia*, from which the presence of the air bladder and the form of the caudal separate it. (ἕτερος, different; στίχος, rank; in allusion to the differentiation of the anterior dorsal spines.)

2688. HETEROSTICHUS ROSTRATUS, Girard.

(Kelpfish.)

Head $3\frac{4}{5}$ in body; depth $4\frac{1}{3}$. D. V–XXXIII, 13; A. II, 34; eye 7 in head; maxillary $2\frac{1}{2}$; pectoral $1\frac{4}{5}$; ventral $2\frac{1}{3}$; first dorsal spine $4\frac{1}{3}$; highest ray of soft dorsal $2\frac{1}{2}$; third anal ray $2\frac{5}{6}$; caudal $2\frac{1}{6}$. Body much compressed, deepest anteriorly; head slender, compressed and pointed; lower jaw strongly projecting, with thick lip; maxillary reaching pupil; width of interorbital a little greater than eye; orbital cirrus minute, usually entirely wanting; cheek and upper edge of opercle with small scales, rest of head naked. Origin of dorsal a little in front of the vertical from gill opening; pectoral under third dorsal spine, reaching to below the eleventh or twelfth; ventrals inserted in front of pectorals in distance equal to length of snout, their tips reaching about $\frac{1}{4}$ of their length beyond base of the pectoral; soft dorsal higher than spinous, ending slightly anterior to the anal; caudal furcate, the middle rays $\frac{2}{3}$ length of outer. Color translucent, reddish brown, varying to blackish or olive, a series of large irregular light spots along sides below lateral line, continuous with a distinct light bar from eye to edge of opercle, bordered with black above, a similar spot on base of pectoral; an irregular line of large spots following outline of body under dorsal and above anal; a clear cut white streak from dorsal to tip of snout and continued on lower lip, the hue and pattern of color varying greatly; young examples most variegated; a translucent spot behind third dorsal spine, generally followed by similar spots for the whole length of the fin. San Francisco to San Diego. The largest of the Clinoid blennies, very abundant in the kelp, with which it agrees in coloration. Here described from a specimen, 16 inches in length, from San Francisco market. (*rostratus*, long-nosed.)

Heterostichus rostratus, Girard, Proc. Ac. Nat. Sci. Phila. 1854, 143, San Diego, California (Type, No. 284. Coll. A. Cassidy); Girard, Pac. R. R. Surv., x, Fishes, 26, pl. 13, 1858; Günther, Cat., 261, 1861; Jordan & Gilbert, Synopsis, 764, 1883.

870. GIBBONSIA, Cooper.

Gibbonsia, Cooper, Proc. Cal. Ac. Nat. Sci., III, 1864, 109 (*elegans*).
Blakea, Steindachner, Ichth. Beiträge, v, 148, 1876 (*elegans*).

Body less elongate and compressed, covered with minute cycloid scales, those along lateral line not enlarged; lateral line complete, abruptly decurved behind the pectoral; head somewhat pointed; snout unequal; conical teeth on jaws and vomer, the teeth mostly in single series, except

in front, where there is a narrow villiform band; no conspicuous posterior canines; maxillary not produced backward from angle of mouth; a tentacle above eye, none at nape; gill membranes united, free from the isthmus. Shoulder girdle with an upturned hook on its inner edge as in *Clinus*. Dorsal fin long and low, chiefly composed of spines, 5 of the anterior spines different from the others, longer and set farther apart; anal fin low, with 2 spines; ventral fins jugular, of 1 spine and 2 or 3 rays; caudal fin truncate; branchiostegals 6; no air bladder; pyloric cæca absent. Viviparous. Pacific coast; bright-colored fishes, inhabiting rock pools among algæ. This genus is very close to *Clinus* (type *C. acuminatus*, Cuvier & Valenciennes), differing chiefly in the form of the dorsal fin and in the pointed snout. In *Clinus* the first 3 dorsal spines are shorter than the others. (Named for Dr. William Peters Gibbons, of Alameda, California, who was one of the early naturalists in the California Academy of Sciences.)

a. Dorsal rays about V–XXXI, 10; anal rays about II, 26; soft dorsal low; coloration comparatively plain, the soft dorsal without pellucid area. EVIDES, 2689.

aa. Dorsal rays about V–XXVIII, 7; anal rays II, 24; soft dorsal high; coloration more or less highly variegated; soft dorsal with a large pellucid blotch posteriorly. ELEGANS, 2690.

2689. GIBBONSIA EVIDES (Jordan & Gilbert).

(KELPFISH; SEÑORITA.)

Head $4\frac{2}{3}$; depth $4\frac{1}{2}$. D. V–XXX or XXXI, 10 or 11; A. II, 26 or 27. Body elongate, compressed; head small, rather pointed; mouth quite small, terminal, the maxillary about reaching pupil, $3\frac{1}{4}$ in head; lower jaw projecting, vomer with teeth; no teeth on palatines; posterior teeth not recurved; eye moderate, shorter than snout, 5 to 6 in head; a small supraocular flap, not higher than pupil; nasal cirrus very small; first spine of dorsal inserted over preopercle, its length more than $\frac{1}{3}$ that of head, the second nearly equal; the third, fourth, and fifth progressively shorter; the sixth about as long as the fourth; the seventh longer; the rest nearly equal to the last, which is lower than the soft rays; the soft dorsal lower and more rounded than in *G. elegans*, the longest ray $2\frac{3}{4}$ in head; pectorals moderate, not reaching vent; ventrals moderate; scales very small, smooth; head naked; no air bladder. Usual color of adult, translucent, reddish or orange, nearly plain or with oblong dark clouds below middle of sides anteriorly; often scattered blackish spots on sides, irregularly placed, forming a broken lateral band, most distinct in the young; a large pellucid spot on the membrane behind third dorsal spine, sometimes some small ones behind it; pectorals nearly plain; dorsal and anal plain, reddish, with a broad dusky shade distally; soft dorsal without pellucid area; caudal plain; a dark streak backward from eye; young examples often variegated, with light and dark shades of red, brown, and white, sometimes with 6 to 8 dark cross bars, sometimes with 4 or 5 lengthwise stripes alternating with paler ones, the hue varying exceedingly and dependent on the surroundings, but never so extravagantly spotted as in *Gibbonsia elegans*. Length 9 inches. Coast of California south to Point Concepcion; abundant in the kelp, rarely in rock pools. Here described from specimens from Monterey. (*εὐειδής*, comely; *εὖ*, well; *εἶδος*, appearance.)

Blakea elegans, STEINDACHNER, Ichth. Beiträge, v, 148, 1876, specimens from San Francisco; not *Myxodes elegans*, COOPER.

Clinus evides,* JORDAN & GILBERT, Synopsis, 763, 1883; specimens from Monterey, exclusive of part of synonymy; name a substitute for *elegans*, preoccupied in *Clinus*.

2690. GIBBONSIA ELEGANS (Cooper).

(SPOTTED KELPFISH.)

Head 4$\frac{1}{3}$; depth 4$\frac{1}{3}$. D. V–XXVIII, 7; A. II, 24. Body rather strongly compressed; head short, rather pointed, mouth small, terminal, rather oblique, the maxillary barely reaching pupil, 3$\frac{1}{5}$ in head; lower jaw projecting; teeth as in *Gibbonsia evides*; eye rather large, 4$\frac{1}{2}$ in head, as long as snout; a small fringed supraocular flap, as long as pupil; a slender nasal cirrus; first dorsal spine 2$\frac{3}{4}$ in head; fins as in *G. evides*, the soft dorsal shorter, higher, and less rounded, its longest ray 2$\frac{1}{4}$ in head; pectorals and ventrals moderate, about as in *G. evides*; caudal fan-shaped on a slender peduncle; scales small and smooth; head naked. Color brown or red, agreeing with rocks or with *Corallina*, usually with eight irregular darker cross bars extending on the dorsal and anal, sometimes nearly plain brown; a dark spot probably always present behind head, and some, 1 or more, along lateral line posteriorly; spinous dorsal with a pellucid spot; usually many pale and dark spots and freckles on head and fins; pectoral and caudal usually barred, but plain in specimens taken in the kelp (*Macrocystis*), these latter much less variegated than tide-pool specimens; soft dorsal always with a large pellucid blotch posteriorly, this wanting or obscure in *G. evides*. Coast of southern California; abundant in rock pools lined with *Corallina* from Point Concepcion to Todos Santos; the specimens here described from Point Loma. Close to the preceding, but smaller and more brightly colored, the fin rays fewer. These differences, though small, seem to be constant; whether the 2 species overlap each other in geographical range is not known. (*elegans*, elegant.)

Myxodes elegans, COOPER, Proc. Cal. Ac. Sci., III, 1864, 109, San Diego and Santa Barbara.

Clinus ocellifer,† MOCQUARD, Bull. Soc. Philom. Paris 1886, 44, California.

Clinus evides, ROSA SMITH, Proc. U. S. Nat. Mus. 1883, 235, specimens from Todos Santos; not of JORDAN & GILBERT.

* The name *evides* may apparently be retained for this species, as the description of Jordan & Gilbert (Synopsis, 763) is based entirely on Monterey specimens, typical of this species. It was intended, however, as a substitute for the name *elegans*, already used in the genus *Clinus*, to which these species were then referred.

† The following is a translation of the description of *Clinus ocellifer* (Mocquard):

Head 4$\frac{1}{3}$; depth 4$\frac{1}{2}$. D. III–XXX, 8; A. II, 24; C. 13; P. 12; V. I, 3. Body strongly compressed, tapering rapidly behind. Eye a little longer than snout, 3$\frac{3}{4}$ in head, twice interorbital space; lower jaw a little longer than maxillary, reaching front of eye; a little tentacle on anterior nostril, elongate, with 4 or 5 unequal branches; a tentacle over eye. Three first dorsal spines nearly double length of those which follow, and separated by an interval equal to that which separates the first spine from the third; last dorsal spines longer than those that precede and stronger than any of the others; dorsal and caudal well separated. Scales very small. Opposite fifth and sixth dorsal spines immediately below lateral line is a lens-shaped spot of brownish black with a dull border; a second ocellus a little before the posterior extremity of the spinous dorsal; this surrounded by a pale brown circle in 1 specimen; the sides also with 5 irregular bands of a paler brown than that of the spots; the anterior is a little behind the corresponding spot; the posterior opposite the third or fourth soft ray of the dorsal; the posterior spot at the upper extremity of the fourth band; other spots of the same color at the base of the dorsal on the right of the caudal; sides with a longitudinal series of small white spots, not surrounded by black circle; other spots on the anal in 6 transverse lines; larger spots on subopercle and about the ventrals. Teeth on the vomer, none on the palatines. Coast of California. Two specimens, 93 mm. long. (Mocquard.)

871. NEOCLINUS, Girard.

Neoclinus, GIRARD, U. S. Pac. R. R. Surv., x, Fish., 114, 1858 (*blanchardi*).
Pterognathus,* GIRARD, Proc. Ac. Nat. Sci. Phila. 1859, 57 (*satiricus*).

Body compressed, rather elongate, covered with minute cycloid scales; lateral line present, incomplete, high anteriorly; head naked, the cheeks tumid; upper jaw protractile; maxillary greatly produced backward, more than ⅔ length of head, reaching far beyond the eye; both jaws, vomer, and palatines with stout, unequal, conical teeth in a single series, besides which, in the front of the jaws, are smaller teeth; nasal and supraocular region with fringed tentacles; gill membranes broadly united, free from the isthmus; gill rakers weak. Dorsal fin long, scarcely emarginate, its anterior ⅔ composed of slender, flexible spines, which are similar to the soft rays, all of which are simple; anal long, its rays all simple; ventrals moderate, I, 3; caudal fin distinct; pectorals rather broad, rounded; no air bladder; no pyloric cæca. Pacific coast, in shallow water; remarkable for the great development of the maxillary, as in *Opisthognathus* and *Gillichthys*. (*νέος*, new; *κλίνος*, *Clinus*.)

NEOCLINUS:

a. Maxillary long, but not reaching beyond head; membrane of jaws white, BLANCHARDI, 2691.

PTEROGNATHUS (*πτερόν*, wing; *γνάθος*, jaw):

aa. Maxillary inordinately developed, reaching gill opening in the adult; maxillary flap blackish, edged with bright yellow. SATIRICUS, 2692.

Subgenus NEOCLINUS.

2691. NEOCLINUS BLANCHARDI, Girard.

Head 4; depth 5¼. D. XXIV, 17; A. II, 30; eye 5 in head; maxillary variable, about 1½; pectoral 2; caudal 1¾ to 2. Upper profile of head convex, snout rather steep; jaws subequal; teeth on jaws, vomer, and palatines, subequal, canine-like; eye set high in head, equal to length of snout. Males with a long thick cirrus over front of middle of eye, twice as long as eye, its end multifid, 3 or 4 short, slender ones behind it over posterior half of eye; females with a much smaller cirrus in front, seldom as long as eye, the posterior ones similar to those of male; both with a multifid flap at anterior nostril; maxillary never reaching past preopercle (in specimens from 6 to 8 inches in length), not longer in males than in females. Head naked; scales on body very small, somewhat embedded; no scales on fins; origin of dorsal directly behind occiput, no notch between spinous and soft dorsals; the tips of last dorsal and anal rays reaching to base of caudal fin; pectorals broad, scarcely reaching to vent; about ½ the length of ventrals in front of base of pectoral. Color varying from dark red or

* "It is more than probable that had we been acquainted with this second species of *Neoclinus* first, we would have been misled as to its real generical characters, and framed a name in allusion to the condition of the upper jaw, such as *Pterognathus*, for example, which would have been most characteristic, for that upper jaw is as truly winged as the upper members of the flying squirrels. We can not help thinking that Cuvier himself would not have coined the name of *Opisthognathus* had he had before him the species which bears his name instead of that which he dedicated to Sonnerat. These two genera (*Opisthognathus* and *Neoclinus*) will furnish one of the best themes to ichthyological studies, as they exemplify the fact that specific characters may be developed to exaggeration, and become more conspicuous than the generic characters themselves." (Girard.)

plum color to olive green; sides mottled and spotted with darker; a dark spot, ocellated with yellow, generally present between first and second dorsal spines; dorsal blackish toward ends of rays; pectorals and anal white in female, slightly dusky in male; unexposed portion of lower lip entirely white; a yellow spot on base of caudal rays below and above. Coast of California, from Monterey to Santa Barbara; not rare; a remarkable fish. Here described from specimens from 6 to 8 inches in length, from Pacific Grove, California. We do not know what variation there may be in maxillary and barbels in larger or smaller specimens. This species differs from *N. satiricus* in having no second spot behind seventh spine of dorsal; in having that part of lower lip which is covered by the maxillary entirely white; barbels in male much longer; maxillary shorter; head slightly shorter; and in having the pectorals and anal lighter. (Named for its discoverer, Dr. S. B. Blanchard.)

Neoclinus blanchardi, GIRARD, U. S. Pac. R. R. Surv., X, Fish., 114, 1858, San Diego (Type, No. 691. Coll. Dr. S. B. Blanchard); GÜNTHER, Cat., III, 259, 1861; JORDAN & GILBERT, Synopsis, 761, 1883.

Subgenus PTEROGNATHUS, Girard.

2692. NEOCLINUS SATIRICUS, Girard.

Head 3½ in body; depth 6. D. XVI, 17; A. 30; eye 5 in head; pectoral 2; caudal 2½. Head bluntish, convex in profile; snout steep; jaws subequal; unequal, small canines on jaws, vomer, and palatines; eye about equal to length of snout, interorbital flattish, about ½ eye in width; 3 or 4 small barbels above eye, seldom as long as eye, the anterior one sometimes absent on one or both sides; cirri not differentiated in the female; a multifid flap on anterior nostril; maxillary always reaching past edge of preopercle (in examples 6 to 9 inches in length), just past in females, longer than head in males. Head naked, scales on body small, partly embedded; no scales on fins. Origin of dorsal directly behind occiput; no notch between spinous and soft dorsals; pectorals in the larger examples reaching to vent; last rays of dorsal and anal reaching base of caudal fin; anterior half of ventrals in front of base of pectorals. Color in spirits, reddish brown or olive green, mottled and spotted with darker; a dark spot ocellated with yellow between first and second dorsal spines, a similar one between seventh and ninth; dorsal blackish, pectoral, anal and ventrals varying from dusky to black, in no case light in our specimens; a yellow spot sometimes present, below and above, on base of caudal rays; the membrane connecting maxillary with lower jaw blackish, broadly and abruptly edged with white (probably yellow in life). Coast of California, from Monterey to Santa Barbara; a rare and most interesting species. Here described from specimens, 6 to 9 inches in length, from Pacific Grove, California. Differing from *N. blanchardi* in length of maxillary; slightly larger head; males without long cirri; a second spot on dorsal; fins darker, and especially in having the membrane of lower lip blackish, edged with white. (*satiricus*, satirical.)

Neoclinus satiricus, GIRARD, Proc. Ac. Nat. Sci. Phila. 1859, 57, Monterey, California, in 30 fathoms (Coll. A. S. Taylor); GÜNTHER, Cat., III., 260, 1861; JORDAN & GILBERT, Synopsis, 761, 1883.

872. MALACOCTENUS, Gill.

Malacoctenus, GILL, Proc. Ac. Nat. Sci. Phila. 1860, 103 (*delalandi*).

This genus is very close to *Labrisomus*, differing in the dentition, the teeth in the jaws being in single series; vomer with a few teeth or with none, none on palatines. The form of the dorsal fin in some species is different, there being usually a notch behind the fourth dorsal spine as well as at front of soft dorsal. Most of the species are not well known, and perhaps more than 1 genus is here included. (μαλακός, soft; κτείς, comb, in reference to the comb of filaments at the nape in the typical species.)

a. Nape without filaments.
b. Orbital tentacle present.
c. D. XXI, 8; spinous dorsal not notched, the first rays shortest; body elongate; snout pointed; scales large, about 38. OCELLATUS, 2693.
cc. D. XX, 12; spinous dorsal weakly notched; body rather robust. VARIUS, 2694.
bb. Orbital tentacle wanting; dorsal rays XXI, 11; spinous dorsal weakly notched; ventrals long. MACROPUS, 2695.
aa. Nape with a single tentacle. D. XVIII, 9; a tentacle above eye. LUGUBRIS, 2696.
aaa. Nape with a comb of slender tentacles; spinous dorsal more or less notched behind fourth or fifth spine.
d. Orbital tentacle present. D. XVIII to XX, 11 or 12 vomer with teeth.
e. Scales 43 or 44.
f. Highest soft ray of dorsal 1½ in head; dorsal without ocelli. GILLII, 2697.
ff. Soft rays of dorsal 1½ in head; dorsal fin with 2 large black ocelli; ventral fins long, as long as head. BIMACULATUS, 2698.
ee. Scales 55; ventrals moderate, shorter than head. DELALANDI, 2699.
dd. Orbital tentacle wanting; (no vomerine teeth?). VERSICOLOR, 2700.
aaaa. Nuchal and other filaments undescribed; a black ocellus on front of dorsal. D. XX, 11; scales 46. BIGUTTATUS, 2701.

2693. MALACOCTENUS OCELLATUS (Steindachner).

Head 4 to 4¼; depth 5 to 5⅔. D. XXI, 8; A. II, 8; scales 38; eye 4½ to 5 in head; snout 5⅔; interorbital width 10. Body elongate; the snout short; profile not steep; tentacle above eye very slender, none on nape. Maxillary ½ long as head, reaching posterior margin of eye. Teeth on jaws and vomer in 1 row, none on palatines. Dorsal with a notch between the spines and soft rays; spines all short, the longest not ½ head, the anterior shortest; the longest soft rays 1½ in head; dorsal slightly joined to base of caudal; ventral and caudal each 1⅓ in head; pectoral almost as long as head. Lateral line complete, strongly arched anteriorly. Color brownish; 8 pairs of narrow dark-brown cross bands on the body, most distinct above, sometimes broken up into cross spots; first membrane of the dorsal fin with black spot behind, sometimes a similar one, oval and indigo, behind eye; numerous sky-blue spots bordered with darker on sides of head and part of body; anal pale violet, edged with white, sometimes spotted; caudal gray, with darker spots in cross rows. Bahama Islands. Length 2 inches. (Steindachner.) Not seen by

us. Perhaps not a member of this genus; the large scales, entire spinous dorsal, and short soft dorsal, indicating affinities with *Starksia*, which has, however, a different dentition. It may prove to be the type of a distinct genus. (*ocellatus*, with eye-like spots.)

Clinus ocellatus, STEINDACHNER, Ichth. Beitr., v, 182, 1876, Bahama Islands.

2694. MALACOCTENUS VARIUS (Poey).

Head $3\frac{3}{5}$ in total length; depth $4\frac{1}{5}$. D. XX, 12; A. 18; C. 14; pectoral $1\frac{2}{5}$ in head; eye $3\frac{2}{3}$ in head, equal to snout. Mouth small; maxillary reaching opposite front of eye; profile prolonged; nostrils small, not tubular. Teeth firm, in 1 row, the points sharp and incurved; no teeth on vomer; a tentacle over eye, none at the nape; head naked; body scaly; lateral line short; dorsal beginning over middle of opercle, the spinous part forming a sinuous curve; the spines firm; the first higher than the 4 which follow; the last low; the next to the last lower than the last; soft dorsal higher than the spines; ventral rays apparently 2, the last one deeply divided. Color clear yellowish; the body spotted with black; an isolated spot at the end of the dorsal fin; vertical fins with all the rays dotted with black; pectoral pale, without specks. Length 52 mm. Cuba. (Poey.) Not seen by us. (*varius*, variegated.)

Myxodes varius, POEY, Enumeratio, 132, pl. 5, f. 2, 1875, Havana. (Coll. Rafael Arango.)

2695. MALACOCTENUS MACROPUS (Poey).

Head $4\frac{1}{4}$ in total length with caudal; depth $5\frac{1}{5}$. D. XXI, 11; A. I, 20; P. 17; eye $\frac{1}{2}$ longer than snout, $3\frac{1}{4}$ in head. Maxillary reaching front of eye. Teeth in 1 series, acute, not close-set; none on the vomer or palatines; no cilia over the eye nor on the nape; lateral line almost complete; ventrals as long as the depth of the trunk; first dorsal spine longest, the others forming a weak curve. Color uniform metallic coppery brown. Cuba (Poey); one specimen 35 mm. long. The type of this species examined by us in the Mus. Comp. Zool. It has scales 35; no hook on the shoulder girdle, and apparently no teeth on vomer or palatines. (*μακρός*, long; *πούς*, foot.)

Myxodes macropus, POEY, Synopsis, III, 99, 1868, Havana. (Coll. Poey.)

2696. MALACOCTENUS LUGUBRIS (Poey).

D. XVIII, 9; A. 20. Tentacle over eye; a filiform appendage on the side of the neck. Ventral very long, extending much beyond the vent. Dorsal fin with 2 depressions, the soft part short and very high, the first spine moderate. Color dark brown, with oblique vertical bands and brown points scattered over the head and trunk; a black spot at the base of the first 3 dorsal membranes; ventral entirely white. Cuba. One specimen 55 mm. long. (Poey.) Not seen by us. (*lugubris*, dismal, from the dark color.)

Myxodes lugubris, POEY, Enumeratio, 131, 1875, Cuba. (Coll. Poey.)

2697. MALACOCTENUS GILLII (Steindachner).

Head 4½; depth 4½; eye 4 in head; snout 3¼; interorbital 6 in head. D. XIX, 11; A. II, 17; P. 14; scales 43. Head pointed, conic anteriorly; snout longer than eye; a rather high tentacle above the eye, slender and split to the base, numerous others on the side of the nape; teeth not described. First three dorsal spines wider apart than others, first longest; eighth to tenth spines highest, ½ head; highest soft ray 1½ in head; dorsal deeply notched. Body greenish gray, with brown spots or faint cross bands; head and dorsal marbled with darker; ventrals white, the longest ray a little longer than head, reaching anal; anal edged with dark; pectoral as long as head. Barbados. Two specimens, the larger 2 inches long. (Steindachner.) This species may be a *Labrisomus*. (Named for Dr. Theodore Gill.)

Clinus gillii, STEINDACHNER, Ichth. Notizen, VI, 46, 1867, Barbados.

2698. MALACOCTENUS BIMACULATUS (Steindachner).

Head 4⅔; depth 4½. D. XX, 10; A. II, 19; ventral 3; scales 44. Near to *M. delalandi*, but the body deeper (said to be 5½ in the latter species, which is not the case). Profile to snout steep; eye a little shorter than snout, 3½ in head; jaws equal, each like the vomer with 1 row of teeth; maxillary reaching about to front of pupil. Interorbital space narrow, more than ½ width of eye; a very slender, rather long, bifid tentacle above eye; tentacles on the nape, upper 1 almost as long as tentacle above eye. Upper margin of dorsal weakly notched between first and fifth spines, more deeply between spines and soft rays, the former as in *M. delalandi*; longest soft rays 1⅕ in head; longest spines 2 in head; first 4 spines more widely separated than the others; pectoral and ventral as long as head; caudal a little shorter. Body brown, with dark-brown bands and numerous blackish spots, only the cross bands on the head strongly marked; tips of the anal rays whitish; above these a bluish violet streak; pectoral with 2 milk-white spots at base; a large black isolated spot at the base of the first 4 dorsal spines, a second on the last 4 spines, extending on the body; anal and caudal thickly spotted with brown. Small rocky islands to the north of Cuba. (Steindachner.) Not seen by us; evidently close to *Malacoctenus delalandi*, but the scales larger. (*bis*, two; *maculatus*, spotted.)

Clinus bimaculatus, STEINDACHNER, Ichth. Beitr., V, 180, 1876, small, rocky islands north of Cuba.

2699. MALACOCTENUS DELALANDI (Cuvier & Valenciennes).

Head 3½; depth 3½ to 4½. D. IV-XVI, 11; A. II, 18; scales 55. Form rather stout, compressed; snout not very short, rather pointed, the profile gibbous above the eyes, thence declining straight to the tip of the snout; mouth rather small, the maxillary reaching front of eye; teeth in a single series in each jaw; vomer with a few teeth, none on palatines; eye large, 3½ in head, as long as snout; small slender cirri above the eyes, and a fringe of moderately long filaments at the nape rather longer than the orbital cirri. Outline of spinous dorsal emarginate; first spine a little longer than eye, the second, third, and fourth progressively shortened, the

fifth again longer; the eighth to eleventh spines longest, thence gradually decreasing to the next to the last, which is much shorter than the last; soft dorsal rays considerably higher than the spines, the longest about $\frac{1}{2}$ length of head; anal long, not very high, the membrane deeply notched between all but the last 6 rays, which are the highest. Pectorals $\frac{5}{6}$ length of head; ventrals as long as from snout to edge of preopercle. Belly naked anteriorly; the scales small, cycloid; lateral line complete. Color olivaceous, darker above, much mottled and speckled with clear dark brown; sides with 5 distinct irregular dark-brown bars, extending from base of dorsal to level of lower margin of pectoral, their lower edges connected by a vague undulating longitudinal band; a blackish blotch on occipital region, and black blotches on cheeks, opercles, and before base of pectoral; opercle with several narrow pinkish streaks; head below with narrow streaks formed by series of dark-brown spots; an interrupted brown bar across lower jaw; belly unspotted; ventrals pale; other fins all barred with narrow series of dark-brown dots; anal somewhat dusky. Coast of Brazil and the west coast of Mexico; common. Here described from the types of *Clinus zonifer*. This is the most abundant denizen of the rock pools around Mazatlan, with the single exception of *Gobius soporator*, reaching a length of 3 to 5 inches. We are unable to separate *M. zonifer* from Mazatlan from Bahia examples of *M. delalandi*, and take our account from specimens of the former. (Named for Delalande, who collected for Cuvier in Brazil.)

Clinus delalandii, Cuvier & Valenciennes, Hist. Nat. Poiss., xi, 378, 1836, Brazil (Coll. Delalande); Günther, Cat., iii, 264, 1861.

Clinus zonifer, Jordan & Gilbert, Proc. U. S. Nat. Mus. 1881, 361, Mazatlan. (Coll. C. H. Gilbert.)

Clinus philipii, Lockington, Proc. Ac. Nat. Sci. Phila. 1881, 114; not of Steindachner.

Labrisomus delalandi, Jordan, Proc. U. S. Nat. Mus. 1888, 333.

2700. MALACOCTENUS VERSICOLOR (Poey).

Head $3\frac{1}{2}$; depth $3\frac{3}{4}$. D. XVIII, 12; A. 20. Body compressed; head moderate; snout prolonged; pectoral $1\frac{1}{5}$ in head; ventral $1\frac{2}{3}$; eye large, as long as snout, $3\frac{1}{2}$ in head; nostrils not tubular; mouth small; maxillary not reaching so far as eye; teeth firm, in 1 row, those above much larger and slightly curved backward; 12 teeth above and 9 below on each side [no teeth on vomer]; no tentacle over eye; no anal papilla; a comb of filiform tentacles on each side of the neck; head naked; body scaly; lateral line short; dorsal with 2 depressions, the first spine higher than the 4 which follow, the depressions much more marked than in *M. varius;* pectoral reaching beyond front of anal. Color yellowish brown; head, trunk, and fins varied with vertical brown bands and large brown spots; ventrals yellowish. On the figure the 5 bands behind the anal cross the body and extend on the vertical fins. Cuba. One specimen known, 53 mm. in length. (Poey.) Apparently very close to *M. delalandi*, but lacking the orbital tentacle, and, according to Poey, vomerine teeth also. (*versicolor*, variegated.)

Myxodes versicolor,* Poey, Enumeratio, 131, pl. 5, f. 1, 1875, Cuba. (Coll. Poey.)

* This species and its affines were referred by Poey to *Myxodes*, a South American genus allied to *Clinus* and *Gibbonsia*, but differing from the latter in its uniserial teeth.

2701. MALACOCTENUS BIGUTTATUS (Cope).

Dorsal XIX–I, 11; anal II, 16; The first dorsal spines the longest, last spine longer than penultimate; length of head without opercular flap, $3\frac{3}{8}$ times in length (exclusive of caudal fin); eye a little less than $\frac{1}{4}$ length of head, $\frac{2}{5}$ greater than interorbital width; pectoral fin reaching to fifth anal; scales large, 4–46–10 [cirri and teeth not described]. Pale reddish brown, humeral red-veined; rufous specks on anterior part of sides; 7 subquadrate brown blotches from nape to caudal fin, continued with interruptions as lateral bands, the fourth near end of spinous dorsal black; a black spot at base of membrane between first to third dorsal spines; 2 small brown spots behind orbit, the posterior on operculum. Length 2.25 inches. This species is well distinguished from *Labrisomus nuchipinnis* by the large scales, form of dorsal fin, coloration, etc. From New Providence, Bahamas; Dr. H. C. Wood's collection. Also a very small specimen from Dr. Rijgersma, St. Martins. (Cope.) Not seen by us. (*bis*, two; *guttatus*, spotted.)

Labrisomus biguttatus, Cope, Trans. Am. Philos. Soc. Phila. 1873, 473, New Providence, Bahama Islands. (Coll. Dr. H. C. Wood.)

873. LABRISOMUS,* Swainson.

Labrisomus, Swainson, Nat. Hist. Class'n Fishes, II, 277, 1839 (*pectinifer*).
Lepisoma, De Kay, New York Fauna: Fishes, 41, 1842 (*cirrhosum*).
Labrosomus, Gill, amended spelling.
? *Blennioclinus*, Gill, Proc. Ac. Nat. Sci. Phila. 1860, 103 (*brachycephalus*).
? *Auchenionchus* (misprinted *Anchenionchus*), Gill, Proc. Ac. Nat. Sci. Phila. 1860, 103 (*variolosus*).
? *Calliclinus*, Gill, Proc. Ac. Nat. Sci. Phila. 1860, 103 (*geniguttatus*).
? *Ophthalmolophus*, Gill, Proc. Ac. Nat. Sci. Phila. 1860, 104 (*latipinnis*).

* Concerning this genus and its affines, Dr. Gill remarks:
"The name *Labrosomus* (or *Labrisomus*) was first published in 1839, in the second volume of the 'Natural History of Fishes, Amphibians, and Reptiles.' At the seventy-fifth page of that volume, Swainson has divided the Cuvieran genus *Clinus* into 5 genera: *Clinus*, of which the *Clinus acuminatus*, Cuvier, is taken as the type; *Labrisomus* with *Clinus pectinifer*, Valenciennes, as type; *Tripterygion*, Risso, *Clinitrachus*, Reese, which is typified by *Blennius variabilis* of Rafinesque, and *Blennophis*, of which the *Clinus anguillaris*, Valenciennes is the only true species. Of these genera, *Clinus* Swainson, and *Clinitrachus* Swainson, are distinguished by false or illusive characters, and cannot be regarded as distinct. The others are valid, but their characters require revision. The only claim to distinction of the genus *Labrosomus* given by Swainson, is founded on the strong, conic, and pointed row of front teeth, behind which are villiform ones; a thicker body than in *Clinus*, and the 'dorsal fin distinctly emarginate toward the caudal.' The genus resting on these characters alone is composed of very incongruous elements. To it are referred, at page 277 of the second volume, the following species, all of which are described as species of *Clinus* by Valenciennes: *Labrosomus gobio*, *L. pectinifer*, *L. capillatus*, *L. delalandii*, *L. linearis*, *L. variolosus*, *L. peruvianus*, *L. microcirrhis*, *L. ? geniguttatus*, *L. elegans*, *L. ? littoreus* and *L. latipinnis*. Of these species, not more than 3 can, with propriety, be regarded as congeners, if the *Labrosomus pectinifer* is taken as the type. These are *Labrosomus pectinifer*, *L. capillatus*, and perhaps *L. delalandii*. The latter is more probably the representative of a distinct genus. That genus is distinguished from *Labrosomus* by the smaller mouth, the presence of only 2 rays to the ventral fins, and perhaps by the undulating margin of the spinous portion of the dorsal fin. It may be named *Malacoctenus*, in allusion to the pectiniform row of filaments. This genus is the nearest ally of *Labrosomus*. All the others are very distinct. *Labrisomus gobio* Swainson, is the type of quite a distinct genus, whose characters consist of a broad, depressed head, with a very short muzzle, large approximated eyes, superciliary and nasal tentacles, 2 ventral rays, and a comparatively short spinous dorsal. The genus may be called *Gobioclinus*. The only species, *Gobioclinus gobio*, is found in the West Indies, and has but 18 dorsal spines. *Labrisomus linearis* Swainson is synonymous with *Clinus brachycephalus*, Valenciennes. This, also, is the type of a

Body oblong, robust; head naked, short, compressed above; mouth rather large, with a row of stout, bluntish teeth in front of each jaw, behind which is a band of smaller teeth, broadest in lower jaw; teeth on vomer, no teeth on palatines; a tentacle above the eye; sides of neck with a tuft or series of fine filaments; dorsal fin continuous, with numerous slender spines and many soft rays, the spines not very unequal; pectorals long; lateral line continuous; scales moderate or small, cycloid; shoulder girdle without upturned hook-like process on its inner edge. Intestinal canal short, shorter than body. The limits of this genus are not well defined, and most of the nominal genera above named will probably be found worthy of recognition. This genus differs from *Clinus* chiefly in the absence of the upturned spine-like processes on the inner edge of the shoulder girdle. This process is found on *Clinus acuminatus*, the type of the genus *Clinus*. (*Labrus; σῶμα*, body.)

a. Scales moderate, about 70 in lateral line (so far as known); soft dorsal with 11 to 13 rays.

b. Dorsal spines 16; anal rays 20; tentacles on nape. HERMINIER, 2702.

bb. Dorsal spines 18; no teeth on palatines; first ray of dorsal not longest; orbital tentacle well developed; nape with a conspicuous comb of fringes.

d. Vomer with a cluster of small teeth. NUCHIPINNIS, 2703.

dd. Vomer with 3 to 5 large blunt teeth arranged in the form of a ∧. XANTI, 2704.

bbb. Dorsal spines 20; teeth on palatines (?); first dorsal spine longest. BUCCIFERUS, 2705.

aa. Scales very small, about 110; a comb of fringes at nape; first dorsal spines low; head with yellow spots. MICROLEPIDOTUS, 2706.

2702. LABRISOMUS HERMINIER (Le Sueur).

D. XVI, 11; A. 20; C. 14; P. 16; V. 3. Body slender, compressed. Cilia on nostrils, above the eye, and on the nape; lips thick, concealing conical teeth, behind each band of smaller teeth; teeth also on the palate and on the base of the gill arches. Scales rather large. Lateral line curved from the pectoral, becoming straight thence to the tail. Color reddish brown with numerous spots; a black spot at front of spinous dorsal.

distinct genus distinguished by its abbreviated and blenniform head, the profile being very convex; by the villiform teeth, the absence of superciliary tentacles, the spinous portion of the dorsal long, and the presence of only 2 rays to the ventral fins. The name *Blennioclinus* is conferred on it; for the species, the specific name of Valenciennes must be retained. *Labrisomus variolosus* is distinguished by a large thick head, with lateral eyes, short superciliary tentacles, and a small nuchal one. The mouth is large; the teeth of the jaws in an outer row strong and conical, behind which are villiform ones; those of the vomer and palate are villiform, in 3 patches, 1 on the vomer and 1 on each palatine bone. The spinous portion of the dorsal is long, and the ventrals have each 3 rays. The species thus characterized is the type of a new genus which may be named *Anchenionchus* (misprint for *Auchenionchus*). *Labrisomus microcirrhis*, *L. elegans* and *L. peruvianus* are nearly related to *Anchenionchus*, and are from the same zoological province. *Labrosomus? geniguttatus* is distinguished from *Anchenionchus* by the more approximated eyes and by the disposition of the vomero-palatine teeth, as well as the small size of the anterior row of maxillary teeth. The dorsal is moderately long, and each of the ventrals has 3 rays. The mouth is comparatively small, and there are superciliary, nasal, and nuchal tentacles. For this species the generic name *Calliclinus* is proposed. *Labrisomus latipinnis* is related to *Blennioclinus*, but is distinguished from the species of that genus by the presence of superciliary tentacles. The generic name of *Ophthalmolophus* may be retained for it." (Proc. Ac. Nat. Sci. Phila. 1860, 102, 103.)

St. Bartholomew, West Indies; known from one specimen taken among madreporic rocks. (Le Sueur.) Not recognized by any recent author; perhaps not distinct from *L. nuchipinnis*.

Blennius herminier, Le Sueur, Jour. Ac. Nat. Sci. Phila., IV, 1824, 361, St. Bartholomew.
Clinus hermineri, Cuvier & Valenciennes, Hist. Nat. Poiss., XI, 380, 1836.

2703. LABRISOMUS NUCHIPINNIS (Quoy & Gaimard).

Head 3½; depth 3½. D. XVIII, 12; A. II, 17; scales 70. Body oblong, rather robust; head naked, thick, short, not very obtuse anteriorly, compressed above; mouth rather large, the maxillaries not prolonged backward, extending to opposite the posterior part of eye, 2½ in head; teeth on vomer and palatines; front teeth of jaws conic, strong, behind them a band of villiform teeth, broadest in lower jaw; vomer with a patch of smallish teeth; eyes large; interorbital space very narrow; each side of neck with a long series of hair-like filaments, nearly as long as eye; orbital tentacle short and broad, multifid; nostril with a tufted barbel; lower jaw slightly projecting, its posterior teeth sometimes recurved; pectorals a little shorter than head, reaching vent. Dorsal spines rather slender, the 3 anterior spines scarcely shorter than the others, all the spines lower than the soft rays; dorsal fin commencing near the nape, the spinous portion long; soft rays higher than the spines; caudal small; pectorals rather large; ventrals moderate; gill-membranes broadly united, free from the isthmus; lateral line complete, high anteriorly, then abruptly decurved; membranes of vertical fins scaly; scales not very small, cycloid. Reddish brown, sometimes with vertical bands; a black spot on opercle, which is often edged with white; cheeks and fins reticulate or dotted. Length 6 to 8 inches. West Indies, north to Florida Keys, south to Brazil; generally common in rock pools; also recorded from the Canary Islands. (*nucha*, nape; *pinna*, fin.)

Clinus nuchipinnis, Quoy & Gaimard, Voy. Uranie et Physicienne, Zool., 255, 1824, Brazil (Coll. M. Freycinet & M. Gay); Günther, Cat., III, 262, 1861; Jordan & Gilbert, Synopsis, 762, 1883.
Clinus pectinifer, Cuvier & Valenciennes, Hist. Nat. Poiss., XI, 374, 1836, Bahia.
Lepisoma cirrhosum, De Kay, N. Y. Fauna: Fishes, 41, 1842, Florida.
Clinus canariensis, Valenciennes, in Webb & Berthelot, Poiss. Iles Canaries, 60, 17, f. 3, Canary Islands.
Clinus capillatus, Cuvier & Valenciennes, Hist. Nat. Poiss., XI, 377, 1836, Martinique.
Labrosomus pectinifer, Gill, Proc. Ac. Nat. Sci. Phila. 1860, 105.
Labrisomus capillatus, Gill, Proc. Ac. Nat. Sci. Phila. 1860, 107.

2704. LABRISOMUS XANTI, Gill.

Head 3½ in body; depth 3½. D. XVIII, 12; A. II, 18; scales 10–64 (pores)–12 (from front of straight portion of lateral line to anal); eye 4½ in head, maxillary 2; highest dorsal spine 2⅔; pectoral 1½; caudal 1⅔. Body not greatly elongate, compressed, anterior profile well rounded from snout to nape; mouth rather large, the maxillary reaching to below middle of eye; teeth small, canine-like, growing gradually larger toward

front of upper jaw; side teeth on lower jaw very small, abruptly enlarged on front half of jaw; teeth on vomer A-shaped, in a single row, the ones at the angles enlarged, 1 or 2 small ones between them at the sides; small multifid dermal flaps at nape, over eye, and above nostril; interorbital concave at the middle, $\frac{3}{5}$ the diameter of eye; gill rakers small and short, 3 + 6 in number. First dorsal spine inserted behind eye a distance equal to diameter of eye, about $\frac{1}{6}$ shorter than longest spine; soft dorsal the higher; origin of anal midway between snout and base of caudal, not running as far back as dorsal; pectoral reaching a little past front of anal; ventrals long and slender, inserted a little in front of pectorals, their ends not reaching vent; caudal rounded. Color in spirits, brownish gray, with about 6 wide irregular cross bars which are darker toward their edges, 2 black streaks running downward and backward from eye; cheeks and opercles with many small light blue spots; spinous dorsal mottled and spotted with darker, other fins with small irregular dark lines running across the rays; ventrals dusky; tentacles on head black. Described from a specimen 5 inches in length from La Paz, Lower California. Pacific coast of Tropical America from Gulf of California to Panama; common in rock pools; representing on the Pacific coast the scarcely different *L. nuchipinnis*. (Named for John Xantus.)

Labrosomus xanti, Gill, Proc. Ac. Nat. Sci. Phila. 1860, 107, Cerro Blanco (Type, Nos. 2334, 2335, 2478. Coll. J. Xantus); Jordan & Gilbert, Proc. U. S. Nat. Mus. 1882, 368.

2705. LABRISOMUS BUCCIFERUS, Poey.

Head $3\frac{3}{4}$ in total length with caudal; depth 5. D. XX, 11; A. II, 19; eye 4 in head, a little longer than snout; anterior nostril with a little tube; lower jaw longer; forehead convex, the snout short; mouth large, reaching beyond middle of eye; a few filaments on nape and 1 above preopercle. Teeth cardiform, the outer ones large; teeth on vomer (and palatines). Body scaly. Lateral line complete. First ray of dorsal longest, the others forming a convex curve up to the 19, which is shortest; pectoral moderate, of 12 rays, the lower thickened. Color brownish yellow, with vertical brown points extending on fins; a series of pale points along sides; the head gray, cheek dark brown. One specimen 55 mm. long. Cuba. (Poey.) Not seen by us. (*bucca*, cheek; *fero*, I bear).

Labrisomus bucciferus, Poey, Synopsis, 399, 1868, Cuba. (Coll. Poey.)

2706. LABRISOMUS MICROLEPIDOTUS, Poey.

Length of head equal to depth; pectoral $1\frac{3}{5}$ in head; eye $1\frac{2}{3}$ in snout, $4\frac{1}{2}$ in head. Maxillary reaching to base of middle of eye, $2\frac{1}{3}$ in head. Mouth oblique, with strong teeth; the lower jaw the longer; small fringe of tentacles at anterior nostril above eye, and a comb of fringes at the nape. First 3 dorsal spines subequal, considerably lower than the second 3, which become progressively longer. Scales in lateral line about 110. Head brown, with small yellow spots scattered over its lower part and on the gill membranes. Pectoral and caudal with some black points. Cuba

(Poey.) Known from an imperfect description, with a drawing of the head of a specimen 180 mm. long. (*μικρός*, small; *λεπιδωτός*, scaly.)

Labrosomus microlepidotus, POEY, Anal. Soc. Esp. Hist. Nat., XIX, 1880, 246, 1, 8, f. 2, Cuba. (Coll. Poey.)

874. MNIERPES, Jordan & Evermann.

Mnierpes, JORDAN & EVERMANN, Check-List Fishes, 468, 1896 (*macrocephulus*).

This genus is close to *Labrisomus*, from which it differs chiefly in the very elongate body and in the absence of an orbital tentacle. The dorsal spines are more numerous, and probably the vertebræ also. The lips are thick and there is no trace of hook on the shoulder girdle. A band of fillitorm teeth in the jaws behind the anterior series; teeth on vomer, none on palatines. (*μνίον*, moss; *ἕρπης*, creeper.)

2707. MNIERPES MACROCEPHALUS (Günther).

Head 4¼; depth 6 to 6¼ (7⅓ in total). D. XXII, 12; A. II, 24; C. 13; P. 13; V. I, 3; scales about 70. The head is depressed, rather short, nearly as broad as long; crown of the head broad and flat; interorbital space concave, narrower than the orbit. Snout very short, obtuse, rounded; the maxillary not extending to behind the posterior margin of the orbit; lips thick. Teeth in jaws forming a band with an outer series of stronger ones; vomerine teeth in a narrow band; palatine teeth none. No orbital tentacles, those at the nostril and on the neck very small. Gill openings wide, the gill membranes being united at the throat. Head naked; scales on the body not very small, cycloid. Dorsal fin commencing at occiput, and terminating near base of caudal, the spines flexible, and much lower than the soft rays; the 3 anterior ones rather more remote from one another than the following; none of the rays of this or of the other fins branched; caudal rounded; anal higher posteriorly than anteriorly, about as high as the spinous dorsal; pectorals rounded, with the middle rays longest, shorter than the head; ventrals jugular, ½ as long as the pectoral, with the spine and the outer ray enveloped in a common thick membrane. Dark grayish olive; head and fins blackish; head, base of the pectoral, anterior part of the body, and dorsal dotted with white. Pacific coast of Central America. (Günther.) Known from a few specimens from Panama. Those examined by us (Mus. Comp. Zool.) have the sides much freckled and mottled with pale. (*μακρός*, long; *κεφαλή*, head.)

Clinus macrocephalus, GÜNTHER, Cat., III, 267, 1861, Pacific coast of Central America (Coll. Capt. John M. Dow); GÜNTHER, Fish. Centr. Amer., 442, pl. 69, fig. 2, 1869.
Labrosomus macrocephalus, JORDAN, Proc. U. S. Nat. Mus. 1885, 389.

875. GOBIOCLINUS, Gill.

Gobioclinus, GILL, Proc. Ac. Nat. Sci. Phila. 1860, 102 (*gobio*).

Body robust; head broad, depressed, with a very short muzzle. Eyes large, approximated, close together; palatine teeth present; a tentacle above eye; no nuchal filaments. Scales very large, about 30 in the lateral

line. Spinous dorsal of 18 spines. This genus seems to differ from *Labrisomus* in the large scales, differently formed head, and in the absence of nuchal filaments. (*Gobio*, the gudgeon; *Clinus*.)

2708. GOBIOCLINUS GOBIO (Cuvier & Valenciennes).

Head 3½ in total length; depth 4½. D. XVIII, 19; A. II, 17; C. 15; P. 14; V. 2; scales 30–10. Head nearly as broad as long, its height a third less. Eye large, 2½ in head, twice interorbital space; a very small tentacle over the eye, another on the nostril. Profile rounded between the eyes, descending vertically to the snout, which is very short. Cheeks inflated; the skull a little rough. Mouth reaching to opposite middle of eye, somewhat black; teeth small, conic, and pointed; upper jaw with 26 equal teeth, the lower with 16, the last 2 larger and more curved; teeth on vomer and palatines, simple, in 2 irregular rows; gill membranes united, free from isthmus. Body posteriorly compressed. Dorsal slightly notched between spines and soft rays of anal; pectorals equal to ventrals, 5 in total length; caudal obtuse, 6 in total length. Lateral line disappearing opposite tip of ventral. Color greenish, with traces of cloudy brownish; the cross bands a deep brown, pointed at base of caudal. Lesser Antilles. Known from several specimens, one 2 inches in length. (Cuvier & Valenciennes.) Not seen by us; apparently a strongly marked species. (*Gobio*, the gudgeon, from its resemblance to *Cottus gobio*, the miller's thumb.)

Clinus gobio, Cuvier & Valenciennes, Hist. Nat. Poiss., xi, 395, 1836, Lesser Antilles. (Coll. Plée.)
Gobioclinus gobio, Gill, Proc. Ac. Nat. Sci. Phila. 1860, 102.

876. STARKSIA, Jordan & Evermann.

Starksia, Jordan & Evermann, Proc. Cal. Ac. Sci. 1896, 231 (*cremnobates*).

This genus is related to *Labrisomus*, differing in the large scales, the presence of palatine teeth, the short soft dorsal fin, and the absence of the comb of nuchal filaments. (Named for Mr. Edwin Chapin Starks, in recognition of his work on the fishes of Western America.)

2709. STARKSIA CREMNOBATES (Gilbert).

Head 3⅛ in length; depth 4⅓. D. XXI or XXII, 8; A. II, 19; scales 37. In appearance resembling very strongly the species of the genus *Auchenopterus*. Body slender, snout sharp, the jaws equal; mouth wide, oblique, the maxillary reaching vertical from posterior margin of orbit, 2¼ in head. Teeth small, villiform, forming a band in front of upper jaw, the outer series enlarged; in lower jaw a single series laterally, becoming double in front; similar teeth on vomer and palatines. Eye longer than snout, 4 in head; interorbital width less than diameter of pupil; opercle terminating in an evenly convex process behind, without spinous points; gill membranes broadly united, free from isthmus; no hook on inner edge of shoulder girdle; nostrils with a flap; a single slender filament above eye and 1 or more on each side of the nape. A slight notch between first

and third dorsal spines and another between the eighteenth and twenty-first spines; the spines are low and strong, the highest equaling the snout and ½ eye; soft rays higher, the longest equaling ½ head; caudal short, rounded, entirely free from dorsal and anal; anal similar to soft dorsal, the first 2 rays spinous; ventrals inserted well in advance of pectorals, each consisting of 1 spine and 2 soft rays, which are joined only at base; pectorals pointed, the lower rays the longest, 1¼ in head. Scales large, cycloid, the lateral line running high in front, descending to middle of sides immediately behind pectorals, thence running straight to tail. In the types, which are probably immature, the pores are not developed on posterior part of body. Color in spirits, uniform light olivaceous, a small dusky spot behind orbit and 1 below and behind it; opercle dusky. In 1 specimen the rays of soft dorsal, anal, and caudal are finely barred with dusky. (Gilbert.) Length 1½ inches. Gulf of California. Two specimens known, from *Albatross* Station 3001, in 71 fathoms. (*Cremnobates; κρημνοβάτης*, one that haunts rocks; a synonym of *Auchenopterus.*)

Labrosomus cremnobates, GILBERT, Proc. U. S. Nat. Mus. 1890, 100, Gulf of California. (Coll. *Albatross*).
Starksia cremnobates, JORDAN, Proc. Cal. Ac. Sci. 1896, 231.

877. CRYPTOTREMA, Gilbert.

Cryptotrema, GILBERT, Proc. U. S. Nat. Mus. 1890, 101 (*corallinum*).

This genus differs from *Labrisomus* chiefly in the absence of nuchal filaments and in the modified anterior portion of the lateral line, which runs on a series of enlarged scales having no externally visible pores. (*κρυπτός*, concealed; *τρῆμα*, pore.)

2710. CRYPTOTREMA CORALLINUM, Gilbert.

Head 3¾ to 4 in length; depth 5½. D. XXVII, 12; A. II, 27. Body elongate, regularly tapering backward to caudal peduncle, whose depth equals length of snout, which is sharp; mouth nearly horizontal; maxillary reaching middle of eye or beyond, 2⅓ to 2½ in head; teeth strong, but none of them enlarged, in a single series in jaws laterally, becoming double anteriorly; teeth on vomer and in a small distinct patch on front of palatines; eyes large, the interorbital space flat, nearly ½ diameter of orbit; orbit slightly exceeding length of snout, 3½ in head; branchiostegal membranes broadly united, free from isthmus, the posterior edge on vertical from preopercular margin; anterior nostril in a short tube, a slender flap arising from its posterior margin; a pair of simple slender filaments arising from the upper edge of each orbit, 1 on each side of nape, none others on head; gill rakers very short and weak; shoulder girdle without hook on its inner edge. Scales rather large, cycloid, the head alone naked; lateral line in its upper anterior portion without externally visible tubes, its position shown by a series of enlarged scales twice the size of the others; on these the tubes are wholly on the under side, each opening anteriorly by a single pore under the edge of the pre-

ceding scale; anteriorly the lateral line runs near the back and parallel with it, becoming suddenly declined behind middle of trunk, thence running on middle of side; the oblique portion of lateral line rests on about 7 scales, and the externally visible tubes of lateral line begin at this point; posterior portion of lateral line contained $1\frac{1}{2}$ to $1\frac{3}{4}$ in dorsal portion; scales of lateral line, 45 in dorsal portion, 7 in oblique portion, and 18 in posterior portion. A slight notch behind fourth dorsal spine, the second and third spines slightly longer than those following, the first little longer than the fourth, the longest spine about $2\frac{1}{4}$ in head; first 2 anal rays spinous, but weak and flexible; last dorsal and anal rays not joined by membrane to caudal peduncle, the depth of the latter equaling the length of its free portion; ventrals long and narrow, nearly reaching vent in males, consisting of 1 spine and 3 simple rays; pectorals with some of the lower rays longest, $1\frac{1}{3}$ in head; all of pectoral rays simple, 14 in number; caudal fin truncate, $1\frac{2}{3}$ to $1\frac{1}{2}$ in head. Length 5 inches. Color dusky olive above, with irregular narrow longitudinal streaks of bright coral red, and 7 round black blotches above middle of sides; reticulating red lines and spots on top and sides of head and snout; branchiostegal membranes dusky in males; 2 red streaks on base of pectorals; dorsal somewhat dusky, marked with lines of red spots; caudal with 3 rather faint cross bars; pectorals, ventrals, and anal largely black in males, pale in females; the red shades persistent in alcohol. Santa Barbara Islands. Three specimens from *Albatross* Station 2945, in 30 fathoms. (Gilbert.) (*Corallina*, a calcareous alga, among which it lives.)

Cryptotrema corallinum, Gilbert, Proc. U. S. Nat. Mus. 1890, 101, off Santa Barbara Islands. (Coll. C. H. Gilbert.)

878. EXERPES, Jordan & Evermann.

Exerpes, Jordan & Evermann, Proc. Cal. Ac. Sci. 1896, 232 (*asper*).

Body slender, much compressed; the snout long, sharp in profile; first dorsal ribbon-shaped, the 3 slender spines close together, inserted at the nape, much in advance of the rest of the fin; ventrals very long and slender. Otherwise as in *Auchenopterus*, the scales large, and but 1 soft ray in the dorsal fin. (ἐξω, without; ἕρπης, creeper.)

2711. EXERPES ASPER (Jenkins & Evermann).

Head 3 ($3\frac{2}{3}$ in total); depth $5\frac{1}{2}$ ($6\frac{1}{3}$); eye $4\frac{1}{3}$ in head; scales 6–43–7, about 40 pores. D. III–XXV, 1; A. II, 20. Body compressed; head narrow, pointed; snout long, lower jaw slightly the longer; mouth a little oblique, cleft moderate, maxillary not reaching nearly to vertical at front of orbit. Teeth in 1 well-defined outer series and a broken inner one, those in the outer series strongest and of pretty uniform size, short and broad; vomerine teeth in a single patch; no palatine teeth. No tentacles of any kind about the head. Profile nearly straight from snout to origin of first dorsal, but very slightly arched from there to base of caudal fin. Scales rather large, cycloid, about 6 rows between origin of second dorsal and lateral line just behind its angle, and about 7 from there to mid-

dle of ventral surface; 9 rows from origin of second dorsal to upper limb of opercle; entire head, opercles, and fins naked. Lateral line beginning at upper limb of opercle on a level with the pupil, almost exactly under the middle of the first dorsal fin, and a little more than ¼ the distance from top of nape to the under side of the throat, arching gently for 7 or 8 scales, leaving but 1 row of scales between it and the first spines of the second dorsal; on the ninth, tenth, and eleventh scales it bears slightly downward until 2 rows are left between it and the dorsal, then a sharp turn is made which puts it 4 scales further down, and from there it pursues a nearly direct line to middle of base of caudal fin. Dorsal fins separate, the first of 3 slender, very close-set, flexible spines, their length about twice in that of head, the fin ribbon-shaped; second dorsal separated from first by a distance somewhat greater than diameter of eye, and composed of 25 rather stout, sharp spines and 1 terminal soft ray; the first 3 are graduated, the first being contained 1½ times in distance between the 2 fins, the second is about ½ longer, and the third still a little longer; the remaining 22 are of approximately equal length, about equaling distance from origin of first dorsal to that of second; the 1 soft ray somewhat shorter than spines, well separated from caudal by a space equal to that between dorsals; pectorals inserted under middle of space separating dorsals, composed of 14 rays, equaling eye and snout in length, and reaching slightly past origin of anal; ventral of 2 rays inserted directly under origin of first dorsal and considerably in front of pectorals, which they somewhat exceed in length, in some specimens reaching vent; anal fin beginning slightly in front of posterior end of pectorals, a little lower than second dorsal and reaching a trifle nearer to caudal fin; first spine longer and more slender than the first regular dorsal spine, while the second equals the third dorsal in length. Caudal rounded, equaling in length the greatest depth of fish. Coloration in alcohol, pale, pretty regularly covered with very fine dark punctulations, thickest on back, palest below; a large dark opercular blotch, 2 similar postocular blotches, and usually a darkish bar extends downward from eye; upper half of preorbital region dark, outer margin of jaws dark; breast and under parts of head pale, top of head and nape dark; first dorsal quite dark, almost black; second dorsal pale, obscurely mottled with brown, which is disposed in about 5 indistinct areas; a large black ocellus upon the twelfth and thirteenth spines of second dorsal, and a similar one upon the twenty-third and twenty-fourth spines; each ocellus is surrounded by a narrow circle of white or pale orange. In the 6 specimens before us there is a slight variation as to the exact position of the 2 ocelli; in 1 example the second ocellus extends back upon the twenty-fifth spine also, but in every case the twelfth and thirteenth and the twenty-third and twenty-fourth are the spines which most evidently locate the spots; pectorals and ventrals plain; anal paler than dorsal, sparsely covered with fine dark points, so grouped as to form 3 or 4 darker areas. Length 2¾ inches. Gulf of California. Known from 6 specimens taken from masses of kelp hauled out by the seine from the bay of Guaymas. (Jenkins & Evermann.) (*asper*, rough.)

Auchenopterus asper, Jenkins & Evermann, Proc. U. S. Nat. Mus. 1888, 154, Guaymas, Mexico. (Type, No. 39643. Coll. Jenkins & Evermann.)

879. AUCHENOPTERUS, Günther.

Auchenopterus, GÜNTHER, Cat., III, 275, 1861 (*monophthalmus*).
Cremnobates, GÜNTHER, Proc. Zool. Soc. Lond. 1861, 374 (*monophthalmus*). Substitute for *Auchenopterus*, regarded as preoccupied on account of its similarity to *Auchenipterus*, a genus of *Siluridæ*.
Corallicola, JORDAN & EVERMANN, new subgenus (*marmoratus*).

Body moderately elongate, compressed, covered with rather large, cycloid scales; head shortish, naked, the snout rather pointed; cheeks full; mouth moderate, with a band of conical teeth in the jaws and about 1 series on the vomer, none on the palatines; lower jaw prominent; gill membranes united, free from the isthmus; upper surface of head with tentacles. Dorsal fin composed of stiff spines, with but a single soft ray, which is lower than the spines; first 3 spines more or less separated from the others, stiff and rather wider set, sometimes higher than the others; anal fin low, with 2 short spines; ventrals jugular, well developed; pectorals broad; lateral line complete, strongly curved anteriorly. Warm seas. This genus differs from *Cristiceps* in having but 1 soft ray in the dorsal fin, and in the large scales. (*αὐχήν*, nape; *πτερόν*, fin.)

CORALLICOLA (*Corallus*, coral; *colo*, I inhabit):
a. First 3 or 4 spines of dorsal forming a separate fin, being much higher than any of the spines in the posterior part of the fin; snout rather acute.
b. Scales 33; dorsal with 1 ocellus, anal with none; a black cross bar at base of caudal; a yellow spot behind eye; snout pointed. NIGRIPINNIS, 2712.
bb. Scales 37 or 38.
c. First dorsal spine longer than second; dorsal with 2 ocelli; anal blackish; D. IV–XXIV, 1. ALTIVELIS, 2713.
cc. First dorsal spine shorter than second; snout slender, very acute; caudal pale; dorsal with 2 ocelli, anal with 1; D. III–XXII, 1. MARMORATUS, 2714.

AUCHENOPTERUS:
aa. First 3 spines of dorsal scarcely forming a separate fin, none of them higher than the posterior spines; snout not very acute; anal without ocellus.
d. Caudal fin pale, usually with a dark bar at its base; a notch between third and fourth dorsal spines.
e. Dorsal spines about 31.
f. Scales 34 to 36; membrane of third spine joining fourth at its base; dorsal and anal plain dusky. AFFINIS, 2715.
ff. Scales 38.
g. Membrane of third spine joining fourth slightly above its base. MONOPHTHALMUS, 2716.
gg. Membrane of third spine joining fourth spine much above its base. INTEGRIPINNIS, 2717.
ee. Dorsal spines about 28; membrane of third spine joining fourth above its base; scales 38; body with distinct cross bars; dorsal with 1 ocellus. FASCIATUS, 2718.
dd. Caudal fin black; body chiefly black; head mottled with whitish; membrane of third dorsal fin joining fourth near its summit, the fin not notched; dorsal spines 30; dorsal with 2 ocelli. NOX, 2719.

Subgenus CORALLICOLA, Jordan & Evermann.

2712. AUCHENOPTERUS NIGRIPINNIS (Steindachner).

Head 4; depth 5⅓. D. XXVIII, 1; A. II, 27; scales 33; eye 4¼ in head; snout 4½, equal to interorbital space; snout pointed. Three first dorsal spines higher than the others and further apart. A tentacle over eye.

Scales of body much largest anteriorly; lateral line arched. A deep black spot with a white ring between the twenty-second and twenty-fourth spines; anal edged with white; black cross band at base of caudal with silvery point at upper base of pectoral; a diffuse yellowish spot below and behind eye. Barbados. One specimen 1 inch and 7 lines long. (Steindachner); not seen by us. (*niger*, black; *pinnis*, fin.)

Clinus nigripinnis, Steindachner, Ich. Notizen, vi, 46, 1867, Barbados.

2713. AUCHENOPTERUS ALTIVELIS (Lockington).

D. IV–XXIV, 1; A. 21; P. 13; C. 13; V. 2; scales 37. Body compressed, greatest depth a little behind pectoral axil; greatest thickness at gill covers; dorsal and abdominal profiles of similar curvature, decreasing regularly to the caudal fin; profile of occiput and superorbital regions convex; snout somewhat produced, its upper outline slightly concave. Head 4 in total length; greatest depth a little less than length of head; caudal peduncle about ¼ of the greatest depth. Eye round, lateral, with a slight direction upward, its diameter less than the length of the snout; interorbital area nearly equal in width to the diameter of the eye, concave transversely, upper orbital borders slightly raised. A short nasal tentacle slightly anterior to the front margin of the eye; a large fimbriated tentacle on each side of the first dorsal ray. Cleft of mouth oblique, the lower jaw the longer; the posterior convex extremity of the club-shaped maxillary about vertical with the center of the pupil. Teeth of the outer row regular, sharp, incurved, the largest in front, gradually decreasing along the lateral portions of the jaws, and not extending much past the middle of their length; a narrow band of small teeth in the rear of the outer row; vomerine teeth present. Branchiostegals 6; gill openings continuous, membranes not attached to the isthmus. Distance from first ray of dorsal to posterior margin of eye equal to length of snout; first 2 rays of dorsal much developed, the first slightly the longer, and nearly equal in height to the distance of its base from the tip of the upper jaw; third ray about ½ the length of the first; fourth very short; succeeding rays to the twenty-sixth longer than the third, the last 3 somewhat decreasing. Anal commencing under eleventh dorsal ray, coterminous with, and equal in height to, the dorsal. Caudal with 13 simple jointed rays, the longest in the center, posterior margin convex. Pectorals narrow, lanceolate, the fifth and sixth rays longest and ⅔ the length of the head. Ventrals inserted in advance of the pectorals. Lateral line with 37 simple pores, parallel with dorsal outline to opposite the origin of the anal, where it is deflected almost perpendicularly downward to the middle of the side of the body, along which it continues to its termination. Scales rather large, about 10 in a transverse row in the central part of the body, their posterior margin membranaceous; no scales on fins; a line of pores around the margin of the orbit, another along the posterior margin of the preoperculum, connected with each other and with the lateral line by a line from the center of the hinder border of the eye. Color in alcohol, bright pink above, becoming dusky below; underside of head light olivaceous, lower lip

blackish; dorsal pink, dusky on its margin, a black spot on the fourth ray, and another on its hinder part upon the twenty-fourth and twenty-fifth rays, the latter spot extending on to the body; membrane of anal black; occipital tentacles black. La Paz, Lower California. A single specimen, $1\frac{9}{10}$ inches long, dredged at a depth of 22 fathoms. (Lockington.) (*altus*, high; *velum*, sail.)

Cremnobates altivelis, LOCKINGTON, Proc. Ac. Nat. Sci. Phila. 1881, 116, La Paz, Lower California. (Coll. W. J. Fisher.)

2714. AUCHENOPTERUS MARMORATUS (Steindachner).

Head $3\frac{3}{5}$ to $3\frac{4}{5}$ in body; depth $3\frac{1}{2}$ to $3\frac{5}{6}$. D. III–XXII, 1; A. II, 19; scales 2–36–9 (28 or 29 anteriorly); eye 4 to 5 in head; first dorsal $1\frac{1}{4}$; pectoral $1\frac{1}{2}$. Body comparatively deep, compressed, the back somewhat arched; head pointed; mouth large, the maxillary extending to behind the eye, 2 in head; opercle with a sharp spine; jaws equal; teeth pointed, in narrow bands, the outer larger; vomerine teeth in 1 row; supraocular tentacle small, about as large as nuchal tentacle; no nasal tentacle. Pectoral a little shorter than head; dorsals separate, the first dorsal higher than second dorsal, the spines of which are about $\frac{1}{2}$ head. Color in life of varying shades of olive gray or sand color, with a series of whitish blotches on head and along sides; markings on dorsal and anal whitish; 2 dark-blue ocelli on dorsal and 1 on anal, these edged with orange and interiorly with black; ventrals, pectorals, and caudal whitish, barred with clear orange red; first dorsal black at tip; a curved blackish line at base of caudal; lower side of head yellowish brown, with whitish bands; specimens from coral reefs more spotted. Florida Keys to Cuba; common in the eelgrass at Key West. Our specimens, 2 to $2\frac{1}{2}$ inches long, taken at Key West and Havana. (*marmoratus*, marbled.)

Cremnobates marmoratus, STEINDACHNER, Ichth. Beiträge, v, 174, pl. 12, f. 6. 1876, a small rocky island north of Cuba; JORDAN & GILBERT, Synopsis, 962, 1883; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1884, 142.

Subgenus **AUCHENOPTERUS.**

2715. AUCHENOPTERUS AFFINIS (Steindachner).

Head 4; depth $4\frac{2}{5}$. D. III–XXVII, 1; A. II, 19; V. I, 2; scales 33 to 35. Form of *A. integripinnis;* maxillary reaching to below posterior margin of eye; a fringed tentacle above eye and 1 on each side of occiput. First dorsal low, its longest (second) ray shorter than the highest of second dorsal; membrane of third spine joining the fourth spine just above its base; last ray of second dorsal joined by membrane to base of caudal. Dark brown, paler than in *A. nox*, but darker and more uniform than in *Auchenopterus fasciatus;* lower side of head pearly gray, thickly speckled with darker; sides with 5 very faint darker cross bands; dorsal and anal dusky, the latter with a pale edge; between the eighteenth and twenty-second dorsal spines a large dark spot ocellated with yellowish; caudal yellowish white, with darker cross streaks, a blackish band at its base; pectoral

dusky at base, its posterior half yellowish, with darker cross streaks; ventral similar; a wedge-shaped, whitish band extending backward from eye to opercle. West Indies; recorded from Key West and St. Thomas. Here described from specimens from Key West. (*affinis*, related,—to *A. monophthalmus*.)

Cremnobates affinis, STEINDACHNER, Ichth. Beiträge, v, 178, 1876, St. Thomas; JORDAN, Proc. U. S. Nat. Mus. 1884, 142; JORDAN, Cat. Fishes N. A., 121, 1885.

2716. AUCHENOPTERUS MONOPHTHALMUS, Günther.

Head $3\frac{4}{5}$; depth 4. D. III–XXVI, 1; A. II, 18; scales 2–32–9; eye 5 in head; maxillary $1\frac{4}{5}$; pectoral $1\frac{1}{5}$; caudal $1\frac{1}{4}$. Body compressed, deepest at middle of pectorals; head moderately pointed, the upper profile slightly and evenly convex; mouth large, maxillary reaching past eye; jaws subequal; teeth villiform, in bands on jaws, vomer, and palatines; interorbital space flat, as wide as eye; a multifid dermal flap over posterior edge of eye, and a smaller one on each side of nape; head naked; body with rather large, regular scales; fins naked. Origin of dorsal over edge of preopercle, the first 3 spines separated from rest of fin by a rather deep notch, the membrane from third spine joining fourth spine at about its middle; spines of posterior part of dorsal the highest; front of anal midway between tip of snout and base of caudal, tips of last rays reaching slightly beyond base of caudal and tips of last dorsal rays; pectorals reaching front of anal; ventrals long and slender, inserted in front of base of pectorals a distance equal to $1\frac{1}{2}$ eye; caudal rounded. Color light grayish red or brown, with about 6 cross bars of darker brown, running up on dorsal; between the bars are scattered milky white irregular spots; a black spot, ocellated with white, on front of dorsal, a similar spot near posterior end, sometimes duplicated; narrow cross bars on anal; a dark bar on base of caudal, and a dark blotch on base of pectoral. Here described from specimens, a couple of inches in length, from La Paz, Lower California. Gulf of California to Panama, abundant in rock pools, creeping about among *Corallina;* close to *A. integripinnis*, but the first dorsal higher and more separate from rest of fin. (μόνος, one; ὀφθαλμός, eye, from the dorsal ocellus.)

Auchenopterus monophthalmus, GÜNTHER, Cat. Fish., III, 275, 1861, Panama; JORDAN, Proc. Cal. Ac. Sci. 1895, 501.

Cremnobates monophthalmus, GÜNTHER, Proc. Zool. Soc. Lond. 1861, 374; GÜNTHER, Fish. Centr. Amer., 442, pl. 69, fig. 1, 1869.

2717. AUCHENOPTERUS INTEGRIPINNIS (Rosa Smith).

Head $3\frac{1}{3}$; depth $4\frac{1}{2}$; eye 4 in head. D. III–XXVII, 1; A. II, 20; scales 2–36–9; pectoral $1\frac{1}{3}$; caudal $1\frac{1}{3}$. Head stout, broad, conical; mouth little oblique, maxillary reaching posterior margin of eye; eyes large; nasal, supraocular and nuchal regions with fringed cirri, those at the nape flaplike. First and second dorsal spines low, a little higher than the third, which, in turn, is higher than the fourth and separated from it by an interspace, the membrane between the third and fourth spines not deeply

emarginate, membrane from third spine attached to the lower $\frac{2}{3}$ of fourth; anterior spines not forming a separate fin; highest anterior spine not higher than the highest of the posterior part of fin. Color dark brown, variegated with different shades of brown and reddish; about 5 indistinct dark cross bars; a distinct ocellated black spot on posterior part of dorsal fin; caudal fin abruptly translucent, speckled, a black bar at its base; base of pectorals violet, bordered with black, the rest of the fin checkered; ventrals barred. Length $2\frac{1}{2}$ inches. Coast of California and southward to Todos Santos; abundant in rock pools among *Corallina*. Here described from a specimen, $1\frac{1}{2}$ inches in length, from San Cristobal, Lower California. (*integer*, entire; *pinna*, fin.)

Cremnobates integripinnis, Rosa Smith, Proc. U. S. Nat. Mus. 1880, 147, La Jolla, near San Diego (Coll. Rosa Smith); Jordan & Gilbert, Synopsis, 764, 1883.

2718. AUCHENOPTERUS FASCIATUS (Steindachner).

Head 4; depth $4\frac{1}{3}$. D. III–XXIV, 1; A. II, 18; scales 37. Body rather slender, a little deeper than in *A. integripinnis*, the snout less acute than in *A. marmoratus*. First dorsal spine rather higher than second and lower than the spines of posterior part of fin; membrane of third spine joining second dorsal at a point above its base, the two parts of the fin therefore separated only by an emargination. Tentacle above eye slender, small; cirri on side of occiput bluish. In life, light pinkish brown, much mottled, and with traces of 6 to 8 faint darker bars; head and its cirri above whitish; 3 blackish spots behind eye, radiating from it, the lower one largest; preopercle with 3 dark dots; dorsal pale, with 9 blackish blotches, in the next to the last of which is a large blue-black ocellus, edged with orange; anal with 5 dark blotches and no ocellus; a blackish bar across base of caudal; rest of caudal and pale part of anal with dark dots; ventrals whitish, barred with black; pectoral similar, its base with a whitish area, which has a brown center, below which is a small black spot. Length 2 inches. Florida Straits; north to Key West. Here described from specimens from Key West. (*fasciatus*, banded.)

Cremnobates fasciatus, Steindachner, Ichth. Beiträge, v, 176, 1876, Florida Straits; Jordan, Proc. U. S. Nat. Mus. 1884, 142; Jordan, Cat. Fishes N. A., 121, 1885.

2719. AUCHENOPTERUS NOX (Jordan & Gilbert).

Head $3\frac{3}{5}$; depth $3\frac{5}{6}$. D. III–XXVII; A. II, 18; lateral line with 34 tubes. Snout not very acute, the upper and lower profiles of head nearly evenly convex; mouth large, maxillary reaching slightly beyond eye, $\frac{1}{2}$ length of head; eye large, equaling length of snout, greater than interorbital width, 4 in head (to end of opercular spine); interorbital width $4\frac{3}{4}$ in head; nasal, supraorbital, and occipital tentacles present, those on snout and above the orbits simple, slender filaments, the latter about as long as diameter of orbit, 1 of them divided to the base, the other simple; the tentacle on each side of nape a compressed slip of skin higher than wide, the margin uneven, but not fringed. Anterior dorsal spines not much elevated,

not higher than some of the posterior spines; the first and second spines about equal, 2½ in head; the third spine shorter, about equal in length to the fourth, from which it is separated by a wide membrane, which is, however, not at all notched; the spines thence increase in length toward the last; caudal 1⅛ in head; pectorals reaching anal, nearly equaling length of head; ventrals not reaching vent, 1⅓ in head. Scales large, 4 series above lateral line and 4 below. Color, body and fins uniform blackish brown; a few small silvery-white specks on dorsal region, mostly along base of dorsal fin; head and base of pectoral fin with light pink areas and mottlings; snout pink above; nape with a pink cross bar; a dark streak upward and backward from eye to nape; a light streak from eye backward to opercle and 1 backward and downward; lower jaw mottled with light and dark; a small round, black spot near base of dorsal between twenty-third and twenty-fifth spines, and 1 between twenty-eighth and thirtieth, both very faintly ocellated with lighter; slight whitish tips on ventrals and lower edge of caudal. Key West; known from a single specimen, 1¾ inches long, taken with the seine in algæ on a rocky bottom at Key West. Its congeners, *A. marmoratus*, *A. fasciatus*, and *A. affinis*, were found in the same waters, *A. marmoratus* being much the most abundant of the 4, and reaching the largest size. (*nox*, night.)

Cremnobates nox, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1884, 30, Key West. (Coll. Jordan.)

88ò. PARACLINUS, Mocquard.

Acanthoclinus, MOCQUARD, Bull. Soc. Philom. Paris 1885, 18 (*chaperi*); name preoccupied.
Paraclinus, MOCQUARD, Bull. Soc. Philom. Paris 1886, 11 (*chaperi*).

Body elongate, compressed, covered with cycloid scales; mouth large, each jaw armed with an external row of conical teeth, with some teeth behind; teeth on the palate; dorsal very long, continuous, composed entirely of spines, anal with 2 spines; ventrals jugular, with few rays; tentacles on head; gill opening very broad; 6 branchiostegals; lateral line interrupted. Evidently very close to *Auchenopterus*, from which it may be distinguished by the continuous dorsal fin, a character which needs verification. (παρά, near; *Clinus*.)

2720. PARACLINUS CHAPERI, Mocquard.

Head 4¼; depth 4¼. D. XXXI; A. II, 19; P. 13; V. 2; scales 35. Body elongate, very strongly compressed; eye large, equal to snout or interorbital width; lower jaw slightly the longer; mouth oblique, reaching front of eye; outer row of teeth strong, canine-like, slenderer and more close set above, below diminishing rapidly in length, the bands of small teeth limited to front of each jaw, a curved group of teeth on palate; dorsal beginning over preopercle, not notched, composed entirely of stout spines; anal equally long; ventrals very narrow, of 2 soft rays, well separated, the inner slightly longer than outer; head with 3 pairs of tentacles, 1 at the nape, filiform, small, ½ as long as eye; the second below the orbit, broadened at base, separated into 3 or 4 branches, progressively

longer from the inner outward, longer than eye; nuchal filament a little in front of dorsal, in form, oblong, entire, laminated, a little broader at its free edge, ¾ as long as eye; scales large, cycloid; lateral line interrupted before front of anal, anterior part rounding over eye with only 2 rows of scales between it and the dorsal, posterior part median; gill membranes broadly united, free from isthmus. Body brownish yellow, fins brown, the base and the caudal darker. Bay of Guanta, near Barcelona, in Venezuela; 1 specimen, 33 mm. long to base of caudal. (Mocquard.) Not seen by us. (Named for its collector, M. Chaper.)

Acanthoclinus chaperi, MOCQUARD, Bull. Soc. Philom. Paris 1885, 19, Bay of Guanta, Venezuela.

Paraclinus chaperi, MOCQUARD, Bull. Soc. Philom. Paris 1886, 41.

881. EMMNION, Jordan.

Emmnion, JORDAN, in Gilbert. Proc. U. S. Nat. Mus. 1896, 454 (*bristolæ*).

Body elongate, covered with caducous, cycloid scales of small size; lateral line straight, ending near base of last dorsal ray. Head moderate, decurved anteriorly, without cirri; mouth moderate; teeth in jaws in bands, the outer enlarged; no teeth on vomer or palatines; dorsal notched, its anterior ⅔ of flexible spines of moderate height; ventrals I, 3, the rays thickish, the fin inserted slightly before pectorals; caudal free. Galapagos Islands. (*ἐν*, in; *μνίον*, sea moss, or alga.)

2721. EMMNION BRISTOLÆ, Jordan.

Head 5⅔; depth 7½. D. XXV, 13; A. I, 27; P. 13; V. I, 3; Br. 5; scales 3–50–11, the count not certain. Body slender, moderately compressed, the dorsal profile forming a nearly straight line from occiput to first dorsal ray, from thence descending very gently to base of caudal; ventral profile about straight. Head broad, slightly convex above, its width 1⅓ in its length; anterior profile from first dorsal spine to a point above eye straight, thence abruptly descending to tip of snout; mouth horizontal, the lower jaw included; maxillary reaching nearly to posterior margin of eye, about 2½ in head. Teeth present on both jaws, canine-like; upper jaw with 8 enlarged teeth in front, about 2 or 3 series of much smaller teeth behind these, only 1 series of which extends into posterior region of mouth; lower jaw with a series of teeth in front and on sides which is greatly enlarged in front; a patch of very small teeth behind the enlarged front teeth; no teeth on vomer or palatines. Premaxillary very protractile; snout blunt, 4⅙ in head; eyes large, round, placed close together, 3¼ in head; interorbital region very narrow, less than pupil; nostrils equal. Caudal peduncle 2⅘ in head; branchiostegal membranes deeply united, free from isthmus; gills 4, a small slit behind the fourth; no cirri above eyes, nor filaments on nape; head naked, body covered with cycloid scales, those on nape much smaller; belly naked. The scales on the body are apparently caducous as all have fallen, but the points are very distinct and they seem to have been embedded on their anterior edge,

as the sac-like fold of skin is prominent. Lateral line simple, straight, running from upper edge of gill opening to last ray of dorsal when it is lost, not reaching the caudal; it is placed very high, and gradually approaches the dorsal fin, from which it is separated only by a very small distance. Dorsal extending from a point a short distance behind occiput nearly to base of caudal, emarginate; last spine shortest, about 2¼ in first soft ray, which is 2½ in head; the longest spines about 3 in head, all the spines slender and flexible. Anal extending from behind vent nearly to base of caudal; similar to soft dorsal, its rays lower. Ventrals well developed with broad base, the rays thickish, inserted very slightly in front of base of pectorals, 1⅓ in head, reaching ⅔ the distance to vent. Caudal subtruncate. Pectorals reaching past vent, about as long as head. Dorsal and anal free from caudal. Color in spirits, dark dull reddish-brown, lighter below; head very dark; dorsals, pectorals, and caudal blackish, pectorals and caudal with lighter blotches; anal and ventrals dusky, anal margined with darker. Length about 3 inches. Galapagos Islands; one specimen known, evidently a rock-pool species. (Named for Miss Susan Brown Bristol, of the department of zoology in Leland Stanford Junior University.)

Emmnion bristolæ, JORDAN, in GILBERT, Proc. U. S. Nat. Mus. 1896, 454, pl. 55, fig. 1, Galapagos Islands. (Coll. *Albatross.*)

882. ATOPOCLINUS, Vaillant.

Atopoclinus, VAILLANT, Bull. Sci. Philom. Paris, serie 8, tome VI, 1894, 73 (*ringens*).

Body elongate, subcylindrical, without visible scales. Head obtuse, the snout short, rounded; mouth inferior, transverse, with compressed trenchant teeth in each jaw, those above at least in a single row, solidly fastened to the skeleton; teeth on vomer and palatines uncertain. Dorsal continuous, extending the whole length of the back, from the nape to the caudal peduncle, its rays mostly simple, only the posterior articulate; anal occupying nearly ½ the length, touching the caudal, which is, nevertheless, distinct; caudal deeply forked; ventrals distinctly jugular, very long, of a spine and a ray; no tentacles; gill membranes apparently rounded at the isthmus. Gulf of California; a singular genus evidently closely allied to *Runula*. (ἄτοπος, strange; *Clinus.*)

2722. ATOPOCLINUS RINGENS, Vaillant.

Head 5; depth 7. D. 24; A. 18; P. 15; V. I, 1. Eye large, 7 in head; interorbital space broad, 3 in head. Caudal a little longer than head. Color clear chamois brown, the belly pale; a brown band before the snout, across the eye to the caudal, on which it extends; a silvery stripe bordering this band above, and below for part of its length. Gulf of California. (Vaillant); known from 1 specimen badly shriveled, 39 mm. in length. (*ringens*, gaping.)

Atopoclinus ringens, VAILLANT, Bull. Sci. Philom. Paris, serie 8, tome VI, No. 2, February 25, 1894, 74, Gulf of California. (Coll. Léon Diguet.)

883. RUNULA, Jordan & Bollman.

Runula, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 171 (*azalea*).

Body slender, its back not elevated; mouth small, inferior, destitute of canines; teeth fixed, upper largest; dorsal fin continuous, its spines and soft rays indistinguishable, most of them articulate; caudal fin lunate; gill openings reduced to a vertical slit in front of pectoral; scales none. This genus is remotely allied to the East Indian genus *Petroskirtes*, but has the mouth and dentition different, and the caudal fin, unlike that of most blennioid fishes, is forked. (Diminutive of *runa*, a dart or javelin.)

2723. RUNULA AZALEA, Jordan & Bollman.

Head $4\frac{4}{5}$; depth $6\frac{1}{5}$. D. 42; A. 26 or 27; V. I, 2. Body moderately elongate, not much compressed; head rather long, its upper outlines convex; snout short and very blunt; mouth entirely inferior, transverse, each jaw provided with long, slender, close-set curved teeth; no evident posterior canines; eye moderate, equal to snout and nearly equal to interorbital width, 4 in head; no tentacles on head; gill membranes fully united to the isthmus, the gill opening reduced to a vertical slit, its lower edge opposite middle of base of pectoral; no scales; lateral line very high, concurrent with the back; dorsal fin very low, continuous; the feeble spines and soft rays indistinguishable, the fin beginning at occiput; anal similar to soft dorsal; caudal lunate behind, well separated from dorsal and anal; pectorals small, rounded, about $1\frac{4}{5}$ in head; ventrals short, before pectorals, about 2 in head. Color reddish brown, silvery below, about 5 dusky cross shades; a dusky lateral streak; a black spot surrounded by paler at base of caudal; dorsal with about 6 black crossbars; anal with 4; other fins pale; lower half of head abruptly pale. Galapagos Archipelago. The type, 2 inches long, taken at Indefatigable Island; 3 more specimens have since been obtained from the same island. (ἀζαλέος, parched, from the brown color.)

Runula azalea, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 171, Indefatigable Island, Galapagos Archipelago (Coll. *Albatross*); JORDAN, Proc. Cal. Ac. Sci. 1896, 233, pl. 37.

884. BLENNIUS (Artedi) Linnæus.

(BLENNIES.)

Blennius, ARTEDI, Genera Piscium, 27, 1738.
Blennius, LINNÆUS, Syst. Nat., Ed. X, 256, 1758 (*galerita*).
Salaria, FORSKÅL, Descr. Anim., 22, 1777 (*basiliscus*).
Pholis, FLEMING, Brit. Anim., 207, 1828 (*lævis=pholis*); not *Pholis* SCOPOLI, 1777.
Adonis, GRONOW, Cat. Fish., Ed. Gray, 93, 1754 (*pavoninus=ocellaris*).
Lipophrys, GILL, American Naturalist, June, 1896, 498 (*pholis*).

Body oblong, compressed, naked; head short, the profile usually bluntly rounded; mouth small, horizontal, with a single series of long, slender, curved, close-set teeth in each jaw, besides which, in the lower jaw at least, is a rather short and stout fang-like canine tooth on each side;

premaxillaries not protractile; gill openings wide, extending forward below, the membranes free from the isthmus, or at least forming a broad fold across it. Dorsal fin entire, or more or less emarginate, the spines slender; pectorals moderate; ventrals well developed, I, 3; no pyloric cæca; lateral line developed anteriorly. Species numerous, lurking under rocks and algæ in most warm seas; some species in the lakes of northern Italy. The European species in general are larger in size than ours, with higher fins. (*Blennius*, the ancient name, from *βλέννα*, slime.)

LIPOPHRYS (λείπω, to disappear; ὀφρύς, eye-brow):

a. Supraorbital cirrus wanting; snout not very blunt in profile.

b. Posterior canine present in each jaw; dorsal slightly emarginate; D. XII, 18. CAROLINUS, 2724.

BLENNIUS:

aa. Supraorbital cirrus present; profile of snout more or less blunt.

c. Canines strong, present in both jaws; no nuchal cirri.

d. Dorsal rays XI or XII, 17 or 18.

e. Supraorbital cirrus bifid; dorsal free from caudal.

f. Supraorbital cirrus as long as head; dorsal emarginate; sides spotted; D. XI, 17. FUCORUM, 2725.

ff. Supraorbital cirrus as long as eye and snout; dorsal continuous; color olivaceous, with dark bars; D. XI, 18. STEARNSI, 2726.

ee. Supraorbital cirrus bifid, nearly as long as head; last ray of dorsal joined to caudal; sides with a network of blue lines; D. XII, 18. FAVOSUS, 2727.

dd. Dorsal rays XII, 21 or 22; supraorbital cirrus long, fringed; dorsal free from caudal; cheeks with network of lines; body nearly plain. PILICORNIS, 2728.

cc. Canines short and stoutish, present in lower jaw only (undescribed in *truncatus* and in *marmoreus*.)

g. Nape without cirrus; snout abruptly decurved; body robust, marbled; D. XII, 20. MARMOREUS, 2729.

gg. Nape with a cirrus on each side.

h. Dorsal and anal free from caudal. Nape with a filiform bifid tentacle on each side; teeth undescribed; supraorbital tentacle simple; color olive, with bright spots. TRUNCATUS, 2730.

hh. Dorsal and anal with the last ray largely joined by membrane to caudal; nape with a small cirrus; posterior canines strong, in lower jaw only; dorsal not notched; color uniform brown; D. XII, 13. VINCTUS, 2731.

ggg. Nape with a comb of many close-set cirri on a fleshy crest; lower jaw only with short posterior canines; dorsal fin continuous, free from caudal; D. XII, 16 or 17. CRISTATUS, 2732.

Subgenus LIPOPHRYS, Gill.

2724. BLENNIUS CAROLINUS (Cuvier & Valenciennes).

D. XII, 18; A. 17. Body rather long and slender, more elongate than in the European species, *Blennius pholis*, more compressed, the head longer; maxillary extending to opposite middle of eye; teeth $\frac{1}{1}^{\frac{0}{0}}$, with strong canines on both jaws; gill membranes free from isthmus; no trace of tentacles above eye; dorsal spines slender, a little lower than the soft

rays, the fin little emarginate; dorsal and anal not joined to the caudal. Greenish, with 4 or 5 irregular dark spots or shades along the back; dorsal with a large black spot in front; anal brown-edged. South Carolina. Only the original type in the museum at Paris known; from this the present description was taken. No later collector has recognized the species and it may not be American.

Pholis carolinus, CUVIER & VALENCIENNES, Hist. Nat. Poiss, XI, 276, 1836, Carolina. (Coll. M. Bosc.)
Blennius carolinus, JORDAN & GILBERT, Synopsis, 760, 1883.

2725. BLENNIUS FUCORUM, Cuvier & Valenciennes.

Head 5 in total length. D. XI, 17; A. 18. Orbitial cirri nearly as long as head, bifid at tip, and fringed at the base. Dorsal fin slightly emarginate, free from the caudal, the spines rather stiff. Head very short and steep, its profile nearly vertical; 24 teeth in each jaw; each jaw with very strong canines; gill membranes free from the isthmus posteriorly. Olive green, becoming darker above, with numerous brown spots on the cheeks and sides of the body; below reddish; dorsal with a large black spot in front, behind which are smaller spots; spinous dorsal edged with paler. (Cuvier & Valenciennes.) Open ocean in floating *Fucus*; the type from near the Azores; recorded by De Kay from the open sea, off New York, in floating seaweed. (*fucorum*, of the seaweed, *Fucus*.)

Blennius fucorum, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 263, 1836, 240 miles south of the Azores (Coll. Claude Gay); GÜNTHER, Cat., III, 217, 1861; DE KAY, N. Y. Fauna: Fishes, 149, pl. 22, fig. 66, 1842; JORDAN & GILBERT, Synopsis, 710, 1883.
Blennius oceanicus,* CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 265, 1836, open sea, 29° N., 50° W.; on a drawing by CLAUDE GAY.

2726. BLENNIUS STEARNSI, Jordan & Gilbert.

Head $3\frac{4}{5}$ ($4\frac{3}{8}$ in total); depth $4\frac{3}{8}$ ($5\frac{3}{8}$); eye $4\frac{1}{3}$; snout $4\frac{1}{3}$. D. XI, 18; A. II, 21. Body much elongate, compressed, tapering regularly behind; anterior profile moderately decurved; snout short and blunt; mouth large, oblique, the jaws even; maxillary reaching slightly beyond middle of orbit, $2\frac{1}{4}$ times in head; teeth in the front of the jaws only, occupying on each side a space equal to $\frac{1}{2}$ length of maxillary; teeth $\frac{3}{2}\frac{0}{4}$, the lateral one on each side much enlarged and canine-like, rather short but strongly curved; canine in upper jaw equaling about $\frac{1}{2}$ diameter of pupil; interorbital space very narrow, not as wide as pupil; upper posterior rim of orbit with a long slender filament, forked at base, its length equaling distance from tip of snout to posterior rim of orbit; no filaments at the nape; gill membranes somewhat united to the isthmus in front, but forming a broad fold across it posteriorly, the gill openings of the two sides therefore continuous below. Dorsals rather high; no notch between the spines and soft portion, the membrane of last ray not reaching base of caudal; spines of

* Very near *Blennius fucorum*, the profile more oblique, the cirri shorter, the spinous dorsal lower, the caudal more truncate; anal shorter. Color brown with brown spots on body and fins; sides clear green; belly silvery. Length 2 inches. (Cuvier & Valenciennes.)

nearly uniform height, all very slender and flexible, the tips almost filamentous; highest spine ½ length of head; highest soft ray 1⅔ in head; anal lower than dorsal, its longest ray very slightly less than ½ length of head; length of caudal peduncle more than ½ its height, about equaling the diameter of orbit; caudal about equaling pectoral, 1⅓ in head; ventrals long, the inner ray much the longest, 1⅓ in head, not quite reaching vent. Color light greenish olive, somewhat mottled; sides with irregular dark bars formed of spots, these extending on the fin; skin everywhere finely punctate; dorsal dark olive, the spinous part darker at tip; anal blackish, with paler edge; ventrals dusky; pectorals and caudal olive. Gulf of Mexico, in deep water. Three specimens known, the largest 3 inches long, taken from the stomach of a Red Snapper, at Pensacola. (Named for Silas Stearns.)

Blennius stearnsi, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 300, Pensacola Snapper Banks. (Type, No. 29669, U. S. Nat. Mus. Coll. Jordan & Stearns.)

2727. BLENNIUS FAVOSUS, Goode & Bean.

Head 3⅔; depth 4¾. D. XII, 18; A. II, 20. Body comparatively elongate and compressed; anterior profile moderately decurved; head nearly ½ longer than deep; snout very short and blunt; mouth large, horizontal; jaws even; maxillary reaching posterior margin of orbit, its length 2½ in head. Each jaw with a long, curved, posterior canine; the canines of lower jaw largest. Preorbital ⅔ diameter of eye, which is 3¾ in head, and equals more than twice interorbital width. An extremely long and slender supraocular cirrus, trifid to the base, the longest branch nearly as long as the head; no nuchal cirri. Gill membranes forming a rather narrow fold across the isthmus. Dorsal low, continuous, the spines very slender and flexible, the longest ½ as long as the head; the longest soft ray ⅔ as long as head; the last ray slightly joined to base of caudal; caudal ¾ as long as head; anal rather high; pectorals ⅔ as long as head; only the straight part of lateral line developed. Color faded, brownish, finely reticulated, a series of obscure bluish blotches along the sides; front and sides of head marked with very distinct blue, reticulating lines surrounding honeycomb-like hexagonal interspaces; top of head with many small blue spots; dorsal with black dots and streaks; a black spot bordered with whitish between the first and second dorsal spines; anal with oblique blue streaks, the fin margined with dusky, tips of rays whitish; base of pectorals with blue reticulations. The whole body was probably reticulated with blue in life. Gulf of Mexico. Known from 2 specimens collected at Garden Key, Florida, by Gustav Würdemann; they are 3⅔ inches and 3 inches long, respectively. (*favosus*, honeycombed.)

Blennius favosus, GOODE & BEAN, Proc. U. S. Nat. Mus. 1882, 416, Garden Key, Florida (Type, No. 2629, U. S. Nat. Mus. Coll. Gustav Würdemann); JORDAN & GILBERT, Synopsis, 961, 1883.

2728. BLENNIUS PILICORNIS, Cuvier & Valenciennes.

Head 4¾ with caudal; depth 5⅕. D. XII, 21 or 22; A. 23 or 24. Snout obtuse, the upper profile very oblique. A strong curved canine in each jaw. Orbital tentacle filiform, with several smaller ones at base. Inter-

orbital space flat, its width ½ vertical diameter of eye; no groove or crest on the neck. Dorsal slightly notched, the spines flexible; caudal separate. Brown, dorsal and caudal spotted with darker. Length 5 to 6 inches. (Günther.) Coast of Brazil north to the West Indies, recorded from Rio Janeiro, Bahia, and the Tortugas, and off the coast of Florida. Mr. Garman gives the following color note on Tortugas specimens, collected by Prof. C. C. Nutting: Small, hexagonal reticulations on cheeks, resembling scales; anal darker toward ends of rays, the tips white; dorsal darker in outer half; basal part of dorsal and anal pale, sides with a few scattered black dots; median rays of caudal longer, the outer margin dark; caudal, pectorals, and ventrals paler than dorsal. (*pilicornis*, with downy horns.)

Blennius pilicornis, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 254, 1836, Rio Janeiro (Coll. Delalande and Gay); CASTELNAU, Anim. Nouv., etc., Amer. Sud, 25, 1855; GARMAN, Bull. Iowa Lab. Nat. Sci. 1896, 89.
Blennius filicornis, GÜNTHER, Cat., III, 216, 1861. (Coll. M. Parzudaki.)

2729. BLENNIUS MARMOREUS, Poey.

Head 4½ in total length with caudal; depth 5. D. XII, 20; A. 16; P. 13. Eye very high, near the profile, twice length of snout. Snout round, falling off abruptly, but less so than in *Blennius truncatus*; posterior nostrils with a distinct tube; superciliary tentacle divided into 3 branches; no cilia at the nape. Teeth undescribed. Gill membranes not described. Pectoral and caudal round; dorsal low, the median spines highest, the soft rays a little higher, the difference slight. Color yellowish brown, darker medially, paler below; under the lens covered with small dots; fins below yellowish. This species differs from *Scartella microstoma* in the stout trunk, the more blunt head, the cilia on the head and in the tube of the nostril. Cuba; 1 specimen 2 inches long. (Poey); not seen by us; perhaps not a *Blennius* as here understood. (*marmoreus*, marbled.)

Blennius marmoreus, POEY, Enumeratio, 130, 1875, Cuba. (Coll. Rafael Arango.)

2730. BLENNIUS TRUNCATUS (Poey).

Head 5¼ in total length with caudal; depth 5¼. D. XII, 19; A. I, 20. Eyes placed very high, profile before them vertical, suggesting the forehead of a bull without horns; mouth small, maxillary reaching below posterior border of eye; anterior nostril divided into 5 at tip; 2 filiform tentacles with a common base on each side of nape; a simple tentacle behind eye; some pores on the head, which is compressed; teeth undescribed; gill membranes undescribed; gill membranes united and free from isthmus; dorsal notched medially; caudal truncate, with 2 faint angles; lateral line long, reaching beyond the point of the pectorals. Color olive, with some bright spots on trunk; the vertical fins darker. Cuba; 1 specimen 3¼ inches long. (Poey); not seen by us; perhaps not a species of *Blennius*. (*truncatus*, cut off short.)

Blennius truncatus, POEY, Memorias, II, 424, 1861, Cuba. (Coll. Poey.)

2731. BLENNIUS VINCTUS, Poey.

Head 3½ to base of caudal; depth 4. D. XII, 13; A. I, 8; V. 3. Eye high, 4 in head, as long as snout. Anterior nostril in a short tube. Jaws equal; 4 pores on the side of the lower jaw; 1 on the opercle; 4 on the suborbital; 4 below eye. A long tentacle above eye; another very small one on the nape. Maxillary reaching to below front of pupil. Teeth large, not pointed, compressed, in 1 series of 10 to 12 on each side of each jaw, feeble, somewhat moveable; gill membranes united, free from isthmus. Dorsal elevated backward, connected by a membrane to the first third of the caudal, as is also the anal, twenty second ray highest, its height ½ depth of body and double length of the dorsal ray above tip of pectoral; anal similar, ⅓ also of the rays of the dorsal and anal simple; the spines flexible, differing from the others in not being articulate; pectoral pointed, its middle rays longest, and also more robust, all simple; ventral not very short; caudal rounded. Lateral line forming a curve anteriorly. Color uniform brown. Cuba. (*vinctus*, bound.)

Blennius vinctus, Poey, Repertorio, 243, 1867, Havana. (Coll. Felipe Poey. Type,* No. 12647, Mus. Comp. Zool.)

2732. BLENNIUS CRISTATUS, Linnæus.

Head 4; depth 4. D. XI, 16; A. 19; maxillary 3. Body moderately elongate, compressed; the head very blunt and deep, almost as deep as long, its anterior profile straight or slightly concave, and nearly vertical. Mouth moderate, the maxillary reaching to past front of eye; lower jaw with 2 short stoutish posterior canines, scarcely longer than the front teeth; upper jaw without canines. Teeth about $\frac{3}{2}\frac{2}{3}$. Preorbital deep, its depth equal to diameter of eye and contained 4½ times in length of head. Interorbital space flat, narrow, ⅔ width of eye. Supraocular cirri small, fringed, their length about equal to that of pupil. Nape with a longitudinal dermal crest reaching to front of dorsal, provided with a series of about 20 filaments, the longest about as long as the eye. Gill membranes forming a broad fold across the isthmus, as in all species of *Blennius*. Dorsal nearly continuous, the last spine a little lower than the first soft ray, not very high, beginning on the nape in front of the vertical of the preopercle, the spines all slender and flexible, the longest ⅔ as long as the head, the longest soft ray ½ as long as the head; caudal free from dorsal and anal, ½ as long as head; anal moderate, ⅔ length of head; pectoral somewhat shorter than head; ventral a little more than ½ length of head. Lateral line forming the usual arch above pectoral, and continued backward on median line to base of caudal, becoming indistinct posteriorly.

*On the type of *Blennius vinctus* we have the following notes: "No. 12647, M. C. Z. Cuba. (Poey.) One and a half inches long, in poor condition. Head ca 3½; depth ca 4. D. XII, 13; A. II, 13. Dorsal joined to caudal as far as tips of the rays, which are high. Dorsal spines high and stiff, the fin not notched, the soft rays higher. A thick scale-like fringed cirrus above each eye, nearly as long as eye, which is small. Gill membrane free. Head blunt. Maxillary to front of pupil. Lower jaw with very strong canines; upper jaw with none. No nuchal cirri."

Color faded, apparently olivaceous, with about 6 dark cross bars, which extend on the dorsal fin; anal and posterior ½ of body with numerous round, whitish, stellate spots, probably bluish in life; bluish streaks from eye across the cheeks; anal edged with dusky; the other fins vaguely marked. Length 2½ to 4 inches. Tropical parts of the Atlantic, among rocks, widely diffused and variable. The above description from the type of *Blennius asterias*, from Garden Key, Florida. We have the following notes on numerous specimens from Abrolhos Islands, off the Coast of Brazil (Coll. *Albatross*): D. XII, 15. Nape with a fringed crest of 10 to 18 filaments. A small trifid tentacle above eye; posterior canines in lower jaw only, short and small; gill membranes broadly united, nearly free from the isthmus. Dorsal slightly notched; nasal tentacle present. Color excessively variable, mostly grayish, with 5 or 6 cross blotches on the back, extending to form quadrate blotches on the side; body mottled; fins also mottled; the anal dark, with a pale edge. Some specimens highly variegated, the caudal banded and with black and white spots; pale streaks from the eye across the cheek; dark bars on sides, extending on dorsal. Most specimens have the region above anal with numerous round whitish spots and some dark ones. These spots sometimes nearly obsolete, most evident on the paler specimens.

The following notes are taken from a specimen, No. 4635, M. C. Z., from Para, Brazil (Coll. Agassiz and Bourgeot): Head 4; depth 4½. D. XII, 14; A. I, 16. Maxillary to front of eye, about equal to eye. Gill membranes free. Lower jaw with a very small canine, not twice the length of the upper teeth. Orbital cirrus quite small; a row of cirri along the nape, longer than the orbital cirrus. Head not very blunt, the anterior profile forming an angle above eye, thence straight and steep. Dorsal spines rather low and flexible, the fin scarcely notched. Color nearly lost; dark marblings on sides and on dorsal fin. This species is evidently the *Blennius crinitus* of Günther and the *B. asterias* of Goode & Bean, probably the *nuchifilis* of Cuvier and Valenciennes, and in all probability the *cristatus* of Linnæus, also. These nominal species are from various localities in the Atlantic. If our specimens are all alike, all these forms most likely belong to 1 species. For this *cristatus* is the oldest name. The very small canines show considerable divergence from the type of *Blennius*, approaching *Scartella*. (Eu.) (*cristatus*, crested.)

Blennius crista setacea longitudinale inter oculos, GRONOW, Museum, I, No. 75; D. 26; A. 16; locality unknown. (Coll. Vosmaer.)

Blennius cristatus, LINNÆUS, Syst. Nat., Ed. X, I, 256, 1758, Indies, after GRONOW; GÜNTHER, Cat. Fish., III, 223, 1861; JORDAN, Proc. U. S. Nat. Mus. 1890, 329.

Blennius crinitus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 237, 1836, La Rochelle, France (Coll. D'Orbigny); GÜNTHER, Cat., III, 224, 1861.

Blennius nuchifilis, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 253, 1836, Isle of Ascension. (Coll. Quoy & Gaimard.)

Blennius asterias, GOODE & BEAN, Proc. U. S. Nat. Mus. 1882, 416, Garden Key, Florida (Type, No. 2620. Coll. G. Würdemann); Garden Key, Florida (Type, No. 2625. Coll. Dr. Whitehurst); Tortugas (Type, No. 6596. Coll. Dr. J. B. Holder); JORDAN & GILBERT, Synopsis, 961, 1883.

Adonis cristatus, GRONOW, Cat. Fish., Ed. Gray, 95, 1854.

885. SCARTELLA, Jordan.

Scartella, JORDAN, Proc. U. S. Nat. Mus. 1886, 50 (*microstoma*).

This genus differs from *Blennius* only in the entire absence of the posterior canine. The relations of this genus with such species of *Blennius* as *Blennius cristatus* are very close. It may be that the groups should be reunited, or that several species here referred to *Blennius* should be placed in *Scartella*. (σκάρτης, one who leaps.)

2733. SCARTELLA MICROSTOMA (Poey).

Head 4 in length (5 with caudal); depth $3\frac{5}{6}$ ($4\frac{3}{8}$); eye $3\frac{1}{4}$. D. XI, 14; A. 15 or 16. Body rather stout, compressed posteriorly; head short, the anterior profile straight and very steep, almost vertical from tip of snout to above eye, where a sharp angle is formed with the straight line of the back. Eye large, longer than snout. Mouth moderate, the maxillary reaching to below front of pupil, its length $3\frac{1}{5}$ in head. Teeth uniform; no posterior canines in either jaw. A small tufted or multifid cirrus over each eye, its length less than diameter of pupil; a row of about 3 short, slender cirri along each side of nape. Gill membranes broadly united, free from isthmus. Lateral line extending about to end of pectoral, each pore with a short, simple branch above and below, directed outward and backward; some conspicuous pores radiating from the eye. Dorsal fin low, subcontinuous, the spines rather slender, lower than the soft rays, the middle spines not much higher than the last; longest rays of dorsal about $\frac{1}{2}$ as long as head; caudal free from dorsal and anal, a little shorter than head; anal low; pectorals slightly longer than head; ventrals $1\frac{1}{3}$ in head. The fins are somewhat shriveled, so that the count of the rays is made with difficulty and may not be perfectly exact. Color very dark olive brown, paler below; head and anterior half of body plain, posterior half sprinkled with sharply defined dots of a vivid sky-blue color, becoming white in alcohol; about 6 obscure round darker blotches in a longitudinal series along sides posteriorly; fins dusky olive, mottled with darker, the caudal obscurely barred, the anal with a pale edge; spinous dorsal nearly black. Length $3\frac{1}{4}$ inches. Cuba. Here described from a specimen taken by Dr. Jordan in Havana. We have also the following notes on Poey's type in the museum at Cambridge: D. XI, 15; A. 17. Dorsal and anal free from caudal. Body rather robust, the head blunt. Last tooth in each jaw a shade longer than its neighbor, but not canine-like. Gill membrane free from isthmus. Dorsal spines low, rather stiff, the fin deeply notched. Color much mottled, with some white spots on posterior half of body; a black ocellus behind first dorsal spine; 5 dark bars along back.

The following is Poey's description: Head $4\frac{1}{3}$ in total length with caudal; depth $5\frac{1}{4}$. D. XII, 15; A. I, 17; P. 14. Snout short; profile falling abruptly; mouth small; eye 3 in head, twice interorbital space. Teeth 15 on each side in each jaw. Gill membrane broadly united, free from isthmus. Lat-

eral line disappearing on middle of back; a row of 6 filaments arranged in pairs on each side of the nape. Membranaceous tentacles over the eye; dorsal somewhat notched, pectoral strongly developed at base. Color brown, with 5 or 6 darker points which form on the back and reach base of the dorsal; pearly spots along sides and some below of the same color; caudal with 3 brown points. Cuba. One specimen, 46 mm. long. (Poey.) (*μικρός*, small; *στόμα*, mouth.)

Blennius microstomus, POEY, Memorias, II, 288, 1861, Cuba. (Coll. Poey.)
Scartella microstoma, JORDAN, Proc. U. S. Nat. Mus. 1886, 50.

886. HYPLEUROCHILUS, Gill.

Hypleurochilus, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 168 (*geminatus*).

This genus differs from *Blennius* in the restriction of the gill-openings to the sides, the gill-membranes being broadly and fully joined to the isthmus; canines well developed. (*υ*, upsilon; *πλεῦρον*, side; *χεῖλος*, lip; in allusion to the V-shaped lateral lips.)

2734. HYPLEUROCHILUS GEMINATUS (Wood).

Head 3½ to 3¾; depth 3½ to 4. D. XI, 15 to XIII, 14; A. II, 18. Head not very blunt, the anterior profile straight, oblique; male (*multifilis*) with the supraocular cirrus very large, each with 4 smaller ones at base; supraocular cirrus in female (*geminatus*) low, shorter than eye, branched at tip; interorbital space concave, not ½ diameter of eye; a slight transverse groove behind eye; canines in both jaws, very strong, hooked backward, the lower considerably stronger than upper; gill openings extending downward to opposite or slightly below lower edge of pectoral. Dorsal fin not emarginate, the spines slender, but rather stiff, lower than the soft rays; pectorals shortish, ventrals rather long. Olive brown, faintly barred with darker; sides plain, or with several pairs of spots of a reddish-brown color, arranged pretty regularly in a double row; vertical fins edged with darker, especially the anal; dorsal black in front. Length 2½ inches. South Atlantic and Gulf coast of the United States, in shallow water; abundant in empty shells and clusters of tunicates. The sexes quite unlike, the male (*multifilis*) distinguished by the high suborbital crest. (*geminatus*, twin.)

Blennius geminatus, WOOD. Journ. Ac. Nat. Sci. Phila., IV, 1824, 278, Charleston, South Carolina, female (Coll. Prof. Bache); CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 265, 1836.
Blennius multifilis, GIRARD, Proc. Ac. Nat. Sci. Phila. 1858, 169, St. Josephs Island, Texas, male (Coll. Gustav Würdemann); GIRARD, U. S. and Mex. Bound. Surv., Zool., pl. 12, fig. 6, 27, 1859; GÜNTHER, Cat., III, 562, 1861
Hypleurochilus multifilis, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 168; JORDAN & GILBERT, Synopsis, 758, 1883.
Hypleurochilus geminatus, JORDAN & GILBERT, Synopsis, 759, 1883.
?*Blennius geminatus*, GÜNTHER, Cat., III, 288, 1861.

887. HYPSOBLENNIUS,* Gill.

Hypsoblennius, GILL, Cat. Fish. East Coast U. S., 20, 1861 (*hentz;* no diagnosis).
Isesthes, JORDAN & GILBERT, Synopsis, 757, 1883 (*gentilis*).
Blenniolus, JORDAN & EVERMANN, new subgenus (*brevipinnis*).

This genus differs from *Blennius* in the absence of canine teeth and in the restriction of the gill openings to the sides, the gill membranes being fully united to the isthmus as far upward as the base of the pectorals; ventral with 1 short, strong spine and 3 simple, articulated rays. The known species are American. (*ὕψι*, high; *Blennius.*)

HYPSOBLENNIUS:
a. Dorsal fin continuous, its margin entire or slightly notched.
b. Dorsal rays XI to XIII, 17 to 19. Pacific species.
c. Orbital cirrus multifid; spines of dorsal stiff; sides blotched or freckled. GILBERTI, 2735.
cc. Orbital cirrus simple or fringed.
d. Spines of dorsal slender and flexible; sides with round dark spots; anal rays 21. GENTILIS, 2736.
dd. Spines of dorsal rather stiff; sides with irregular dark cross bands rather than spots; anal rays 19. STRIATUS, 2737.
bb. Dorsal rays XII, 14 or XII, 15. Atlantic species.
e. Orbital cirrus simple, large or small; body everywhere with dark spots; dorsal spines rather low, stiffish. IONTHAS, 2738.
ee. Orbital tentacle forked at tip, long in males; dorsal spines stiff; body spotted. HENTZ, 2739.
BLENNIOLUS, (diminutive form, from *Blennius*):
aa. Dorsal fin deeply notched, very short, its rays XI, 12 or XII, 12; orbital tentacle slender, fringed; a dark lateral shade. BREVIPINNIS, 2740.

Subgenus HYPSOBLENNIUS, Gill.

2735. HYPSOBLENNIUS GILBERTI (Jordan).

Head 4 in length ($4\frac{3}{8}$ with caudal); depth 4 ($4\frac{3}{8}$). D. XII, 19; A. II, 21. Body comparatively robust, deep, and compressed. Head large, rounded, the anterior profile less blunt than in *H. gentilis* and less rounded, nearly straight from tip of snout to above eye, thence again nearly straight to front of dorsal. Length of snout about equal to diameter of eye, $4\frac{1}{4}$ in head. Mouth rather small, terminal, the maxillary reaching to opposite middle of eye, $2\frac{3}{8}$ in head. Teeth subequal, with no trace of posterior canines. Superciliary tentacle large, multifid, much branched from near the base, the principal division $3\frac{3}{8}$ in head. Gill openings larger than in *H. gentilis*, extending downward to the level of lower edge of pectoral, the length of the slit $1\frac{3}{4}$ in head. Lateral line developed beyond the straight part, its posterior portion curved downward. Dorsal fin continuous, with a slight but distinct depression between the spinous and soft parts, the spines somewhat curved, but stiff and strong, the longest spine about $2\frac{1}{8}$ in head; longest soft rays 2 in head. Caudal fin free from dorsal and anal, $1\frac{1}{2}$ in head; ventrals $1\frac{1}{8}$ in head; pectorals about as long as head. Males,

* The recent identification of *Blennius hentz* with *Isesthes punctatus* enables us to understand the undefined genus *Hypsoblennius*, and to substitute it for the later *Isesthes*. Our judgment is opposed to the recognition of such unexplained "typonyms," but we defer to the custom of the American Ornithologists' Union.

as usual in this genus, with the anal spines partly detached, and provided with fleshy tips. Coloration olivaceous, the body and fins everywhere profusely mottled and reticulated with darker; obscure dark shades extending downward from eye across, or partly across, lower side of head; head without distinct spots or other sharply defined markings, except faint streaks radiating from eye; no pale bars on side of head in either sex; some yellowish markings on anterior part of dorsal. Length 5 inches. California, from Point Concepcion southward to Todos Santos or beyond; common among rocks in the kelp; our specimens from Santa Barbara and Point Loma. (Named for Charles Henry Gilbert.)

Isesthes gilberti, JORDAN, Proc. U. S. Nat. Mus. 1882, 349, Santa Barbara, California (Type, Nos. 26916 and 26917. Coll. Jordan & Gilbert); ROSA SMITH, Proc. U. S. Nat. Mus. 1883, 235, specimens from Todos Santos Bay; D. XI or XII, 16 to 21; A. 19 to 21; head $4\frac{1}{8}$; depth $4\frac{1}{8}$.

2736. HYPSOBLENNIUS GENTILIS (Girard).

Head $3\frac{3}{4}$ in length ($4\frac{1}{4}$ with caudal); depth 4 ($4\frac{1}{4}$). D. XIII, 17; A. II, 19. Body rather robust, deep and compressed, the head large, very bluntly and evenly rounded in profile, more obtuse and more evenly curved than in *H. gilberti*, the snout shorter, about equal to eye, $4\frac{1}{2}$ in head. Mouth rather small, terminal, the maxillary reaching to opposite middle of eye, its length 3 in head. Teeth subequal, the hindmost on each side of upper jaw shorter than the others, and a little apart from them but not forming "a small canine," as stated by Girard. Superciliary tentacle long and simple in the male, its edge fringed with short branchlets, its length about 3 in head; tentacles much smaller in the female, where they are scarcely visible. Gill opening extending downward not quite to lower edge of pectoral, its length (vertical) $2\frac{1}{6}$ in head. Lateral line with only the straight anterior portion developed, not curved downward posteriorly. Dorsal fins continuous, with scarcely a trace of emargination between the spinous and soft parts. Dorsal spines comparatively low and flexible, much less strong than in *H. gilberti*, the longest spines 3 in head; longest soft rays $1\frac{3}{4}$. Caudal free from dorsal and anal, $1\frac{3}{4}$ in head; ventrals $1\frac{3}{4}$ in head; pectorals $1\frac{1}{3}$. Coloration in spirits, brown, the whole body closely mottled and blotched with darker brown, so that the light ground color forms, especially anteriorly, light reticulations around darker spots; on the head the dark spots are small and close together, smallest anteriorly, the lower parts of the head being immaculate, extending from the curve of the preopercle downward, across the interopercle and branchiostegals, in a sharply defined white bar (said to be golden yellow in life), edged with black; behind this and parallel with it across subopercle and isthmus is a similar bar, these bars present only in the males; a few pale spots or bars in front of these; back with about 6 dusky cross shades, below each of these is an oblong dark blotch, the anterior placed along the lateral line, altogether forming an interrupted dark stripe; a similar dark stripe near the median line of the body, interrupted by some pale blotches. Fins all blotched and spotted by light and dark colors, but without distinct markings (a bluish spot in front of dorsal in life);

ventrals and anal nearly blackish in males, the base of the anal with a pale streak. Females more distinctly blotched, with a black spot in front of dorsal and white spots on middle of sides; head lacking the pale bars and black spots, but much mottled with brown and whitish; a very distinct blackish blotch on front of spinal dorsal; pectoral and caudal pale, a dark blotch on base of pectoral. Length about 4 inches. Monterey to Cape San Lucas; common southward in rock pools. Here described from specimens from Angel Island, Gulf of California, from Cape San Lucas, and from Monterey and San Diego. (*gentilis,* related.)

Blennius gentilis, GIRARD, Proc. Ac. Nat. Sci. Phil. 1854, 149, Monterey, California. Types, Nos. 690 and 785 (Coll. A. Cassidy; No. 489, Lieut. Trowbridge); GIRARD, Pac. R. R. Surv., x, Fishes, 113, pl. 25a, fig. 4, 1858; GÜNTHER, Cat., III, 217, 1861.

Isesthes gentilis, STEINDACHNER, Ichth. Beiträge, v, 150, 1876; JORDAN, Proc. U. S. Nat. Mus. 1882, 350; JORDAN & GILBERT, Synopsis, 956, 1883; JORDAN, Proc. U. S. Nat. Mus. 1882, 349.

2737. HYPSOBLENNIUS STRIATUS (Steindachner).

Head 4 to $4\frac{1}{3}$; depth $4\frac{2}{5}$ to 5. D. XI or XII, 17; A. 19; P. 15; V. I, 3. Snout steep, and slightly concave in older examples; interorbital narrow, equal to $\frac{1}{2}$ eye; origin of dorsal a little before the edge of preopercle; second and third dorsal spines equal to the distance from tip of snout to edge of preopercle; dorsal and anal free from caudal; pectoral reaching nearly to front of anal. Color yellowish below, sides brownish, irregular dark-brown cross bars on back and sides; toward the caudal are rows of spots, 4 or 5 wider cross bars of dark brown or violet; a dark blotch from the third to the fifth dorsal spine, behind which are irregular longitudinal dark stripes; anal edged with white, behind which runs a violet line; pectoral and caudal spotted; a dark oval spot behind eye; a brown line from first dorsal spine to eye. Panama (Steindachner), where specimens were also taken by Dr. Gilbert, none of these showing posterior canines, although Steindachner notes the presence of a small canine in 1 specimen. (*striatus,* striped.)

Blennius striatus, STEINDACHNER, Ichth. Beiträge, v, 15, 1876, with plates, Panama.
Isesthes striatus, JORDAN & GILBERT, Bull. U. S. Fish Comm. 1882, 111.

2738. HYPSOBLENNIUS IONTHAS (Jordan & Gilbert).

Head $3\frac{3}{5}$ to 4 ($4\frac{1}{8}$ to $4\frac{3}{8}$ in total); depth $3\frac{1}{6}$ to $3\frac{1}{2}$ ($3\frac{4}{5}$ to $4\frac{2}{7}$). D. XII, 13, or XII, 14; A. II, 13, or II, 14. Body rather deep, moderately compressed, the back little elevated. Head short, blunt, but less so than in *H. punctatus;* the profile prominent above the eye, thence descending abruptly but not vertically to the tip of the snout; length of snout $3\frac{1}{5}$ in head. Mouth small, low, its cleft largely anterior, the short maxillary scarcely reaching past the front of the eye, 4 in head. Eyes large, placed high, 5 in head, the interorbital space about $\frac{1}{2}$ their diameter. Female (*ionthas*) with the orbital cirrus low, scarcely larger than nasal cirrus, which is about equal to diameter of pupil. Teeth moderate, equal; no posterior canines. Gill opening extending

downward to a point varying from a little above to a little below middle of base of pectoral, the height of the slit 3 in head. Lateral line not reaching tip of pectoral. Dorsal fin continuous, the spines low and rather stiff, slenderer than in *H. punctatus*, the longest spines a little lower than the soft rays, which are about $1\frac{1}{3}$ in head. Caudal free from anal, slightly connected with dorsal; a little shorter than head; pectoral about as long as head; ventrals shorter than head. Color of female clear olive green, with only traces of darker bars; body everywhere densely freckled with small round blackish spots, smaller than the pupil; on the sides and lower part of head these spots are reduced to close-set dots; 2 dark lines, separated by a golden area, downward from eye; a vertical curved blackish patch behind eye, in front of which is a golden area; vertical fins olive green, dorsal and caudal usually mottled with dusky; paired fins dusky olive; lower parts of head tinged with golden, sometimes with dusky cross bars; cirri green.

The male (*scrutator*) is thus described: Head 4 ($4\frac{3}{8}$ in total); depth $3\frac{3}{4}$ ($4\frac{1}{2}$). D. XII, 14 or 15; A. II, 15 or 16. Body rather deep, compressed, the back not elevated. Head short, very blunt, almost as deep as long, the profile abruptly descending before eye, the snout about $\frac{1}{3}$ length of head. Mouth very small, anterior, the maxillary extending to opposite front of eye, $3\frac{1}{3}$ in head; teeth subequal, without canines. Orbital cirri very long, reaching when depressed about to the front of dorsal, their length more than $\frac{1}{2}$ head in adult, somewhat shorter in young; a short branch near its middle. Nasal barbel minute. Eye large, much broader than the concave interorbital space, about $4\frac{1}{2}$ in head. Lower edge of gill opening a little below middle of base of pectoral, the depth of the slit $2\frac{1}{2}$ to 3 in head. Dorsal fin scarcely emarginate, the spines rather stiff, lower than the soft rays, the longest spine 2 in head. Caudal slightly connected at base with dorsal, $1\frac{1}{3}$ in head; pectoral about as long as head, reaching past front of anal; ventrals $1\frac{2}{3}$ in head. Lateral line extending to base of eighth spine, not to tip of pectoral. Color in life, deep olive green, almost immaculate, or with faint traces of darker vertical bars; a golden blotch behind eye, behind which is a dusky crescent; 2 dark bars downward from eye, separated by a yellowish area; fins all dusky greenish, nearly or quite immaculate; front of spinous dorsal blackish. South Carolina to Texas, in rock pools; numerous specimens, the largest about $2\frac{1}{2}$ inches long, were obtained with hook and line from the wharves at Pensacola. (*ἰονθάς*, freckled.)

Isesthes ionthas, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 299, Pensacola, Florida (Type, No. 30856, U. S. Nat. Mus. Coll. Jordan & Stearns), female; JORDAN & GILBERT, Synopsis, 960, 1883.

Isesthes scrutator,* JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 300, Pensacola (Type, No. 30850. (Coll. Jordan & Stearns); Galveston (Coll. Dr. August Galny); JORDAN & GILBERT, Synopsis, 960, 1883.

* The form called *scrutator* agrees very closely with *Hypsoblennius ionthas* in all respects except the great length of the orbital cirrus and the different coloration of the body. In both the golden blotch and dark crescent behind the eye are distinct, as also the 2 dark bars separated by a yellow one below the eye. Renewed comparison strengthens our impression that *Hypsoblennius scrutator* is the male of *Hypsoblennius ionthas*.

2739. HYPSOBLENNIUS HENTZ (Le Sueur).

Head $3\frac{2}{3}$; depth 3. D. XII, 15; A. 18; pectoral $1\frac{1}{2}$ in head; ventral $1\frac{2}{3}$; gill slit $2\frac{1}{4}$; eye $4\frac{1}{2}$; maxillary $2\frac{2}{3}$. Orbital tentacle very slender, once forked, 3 in head. Body rather deep; head large, obtuse; interorbital space concave, $\frac{1}{2}$ the diameter of orbit; orbital cirrus as long as dorsal spines, bifid at tip, branched below; a minute nasal cirrus; no canines; gill openings extending to about lower fourth of base of pectoral, thus narrower than in most related species. Dorsal fin high, little notched, the soft part highest, the spines stiff, $2\frac{2}{3}$ in head. Tip of each dorsal spine with a filiform, articulated, ray-like appendage. Color in spirits, olivaceous, back and sides of head and body everywhere covered with brown spots, very irregular in size and shape; on posterior part of body the spots are larger, and show a tendency to form vertical bars; cheeks dark; lower side of head with traces of 3 cross bars; spinous dorsal with an elliptical black spot on membrane of first 3 spines; soft dorsal and caudal obscurely barred; anal, ventrals, and lower rays of pectorals dusky; pectorals olivaceous, spotted with brown. Coasts of North and South Carolina, south to Indian River, Florida; locally common. (Named for its collector, Dr. Nicholas Marcellus Hentz, "the father of American Araneology.")

Blennius punctatus, WOOD, Journ. Ac. Nat. Sci. Phila., IV, 1825, 278, Charleston, South Carolina (Coll. Prof. Bache); CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 267, 1836; GÜNTHER, Cat., III, 228, 1861; not *Blennius punctatus*, Fabricius, 1780, which is a *Stichæus*.

Blennius hentzi,* LE SUEUR, Journ. Ac. Nat. Sci. Phila., IV, 1825, 363, Charleston, South Carolina. (Coll. Dr. Hentz.)

Hypsoblennius hentzi, GILL, Cat. Fish. East Coast N. A., 1861 (*nomen nudum*).

Hypleurochilus punctatus, GILL, Cat. Fish. East Coast N. A., 20, 1873.

Isesthes punctatus, JORDAN & GILBERT, Synopsis, 758, 1883; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1883, 616.

Isesthes hentzi, JORDAN & GILBERT, Synopsis, 960, 1883.

Subgenus BLENNIOLUS, Jordan & Evermann.

2740. HYPSOBLENNIUS BREVIPINNIS (Günther).

Head $3\frac{1}{5}$; depth 4. D. XII, 12; A. II, 14; pectoral $1\frac{2}{3}$ in head; ventral $1\frac{2}{4}$; gill slit $2\frac{1}{2}$; eye $3\frac{1}{3}$ in head; snout $2\frac{2}{4}$ in head; maxillary $2\frac{1}{2}$. Orbital tentacle slender, less than eye. Body rather deep, compressed, back not elevated; anterior profile from first dorsal spine to above eye almost horizontal or slightly decurved, from thence to tip of snout abruptly decurved; head large, its width not quite 2 in its length; interorbital space narrow, grooved, about equaling pupil; eyes large, placed high and close together. Mouth small, low, the maxillary reaching to pupil; teeth subequal, pectinate; no canines; dorsal fin continuous, deeply emarginate, the spines lower than the soft rays, the longest spine about $2\frac{1}{2}$ in head; caudal free

* The following is the substance of the account of "*Blennius hentzi*:" Depth $3\frac{1}{3}$ (in total). D. XI, 14; A. 16. Body little elongate; snout very short, but not vertically truncate; eyes above angle of mouth, placed high; gill slit extending from level of base of pectoral fin to height of eye; teeth equal; dorsal slightly depressed in the middle; pectorals large; a short cirrus above each eye and a smaller one over each nostril. Light bluish ash, mixed with rufous, with numerous irregular black and rufous spots; dorsal black, with whitish spots; soft dorsal with 5 dark bands; ventrals blackish, with pale bands; caudal with 3 or 4 dark bands. Charleston Harbor, South Carolina. (Le Sueur.)

from anal and dorsal; lateral line not reaching soft dorsal. Olive brown, lighter below; back and upper half of sides irregularly marked with about 6 distinct dark-brown cross bars, these uniting at their lower edges and forming a continuous line from head to base of caudal; the bars nearly confluent on the back at base of dorsal fin; a dark lateral band nearly as wide as eye from opercle to base of caudal, containing 5 or 6 light-yellowish spots corresponding to the pale interspaces along the back; fins dusky, anal margined with black; head with a dark spot behind each eye, and 2 smaller blotches in the median line, 1 immediately behind the eyes, the other a short distance in front of dorsal. Pacific coast of Mexico, from Mazatlan to Panama; rather common. The specimens here described from Mazatlan. (*brevis*, short; *pinna*, a fin.)

Blennius brevipinnis, GÜNTHER, Cat. Fish., III, 226, 1861, Pacific coast Central America (Coll. Capt. John M. Dow); one specimen wrongly attributed to Hawaiian Islands.

888. CHASMODES, Cuvier & Valenciennes.

Chasmodes, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 295, 1836 (*bosquianus*).
Blenitrachus, SWAINSON, Class'n Fishes, etc., II, 78, 274, 1839 (*quadrifasciatus*).

Body oblong, compressed, naked; head triangular in profile, the snout somewhat pointed; mouth large, with lateral cleft, the maxillary usually, but not always, extending to beyond eye; premaxillaries not protractile; teeth in a single series, long and slender, comb-like, confined to the front of each jaw; no canines; cirri very small or wanting; gill openings very small, their lower edge above the middle of the base of the pectorals; lateral line incomplete. Fins as in *Blennius*. American. The species with smaller mouth approach *Hypsoblennius*, which genus is not far separated from *Chasmodes*. (χασμώδης, yawning.)

a. Dorsal and anal free from caudal.
b. Anal rays 18 or 19; body not banded. JENKINSI, 2741.
bb. Anal rays 15; body with 4 dark cross bands. QUADRIFASCIATUS, 2742.
aa. Dorsal joined to base of caudal.
c. Mouth moderate, the maxillary not extending to posterior border of eye, 2⅓ in head. SABURRÆ, 2743.
cc. Mouth large, maxillary reaching posterior border of eye. NOVEMLINEATUS, 2744.
ccc. Mouth very large, the maxillary extending to beyond eye. BOSQUIANUS, 2745.

2741. CHASMODES JENKINSI (Jordan & Evermann).

Head 3⅓ (4 in total); depth 4 (5). D. XII, 17; A. 18 or 19; eye 4 to 5 in head. Body more robust than in related species, resembling *Hypsoblennius;* head large, gently rounded in profile, the snout steep, 4 in head; interorbital space narrow, grooved; orbital tentacle (male) much as in *Hypsoblennius gilberti*, about 3 in head, branched, the branches usually 4; mouth much larger than in *Hypsoblennius*, the maxillary 2¾ to 3 in head, reaching to below posterior margin of eye; teeth even, comb-like; gill opening 2 in head, extending downward nearly to lower edge of pectoral, much larger than in *Chasmodes saburræ*. Dorsal little notched, the spines

slender, 2½ in head, the rays a little higher; anal lower, the rays 3½ to 4 in head; pectorals reaching anal, 1⅓ in head; ventrals 2¼; dorsal and anal free from caudal. Color in life, according to Evermann & Jenkins, yellowish; 5 quadrate spots of darker extending from dorsal to a line drawn from middle of eye to lower base of caudal, the anterior one above tip of pectoral; median line of side with a more or less distinct series of small spots; a short dark vertical line behind the eye; a dark blotch in front of origin of dorsal fin and another on humeral region; underside of head with 2 ill-defined dark bands; dorsal fin more or less speckled with black, the anal with a narrow white border above which is a broader band of deep brown. Six specimens, the largest about 3 inches long, were obtained at Guaymas, Sonora, by Drs. Evermann & Jenkins, in 1887. One of these, (No. 412, L. S. Jr. Univ. Mus.), examined by us, is the type of the present description. The large mouth distinguishes this species at once from *Hypsoblennius striatus*, with which it has been identified. The species is intermediate between typical *Chasmodes* and *Hypsoblennius*, and its discovery may make it necessary to merge the latter in *Chasmodes*. (Named for Dr. Oliver Peebles Jenkins.)

Hypsoblennius striatus, Evermann & Jenkins, Proc. U. S. Nat. Mus. 1891, 163; not of Steindachner.

Chasmodes jenkinsi, Jordan & Evermann, Proc. Cal. Ac. Sci. 1896, 232, pl. 39, Guaymas. (Coll. Evermann & Jenkins.)

2742. CHASMODES QUADRIFASCIATUS (Wood).

D. 27; A. 15. Form of *Chasmodes bosquianus:* Lower jaw slightly longer than the upper. Dorsal and anal free from caudal; anal fin highest anteriorly. Body with 4 distinct brownish bands, a fifth broader and less marked on the neck; 4 round yellowish spots along base of anal; head spotted with blackish. (Wood.) *Habitat* uncertain, probably South Atlantic coast of the United States; not recognized by recent collectors; very likely based on the female of *C. bosquianus*, with the caudal torn from the other vertical fins. (*quadri-*, four; *fasciatus*, banded).

Pholis quadrifasciatus, Wood, Journ. Ac. Nat. Sci. Phila., iv, 1825, 282; locality unknown, probably South Carolina. (Coll. Rubens Peale.)

Chasmodes quadrifasciatus, Cuvier & Valenciennes, Hist. Nat. Poiss., xi, 298, 1836; Günther, Cat. Fish., iii, 229, 1861; Jordan & Gilbert, Synopsis, 757, 1883.

2743. CHASMODES SABURRÆ, Jordan & Gilbert.

Head 3½ to 3⅔; depth 3½ to 3¾. D. XI or XII, 17 to 19; A. II, 18 or 19. Body rather deep and compressed, less elongate than in *C. bosquianus;* the back somewhat arched. Head comparatively short, much shorter than in *C. bosquianus*, not ¼ longer than deep; profile forming a nearly even curve from the base of the dorsal to the tip of the snout; mouth notably smaller than in *C. bosquianus;* maxillary not reaching posterior margin of eye, its length 2⅓ in head; teeth occupying about ½ of lower jaw; height of gill slit 3⅔ in head, its lower ray opposite third ray of pectoral. A minute cirrus, shorter than pupil, above each eye and each nostril. Dorsal con-

tinuous, with slender rays, the last one joined to the caudal. First two rays of anal short, thick, and fleshy in the males. Male deep olive, with dark cross shades; numerous pale spots on the sides which form undulating lines converging backwards; dark stripes downward and forward from eye; top of head and upper part of dorsal fin usually with fine black spots; spinous dorsal with a median orange longitudinal band; other fins mostly dusky olive. Some specimens with the outer part of both dorsals and the top of head dusted with black spots, others with these spots obsolete; soft dorsal and caudal light orange, barred with light greenish; anal dull orange, with an obscure blackish median band, the exserted tips of the rays abruptly whitish; pectorals dusky olive, strongly tinged with orange; ventrals blackish, orange at tip. Female with about 8 blackish cross bands extending on the dorsal fin; the body everywhere with pale spots; fins all sharply barred with blackish and olive. Pensacola Bay, Florida; common about the wharves and ballast rocks in shallow water; taken with seines and pinhooks. Allied to *Chasmodes bosquianus*, but with the mouth smaller, the form less elongate. (*saburra*, ballast.)

Chasmodes saburræ, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 298, Pensacola, Florida (Type, No. 30824. Coll. Jordan & Stearns); JORDAN & GILBERT, Synopsis, 958, 1883.

2744. CHASMODES NOVEMLINEATUS (Wood).

Head $3\frac{2}{5}$; depth $3\frac{2}{5}$; eye $4\frac{1}{4}$; snout $3\frac{1}{4}$; maxillary reaching posterior border of eye. D. XI, 18; A, III, 17. Head and shoulders heavy, the body lance-shaped, tapering gradually to tail; snout short, blunt, profile nearly vertical to eye, thence gently rounded; mouth rather large, somewhat oblique, the maxillary reaching posterior border of eye; dorsal and anal high, longest dorsal rays 2 in head; anal considerably lower; pectoral nearly as long as head; ventrals $1\frac{1}{4}$ in head. Color, side with 6 broad, dark, vertical bars, the anterior 4 extending on the dorsal fin, these bars separated by irregular narrow pale spaces; entire side profusely covered with small white spots; a small black spot at base of caudal; head mottled with light and dark; 2 small dark spots on under side of lower jaw; just behind these and extending downward from the angles of the mouth are 2 other larger, blacker spots, while behind these, extending downward and backward from middle of cheek, is an irregular black line; whole head with numerous fine dark punctulations; dorsal and anal variously spotted or barred with light and dark; spinous dorsal with a large dark area at top of anterior spines; caudal faintly barred; pectorals and ventrals more plainly barred. Length 2 inches. South Atlantic coast of the United States, South Carolina to Florida; abundant in Indian River, Florida, where numerous specimens were taken in January, 1896, by Evermann & Bean. (*novem*, nine; *lineatus*, lined.)

Pholis novemlineatus, WOOD,* Journ. Ac. Nat. Sci. Phila., IV, 1825, 280, Charleston Harbor, South Carolina.

*The following is the substance of Wood's original description of this species: "Body with 9 whitish longitudinal bands; dorsal fin with an irregular blackish spot between the first and second rays; remainder of the fin clouded with dusky brown. Head descending somewhat abruptly, tuberculated anteriorly; nostrils with a small appendage; head, lips,

Chasmodes saburræ, EVERMANN & BEAN, Fishes of Indian River, Florida, in Rept. U. S. Fish Comm. 1896, 247; not of JORDAN & GILBERT.
Chasmodes novemlineatus, GÜNTHER, Cat., III, 229, 1861.

2745. CHASMODES BOSQUIANUS (Lacépède).

Head 3½; depth 3½. D. XI, 19; A. II, 19. Orbital tentacle very minute or wanting; maxillary extending to rather beyond eye; interocular space very narrow, not concave. Dorsal fin not emarginate, the spines slender. Dorsal joined to base of caudal; anal free. Color (in male) olive green, with about 9 horizontal narrow blue lines, these somewhat irregular and interrupted, converging backward; opercular membrane and a broad stripe through middle of spinous dorsal deep orange yellow; anal fin dark, the rays with white membranaceous tips; female dark olive green, reticulated with narrow, pale green lines, and with several broad dark bars, which are more distinct posteriorly; vertical fins similarly marked; head finely dotted with black; a dusky spot at base of caudal in both sexes. New York to Florida; common southward in shallow water. (Named for M. Bosc, who collected at Charleston for Lacépède.)

Blennius bosquianus, LACÉPÈDE, Nat. Hist. Poiss., II, 493, 1800 (female), South Carolina. (Coll. Bosc.)
?Pholis quadrifasciatus, WOOD, Journ. Ac. Nat. Sci. Phila., IV, 1824, 282, locality unknown, probably South Carolina. (Coll. Rubens Peale.)
Chasmodes boscianus, GÜNTHER, Cat., III, 229, 1861.
Chasmodes bosquianus, JORDAN & GILBERT, Synopsis, 756, 1883.

889. HOMESTHES, Gilbert, new genus.

Homesthes, GILBERT, new genus (*caulopus*).

Differing from *Hypsoblennius* chiefly in the presence of 4 articulated ventral rays instead of 3, as usual in *Blenniinæ*. We have examined the ventrals of *Hypsoblennius striatus*, *punctatus*, *ionthas*, *gentilis*, and *gilberti*, and have found them to consist constantly of 1 short, strong spine and 3 simple articulated rays. In *Homesthes caulopus* there is 1 strong, short spine and 4 well-developed simple jointed rays. (ὁμός, uniform; ἐσθίω, to eat.)

2746. HOMESTHES CAULOPUS, Gilbert, new species.

Head 3⅔ in length; depth at base of ventrals 4, at middle of abdomen 3⅔; least depth of caudal peduncle ⅓ length of head; snout 4; eye 4 to 4⅕. D. XII, 15 or 16; A. II, 17; P. 14. Longest dorsal spine 2⅔; last dorsal

opercula, etc., and base of the pectoral fins, finely spotted with bluish black, the spots being larger on the front and opercula; branchial opening extremely small, extending ⅓ of the length of the external curve of the operculum; mouth descending little; gape moderate; sides of the head fleshy; body compressed; rib spaces evident; sides with 9 longitudinal whitish lines, some of which are interrupted; behind the eye and under the dorsal fin are 2 irregular whitish patches; dorsal fin commencing before the pectoral fins; between the first and second rays is an irregular blackish spot, several of the following rays are also spotted, the color of the spots becoming lighter as they recede toward the tail, where they mingle with the dusky color of the fin and are lost; fin rising posteriorly, and joining the caudal fin at about ⅓ the distance from its extremity; anal fin commencing under the termination of the pectoral fin, and extending nearly to the tail; caudal fin rounded; ventral fins 2-rayed; pectoral fins rather large, the base thick and fleshy, finely spotted with bluish black; anus small, tubercle small; color brownish, fins dusky. D. 30; C. 12⅔; A. 20; V. 2; P. 13. Length 3¼ inches; depth, exclusive of the dorsal fin, hardly 1 inch."

spine $3\frac{1}{3}$; longest (tenth) dorsal ray 2; longest (fifteenth) anal ray $2\frac{1}{2}$; ventrals $1\frac{5}{6}$; longest pectoral ray $1\frac{3}{4}$ to $1\frac{2}{3}$; caudal $1\frac{1}{2}$. Robust, moderately compressed, with wide heavy head and short, bluntly rounded snout, the anterior profile of which is nearly vertical. In shape and general appearance much resembling *Hypsoblennius gilberti*. Mouth very wide, horizontal, short, the maxillaries reaching vertical from hinder edge of pupil, 3 to $3\frac{1}{3}$ in head. Teeth, as usual in this group, the posterior not enlarged or canine-like. Nostrils with slightly elevated margins, scarcely tubular, the hinder edge of anterior nostril produced into a conspicuous laciniate flap, about $\frac{2}{3}$ as long as the diameter of orbit. A similar but larger orbital cirrus, divided nearly to the base into 6 or 8 slender filaments. Interorbital space deeply grooved, without median ridge, opening posteriorly into the deep transverse groove which separates the orbital region from the somewhat swollen occiput, its width $1\frac{1}{3}$ eye. The mucous canals of head give off transverse branches which open by numerous pores. These thickly beset the snout, subocular region, top of head, preopercle, and upper portion of opercle. Width of gill slit equaling or slightly exceeding $\frac{1}{2}$ length of head, confined to area above lower base of pectorals. First dorsal spine over margin of preopercle; spinous dorsal low, of nearly uniform height, much lower than second dorsal, the spines rather strong at base, with weak reflexed tips; membrane of last dorsal ray joined to extreme base of rudimentary caudal rays; anal low, rising slightly posteriorly, leaving a short free interval between its last ray and caudal. Lateral line strongly developed anteriorly for a distance equaling length of head; from that point it is only faintly visible, declining abruptly to middle of sides, along which it may be traced to base of caudal; the anterior portion gives off numerous pairs of short transverse lines, each of which ends in a pore; no pores or lines are visible posteriorly. Blackish, without sharp markings, the sides with irregular light blotches, some of which are subcircular in outline and contain 1 or more black central specks; the light markings near the back elongate and vertically placed, faintly outlining dark bars of the ground color; a vertical black blotch on cheek behind eye; lower parts lighter; no distinct bars on head; fins all blackish, the anal, the ventrals, the lower caudal and pectoral rays deeper black; anal and caudal margined with white, some of the dorsal rays narrowly tipped with white; tentacles whitish. Two specimens, 4 and $4\frac{1}{2}$ inches long, from Panama Bay. (Gilbert.) (*καυλός*, stem; *πούς*, foot.)

Homesthes caulopus, GILBERT MS., Fishes of Panama, Panama. (Coll. Gilbert. Type, No. 5623, L. S. Jr. Univ. Mus.)

890. SCARTICHTHYS, Jordan & Evermann, new genus.

Scartes, JORDAN & EVERMANN, Check-List Fishes, 471, 1896 (*rubropunctatus*); preoccupied by *Scartes*, Swainson, a genus of mammals.
Scartichthys, JORDAN & EVERMANN, new genus (*rubripunctatus*).

Body elongate, slowly declining to the caudal. Head obliquely compressed, oblong, the profile more or less vertical. Eyes lateral, closely approximated, situated at the angle of the profile with the postocular

region. Gill apertures continuous under the throat, gill membrane free from isthmus. Branchiostegals 6. Mouth moderate, the contour of the upper jaw semicircular; upper jaw protruding beyond the lower; lips moderate, uniform and free, concealing the teeth. Teeth labial and movable, very slender and recurved, contiguous and uniserial; no posterior canines. Dorsal fin divided; anal similar to soft dorsal; caudal obtusely rounded; pectorals moderate, angularly rounded; ventrals approximated, each with 3 simple rays, the internal of which is smallest. This genus is very close to the Old World genus, *Salarias*,* Cuvier, which differs in having the dorsal fin continuous, as in *Rupiscartes*. (σκάρτης, one who leaps; ἰχθύς, fish.)

2747. SCARTICHTHYS RUBROPUNCTATUS (Cuvier & Valenciennes).

Head 4; depth 4 (5 with caudal); D. XI–16; A. 20; eye $4\frac{1}{3}$ in head; teeth less flexible than in *Rupiscartes atlanticus;* no canine teeth; the forehead not projecting beyond the mouth; a very small tentacle on the neck, a longer fringed one above the orbit; dorsal fin deeply notched, not extending on to the caudal. Color brown, marbled with black, and dotted with reddish; a black spot on the anterior part of the dorsal; throat with 2 or 3 brownish cross bands; a jet-black spot behind eye, with a narrow edge posteriorly. (Günther.) Coast of Peru and Chile, north to Panama. Specimens examined by us collected by Prof. Frank H. Bradley at Pearl Islands, near Panama, and at Callao. Length 3 inches. (*ruber*, red; *punctatus*, spotted.)

Salarias rubropunctatus, Cuvier & Valenciennes, Hist. Nat. Poiss., xi, 348, 1836, Juan Fernandez (Coll. Claude Gay); Günther, Cat., iii, 249, 1861; Jordan & Gilbert, Proc. U. S. Nat. Mus. 1882, 628; not of Kner, Novara-Fische, 198.

891. RUPISCARTES, Swainson.

Alticus,† Commerson, in Lacépède, Hist. Nat. Poiss., ii, 458, 1800 (*saliens*).
Alticus,‡ Cuvier & Valenciennes, Hist. Nat. Poiss., xi, 337, 1836 (*alticus*).
Rupiscartes, Swainson, Nat. Hist. Class'n Anim., ii, 275, 1839 (*alticus*).

This genus, as here understood, differs from *Salarias* only in the presence in 1 or both jaws of posterior canines. Dorsal fin continuous, without deep notch. Vertebræ $12+22=34$ (*atlanticus*). (*rupes*, rock; σκάρτης, one who leaps; *Rupiscartes tridactylus* (*alticus*), "said to jump on the sea rocks like a lizard." Swainson.)

* *Salarias*, Cuvier, Règne Anim., Ed. 2, ii, 175, 1829 (*quadripinnis*). *Erpichthys*, Swainson, Nat. Hist. Class'n. Anim. ii, 275, 1839 (*quadripinnis*, etc). (σαλάρια, a modern Greek name of *Blennius basilicus*.)

† We do not think that the name *Alticus* can be substituted for *Rupiscartes*, because Lacépède does not adopt this genus of Commerson, but merges it in *Blennius*, quoting Commerson's account as a footnote. This is as follows: "*Alticus saltatorius*, pinna spuria in capitis vertice; seu pinnula longitudinali pone oculos cartilaginea; seu alticus desultor, occipite cristato, ore circulare deorsum patulo." Apparently this quotation of a generic description not approved, does not give priority to the latter.

‡ This genus *Alticus* is not adopted by Cuvier & Valenciennes. Valenciennes speaks of "un petit *Salarias* que nous parait être celui-là même sur lequel Commerson avait établi son genre *Alticus*." But a genus is not established until it is accepted by some authority as well as defined.

2748. RUPISCARTES ATLANTICUS (Cuvier & Valenciennes).

Head 4 to $4\frac{1}{2}$; depth $3\frac{1}{3}$ to $3\frac{2}{3}$. D. XII, 21 or XIII, 20; A. 24 or 25; vertebræ $12+22=34$; eye 4 to $4\frac{1}{2}$ in head. Body rather high, compressed. Head short, very blunt, its width about 2 in the length; anterior profile from first dorsal spine to above eye straight or slightly convex; from thence to tip of snout abruptly decurved, in some specimens nearly vertical. Mouth inferior, lower jaw included; maxillary about reaching posterior border of eye. Teeth small, pectinate, the lower canines exceedingly large and entering the cavity in the palate. Supraorbital tentacle well developed, slender; a group of 5 or 6 short tentacles on either side of head in front of nostrils and on either side of neck, these shorter than pupil. Dorsal fin not emarginate, extending from a point above middle of operculum to base of caudal; anal lower than soft dorsal, $1\frac{1}{2}$ to 2 in dorsal; pectorals reaching past vent, about equaling head; ventrals about 2 in head. "The intestinal tract is more than 3 times as long as the entire body. The structure of the skeleton is very similar to that of the Blennies; the jaw bones, however, are still shorter, and the intermaxillary and mandibulary are deeply concave anteriorly. There are 12 abdominal and 22 caudal vertebræ, the former portion being only $\frac{1}{2}$ as long as the caudal." (Günther.) Some specimens, apparently males, with the anterior profile vertical and very high; fins high; caudal lanceolate, the black median rays much exceeding the outer pale ones. Females with the anterior profile a nearly even curve, the caudal lunate, its median black rays shorter than the outer pale ones. Body liver brown, paler below, with usually 5 or 6 darker cross bars extending on the dorsal; a black spot behind eye in all; fins mostly blackish, an orange area on upper edge of caudal; a yellow one tinged reddish below; eye red posteriorly. Length 6 to 8 inches. Tropical America, on both coasts, very abundant in rock pools, north to West Indies and to Todos Santos. Here described from specimens from Mazatlan.

Punaru, Marcgrave, Hist. Brazil, 165, 1648, Brazil.
Salarias atlanticus, Cuvier & Valenciennes, Hist. Nat. Poiss., xi, 321, 1836, Madeira (Coll. Richardson) Antilles (Coll. Plée); Günther, Cat. Fish., iii, 242, 1861.
Rupiscartes atlanticus, Jordan, Proc. U. S. Nat. Mus. 1888, 333.

892. ENTOMACRODUS,* Gill.

Salarias, Swainson, Nat. Hist. Class'n Fishes, ii, 274, 1839 (*vermicularis;* not of Cuvier).
Entomacrodus, Gill, Proc. Ac. Nat. Sci. Phila. 1859, 168 (*nigricans*).

This genus has large posterior canines as in *Rupiscartes*, but the dorsal fin is divided into 2 fins as in *Scartichthys*. (ἐν, in; τόμος, cutting; ἀκρός, sharp; ὀδούς, tooth.)

a. Orbital cirrus present; dorsal rays XII or XIII–15; canines small.
b. Cirrus above eye divided; anal rays 15. CHIOSTICTUS, 2749.
bb. Cirrus above eye simple or nearly so; anal rays 18; body with pearly spots. MARGARITACEUS, 2750.

* This genus is equivalent to *Salarias* of Swainson, but the generic name *Salarias* was based on *Salarias quadripinnis*, before either of the species referred to it by Swainson was made known.

aa. Orbital cirrus wanting; no cirri at nape.

c. Dorsal rays XII–19; anal 15; body rather slender, the depth about 5 in length; body with bands and spots. DECORATUS, 2751.

cc. Dorsal rays XI–15; anal 17; body very slender, the depth about 6 in length; color blackish, nearly plain. NIGRICANS, 2752.

2749. ENTOMACRODUS CHIOSTICTUS (Jordan & Gilbert).

Head 4½ in length; depth 5¼. D. XII–15; A. 15; eye 3⅓ to 4⅔ in head, varying with age. Body moderately elongate, compressed, the head short, blunt, almost globular, about as broad as deep, and a little longer than broad. Mouth inferior, with little lateral cleft, the lower jaw included; width of cleft of mouth ⅔ length of head. Teeth small, weak, finely pectinate; canine teeth small, not so long as diameter of pupil. Supraorbital cirrus divided into 4, its height ⅔ that of eye; a few minute slips at the nape. Interorbital space channeled, narrower than eye. Maxillary extending to behind middle of eye. No crest on top of head. First dorsal low and even, its spines rather slender, the last spines short, scarcely connected by membrane with the soft rays; soft dorsal well separated from caudal; caudal subtruncate, with rounded angles; anal lower than soft dorsal, with a little longer base; pectorals a little longer than head; ventrals about ⅓ as long. Color in life, olive brown above, lighter below; 5 broad, dark bars from dorsal fin to middle of sides, each terminating above on the fin, and below on sides in a pair of black spots; sometimes only the dark spots are distinguishable, the bars being obscure; sides below spinous dorsal with numerous black specks, and with numerous oblong spots of bright silvery; sometimes a silvery streak from upper portion of base of pectorals to base of caudal; a broad salmon-colored streak on each side of ventral line; sometimes the space between the silvery lateral band and the base of the anal is darker, the vertical bars again appearing as pairs of black, vertical blotches; head yellowish olive, darker above, and reticulating with narrow brown lines, these appearing as parallel bars on the upper lip, and radiating from the median line on the upper side of the head; vertical fins light grayish, with black spots, which appear as wavy bars on the caudal fin; pectorals and ventrals pale, the former with a yellowish shade at base; orbital tentacles bright red. Pacific coast of Mexico. Known from 4 specimens (the largest 2¼ inches in length), taken in a deep rock pool at Mazatlan. Two others taken by the *Albatross* from Clarion Island. (χιών, snow; στικτός, spotted.)

Salarias chiostictus, JORDAN & GILBERT, Synopsis, 363, 1883, Mazatlan, Mexico. (Coll. Jordan & Gilbert.)

2750. ENTOMACRODUS MARGARITACEUS (Poey).

Head 5 in total length with caudal; depth 6½. D. XII–14; A. I, 14; eye 4⅓ in head, well forward. Body large, snout abruptly decurved; mouth very low, maxillary reaching anterior nostril, which has a little tentacle; (canines small); small tentacle over eye; gill membranes broadly connected, free from isthmus; dorsal deeply emarginate, almost divided; anal beginning under middle of body without caudal, and anal papillæ and caudal

rounded; ventrals short; lateral line present anteriorly, no tentacles on nape. Color brown, with 2 vertical bands of a dusky silvery; a central point in each band shining bright. One specimen, 2½ inches long. Cuba. (Poey.) Perhaps a *Salarias*.

We have the following notes on a specimen, possibly the type of this species, sent by Poey to the museum at Cambridge: Head 4⅜; depth 5. D. XII–15; A. 18. Body slender. Interorbital concave. Head short, blunt, almost round; a small cirrus over the eye, none on nape. Canines present, small. Body with about 6 dark cross bars besides pearly spots and various markings. Dorsal divided nearly to base. Closely resembles *Salariichthys textilis*. (*margarita*, μαργαρίτης, pearl.)

Salarias margaritaceus, Poey, Memorias, II, 289, 1861, Cuba. (Coll. Poey.)

2751. ENTOMACRODUS DECORATUS, Poey.

Head 5 in total length with caudal; depth 5. D. XII–19; A. 15; P. 14. Eye very high; anterior nostril prolonged in a tube; nape following a straight line to the posterior nostril, profile thin, following a straight and oblique line to mouth, which is very low and short, the maxillary reaching posterior nostril. Lower jaw shorter. Teeth movable, numerous, incurved, close set, in 1 row. (Canines not described.) No cilia on head. Dorsals of equal length, the soft rays more elevated; anal similar to second dorsal; pectoral broad, its lower rays thickened; caudal rounded. Color brownish yellow; the body with darker cross bands, which begin below the middle of the first dorsal, alternating with narrower spaces of the ground color; along the middle and edges of the bands vertical rows of sky-blue spots; in the pale interspaces below the lateral line, which is much curved, a white spot; 3 pale spots placed obliquely below the eye; rays of dorsal and caudal dotted with black. One specimen, 2 inches long. Cuba. (Poey.) Not seen by us; perhaps a *Salarias*. (*decoratus*, decorated.)

Entomacrodus decoratus, Poey, Synopsis, 398, 1868, Cuba. (Coll. Poey.)

2752. ENTOMACRODUS NIGRICANS, Gill.

The elongated body, from the snout to the end of the caudal fin, is between 7 and 8 times longer than it is high at the pectorals. Its height at the caudal is about ⅓ of the same length. The head is subquadrate, and forms ⅔ of the total length. Its greatest height equals ⅔ of its length. Its sides decline obliquely outward and downward. The first dorsal commences near the nape, and 2 of its rays are in advance of the pectorals. The second dorsal commences immediately behind the first, and nearly over the fourth ray of the anal, it ceases some distance from the base of the caudal. The anal is more uniform in height than the dorsal, and ceases before it does. The caudal forms less than ⅕ of the total length. D. XI–15; A. 17; P. 15; V. 3. The general color of the body and fins is blackish. West Indies. A single specimen was caught in shallow water, at the island of Barbados, near Bridgetown. (Gill.) Not seen by us. (*nigricans*, blackish.)

Entomacrodus nigricans, Gill, Proc. Ac. Nat. Sci. Phila. 1859, 168, Barbados. (Coll. Dr. Gill.)

893. SALARIICHTHYS, Guichenot.

Salariichthys, Guichenot, Mém. Soc. Sci. Nat. Cherbourg, xiii, 1867, 96 (*textilis*).

This genus differs from *Entomacrodus* in the presence of teeth on the vomer; dorsal deeply notched; cirri present over eye and on nape; posterior canines small. (*Salarias; ἰχθύς*, fish.)

2753. SALARIICHTHYS TEXTILIS (Quoy & Gaimard).

D. XII, 16; A. 18. A few bluntish teeth on vomer; tentacles very small, fringed over nostril and eye, simple on neck; canines quite short; depth $4\frac{3}{5}$; head $4\frac{3}{5}$; pectoral short, little longer than head; gill membranes broadly united, free from isthmus; dorsal notched almost to base, free from caudal; orbital filament $\frac{1}{4}$ eye. Olive, with 13 silvery cross streaks, not $\frac{1}{2}$ as wide as the dark interspaces, some of the cross streaks Y-shaped; both dorsals with cross markings, the second with 12 or 13 streaks of dark obliquely upward and backward, alternately with similar pale streaks; cross bars on sides bent in middle, extending up and back and down and back from middle line parallel with muscular impressions; sides with some obscure pale dots; caudal barred with 7 dark bars; anal darkest mesially; lower side of head with dark streaks radiating from the isthmus; bars at chin Y-shaped, upper part of head with darker markings; pectoral nearly plain; a dusky area at base below which is a dusky spot; marblings at base of dorsal. West Indies, from Bermudas to Brazil. Here described from a specimen from Abrolhos Islands (Coll. *Albatross*). This specimen agrees fairly with the account given by Jenyns, but Jenyns describes 5 bars on the tail. It also agrees fairly with the account of the Bermuda specimens given by Goode. It is evidently the *Salarias vomerinus* of Cuvier & Valenciennes, and probably their *textilis* also; but their description of the latter does not apply very well to the coloration of our specimen. (*textilis*, woven.)

Salarias textilis, Quoy & Gaimard MS., Cuvier & Valenciennes, Hist. Nat. Poiss., xi, 307, 1836, Ascension Island (Coll. Quoy & Gaimard); Günther, Cat., iii, 248, 1861; Goode, Bull. U. S. Nat. Mus., v, 29, 1876.

Salarius vomerinus, Cuvier & Valenciennes, Hist. Nat. Poiss., xi, 349, 1836, Bahia. (Coll. Blanchet.)

Salariichthys textilis, Jordan, Proc. U. S. Nat. Mus. 1890, 329.

894. OPHIOBLENNIUS, Gill.

Blennophis, Valenciennes, in Webb & Berthelot, Poiss. Îles Canar., 60, 1844 (*webbii*; not *Blennophis*, Cuvier & Valenciennes, a genus of *Clininæ*).

Ophioblennius, Gill, Proc. Ac. Nat. Sci. Phila. 1860, 103 (*webbii*; substitute for *Blennophis*).

Body oblong, strongly compressed, scaleless; snout short, high, abruptly decurved anteriorly; symphysis of lower jaw of 4 hooked canines, the outer strongest and bent backward, almost forming a right angle; sides of lower jaw with 2 or 3 still larger canines, the hindermost very large and bent backward; upper jaw with 4 slender canines in front, followed by a long row of shorter, slender, movable teeth, which are set close together; nasal tentacle digitate; a low, simple tentacle above eye; gill openings wide. Dorsal fin long, the spines slender, separated by a slight notch from

the soft rays; caudal lunate or forked, free from dorsal and anal; ventrals small, I, 2; lateral line incomplete; pectorals large. A strongly marked genus, perhaps more nearly allied to *Blennius* than to *Emblemaria* or *Chænopsis*. (ὄφις, snake; *Blennius*, in allusion to the fang-like teeth.)

a. D. X, 20; A. 20; depth 5⅓ in length. WEBBII, 2754.
aa. D. XI, 22; A. II, 23; depth 4¼ in length. STEINDACHNERI, 2755.

2754. OPHIOBLENNIUS WEBBII (Valenciennes).

Head 5; depth 5½. D. X, 20; A. 20; P. 16. A slender tentacle above eye in front, and a much broader one, divided into 4 to the base, above the nostril. Snout obtuse, nearly vertical at tip; eye large; 4 teeth at end of upper jaw, strongly pointed, curved backward like hooks; lower jaw with 4 teeth at tip, the two middle ones like upper teeth, the two outer hidden and turned backward; a little recurved tooth on side of lower jaw; caudal fin forked; dorsal somewhat notched at the last spine; lateral line ending near middle of body. Olive green, light or dark; dorsal and anal dusky violet, the base pale; back and sides often with fine points; a dark spot behind eye; the silvery swim bladder showing through sides of belly. (Steindachner). Tropical Atlantic; known only from the Canaries and Barbados; not seen by us. (Named for P. B. Webb, one of the explorers of the Canary Islands.)

Blennophis webbii, Valenciennes, in Webb & Berthelot, Îles Canar., Poiss., 60, pl. 20, f. 1, 1844, Fortaventura, Canary Islands (Coll. Webb); "caught in myriads at Puerto de Cabras in August, eaten as Anchovias" (Webb); Günther, Cat., III, 259, 1861; Steindachner, Ichth. Notizen, VI, 48, 1867.
Ophioblennius webbi, Jordan & Gilbert, Synopsis, 756, 1883.

2755. OPHIOBLENNIUS STEINDACHNERI, Jordan & Evermann, new species.

Head 4 to 4⅕; depth 4 to 4⅓. D. XI, 22; A. II, 23; V. I, 2; P. 15. Head much compressed; eye 3¼ in head; snout 4½. Dorsal beginning above gill opening, ending just before caudal, its soft rays somewhat higher than the spines, the highest spine 1⅔ in head; caudal and pectorals each about as long as head; ventrals 1½ in head. Dark golden brown, sometimes with a broad cross band of dusky violet on back and dorsal fin; caudal with 2 dark longitudinal stripes; dorsal and anal purplish or orange; an intense, round, dark, ocellated spot behind eye. (Steindachner.) West coast of Mexico; not seen by us; recorded from near Mazatlan and the Tres Marias Islands. (Named for Dr. Franz Steindachner.)

Blennophis (Ophioblennius) webbi, Steindachner, Ich. Beitr., VIII, 41, 1879, 5 specimens 70 mm. long, from Navidad near Mazatlan and the Tres Marias Islands.
Ophioblennius steindachneri, Jordan & Evermann, Check-List Fishes N. and M. A., 472, 1896, name only, Tres Marias Islands; after Steindachner.

895. EMBLEMARIA, Jordan & Gilbert.

Emblemaria, Jordan & Gilbert, Proc. U. S. Nat. Mus. 1883, 627 (*nivipes*).

Body slender, not eel-shaped, compressed, scaleless. Ventrals present, jugular, each of 1 spine and 2 soft rays. A single high dorsal fin beginning on the nape and extending to the caudal, with which it is not conflu-

ent; no notch between spinous and soft parts. Head cuboid, compressed, narrowed anteriorly. Symphysis of lower jaw forming a very acute angle. A single series of strong, blunt, conical teeth on each jaw, and on vomer and palatines. Vomer and palatine teeth larger, their series continuous, parallel to the series in upper jaw. No cirri at the nape; sometimes a cirrus on upper part of eyeball. Gill openings very wide, the membranes broadly united below, free from the isthmus. Lateral line obsolete. This genus bears some resemblance to *Blennius*, but the dentition is entirely different, approaching that of *Chænopsis*. Tropical America, in rather deep water. (*Emblema;* ἔμβλημα, a banner.)

a. Eye without cirrus.
  b: Depth 5 in length; dorsal rays 33; ventrals not pure white. ATLANTICA, 2756.
  bb. Depth 7 in length; dorsal rays 37; ventrals pure white. NIVIPES, 2757.
aa. Eye with a long cirrus on eyeball above pupil; ventrals dusky; maxillary not extending beyond eye. OCULOCIRRIS, 2758.

2756. EMBLEMARIA ATLANTICA, Jordan & Evermann, new species.

Head $3\frac{2}{3}$; depth 5. D. 35; A. 24; P. 15; V. 3. Body slender, compressed; head heavy; snout evenly decurved; mouth large, horizontal, reaching back of eye. Jaws with short, strong, incurved conical teeth. Fin rays long and filamentous, the longest dorsal rays as long as head; anal rays shorter. Coloration faded in the type, but traces of about 7 broad brown vertical bars as broad as eye and twice as broad as the pale interspaces, the dark bars extending upon dorsal fin; ventrals pale. Gulf of Mexico. Known from 1 specimen, $3\frac{1}{8}$ inches long, taken from the stomach of *Ncomænis aya*, on the Snapper Banks off Pensacola, Florida; very close to *E. nivipes*, but more robust, with fewer dorsal rays.

Emblemaria atlantica, JORDAN & EVERMANN, Check-List Fishes, 472, 1896, name only, Snapper Banks off Pensacola, Florida. (Type, No. 33915. Coll. Silas Stearns.)

2757. EMBLEMARIA NIVIPES, Jordan & Gilbert.

Head $3\frac{3}{4}$ in length; depth 7. D. XXIII, 14; A. 25. Body everywhere equally compressed, posteriorly tapering; head wider than body, of about equal depth, with very short, subvertical, sharply compressed snout; eyes very large, approximated above, with some vertical range; orbital ridges sharply raised above, the interorbital region very narrow, channeled, about equaling diameter of pupil; eye $3\frac{3}{8}$ in head. Gape very wide, horizontal, low, reaching much beyond eye, the maxillary about $\frac{1}{2}$ head, not produced beyond angle of mouth; intermaxillaries separated by a groove from the snout, this groove continuous for the entire length of the upper jaw, maxillary not evident, apparently adnate to the skin of the preorbital. First dorsal spine inserted over margin of preopercle; spines all very slender and flexible, the posterior but weakly differentiated from the soft rays, the anterior portion of fin very high, the spines filiform, not exserted beyond the membrane; the longest dorsal spine about $\frac{1}{3}$ length of body, the last spine about $\frac{1}{2}$ head; membranes of last rays of both dorsal and anal slightly joined to base of caudal. Front of anal nearer snout than

base of caudal by a distance equaling $\frac{1}{8}$ length of head. Caudal $\frac{3}{8}$ length of head; ventrals and pectorals slightly less. Color in spirits, sides dark brown, with 8 to 10 lighter vertical bars of variable width; body lighter below; obscure cross bands on lower side of head; dorsal blackish anteriorly, whitish behind, with membrane at intervals of every second, third, or fourth ray dusky; caudal light at base, its tip blackish; anal dusky translucent; ventrals bright white, the basal portion dusky. Pearl Islands, near Panama. A specimen 2 inches long is the type of the species. Numerous smaller specimens were obtained at the same time. (*nix, nivis,* snow; *pes,* foot.)

Emblemaria nivipes, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1883, 627, Pearl Islands, near Panama. (Type, No. 29676. Coll. Prof. Frank H. Bradley.)

2758. EMBLEMARIA OCULOCIRRIS, Jordan.

Head $3\frac{3}{4}$; depth $6\frac{3}{4}$. D. about 35; A. 25. Upper part of eyeball above pupil (sclerotica) with a slender cirrus tipped with black, this nearly as long as eye; eye longer than snout, about $3\frac{3}{4}$ in head, the maxillary extending to below posterior part of pupil; snout sharper than in *Emblemaria nivipes*, $\frac{2}{3}$ eye; teeth small, rather sharp, directed backward; longest dorsal spines as long as head; pectorals $1\frac{1}{3}$ in head; ventrals $1\frac{2}{3}$, inserted before pectorals. Color in spirits, brown, with traces of about 9 blackish cross bars, which are separated on the back by whitish, quadrate interspaces; a white spot at nape; some dusky below eye; dorsal dusky, the pale bars of back extending on its base; anal dusky; ventrals blackish; caudal pale, its tip black; pectorals pale. Gulf of California. Known from 1 specimen, $1\frac{1}{2}$ inches long, from La Paz. It is shriveled and in poor condition. It seems to be very close to *Emblemaria nivipes*, but differs in the presence of an ocular cirrus, in the sharper snout, smaller mouth, and dusky ventrals. The teeth seem rather more slender, but can not be well examined. (*oculus,* eye; *cirrus,* filament.)

Emblemaria oculocirris, JORDAN, in GILBERT, Proc. U. S. Nat. Mus. 1896, 456, La Paz. (Type, No. 47749. Coll. *Albatross.*)

896. CHÆNOPSIS, Gill.

Chænopsis, GILL, Ann. Lyc. Nat. Hist. N. Y., VIII, 1865, 141 (*ocellatus*).

Body naked, eel-like. Head much elongate, quadrate behind, conic in front, profile straight; snout acute, jaws produced; no teeth on vomer, teeth in front of jaws strong, with villiform teeth behind them. Dorsal and anal long, continuous, confluent with the caudal. Dorsal rays about XVIII, 38; anal II, 38. Ventrals inserted slightly before pectorals. West Indies. (*χαίνω,* to yawn; *ὄψις,* face.)

2759. CHÆNOPSIS OCELLATUS, Poey.

D. XVIII, 38; A. II, 38; C. 15. Body naked, eel-like; anus submedian. Head much elongate, quadrate behind at the opercular region, conic in front, with the profile rectilinear and the snout acute; eyes moderate;

mouth large, with the cleft wide and nearly horizontal. Teeth subcylindrical, in a uniform row, behind which, in front, there is a broad band of villiform teeth; on the palatine bones, uniserial and obtusely subcylindrical like those of the jaws; the palatine rows are parallel; vomer edentulous. Gill membranes confluent below, free from the isthmus. Dorsal and anal long, confluent with caudal; ventrals slightly in advance of pectorals, with 2 or 3 rays. (Gill.) Matanzas, Cuba; 1 specimen, examined by us in the National Museum. (*ocellatus*, having eye-like spots.)

Chænopsis ocellatus, POEY, in GILL, Ann. Lyc. Nat. Hist. N. Y., VIII, 1867, 143, Matanzas, Cuba. (Coll. Poey.)

897. LUCIOBLENNIUS, Gilbert.

Lucioblennius, GILBERT, Proc. U. S. Nat. Mus. 1890, 103 (*alepidotus*).

Body very elongate, wholly naked; gill membranes broadly united, free from isthmus; dorsal fin single, extending along the entire back, its anterior half spinous. Ventrals in front of pectorals, I, 2. First two anal rays spinous. Last rays of dorsal and anal joined to caudal. Teeth conic, not movable, in jaws and on vomer and palatines. Lateral line not described. A strange genus, evidently very close to *Chænopsis*. (*Lucius*, pike; *Blennius*, blenny.)

2760. LUCIOBLENNIUS ALEPIDOTUS, Gilbert.

Head 3 in length; depth 3¼ in head. D. XVIII, 32; A. II, 30. Body much compressed, slender throughout, the head rather deeper and wider than body. Snout long, depressed, and rather wide, the anterior profile descending very gradually. Mouth nearly horizontal, the lower jaw protruding, the gape extending to much behind orbit, the entire physiognomy remarkably pike-like. Snout 4 in head; maxillary 1¾; eye4⅔ to 5. Teeth in a villiform band in upper jaw, the outer series slightly larger; in lower jaw in a single series laterally, widening into a patch anteriorly, the outer enlarged; a few teeth only on vomer; palatines with a long and rather broad patch similar to those in jaws. Dorsal fin beginning on the nape in advance of middle of opercle, the fin uniformly low, extending the whole length of back, the posterior ray joined by membrane with the caudal; the spines and rays are similar in appearance, flexible and simple, none of the soft rays branched; the spines are more slender, and show no joints, the articulations being present in small number on all the soft rays; the highest ray is less than diameter of orbit; anal and caudal rays similar to those of soft dorsal; caudal short, rounded; origin of anal midway between tip of snout and end of caudal fin, its first 2 rays spinous; ventrals under opercular margin, of 1 spine and 2 well-developed rays, nearly ½ as long as head; pectorals narrow, of apparently unbranched rays, about ⅓ as long as head. Color light olivaceous, with 11 vertical dark blotches on sides, most of which divide to form on middle of sides double vertical bars; top and sides of head with dark cloudings, and with numerous black specks of varying size; middle of sides and base

of dorsal with numerous pearly dots nearly as large as pupil; branchiostegal membrane black posteriorly; the lateral bars extended to base of dorsal, the anterior ones usually forming conspicuous black blotches which extend well up on the fin; other fins unmarked. Length 1½ inches. Gulf of California; two specimens from *Albatross* Station 3005, in 21 fathoms. (*ἀλεπιδωτός*, scaleless.)

Lucioblennius alepidotus, GILBERT, Proc. U. S. Nat. Mus. 1890, 103, Lower California (Coll. *Albatross*); JORDAN, Proc. Cal. Ac. Sci. 1896, 233, pl. 37.

898. PHOLIDICHTHYS, Bleeker.

Pholidichthys, BLEEKER, Boeroe, 406, 1856 (*leucotænia*).

Body elongate, tapering, naked; snout obtuse; no cirri; teeth unequal, on jaws only; dorsal, anal, and caudal fins distinct, but connected by a membrane; the dorsal formed of flexible spines; the soft rays, if present, not distinguishable from them; ventrals inserted scarcely before the pectorals, of 2 rays. Lateral line and vertebræ undescribed. Tropical parts of the Pacific. (*Pholis; ἰχθύς*, fish.)

2761. PHOLIDICHTHYS ANGUILLIFORMIS, Lockington.

Head 6⅔ in total length with caudal; depth 16. Body exceedingly elongate, much compressed, naked; upper profile of head forming a continuous convex curve to the tip of the snout, which is about equal in length to the eye. Eye lateral, round; interorbital space about ⅜ of the diameter of the eye, convex transversely. Posterior extremity of maxillary vertical with the hinder margin of the eye. Tip of snout a little below the level from the center of the eye; mouth moderately oblique, lower jaw slightly the longer. Teeth of lower jaw in a close-set row, the largest in front, diminishing along the sides; teeth of upper jaw similar, but smaller; palate smooth. Vertical fins continuous, but distinct; dorsal entirely spinous; anal commencing a little behind the middle of the entire length of the fish; ventrals 2-rayed, very slightly in advance of the pectorals, which are about equal in length to the distance of their base from the eye. Color in spirits, dark blackish brown mingled with white upon top, sides, and lower parts of head; interorbital area and top of snout white. Gulf of California; a single specimen dredged off San Jose Island, Amortiguado Bay. Total length 1 5/12 inches. Head ¼ inch. The example is broken across, the branchiostegals are defective, the caudal fin broken and some fin rays missing, so that the fin formula can not be exactly given. The dorsal fin has above 60 rays. The body is much more slender than that of *P. leucotænia*, Bleeker, and there is no trace of the longitudinal bluish-white band of that species. (Lockington.) (*Anguilla*, eel; *forma*, shape.)

Pholidichthys anguilliformis, LOCKINGTON, Proc. Ac. Nat. Sci. Phila. 1881, 118, San Jose Island, Lower California. (Coll. W. J. Fisher.)

899. PSEDNOBLENNIUS, Jenkins & Evermann.

Psednoblennius, JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 156 (*hypacanthus*).

Body compressed, elongate, naked; head short, blunt; no cirri; mouth large, the jaws subequal; teeth in a single series in each jaw, none on vomer or palatines; lateral line not developed. Dorsal fins 2, the first at the nape, of three flexible spines; second dorsal with a few slender spines which pass into the soft rays; anal much shorter than second dorsal, both fins joined to base of caudal; dorsal rays III–34; anal 27; ventral rays 2, the fin directly below pectorals. Apparently close to *Pholidichthys*, but with the dorsal divided and changing gradually from spines to soft rays. (*ψεδνός*, naked; *Blennius*.)

2762. PSEDNOBLENNIUS HYPACANTHUS, Jenkins & Evermann.

Head 4⅛ (5 in total); depth 7 (8); eye 4, equal to snout; B. 6. D. III-34; A. 27. Body greatly compressed, elongate; head short, snout blunt, about equal to eye; anteorbital profile very steep, gently rounded from front of eye to first dorsal, from there nearly straight to caudal; ventral line nearly straight. Body naked, no membranaceous appendages. Mouth large, horizontal, jaws subequal, extending to beyond middle of eye. Teeth in a single series in each jaw, well developed, pretty uniform in size, slightly projecting backward; vomer and palatines apparently smooth. Eye large, equal to twice interorbital space, high up. Dorsal fins 2, the first of 3 very slender, flexible spines, hard to distinguish from soft rays, but they do not appear to be at all jointed. This fin is inserted upon the nape immediately above the posterior edge of the preopercle, and a distance in front of second dorsal nearly equal to length of snout, its very soft spines equal distance from end of snout to posterior rim of orbit; second dorsal begins directly over origin of pectorals and extends to caudal, with which it is slightly connected; first few rays of second dorsal very weak, flexible spines, the last few pretty evidently soft, jointed rays, while the intermediate ones are not distinguishable as definite spines or soft rays—in short, there seems to be a gradual change from spines to soft rays from the anterior to the posterior part of the fin. This character, if we mistake not, is entirely unique. The fin is of nearly uniform height, the rays about equaling those of the first dorsal in length; anal similar to second dorsal in shape and height, but much shorter, its origin being much behind that of the second dorsal or nearly halfway from the snout to base of caudal; posteriorly it extends coterminously with the dorsal, and, like it, is slightly joined to the caudal fin; caudal fin apparently rounded, fan-shaped, but its shape can not be exactly made out, as some of its rays are broken off; pectorals inserted below axis of body, directly over ventrals, their length about ¾ that of head; ventrals of 2 rays, inserted under pectorals, about equal to pectorals in length; body entirely scaleless. Coloration in alcohol, pale, mottled with fine dark points so arranged as to inclose circular areas with fewer spots; a long dark blotch behind the axil, inclining downward and backward; head covered with similar punctulations; opercles dusky; chin with 2

dark cross lines, separated by 1 of white, extending onto upper jaw on each side; top of head with a purple spot; sides with a series of about 6 short black lines, the last broadest and plainest; base of caudal with a distinct black blotch; first dorsal quite dark, almost black; second dorsal with about 8 pretty well-defined dark blotches at its base, rest of fin with numerous dark spots of different sizes; anal with about 12 dark blotches extending somewhat regularly from the base slightly forward, these separated by plain unmarked spaces of a little greater width; caudal sparingly marked with dark points arranged in wavy cross bars; pectorals and ventrals unmarked. Gulf of California at Guaymas. A single specimen, $1\frac{3}{4}$ inches long, obtained from a shallow arm of the bay. (Jenkins & Evermann.) (*ὑπό*, below (imperfect); *ἄκανθα*, spine.)

Psednoblennius hypacanthus, JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 156, Guaymas, Mexico. (Type, No. 39638. Coll. Jenkins & Evermann.)

900. STATHMONOTUS, Bean.

Stathmonotus, BEAN, Proc. U. S. Nat. Mus. 1885, 191 (*hemphillii*).

Body moderately long and low, much compressed; head small, compressed, naked; mouth small, oblique; conical teeth in both jaws, in 2 series, the outer slightly enlarged and, in the upper jaw, somewhat recurved; a few teeth on the vomer. Gill membranes, as in *Pholis*, broadly united, free from the isthmus. Scales none. No lateral line. Dorsal fin long and low, beginning near the head, and consisting entirely of stiff, sharp spines, which are very short anteriorly and gradually increase in size posteriorly. Anal similar to dorsal, with 2 spines and many soft rays. Caudal short, rounded, scarcely separated from the dorsal and anal; pectorals small, much smaller than in *Pholis*, containing only a few rays; ventrals better developed than in *Pholis*, their position more anterior, consisting of a spine and 2 rays. Pseudobranchiæ absent. Branchiostegals 5. Coast of Florida. (*στα θμή*, a carpenter's rule; *νῶτος*, back.)

2763. STATHMONOTUS HEMPHILLII, Bean.

Head 7; depth 8 to $8\frac{1}{2}$; D. LI; A. II, 27; V. I, 2; P. 5 or 6; eye 6 in head. Maxillary extending about to vertical through hind margin of eye; jaws subequal, or the lower projecting very slightly beyond upper; eyes small, separated by an interspace about equal to their own length, and very slightly greater than length of snout; pectoral very little more than $\frac{1}{6}$ as long as head, and scarcely as long as ventral; dorsal beginning over posterior end of pectoral, its anterior spines very much shorter than the posterior ones; length of caudal about equal to length of postorbital part of head; vent slightly in advance of middle of total length to base of caudal, and about under the twentieth dorsal spine. Colors from the alcoholic specimen: A white line extending from tip of snout to caudal, divided into small segments by short cross bars, the first 2 on the head, and the last at origin of caudal; posteriorly, these short bars extend downward, terminating slightly below the base of the dorsal fin; several white blotches, simulating bars, on posterior half of anal fin; edge of

caudal white; sides and under surface of head with several whitish oblique bands forming V-shaped markings; a few roundish white blotches on sides of head, the most conspicuous behind eye; general color darkish brown, nearly black. Length about 2 inches. Key West; 2 specimens known. (Bean.) (Named for the collector, Henry Hemphill.)

Stathmonotus hemphillii, BEAN, Proc. U. S. Nat. Mus. 1885, 191, pl. 13, Key West, Florida. (Coll. Henry Hemphill. Type, No. 37193, U. S. Nat. Mus.)

901. BRYOSTEMMA, Jordan & Starks.

Bryostemma, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 841 (*polyactocephalum*).

Body moderately elongate, covered with small scales; snout short; no teeth on vomer or palatines; teeth in jaws small; gill membranes united, free from the isthmus; nostrils, orbital regions, and neck with dermal flaps, the supraorbital flaps high. Dorsal fin long, of spines only; pectorals well developed, more than half length of head; ventrals well developed, jugular; caudal fin distinct. No air bladder or pyloric cæca. No true lateral line; a short series of large pores above pectoral. North Pacific, representing *Chirolophis* of the Atlantic. This genus differs from the European genus, *Chirolophis*, Swainson (*Blenniops*, Nilsson), in the absence of a true lateral line. Dr. Boulenger informs us that a true median lateral line is developed in *Chirolophis ascanii*. (βρύον, moss; στέμμα, crown.)

a. Dorsal with about 60 spines; anal with about 55 soft rays; a black spot on anterior part of dorsal, but no ocelli posteriorily. POLYACTOCEPHALUM, 2764.

aa. Dorsal with about 54 spines; anal with 40 soft rays; dorsal with several black ocelli, most distinct posteriorly. NUGATOR, 2765.

2764. BRYOSTEMMA POLYACTOCEPHALUM (Pallas).

Head 6½; depth 6. D. LXI; A. 55 (51 to 57); P. 14; V. I, 3; lateral series with 9 to 15 pores. Body elongate, much compressed, covered with small, smooth, embedded scales. Head very short, blunt in profile; mouth short, terminal, the maxillary 3 in head; lower jaw heavy, projecting, its tip with 2 small slender cirri, which are pale in color; teeth subequal, small, bluntish, close set, in 1 row in each jaw; eyes 4 in head, near together; the snout 4; supraorbital cirri 2⅛ in head; interorbital space flat; a flat fringed cirrus over front of eye, these 2 joined at base, about 3 in head; a small cirrus about ½ length of this over posterior part of each eye, these 5 to 6 in head; top of head and nape covered with series of erect cirri, the longest nearly as long as eye; about 15 minute cirri along dorsal edge of lateral line, 1 on each pore. Rows of pores running around eye, under preopercle, and along entire length of the short lateral line; lateral series of pores ½ length of head; gill rakers not developed. Dorsal fin beginning over pectoral and running to caudal; anterior rays fringed with fleshy cirri; first ray, including cirri, 2 in length of head; anal beginning close behind vent and running to caudal, to which it is joined at base; distance from tip of snout to vent nearly 3 in body; pectoral fin but little shorter than head, its breadth at base not ½ its length. Color in spirits, pale

brownish, plain or mottled with darker, with about 13 dark blotches along dorsal and anal fins, more distinct on dorsal; a black spot on fourth to sixth dorsal spines very distinct; a faint one on anterior part of anal; a few dark markings about head and nape; cirri mostly pale. Bering Sea, south to Puget Sound and Yezo. Here described from a fine specimen, 6½ inches long, from Port Orchard, near Seattle, collected by Prof. O. B. Johnson. Other specimens before us from St. Paul (Pribilof Islands), from *Albatross* Stations 3213 and 3274, south and north of the Peninsula of Alaska, and from Petropaulski Harbor, Kamchatka. These specimens show a great deal of variation, and possibly represent 3 different species. It is more likely, however, that they represent extremes of variation. Young examples, collected by the *Albatross* in eastern Bering Sea, are more elongate and less compressed; body much mottled and vaguely barred; ventral fins checkered in fine pattern; head sand color; a black blotch on fourth to sixth dorsal spine; anterior dorsal spine little elevated and with few fringes; sides of head without cirri; anterior cirri joined almost to the tip, a little shorter than the posterior cirri, which are long and very slender. In 1 specimen of these, however, the cheeks are covered with densely matted cirri extending from the angle of the mouth to the dorsal. In these examples the anterior cirri are short and separate, about as long as the posterior cirri. The larger example, 75 cm. long, from Petropaulski, is evidently the typical *polyactocephalum*, and corresponds perfectly to Herzenstein's account of *B. japonicum*. It shows the following characters: Head 6⅓; depth 5⅓. D. LXI; A. 45; P. 14; V. I, 3; lateral series with 6 pores. Body a little deeper than in Puget Sound examples; head short, blunt in profile; mouth short, terminal, oblique, the maxillary 2⅔ in head; lower jaw heavy, projecting, its tip with 2 broad fringed flaps of a dark color; eyes 4 in head, close together, the interorbital space concave; a fringed cirrus above each eye in front, the 2 connected with each other only in the thickened skin at base; a similar cirrus over each eye behind; the posterior cirri ⅓ longer than the anterior ones, 2⅓ in head; top of head and nape with similar cirri, none of them longer than pupil; a few small cirri on cheeks and opercles; some along lateral series of pores, which is 2½ in head; anterior rays of dorsal fringed with fleshy cirri, the first 2 in head; distance from snout to vent 2¾ in body; pectorals nearly as long as head, the rays thickened in the adult, the base of the fin about ½ its length. Color very dark brown, with vague cross bands and many spots; dorsal and anal each with a broad black edge; other fins all black, the caudal barred. Perhaps the dark coloration and long cirri are characters of the adult male. (*πολύς*, many; *ἀκτίς*, ray; *κεφαλή*, head.)

Blennius polyactocephalus, Pallas, Zool. Rosso-Asiat., III, 179, 1811, Kamchatka.

Chirolophus japonicus, Herzenstein, Mélanges Biologiques Soc. Sci. Petersb., XIII, 1890, 123, Yezo.

Chirolophis polyactocephalus, Jordan & Gilbert, Synopsis, 765, 1883; Bean in Nelson, Rept. Nat. Hist. Coll. Alaska, 305, pl. 15, f. 2, 1887.

Bryostemma polyactocephalum, Jordan & Starks, Proc. Cal. Ac. Sci. 1895, 841; Jordan & Gilbert, Rept. Fur Seal Invest., 1898.

2765. BRYOSTEMMA NUGATOR, Jordan & Williams.

Head 5½; depth 5½; D. LIV; A. 41; V. I, 3; pores of lateral line 25. Body elongate, less compressed than in *Bryostemma polyactocephalum,* covered with small, smooth, embedded scales. Head short, very obtuse, almost truncate; top of head from nostrils to near front of dorsal covered with fleshy cirri, much smaller than in *B. polyactocephalum;* only 2 or 3 small ones extending on first dorsal spine; supraorbital cirrus short, 4 to 5 in head; 2 small cirri placed at the sides of snout with a larger median one behind them, forming a triangle; jaws equal; mouth horizontal, the angle extending to below pupil; eyes small, 4 in head; snout very short, almost vertically truncate, ⅜ in eye; teeth of both jaws subequal, short, bluntish, and close set. Lateral line short, 7½ in length of body, concurrent with the dorsal outline of body. A line of pores begins in front of eye on a level with pupil, runs under eye and to a level with pupil again, then back to and along the entire length of the short lateral line. Gill rakers not developed; gill membranes free from isthmus. Vent ⅓ distance from tip of snout to tip of caudal; distance from origin of ventral to anus 4½ in length of body Pectoral fin 5½ in body, as long as head. Dorsal fin beginning in front of the pectoral, highest along the posterior half, the longest spine 2⅜ in head, the fin higher than anal; dorsal slightly joined to caudal; anal separated from caudal; caudal rounded, 1⅔ in head; first dorsal spine 4½ in head, its surface with 2 or 3 small cirri. Color in spirits of 1 specimen, probably male, dark brown, with 13 pale cross bars along back, extending on dorsal fin; along sides these become obsolete; on belly they become increased in number and broadened below; dorsal fin with 13 large, very distinct black ocelli with yellowish rings, 1 between each pair of the pale blotches; anal with about 7 small blackish spots at base on posterior part, the fin otherwise nearly plain; caudal faintly barred with light and dark; pectorals pale, with 2 dark pale-edged oblique bars before it; sides of head with irregular dark vertical bars, 1 of them forming an inverted Λ below eye, this and others extending across lower jaw; cirri mostly black. The other specimen, probably the female, has the body nearly plain brown, the dorsal with but 4 ocelli, the anterior 9 being replaced by dark bars on the fin; anal with dark oblique cross bars; pectorals barred with black; markings on head more sharply defined, coloration otherwise similar. This second specimen is 4¾ inches in length, the other 4. Puget Sound; the above account from the 2 original types from near Seattle. Three others since obtained near Channel Rocks, Port Orchard, show the following life coloration: Dark red above, orange brown below, belly cream color; sides below with cream-colored cross bars, wider than eye, running from the axis of body downward and fading into the general color below; a Λ-shaped mark downward from eye across branchiostegals to isthmus, a similar mark behind eye across edge of preopercle, this last sometimes broken up and chain-like; top of head dark; snout light; 2 oblique dark bars at base of pectoral; dorsal with 12 or 13 sharp dark brown spots as large as eye, edged with bright red, these arranged regularly along the whole length of fin; pectorals and caudal bright red with wavy, irregular, brown

lines running across the rays; anal red, with dark brown bars as wide as the interspaces running obliquely downward and forward; ventrals light brown. (*nugator*, a fop.)

Bryostemma nugator, JORDAN & WILLIAMS, Proc. Cal. Ac. Sci. 1895, 843, pl. 101, Seattle, Washington. (Coll. Young Nat. Soc. Type, No. 3134, L. S. Jr. Univ.)

902. APODICHTHYS, Girard.

Apodichthys, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 150 (*flavidus*).

Body elongate, compressed, covered with very small scales; no lateral line; snout short; mouth moderate, oblique; teeth in the jaws moderate, stouter anteriorly; vomer with teeth; gill membranes united, free from the isthmus. Dorsal fin long, low, even, of spines only; anal fin similar, preceded by a very large pen-shaped spine channeled along its anterior surface and hidden in a pouch of skin; caudal fin short, connected with dorsal and anal; no ventral fins; pectoral fins moderate; intestinal canal short, without pyloric cæca. Small, bright-colored fishes of the Pacific, living among rocks near shore. (ἄπους, without feet; ἰχθύς, fish; in allusion to the want of ventral fins.)

a. Color various, green, olive, or scarlet; sides of head without silvery band; depth 7 to 8 in length; head 9. FLAVIDUS, 2766.
b. General color olivaceous. var. *flavidus*, 2766a.
bb. General color scarlet. var. *sanguineus*, 2766b.
bbb. General color grass green. var. *virescens*, 2766c.
aa. Color reddish; a bluish silvery stripe on side of head; depth 9 to 10 in length; head 7. UNIVITTATUS, 2767.

2766. APODICHTHYS FLAVIDUS, Girard.

Head 9⅛; depth 7⅓. D. XCIII; A. I, 40. Head short; mouth very oblique; maxillary reaching pupil; upper jaw with a series of conical teeth, behind which is a patch of smaller teeth; sides of mandible with conical teeth in a single series, forming a patch in front; vomer with 3 conical teeth; palatines toothless; nape equidistant between front of dorsal and pupil. Anal spine very large, ⅔ length of head, shaped like a pen, deeply excavated on its anterior side, and very convex behind, very thin, flexible, and with sharp edges, entirely included in a pouch of skin; pectoral fins about ⅔ length of head. Color orange, varying with the surroundings to intense grass-green (var. *virescens*), yellowish brown (var. *flavidus*), crimson and dark purple (var. *sanguineus*); a few light round spots along axis of body posteriorly; a narrow black bar downward and backward from eye; a shorter, less distinct bar from upper margin of orbit backward to occiput; anal fin obliquely barred with brownish. Length 18 inches. Pacific coast, Vancouver Island to the Santa Barbara Islands; abundant; usually found below low tide mark. The following color notes are from specimens taken in Puget Sound belonging to the green form (var. *virescens*), the larger 10 inches in length, the smaller 3 inches. The large one is a bright grass-green, mottled with light gray; a series of blended white spots, as large as eye, along the axis of body

from the pectoral fin to the middle of caudal peduncle; belly with many similar spots smaller in size and somewhat sharper in outline; a row of conspicuous black spots, irregular in size, shape, and position, along back at the base of dorsal spines; a black line as wide as pupil from nape to eye, a similar line from eye to posterior end of maxillary; a faint light streak across cheek posteriorly; cheek and base of pectoral dusted with fine dark points. The small one is bright green without distinct markings on body; a silvery bar, running posteriorly from tip of snout through eye, across cheek, to the middle of opercle; no bar downward from eye to maxillary, or from eye to nape as in the large one. (*flavidus*, yellowish.)

Apodichthys flavidus, GIRARD, Proc. Ac. Nat. Sci. 1854, 150, Presidio, San Francisco Bay (Coll. Dr. Kennerly. Type, No. 494, U. S. Nat. Mus.); GIRARD, Pac. R. R. Surv., x, Fishes, 117, 1858; GÜNTHER, Cat., 290, 1861; JORDAN & GILBERT, Synopsis, 769, 1883.

Apodichthys virescens, AYRES, Proc. Cal. Ac. Nat. Sci. 1855, 55, San Francisco; GIRARD, Pac. R. R. Surv., x, Fishes, 118, 1858.

Apodichthys inornatus, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 279, Puget Sound, probably (Coll. Northwestern Boundary Commission); D. XC; A. 38.

Apodichthys sanguineus, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 279, California. (Coll. Dr. Samuel Hubbard.)

2767. APODICHTHYS UNIVITTATUS, Lockington.

D. about XCV; A. about I, 40. Body elongate, much compressed, band-like, preserving almost same depth to about posterior fifth of body, thence tapering more rapidly to caudal fin. Head 7; depth nearly 10 times in total length; depth of caudal peduncle about $\frac{1}{2}$ of that of body; snout obtuse, about $\frac{2}{3}$ as long as diameter of eye, upper profile of head a continuous curve from snout to occiput. Interorbital area highly convex transversely, about equal in width to $\frac{1}{2}$ diameter of eye. Eye entirely lateral, round, contained entirely in anterior half of head; iris golden. Mouth small, posterior extremity of the maxillary reaching to anterior margin of eye. Teeth small. Branchiostegals 5. Dorsal continuous with, but distinct from, anal, arising vertically from tip of operculum, and composed of spines only. Anal preceded by a long, sharp, slender spine of V-shaped transverse section, hollow side anterior, length of spine equal to about $\frac{1}{2}$ depth of fish. Distance from anal spine to tip of operculum a little more than to tip of caudal. Caudal with numerous accessory rays, so that its sides are almost straight, posterior margin broken in the type, all rays simple. General color in spirits, light reddish, vertical fins rather bright, and top of head reddish brown; tip of snout brown; a silvery band (possibly bluish in life) from tip of snout, across lower part of eye, cheek, and opercles, terminating at about middle of length of operculum, this band bordered above by a narrower brown band. Lower California, probably from the gulf. A single specimen. Length 1.88 inches. The peculiar vitta upon each side of the head at once distinguishes this species from the other described forms. (Lockington.) Not seen by us. (*uni-vittatus*, having one band.)

Apodichthys univittatus, LOCKINGTON, Proc. Ac. Nat. Sci. Phila. 1881, 118, Gulf of California.

903. XERERPES, Jordan & Gilbert.

Xererpes, JORDAN & GILBERT, Proc. Cal. Ac. Sci. 1895, 846 (*fucorum*).

This genus differs from *Apodichthys* in the moderate size of the anal spine, which is rounded and not channeled on its anterior edge, and in the small size of its pectoral fins. The single known species lives in *Fucus* chiefly above low-tide mark and may often be shaken out of half-dry mats of seaweed on rocks well above the water. (ξερός, dry; ἕρπης, creeper.)

2768. XERERPES FUCORUM (Jordan & Gilbert).

Head 10; depth 9½. D. LXXXIII; A. 35. Form and dentition as in *Apodichthys flavidus*. Mouth very oblique, the maxillary reaching center of pupil; nape nearer front of dorsal than end of snout. Anal spine comparatively small, about ⅓ length of head, transversely very convex in front, and slightly concave or grooved behind, the pouch of skin at its base little developed; pectorals very small, shorter than eye; anal fin beginning nearer tip of caudal than tip of snout by about 3 times length of head. Bright olive green or deep red, the color varying with the surroundings; a row of dark spots along axis of body, these sometimes with light-bluish center, and connected by a very narrow dark streak; generally a dark streak downward from eye, but no other markings about head. Length 6 inches. Monterey to Puget Sound; abundant in rock pools and bunches of *Fucus*; remarkable for its active movements. It is found mostly in masses of *Fucus* attached to rocks between tide marks, and it is often found at low tide at a considerable distance from any water, kept damp by the masses of algæ. Sometimes a dozen of them can be shaken from a bunch of algæ attached to a dry rock. It is, like the species of *Xiphidion*, very active, moving over stones or sand, and showing less anxiety about the presence of its native element than any other fish known to us. (*fucorum*; of the *Fucus* or seaweed.)

Apodichthys fucorum, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 139, Monterey (Coll. Jordan & Gilbert); JORDAN & GILBERT, Synopsis, 770, 1883.
Xererpes fucorum, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 846.

904. ULVICOLA, Gilbert & Starks.

Ulvicola, GILBERT & STARKS, Proc. U. S. Nat. Mus. 1896, 455 (*sanctæ-rosæ*).

This genus is allied to *Xererpes*, but differs in having the opercle above angle adnate to shoulder girdle, in the smaller size of the anal spine, and especially in the entire absence of pectoral fins. (*Ulva*, sea lettuce; *colo*, I inhabit.)

2769. ULVICOLA SANCTÆ-ROSÆ, Gilbert & Starks.

Head 10 in body; depth 13. D. XCVII; A. I, 40; eye 4½ in head; caudal 1½. Body elongate, as in *Apodichthys*, strongly compressed, upper profile of head slightly convex, no construction at nape; mouth very small, oblique, the maxillary reaching about to front of eye; teeth very small in

a single row on jaws; vomer with teeth; interorbital a narrow, sharp ridge; snout about equal to length of eye; gill opening short, limited to the part below angle of opercle, above adnate to shoulder girdle. Origin of dorsal above upper end of gill opening, much nearer occiput than tip of snout; anal spine small, not channeled as in *Apodichthys flavidus*; origin of anal nearer base of caudal than tip of snout by a distance equal to twice length of head; pectorals and ventrals obsolete; caudal rather long, confluent with dorsal and anal. Color in spirits, light brown, slightly lighter under head and on belly; no markings. The type is a specimen 4½ inches in length, collected by the *Albatross* at Santa Rosa Island, off Santa Barbara, January 6, 1889. (Type, No. 47579. Coll. *Albatross*.)

Ulvicola sanctæ-rosæ, GILBERT & STARKS, Proc. U. S. Nat. Mus. 1896, 455, pl. 55, fig. 2, Santa Rosa Island, California.

905. PHOLIS (Gronow) Scopoli.

(GUNNELS.)

Pholis, GRONOW, Zoophylaceum, 78, 1765 (not binomial).
Pholis, SCOPOLI, Introd. Hist. Nat., 456, 1777 (*gunnellus*).
Murænoides, LACÉPÈDE, Hist. Nat. Poiss., II, 324, 1800 (*sujef*).
Centronotus, BLOCH & SCHNEIDER, Syst. Ichth., 165, 1801 (*fasciatus*).
Dactyleptus, Rafinesque Anal. de la Nature 1815, 82; substitute for *Murænoides*.
Centronotus, CUVIER, Règne Animal, Ed. 2, II, 239, 1829 (*gunnellus*).
Ophisomus,* SWAINSON, Nat. Hist. Class'n. Anim., II, 277, 1839 (*gunnellus*).
Urocentrus, KNER, Sitzber. K. Akad. Wiss. Wien, LVIII, 1868, 51 (*pictus*).
Rhodymenichthys, JORDAN & EVERMANN, Check-List Fishes, 474, 1896 (*ruberrimus* = *dolichogaster*).

Body long and low, considerably compressed, somewhat band-shaped, the tail slowly tapering; head small, compressed, naked;† mouth rather small, oblique; jaws with rather small teeth in narrrow bands or single series; vomer and palatines usually toothless; gill membranes broadly united, free from the isthmus; scales very small, smooth; no lateral line. Dorsal fin long and low, beginning near the head, composed entirely of stiff, sharp, subequal spines; anal similar in form, of 2 spines and many

* Substitute for *Gunnellus*, the latter being a barbarous word derived from "gunwale." "Nomina generica quæ ex Græca vel Latina lingua radicem non habent, regicienda sunt." This rule has never been generally adopted.

† In *Pholis nebulosus*, a Japanese species, the head is scaly. This species is the type of a distinct genus; which may be called

ENEDRIAS, Jordan & Gilbert, new genus.

Enedrias, JORDAN & GILBERT, new genus (*nebulosus*).

This genus differs from *Pholis* in the scaly head. (*ἐνέδρα*, lurking place.)

Enedrias nebulosa (SCHLEGEL).

Head 7½ to 8; depth 8⅓ to 9⅓. D. LXXX; A. II, 39. Dorsal and anal somewhat connected to caudal; pectoral 2⅔ to 2⅘ in head. Head small. Body everywhere freckled with dark blotches; 12 dark triangular blotches along base of dorsal; a row of dusky blotches on middle of side posteriorly; 10 or 12 dark blotches on base of anal; caudal dusky, edged with pale, 2 pale cross streaks on top of head; pectoral pale. Northern Japan to Okhotsk Sea, Gulf of Strietok; our specimens from Hakodate. (*nebulosus*, clouded.)

Gunnellus nebulosus, SCHLEGEL, Fauna Japonica, Poiss., 138, 1850, Bay of Magi, Japan.
Centronotus nebulosus, STEINDACHNER, Ichth. Beitr., IX, 24, 1880.
Enedrias nebulosus, JORDAN & GILBERT, Rept. Fur Seal Invest., 1898, with plate.

soft rays; caudal fin short and small, more or less joined to dorsal and anal; pectorals short, rather shorter than head; ventrals very small, of 1 spine and a rudimentary ray; intestinal canal short, without cæca. Shore fishes of the Northern seas. (*φωλίς*, name of some fish said to shelter itself when lying in wait by producing a cloud of mucus; *φωλάς*, one who lies in wait.)

UROCENTRUS (*οὐρά*, tail; *κέντρον*, spine):

a. Pectoral fin small, 3½ to 4 times in length of head; dorsal spines about 93; anal rays 48; body with 2 rows of dark blotches; fins nearly plain. PICTUS, 2770.

aa. Pectoral fin moderate, 2 to 2½ times in length of head.

RHODYMENICHTHYS (*Rhodymenia*, a large red alga; *ῥόδον*, rose; *ὑμήν*, membrane; *ἰχθύς*, fish).

b. Dorsal and anal joined to the caudal to the full height of the spines, without constriction at base of caudal; body greatly compressed, ribbon-like. Dorsal spines about 93; anal about 47; pectorals short, 2⅔ in head; no ocelli along base of dorsal. DOLICHOGASTER, 2771.

PHOLIS:

bb. Dorsal and anal slightly connected with caudal, leaving a constriction of outline at base of caudal; body less compressed; dorsal fin with dark blotches or ocelli.

c. Pectoral fins well developed, about ½ length of head. Dorsal spines about 88; anal rays about 42; pectoral 2½ in head; dorsal fin with dark quadrate blotches rather than ocelli; sides scarlet in adult, bounded with black. FASCIATUS, 2772.

cc. Dorsal spines about 80 (76 to 85); anal rays about 40; pectoral 2 in head; dorsal fin with small rounded black blotches. GUNNELLUS, 2773.

ccc. Dorsal spines about 77; anal rays about 35; pectoral 2 in head; dorsal fin with ocelli, or lunate, dark blotches. ORNATUS, 2774.

Subgenus UROCENTRUS, Kner.

2770. PHOLIS PICTUS (Kner).

Head 9½ to 10½; depth 8 to 10. D. XCIII or XCIV; A. II, 46 to 48 (misprinted 40 in Kner's account). Eye as long as snout; mouth oblique, the upper jaw the longer, reaching to front of eye; pectorals very short, scarcely longer than eye, 3 to 4 in head; anal said to have an isolated channeled spine hidden in the skin, but our specimens show no peculiar structure. Color yellowish, with 2 lengthwise series of large oblong blackish blotches, the one along base of dorsal, but not on the fin, of 21 or 22 blotches, the other on lower part of sides, of about 25; a series of fainter blotches along base of anal; in other specimens the lower row becomes obscure, the upper more distinct, and the series above anal disappears; a black bar downward from eye, a whitish band behind it; opercles dusky. West side of Bering Sea; our specimens from Shana Bay, Iturup Island, Kuril Group.

As already shown by Steindachner, this is a typical *Pholis*, Kner having been in error in ascribing to it an isolated and channeled first anal spine. The ventral spines are bound down by the integument more closely than usual, but they are in other respects not peculiar. Each is accompanied by 2 short spinous rays concealed in the membrane, and difficult to detect.

The latter are stiff and pungent, and seem to be not articulated. The ventrals of *P. ornatus* show the same structure. Kner gives the anal formula as II, 40. This must be a misprint for II, 49, as the artist figures 51 rays in the fin, not differentiating the 2 anterior ones. (*pictus*, painted.)

Urocentrus pictus, KNER, Sitzungsb. d. k. Akad. D. Wissench., LVIII, 1868, 51, taf. 7, fig. 21, Singapore; an error.
Centronotus pictus, STEINDACHNER, Ichth. Beiträge, IX, 25. 1880.
Pholis pictus, JORDAN & GILBERT, Rept. Fur. Seal Invest., 1898.

Subgenus RHODYMENICHTHYS, Jordan & Evermann.

2771. PHOLIS DOLICHOGASTER* (Pallas).

(BUTTER-FISH.)

Head 9⅓ in length; depth 8. D. XCII; A. II, 44; pectoral 14; eye 5 in head; maxillary 2¾; pectoral 2½; caudal 2; ventral spines 1⅔ in eye. Body elongate, much compressed; head small, its upper profile convex; mouth moderate, very oblique, the maxillary reaching to below middle of eye; teeth rather large and blunt, arranged in a single row, the anterior one not enlarged; interorbital space narrow, without a sharp ridge, its width less than eye; snout equal in length to eye; distance from tip of snout to occiput 1⅕ in head. Head entirely naked; body covered with small, cycloid, inconspicuous scales. Origin of dorsal over upper end of gill slit, its distance from nape equal to distance from nape to front of eye, the spines toward the anterior end of the fin the highest; origin of anal a little nearer tip of caudal than snout; dorsal and anal confluent with caudal, the anal more broadly connected than dorsal; pectorals small, rounded behind; ventral spines inserted directly under base of pectorals, their length little greater than their distance apart; caudal short and broad, well rounded in outline. Bering Sea; recorded from the Kurils, and from Robben, Bering, and Medni islands, and from Kigiktowik Bay. The specimen above described was taken at Robben Island by Capt. J. G. Blair, then in command of the guardship *Leon*. It is 9 inches long and is uniform red in color, with a few pale dots. Another specimen, 18 cm. long, taken by Mr. Gerald E. H. Barrett-Hamilton at Bering Island, shows the following characters: The color is cherry red on the body and fins, lighter on belly, lower half of cheek and under side of head; lips blackish anteriorly, a narrow black streak running from them along snout to eye and from eye across cheek and opercles toward upper edge of pectoral base; this line separates the deep-red upper part of the head from the lighter area below;

* The following species is allied to *Pholis dolichogaster:*

Pholis taczanowskii (Steindachner).

Head 9; depth 10; D. LXXXII; A. II, 45; teeth bluntly conical; dorsal very low, joined to the caudal without constriction. Snout scarcely longer than eye, which is 5⅓ in head. Pectoral 3 in head. Scales very small, the head naked. Clear, yellowish gray, finely dotted, fins grayish, the pectoral yellowish; a yellowish streak edged with darker from eye to axil. Gulf of Strietok. (Steindachner.) (A personal name.)

Centronotus taczanowskii, STEINDACHNER, Ichth. Beitr., IX, 24, pl. 3, fig. 1, 1880, Gulf of Strietok, Okhotsk Sea. (Coll. Prof. Dybowsky.)

sides of body with a number of minute scattered black spots; along middle of sides is a distant series of light spots as large as pupil, the margin of each with 2 to 4 black specks like those scattered over sides. The dorsal and anal are more widely joined to the caudal than in other species, the fins being higher posteriorly and without perceptible notch. The dorsal contains 93 spines, the anal 2 spines and 47 rays, the pectorals 15 rays. Head $9\frac{1}{2}$ in length; depth $7\frac{5}{6}$. Eye 5 in head; maxillary $3\frac{3}{4}$; pectorals $2\frac{1}{2}$; caudal $2\frac{1}{3}$; ventral spine $2\frac{1}{3}$ in eye. *Blennius dolichogaster*, Pallas, is undoubtedly identical with *Gunnellus ruberrimus*, Cuvier & Valenciennes. They agree in the very long dorsal and anal fins (D. XCIII, A. II, 50 in *dolichogaster*), and in the color. *P. dolichogaster* is described as having the color brownish olive, shaded with greenish and yellowish, spotted with green above the lateral line; belly yellow; anal, caudal, and pectorals yellowish; dorsal and anal dusky, with transverse pale bars. Compare with this, details of coloration recently published concerning *P. ruberrimus* by Bean & Bean (Proc. U. S. Nat. Mus. 1896, 248): "Color olive brown, with minute black spots; belly yellowish." In another specimen, "Across the spinous dorsal there are 20 narrow, nearly vertical pale streaks. Similar streaks to the number of 12 cross the anal." The species is evidently not always red in life. (*δολιχός*, long; *γαστήρ*, belly.)

Blennius dolichogaster, Pallas, Zoogr. Rosso-Asiat., III, 175, 1811, Kamchatka. (Type in Mus. Berlin.)

Gunnellus ruberrimus, Cuvier & Valenciennes, Hist. Nat. Poiss., XIV, 440, 1839, Kuril Islands; after notes of Pallas,* Zoogr. Rosso-Asiat., III, 178, 1811.

Gunellus dolichogaster, Cuvier & Valenciennes, Hist. Nat. Poiss., XI, 436, 1836.

Centronotus dolichogaster, Günther, Cat., 288, 1861.

Murænoides dolichogaster, Jordan & Gilbert, Synopsis, 768, 1883.

Pholis dolichogaster, Jordan & Gilbert, Rept. Fur Seal Invest., 1898.

Rhodymenichthys ruberrimus, Jordan & Evermann, Check-List Fishes North and Middle America, 474, 1896.

Pholis ruberrimus, Bean & Bean, Proc. U. S. Nat. Mus. 1896, 248.

Subgenus PHOLIS.

2772. PHOLIS FASCIATUS (Bloch & Schneider).

Head 8 to $9\frac{1}{2}$; depth 7 to 9. D. LXXXVI to LXXXIX; A. II, 42 to 44; V. I, 1. Head scaleless; mouth decidedly oblique, the tip of lower jaw on level of middle of the eye; width of mouth nearly $\frac{1}{2}$ head. Eye equal to snout, a little more than interorbital width; ventral spine $\frac{3}{8}$ eye, $\frac{1}{2}$ length of mandible; caudal $\frac{1}{2}$ head; pectoral $2\frac{1}{4}$ in head; vertical fins slightly joined at base. Ground color yellowish gray in life, the sides of a brilliant scarlet; base of dorsal occupied by 10 or 11 oblong blotches of dark brown, which extend to the tips of the fins; these blotches each divided upon the fin by a median spot of the ground color; the areas of the ground color alternating with these blotches are finely speckled with brown, a large spot of brown usually occupying a median

* Bright red. Form of *tænia;* scales inconspicuous; ventrals each a single scarcely projecting spine; caudal broad, rounded, distinct. D. CXV. Kuril Islands. (Pallas.) *Murænoides ruberrimus*, Bean, in Nelson, Rept. Nat. Hist. Collections made in Alaska, 305, pl. XIV, fig. 1, 1887.

position upon the fin; middle and lower part of sides occupied by vermiculating brown lines on the ground color, these vermiculations arranged in more or less distinct cross bars, about 20 in number, reaching to or nearly to the midventral line, the posterior ones often continued faintly onto the anal fin; pectoral and caudal fins yellow, unmarked; a brown blotch across snout and tip of mandible, followed by a narrow yellowish bar descending to front of eye; interorbital space crossed by a broad brown bar with blackish margins, which become much narrower below and traverse the eye and the cheek; behind this a broader yellow bar, margined behind with a narrow brown line. In life the coloration is extremely brilliant, the pale markings being bright orange or scarlet. Bering Sea and Arctic Ocean, from Greenland to the Kurils; locally abundant; numerous fine large specimens taken from the stomachs of cormorants on St. Paul Island, Pribilof Group; others dredged in shallow waters. Our specimens from St. Paul, Bristol Bay, and Upernavik, Greenland. Three large specimens from St. Paul Island, the type locality of *P. maxillaris*, have been compared with a number of individuals of *P. fasciatus* from Upernavik, Greenland. We can appreciate no differences between the two. The size of the mouth and the length of the head are the same in specimens of equal length, and no difference exists in the development of the ventrals. The agreement seems to be perfect in the fin rays, relative proportion and coloration. Pallas's short account of *Blennius tænia* contains nothing distinctive except the number of fin rays and the statement that the body is banded. As both of these items agree with the present species, we may safely follow Bean & Bean in making the identification. In a specimen from St. Paul, 29 cm. long, the length of the maxillary is contained $2\frac{2}{3}$ times in distance from tip of snout to origin of dorsal; the mandible equals the length of the pectoral. In a younger example, 15 cm. long, from Bristol Bay, the maxillary is contained $3\frac{1}{6}$ in predorsal length; the mandible approximately equals length of pectoral. (*fasciatus*, banded.)

Centronotus fasciatus, Bloch & Schneider, Syst. Ichth., 165, pl. 37. fig. 1, 1801, Tranquebar; an error? Günther, Cat., iii, 287, 1861.

Gunnellus grœnlandicus, Cuvier & Valenciennes, Hist. Nat. Poiss., xi, 442, 1836, Greenland, after Bloch & Schneider; Reinhardt, Dansk. Vidensk. Selsk. Nat. og Mathem. Afh., vii, 122, 1838.

Gunnellus murœnoides, Valenciennes, in Cuvier, Règne Animal, Poiss., pl. 78, fig. 2, 916; after Bloch & Schneider.

Blennius tænia,* Pallas, Zoogr. Rosso-Asiat., iii, 1811, 178, Kuril Islands.

Murœnoides maxillaris, Bean, Proc. U. S. Nat. Mus. 1881, 147, St. Paul Island, Alaska (Type, No. 23999. Coll. Henry W. Elliott); Jordan & Gilbert, Synopsis, 768, 1883.

Gunnellus fasciatus, Cuvier & Valenciennes, Hist. Nat. Poiss., xi, 441, 1836.

Murœnoides fasciatus, Jordan & Gilbert, Synopsis, 767, 1883.

Murœnoides tænia, Jordan & Gilbert, Synopsis, 766, 1883.

Pholis fasciatus, Gilbert, Rept. U. S. Fish Comm. 1893, 449; Jordan & Gilbert, Rept. Fur Seal Invest., 1898.

**Pholis tænia* is thus described: Body banded; teeth obtuse, subdistinct; head subtriangular, compressed; body ensiform, covered with minute embedded scales; vent median. Dorsal fin extending from near the head to the tail, the spines subequal; caudal subdistinct; pectorals small; ventrals represented by 2 recurved spines. Body banded. D. LXXXVII; A. 47. Kuril Islands. (Pallas.)

2773. PHOLIS GUNNELLUS (Linnæus).

(GUNNEL; BUTTER FISH.)

Head from 7 to 8 in body; depth 7 to 8; D. LXXVI to LXXXV; A. II, 38 to 44; V. I, 1; eye 5 in head; maxillary 3; P. 2; C. $1\frac{1}{3}$. Head compressed, naked; mouth oblique, the maxillary reaching to front of pupil; teeth blunt, in a single row, somewhat enlarged anteriorly; interorbital a narrow ridge about $\frac{1}{2}$ eye. Distance from origin of dorsal to nape equal to distance from nape to middle of eye; pectoral rather large, about 2 in head, inserted directly under front of dorsal. Color olive brown, sides with numerous obscure darker bars; base of dorsal with blackish spots, generally bordered with a narrow yellow line, a dark bar running downward and backward from eye; anal with dusky bars across the rays. This species differs from *Pholis ornatus* in the more numerous fin rays and in coloration; the spots on dorsal are black, edged with yellow; in *P. ornatus* they are yellow with a black bar before and behind, each partly encircling it; no black bordered light streak from eye to occiput. Length 12 inches. North Atlantic, from Labrador south to Woods Hole and Norway to France; abundant on rocky shores among algæ, both in America and Europe. Here described from specimens from Salem, Massachusetts. (Eu.) (*grunnellus*, English gunnel, said to be corrupted from gunwale.)

Blennius pinna dorsalis ocellis X nigris, LINNÆUS, Mus. Adolph-Fred., I, 69.

Blennius gunnellus, LINNÆUS, Syst. Nat., Ed. X, 257, 1758, Atlantic Ocean; after *Blennius pinna dorsalis*, etc.

Ophidion imberbe, LINNÆUS, Syst. Nat., Ed. X, 259, 1758, Europe; after *Oph. cirris careus*, ARTEDI.

Centronotus gunnellus, BLOCH & SCHNEIDER, Syst. Ichth., 167, 1801; GÜNTHER, Cat., III, 285, after RÜPPELL'S type.

Muræoides gunnellus, JORDAN & GILBERT, Synopsis, 767, 1883.

Blennius europæus, OLAFSEN, Reisei Island, I, 81, 1772, Iceland.

Blennius muræoides, SUJEF, Act. Petrop. II, 1779, 195, no locality, probably the Baltic; GMELIN, Syst. Nat., 1184, 1788.

Muræoides sujef, LACÉPÈDE, Hist. Nat. Poiss., II, 324, 1800; after SUJEF.

Ophidium mucronatum, MITCHILL, Trans. Lit. & Phil. Soc. N. Y., II, 1815, 361, pl. 1, f. 1, New York; earliest American name.

Gunellus vulgaris, FLEMING, British Anim., 207, 1828, England.

Muræoides guttatus, LACÉPÈDE, Hist. Nat. Poiss., II, 324, 1800; YARRELL, Brit. Fish., I, 269.

Gunellus ingens, H. R. STORER, Bost. Journ. Nat. Hist., VI, 1850, 261, pl. 8, f. 1, Labrador. (Coll. H. R. Storer.)

Gunellus macrocephalus, GIRARD, in H. R. STORER, Bost. Journ. Nat. Hist., VI, 1850, 263, Chelsea Beach, Massachusetts (Coll. Chas. Girard); D. H. STORER, Rept. Fish. Mass., 261, pl. 17, f. 3.

2774. PHOLIS ORNATUS (Girard).

Head 8; depth 8. D. LXXVII to LXXIX; A. II, 35 to 37. Head naked, very narrow above; nape nearly equidistant between origin of dorsal and front of orbit; origin of anal equidistant between base of caudal and base of pectoral; pectoral 2 in head. Coloration, usually olive green above, yellow or orange below,* but varying with the surroundings to brown

* A specimen from near Seattle varies much in color from all the others before us. It is purplish red, paler below; 2 conspicuous white spots bordered with white on front of dorsal; a pale streak bordered with black from eye to nape.

and cherry red; traces of about 20 darker bars along sides; a dark bar downward from eye; fins reddish; a V-shaped mark from eye to occiput, grayish, bordered by jet-black; the common form with about 14 red spots along base of dorsal, each with a curved black bar in front and behind, partly encircling it; others with about as many broad ⌒-shaped darker blotches, which extend on the fin, the first one or two blotches often shaped as in the former case; anal white, unmarked. Length 12 inches. San Francisco to Bering Sea; very common northward, its range extending to Kamchatka;* very common at Unalaska; always in shallow water. (*ornatus*, ornamented.)

Gunnellus ornatus, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 149; GIRARD, Pac. R. R. Surv., x, Fishes, 116, pl. 25b, figs. 6 and 7, 1858 (Type, No. 490, Presidio, California, Coll. Lieut. Trowbridge; No. 491, Shoalwater Bay, Washington, Coll. Dr. J. G. Cooper; No. 492, Fort Steilacoom, Washington, Coll. Dr. Geo. Suckley).
Centronotus lætus, COPE, Proc. Amer. Phil. Soc. Phila. 1873, 27, Sitka or Unalaska (Coll. George Davidson); A. II, 33.
Murænoides ornatus, JORDAN & GILBERT, Synopsis, 767, 1883.
Pholis ornatus, GILBERT, Rept. U. S. Fish Comm. 1893, 450.

906. GUNNELLOPS, Bleeker.

Gunnellops, BLEEKER, Versl. Ak. Amst., 2, VIII, 1874, 368 (*roseus*).

This genus is apparently distinguished from *Pholis* by the tapering tail, around which the vertical fins are confluent; palatine teeth present. (*Gunnellus*, Gunnel, an old name of *Pholis gunnellus*; ὤψ, appearance.)

2775. GUNNELLOPS ROSEUS (Pallas).

D. ca. C; A. ca. 90; P. 9; V. I. Head obtuse, the lower jaw projecting; eyes large; body very long, compressed, tapering into a slender tail; pectorals small, ovate, hyaline; 2 spines in place of ventrals; dorsal extending from the nape to the end of the tail; anal joined to caudal. Color intensely red. Kuril Islands. (Pallas.) Not seen by any recent collector. (*roseus*, rosy.)

Blennius roseus, PALLAS, Zoogr. Rosso-Asiat., III, 177, 1811, Kuril Islands.
Centronotus roseus, GÜNTHER, Cat., III, 290, 1861.
Gunnellops roseus, JORDAN & EVERMANN, Check-List Fishes N. and M. A., 474, 1896.

907. ASTERNOPTERYX, Rüppell.

Asternopteryx (RÜPPELL MS.) GÜNTHER, Cat. Fishes Brit. Mus., III, 288, 1861, name only; JORDAN & GILBERT, Synopsis, 769, 1883 (*gunelliformis*).

This genus is closely allied to *Pholis*, differing chiefly in the entire absence of ventral fins. From *Pholidapus* it is distinguished by the shorter pectorals and by the more broadly united gill membranes. Greenland. A single species known. (ἀ-, without; στέρνον, breast; πτέρυξ, fin.)

2776. ASTERNOPTERYX GUNELLIFORMIS, Rüppell.

Head 9; depth 8½. D. LXXXVII (LXXXI, according to Günther); A. II, 40. Head and body strongly compressed; head bluntish, snout short,

*We have specimens collected at Tareinsky Bay by Mr. Barrett-Hamilton.

jaws equal; maxillary reaching pupil, 3 in head; eye 5½; gill membranes broadly united, their outline not notched; no trace of ventral fins; pectoral large, 2 in head (3 according to Günther). Dorsal and anal joined to the caudal, the anal with a slight notch behind the last ray; dorsal spines short and all pungent. Color dark brown, clouded with darker; about 11 quadrate pale areas along dorsal fin extending on the sides, these areas each with a black central spot at tip and faintly marked with dark blotches; dorsal with dark spots; a dark band from eye downward, a pale band behind it; lips dark; anal fin bright orange; pectorals and gill membranes pale orange; caudal orange. Greenland. Here described from a fine specimen, 9½ inches long, in the U. S. National Museum, from Omanak Fjord, Karsak, Noursoak Peninsula, taken in 1897 by Schuchert and White; only the original type in the Senckenburg Museum hitherto known. (*Gunellus; forma*, shape.)

Asternopteryx gunelliformis, RÜPPELL MS.: type (in Senckenburg Museum) from Greenland.

Centronotus gunelliformis, GÜNTHER, Cat., III, 288, 1861.

Murænoides gunelliformis, JORDAN & GILBERT, Synopsis, 769, 1883.

908. ANOPLARCHUS, Gill.

Anoplarchus, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 261 (*atropurpureus*).

Body elongated, compressed, covered with very small, embedded scales which are obsolete or concealed anteriorly; lateral line obsolete. Head small, compressed; eyes small; mouth oblique; teeth in each jaw in a narrow band, the outer somewhat enlarged; narrow bands of teeth on vomer and palatines; gill membranes attached to the isthmus; sometimes with a free fold behind; branchiostegals 5. Dorsal fin not very low; no anal spines; ventrals wanting; caudal fin small, entire; pectoral fins moderate or small; pyloric cæca present, few. Pacific. (ἄνοπλος, unarmed; ἄρχος, anus; the anal fin being without spines.)

a. Gill membranes narrowly joined to the isthmus, with a free fold behind; dorsal with about 63 spines. ALECTROLOPHUS, 2777.

aa. Gill membranes broadly joined to the isthmus, without free fold behind; dorsal with 54 to 57 spines. ATROPURPUREUS, 2778.

2777. ANOPLARCHUS ALECTROLOPHUS (Pallas).

Head 6⅜ in length; depth 7⅗. D. LXII or LXIII; A. 43. Mouth oblique, maxillary reaching vertical behind pupil, 2¼ in head. Teeth in narrow bands on the jaws, the outer series in upper jaw somewhat enlarged; vomer and palatines with narrow bands of teeth; dentition similar to that in *A. atropurpureus*, which has been erroneously described as having the teeth in the jaws in single series and the vomer and palate toothless; gill membranes rather narrowly joined to the isthmus and with a free posterior edge slightly wider than pupil. *A. atropurpureus* has the gill opening somewhat more restricted and the gill membranes without free fold. Large pores on head arranged similarly in the two species. Spinous dorsal beginning slightly in advance of base of pectoral, its distance

from snout less than length of head; distance from origin of anal to tip of snout $2\frac{2}{5}$ in length to base of caudal; pectoral short and broad, rounded, $2\frac{1}{6}$ in head. Scales small, embedded, those on the anterior part of the body concealed by the thickened integument, as in *A. atropurpureus*. Coloration in our specimens nearly uniform dark olive, with obscure dusky mottlings on the side. In 1 specimen there is a light bar extending obliquely downward and backward from eye, with a dark bar above and below it, the 3 separated by narrow light gray lines; caudal narrowly cross-banded with light and dark as in *A. atropurpureus*, and the anal obliquely barred with the same. In the smallest specimen is a series of roundish spots about as large as eye along back just below dorsal fin; each spot seems to have a narrow dark margin, a light ring, a dusky ring, and a light center; a series of similar but smaller spots along middle of sides posteriorly; the colors were probably brighter and more varied in life. Western part of Bering Sea and Sea of Okhotsk. Here described from 3 small specimens, $3\frac{1}{8}$ to 9 inches long, taken at Tareinsky Bay, Kamchatka, by Mr. Barrett-Hamilton; 2 other fine specimens since taken by Arthur W. Greeley in Monterey Bay; the only ones recorded since Pallas. They differ from specimens of *A. atropurpureus* in the higher crest, the more numerous fin rays, and in having the gill membranes with a distinct free margin. (*ἀλέκτωρ*, cock; *λόφος*, crest.)

Blennius alectrolophus, Pallas, Zoogr. Rosso-Asiat., III, 174, 1811, Island of Talek, Gulf of Penshin, Okhotsk Sea.
Gunnellus alectrolophus, Cuvier & Valenciennes, Hist. Nat. Poiss., XI, 447, 1836.
Centronotus alectrolophus, Günther, Cat., III, 289, 1861.
Anoplarchus alectrolophus, Jordan & Gilbert, Rept. Fur Seal Invest., 1898.

2778. ANOPLARCHUS ATROPURPUREUS (Kittlitz).

Head $6\frac{1}{2}$ in body; depth 7. D. LV; A. 40; eye $5\frac{1}{3}$ in head; maxillary $2\frac{1}{2}$; pectoral $2\frac{1}{2}$; caudal $1\frac{3}{4}$. Head with a fleshy crest, which rests on a ridge of bone, its height in older examples about equal to eye; mouth rather large, the maxillary reaching beyond the orbit. Dorsal and anal comparatively high, barely connected with the base of caudal; nape midway between origin of dorsal and pupil. Body naked anteriorly, scaled behind. Color grayish olive, varying to brown; everywhere above finely marked with blackish reticulations; along each side of back a series of small, irregular, sharply defined grayish spots; a series of small pale spots along lateral line; belly pale; crest and middle line of back rather pale; under parts of head yellowish; an oblique, wedge-shaped, pale streak extending downward and backward from the eye, bounded on each side by a sharp light-red line, and then by a dusky area; lower jaw mottled; dorsal olivaceous, speckled, a blackish spot on front; anal olive, tinged with red; pectorals dull orange, barred at base; caudal reddish, with narrow pale streaks, and a light bar at base; color sometimes nearly plain purplish, but more often grayish and mottled. Alaska to San Francisco; abundant northward; common in Bering Sea. Here described from specimens from Neah Bay, Straits of Fuca, Washington. We have also specimens from the Pribilof Islands. (*ater*, black; *purpureus*, purple.)

Ophidium atropurpureum, KITTLITZ, Denkwürd einer Reise Russ.-Amer., I, 225, 1858, Alaska.

Centronotus cristagalli, GÜNTHER, Cat., III, 289, 1861, Vancouver Island.

Anoplarchus purpurescens, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 261, Washington Territory. (Coll. Dr. Kennerly.)

Anoplarchus cristagalli, GÜNTHER, Cat., III, 564, 1861.

Anoplarchus atropurpureus, JORDAN & GILBERT, Synopsis, 771, 1883; JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 846.

909. XIPHISTES, Jordan & Starks.

Xiphistes, JORDAN & STARKS, Proc. Cal. Ac. Sci. 1895, 846 (*chirus*).

This genus is very close to *Xiphidion*, differing in the well-developed pectoral fins, which are longer than eye; lower lateral line not connected with abdominal line. (ξιφιστής, a sword belt.)

a. Anal spines 3; branches of upper lateral line extending on dorsal fin; color grass-green. ULVÆ, 2779.

aa. Anal spines 2; branches of upper lateral line shorter; color brownish, marbled, and with red blotches. CHIRUS, 2780.

2779. XIPHISTES ULVÆ, Jordan & Starks.

Head 8; depth 10. D. LXXIV; A. III, 48; eye 5 in head; maxillary $2\frac{3}{4}$; pectoral $3\frac{1}{2}$. Body eel-shaped, as in the related species; head short; mouth small, oblique, maxillary extending to below posterior margin of eye; jaws subequal, with canine teeth; 4 enlarged canines in front of lower jaw; teeth in upper jaw gradually enlarged from behind forward; eye moderate, equal to length of snout; interorbital space prominent, sharply convex, narrower than width of eye; nape not constricted. Five mucous canals radiating downward and backward from eye, not reaching to edge of preopercle, the branches running upward from upper lateral line ending on the membrane of dorsal, the lower lateral line not connected with the abdominal line. Lateral line otherwise as in *Xiphistes chirus*. Origin of dorsal at a distance behind nape equal to distance from nape to middle of eye, the fin posteriorly barely connecting with caudal; anal with 3 spines, its origin about a head's length nearer snout than base of caudal, connected with caudal posteriorly; pectorals equal in length to snout and $\frac{1}{2}$ eye, slightly shorter than caudal; caudal rounded, fan-shaped. Color olive-green above, very bright green below; middle and lateral line posteriorly, with conspicuous white spots, $\frac{1}{2}$ as large as pupil, each with a black spot before and behind it; a black streak from tip of snout, through eye, to nape, a streak starting from eye behind quickly fading out; dorsal darker than body, unmarked; the anterior third of anal green, without markings, behind this, faint cross bars of brown appear, growing broader and darker posteriorly; caudal olive green, with a light bar across base; pectorals green, without markings. One specimen obtained at Waadda Island, Neah Bay. It was found high on the rocks, among algæ, just below high water mark. Length 5 inches. This species is very closely related to *Xiphistes chirus;* it differs from it chiefly in having 3 anal

spines, in the branches of the upper lateral line running higher, and in coloration. (*Ulva*, the green sea lettuce.)

Xiphidion ulvæ, JORDAN & STARKS, Fishes of Puget Sound, 847, 1895, Waadda Island, Neah Bay. (Type, No. 3132, L. S. Jr. Univ. Mus. Coll. E. C. Starks.)

2780. XIPHISTES CHIRUS (Jordan & Gilbert).

Head 7; depth 9. D. LXX; A. II, 50. Head short; nape not constricted; mouth small; maxillary extending to middle of pupil; teeth strong, the anterior canine-like, bluntish; about 4 canines in lower jaw, 5 or 6 in the upper, similar to the teeth behind them, but somewhat larger. Abdominal lines meeting on the breast, but not connected with the lower lateral line. Dorsal fin beginning close behind pectoral; nape midway between middle of eye and front of dorsal; anal beginning about a head's length nearer snout than base of caudal; pectoral fin comparatively large, longer than the eye, its length about equal to distance between middle and lower lateral lines. Color olive brown, yellowish below; sides with marblings of different shades of brown, sometimes with short blackish vertical bars; some round black spots along the back and sides; a black spot behind opercles; numerous black spots on sides of head, forming in older individuals light and dark streaks, which radiate from eye across cheek and opercles, the pale streaks forming reticulations; dorsal with black spots and a series of bright reddish brown cross blotches; pectorals and caudal plain. Monterey to Alaska; smaller than the other species, and living in deeper water; abundant about Cape Flattery. (Jordan & Gilbert.) (*χείρ*, hand.)

Xiphister chirus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 135, Point of Los Pinos, near Monterey, California (Coll. Jordan & Gilbert); JORDAN & GILBERT, Synopsis, 772, 1883; JORDAN & STARKS, Fishes of Puget Sound, 846, 1895.

910. XIPHIDION, Girard.

Xiphidion, GIRARD, Pac. R. R. Surv., X, Fishes, 119, 1858 (*mucosum*); not *Xiphidium*, Serv., a genus of Grasshoppers.

Xiphister, JORDAN, Proc. U. S. Nat. Mus. 1879, 241 (*mucosum*); substitute for *Xiphidion*, regarded as preoccupied by *Xiphidium*.

Body elongate, eel-shaped, covered with small scales; lateral lines several: 1 along the median line of the side, 1 above this, and 1 below it; 1 on each side of the abdomen, the 2 meeting in front, and 1 from the occiput toward the base of the dorsal fin. Each of these has on each side series of short branches, placed at right angles to the main line, those on opposite sides alternating. Each of these branches has about 2 open mucous pores. Lower lateral line connected with the abdominal line. Head short, bluntish, scaleless; mouth moderate, oblique; jaws with rather strong teeth, the anterior canine-like; no teeth on vomer or palatines. Branchiostegals 6; gill membranes separate, free from the isthmus. A single long, low, uniform dorsal fin, consisting of spines only; anal fin similar in form, with small spines, indistinct or obsolete; caudal short, joined to dorsal and anal; no ventral fins; pectoral fins very small, shorter

than eye. Intestinal canal moderately elongate, with 4 to 6 well-developed pyloric cæca. Herbivorous, feeding on algæ. Active fishes, inhabiting tide pools and crevices among rocks in the North Pacific. (*ξιφίδιον*, a small sword.)

a. Distance from origin of dorsal to occiput less than that from occiput to tip of snout; streaks radiating from eye paler in the center, edged above and below with blackish. MUCOSUM, 2781.

aa. Distance from origin of dorsal to occiput greater than that from occiput to snout; streaks radiating from eye black, abruptly margined with pale olive. RUPESTRE, 2782.

2781. XIPHIDION MUCOSUM, Girard.

Head 7 in body; depth 8¼. D. LXXIV; A. 46; eye 7½ in head; maxillary 2⅔; caudal 2⅔; pectoral a little longer than eye. Lower jaw with a series of short, stout conical teeth; upper jaw with a narrow band of similar teeth; 2 strong canines in upper jaw, 4 in the lower. Lower lateral line sending a branch to the abdominal line; nape not constricted. Dorsal beginning anteriorly, distance from its origin to occiput less than that from occiput to tip of snout; origin of anal nearer snout than tip of caudal by about ½ length of head. Blackish green, pale on belly and sides of head, marked posteriorly with olive green in various pattern; a transverse light-greenish bar at base of caudal; 3 olive-brown streaks radiating backward from eye, paler in the center and edged above and below with blackish, outside of which is sometimes a streak of pale olive; these streaks all merge backward into the color of the head; middle streak broadly wedge-shaped, the third streak terminating before reaching margin of preopercle; old individuals often coarsely blotched with yellow. Length 18 inches. Monterey to Alaska; very abundant among rocks and algæ. Here described from specimens, 9 or 10 inches in length, from Neah Bay, Straits of Fuca, Washington. (*mucosus*, slimy.)

Xiphidion mucosum, GIRARD, Pac. R. R. Surv., X, Fishes, 119, 1858, South Farallones, California (Coll. Lieut. Trowbridge. Type, No. 493, U. S. Nat. Mus.); GÜNTHER, Cat., III, 291, 1861; JORDAN & STARKS, Fishes Puget Sound, 848, 1895.

Xiphidion cruoreum,* COPE, Proc. Amer. Phil. Soc. Phila. 1873, 27, Sitka (Coll. Prof. George Davidson); JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 137.

Xiphister mucosum, JORDAN & GILBERT, Synopsis, 772, 1883.

* The following is the original description of *Xiphidion cruoreum:* Head 8¼ in total length; depth 9⅓; eye 7 in head, equal to length of pectoral fin. D. about 70; A. 48; Br. 5. Teeth, 2 canines above, 4 below, subequal. Dorsal spines not commencing near the head, the anterior buried in a soft fold of skin; caudal fin not distinct. Three lateral mucous canals extending entire length of caudal fin, which have numerous alternating transverse branches, those of the superior reaching base of dorsal, those of inferior reaching base of anal; each of the cross branches with several excretory pores, none on the main stem; a similar but short tube extending from near base of dorsal fin to supra-occipital region, and not branching anteriorly; the superior lateral canal descending to near the median, but not joining it, nor does the latter extend into the inferior; another tubular line on each side of abdomen, these uniting on jugular region by a continuation of the inferior lateral tube. Vent nearer end of muzzle than end of caudal fin, by length of head. Color maroon, more reddish below; a vertical, broad, reddish bar at base of tail, beyond which is a dark spot; 2 brown radii, black-edged, extending backward and downward from eye. Body covered with small scales, except on the jugular and abdominal regions, which are naked. Length 8 inches. This fish is not very different from *X. mucosum*, Girard. It differs in the smaller eye, the more remote origin of the dorsal fin from the head, the lack of anterior union of the mucous canals, and the coloration. (Cope.)

2782. XIPHIDION RUPESTRE (Jordan & Gilbert).

Head 7⅓ in body; depth 9. D. LXVIII; A. 50; eye 6 in head; maxillary 2⅓; caudal 2⅘. Teeth essentially as in *X. mucosum*. The lower lateral line sends a branch to the abdominal line; a constriction at the nape. Distance from origin of dorsal to the occiput greater than the distance from the occiput to the snout. Anal fin beginning much in advance of middle of body, the distance from the first ray to tip of caudal exceeding the distance to snout by nearly twice length of head; pectoral very short, its length less than diameter of eye. Reddish brown, uniform or variously shaded with lighter; a light olivaceous bar at base of caudal, extending on dorsal and anal, behind this a blackish area; tip of caudal usually pale; 3 long, well-defined stripes radiating backward from eye, these stripes uniform black, abruptly margined with very light olive; the central stripe proceeding straight backward from the eye, ½ breadth of cheek, at which point it is broadest; it is then narrowed and bent abruptly downward; both the middle and lower stripes reach the margin of preopercle. Length 12 inches. Smaller than the preceding, and equally abundant; among rocks and algæ, from Vancouver Island to Monterey. Here described from specimens, 6 or 7 inches in length, from Neah Bay, Straits of Fuca, Washington. (*rupestris*, living among rocks.)

Xiphister rupestris, Jordan & Gilbert, Proc. U. S. Nat. Mus. 1880, 137, Monterey Bay, California (Coll. Jordan & Gilbert); Jordan & Gilbert, Synopsis, 773, 1883.
Xiphidion rupestre, Jordan & Starks, Fishes Puget Sound, 848, 1895.

911. CEBEDICHTHYS,* Ayres.

Cebedichthys, Ayres, Proc. Cal. Ac. Nat. Sci., I, 1855, 59 (*violaceus*).

Body comparatively short, compressed, covered with minute scales; lateral line distinct, running very high, with very short branches, each ending in a pore, as in *Xiphistes*, but the branches more oblique and less regular. Head short; crown with a conspicuous fleshy longitudinal crest in the adult; jaws subequal, with conical teeth; villiform teeth on vomer and palatines; gill membranes united, free from the isthmus. Dorsal fin continuous, long and low, the anterior part composed of sharp spines, which are rather lower than the soft rays; caudal fin rounded, connected with dorsal and anal; anal fin similar to soft dorsal, with 1 or 2 small spines; pectorals small; ventrals wanting. Intestinal canal elongate, with several pyloric cæca. Pacific Ocean. Herbivorous; similar in habits

* The following remarkable genus may be allied to *Cebedichthys:*

NEOZOARCES, Steindachner.

Neozoarces, Steindachner, Ichth. Beitr., IX, 26, 1880 (*pulcher*).

Neozoarces pulcher, Steindachner.

Body elongate, tapering backward, the dorsal and anal united at the tail without distinct caudal. Scales small, embedded, no lateral line. Mouth very large, the maxillary extending far beyond eye; lower jaw slightly longer than upper; blunt, conical teeth in many rows on jaws, vomer, and palatines. A thick tentacle above nostril; gill membranes united, free from isthmus. Dorsal low, the anterior portion of short, stiffish spines; no anal spine; ventrals wanting; pectorals moderate; pseudobranchiæ present. Head 6; depth 9. D. XLI, 50; A. I, 75. Color highly variegated. Gulf of Strietok, Okhotsk Sea. (νέος, new; *Zoarces*; but it has little affinity with the latter genus.)

Neozoarces pulcher, Steindachner, Ichth. Beitr., IX, 27, taf. 6, f. 2, 1880, Gulf of Strietok. (Coll. Professor Dybowski.)

to the species of *Xiphidion*. (*κῆβος*, the Sapajou, a kind of monkey; *ἰχθύς*, fish; in allusion to the "peculiar monkey-like" physiognomy as seen from the front.)

2783. CEBEDICHTHYS VIOLACEUS (Ayres).

Head 6½; depth 6. D. XXIII, 41; A. I, 41. Maxillary extending to or beyond orbit. Dorsal scaly at base; vent nearer snout than base of caudal; pectoral ⅔ length of head; nape midway between dorsal and eye. Dull olive grayish, mottled with lighter, sometimes reddish tinged; vertical fins all edged with reddish; cheek with 3 darker stripes, edged with paler, 1 downward and backward from the eye, close behind angle of mouth; another above it to root of pectoral; another running upward and backward from the eye, and meeting its fellow over the crest. Length 30 inches. San Francisco to Point Concepcion; abundant; often brought into the markets. (*violaceus*, violet.)

Apodichthys violaceus, Girard, Proc. Ac. Nat. Sci. Phila. 1854, 150, San Luis Obispo, California. (Coll. Dr. Kennerly.)

Cebedichthys cristagalli, Ayres, Proc. Cal. Ac. Nat. Sci., I, 1855, 58, San Francisco.

Cebedichthys violaceus, Girard, Pac. R. R. Surv., X, Fishes, 121, pl. 26, figs. 4 and 5, 1858; Jordan & Gilbert, Synopsis, 774, 1883.

912. PLAGIOGRAMMUS, Bean.

Plagiogrammus, Bean, Proc. U. S. Nat. Mus. 1893, 699 (*hopkinsi*).

Body moderately elongate, compressed, covered with very small scales; lateral lines 2, viz, 1 beginning above and slightly in advance of the upper angle of the gill opening and extending along the upper part of the body, but not reaching the tail, and 1 beginning in advance of the end of this and reaching to the caudal; numerous lateral ridges on the sides, similar to those on *Dictyosoma* of Temminck & Schlegel; a series of subpentagonal plate-like bodies along the abdominal edge on each side between the ventral and the anal. Head moderately long, naked, with pointed snout; mouth oblique and rather large; jaws subequal, or the lower slightly projecting; jaws with strong teeth in broad bands, the intermaxillaries with an outer series of enlarged canine-like teeth; teeth on vomer and palate; a pair of large canines near the symphysis in each jaw, the canines of the upper jaw fitting into an interspace behind the mandibulary canines. A series of pores on the ramus of the mandibula continuing around the preopercular edge; a series of similar pores along the lower margin of the preorbital continued backward and upward toward the nape. Anterior nostril tubular; posterior without tube. Maxillary broadly expanded posteriorly; lips well developed. Branchiostegals 5; gill membranes partly united, but free from the isthmus behind. Gill rakers minute, tubercular, in moderate number. A single long dorsal fin consisting of spines only, the spines longest in the posterior portion; anal fin lower than the dorsal, but similar in shape. Pectoral large, entirely below median line. Ventrals well developed, in advance of pectorals; caudal rounded, distinct. Intestinal canal short, with 5 small pyloric cæca. (*πλάγιος*, oblique; *γραμμή*, line.)

2784. PLAGIOGRAMMUS HOPKINSI, Bean.

Head 4; depth 5⅓; eye 5. D. XLI; A. II, 29; V. I, 5; B. 5; scales about 95; ridges on side 32. Snout acute; anterior nostril tubular and nearer eye than tip of snout; posterior nostril close to upper anterior margin of eye; maxillary extending almost to vertical through hind margin of eye; intermaxillary long, slender, and reaching nearly as far back as maxillary; intermaxillary teeth in broad bands, with an outer series of 5 or 6 large canines, those near the symphysis largest; teeth in mandible in broad bands in front, followed by several enlarged canine-like teeth; a large canine on each side of symphysis, the interspace between the 2 mandibulary canines receiving the canines of the intermaxillary when the jaws are closed. A row of 8 pores along ramus of mandible and edge of preopercle; another series around lower margin of preorbital bone as described for the genus; about 8 gill rakers on first arch below angle. Distance of dorsal origin from snout nearly equal to length of head; spines lowest in front, the longest spine ⅔ length of head; longest rays of anal near end of fin and scarcely exceed length of eye; length of pectoral equaling that of postorbital part of head; ventrals close together; inner rays longest, ⅔ as long as head; caudal rounded, its length nearly ½ that of head; vent under eleventh spine of dorsal. Upper lateral line beginning above and slightly in advance of upper angle of gill opening, curving very slightly over pectoral and extending to below twenty-fifth spine of dorsal, its distance from dorsal edge equal to diameter of eye and also equal to its distance from lower lateral line; lower lateral line beginning under sixteenth spine of dorsal and extending to caudal. On each side of the abdominal ridge, between the ventrals and the vent, are about 10 subpentagonal plate-like bodies, the largest about ½ as long as eye. Color dusky brown, the fins black. Monterey, California; a few specimens dredged among rocks. Little is known about the habits of the species, beyond the fact that in the aquarium it hides in rock crevices and seldom ventures from its hiding place. (I take pleasure in associating with this blenny the name of Mr. Timothy Hopkins, of Menlo Park, California, the founder of the Seaside Laboratory at Pacific Grove, Monterey Bay, in commemoration of his services in behalf of science. Bean.)

Plagiogrammus hopkinsi, Bean, Proc. U. S. Nat. Mus. 1893, 699, Monterey Bay, California. (Type, No. 44721, U. S. Nat. Mus.)

913. OPISTHOCENTRUS, Kner.

Opisthocentrus, Kner, Sitzber. Akad. Wiss. Wien 1868, 49 (*quinquemaculatus*).
Blenniophidium, Boulenger, Proc. Zool. Soc. Lond. 1892, 583 (*petropauli*).

Body moderately elongate, compressed, covered with very small cycloid scales. Mouth small, horizontal, protractile, with fleshy lips; small conical teeth in jaws and on vomer and palatines. No cirri. Gill membranes broadly connected, but free from isthmus; branchiostegals 4. Dorsal fin very long, extending from the nape to the caudal, with which it is subcontinuous; a few of the posterior rays are stiff spines, the rest being

simple and not articulate, but flexible; anal fin extending from the anus, which is a little nearer the anterior than the posterior extremity, to the caudal, formed exclusively of soft rays; no ventrals. No lateral line. No prominent anal papillæ. Pyloric appendages present. A remarkable genus, allied to *Lumpenus*, or rather to *Plectobranchus*, distinguished by having only the posterior spines rigid. North Pacific. (ὄπισθε, behind; κέντρον, spine.)

2785. OPISTHOCENTRUS OCELLATUS (Tilesius).

Head 6¼; depth 6¼ (without caudal). D. LV to LXI, usually LIX; A. 36 to 39; 5 to 7 of the posterior dorsal spines rigid; Eye as long as snout, 4 in head, and a little more than interorbital width; maxillary extending to below anterior fourth of eye; some wide pores on the head; cheeks, opercles, and occiput closely scaled; strips of small scales on the branchiostegal membrane between the rays. Dorsal rays continuous and subequal in depth, the longest spine 2⅕ in head in females, 1⅙ in males; pectoral 1⅓ in head, about as long as caudal. Anus twice as far from caudal as from base of pectoral. Yellowish brown, with ill-defined darker marblings; a crescentic black line on the top of the head from eye to eye; a black line, obliquely directed forward, below the eye, and another, in opposite direction, from the eye to the opercle; 2 dark-brown streaks across the nape, the second crossing the origin of the dorsal fin and extending to the base of the pectoral; dorsal and caudal fins grayish olive, lighter at the base, the dorsal with 5 to 9 (usually 6) large black spots at regular intervals, these wanting in the males; pectorals and anal colorless. Numerous specimens are from Tareinsky Bay, Kamchatka; Petropaulski Harbor, and Shana Bay, Iturup Island. The number of dorsal ocelli varies from 5 to 9 in our specimens, 6 being the prevailing number. Of 24 specimens whose fins we have enumerated, 4 have 58 dorsal spines, 10 have 59, 5 have 60, and 5 have 61. In addition, 1 specimen has but 55 spines. The latter is the only male in the collection and is conspicuous by the absence of distinct dorsal ocelli and the great height of the vertical fins, the longest dorsal spine exceeding the length of the pectoral and contained 1⅙ times in head. In females the longest spine is 2⅓ in head. The anal contains 36 to 39 rays in all our specimens. The dorsal fin is composed exclusively of spines, the anterior flexible ones passing into the strong pungent ones near the posterior end. The stronger spines vary from 7 to 12 in number in our specimens. Our material answers the description of the type of *O. quinquemaculatus* which had 57 dorsal spines and 36 anal rays. It also agrees with specimens from Petropaulski, reported on by Bean & Bean (Proc. U. S. Nat. Mus. 1896, 391), with dorsal spines 58 in number. *Blenniophidium petropauli*, Boulenger, has but 52 dorsal spines, but it is otherwise not to be distinguished from *O. ocellatus*. Still more aberrant are 4 specimens from Gulf of Strietok, northern Japan, mentioned by Steindachner (Ichth. Beiträge, IX, 25), with but 50 to 53 spines and 32 to 34 anal rays. These may represent a distinct species. *Ophidium ocellatum* of Tilesius must be this species, but the count of fin rays is incorrect and may be taken from

the rough figure. *Opisthocentrus tenuis* is probably also identical with *O. ocellatus*, though the writers did not think so until after examination of the present large material. Coast of Kamchatka, southwestward to Okhotsk Sea, generally common from Komandorski Islands to Yezo. (*ocellatus*, with eye-like spots.)

Ophidium ocellatum, TILESIUS, Mém. Ac. St. Petersb., II, 1811, 237, Kamchatka. D. 80; A. 50; evidently an error. The rude figure shows D. 73; A. 50, the spines low; the dorsal with 5 ocelli.

Centronotus (Opisthocentrus) quinquemaculatus, KNER, Sitzber. Akad. Wiss. Wein 1868. 48, taf. 7, f. 20, "Pinang." Described from a young specimen 2 inches long, No. 6353, Mus. Wien.

Gunnellus apos, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XIV, 426, 1839; after TILESIUS.

Centronotus apus, GÜNTHER, Cat., III, 288, 1861.

Blenniophidium petropauli, BOULENGER, Proc. Zool. Soc. London 1892, 584, with plate, Petropaulski (Coll. George Baden-Powell); D. 52; A. 37; 5 ocelli.

Opisthocentrus tenuis,* BEAN & BEAN, Proc. U. S. Nat. Mus. 1897, Volcano Bay, Port Morusan, Japan. (Coll. Col. Nicolai A. Grebnitski. Type, No. 47565, U. S. Nat. Mus.)

Opisthocentrus quinquemaculatus, STEINDACHNER, Ichth. Beitr., IX, 25, 1880; BEAN & BEAN, Proc. U. S. Nat. Mus. 1896, 381, 392.

Opisthocentrus ocellatus, JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

914. PHOLIDAPUS, Bean & Bean.

Pholidapus, BEAN & BEAN, Proc. U. S. Nat. Mus. 1896, 389 (*grebnitskii*).

Body moderately elongate, compressed, covered with very small, smooth scales. Mouth small, horizontal; bands of small teeth on jaws and vomer, none on palatines. Head naked; gill membranes broadly connected, free from the isthmus; dorsal very long, composed entirely of flexible spines; anal of soft rays; caudal short, rounded, separate; no ventral fins; no lateral line; pyloric cæca present. This genus is close to *Opisthocentrus*, but has no pungent spines, and the head is naked. Okhotsk Sea. (φολίς, *Pholis*; ἄπους, without feet, i. e., ventral fins.)

2786. PHOLIDAPUS DYBOWSKII (Steindachner).

Head $5\frac{1}{2}$ to $6\frac{2}{3}$; depth 6 to $6\frac{1}{4}$. D. LXII or LXIII; A. II, 39. Eye $3\frac{3}{5}$ to to $4\frac{2}{3}$ in head; snout a little longer than eye; lower jaw scarcely included; 1 or 2 strong conical teeth on each side behind the narrow premaxillary band of teeth; teeth on vomer, none on palatines; no cirri; large pores

* *Opisthocentrus tenuis* is thus described:

D. 39, XV; A. 38. Length of fish to caudal base $5\frac{1}{8}$ inches; length of head 1; depth of body $\frac{1}{6}$; the greatest width of the body is contained $2\frac{1}{3}$ times in the length of the head. The diameter of the eye is nearly equal to the length of the snout and is contained $4\frac{1}{2}$ times in the length of the head; the width of the interorbital space is almost equal to the long diameter of eye. The maxilla reaches to the vertical past front of eye. Teeth bluntly rounded, embedded in flesh; vomerine teeth present; palatines none. The origin of the dorsal fin is over the end of the gill cover, its first 39 rays are simple and flexible, the last 15 are strong spines and end slightly above the membrane in stiff points, the longest spine is almost $\frac{1}{2}$ as long as the head. The anal originates under the twentieth ray of the dorsal; its rays are divided and articulated; the longest ray is $\frac{1}{2}$ as long as the head. The general color is brown with cross reticulations of black. Sides of head and body along base of anal, orange; anal, caudal, and pectorals light with dusky shadings; dorsal finely mottled with black and bearing 6 black spots on areas of white, the first of these spots being on the sixth ray and the last on the next to last spine; a black bar from front of eye downward, and another from posterior margin obliquely down and backward. This species differs from the typical form in its greater compression of the body and its increased number of dorsal spines. (Bean & Bean.)

about eye and on opercles; longest dorsal spines 2½ to 3 in depth of body, last spines shorter and stiffer than others; dorsal and anal slightly joined to caudal; pectoral as long as caudal, about 1¼ in head. Head naked. Brown or grayish, with faint spots or marblings; 1 or 2, rarely 3, dark ocelli on the dorsal; 3 or 4 dark streaks radiating from eye, the uppermost joining its fellow. Length 10 to 15 inches. Coast of northern Japan and sea of Okhotsk, north to the Kuril Islands. Our specimens, 5 in number, the largest 25 cm. long, from Shana Bay, Iturup Island. Steindachner's excellent and detailed description leaves nothing to be desired, and corresponds perfectly with our material except in the character of the scales. A careful examination of these under high magnification fails to show that they are "am hinteren Rande mit kurzen Zähnchen bewaffnet." The posterior border is entire and the scales strongly marked with concentric striæ. Dorsal spines number 62, 63, 63, 64, 64. Dorsal ocelli present in all our specimens, 2 of them being faintly visible even in the youngest, 55 mm. long. *Pholidapus grebnitskii* seems to differ only in the shorter dorsal fin (57 spines). (Named for Professor Dybowski, its first collector.)

Centronotus dybowskii, STEINDACHNER, Ichth. Beiträge, IX, 22, 1880, Gulf of Strietok, northern Japan (Coll. Prof. Dybowski); JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

?*Pholidapus grebnitskii*,* BEAN & BEAN, Proc. U. S. Nat. Mus. 1896, pl. 34, 390, Yezo, Japan. (Coll. Col. Nicolai A. Grebnitski.)

915. PLECTOBRANCHUS, Gilbert.

Plectobranchus, GILBERT, Proc. U. S. Nat. Mus. 1890, 102 (*evides*).

Teeth conic, on jaws, vomer, and palatines, some of them canine-like. Body scaly; lateral line obsolete, its course indicated by a lighter streak on middle of sides. Gill slits not continued far forward, the membranes

**Pholidapus grebnitskii*, Bean & Bean, is thus described: The specimens are 141 mm. long, including caudal; 126 mm. to base of caudal. The head (22 mm.) is equal to the greatest depth of body. The eye is slightly longer than the snout and ¼ as long as the head. The interorbital space is narrow, ⅓ of the length of the eye. The naked head resembles that of *Pholis*, its length is contained about 5½ times in total length without the caudal. The mouth is small and very oblique; the mandible is slightly included and has a well developed lip. The maxilla is partly concealed under the preorbital bone; it does not quite reach to below the anterior margin of the pupil. The anterior nostril is midway between the eye and the tip of the intermaxilla. Seven mucous pores around the orbit: 3 on the preorbital bone. The pore in the origin of the semicircular dark band around the nape is continued backward by a series of 6 similar ones ending near the upper angle of the gill opening. A series of 10 or 11 pores beginning near the front of the chin on each side, extending backward and curving upward to the upper anterior edge of the operculum. The gill membranes are broadly united, but they are not joined to the isthmus. The dorsal origin is over the end of the head; the fin is low, and consists of spines, the longest and strongest in the posterior third being slightly longer than the eye. The distance of the vent from the tip of the snout contains the head length 2⅔ times. The anal is slightly lower than the dorsal, the rays longest posteriorly. The caudal is rounded, and is barely separated from the dorsal and anal. The pectoral base is broad, and the fin is ⅔ as long as the head. The intestine is slender, and is more than twice as long as the head. Stomach short, pear-shaped, with 6 slender pyloric cæca of unequal length, the longest about twice as long as the eye. The body is completely scaled, the scales very small, cycloid, closely imbricated, with numerous concentric striæ, and they extend halfway up the membrane connecting the dorsal spines. The general body color is brown, the sides sparsely and vaguely mottled. The pectorals are pale. A narrow, dark band extends from the middle of the eye downward and forward, a similar band running backward from the eye on the preopercle; an interrupted semicircular band from eye to eye across the nape. D. LVII; A. II, 39 or 40. (The species is named for Mr. N. Grebnitski, to whose industry and zeal the Museum is indebted for many valuable collections. Bean & Bean.)

broadly united, wholly free from isthmus. Dorsal of spines only. Anal with 2 spines. Ventral with 1 spine and 3 well-developed rays. Lower pectoral rays longest, as in *Leptoclinus*. North Pacific. (*πλεκτός*, enfolded; *βράγχος*, gill.)

2787. PLECTOBRANCHUS EVIDES, Gilbert.

Head rather long, $4\frac{3}{4}$ in length, extending well beyond origin of dorsal fin; depth about 11. D. LVI; A. II, 34. Body very slender, the depth nearly constant throughout. Caudal peduncle without free portion, its depth $2\frac{1}{2}$ in that of body. Upper jaw with a broad inner band of minute teeth in front and on the sides, the outer series enlarged, 2 in the front of the jaw distinctly canine-like; teeth in the lower jaw similar to the outer series above, in a single series laterally, forming a patch in front of jaw, where 2 of them are much enlarged canines, the largest teeth in the jaws; vomer and palatines with bands of small but very evident teeth. Eyes large, close together, the interorbital space $\frac{1}{2}$ pupil. Orbit $3\frac{1}{3}$ in head, longer than snout. Posterior nostril with a short flap, the tube obsolete. Mouth large, somewhat oblique, maxillary reaching middle of orbit, $2\frac{2}{3}$ in head. Top of head with very large pores, a series running backward from each eye, the two joined by a cross series on occiput. Body covered with very small cycloid scales, including belly, nape, breast, and cheeks, those on breast and cheeks not imbricated; lateral line without visible pores. Spinous dorsal beginning well forward, the distance from its origin to nape less than from latter to posterior margin of orbit. Anterior spines short, but fully united by membrane, the longest spine $3\frac{1}{2}$ in head; membrane of last spine reaching base of upper caudal rays; origin of anal very slightly in advance of middle of body; anal with 2 short, sharp spines, the rays longer, their terminal $\frac{1}{3}$ free from membrane; last anal ray connected with base of lower caudal ray; ventrals well developed, nearly $\frac{1}{2}$ head; pectorals with lowermost rays abruptly lengthened, $\frac{2}{3}$ head; caudal short, rounded, little more than $\frac{1}{2}$ head. Color dusky olive above, lighter below; sides crossed by about 25 narrow white bars, narrower than interspaces; 3 equidistant dark blotches near back, each double, the two halves occupying contiguous interspaces between white bars; branchiostegal membrane black; head without markings; pectorals white at base, the distal half black, margined with white; ventrals white; dorsal with alternating oblique bars of white and blackish, 2 jet-black roundish spots on its posterior portion; caudal whitish at base, then dusky, margined with white, its upper ray jet-black; anal light at base, becoming black at edge of membrane, the free tips of rays white. Coast of Oregon. A single specimen, 4 inches long, from *Albatross* Station 3064, in 46 fathoms. (*εὐειδής*, comely.)

Plectobranchus evides, Gilbert, Proc. U. S. Nat. Mus. 1890, 102, coast of Oregon, at Albatross Station 3064. (Coll. *Albatross*.)

916. LEPTOCLINUS, Gill.

Ctenodon, Nilsson, Skandinav. Faun., IV, 190, 1853 (*maculatus*) (name three times* preoccupied).

Leptoclinus, Gill, Proc. Ac. Nat. Sci. Phila. 1864, 209 (*aculeatus*).

* *Ctenodon*, Wagler, 1830, a lizard; Ehrenberg, 1838, a rotifer, and Swainson, 1839, a fish.

Body much elongated; lateral line obsolete; teeth on jaws, vomer, and palatines; pectoral fins with the upper rays shortened; caudal fin subtruncate. Arctic seas. This genus is close to *Lumpenus*, differing mainly in the form of the pectoral. (*λεπτός*, slender; *Clinus*.)

2788. LEPTOCLINUS MACULATUS (Fries).

(LANGBARN.)

Head 5; depth 8. D. LX (LVIII to LX); A. 36 (35 to 38). Eye large, $3\frac{1}{2}$ in head; snout short and blunt, $4\frac{3}{4}$ in head, maxillary reaching past middle of eye, $2\frac{1}{3}$ in head. Teeth in jaws, vomer, and palatines; jaws each having 2 strong canines in front. Scales small, cycloid. First 3 or 4 dorsal spines short and free; longest dorsal spines as long as eye; caudal fin free from dorsal and anal; ventrals 3 in head; pectorals rather large, $1\frac{1}{2}$ in head. Color yellowish, irregularly marked with dark spots, a series of about 6 of these spots extending along sides close to base of dorsal fin; a series of smaller spots extending along center of sides from upper base of pectoral to caudal; dorsal irregularly covered with dark spots; caudal with 4 dark cross bands; anal, ventral, and pectorals plain yellowish. Bering Sea to Spitzbergen, south to Aleutian Islands and the coasts of Sweden and Norway. This description is taken from a specimen, $5\frac{1}{4}$ inches long, from Alaska, near Unimak Pass (*Albatross* Station 3309). A few young individuals of this species, hitherto known only from the North Atlantic, were taken in Unimak Pass and in Bristol Bay, in $29\frac{1}{2}$ to 70 fathoms. Three small specimens were also taken off Robben Reef, near the Kamchatka coast, in 28 fathoms, and one off Karluk, Kadiak Island. Having no Atlantic specimens of this species, we are unable to satisfy ourselves of the identity of the two, but no difference is evident from descriptions. The lateral line is much more distinct than in our specimens of *Lumpenus medius*, in which it can be made out with difficulty on scattered scales along middle of sides. (Eu.) (*maculatus*, spotted.)

Clinus maculatus, FRIES, Kgl. Vet. Ak. Handl. 49, 1837, Bohüslän, Sweden.
Lumpenus aculeatus, REINHARDT, Kong. Dansk. Vid. Selsk., VI, 1837, 190, no description.
Clinus aculeatus, REINHARDT, Dansk. Vidensk. Selsk. Natur. Afh., VII, 1838, 114, 122, 194, Spitzbergen.
Ctenodon maculatus, NILSSON, Skand. Fauna, IV, 190, 1853.
Stichæus maculatus, GÜNTHER, Cat., III, 281, 1861.
Lumpenus aculeatus, KRÖYER, Naturhist. Tidsskr., I, 377, 1862.
Stichæus aculeatus, GÜNTHER, Cat., III, 282, 1861; COLLETT, Norske Nord-Havs Exp., 67, 1880.
Lumpenus maculatus, JORDAN & GILBERT. Synopsis, 777, 1883; Lilljeborg, Sveriges Och Norges Fish., 500, 1891.
Leptoclinus maculatus, GILBERT, Rept. U. S. Fish Comm. 1893, 450.

917. POROCLINUS, Bean.

Poroclinus, BEAN, Proc. U. S. Nat. Mus. 1890, 40 (*rothrocki*).

Body elongate, moderately compressed, covered with small scales; lateral line obsolete. Head moderately long; snout short; eyes large; interorbital space narrow. Mouth small, lower jaw slightly included; teeth on vomer and palate; narrow bands of teeth in jaws, the outer series

enlarged. Gill openings slightly prolonged forward below, narrowly attached to the isthmus anteriorly. Dorsal composed of many sharp, flexible spines, diminished in length anteriorly. Caudal long, pointed; anal with 3 spines and many rays; pectorals large, the middle rays longest; ventrals jugular, with 1 spine and 3 rays. Intestine short; pyloric cæca 1 or 2; no air bladder. Northern Seas. (*πόρος*, pore; *Clinus*.)

2789. POROCLINUS ROTHROCKI, Bean.

Head $6\frac{1}{4}$ in length; depth at nape 12. D. LVII to LX; A. III, 40 to 42. Body tapering uniformly backward. Vent placed anteriorly, its distance from snout $1\frac{3}{5}$ to $1\frac{3}{4}$ in its distance from base of caudal. Snout compressed, slightly projecting, the lower jaw included; maxillary reaching vertical from front of pupil, $3\frac{1}{3}$ to $3\frac{1}{2}$ in head. Teeth acute, in narrow bands in the jaws, a single well-marked series on vomer, and a patch on front of palatines, those on vomer and palatines fully as large as those on jaws, and equally developed in young and adults. Eyes large, close together, the interorbital space convex, its width about $\frac{1}{2}$ pupil. Diameter of orbit equaling length of maxillary, about $3\frac{1}{2}$ in head. Nostril tubes well developed, $\frac{1}{2}$ diameter of pupil. Gill openings narrower than in other described members of this group, extending forward below the vertical from posterior part of cheek, where they are firmly joined to isthmus, across which they do not form a fold. Gill rakers obsolete. Dorsal beginning over end of opercular flap, its distance from nape equaling distance of latter from posterior margin of pupil; membrane of last spine slightly joined to base of caudal; anterior dorsal spines short, but well connected by membrane; anal with 3 distinct spines, shorter than the rays that follow, the second the longest, all as strong as dorsal spines, and fully connected by membrane, rays all branched at tip, membrane of last ray joined only slightly to base of caudal; caudal sharply pointed in all our specimens, the median rays longest, about as long as head; pectorals evenly rounded, the median rays longest, 14 or 15 in number, all branched; ventrals well developed, about $\frac{2}{3}$ as long as head, consisting of 1 short, sharp spine and 3 rays, the spine not closely joined to rays. Lateral line indistinct, usually appearing obsolete, more evident toward head, consisting of a series of distant pores along median line; scales very small, cycloid, imbricated, covering body, inclosing abdomen, breast, and nape; cheeks scaled, the head otherwise naked, or sometimes with a small patch of scales on upper part of opercles. Color, sides with a series of 10 to 12 narrow white cross bars, the first in front of dorsal fin, the last under last dorsal spine, the bars about $\frac{1}{5}$ interspaces; above lateral line scales conspicuously margined with darker, below lateral line they broaden out and become forked; upper caudal rays at base with an oval white ring inclosing a darker area, this mark more conspicuous in the young; belly and ventrals white, other fins dusky, but without definite markings. (Gilbert.) Bering Sea. Known from 2 specimens; the type, 7 inches long, was taken August 4, 1888, at *Albatross* Station 2852, north latitude 55° 15′, west longitude 159° 37′, at a depth of 58 fathoms, between Nagai and Big Koniushi Islands. The spec-

imen here described from Unalaska. (Named for Dr. J. T. Rothrock, professor of botany, University of Pennsylvania.)

Poroclinus rothrocki, BEAN, Proc. U. S. Nat. Mus. 1890, 40, 55° 15′ N., 159° 37′ W., between Nagai Island and Koniushi Islands. (Coll. *Albatross.*)

918. LUMPENUS, Reinhardt.

(SNAKE BLENNIES.)

Lumpenus, REINHARDT, Dansk. Vidensk. Selsk. Natur., VI, 1837, 110 (*lumpenus = fabricii*).
Leptogunnellus, AYRES, Proc. Cal. Ac. Nat. Sci., I, 1854, 26 (*gracilis*).
Centroblennius, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 209 (*nubilus*).
Leptoblennius, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 209 (*serpentinus*).
Anisarchus, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 209 (*medius*).

Body greatly elongate, moderately compressed, covered with small scales; lateral line indistinct or obsolete. Head long; snout short; no cirri; eyes large, placed high; mouth moderate, with a single row of rather small conical teeth on each jaw; palatine teeth present or absent; gill openings prolonged forward below, very narrowly united anteriorly to the isthmus, not forming a free fold across it. Dorsal composed of numerous, sharp, flexible, rather high, spines; caudal fin long; anal many-rayed; pectorals large, more than ¼ length of head, the middle rays longest; ventrals well developed, jugular, I, 3 or I, 4; intestinal canal long; pyloric cæca present; no air bladder. Chiefly herbivorous. Northern seas. (*Lumpen*, a Danish name of *Zoarces viviparus*, with which these fishes were at first confounded.)

ANISARCHUS (ἄνισος, unequal; ἄρχος, for anal):
a. Anal fin very low in front, the rays gradually lengthened; dorsal spines 61; anal rays 42. MEDIUS, 2790.
aa. Anal fin not much lower in front than behind.
LUMPENUS:
b. Teeth on palatines more or less developed, at least in the adult; anal rays 40 to 46; dorsal spines 63 to 71.
c. Dorsal spines 69 to 71.
d. Anal rays 46; dorsal separate from caudal. ANGUILLARIS, 2791.
dd. Anal rays 41; dorsal slightly joined to caudal. MACKAYI, 2792.
cc. Dorsal spines about 63; anal rays 43. FABRICII, 2793.
LEPTOBLENNIUS (λεπτός, slender; *Blennius*).
bb. Teeth on palatines wanting; dorsal spines 72 to 75; anal rays about 50. LAMPETRÆFORMIS, 2794.

Subgenus ANISARCHUS, Gill.

2790. LUMPENUS MEDIUS (Reinhardt).

Head 6; depth 10. D. LXI; A. 42; V. I, 3. Lower jaw scarcely included, the maxillary reaching front of eye; teeth on palatines, none on vomer; ventrals slender, ½ length of head; lower rays of pectoral shorter than middle ones, the fin shorter than head. Dorsal and anal slightly joined to the truncate caudal; anterior half of anal with the rays shortened. (Collett.) Yellowish, nearly plain. Greenland to Norway and Spitzbergen and westward to Bering Sea and Kamchatka. Specimens from the

coast of Kamchatka are not evidently different from the current figures and descriptions of Atlantic specimens. (Eu.) (*medius*, middle.)

Clinus medius, REINHARDT, Dansk. Vidensk. Afh., VII, 1838, 194, Greenland.
Lumpenus medius, KRÖYER, Naturh. Tidsskr., I, 377, 1837; JORDAN & GILBERT, Synopsis, 777, 1883.
Stichæus medius, GÜNTHER, Cat., III, 281, 1861.
Anisarchus medius, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 210.
Lumpenus medius, COLLETT, Norske Nord-Havs Exp., 62, 1880; JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

Subgenus **LUMPENUS.**

2791. LUMPENUS ANGUILLARIS (Pallas).

Head 8; depth 14. D. LXXI; A. 46 (45 to 50); V. I, 4; B. 7. Cheeks scaly; mouth somewhat oblique, the lower jaw included; maxillary reaching front of pupil; teeth on palatines, none on the vomer; a single series of rather long, conical, and not very closely-set teeth in each jaw. Gill openings prolonged forward a distance greater than length of snout; pyloric cæca 4, unequal. Fins all comparatively high, pectorals $\frac{2}{3}$ length of head, the middle rays longest; ventrals $\frac{1}{3}$ length of head; dorsal and anal distinct from the pointed caudal, which is nearly as long as head. Olive green above, pale below; sides marked above with dark olive brown; a series of more or less distinct oblong blotches of olive brown along middle of sides; dorsal barred or spotted; anal pale; opercle with a dark blotch; head dusky above. Length 18 inches. San Francisco to Alaska; very abundant northward to Sitka and Unalaska; originally recorded from Kamchatka. (*anguillaris*, eel-like.)

Blennius anguillaris, PALLAS, Zoogr. Rosso-Asiat., II, 176, 1811, Kamchatha and Aleutian Islands. (Coll. Billings and Merk.)
Septogunnellus gracilis, AYRES, Proc. Cal. Ac. Nat. Sci., I, 1855, 26, San Francisco.
Gunnellus anguillaris, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 434, 1836.
Lumpenus anguillaris, GIRARD, Pac. R. R. Surv., X, Fishes, 123, pl. 25*b*, figs. 1 to 3, 1858; STORER, Synopsis, 121, 1846; JORDAN & GILBERT, Synopsis, 777, 1883; JORDAN & STARKS, Fishes Puget Sound, 848, 1895.
Stichæus anguillaris, GÜNTHER, Cat., III, 282, 1861.

2792. LUMPENUS MACKAYI (Gilbert).

Head $6\frac{2}{3}$; depth 13 or 14; eye 8 in head; snout 4. D. LXIX; A. II, 41. Very elongate. Head compressed and high, especially anteriorly, the upper profile of snout very convex, the upper jaw decidedly longer than the lower. Mouth nearly horizontal. Maxillary reaching vertical from front or middle of pupil, its length $3\frac{3}{5}$ to $3\frac{4}{5}$ in head. Teeth small, in a narrow band in jaws; a single series of weak teeth on palatines; vomer toothless. Gill openings continued forward to below middle of cheeks, the membranes then narrowly joined to isthmus; gill rakers short and weak, about 10 on horizontal limb of arch. Eye small, its horizontal diameter $\frac{1}{5}$ longer than its vertical, slightly longer than interorbital width. Distance from snout to nape equaling length of postorbital part of head. Opercles large, continued to beyond base of pectorals. Dorsal

beginning immediately above upper end of gill slit, the spines short, strong, and pungent, not flexible; some of the anterior spines short, but not free, the fin increasing in height to opposite front of anal, the longest spine equaling length of snout, the membrane of last spine joining base of upper rays of caudal; anal with 2 strong spines similar to those of the dorsal fin, the second twice length of first and ¾ that of highest dorsal spines; anal rays all forked, the posterior longest, equaling length of snout and eye, free from the caudal; caudal fin rounded in younger specimens, lanceolate in adults, becoming in the latter ⅘ as long as head; ventrals short, of 1 short spine and 3 simple rays, the fin ¼ length of head; pectorals large, the middle rays longest, ⅔ length of head. Scales small, smooth, elongate, imperfectly imbricated, partially embedded or altogether wanting on anterior part of back; cheeks scaled, head otherwise naked; faint traces of a lateral line sometimes visible on middle of sides anteriorly. Color in spirits, light olivaceous (light yellowish in life); a continuous jet-black streak from occiput along each side of dorsal to base of caudal, with 2 interrupted black streaks below it, the lowermost running on middle of side; top and sides of head darker, variously marked with anastomosing black lines and spots; opercles blackish; dorsal and caudal fins dusky translucent, without distinctive markings; anal and ventrals white; pectorals white or dusky; roof of mouth black; peritoneum black dorsally, white ventrally. Bering Sea. Several specimens were seined near the mouth of the Nushagak River, Alaska. (Gilbert.) (Named for Charles Lesley McKay, of Appleton, Wisconsin, a very able young ichthyologist, who was drowned at Nushagak, in Bristol Bay, in 1883.)

Lumpenus mackayi, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 450, pl. 32, mouth of Nushagak River, Bristol Bay. (Coll. Gilbert.)

2793. LUMPENUS FABRICII (Cuvier & Valenciennes).

Head 8 or 9; depth 11 to 15. D. LXIII to LXV; A. 41 to 43; V. I, 3; P. 15. Upper jaw scarcely longer than lower; teeth on palatines few and small, often really or apparently wanting, especially in the young; maxillary not reaching eye; vertical fins distinct; pectorals large, ovate. Color light brown, with large pale rounded blotches separated by brown shades; head yellowish; pectorals yellowish mottled, with a dusky spot at base. Arctic seas; recorded from Spitzbergen, Greenland, the Gulf of St. Lawrence, Wellington Sound, Bristol Bay, and other localities in Bering Sea (Petropaulski and Plover Bay, as *L. anguillaris*). We have specimens from Bristol Bay, Disco, Upernavik, and the Gulf of St. Lawrence. These are apparently identical, and they show clearly the identity of *L. nubilus* with *L. fabricii*.

The following notes are from specimens taken in Bristol Bay, in 4½ to 14 fathoms: These specimens seem to agree in structural details with specimens of *Lumpenus fabricii* from the North Atlantic. The Pacific specimens are lighter in color, with the dusky mottlings confined to the dorsal region, and with a very distinct series of oblong brown blotches along lateral line, alternating with a lower series of small, faint, round spots.

Under parts immaculate; the mottlings along base of dorsal frequently uniting to form a series of oblong blotches alternating with those of lateral line; other specimens show no traces of dorsal blotches; dorsal fin translucent, faintly mottled with darker; caudal with brownish cross bars; pectoral with a round dusky shade at base; fins otherwise unmarked. Mandible with a single series of conical teeth, which widens at symphysis into an irregular double series or narrow patch; a similar series of conical teeth in premaxillaries, within which is a band of fine villiform teeth. A number of small specimens from Disco, Greenland, are entirely similar except for the darker coloration. This species is near *L. anguillaris*, but the latter has a larger mouth, larger teeth, and more numerous fin rays. (Named for Otho Fabricius, the first student of the fishes of Greenland.)

Blennius lumpenus, FABRICIUS, Fauna Grönlandica, 151, 1780, Greenland; not *Blennius lumpenus*, LINNÆUS, which is a species of *Gaidropsarus*, with 2 barbels at the chin.
Gunnellus fabricii, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 431, 1836, Greenland, after FABRICIUS; KRÖYER, Naturhist. Tidsskr., I, 377, 1837; GAIMARD, Voy. Scand., Zool., Poiss., pl. 14, fig. 1.
Lumpenus nubilus, RICHARDSON, Last Arctic Voyage, Fishes, 13, pl. 28, 1855, Wellington Sound. (Coll. Edward Belcher.)
Blennius (Clinus) lumpenus, RICHARDSON, Fauna Bor.-Amer., 90, 1836.
Clinus lumpenus, REINHARDT, Dans. Vidensk. Selsk. Nat. Afh., VII, 194, 1838.
Stichæus lumpenus, GÜNTHER, Cat., III, 280, 1861.
Stichæus nubilus, GÜNTHER, Cat., III, 564, 1861.
Centroblennius nubilus, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 209.
Lumpenus fabricii, JORDAN & GILBERT, Synopsis, 778, 1883.
Leptoblennius nubilus, JORDAN & GILBERT, Synopsis, 778, 1883; GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 451.

2794. LUMPENUS LAMPETRÆFORMIS (Walbaum).

(SNAKE BLENNY; TANGBROSME.)

Head 9; depth about 15. D. LXXIII (LXVIII to LXXIV); A. 50 (49 to 52); V. I, 3. Body elongate, head slender; lower jaw little shorter than upper; maxillary reaching front of eye. Vent well forward, near end of first third of body; pectoral convex, somewhat shorter than head; first 3 or 4 rays of dorsal short, little connected; caudal acuminate, free from dorsal and anal. Yellowish or greenish, with numerous (about 20) faint brown blotches of different sizes, some of them confluent and extending obliquely upward on dorsal; caudal with transverse dark shades. (Collett.) North Atlantic and Arctic on both shores, south to Sweden and Norway, east to Spitzbergen; rare south to Cape Cod, if *L. serpentinus* is the same. We can find no difference on a comparison of our notes with published figures and descriptions, except that Storer describes *serpentinus* as having the caudal plain yellowish. (Eu.) (*Lampetra*, lamprey; *forma*, form.)

Blennius capiti lævi, etc., MOHR, Hist. Nat. Islandiæ, 85, taf. 4, 1786, Iceland; D. 72; A. 54.
Blennius lampetræformis, WALBAUM, Artedi Piscium, III, 184, 1792, Iceland; after MOHR.
Centronotus islandicus, BLOCH & SCHNEIDER, Syst. Ichth., 157, 1801, Iceland; after MOHR.
Clinus nebulosus, FRIES, Vet. Akad. Handl., 55, 1837, Bohuslän, Sweden.
Clinus mohri, KRÖYER, Naturh. Tidsskr., 1 R, 1837, 32, Iceland.
Blennius gracilis, STUVITZ, Nye Mag., Naturvid., I, 406, 1838, west coast of Norway.

Blennius serpentinus, STORER, Proc. Bost. Soc. Nat. Hist., III, 1848, 30, Massachusetts Bay, from the stomach of a codfish (Coll. Capt. Nathaniel E. Atwood); STORER, Hist. Fish. Mass. 169, pl. 18, f. 1, 1867.
Gunnellus islandicus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 433, 1836.
Stichæus islandicus, GÜNTHER, Cat., III, 281, 1861.
Lumpenus lampetræformis, COLLETT, Norske Nord-Havs Exp., 71, 1880; JORDAN & GILBERT, Synopsis, 778, 1883.
Leptoblenninus serpentinus, JORDAN & GILBERT, Synopsis, 778, 1883.

919. STICHÆUS, Reinhardt.

Stichæus, REINHARDT, Dansk. Vidensk. Natur. og Math. Afhandl. 1837, 109 (*punctatus*).
Notogrammus, BEAN, Proc. U. S. Nat. Mus., IV, 1881, 147 (*rothrocki*); young.

Body moderately elongate, covered with small scales; teeth on jaws, vomer, and palatines. Lateral line present, single, running along side of back; pectorals and ventrals well developed. Dorsal moderately high, of spines only; gill openings continued forward below, the membranes scarcely united to the isthmus; pyloric cæca present. Arctic seas. (στιχάω, to set in rows.)

2795. STICHÆUS PUNCTATUS (Fabricius).

Head 4½; depth about 7. D. XLVIII or XLVIX; A. 32 to 35; eye twice interorbital width, 4½ in head; snout subconical, 4 in head. Maxillary about equal to snout, 3⅔ in head, reaching slightly beyond front of eye. Narrow bands of teeth in the jaws and present on vomer and palatines, the outer series in the upper jaw and the inner series in lower jaw enlarged. Scales small, cycloid; head and cheeks scaleless; longest dorsal spines slightly longer than snout. The membrane from last dorsal spine joining extreme base of upper caudal ray; anal wholly distinct; pectorals rather long, reaching vent, 1⅓ in head; ventrals 2½ in head. Numerous large pores scattered over top and sides of head. Lateral line rather close to back, running along the upper fourth of height of body and ending abruptly at about ⅔ the length of body. Color bright scarlet, the head marked below with 5 or 6 brown reticulations; a brown streak from snout through eye; fins irregularly marked by dark bars or spots; a narrow row of 5 large round black spots, each with a white band near its posterior margin, occurring at regular instances along dorsal fin; a row of about 8 large dark spots on anal. Arctic seas, from Greenland to northern Siberia, south to Bristol Bay and Newfoundland. Our description (from Dr. Gilbert) taken from a specimen, about 5 inches in length, from Karta Bay, Alaska. It agrees very closely with the account by Ensign H. G. Dresel, of 2 examples from Godhavn, Disco Island, Greenland. The Alaska species must be the same as the other. Dresel finds the depth 7⅓ in length. Dr. Gilbert further observes: A single specimen, 86 mm. long, was dredged in Bristol Bay, Alaska, Station 3239, depth 11½ fathoms. Several larger individuals were seined in Karta Bay, Prince of Wales Island, Alaska, July 12, 1889. The position of the lateral line in this species is incorrectly given as "median" by Jordan & Gilbert in the Synopsis, pp. 755 and 775. Cuvier and Valenciennes, in their description, drawn from the writings of Fabricius, state

that the lateral line runs along the upper fifth of the height of the body and terminates at about the middle of the length. This correctly describes its position in all our specimens, where it originates immediately above the opercle, exhibiting at first rather a strong upward convex curve, then running nearly parallel to the back, separated from base of dorsal fin by $\frac{1}{5}$ height of body. It is very distinct throughout its course, and terminates at about the middle of the length. The narrow brown streak bounding the lateral line above, in *Notogrammus rothrocki*, is conspicuous in our smallest specimen (86 mm.). Branchiostegal membranes very narrowly joined anteriorly, forming a narrow free fold across the isthmus, from which they are entirely distinct. Narrow bands of teeth in the jaws, and distinctly present on vomer and palatines; the outer series in upper jaw and the inner series in the lower jaw enlarged. D. XLVII or XLVIII; A. I, 32 to 35. The membranes from last dorsal spine join extreme base of upper caudal ray; anal wholly distinct. We have not the material for a comparison of Pacific with Atlantic representatives of this species, and the published descriptions of the latter lack detail. (*punctatus*, spotted.)

Blennius punctatus, FABRICIUS, Fauna Grönlandica, 153, 1780, Greenland; REINHARDT, Naturhist. Selsk. Skrift., II, pt. 2, pl. 10, fig. 3.

Notogrammus rothrocki, BEAN, Proc. U. S. Nat. Mus., IV, 1881, 146, Plover Bay and Cape Lisburne, Siberia; young. (Types, Nos. 27565, 27580, and 27573. Coll. Dr. Bean.)

Clinus punctatus, RICHARDSON, Fauna Bor.-Amer., III, 88, 1836.

Gunnellus punctatus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 428, 1836.

Stichæus punctatus, KRÖYER, Naturhist. Tidsskr., I, 377, 1837; GAIMARD, Voy. en Scand. et Lapon., Zool., Poiss., pl. 20, fig. 2; GÜNTHER, Cat., III, 283, 1861; JORDAN & GILBERT, Synopsis, 775, 1883; DRESEL, Proc. U. S. Nat. Mus. 1884, 249; GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 450.

920. ULVARIA, Jordan & Evermann.

Ulvaria, JORDAN & EVERMANN, Check-List Fishes N. and M. A., 475, 1896 (*subbifurcatus*).

This genus is very close to *Eumesogrammus*, from which it differs in the absence of the lowermost or third lateral line, the median line being bifurcate. (*Ulva*, the sea lettuce, in which many Blennioid fishes live.)

2796. ULVARIA SUBBIFURCATA (Storer).

Head $4\frac{1}{3}$; depth nearly 5. D. XLIV; A. 30. Mouth rather large; maxillary reaching to below orbit; back somewhat arched; ventral outline nearly straight; eyes large; lateral lines 2 (the lowermost lateral line wanting); median lateral line forked; upper branch of median lateral line about $\frac{2}{3}$ length of the head. Brownish, with several round paler blotches above at the base of the dorsal fin; spaces between these blotches darker, appearing like bars; a broad black bar crossing the opercle obliquely from below the orbit, and 2 parallel dark bars running backwards from orbit; belly yellowish white; dorsal fin with numerous black dots. North Atlantic, south to Cape Cod; very rare. (*subbifurcatus*, somewhat forked.)

Pholis subbifurcatus, STORER, Rep. Fish. Mass., 63, 1839, Nahant, Mass. (Coll. Dr. Thos. M. Brewer); DE KAY, N. Y. Fauna: Fishes, 150, 1842; STORER, Hist. Fish. Mass., 258, 1867.

Eumesogrammus subbifurcatus, JORDAN & GILBERT, Synopsis, 775, 1883.

921. EUMESOGRAMMUS,* Gill.

Eumesogrammus, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 210 (*præcisus*).

Body comparatively short, the back somewhat arched; mouth rather large, the jaws with villiform teeth; teeth on vomer and palatines. Scales small; lateral lines 3, without accessory branches; pectorals and ventrals well developed. Dorsal moderately high, of spines only, free or slightly connected with the rounded caudal; gill openings continued forward below, the membranes narrowly joined to the isthmus; pyloric cæca present. Northern seas. (*εὖ*, well; *μέσος*, middle; *γραμμή*, line; the longest lateral line being the middle one.)

2797. EUMESOGRAMMUS PRÆCISUS (Kröyer).

Head 4; depth nearly 6. D. XLIX; A. 34; V. 3. Snout subconical; cleft of mouth slightly oblique; vomerine and palatine teeth present; 3 lateral lines on each side, the median one continued to the base of the caudal; ventral fin $\frac{1}{2}$ as long as the pectoral, which is much shorter than head; dorsal fin terminating just at root of caudal. An ovate, black, white-edged spot between the sixth and tenth dorsal spines. Coasts of Greenland. (Günther.) (*præcisus*, exact.)

Clinus præcisus, KRÖYER, Naturh. Tidsskr., I, 25, August, 1836,† Greenland.
Clinus unimaculatus, REINHARDT, Dansk. Vidensk. Selsk., VII, 114, Feb., 1837, Greenland.
Stichæus unimaculatus, REINHARDT, Dansk. Vidensk., 109, 1837; GÜNTHER, Cat., 283, 1861.
Eumesogrammus præcisus, JORDAN & GILBERT, Synopsis, 774, 1883.

* The 2 following species from the Okhotsk Sea seem to represent 2 new genera (*Ernogrammus* and *Ozorthe*) closely related to *Eumesogrammus:*

ERNOGRAMMUS ENNEAGRAMMUS (Kner).

Head $3\frac{3}{4}$; depth $6\frac{3}{4}$. D. XLI; A. 33 or 34; P. 14 or 15. Eye 4 in head, as long as snout. Mouth large, nearly horizontal, the maxillary reaching middle of eye; lower jaw projecting; profile of snout nearly horizontal; fine pointed teeth in bands on jaws and across the vomer. Head naked; dorsal of high, slender spines; caudal separate, rounded; anal high; pectoral long, $1\frac{1}{3}$ in head; ventrals $\frac{1}{2}$ as long as pectorals; scales very small, smooth; lateral lines each with short oblique branches, each ending in a wide pore; 1 lateral line along base of dorsal from head to caudal, 1 along middle of side, 1 along base of anal to caudal, this forking at the vent and sending 2 parallel branches forward to the breast; brownish, 2 rows of small dark spots along middle lateral line; dorsal and anal with dark spots and a broad dark margin; pectorals with 3 black cross bands; a dark bar at base of caudal; 3 black bars from eye. Okhotsk Sea. Known from a specimen, $1\frac{3}{4}$ inches long, from Decastris Bay. (Kner.) (*ἐννέα*, nine; *γραμμή*, line.) *Ernogrammus*, new genus (*ἔρνος*, branch), is distinguished from *Eumesogrammus* by the branching lateral line.

Stichæus enneagrammus, KNER, Sitzber. Akad. Wiss. Wien 1868, 16, taf. VI, f. 19 Decastris Bay. (No. 1401c Mus. Wien.)

OZORTHE HEXAGRAMMA (Schlegel).

Head $5\frac{1}{4}$; depth $5\frac{1}{4}$. D. XLIII; A. 24. Snout pointed; mouth little oblique; the maxillary reaching front of eye; bands of fine teeth on vomer and palatines; a few large canine-like teeth in front; eye 5 in head, dorsal spines all stiff, the middle ones longest; dorsal joined to caudal by membrane; lateral lines 3, the upper partly interrupted, sending at right angles upward and downward lines which join the middle line; third lateral line along base of anal only. Scales small. Dorsal with large dark brown spots obliquely placed; 3 brown stripes across cheek; anal colored like dorsal; caudal pectoral and ventrals each with 3 dark cross bands. Northern coast of Japan to Okhotsk Sea. This description (after Kner) from a specimen from Decastris Bay (No. 5575, Mus. Wien.). This differs somewhat from the type of the species and may be different. (*ἕξ*, six; *γραμμή*, line.) The new genus *Ozorthe* (*ὄρος*, branch; *ὀρθη*, right angle) is distinguished by the form of its lateral lines as above described.

Stichæus hexagrammus, SCHLEGEL, Fauna Japonica, Pisces, 136, pl. 3, f. 1., 1850. Bay of Simabara, Japan. Head $4\frac{1}{3}$; depth $6\frac{1}{2}$. D. XL; A. 29. GÜNTHER, Cat., III, 284, 1861; KNER, Sitzber. Akad. Wiss. Wien 1868, 45.

† These dates are thus given by Kröyer, as quoted by Dr. Gill, Proc. Ac. Nat. Sci. Phila. 1864, 210. We have been unable to verify them.

Family CCI. CRYPTACANTHODIDÆ.

(The Wry-mouths.)

Body very long and slender, compressed, naked or covered with small, cycloid scales; lateral line obsolete or composed of open pores without tubes; head oblong, cuboid, with vertical cheeks; conspicuous muciferous channels in mandible and preopercle; head flattish above, with deep rounded pits between and behind eyes; mouth large, very oblique; lower jaw very heavy, its tip projecting; premaxillary not protractile; jaws with rather sharp, conical teeth; larger teeth on the vomer and sometimes on palatines. Gill membranes joined to the isthmus, the gill openings prolonged forward below. Pyloric cæca 5. Pseudobranchiæ small. Dorsal fin long, composed entirely of spines, which are rather strong, but enveloped in the skin; dorsal and anal joined to the caudal; no ventral fins; pectorals short. Blennies of large size, of the Northern shores of America. Three species known, forming 3 genera. (*Blenniidæ*, genus *Cryptacanthodes*, Günther, Cat., III, 291, 1861.)

a. Body scaly; lateral line present, composed of open pores; isthmus narrow; teeth on palatines. Delolepis, 922.

aa. Body naked; lateral line obsolete.

c. Palatines with teeth; isthmus narrow. Cryptacanthodes, 923.

cc. Palatines toothless; isthmus rather broad. Lyconectes, 924.

922. DELOLEPIS, Bean.

Delolepis, Bean, Proc. U. S. Nat. Mus. 1882, 465 (*virgatus*).

Body anguilliform, moderately compressed posteriorly, covered with small, imbricated, cycloid scales; vent nearly median; a small anal papilla; lateral line continuous, straight, nearly median, composed of open pores, without prominent tubes. Head oblong, subquadrangular, naked, the muciferous channels well developed, the vertex shallow concave; snout short, obtuse; nostril tubular, close behind premaxillary; eyes small, high, separated by an interspace of moderate width, surrounded by a series of shallow pits; mouth wide, oblique, terminal, the lower jaw projecting beyond the upper; lips fleshy; premaxillaries slightly protractile, with 2 rows of small conical teeth; a few larger teeth at the symphysis; vomer and palatines with a few rather large teeth; tongue smooth, adherent; mandible with a few shallow pits, the series continued on the posterior border of preopercle; opercles unarmed. Gill membranes attached to a narrow isthmus; gill rakers very short; pseudobranchiæ present. Branchiostegals 6. Pectorals short, placed low, their bases vertical; ventrals none; dorsal beginning above gill opening, composed entirely of spines; anal with 2 spines and many split rays; dorsal and anal continuous with the caudal, which is rather long and pointed. Intestine short, with a few pyloric cæca. (*δῆλος*, visible; *λεπίς*, scale.)

2798. DELOPLEPIS VIRGATUS, Bean.

Head 6; depth 10. D. LXXVI; A. II, 46; P. 13; cæca 6. Width of head equal to greatest depth of body; interorbital area equal to snout, or $\frac{1}{3}$ length of mandible; maxillary reaching a little behind eye, its length 3 in

distance from snout to front of dorsal; eye 2 in snout, 11 in head. Beginning at a short distance behind origin of dorsal, small, oblong, cycloid scales, closely imbricated, cover a strip of the body along the lateral line; the scaled area gradually widens backward until, behind the vent, only a very narrow strip along bases of dorsal and anal is naked. Dorsal beginning over upper angle of gill opening; first spine ½ as long as the seventy-first or longest; caudal 11 in length; pectoral 3 in head. Brownish yellow; a brown stripe along lateral line, another along back, a third along base of anal. Length 30 inches. Coast of southern Alaska to Puget Sound; not rare about Seattle. (*virgatus*, striped.)

Delolepis virgatus, BEAN, Proc. U. S. Nat. Mus. 1881, 466, Kingcombe Inlet, British Columbia; Port Wrangel, Alaska (Coll. Capt. H. E. Nichols. Types, Nos. 29149 and 29150, U. S. Nat. Mus.); JORDAN & STARKS, Fishes Puget Sound, in Proc. Cal. Ac. Sci. 1895, 848.

923. CRYPTACANTHODES, Storer.

Cryptacanthodes, STORER, Rept. Fish. Mass., 28, 1839 (*maculatus*).

Body long and slender, compressed, naked, without lateral line; head cuboid, with vertical cheeks and conspicuous muciferous cavities; eyes small, placed high; mouth large, very oblique, the very heavy lower jaw prominent in front; jaws, vomer, and palatines with stoutish conical teeth, in few series. Gill openings prolonged forward below, narrowly attached to the isthmus. Dorsal fin of stoutish spines, hidden in the skin; dorsal and anal joined to caudal; pectorals short; ventrals wanting. (κρυπτός, hidden; ἀκανθώδης, spined.)

2799. CRYPTACANTHODES MACULATUS, Storer.

(WRY-MOUTH; GHOST-FISH.)

Head 6½; depth 13. D. LXXIII; A. 50. Eyes small, placed high, not so wide as the interorbital space, which has 2 ridges and 3 pits; orbital rim raised; 2 deep pits behind each eye at the temples; a deeper pit on the top of head between them; a raised ridge continued backward on each side of head behind orbital rim; maxillary extending to beyond eye; pseudobranchiæ small; pectorals short, 3 in head, their tips reaching beyond front of dorsal; vent a little in front of middle of body. Light brownish, with several series of smallish dark spots, arranged in more or less regular rows, from head to base of caudal; vertical fins closely spotted with darker; head above thickly speckled; body sometimes ("*inornatus*") entirely immaculate. Length 24 inches. Labrador to Long Island Sound; not very common; a few specimens have been taken at Woods Hole. The ghost-fish form (*inornatus*) occasionally seen, is doubtless an albino. (*maculatus*, spotted.)

Cryptacanthodes maculatus, STORER, Rept. Fish. Mass., 28, 1839, coast of Massachusetts; DE KAY, N. Y. Fauna: Fishes, 63, pl. 18, fig. 50, 1842; GÜNTHER, Cat., III, 291, 1861; JORDAN & GILBERT, Synopsis, 780, 1883.

Ophidium imberbe, PECK, Amer. Acad., 2d part, II, 1804, 46, pl. 4, New Hampshire; A. 49; P. 14; C. 22; not of LINNÆUS.

Fierasfer borealis, DEKAY, New York Fauna: Fishes, 316, 1842, New York; after PECK.

Cryptacanthodes inornatus, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 332, Coast of Massachusetts; albino form.

924. LYCONECTES, Gilbert.

Lyconectes, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 452 (*aleutensis*).

Mouth subvertical; lower jaw projecting; premaxillary protractile. Teeth strong, conic, wide set, in more than 1 series. Mucous pits prominent on head. Gill opening narrow, ceasing opposite middle of base of pectorals, the membranes widely joined to isthmus. Dorsal and anal wholly joined to caudal, the latter extending well beyond them; dorsal fin composed of spines only; no ventral fins. Body naked; no lateral line. This genus differs from *Cryptacanthodes* principally in the absence of palatine teeth, agreeing with it in general appearance and in most details of structure. Alaska. (λύκος, wolf; νηκτήρ, swimmer.)

2800. LYCONECTES ALEUTENSIS, Gilbert.

Head 7⅓; depth 14⅓. D. LXIX; A. 49; P. 13; caudal 18. Head square in cross section, the upper and lower surfaces plain, the cheeks vertical, the depth and width equal. Mouth still more oblique than in *Cryptacanthodes maculatus*, with much heavier mandible and less expanded maxillary, the exposed portion of the latter lying vertically, and not extending beyond vertical from middle of eye. Teeth all similar, few in number, those in premaxillary arranged in 2 series, the inner of which are smaller than the outer, from which they are separated by a wide interspace; teeth in mandible in a single series laterally, becoming a sparsely filled patch toward symphysis; 4 or 5 similar conical teeth on head of vomer; palatines toothless. A long nostril tube overhangs the upper lip. Upper lip separated by a fold from forehead, the upper jaw protractile. Eye extremely small, sunken in the socket, which it does not nearly fill, its diameter slightly less than ⅓ interorbital width; supraorbital rim not elevated, and containing no conspicuous projections; suborbital rim swollen, with an enlarged mucous channel; a conspicuous series of mucous pits along each mandible and the margin of preopercle; 2 series on top of head diverging backward from above the eyes; otherwise no pits or projections on head; a shallow triangular depression on occiput. Gill slit much less oblique than margin of preopercle, its length 1⅓ times the distance between lower ends of gill slits, the latter reaching the vertical from middle of opercles. Dorsal fin of rather flexible spines, not concealed in heavy fin membranes; origin of dorsal immediately behind axil of pectorals. Hinder margin of occiput midway between front of dorsal and middle of eye. Origin of anal well in advance of middle of length, its distance from tip of snout contained 1⅔ times in its distance from base of caudal. Pectoral short, rounded, its base separated by a wide prepectoral area from gill slit, the width of area ¾ length of fin, the latter equaling distance from tip of snout to middle of eye. No ventrals. Body covered with lax naked skin, which also covers but does not obscure rays of anal fin; no pores to lateral line. Color in life, reddish on head, body, and fins, due to the blood vessels in the skin. Aleutian Islands. A single specimen, 180 mm. long, known. (Gilbert.)

Lyconectes aleutensis, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 452, pl. 34, fig. 3, Albatross Station 3312, north of Unalaska Island, in 45 fathoms. (Coll. *Albatross*.)

Family CCII. ANARHICHADIDÆ.

(The Wolf-Fishes.)

Body oblong or elongate, covered with rudimentary scales; no lateral line. Head scaleless, without cirri, its bones very thick and strong, the profile strongly decurved. Mouth very large, oblique, the jaws anteriorly with very strong conical canines; sides of lower jaw with very strong molar teeth, which shut against a series of very coarse molars on the palatines; vomer solid, armed with strong molar teeth, the dentition adapted for crushing sea-urchins and mollusks. Gill membranes broadly united to the isthmus; no pyloric cæca. Dorsal fin high, composed entirely of flexible spines; no ventral fins; pectoral fins broad, placed low. Large carnivorous fishes of the northern seas. Two genera and about 6 species known. (*Blenniidæ*, pt., Günther, Cat., III, 208–211, 1861.)

Anarhichadinæ:

a. Body moderately elongate, the tail not tapering to a point; dorsal and anal separate from the caudal. Anarhichas, 925.

Anarrhichthyinæ:

aa. Body eel-shaped, excessively elongate; the dorsal and anal joined with the caudal at the end of the long and tapering tail. Anarrhichthys, 926.

925. ANARHICHAS (Artedi) Linnæus.

(Wolf-Fishes.)

Anarhichas (Artedi) Linnæus, Syst. Nat., Ed. x, 247, 1758 (*lupus*).

Body moderately elongate, covered with rudimentary scales; head scaleless, without cirri, compressed, narrowed above, the profile strongly decurved; mouth wide, oblique; premaxillary not protractile; jaws with very strong conical canines anteriorly; lateral teeth of lower jaw either molar or with pointed tubercles; upper jaw without lateral teeth; vomer extremely thick and solid, with 2 series of coarse molar teeth; palatines with 1 or 2 similar series. Gill membranes broadly joined to the isthmus; no lateral line. Dorsal fin rather high, composed entirely of flexible spines, which are enveloped in the skin; anal fin lower; caudal fin developed, free from dorsal and anal; no ventral fins; pectoral fins broad, placed low; air-bladder present; no pyloric cæca. Northern seas. (*Anarhichas* (or *Scansor*), the climber; an ancient name of *Anarhichas lupus;* from ἀναρριχάομαι, to climb or scramble up; the allusion not evident, the word spelled with a single *r* by Artedi and Linnæus.)

a. Dorsal spines 60 to 70.

b. Vomerine teeth not extending farther backward than the palatine teeth.

c. Back and sides vaguely mottled, without spots or bands; vomerine teeth not extending nearly as far backward as palatine teeth.

latifrons, 2801.

cc. Back and sides profusely covered with roundish black spots; vomerine teeth extending nearly as far backward as palatine teeth.

minor, 2802.

bb. Vomerine teeth extending much farther backward than the short band of palatine teeth; sides of body with 9 to 12 darker cross bars; nape and shoulder with dark spots. lupus, 2803.

aa. Dorsal spines 80 to 85; body without bands or spots; vomerine teeth extending farther backward than palatine band.

d. Head moderate, 4½ in length; caudal rays 20; upper canines 4. LEPTURUS, 2804.

dd. Head very large; caudal rays 17; upper canines 6. ORIENTALIS, 2805.

2801. ANARHICHAS LATIFRONS, Steenstrup & Hallgrimossn.

Head 5; depth 4. D. LXVII; A. 45. Body more robust than *A. lupus,* the dorsal fin lower. Head broad, the profile not strongly decurved; teeth much smaller than in *A. lupus;* vomerine teeth not extending nearly as far back as the palatine series. Pectorals ⅗ length of head; dorsal fin not very high, beginning above the gill opening, the longest spine less than ½ head; caudal 2½ in head. Brown, obscurely spotted with darker; the sides without dark bars or black spots. (Collett.) North Atlantic on both coasts, chiefly north of the Arctic Circle, south to Banquereau on our coast. (Eu.) (*latus,* broad; *frons,* forehead.)

Anarrhichas latifrons, Steenstrup & Hallgrimsson, Förh. Skand. Naturf. 3 die Möte 1842, 647; Bean, Proc. U. S. Nat. Mus., II, 1879, 218; Collett, Meddelsk. Norges Fiske 1879, 46; Jordan & Gilbert, Synopsis, 782, 1883; Goode & Bean, Oceanic Ichthyology, 301, fig. 271, 1896.

Anarrhichas denticulatus, Kröyer, Overs. Vidensk. Selsk. Kjöb. 1844, 140.

2802. ANARHICHAS MINOR, Olafsen.

Head 5⅛; depth 5½. D. LXXVIII; A. 46. Form of *Anarhichas lupus* or a little more slender; fins similarly formed, the dorsal a little lower. Vomerine teeth extending nearly or quite as far back as the palatines. Body pale olivaceous or yellowish; sides without vertical bars; round, black spots covering dorsal and caudal fins as well as back and sides down to the level of the pectoral; head spotted; belly immaculate. North Atlantic, on both coasts, chiefly north of the Arctic Circle, south to Eastport, Maine; Gloucester; and Norway. (Eu.) (*minor,* smaller.)

Anarrhichas minor, Olafsen, Reise i Island, 592, 1772, Iceland.

Anarrhichas pantherinus, Zuiew, Nov. Act. Petrop. 1781, 271; Bean, Proc. U. S. Nat. Mus., II, 1879, 217; Jordan & Gilbert, Synopsis, 781, 1883; Goode & Bean, Oceanic Ichthyology, 301, fig. 270, 1896.

Anarrhichas karrak, Bonnaterre, Encycl. Ichth., 38, 1788, Iceland; after Olafsen.

Anarrhichas maculatus, Bloch & Schneider, Syst. Ichth., 496, 1801, Iceland; after Olafsén.

Anarrhichas leopardus, Agassiz in Spix, Pisc. Brasil., tab. 51, 1829, "Atlantic Ocean."

2803. ANARHICHAS LUPUS, Linnæus.

(Wolf-Fish.)

Head 6; depth 5¼. D. LXII; A. 42. Maxillary reaching beyond orbit; band of vomerine teeth extending much farther back than the short palatine band. Pectorals large, rounded, ⅔ length of head. Dorsal high, beginning over the gill opening, its longest rays about ½ length of head. Brownish or bluish gray; sides with numerous (9 to 12) very dark transverse bars, which are continued on the dorsal fin; besides these

numerous dark spots and reticulations, the spots most distinct below front of dorsal; fins dark; caudal tipped with reddish. Length 3 to 4 feet. North Atlantic, south to Cape Cod and France; rather common both in America and Europe. A large voracious fish, not valued as food. The American form, *vomerinus*, seems to be fully identical with the European. (Eu.) (*Lupus*, a wolf.)

Anarhichas lupus, LINNÆUS, Syst. Nat., Ed. x, 247, 1758, no definite locality; after ARTEDI; GÜNTHER, Cat., III, 208, 1861; JORDAN & GILBERT, Synopsis, 781, 1883; GOODE & BEAN, Oceanic Ichthyology, 299, 1896.

Anarhichas strigosus, GMELIN, Syst. Nat., I, 1144, 1788, British Sea.

Anarrhichas vomerinus, AGASSIZ in STORER, Hist. Fish. Mass., 265, pl. 18, fig. 1, 1867, Cusk Rocks, between Boston and Cape Ann.

2804. ANARHICHAS LEPTURUS, Bean.

(ALASKA WOLF-FISH.)

Head 4½; depth 5. D. LXXXI; A. 52; C. 20 or 21. Head moderate; maxillary ½ as long as head; nostril nearer eye than mouth. Four large canines in the upper jaw and 5 in the lower, all of them strongly recurved; behind the canines in each jaw are a few sharp, conical teeth, also recurved; palatine teeth in 2 series, 4 in the outer and 5 in the inner series, those in the outer series the longer; vomerine teeth in 2 series, the vomerine patch beginning in advance of the palatine, and extending farther back than the latter; head and fins scaleless; median line of body and all of tail with small, widely separated scales. Dark brown, without bands or spots; belly pale, clouded with very dark brown. (Bean.) Coasts of Alaska, south to Vancouver Island; common about the Aleutian Islands, and perhaps identical with *Anarhichas orientalis*. (λεπτός, slender; οὐρά, tail.)

Anarrhichas lepturus, BEAN, Proc. U. S. Nat. Mus., II, 1879, 212, St. Michaels, Alaska; JORDAN & GILBERT, Synopsis, 782, 1883; GOODE & BEAN, Ocean. Ichth., 299, 1896.

? Anarrhichas orientalis, PALLAS, Zoogr. Rosso-Asiat., III, 77, 1811, Kamchatka.

2805. ANARHICHAS ORIENTALIS, Pallas.

This species, if correctly described, would differ from *Anarhichas lepturus* in the very large head, 2⅓ times in total length of body; in the absence of scales; in having the nostril midway between eye and mouth, and in having 6 canines in the upper jaw. Color plain brown. D. LXXXIV; C. 17. Coast of Kamchatka. (Pallas.) As the first of these characters is certainly erroneous, it is likely that the others are also, and that this species is not distinct from *Anarhichas lepturus*. (*orientalis*, eastern.)

Anarrhichas orientalis, PALLAS, Zoogr. Rosso-Asiat., III, 77, 1811, Kamchatka.

? Anarrhichas lepturus, BEAN, Proc. U. S. Nat. Mus., II, 1879, 212, St. Michaels.

926. ANARRHICHTHYS, Ayres.

Anarrhichthys, AYRES, Proc. Cal. Ac. Nat. Sci., I, 1855, 32 (*ocellatus*).

Body elongate, tapering backward into a very long and compressed tail, around which the dorsal and anal are confluent with the caudal. Scales rudimentary; no lateral line. Dorsal high, composed entirely of

flexible spines; pectoral fins broad, placed low; no ventral fins. Head very large, compressed, the snout rather short; mouth large; jaws with very strong, conical canines anteriorly; vomer and palatines each with about 2 rows of coarse molars, the palatine band shutting against similar teeth on the sides of the lower jaw. Gill membranes broadly united to the isthmus. No pyloric cæca. Large, eel-shaped fishes of the North Pacific, remarkable for the tremendous dentition, the head essentially as in *Anarrhichas*, the body strikingly different. (*Anarrhichas; ἰχθύς*, fish.)

2806. ANARRHICHTHYS OCELLATUS, Ayres.

(WOLF-EEL.)

Head 11; depth 15. D. CCL; A. 233; P. 19. Body elongate, formed as in an eel; the head and jaws very strong. Pectorals broad, more than ½ head; longest dorsal spine ½ head. Color dark greenish, the body and dorsal fin everywhere covered with round, ocellated black spots of various sizes, the light markings forming reticulations around the spots; head paler, with the reticulations in much finer pattern; anal pale-edged. Length 5 to 8 feet. Pacific coast, from Monterey north to Puget Sound; generally common. One of our most remarkable fishes; rarely used as food. It feeds chiefly on sea-urchins and sand dollars. (*ocellatus*, with eye-like spots.)

Anarrhichthys ocellatus, AYRES, Proc. Cal. Ac. Nat. Sci., I, 1855, 31, San Francisco; JORDAN & GILBERT, Synopsis, 782, 1893; JORDAN & STARKS, Fishes Puget Sound, 848, 1895.

Anarrhichthys felis, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 150, San Francisco (Type, No. 511. Coll. W. O. Ayres), name only, no description; GIRARD, U. S. Pac. R. R. Surv., X, Fish., 125, pl. 25*a*, figs. 1 to 3, 1858; GÜNTHER, Cat., III, 211, 1861.

Family CCIII. CERDALIDÆ.

Body elongate, compressed, covered with small scales; no lateral line; head small; gill openings reduced to small slit-like openings more or less horizontal in position; dorsal fin very long and low, anteriorly of slender spines, which pass gradually into the soft rays; no free spines; no cirri; tail not isocercal; pseudobranchiæ well developed. Three species known, from the west coast of tropical America in rock pools near the shore. The presence of some spines in the dorsal separates them from the *Scytalinidæ*, while the small gill openings distinguish them from the *Blenniidæ*, to which they are more nearly allied.

a. Ventral fins each with 2 rays; dorsal rays 41; body moderately elongate; greatest depth 10¾ in length; distance from insertion of dorsal to occiput equal to length of head. CERDALE, 927.

aa. Ventral fins each with 1 ray; dorsal rays 48 to 55; body very elongate, eel-like, its depth 15 to 18 in length. MICRODESMUS, 928.

927. CERDALE, Jordan & Gilbert.

Cerdale, JORDAN & GILBERT, Bull. U. S. Fish Comm. 1881, 332 (*ionthas*).

This genus differs from *Microdesmus* in the presence of 2 rays in the ventral fin. Its body is much less elongate than in *Microdesmus*. The gill openings are reduced to small, nearly horizontal slits below and in front of the pectoral fins. (*κερδαλῆ*, the wary one, the fox-like.)

2807. CERDALE IONTHAS, Jordan & Gilbert.

Head $7\frac{3}{4}$ in length; depth $10\frac{3}{4}$. D. 41; A. 36 to 38; C. 4-17-4; P. 12; V. 2. Body considerably elongate, compressed, of nearly equal depth throughout, the head tapering rapidly from occiput to snout; snout short, not obtuse, but the lower jaw heavy and blunt, much projecting beyond the premaxillaries; gape very short and oblique, the tip of the premaxillary not reaching ventral from orbit. Margin of upper jaw formed entirely by the premaxillaries, which are free laterally, but scarcely movable mesially. Maxillary not distinguishable, probably enveloped in the integument of the snout. Teeth rather strong, short, blunt, in a double series in each jaw, apparently wanting on the vomer and palatines. Lips developed laterally, where they form a fold around the angle of the mouth; lower lip adnate mesially, the upper reduced to an obsolete fold. Length of gape $\frac{1}{5}$ length of head. Nostrils 2, distant, the anterior at the end of the snout, almost labial, the posterior above front of orbit, both circular. Eye very small, somewhat less than interorbital width or than length of snout. Distance from snout to past margin of orbit contained $2\frac{3}{4}$ times in length of head. Pseudobranchiæ well developed. Gill openings very narrow, reduced to a short, nearly horizontal slit, extending forward from a point just below the lower base of the pectoral fin. Branchiostegals evident, apparently 4 in number. Vertical fins well developed; dorsal and anal both long, the membrane of the last ray of each joining the base of the rudimentary rays of the caudal. Distance from occiput to the origin of dorsal fin equal to the length of the head; rays of dorsal fin very slender, distinct, the membrane thin and transparent, the rays all, or nearly all, articulate, the anterior simple, the posterior bifid at tip. Vent slightly in advance of middle of length of body, the anal fin beginning immediately behind it; anal rays bifid at tip, excepting the first 2, which appear simple; tail not isocercal, truncate at base of caudal, most of the rays of the caudal springing from the expanding last vertebra; caudal fin rounded, $\frac{4}{5}$ length of head, its rays much branched, more closely set than the rays of the dorsal and anal; rudimentary rays very numerous; ventral fins small, close together, inserted slightly in advance of the lower end of the pectoral, each fin composed of 2 rays, the inner prolonged beyond the outer, and bifid at tip, about as long as pectoral fin and $\frac{2}{3}$ length of head; pectorals well developed, broad, the rays branched at tip. Head and body entirely covered with small scales, which are close set but hardly imbricate, not arranged in series; mandible, snout, and gill membrane scaly; scales on belly and breast smaller than the others and more thickly set; base of caudal and pectoral fins scaled. Coloration in life, body translucent light olive, immaculate below; back and sides very finely marked with clusters of fine dots, the ground color appearing as reticulations between the clusters, which are of irregular size and form; on the sides of the head these dots form bars, which radiate from the eye to the snout and lower side of the head. This species is known from 3 specimens, $2\frac{1}{2}$ to 3 inches in length, taken in a rock pool at Panama. (*ἰονθάς*, freckled.)

Cerdale ionthas, Jordan & Gilbert, Bull. U. S. Fish Comm. 1881, 332, Panama. (Coll. Chas. H. Gilbert.)

928. MICRODESMUS, Günther.

Microdesmus, GÜNTHER, Proc. Zool. Soc. London 1864, 26 (*dipus*).

Body anguilliform, covered with rudimentary scales; head small, with short, obtuse snout and small mouth; lower jaw projecting; teeth minute, in jaws only; eyes very small; gill opening reduced to a very narrow, somewhat oblique slit, in front of lower part of pectorals; vertical fins well developed, the dorsal and anal joined to the caudal by a thin membrane; rays of dorsal mostly articulate, all but a few of the last simple; ventral fins very small, reduced to a single ray; pectorals moderate; vent normal, in middle of body. Pacific coast of tropical America. (*μικρός*, small; *δεσμός*, a band.)

a. Dorsal rays 55, the fin beginning less than a head's length behind occiput.
DIPUS, 2808.

aa. Dorsal rays 48, the fin beginning more than a head's length behind the occiput.
RETROPINNIS, 2809.

2808. MICRODESMUS DIPUS, Günther.

Head about 11 in total length; depth about 18. D. 55; A. 34; C. 16; P. 12; V. 1. Head rather compressed, snout short, mouth very narrow, lower jaw very prominent. Eye minute, lateral, and in anterior third of head. Dorsal fin commencing at a distance from occiput which is somewhat less than length of head, nearly even, the rays very distinct, the interradial membrane being thin and transparent; anal fin commencing immediately behind vent. Caudal rays much more slender and more closely set than those of dorsal and anal; caudal fin rounded, $\frac{2}{3}$ length of head; pectorals as long as ventrals, and $\frac{1}{2}$ as long as head; ventrals close together, and inserted a little behind root of pectoral. Upper parts uniformly brownish olive. Panama. Known from a single specimen, $4\frac{1}{2}$ inches long. (Günther.) (*δίς*, two; *πούς*, foot.)

Microdesmus dipus, GÜNTHER, Proc. Zool. Soc. London, January 26, 1864, 4, pl. 3, fig. 2, Central America (Coll. Capt. Dow); JORDAN, Cat., 126, 1885.

2809. MICRODESMUS RETROPINNIS, Jordan & Gilbert.

Head $14\frac{1}{2}$ in length; greatest depth $15\frac{2}{3}$. D. 48; A. 29; C. 3–17–3; P. 13; V. 1. Body very elongate, compressed, tapering somewhat from front of dorsal to caudal peduncle. Head very small, rapidly tapering forward from occiput; upper profile with a noticable depression behind the orbits, the outline thence to snout strongly convex. Mouth very small, somewhat oblique, the fleshy tip at symphysis of lower jaw projecting much beyond the premaxillaries; gape scarcely reaching vertical from orbit. Teeth small, apparently in a single series in each jaw only. Nostrils double, distant, the anterior near the end of snout, the posterior above anterior margin of orbit. Gill openings a very narrow, somewhat oblique slit, from front of lower third of pectoral fin downward and forward. Branchiostegals evident, 4 or 5 in number. Eye very small, lateral, situated near the upper profile of the head, its diameter nearly $\frac{1}{2}$ the length of the short snout. Vertical fins well developed; dorsal and anal

connected with the caudal by a very delicate membrane. Distance from origin of dorsal fin to occiput 3 times the length of the head, its rays distinct, connected by thin transparent membrane, as are the rays of the anal; most of the rays simple and undivided (but mostly articulate), a few of the posterior only forked at tip; origin of anal fin nearly equidistant between gill rakers and tip of caudal, its rays mostly forked at tip; caudal rays much divided and more closely set than those of dorsal and anal, the fin somewhat pointed in outline, as long as the head; tail not isocercal, truncate at base of caudal fin; ventral fins very small, close together, inserted slightly behind base of pectorals; each fin reduced to a single undivided filament; pectoral fin small, pointed, the middle rays longest, much shorter than the ventrals, and ½ the length of the head. Vent considerably behind middle of total length of the fish (with caudal). Head and body covered with scattered rudimentary scales. Color in life, translucent light olive, with a series of irregular quadrate dark blotches along the back and a series along each side, these blotches formed of clusters of dark points. One specimen, nearly 4 inches in length, was taken in a rock pool at Panama; others since taken by Dr. Gilbert. This species differs from the description of the previously known *Microdesmus dipus*, Günther, in the posterior insertion of the dorsal and the posterior position of the vent, the smaller number of fin rays, the shorter head, longer ventrals, and mottled coloration. (*retro*, backward; *pinna*, fin.)

Microdesmus retropinnis, Jordan & Gilbert, Bull. U. S. Fish Comm. 1881, 331, Panama. (Coll. C. H. Gilbert.)

Family CCIV. PTILICHTHYIDÆ.

(The Quill-Fishes.)

Body extremely elongate, serpentiform, little compressed, the tail tapering to a point. Skin with a few thin, loose, scattered scales; no lateral line. Head unarmed, rather small; upper jaw not protractile; snout short; mouth oblique; lower jaw projecting considerably beyond the upper, with a protruding fleshy appendage at tip. Maxillary reaching front of eye. Mandible little movable. Both jaws with fine, close-set, sharp teeth, in 1 row, the posterior teeth a little the largest; no evident teeth on vomer or palatines. Gill openings restricted to below the most convex part of the opercle, the membranes broadly united below, free from the isthmus. Gills 4, a slit behind the fourth. Pseudobranchiæ very small, almost obsolete. Gill rakers short and stout. Pectorals short; ventrals wanting; dorsal beginning close behind the nape, the anterior portion for about ½ the length of the body composed of very low, stiff, free spines, hooked backward, the posterior portion higher, of slender soft rays connected by thin membrane. No caudal fin, the tip of the tail free. Anal similar to the soft dorsal, without spines. Vent at considerable distance from the head. North Pacific. A single species known.

Concerning the relationships of this interesting group, Dr. Gilbert observes:

"The genus *Ptilichthys*, of which this species [*P. goodei*] is the sole representative, has been doubtfully referred by Dr. Bean to the *Mastacembelidæ*, a

family of fresh-water fishes inhabiting the East Indies, characterized by having the shoulder girdle posteriorly placed and not articulating with the cranium (Order *Opisthomi*, Gill). The necessity for preserving intact the unique type of the species prevented Dr. Bean from making any anatomical examination of *Ptilichthys*, and it was reserved for Dr. Theodore Gill, in the Standard Natural History, III, 259, 1885, to express his disbelief in the relationships which have been suggested, and to make the fish the type of a peculiar family, the *Ptilichthyidæ*, to be placed provisionally among the Blennioid series. His adherence to this view is again expressed in his list of 'Families and Subfamilies of Fishes,' appearing as the Sixth Memoir of Volume VI, of the National Academy of Sciences. He has doubtless indicated the proper position of this peculiar fish as nearly as we are now able to determine it. An examination of its shoulder girdle shows it to be entirely normal. The post-temporal is not furcate, but is a very slender bony rod attaching to the epiotic region of the skull, and giving loose attachment posteriorly to the almost equally slender postero-temporal. The latter overlaps the upper end of the clavicle in the usual manner. A postclavicle was not detected. The coracoid portion consists of a roundish, oblong, perforated hypercoracoid meeting the hypocoracoid directly, without intervening cartilage. The curved line separating the two bones corresponds distally with the interspace between the first (upper) and second actinosts. The hypocoracoid is broad and short; its mesially directed (i. e., inferior) process joins at its tip the clavicle, but is elsewhere separated from the latter by the usual elongate membranaceous interspace. The actinosts are 4 in number, of large size, hourglass-shaped. The jaws are normal, the premaxillary alone occupying the front and sides of upper jaw and bearing the teeth, while the maxillary is a broad bone lying behind it, overlapped proximally by the maxillary process of the palatines. Both vomer and palatines seem to be toothless. The alimentary canal is almost perfectly straight, with the anterior portion entirely enveloped in the long, narrow liver. At the pylorus occurs a short and abrupt U-shaped flexure, scarcely noticeable on account of the closeness with which the sides are joined, and the fact that the width of the flexure is no greater than the cross diameter of the tube. Pyloric cæca are not evident. Air bladder is entirely wanting. The ovary is single, apparently without viaduct, and contains in our specimen eggs which are comparatively very large." (Gilbert.) (*Ptilichthyidæ*, Gill, Standard Nat. Hist., III, 259, 1885.)

929. PTILICHTHYS, Bean.

Ptilichthys, Bean, Proc. U. S. Nat. Mus., IV, 1881, 157 (*goodei*).

Characters of the genus included above. (πτίλον, quill; ἰχθύς, fish.)

2810. PTILICHTHYS GOODEI, Bean.

D. XC, 145; A. about 185; P. 12. Eye rather large, as long as snout, 5 in head; cheeks and opercles long; pectoral fin ½ as long as head; soft dorsal and anal deeper than body posteriorly, anal a little lower than dorsal. Vent near end of anterior third of body; distance from vent to

beginning of soft dorsal 3¼ times length of head; length of head twice its greatest depth, 5½ in distance to vent; appendage of mandible ½ as long as eye; free tip of caudal ⅔ eye. Orange or yellowish, body with a blackish longitudinal stripe; anal darker in color than dorsal. Length about 12 inches. Aleutian Islands; rare; in water of moderate depth. Here described from the original type from Unalaska; 2 other specimens known, the one studied by Dr. Gilbert taken in the entrance to the harbor of Unalaska. (Named for Dr. George Brown Goode.)

Ptilichthys goodei, BEAN, Proc. U. S. Nat. Mus., IV, 1881, 157, Port Levachef, Unalaska (Col. Sylvanus Bailey. Type, No. 26619, U. S. Nat. Mus.); JORDAN & GILBERT, Synopsis, 369, 1883; GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 453.

Group OPHIDIOIDEA.

(THE EEL-POUTS.)

This group, as a whole, agrees with the *Blennioidea* in all respects, except that no spines are developed in any of the fins, save sometimes in the posterior part of the dorsal. From the *Anacanthini*, with which the *Ophidioidea* agree in the jugular ventrals and in the absence of spines, they are separated by the form of the hypercoracoid, which is perforate, as in ordinary fishes. The group is a very large and varied one, widely distributed in all seas.

a. Pseudobranchiæ well developed, very rarely small or obsolete.
  b. Ventral fins jugular, inserted much behind the eye, often wanting, never filamentous.
    c. Gill membranes broadly united, free from the isthmus; ventrals wanting; no scales. SCYTALINIDÆ, CCV.
    cc. Gill membranes united to the isthmus, the gill openings lateral. ZOARCIDÆ, CCVI.
  bb. Ventral fins developed as slender filaments attached at the throat not far behind eye.
    e. Gill membranes broadly attached to the isthmus; no scales. DEREPODICHTHYIDÆ, CCVII.
    ee. Gill membranes nearly separate, free from the isthmus; body scaly. OPHIDIIDÆ, CCVIII.
aa. Pseudobranchiæ absent (or rudimentary in some *Brotulidæ*).
  f. Ventral fins wanting; no scales.
    g. Vent normal, well behind pectorals. LYCODAPODIDÆ, CCIX.
    gg. Vent at the throat. FIERASFERIDÆ, CCX.
  ff. Ventral fins well developed; vent posterior, normal.
    h. Dorsal fin single, low; ventral fins short. BROTULIDÆ, CCXI.
    hh. Dorsal fins 2, the anterior, at the nape, of a single long ray; ventral fins elongate. BREGMACEROTIDÆ, CCXII.

Family CCV. SCYTALINIDÆ.

Body elongate, compressed, eel-shaped, naked. Head depressed, with tumid cheeks, like the head of a snake. Mouth moderate, horizontal, the lower jaw the longer; teeth in a single series in the jaws, vomer, and palatines; no barbels. Gills 4, a slit behind the fourth; pseudobranchiæ present. Gill membranes broadly connected, free from the isthmus. Dor-

sal fin long and low, beginning near middle of body, of slender rays embedded in the skin; anal similar to dorsal, both connected to the caudal fin; tail diphycercal; pectoral fins small, ventral fins wanting. Vent remote from the head, without papilla. Air bladder none; cæca none. Vertebræ numerous, small. The skeleton does not differ essentially from that of *Lycodopsis pacificus*, with which it has been compared. The skull is not at all depressed, the wide depressed form of the head of the fish is due to the fleshy cheeks. The frontal takes up the greater part of the top of the skull, the parietals are separated by the supraoccipital, which extends forward to the frontals. Opercles all present. Lower jaw large and strong. Post-temporal scarcely so firmly attached as in *Lycodes;* the clavicle long and slender. As here understood, this family consists of a single species, a shore fish of the Northern Pacific, living in the gravel between tide marks, and diving with great activity into the wet gravel when disturbed. Its relations are apparently with the *Zoarcidæ*. It is not certain that *Scytalina* has any special affinity with the *Congrogadidæ*, in which group it was at first placed by Jordan & Gilbert.

930. SCYTALINA, Jordan & Gilbert.

Scytalina, Jordan & Gilbert, Proc. U. S. Nat. Mus. 1880, 266 (*cerdale*).
Scytaliscus, Jordan & Gilbert, Proc. U. S. Nat. Mus. 1883, 111 (*cerdale*); substitute for *Scytalina* on account of the earlier *Scytalinus*, Erichson, a genus of Coleoptera.

Body very long and slender, covered with small scales. Head depressed, shaped like the head of a snake, with tumid cheeks and a distinct neck. Eyes small, superior. Mouth rather large, the lower jaw slightly projecting. Teeth conic, in single series on jaws, vomer, and palatines. Each jaw with 2 canines in front. No lateral line; pseudobranchiæ small. Gill rakers almost obsolete. Dorsal fin very low, its first ray near the middle of the body. Anal fin similar to dorsal, nearly as long. Tail diphycercal, the caudal well developed. (Diminutive of *Scytale,* from σκυτάλη, a viper.)

2811. SCYTALINA CERDALE, Jordan & Gilbert.

Head 8; depth 14. D. 41; A. 36. Head broader than body; body much deeper behind vent than anteriorly; snout depressed, rounded at tip; cheeks very long; opercle short; interorbital space rather broad, concave posteriorly; eyes very small, anterior and superior, 10 to 12 in head, 2 in snout, 3 to 4 in interorbital width; upper lip separated by a crease from the skin of the forehead; lower jaw scarcely projecting; edge of lower lip with pores, and small dermal flaps and fringes; maxillary extending somewhat beyond eye; anterior nostrils with small flaps. Lower jaw with a series of close-set, even, conical teeth, besides 2 divergent canines in front; upper jaw with similar teeth in several series in front, the canines smaller and closer together. Pectorals inserted high, little longer than eye; insertion of dorsal slightly in front of anal, a little in front of middle of body; rays of vertical fins low and weak, those of caudal most developed; dorsal and anal joined to caudal; vent close in front of anal,

which is similar to dorsal. Flesh colored, with much mottling of purplish in fine pattern; belly nearly plain; caudal reddish-edged. Length 6 inches. Straits of Juan de Fuca; burrowing among rocks near tide mark. The 2 original types came from the shore of Waadda Island, near Cape Flattery, where the species lives in wet shingle and shows extraordinary activity in hiding among rocks when disturbed. In the same locality 25 additional specimens have been dug out of the gravel by Mr. E. C. Starks in 1895. The species is still unknown from any other locality. (*κερδαλῆ*, the wary one, the fox.)

Scytalina cerdale, JORDAN & GILBERT, Proc. U. S. Nat. Mus., III, 1880, 266, Waadda Island (Type No. 27400. Coll. Jordan & Gilbert); JORDAN & GILBERT, Synopsis, 791, 1883; JORDAN & STARKS, Fishes of Puget Sound, in Proc. Cal. Ac. Sci. 1895, 849, pl. 104.

Family CCVI. ZOARCIDÆ.

(THE EEL-POUTS.)

Body elongate, more or less eel-shaped, naked or covered with very small, embedded, cycloid scales; head large; mouth large, with conical teeth in jaws, and sometimes on vomer and palatines; bones of head unarmed. Gill membranes broadly united to the isthmus, the gill opening reduced to a vertical slit; pseudobranchiæ present; gills 4, a slit behind the fourth. Dorsal and anal fins very long, of soft rays only, or the dorsal with a few spines in its posterior portion; vertical fins sometimes confluent around the tail; pectorals small; ventrals jugular, very small or wanting, if present, inserted behind the eye. Lateral line obsolete or little developed, sometimes bent downward behind pectorals, sometimes sending a branch on median line backward. Gill rakers small; pyloric cæca rudimentary; vent not near head. Pseudobranchiæ present. Genera about 15; species 50. Bottom fishes, chiefly of the Arctic and Antarctic seas; some of them, at least, are viviparous, and some descend to considerable depths. Dr. Gill thus enumerates the skeletal characters of the *Zoarcidæ:*

Orbito-rostral portion of the cranium contracted and shorter than the posterior, the cranial cavity open in front, but bounded laterally by the expansion of the annectant parasphenoid and frontals, with the supraoccipital declivous and tectiform behind, the occipitals above inclined forward along the sides of the supraoccipital, and the exoccipital condyles distant, with the hypercoracoid foraminate about its center and the hypocoracoid with an inferior process convergent to the proscapula. These characters are formulated from the skeleton of *Zoarces anguillaris*. (Gill, Proc. Ac. Nat. Sci. Phila. 1884, 179.) *Zoarchidæ*,* SWAINSON, Nat. Hist. Class. Fishes, II, 82, 184, and 283, 1839. *Lycodidæ*, GÜNTHER, Cat., IV, 319–326, 1862; genus *Zoarces*, GÜNTHER, Cat., III, 295. 1861.

ZOARCINÆ:

I. Dorsal fin low behind, some of its posterior rays short and spine-like; ventrals small; scales present; teeth strong, conic, in jaws only; lateral line present, along middle of side. ZOARCES, 931.

II. Dorsal fin continuous, of soft rays only.

* The name *Zoarchidæ* or *Zoarcidæ* is prior to that of *Lycodidæ*.

LYCODINÆ:
a. Ventral fins present.
b. Vomer without teeth; body scaly.
c. Palatines without teeth.
d. Body very slender, the depth 12 to 16 times in length; lateral line short and faint, ventral in position. EMBRYX, 932.
dd. Body rather robust, the depth 8 to 9 in length; lateral line rather faint, lateral in position. LYCODOPSIS, 933.
cc. Palatines with teeth; lateral line distinct, running along middle of side. APRODON, 934.
bb. Vomer and palatines with teeth.
e. Lower jaw without barbels.
f. Dorsal fin without sculptured scutes at base.
g. Body rather deep, the depth 6 to 8 times in the length.
h. Body more or less scaly. LYCODES, 935.
hh. Body entirely naked, or with a few scales on tail only; none on body or fins. LYCODALEPIS, 936.
gg. Body more slender, the depth 12 to 20 in the length; lateral line lateral in position.
i. Pectoral fin with rounded outlines, the lower rays not greatly produced. LYCENCHELYS, 937.
ii. Pectoral fin deeply notched, the lower rays much produced; lateral line ventral in position. FURCIMANUS, 938.
ff. Dorsal fin with the rays each provided with a sculptured scute or appendage at base; no lateral line; body elongate. LYCODONUS, 939.
ee. Lower jaw with many barbels; body slender, scaly. LYCONEMA, 940.
aa. Ventral fins entirely wanting.
GYMNELINÆ:
j. Teeth moderate, nearly uniform, on jaws, vomer, and palatines.
k. Body scaly; vomer and palatines with teeth; body compressed, not very slender; skull cavernous. BOTHROCARA, 941.
kk. Body scaleless.
l. Lower jaw not very prominent; body very slender; gill openings very narrow. GYMNELIS, 942.
ll. Lower jaw very prominent; body slender, tapering behind; scales undescribed. LYCOCARA, 943.
MELANOSTIGMATINÆ:
jj. Teeth long, unequal, on jaws, vomer, and palatines; skin lax; gill openings reduced to a small foramen; body very slender; scales obsolete. MELANOSTIGMA, 944.

931. ZOARCES, Gill.

(EEL-POUTS.)

Enchelyopus, KLEIN, Ichthyologia Missus, IV, 52, 1747; not as restricted by BLOCH & SCHNEIDER.
Zoarces, CUVIER, Règne Animal, Ed. 2, II, 240, 1829 (*viviparus*).
Zoarchus, SWAINSON, Nat. Hist. Class'n Fishes, II, 283, 1839 (*viviparus*).
Enchelyopus, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 258 (*viviparus*); not of BLOCH & SCHNEIDER.
Macrozoarces, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 258 (*anguillaris*).

Body elongate, compressed, tapering posteriorly; head oblong, heavy, narrowed above, the profile decurved; mouth large; teeth strong, conic, bluntish, in 2 series in the front of each jaw and 1 series on the sides;

teeth in outer series larger; no teeth on vomer or palatines; dorsal fin very long, low, some of its posterior rays much lower than the others, developed as sharp spines; pectoral fins broad; ventrals jugular, of 3 or 4 soft rays; scales small, not imbricated, embedded in the skin; lateral line slender, lateral in position; size large; species viviparous. The American and Asiatic species (subgenus *Macrozoarces*) differ from the European type of *Zoarces*, Cuvier, in the increased number of fin rays and vertebræ. In *Zoarces viviparus* (Linnæus), the European eelpout, the dorsal rays are about 100, the anal about 85, and the number of vertebræ is proportionally diminished. (ζωαρκής, viviparous.)

Subgenus MACROZOARCES, Gill.

2812. ZOARCES ANGUILLARIS* (Peck).

(EEL-POUT; MUTTON-FISH; MOTHER OF EELS.)

Head 6; depth 7. D. 95, XVIII, 17; A. 105. Mouth moderate, lower jaw included; maxillary reaching beyond orbit; pectoral long, about $\frac{3}{4}$ length of head; ventrals $\frac{1}{5}$ head; highest ray of dorsal about equal to snout, the posterior spines about $\frac{1}{3}$ length of eye; first ray of dorsal above preopercle. Reddish brown, mottled with olive, the scales paler than the skin about them; dorsal fin marked with darker; a dark streak from eye across cheek and opercles. Length 20 inches. Delaware to Labrador; rather common north of Cape Cod. Two forms occur, distinguished by the size of the jaws. These have been regarded as distinct species, but the large-mouthed form (*ciliatus; labrosus*) is doubtless the male, as a similar variation occurs in *Lycodopsis pacificus*, and exists in some degree in species of *Lycodes*. (*anguillaris*, eel-like.)

? Encheliopus, GRONOW, Zoophyl., 77, No. 266, 1763, America (*unicolor*); dorsal and anal united with the caudal.

? Blennius americanus, BLOCH & SCHNEIDER, Syst. Ichth., 171, 1801, America; after Gronow.

Blennius anguillaris, PECK, Mem. Amer. Ac. Sci., II, 1804, 46, New Hampshire.

Blennius fimbriatus, MITCHILL, Trans. Lit. and Phil. Soc. N. Y. 1815, 374, pl. 1, fig. 6, New York.

Blennius ciliatus, MITCHILL, Trans. Lit. and Phil. Soc. N. Y. 1815, 374, pl. 1, fig. 7, New York.

Zoarces labrosus, CUVIER, Règne Anim., Ed. II, vol. 2, 240, 1829, America; CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 466, 1836.

Zoarces gronovii, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 469, 1836; after Gronow.

? Enchelyopus americanus, GRONOW, Cat. Fishes, Ed. Gray, 101, 1854, American Ocean.

Zoarces fimbriatus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XI, 468, 1836.

Blennius labrosus, MITCHILL, Trans. Lit. and Phil. Soc. N. Y. 1815, 375.

Zoarces anguillaris, STORER, Fishes Mass., 66, 1839; STORER, Synopsis Fishes N. A., 375, 1845; GÜNTHER, Cat., III, 296, 1861; JORDAN & GILBERT, Synopsis, 784, 1883.

* Allied to *Zoarces anguillaris* is the following species from the Ochotsk Sea:

ZOARCES ELONGATUS, Kner.

Head $5\frac{2}{3}$; depth $11\frac{1}{3}$. D. 80, XII, 22. A. 90 or more. Lateral line extending somewhat beyond pectorals. Color brownish, no brown streak behind eye; dorsal with 12 to 14 large dark spots which extend on the back as faint bands, between which are smaller ones. Known from 1 specimen, $10\frac{1}{2}$ inches long, from Decastris Bay, near the mouth of the Amur. (Kner). (*elongatus*, elongate.)

Zoarces elongatus, KNER, Sitzber. k. k. Akad. Wien 1868, 52, taf. 7, f. 2, Ochotsk Sea. (No. 1502, Wien Mus.)

932. EMBRYX, Jordan & Evermann.

Embryx, JORDAN & EVERMANN, new genus (*crotalinus*).

This genus differs from *Lycodopsis* in the very slender body, the depth being 12 to 16 times in the length, and especially in the ventral position of the lateral line which is faint and incomplete, only the anterior descending portion developed. Deep seas. (ἔν, in; βρύξ, abyss.)

a. Ventrals nearly as long as eye; head 6¾ in length; no scales on head.
CRASSILABRIS, 2813.

aa. Ventrals shorter than pupil; head 5½ in length; head with some scales.
CROTALINUS, 2814.

2813. EMBRYX CRASSILABRIS (Gilbert).

Head 6¾; depth 16; maxillary reaching vertical from front of pupil, 3 in head; exposed portion of eye 6; snout 4; width of snout 3. Body exceedingly slender. Occiput flat, forming a right angle with the descending cheeks, the snout short and wide, the upper lip conspicuously thickened and fleshy on the sides. Upper jaw with a single series of rather large, distant teeth; mandible with a broad patch of cardiform teeth anteriorly, which becomes abruptly constricted on middle of lateral portion of jaw, the inner series alone continued backward toward angle. Palate smooth. Head not conspicuously excavated with mucous canals; series of pores present on mandible and sides of head. Gill openings continued forward to below pectorals, and about to vertical from middle of opercle; the width of the isthmus ¼ the length of slit. Opercular flap with a wide membranaceous border, produced backward and largely covering base of pectorals. Gill rakers very little developed, about 12 movable rudiments on horizontal limb of arch. Origin of dorsal in front of middle of pectorals, slightly farther from occiput than is the latter from front of eye; distance from origin of anal to tip of snout 3¾ in total length; ventrals nearly as long as eye, inserted under middle of opercle; pectorals with 14 or 15 rays, the upper portion of fin longest, the lower rays rapidly shortened, the longest rays ½ as long as head. Scales small, circular, covering nape, breast, and under side of pectorals, but absent on head. Lateral line single, inconspicuous, running below middle of sides, ventral in position, the pores not developed on the scales. Color light brownish above, dark below; lower side of head, margins of snout, gill membranes, part of opercles, and margins of vertical fins jet black; ventrals and posterior face of pectorals black; anterior face of pectorals light glaucous blue, margined with black; lining of mouth and gill cavity and peritoneum black. Pacific coast of southern California. A single specimen, 12 inches long, from *Albatross* Station 2839. (Gilbert.) (*crassus*, thick; *labrum*, lip.)

Lycodopsis crassilabris, GILBERT, Proc. U. S. Nat. Mus. 1890, 106, off southern California. (Type, No. 44280. Coll. *Albatross*.)

2814. EMBRYX CROTALINUS (Gilbert).

Head 5⅛; depth 12; maxillary reaching to behind middle of pupil, 2⅔ in head; eye 6; interorbital width 14; snout 4. Body very slender, with

much the appearance of *Lycenchelys paxillus*, the cheeks tumid, much projecting laterally, the greatest width of head more than ½ its length. Snout short and broad, much depressed, the head scarcely constricted opposite orbits. Eyes with little lateral range. In the single type specimen the upper jaw greatly overlaps the lower, the mandibular band of teeth shutting entirely within those on premaxillaries. Teeth in upper jaw in a single series, 2 or 3 small teeth sometimes present anteriorly, giving traces of an inner series. In lower jaw the teeth are sparsely set in a broad band anteriorly, becoming suddenly contracted to a single series on middle of sides. None of the anterior mandibular teeth enlarged, 2 or 3 of posterior teeth on sides larger and hooked backward. No teeth on vomer or palatines. Nostril in a short tube. Gill slits wide, reaching to below pectorals, but not extending farther forward below than above. Width of isthmus ⅓ length of slit. A series of 7 pores along mandible and preopercle; a second series of 7 or 8 extending from snout along sides of head above premaxillaries. Lateral line faint, descending, its position ventral. Dorsal inserted over middle of pectorals, its origin as far from occiput as is the latter from front of pupil; distance of front of anal from snout equals ⅓ length of body; ventrals short, less than length of pupil; pectorals with posterior margin obliquely truncate, the upper rays longest, the lower growing regularly shorter, thickened at tips, the rays 15 or 16, the longest 2⅔ in head. Scales small, embedded, covering body and most of vertical fins. A few very small, scattered scales on nape, posterior part of occiput, and contiguous parts of cheeks and opercles. Lateral line single, indistinct, running obliquely downward to near base of anal, thence backward, not reaching base of caudal fin. Color dark brown, black on opercles, sides of snout, fins, and lower parts generally; a broad light bar across head behind eyes, extending down on cheeks; some light mottling on mandible and gular membrane; lower rays of pectorals margined with whitish; lining of mouth, gill cavity, and peritoneum jet black. North Pacific. Two specimens known; the type above described from Santa Barbara Islands, the second from *Albatross* Station 3210, south of Saanak Islands, Alaska, depth 483 fathoms.

On this Dr. Gilbert has the following notes:

"The stomach contained remains of Crustacea. Colors in life, head and body light brown, the lower parts darker; snout, suborbital region, and a band across pectorals greenish gilt; no light bar on head. Depth 12¾ in length; head 5⅓; maxillary 2⅓ in head; eye 7, equal to interorbital width. Width of bone between orbits 17 in head. Snout 3⅓ in head. Teeth above in a narrow band, reaching only about halfway of gape. In the mandible, teeth are absent on posterior ⅔ of gape. The gill slit extends a little farther forward below than above. Ventrals as long as pupil. Longest pectoral ray 2¾ in head. Head wholly scaled behind eyes. Lateral line not evident."

(*crotalinus*, from *Crotalus*, κρόταλος, a rattlesnake.)

Lycodopsis crotalinus, GILBERT, Proc. U. S. Nat. Mus. 1890, 105, Albatross Station, 2980, off Santa Barbara Islands. (Coll. *Albatross*.)

933. LYCODOPSIS, Collett.

Lycodopsis, COLLETT, Proc. Zool. Soc. London 1879, 381 (*pacificus*).
Leurynnis, LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 326 (*paucidens*).

Body moderately elongate, the depth 8 to 9 times in length, covered with small, smooth, embedded scales. Lateral line rather faint, extending along middle of side. Head large; snout broad and long; interorbital space very narrow; mouth large, horizontal; teeth conical, those of the upper jaw in a single row; those of the lower in a band in front, the inner series enlarged, larger than the upper teeth; no teeth on vomer or palatines. Ventral fins very small; vertical fins continuous, without spines. Sexes more or less unlike, the mouth larger in the male. Pacific Ocean. (*Lycodes; ὄψις*, appearance.)

2815. LYCODOPSIS PACIFICUS (Collett).

Head 4½ (male) to 5½ (female); depth 8 (male) to 8½ (female). D. 100; A. 85. Female (*pacificus*), head comparatively short; orbital region not restricted, nor cheeks tumid; mouth comparatively small, the maxillary reaching center of pupil. Male (*paucidens*), with the head and mouth large, the snout very broad, the interorbital region constricted; maxillary reaching posterior edge of orbit. Head, nape, and axil of pectoral naked. Dorsal and anal fins enveloped in thick skin, which is covered with embedded scales like those on the body; pectoral ⅓ the length of head in female, ⅔ in male; ventrals ½ length of orbit; mandible ⅓ length of head in female, ⅔ in male; distance from snout to base of dorsal 4½ in length in female, 3¾ in male. Lateral line lateral in portion. Light reddish olive, becoming lighter below; vertical fins margined with black; the scales paler than skin, forming light spots; pectorals dusky. Length 12 to 18 inches. San Francisco to Puget Sound; rather common in water of moderate depth offshore. Sexes markedly different.

Lycodes pacificus,* COLLETT, Proc. Zool. Soc. London 1879, 381, female, Japan. (Coll. Peters) the locality given probably an error.

Leurynnis paucidens,† LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 326, off San Francisco, California, male (Type, No. 23502, U. S. Nat. Mus. Coll. W. N. Lockington); JORDAN & GILBERT, Synopsis, 785, 1883.

Lycodopsis paucidens, GILL, Proc. U. S. Nat. Mus. 1880, 248.

Lycodopsis pacificus, JORDAN & GILBERT, Synopsis, 785, 1883.

934. APRODON, Gilbert.

Aprodon, GILBERT, Proc. U. S. Nat. Mus. 1890, 106 (*cortezianus*).

This genus differs from *Lycodes* only in the dentition, the teeth being present in a single strong series on the palatines, but none on the vomer.

* In regard to the type specimen of *Lycodes pacificus*, Professor Collett writes us as follows (December 2, 1895):

"I got the specimen for describing from the Museum of Berlin from the hands of Professor Peters himself, and he told me that the specimen was from Japan. It is not impossible that he was mistaken; but I can not have any opinion about that."

In view of the fact that the species is abundant off the California coast, whence Professor Peters had obtained collections, that it has not been found in Japan nor in Alaska, we have no doubt that the locality given by Professor Peters is erroneous, and that the fish really came from California.

† The examination of many specimens leaves no room for doubt that *L. pacificus* is the female and *L. paucidens* the male of the same species.

The genus is thus intermediate between *Lycodes* and *Lycodopsis*. (*ἀ*, without; *πρό*, before; *ὀδούς*, tooth.)

2816. APRODON CORTEZIANUS, Gilbert.

Head $4\frac{1}{3}$ to $4\frac{1}{2}$; depth $7\frac{1}{2}$ to 9 in length; head high and narrow, snout broader, but long and very convex. Mouth large, maxillary reaching vertical from middle of orbit, $2\frac{1}{3}$ in head; eye $4\frac{1}{5}$; snout 3; depth of head 2. Teeth in premaxillaries strong, conical, in a single series; lower jaw with the teeth mainly in 2 series, an outer row of slightly enlarged teeth, and an inner row directed backward, a wide interspace between the two series with occasional scattered teeth only posteriorly; on sides of mandible a single series of teeth similar to those in upper jaw; vomer toothless; palatines with a single series of strong conical teeth. Head without conspicuous mucous pores; a strong ridge on middle of occiput anteriorly; gill slit wide, continued forward to vertical from preopercle, the width of isthmus 5 times in length of slit; gill rakers short, better developed than usual, 15 on horizontal limb of outer arch. The vertical limb of arches joined to gill cover by a fold of the lining membrane of the latter, as in *Macrourus*. Pseudobranchiæ well developed. Origin of dorsal but little behind base of pectorals; the hinder margin of occiput midway between dorsal and front or middle of eye; distance from snout to origin of anal $2\frac{2}{5}$ in total length; ventrals inserted under front of opercles, their length about $\frac{3}{5}$ of orbit; pectorals very large, broadly rounded, the upper portion of fin longest, the lower rays rapidly shortened, the lowermost with broad, fleshy tips; rays 20 or 21 in number; scales of the usual type, those on abdomen so deeply embedded as to be almost invisible; head, anterior half of nape, breast, and base of pectorals naked; pectorals and ventrals not scaled, other fins partly covered; lateral line little developed, running along middle of sides and tail. Color light brownish, lighter below; vertical fins broadly margined with black, becoming almost wholly black behind; pectorals light at base, black distally, with a conspicuous white edge; ventrals white; lining of mouth white, of gill cavity dusky; peritoneum black. Cortez Banks, near San Diego, California. The types, 6 specimens, the longest 15 inches, from *Albatross* Stations 2925 and 2948, in 339 and 266 fathoms. Dr. Gilbert also records 1 specimen from *Albatross* Station 3349, off the coast of northern California, depth 239 fathoms. (*cortezianus*, from Cortez Banks.)

Aprodon corteziana, Gilbert, Proc. U. S. Nat. Mus. 1890, 107, Cortez Banks, off San Diego. (Type, No. 46457. Coll. *Albatross*.)

935. LYCODES, Reinhardt.

Lycodes, Reinhardt, Kongl. Dansk. Vidensk. Selsk. Naturv., vii, 1838, 153 (*vahli*).
Lycias, Jordan & Evermann, new subgenus (*seminudus*).

Body moderately elongate, more or less eel-shaped, tapering behind, the depth from 6 to 10 times in the length; head oblong; mouth nearly horizontal; lower jaw included; conical teeth on jaws, vomer, and palatines, those on jaws and palatines mostly in a single series. Dorsal fin beginning behind base of pectoral, without any spines; the rays all soft and articu-

late; pectorals moderate, inserted rather high, its outline rounded; ventral fins small, of 3 or 4 rays. Scales small and embedded, present on part or all of the body, the scaly area more extensive in the adult than in the young. Lateral line faint, sometimes obsolete, normally bent downward behind pectorals and following ventral outline, sometimes with an accessory branch following middle of side; the median branch usually wanting. No air bladder; no anal papilla; pyloric cæca 2 or none. Species numerous, chiefly of the northern seas, inhabiting considerable depths. In general, the male has the head and mouth larger than the female, and the lips thickened. (*λυκώδης*, wolfish.)

LYCODES:*

I. Trunk more or less completely scaled.

a. Dorsal rays about 115; anal rays 90 to 105.

b. Head 4½ to 5 in length; depth 7 to 8.

c. Nape wholly scaly.

d. Lateral line double, with a median and a ventral branch; pectoral rays 22; body blackish with yellowish cross bands or series of spots. ESMARKII, 2817.

dd. Lateral line simple, ventral; body blackish, the young with 6 darker cross bands. VAHLII, 2818.

cc. Nape naked; lateral line obsolete; color plain brown, the fins edged with darker, pectoral rays 21; ventrals short. CONCOLOR, 2819.

bb. Head 5⅔ in length; depth 9; a naked area around dorsal; pectoral rays 19; lateral line ventral; color brownish mottled, the young barred; a black blotch at front of dorsal. ZOARCHUS, 2820.

aa. Dorsal rays 85 to 105; anal rays 68 to 93.

e. Head large, 3⅔ to 4⅓ in length; ventrals about as long as eye; depth 8 to 9½ in length; body chiefly scaly, the fins naked.

f. Body brownish, with a fine network of black lines on head and body, those on body in 5 groups; dorsal edged with black; lateral line probably developed anteriorly only, figured as median; pectorals broad, of about 23 rays. RETICULATUS, 2821.

ff. Body not covered with a network of black lines.

g. Color pale, with dark bands and 2 ocellated spots on the forehead; pectoral rays about 17; lateral line figured as lateral. PERSPICILLUM, 2822.

gg. Color grayish, without bands or spots; pectoral rays about 20; lateral line single, ventral. FRIGIDUS, 2823.

ee. Head short, 5 to 5⅓ in length.

h. Pectoral broad, of 23 or 24 rays; lateral line single, ventral; color plain. TERRÆ-NOVÆ, 2824.

hh. Pectorals narrow, of about 18 rays; ventral fins shorter than eye; lateral line obsolete, or nearly so.

i. Dorsal rays 101 to 105; anal rays 81 to 90; dorsal and anal without dark markings; ventrals more than ½ length of eye; jaws with enlarged flaps of skin.

j. Body in adult not barred, but with 4 dark longitudinal stripes. DIGITATUS, 2825.

jj. Body with 14 to 16 pale crossbars above, which disappear in the adult. PALEARIS, 2826.

ii. Dorsal rays 85; anal 74; ventrals minute, not ½ length of eye; flaps of jaws narrow or obsolete. BREVIPES, 2827.

* The analytical key to the species here given is far from satisfactory. The species should be divided into groups distinguished by the development of the lateral line and the breadth of the pectoral; unfortunately the last named character has been neglected in most of the current descriptions; we have examined all the species accessible to us.

LYCIAS (λύκος, wolf):

II. Trunk naked anteriorly, scaled only on the tail or posterior half.

k. Dorsal fin scaled posteriorly; color brown, with faint yellow transverse bands on back. NEBULOSUS, 2828.

kk. Dorsal fin naked; color uniform pale grayish brown without spots or bands; pectoral rays 21; lateral line single, median. SEMINUDUS, 2829.

Subgenus LYCODES.

2817. LYCODES ESMARKII, Collett.

Head 4½; depth 8. D. 110 to 116; P. 22; A. 95; V. 4. Body behind front of dorsal scaled; vertical fins scaly; nape scaly; snout obtuse; maxillary not more than ½ head; lateral line indistinct, divided, having a median branch besides the ventral series of pores, the median series faint, soon obsolete; pectorals 8 in length; vertebræ 25 + 87. Brownish black, with a whitish-yellow patch on the nape, and 5 to 8 transverse bands of the same color across the dorsal and posteriorly across the anal, these bands becoming broken into annular spots with age. North Atlantic; recorded from Finmark and Spitzbergen. American specimens from the Gulf Stream in about lat. 40°. (Collett.) (Named for Professor Lauritz Esmark, of Copenhagen.)

Lycodes esmarkii, COLLETT, Norges Fiske, 95, 1874, Varanger Fjord, Finmark (Coll. Lensmand Klerk and Prof. Esmark); COLLETT, Norske Nord-Havs Exp., Fiske, 84, pl. 3, fig. 22, 1880.

Lycodes vahli, GOODE & BEAN, Proc. U. S. Nat. Mus. 1879, 209, not of REINHARDT; JORDAN & GILBERT, Synopsis, 786, 1883; GOODE & BEAN, Oceanic Ichthyology, 303, 1896.

2818. LYCODES VAHLII, Reinhardt.

Head 4⅓; depth 8. D. 116; A. 93; V. 4. Head nearly twice as long as high; snout long, maxillary reaching to opposite middle of eye; distance of vent from ventrals nearly equal to length of head; ventral fins less than ½ as long as pectorals; vertical fins scaly; body wholly scaly; lateral line distinct, ventral in position; vertebræ 25 + 87. Brownish yellow, with 6 blackish cross bands extending on the dorsal fin and confluent on the belly, the first cross band on and below the anterior dorsal rays, the second above the vent; adults nearly uniform blackish. Coast of Greenland. (Günther.) (Named for Martin Vahl, an early Danish naturalist.)

Lycodes vahlii, REINHARDT, Kon. Dan. Vidensk. Selsk. Nat. Math. Afh., VII, 1838, 153, tab. V, Greenland; GILL, Cat. Fishes East Coast N. A., 46, 1861; GÜNTHER, Cat., IV, 319, 1862; JORDAN & GILBERT, Synopsis, 786, 1883; GOODE & BEAN, Oceanic Ichthyology, 303, 1896.

2819. LYCODES CONCOLOR, Gill & Townsend.

Head 5 in total; depth about 7½; eye 7¼ in head; snout 3; ventral fin 2 in eye; pectoral 2 in head. D. 118; A. 98; P. 21. Body rather elongate, covered with very small, entirely separated embedded scales which become more distinct anteriorly and extend in advance of the dorsal fin and scapular region, as well as on the vertical fins; lateral line obsolete; pectorals with scattered scales on external and internal surfaces near base; a specialized area of smaller scales behind base of pectoral and a naked area around

upper axilla of pectoral; head moderate, entirely naked; nape naked. Upper teeth in a cardiform band in front, thinning out behind. Lips rather thin. Color nearly uniform, only relieved by the apparently lighter hue of the scales and the somewhat darker margins of the fins; the scales paler than the ground color, which is thus covered with whitish or silvery specks. Bering Sea. Only the type known, its length 22 inches, from which we have taken the above description. (*concolor*, uniformly colored.)

Lycodes concolor, GILL & TOWNSEND, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17, 1897), 233, Bering Sea, lat. 55° 19′ N., long. 168° 11′ W., Albatross Station 3608, (Aug. 12, 1895), in 276 fathoms. (Type, No. 48764, U. S. Nat. Mus. Coll. *Albatross*.)

2820. LYCODES ZOARCHUS, Goode & Bean.

Head nearly 5⅔ in total length; depth 9; eye 4 in head = snout. D. 116; A. 102; P. 19. Body covered with conspicuous embedded scales which extend behind the dorsal and anal, leaving only a narrow naked margin around these fins; head and pectorals naked. A lateral line begins slightly above the upper angle of the gill opening, rapidly curving downward and extending along the lower part of the body not far from base of anal fin; it can be traced above the anterior ⅔ of the anal. Interorbital distance, measured on the bone, 4 in eye; nostrils placed close to upper lip and as far from each other as from the eye; maxillary reaching to vertical through middle of eye; upper jaw 2⅓ in head; mandible nearly ½ head; mandible with a conspicuous flap on each side, about as long as eye, beginning at a distance from the symphysis equal to ½ length of eye; inner edge of mandible also with a slightly elevated ridge of skin. Length of intermaxillary series of teeth equal to ⅓ length of head; length of palatine series nearly equal to that of intermaxillary; vomerines in a round patch; mandibular teeth in 3 series; width of gill opening ⅔ length of head; ventrals in front of base of pectorals, their length 8 in head. Distance between lower angles of gill opening nearly ⅓ length of head; origin of dorsal distant from the head a space equal to ⅓ length of head, slightly behind middle of pectoral; pectoral, when extended, reaching to about vertical from sixth dorsal ray; longest ray of dorsal about ¼ length of head; anal origin under seventeenth ray of dorsal; vent under fifteenth ray of dorsal; longest pectoral ray contained about 9⅓ times in total length. Lateral line distinct, ventral in position, the median pores absent. Color grayish brown, lighter on the belly and under surface of the head; sides irregularly mottled with darker, a narrow dark edge at tip of first 4 dorsal rays. In a young example (No. 39299, U. S. Nat. Mus.) the mottlings on the sides are band-like, the bands not extending below the middle of the body entirely. This example is from lat. 44° 26′ N., long. 57° 11′ 15″ W., 190 fathoms. The type of the description is a specimen 366 mm. long, obtained by the *Albatross* in lat. 44° 46′ 30″ N., 130 fathoms, off Nova Scotia. (*Zoarchus*, a synonym of *Zoarces;* from ζωαρκής, viviparous.)

Lycodes zoarchus, GOODE & BEAN, Oceanic Ichthyology, 308, 1896, off Nova Scotia, in 130 fathoms. (Type, No. 39298. Coll. *Albatross*.)

2821. LYCODES RETICULATUS, Reinhardt.

Head 4; depth about 8. D. 94; A. 75; V. 4. Body entirely scaly; lateral line faint, developed anteriorly (fide Günther's plate), probably becoming ventral; vertical fins naked. Head twice as long as high; snout long; maxillary extending to behind middle of eye; distance from vent to ventrals more than length of head; cæca 2. Brownish, with reticulated black lines on the head and body, those on the body disposed in 5 groups or cross bands, the 3 anterior of which emit 1 or 2 vertical streaks on the dorsal fin; dorsal dark edged. Length 14 inches. North Atlantic, from Greenland south to Narragansett Bay, in 17 to 140 fathoms; abundant also in northern Europe. (Eu.) (*reticulatus*, netted.)

Lycodes reticulatus, Reinhardt, Kong. Dansk. Vid. Afh., vii, 1838, 167, Greenland; Günther, Cat., iv, 320; Gill, *l. c.*, 260; Collett, Nord-Havs Exp., 84; Jordan & Gilbert, Synopsis, 787, 1883; Goode & Bean, Oceanic Ichthyology, 305, 1896.
Lycodes rossi. Malmgren, Om Spetsbergen Fiskfauna, 516, 1864, Spitzbergen.
Lycodes gracilis, Sars, Christ. Vid. Selsk. Forh. 1866, Dröbak.

2822. LYCODES PERSPICILLUM, Kröyer.

This species is distinguished by a light body color and dark bands, also 2 ocellated spots on the forehead, which have suggested the specific name. Still further separated from the previously known species of *Lycodes* by the smaller number of fin rays, larger eye, etc. (Kröyer.) Greenland and southward in deep water. Specimens were obtained by the *Albatross* from Station 2491, in 45° 24′ 30″ N. lat., 58° 35′ 15″ W. long., at a depth of 59 fathoms, and from Station 2456, in 47° 29′ N. lat., 52° 18′ W. long., at a depth of 86 fathoms.

The following is the substance of Dr. Günther's description:

Head 4 in total length; depth nearly 8. Head not quite twice as long as high; snout long; upper maxillary extending to below middle of eye. Distance of vent from ventrals nearly equal to length of head. Yellowish, with 9 or 10 brownish cross bands, edged with dark brown, and broader than the interspaces, the first occupying the upper parts of the head and inclosing a pair of roundish, yellowish spots situated behind the level of the eyes; the second cross band is on and before the anterior dorsal rays.

(In the figure of Goode & Bean the lateral line is represented as median, which is probably not correct.) (*perspicillum*, eyebrow, from the spot above the eye.)

Lycodes perspicillum, Kröyer, Dansk. Vidensk. Selsk. Afhandl., xi, 1845, 233, Greenland; Günther, Cat., iv, 320, 1862; Gill, Proc. Ac. Nat. Sci. Phila. 1863, 260.

2823. LYCODES FRIGIDUS, Collett.

Head 4 to 4½ in total length; depth 6½ (to 9½, young). D. 93 to 98 (including ½ of the caudal, 99 to 104); A. 80 to 85 (including ½ of the caudal 86 to 90); P. 20 to 21; V. 3. Head wide and flat. Scales with very conspicuous mucous cavities below, small, covering the entire body, but not the head, nor the base of the dorsal and anal fins. In the young the middle of the belly, the base of the fins, and the fins themselves are usually naked.

Teeth present on intermaxillary, mandible, palatines, and vomer; lateral line low, extending from upper end of gill opening in a curved direction down toward vent from which it runs close along anal to end of tail. (Goode & Bean.) Pectoral fin obliquely truncate at tip, appearing furcate when not spread open. North Atlantic and Arctic Ocean, from Spitzbergen south to the New England coast, where many specimens were taken in 516 to 1,423 fathoms. (Eu.) One of Collett's specimens from Hammerfest, examined by us.) (*frigidus*, frozen.)

Lycodes frigidus, COLLETT, Forh. Selsk. Christ. 1878, Nos. 14 and 15, Beeren Island and Spitzbergen; COLLETT, Norske Nord-Havs Exp., 96, pl. 3, f. 23, 24, 1880; GOODE & BEAN, Oceanic Ichthyology, 305, 1896.

2824. LYCODES TERRÆ-NOVÆ, Collett.

Head 5 to $5\frac{1}{2}$; depth 8 to 11. D. 106 to 108; A. 89 to 93; P. 23 or 24. Body slender, head small; pectorals broad; maxillary reaching to middle of eye; band of palatine teeth very short, scarcely $\frac{1}{4}$ length of maxillary band; body entirely scaly, head naked; lateral line ventral, extending along edge of belly, the median branch wanting; vent before middle of body. Color lost in type, the only specimen known. Banks of Newfoundland, in 155 fathoms. (Collett.) (*terra*, land; *novus*, new, from Newfoundland.)

Lycodes terræ-novæ, COLLETT, Campagnes Scientifiques, L'Hirondelle, X, 1896, 54, Bank of Newfoundland, Hirondelle Station 162, in 155 fathoms. (Coll. Albert, Prince of Monaco.)

2825. LYCODES DIGITATUS, Gill & Townsend.

Head 5 in total; depth about $8\frac{1}{2}$; eye $6\frac{1}{2}$ in head; snout 3; ventral fin $1\frac{2}{3}$ in eye; pectoral $1\frac{3}{10}$ in head. D. 101; A. 81; P. 18. Body moderately elongate; covered with small, entirely separated embedded scales, which become nearer anteriorly and extend in advance of the dorsal fin as well as on the vertical fins; no specialized area of smaller scales behind base of pectorals; pectorals scaleless; head moderate, entirely naked; nape naked; upper jaw with outer row of close-set teeth, broader in front; teeth on vomer and palatines; lips rather thick. Color in alcohol, brownish yellow, suffused with reddish in front, variegated, darker anteriorly, with 4 dark longitudinal stripes most distinct about middle of body, fading out backward; fins light and without dark margins; head dark above and laterally light below. Bering Sea. Only the type known, from which we have taken this description, its length 18 inches; possibly the adult of *L. palearis*, but the pectoral fins are shorter than in the latter. (*digitatus*, fingered.)

Lycodes digitatus, GILL & TOWNSEND, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17, 1897), 232, Bering Sea, lat. 56° 14′ N., long. 164° 8′ W., at Albatross Station 3541, in 49 fathoms. (Type, No. 48765, U. S. Nat. Mus. Coll. *Albatross*.)

2826. LYCODES PALEARIS, Gilbert.

Head $5\frac{1}{4}$ in length; depth $9\frac{1}{2}$ to 11 in length, $2\frac{1}{6}$ in head; eye 5 to 6 in head, $1\frac{1}{2}$ to 2 in snout. Dorsal with about 105 rays, counted to middle of caudal; anal about 90; pectoral 18; ventrals $1\frac{1}{2}$ to $1\frac{1}{3}$ in eye, twice as long

as in *L. brevipes*; pectorals 1⅓ in head. Head naked; nape more or less naked, the scaleless area variable in extent, sometimes confined to its anterior third, sometimes reaching nearly to front of dorsal; body sparsely covered with embedded scales; axil naked; lateral line short, decurved, extending scarcely beyond middle of pectorals. Anal origin under eighteenth dorsal ray. Teeth present in jaws, vomer and palatines, those in premaxillaries laterally in a single series which widens anteriorly into a rather broad patch, the outer teeth somewhat enlarged, especially in front; all the premaxillary teeth shut outside on the mandibular series which are opposed to those on vomer and palatines; mandibular teeth arranged similarly to those in upper jaw, the lateral series somewhat enlarged, continuous with the inner edge of the symphyseal patch; vomerine teeth bluntly conic, 3 or 4 in number; palatines in a single series. Snout long, prominent, the upper jaw projecting beyond the lower for a distance equaling ⅔ of orbit; upper lip thin, much expanded laterally, continuous posteriorly with the lower lip which forms a wide free membranaceous lobe opposite middle of each mandible; anteriorly the lower lip becomes abruptly contracted and adnate to the jaw, leaving the symphyseal portion without free margin; inner edge of mandible with wide membranaceous borders, which increase in width anteriorly where they terminate in a pair of acutely pointed free flaps; these and the membranaceous margins very conspicuous in both young and old individuals. In *L. brevipes* they are very inconspicuous, becoming evident in adults only. General color brownish olive, growing lighter on the lower parts; dorsal with 14 to 16 white vertical bars, extending in young examples across back and sides and onto anal fin, in adults confined to the fins, and frequently indistinct or wanting; anterior dorsal angle frequently black, separated from remainder of fin by a curved white bar; dorsal and anal not black margined as in *L. brevipes;* in the latter, the white lateral bars are 9 to 12 in number, and are usually confined to upper half of body; there is also no black spot on anterior dorsal rays. This species is very close to *L. brevipes* Bean, differing constantly in the longer ventrals, the greater development of mandibular and labial folds, the more numerous white bars, and the smaller eye. Bering Sea. Three specimens, 113 to 166 mm. long, from *Albatross* Stations 3253 and 3254, in Bristol Bay, in 36 and 46 fathoms. (Gilbert.) (*palea*, the wattles of a cock.)

Lycodes palearis, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 454, Bristol Bay, Alaska. (Coll. *Albatross.*)

2827. LYCODES BREVIPES, Bean.

Head 5 in total length; depth 10. D. 85 to middle of caudal; A. 74; P. 21. Body covered with scales except immediately behind pectoral fins; head naked; dorsal and anal fins minutely scaled; diameter of eye equals the length of the snout, 4 in head; dorsal origin nearly over middle of pectoral; anal origin under eighteenth ray of dorsal; ventrals minute, scarcely more than ⅓ diameter of eye; pectorals 9 in length of the body; lateral line single, very faint, ventral in position, abruptly decurved and becoming obsolete over about the tenth anal ray. A narrow light band

across the nape and from 9 to 11 across the back extending downward about to median line and becoming obscure in adults; dorsal and anal each with a narrow dark margin. (Bean.) Aleutian Islands to Kadiak; abundant; taken by us in large numbers off Karluk in 1897. (*brevis*, short; *pes*, foot.)

Lycodes brevipes, BEAN, Proc. U. S. Nat. Mus. 1890, 38, between Unga and Nagai islands, at Albatross Station 2848, in 110 fathoms. (Type, No. 45362. Coll. *Albatross*.)

Subgenus LYCIAS, Jordan & Evermann.

2828. LYCODES NEBULOSUS, Kröyer.

D. 87; A. 68; P. 19; V. 3. Body naked anteriorly, the posterior part of dorsal fin scaly; the anal naked or nearly so. Brown, with small, faint, yellow, transverse bands across the back. Greenland. (Kröyer.) An imperfectly described species, not recognized by any recent writer. This species and the next should perhaps be placed in *Lycodalepis*. (*nebulosus*, clouded.)

Lycodes nebulosus, KRÖYER, Kong. Dan. Vidensk. Sel. 1844, 140, Greenland; GILL, Proc. Ac. Nat. Sci. Phila. 1863, 261; JORDAN & GILBERT, Synopsis, 787, 1883.

2829. LYCODES SEMINUDUS, Reinhardt.

Head 3½; depth 7. D. 91; A. 74; P. 21. Body naked in front of vent, scaly behind; fins naked. Head large. Distance of ventrals from vent somewhat more than length of head; cæca 2. Color uniform pale grayish brown, without spots or bands. North Atlantic, from Greenland to Spitzbergen; rare. (Collett.) (*semi-*, half; *nudus*, naked.)

Lycodes seminudus, REINHARDT, Kong. Dansk. Selsk., etc., 1838, 221, Omenak, Greenland; GÜNTHER, Cat., IV, 320, 1862; JORDAN & GILBERT, Synopsis, 787, 1883; GOODE & BEAN, Oceanic Ichthyology, 307, 1896.

936. LYCODALEPIS, Bleeker.

Lycodalepis, BLEEKER, Verl. Akad. Amst., Ed. 2, VIII, 1874, 369 (*mucosus*).

This genus differs from *Lycodes* in the absence of scales on trunk and fins; scattered scales sometimes present on the tail only. (λυκώδης, *Lycodes*; ἀλεπίς, without scales.)

a. Color brownish, with many cross bands and streaks of cream color; head 4⅛ in length; depth 8; lateral line obsolete; tail sometimes with a few scales. POLARIS, 2830.

aa. Color blackish, with about 5 narrow pale cross bars on back; head 3¾ in length; depth 8; lateral line double, a median and a ventral series of pores being faintly developed. MUCOSUS, 2831.

2830. LYCODALEPIS POLARIS (Sabine).

Head 4⅛; depth 8. D. 85; A. 67; P. 18; V. 3; Br. 6. Head depressed, its greatest width ¾ of its length; distance from tip of snout to nape nearly equaling greatest width of head, 6 in length; upper jaw 1¾ to 2 in head, extending to vertical of hind margin of orbit, larger in male than in the female; a full series of teeth on premaxilliaries, and in front of these a few smaller teeth form an outer imperfect series; a toothless space at symphysis,

first tooth on each side of this larger than any of the rest; 1 complete series of teeth on mandible, and in front of it, about the symphysis, 2 irregular short series; a few teeth in a cluster on head of vomer; palatines with a short single series; teeth all slender and slightly recurved; long diameter of eye 9 in head. Pectoral $1\frac{1}{2}$ in head; ventral about as long as eye; longest dorsal ray $3\frac{2}{3}$ in head; vent in middle of total length, immediately behind third cross band; longest anal ray $4\frac{1}{3}$ in head; scattered scales present on posterior two-thirds of tail in 1 specimen (type of *L. coccineus*), wholly wanting in the others, typical of *L. turneri;* no scales on the fins; no trace of lateral line. Color light brown; abdomen grayish brown; lower parts of head cream; a band of cream on the anal from origin of rays to about their middle; a crescentic V-shaped band of same color, mottled with umber, crossing nape and continuing behind pectorals, extending backward to the first cross bar; a streak of cream, more or less interrupted by umber, extending backward from eye across cheek almost to end of operculum; 10 bands of cream color, bordered with dark umber, from tips of dorsal rays extending on lower half of body, becoming wider and somewhat broken below middle of body; a very indistinct caudal tip of cream color. In young examples these markings are very distinct; in older ones they grow progressively more obscure, the oldest having scattering blotches of cream color instead of bands, the V-shaped nuchal band persisting longest. The type of *Lycodes coccineus* is described as brown, red below; pectorals reddish brown above, carmine below; 9 bluish-white bands on the dorsal; a few whitish blotches on sides and on head; anal brownish red; head white below; a whitish blotch as large as eye at upper angle of gill opening. Length 18 to 20 inches. Arctic Ocean, Bering Straits, and adjacent waters south to St. Michaels. Here described from the type of *Lycodes coccineus* and from a number of specimens from Point Barrow referred to *Lycodes turneri*. Evidently all belong to the same species, but 1 has a scaly tail while the others are wholly naked. In 2 large examples, supposed to be males, the head is very much depressed, broad and flat, and the maxillary is more than $\frac{1}{2}$ head. In the others the head is smaller, less flattened, with smaller mouth, the maxillary 2 in head. These are doubtless females and young. The species should probably stand as *Lycodalepis polaris*. (*polaris*, polar.)

?Blennius polaris,* SABINE, Parry's Journal, Voyage 1819–20, Supplement, 212, North Georgia.

Lycodes turneri, BEAN, Proc. U. S. Nat. Mus. 1878, 464, St. Michaels, Alaska (Type, No. 21529. Coll. Dr. Lucien M. Turner); TURNER, Contr. Nat. Hist. Alaska, 93, pl. 4, 1886.

Lycodes coccineus, BEAN, Proc. U. S. Nat. Mus., IV, 1881, 144, Big Diomede Island, Bering Strait (Coll. Dr. Bean. Type, No. 27748, 20 inches long, with scales on the tail); JORDAN & GILBERT, Synopsis, 787, 1883.

Blennius (Zoarches?) polaris, RICHARDSON, Fauna Bor.-Amer., III, 94, 1836.

Lycodes polaris, GÜNTHER, Cat., IV, 321, 1862.

Lycodalepis turneri, JORDAN & GILBERT, Synopsis, 788, 1883; SCOFIELD, in JORDAN & GILBERT, Fur Seal Invest., 1898.

Lycodalepis polaris, JORDAN & GILBERT, Synopsis, 788, 1883.

* *Blennius polaris* is thus described: Without any scales; length of the pectoral exceeding twice its breadth, having 15 rays. Yellowish, lighter on the belly, with 11 large saddle-like markings across the back, the middle of these markings being much lighter than their edges; the whole back and the sides marbled. (Sabine.) Coast of North Georgia.

2831. LYCODALEPIS MUCOSUS (Richardson).

Head 3⅐; depth 8. D. (including ½ of caudal) 90; A. (including ½ of caudal) 71; P. 18; V. 3. Body robust, head very large; snout 3 in head; interorbital area 6 in head; nostrils much farther from eyes than from each other, their distance from eyes 4⅛ in head; upper jaw 6¼ in total length; lower jaw 6⅜; eyes small, close together, their long diameter 11 in the head; distance from tip of snout to base of pectoral fin 3½ in total length; pectoral fin 6⅜; length of ventrals equaling long diameter of eye. (Goode & Bean.) Lateral line (in specimens from Cumberland Gulf) very faint, but with both median and ventral branch. Blackish, with irregular white markings in the form of 5 faint and narrow bars across the back. Arctic America. (*mucosus*, slimy.)

Lycodes mucosus, Richardson, Last Arctic Voyage, 362, pl. 26, 1855, Northumberland Sound; Bean, Bull. U. S. Nat. Mus., No. 15, 112, 1879; Goode & Bean, Oceanic Ichthyology, 306, 1896.

Lycodalepis mucosus, Jordan & Gilbert, Synopsis, 788, 1883.

937. LYCENCHELYS, Gill.

Lycenchelys, Gill, Proc. Ac. Nat. Sci. Phila. 1884, 110 (*muræna*).

This genus contains small and very slender species differing from *Lycodes* in the elongation of the body, the depth being from 10 to 20 times in the length. The lateral line is single and median in all known species. The genus is very close to *Lycodes*, but the position of the lateral line sufficiently defines it, especially in connection with the slender eel-like form. (λύκος, wolf; ἔγχελυς, eel.)

a. Lower half of pectoral not notably longer than upper; depth 12 to 16 in length.
b. Dorsal rays 92; anal 88; color grayish, with irregular brown patches. VERRILLII, 2832.
bb. Dorsal rays 118; anal 110; color brown, the head darker. PAXILLUS, 2833.
aa. Lower half of pectoral considerably longer than upper; head with large pores; depth 14 times in length; color dusky brown. PORIFER, 2834.

2832. LYCENCHELYS VERRILLII (Goode & Bean).

Head 5⅝; depth about 13; eye 2 in snout. D. 92; A. 88; P. 15; V. 5. Body elongate; head much depressed. Distance of vent from ventrals slightly greater than head, its distance from snout about 3 in body; distance of dorsal fin from snout ½ greater than head; distance of anal from snout twice head; dorsal and anal fins about equal in height, with even margins, not differentiated from caudal, the rays increasing somewhat in length posteriorly; distance of pectoral from snout about equal to head, twice length of pectoral; pectoral reaching vertical from base of second dorsal ray; distance of ventrals from snout less than head, their length less than ¼ that of pectorals. Head, body, and fins enveloped in tough, lax skin. Scales cycloid, circular, and ovate, with numerous concentric striæ, and about 18 lobes on margin, the whole perimeter being lobed; scales deeply embedded in the skin at distances from each other equal to their own diameters, most numerous on upper part of body and

extending upon base of dorsal; very few scales upon lower half of body, none on anal fin. Upper jaw far overlapping the lower; gape reaching orbit. A series of 6 large pores on each side, extending backward from nostril toward angle of opercle, the fourth of the series under center of orbit; a similar series, 7 on each side, along line of lower jaw from its symphysis to angle of opercle, all slit-like, the others circular. Nostrils at extremities of fleshy tubes. Teeth in lower jaw in 2 rows, nearly uniform in size; teeth of upper jaw in a single series, somewhat enlarged near the symphysis; patches of smaller teeth behind; about 7 teeth on vomer; a single row on palatines; all the teeth curved. Gill opening narrow, the membranes attached to the isthmus. Color, body above lateral line light grayish brown with numerous minute circular dots marking the position of the scales; pearly white below lateral line; brown irregular patches upon sides, bisected by lateral line, the lower half color of dorsal, that above darker and with the white dots, these brown patches 7 to 10 in number; a brown spot on tip of tail; abdominal region livid blue. Coast of Massachusetts, in deep water; a dwarf species very small in size. (Named for Prof. Addison E. Verrill of Yale University.)

Lycodes verrillii, GOODE & BEAN, Amer. Journ. Sci. Arts, XIV, 1877, 474, off coast of New England in the Gulf Stream; JORDAN & GILBERT, Synopsis, 786, 1883.

Lycenchelys verrilli, JORDAN, Cat., 124, 1885; GOODE & BEAN, Oceanic Ichthyology, 309, figs. 277 and 277 A, 1896.

2833. LYCENCHELYS PAXILLUS (Goode & Bean).

Head 8; depth 16; eye 3½ to 4 in head, equal to snout, which is 4 times interorbital width. D. (with ½ of caudal) 118; A. 110; P. 16; V. 3. Body attenuate, head broad, flat above, with declivous profile; cheeks full and protuberant; teeth stout, recurved, and sharply pointed, in a single series in each jaw, except at the symphysis; a few teeth clustered at the head of the vomer; palatines with a single series; the tubular nostril much nearer tip of snout than eye. Lateral line median, faint and short (in specimens examined by us). Dorsal beginning over tip of pectoral; ventral little longer than pupil. Scales very small, present everywhere except on head and pectorals, nearly covering vertical fins. Light brown, the head somewhat darker. Gulf stream, lat. 35° to 41° N., in deep water, 263 to 904 fathoms. (Goode & Bean.) (*paxillus*, a peg.)

Lycodes paxillus, GOODE & BEAN, Proc. U. S. Nat. Mus. 1879, 44, between LaHave and Sable Island Banks (Type, No. 22177. Coll. Capt. J. W. Collins), a male in breeding form; JORDAN & GILBERT, Synopsis, 785, 1883.

Lycodes paxilloides, GOODE & BEAN, Bull. Mus. Comp. Zool., X, No. 5, 207, 1883, off Newfoundland (Type in M. C. Z. Coll. *The Blake*); a normal, not sexually distorted individual.

Lycenchelys paxillus, JORDAN, Cat., 124, 1885; GOODE & BEAN, Oceanic Ichthyology, 311, figs. 279 and 282, 1896.

2834. LYCENCHELYS PORIFER (Gilbert).

Head 5⅔; depth 14. Body very slender. Head much contracted opposite orbits, the snout expanded, as in *Lycodopsis paucidens*. Mouth moderate, the maxillary reaching vertical from front of pupil, 3⅔ in head; eye 5⅔; snout 3⅔; interorbital width ⅓ eye. Teeth in front of premaxillaries

in 2 series, merging into 1 laterally, the outer series anteriorly somewhat enlarged; teeth in front of mandible in a broad band, narrowing laterally to a single series, none of them enlarged; vomer and palatines with single series. Head with 2 series of large and very conspicuous elongate pores, 1 series on mandible and subopercle, the second parallel with it on level of snout. Gill openings wide, extending forward beyond preopercular margin, the width of isthmus less than $\frac{1}{4}$ length of slit. Distance from origin of dorsal to tip of snout $4\frac{1}{8}$ in length. Median dorsal rays simply forked near base, those posteriorly in both dorsal and anal repeatedly subdividing. Distance of anal from snout $2\frac{7}{8}$ in length; pectorals rounded, the lower half of fin longer than the upper, the rays thickened, the fin containing 15 or 16 rays, its length less than $\frac{1}{2}$ head; ventrals longer and slenderer than usual, each apparently composed of 2 rays closely joined, their length $\frac{3}{4}$ orbit, inserted unusually far forward, being in advance of preopercular margin. Scales very small, circular, partially embedded, covering body and vertical fins; head, antedorsal region, breast, and a strip connecting the two latter embracing base and axil of pectorals, naked; lateral line median. Color dusky brown, the fins, sides of head, and belly blackish; lining of mouth and gill cavity and peritoneum black. Off Lower California. A single specimen, 12 inches long, from *Albatross* Station 3009, in 857 fathoms. A transitional species approaching *Furcella*. (*porus*, pore; *fero*, I bear.)

Lycodes porifer, GILBERT, Proc. U. S. Nat. Mus. 1890, 104, off Lower California, in 857 fathoms. (Type, No. 44384. Coll. Dr. Gilbert.)

938. FURCIMANUS, Jordan & Evermann.

Furcimanus, new genus (*diapterus*); JORDAN & EVERMANN, Check-List Fishes, 480. 1896 (*diapterus*); preoccupied by Furcella, Lamarck, 1801, a genus of mollusca.

This genus differs from *Lycenchelys* in the forked pectorals, the upper and lower rays being much longer than the middle ones. The lateral line is single and ventral in position (not lateral as in *Lycenchelys*). (*furca*, a fork; *manus*, hand.)

2835. FURCIMANUS DIAPTERA (Gilbert).

Head $5\frac{3}{8}$ to 6; depth 12; eye large, usually longer than snout, 3 to $3\frac{3}{8}$ in head; snout $3\frac{1}{4}$ to $3\frac{3}{8}$; interorbital width about 10. Body slender. Mouth small, somewhat variable in length, the maxillary reaching vertical from between front and middle of pupil, $2\frac{1}{2}$ to 3 in head. Teeth in premaxillaries in a double row throughout, the 2 series well separated, rarely with 1 or 2 teeth intercalated, showing traces of a third row; the teeth of inner series small and directed obliquely inward; those of outer series anteriorly enlarged, becoming smaller on sides of jaw; on front of mandible the teeth are in a broad band, in which traces of 3 or 4 irregular series can be made out; none of these enlarged; laterally the teeth are arranged in a single series, those opposite middle of cleft considerably enlarged; a small patch of from 2 to 5 teeth on vomer; palatines with a single row much shorter than premaxillary patch. Nostril with a short inconspicuous tube. Mandible and preopercular border with deep pit-like excavations, which are not evident in fresh specimens; no evident mucous pores on the head.

Gill openings wide, extending below the base of the pectorals; the gill membranes joined to isthmus for a distance equaling $\frac{2}{3}$ length of slit; gill rakers very short, almost tubercular, but compressed and slightly movable, about 15 present on anterior limb of outer arch; a wide slit behind fourth gill. Ventrals short, inserted under middle of opercle. Pectorals deeply notched in both young and adults, the median rays much shorter than either upper or lower, the lobe produced by the elongate lower rays varying in length, being sometimes shorter than upper lobe, sometimes longer; the rays of lower lobe are thickened, and undoubtedly serve as a support to the fish when resting on the bottom, as has been observed in so many other forms; the pectorals contain 20 or 21 rays; in the structure of this fin the present species seems to differ from all previously described forms, with the exception of *L. esmarkii*, in which the notched condition of the fin does not persist in the adults. Scales small, embedded, covering entire body and vertical fins; the scales on nape are much reduced in size, and in 2 specimens ($11\frac{1}{2}$ and $7\frac{1}{2}$ inches long) are continued onto occiput, which they entirely cover; in another specimen, 9 inches long, the occiput is naked, and in another, 5 inches long, the anterior part of nape is likewise naked; in the latter, as in other specimens, the dorsal and anal are well scaled. Lateral line single, wavy, ventral in position, extending from above gill slit obliquely downward to near base of anal, along which it is continued for a variable distance, not reaching base of caudal. Color dusky brownish, blue-black on belly and along anterior portion of base of anal; 8 or 9 narrow white bars on sides, most conspicuous in the young, in which they are continued up on dorsal fin and become forked below on middle of sides, forming Λ-shaped marks; in adults these bars become faint or wholly disappear; when present, they are not continued on dorsal, and are usually vertically divided by a streak of the ground color; in the small specimen there is a distinct black blotch on margin of anterior dorsal rays; in adults, the vertical fins are brownish on basal portion, their distal half black; pectorals and ventrals deep blue-black; mouth, gill cavity, and peritoneum dusky or black. Several specimens, from *Albatross* Stations 2892, 2896, 3067, and 3077, in depths from 82 to 376 fathoms, off the coasts of California and Oregon. (Gilbert.) A remarkable species. (*διά*, divided; *πτερόν*, fin.)

Lycodes diapterus, Gilbert, Proc. U. S. Nat. Mus. 1891, 564, off the coast of Oregon, in 685 to 877 fathoms. (Type, No. 44385. Coll. Dr. Gilbert.)

939. LYCODONUS, Goode & Bean.

Lycodonus, Goode & Bean, Bull. Mus. Comp. Zool., x, No. 5, 208, 1883 (*mirabilis*).

Body elongate, formed as in *Lycenchelys* and *Lyciscus;* scales small, circular, embedded in the skin; lateral line very short or obsolete; jaws without fringes, lower jaw included; fin rays all articulated, each ray of dorsal and anal supported laterally by a pair of sculptured scutes; caudal distinct, not fully connate with dorsal and anal; ventrals present; gill opening narrow; teeth as in *Lycodes*. Deep water. (*Lycodes*, with a meaningless change of termination.)

2836. LYCODONUS MIRABILIS, Goode & Bean.

Head 7 in total length; depth about 18. D. about 80; A. about 70; C. 9; P. 18; V. 3; scales as in *Lycodes*, the scales not extending out upon the fins; no scales on head and nape. Lateral line apparently obsolete posteriorly; not extending back of the extremity of the pectoral, its position median; eye high up, $2\frac{1}{2}$ in head, equal to postorbital portion of the head; the width of interorbital space less than diameter of pupil, $3\frac{1}{2}$ times in long diameter of eye; nostrils immediately in front of eye; maxillary extending to vertical through anterior margin of pupil; mandible, to a little behind vertical through posterior margin of the pupil; dorsal fin inserted slightly behind vertical through base of pectoral (the portion of the fin present in the mutilated specimen before us contains 80 articulated rays; the first 10 or 11 scutes do not support rays, but whether rays were originally present or not can not be ascertained); longest dorsal ray about equal to longest anal ray, its length about 3 in head; distance of vent from snout twice length of head; anal beginning immediately behind vent, of about 70 articulated rays; caudal rays extending beyond tips of ultimate dorsal and anal rays, about 9 in number; distance of ventral from snout equal to twice length of upper jaw; middle ventral ray longest, it being $\frac{1}{2}$ as long as postorbital part of head; length of pectoral equaling 3 times that of snout. Off the New England coast, in depths of 721 to 1,309 fathoms; a most remarkable little fish. (*mirabilis*, wonderful.)

Lycodonus mirabilis, Goode & Bean, Bull. Mus. Comp. Zool., x, No. 5, 208, 1883, New England Coast, lat. 38° 20′ 8″ N., long. 73° 23′ 20″ W., in 740 fathoms (Type in M. C. Z.); Jordan, Cat. Fishes, 124, 1885; Goode & Bean, Oceanic Ichthyology, 312, 1896.

940. LYCONEMA, Gilbert.

Lyconema, Gilbert, Rept. U. S. Fish Comm. 1893 (1896), 471 (*barbatum*).

Generic characters as in *Lycodes*, but the lower jaw covered with a dense mass of slender filaments or barbels, between which can be seen the mucous pores of the mandible. In *Iluocetes*, a related genus from the Antarctic, the mandible is provided with a series of hollow tubes, which are doubtless the produced margins of the pores. Alaska. (λύκος, wolf; νῆμα, thread.)

2837. LYCONEMA BARBATUM, Gilbert.

Head $6\frac{1}{3}$; depth $11\frac{1}{2}$; maxillary 3 in head; eye $3\frac{1}{3}$; snout $4\frac{1}{2}$. D. 103; A. 90 (each counted to middle of caudal); P. 15, its length $1\frac{9}{10}$ in head; ventrals very short, $\frac{1}{2}$ to $\frac{2}{3}$ diameter of orbit. A dense fringe of filaments covers the entire under surface of lower jaw, extending to behind angle of mouth; another series laterally on the throat, and a few scattering ones sometimes present on the branchiostegal membranes; upper jaw without barbels. Body slender; upper jaw overlapping the lower; mouth small, maxillary reaching vertical from front of pupil; teeth all conical, none of them much enlarged, those in lower jaw in a patch or irregular double series, narrowing to a single series laterally; in upper jaw, a single series, the teeth of which increase in size toward the middle line, the mid-

dle teeth being almost canine-like; behind the latter, a short inner series of small teeth directed backward; teeth on vomer and palatines in a single series. Gill slits continued forward to slightly beyond bases of ventrals, and to level of lower edge of base of pectorals; width between gill slits ½ diameter of eye; pseudobranchiæ well developed; posterior line of occiput midway between origin of dorsal and front of pupil or front of eye; origin of anal fin at end of first third of length of body; pectorals broad, with the posterior edge emarginate, some of the upper and the lower rays longer than the intermediate ones. Scales showing traces of definite arrangement in series, widely separated anteriorly, becoming crowded toward end of tail, continued up on the vertical fins, but not on head, on anterior half of nape, nor on the pectoral fins; lateral line very faintly shown, and for only a short distance behind head, where its course is obliquely downward; the usual series of mucous pores present, but not conspicuous. In spirits this species has an olive-brown ground color, becoming white on underside of head and on abdomen; a series of 8 or 9 brown spots ½ as large as eye, along middle of sides, those posteriorly continued downward onto base of anal, the last 2 or 3 reaching edge of fin and there developing into intense black blotches; a similar series of smaller spots corresponding in position to those just described occurs along the base of dorsal, these continued as faint bars on the fin, at the margin of which they develop into a black blotch, those posteriorly wider and more intense; an intermediate series of spots alternating with the 2 just described; an elliptical jet-black spot occupies the greater part of caudal fin, and is narrowly margined all around with white; peritoneum jet-black; the mouth and gill cavities white. Coast of Alaska, in rather deep water; known from 12 specimens, the longest 6½ inches; depth 204 fathoms. (*barbatus*, bearded.)

Lyconema barbatum, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 471, coast of Alaska, at Albatross Station 3129, lat. 36° 39′ 40″ N., long. 122° 01′ W., in 204 fathoms.

941. BOTHROCARA, Bean.

Bothrocara, BEAN, Proc. U. S. Nat. Mus. 1890, 38 (*mollis*).

Body elongate, compressed, semitranslucent, covered with small scales; small teeth in jaws and on vomer and palatines; mucous pores about head largely developed. No ventral fins; dorsal and anal joined to caudal. Deep-sea fishes, allied to *Lycodes*, but lacking ventrals. The species have been referred to the Antarctic genus *Maynea*, Cunningham. From the latter, however, *Bothrocara mollis* seems to be distinct, differing in the larger mouth, more cavernous head, and lower dorsal. In some regards *B. pusilla* is intermediate, and it may belong to *Maynea*. (*βόθρος*, cavity; *κάρα*, head.)

a. Body elongate, with the head short, 6 in length; depth 9; mouth small; mucous cavities small; color light brown, the dorsal dark-edged. PUSILLA, 2838.

aa. Body deeper and more compressed; the large head 4¼ in length (5½ in young); depth 6⅔; mouth large; mucous cavities large; color uniform brown, the vertical fins dark-edged. Size large. MOLLIS, 2839.

2838. BOTHROCARA PUSILLA (Bean).

Head 6 in the total length; depth 9. D. 95, including ½ caudal; A. 81, including ½ of caudal; P. 17; eye 3 in head; snout 4. Body elongate, little compressed; head short; mouth small; maxillary extending to below front of pupil; gill clefts narrow, the anterior end below margin of preopercle; width of isthmus rather less than ½ of orbit; the low dorsal beginning nearly over axil of pectoral; pectoral nearly ⅔ as long as head; vent as far from end of head as dorsal origin from tip of snout. Color light brown; dorsal and anal with a narrow dark margin. Size small; length 6½ inches. Eastern parts of Bering Sea, and about the Alaskan Peninsula, in rather deep water. Besides the original types Dr. Gilbert records a few specimens from north of Unalaska, at depths of 121 to 351 fathoms. (*pusillus*, weak.)

Maynea pusilla, BEAN, Proc. U. S. Nat. Mus. 1890, 39, off Nagai Island, lat. 55° 10′ N., lon. 160° 18′ W., in 110 fathoms (Type, No. 45360. Coll. *Albatross*); GILBERT, Rept. U. S. Fish Comm. 1893 (1895), 455.

2839. BOTHROCARA MOLLIS, Bean.

Head 4¼ in total length in adult, 5⅓ to 5½ in young; depth 6⅔ in adult, 10 in young. D. 100 to 105 to middle of caudal; A. 89 to 95 to middle of caudal; eye 4 in head in adult, 3½ in young. Body covered with embedded scales, which extend on dorsal and anal fins. Head naked, breast and nape scaly; snout blunt, the lower jaw included; maxillary reaching middle of pupil, 2½ in head; large mucous cavities conspicuous along mandible, suborbital ring, and top of head; vomerine and palatine teeth present, the latter in a narrow band, obscure in the young. Pectoral 2 in head; origin of dorsal slightly behind base of pectoral, its distance from tip of snout 4 in total length; origin of anal under seventeenth dorsal ray; longest dorsal ray 5 in head; longest anal ray 8½ in head. Gill openings wider than in *Bothrocara pusilla*, the anterior end of the cleft under posterior margin of eye, the width of the isthmus less than ½ diameter of pupil. Color uniform brown, fins lighter; dorsal and anal margined with black, more prominent posteriorly where it covers the entire fins. North Pacific. Adult examples from southern California, 18 inches long, were described as *Maynea brunnea*, while a young individual, 5½ inches long, from Queen Charlotte Islands, with the vomerine and palatine teeth not evident, was made the type of a distinct genus, as *Bothrocara mollis*. The two are identical and apparently belong to the same genus as *B. pusilla*. Similar specimens, 1 adult and 2 young, were dredged by us (*Albatross*) off Bogoslof Island in 664 fathoms. Dr. Gilbert records also specimens from near Unalaska, depth 316 fathoms. The teeth on the palatines are in a single series instead of a wide band, as stated in the original description. (*mollis*, soft.)

Bothrocara mollis, BEAN, Proc. U. S. Nat. Mus. 1890, 38, off Queen Charlotte Islands, in 876 fathoms (Type, No. 45359. Coll. *Albatross*); JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

Maynea brunnea, BEAN, Proc. U. S. Nat. Mus. 1890, 39, lat. 33° 8′ N., lon. 118° 40′ W., off San Clemente Island, southern California, in 414 fathoms. (Coll. *Albatross*.)

942. GYMNELIS, Reinhardt.

Gymnelis, Reinhardt, Dansk. Vidensk. Selsk. Afhandl., vii, 131, 1838 (*viride*).
Cepolophis, Kaup, in Archiv fur Naturgesch. 1856, 96 (*viridis*).

Body elongate, naked. Vertical fins without spines; ventral fins none. Small conical teeth on the jaws, vomer, and palatines. Gill openings very narrow. No air bladder; pyloric cæca none; no anal papilla. Size small. Cold seas. Two or 3 species known: *G. pictus*, from the Antarctic, and *G. viridis*, which ranges widely in Arctic waters, and with which the very dubious *G. stigma* is probably identical. (γυμνός, naked; ἔγχελυς, eel.)

a. Dorsal fin inserted close behind pectoral, its distance from it much less than diameter of eye; no ocellus on dorsal fin. VIRIDIS, 2840.

aa. Dorsal fin inserted an eye's diameter behind pectoral; a large black spot, ocellated with white, on dorsal fin above vent; other ocelli sometimes present. STIGMA, 2841.

2840. GYMNELIS VIRIDIS (Fabricius).

Head about 6½; depth about 13; eye 7 in head. D. 100; A. 80. Snout subconical, longer than the eye; jaws equal; mouth oblique; maxillary reaching beyond eye; teeth rather small, conical, in a single series on each side, forming a patch anteriorly; distance from snout to vent 2⅓ times length of head. Pectoral rounded, inserted low, its length less than ½ that of head. Dorsal fin inserted close behind pectoral, its distance from it much less than diameter of eye. Body pale, with faint dark cross shades; dorsal clouded but without black spot; anal dusky. Arctic seas, Alaska to Greenland and Nova Scotia; abundant in the Arctic waters south to Unalaska and Bristol Bay, where specimens were taken in shallow water; our specimens from Bristol Bay.

Ensign H. G. Dresel records 1 small specimen (No. 28636, U. S. Nat. Mus.), badly preserved, obtained by Mr. Newton Pratt Scudder in Davis Straits, July, 1879. Length 100 mm. D. ca. 97; A. ca. 80. In this specimen the maxillary does not extend to the posterior margin of the eye, which is comparatively very large. Its diameter is longer than distance from tip of snout to orbit, and is contained 4 times in head. Head 7 in total length; depth 12. Pectoral 2 in head. (*viridis*, green.)

Ophidium viride, Fabricius, Faun. Grœn., 141, 1780, Greenland.
Ophidium unernak, Lacépède, Hist. Nat. Poiss., ii, 280, 1800, Greenland; after Fabricius.
Gymnelis viridis, Richardson, Last Arctic Voyage, 321, pl. 29, 1854.
Gymnelis viridis, Reinhardt, Dansk. Vidensk. Selsk. Afh., vii, 1838, 131; Günther, Cat., iv, 323, 1862; Krøyer, Poissons du Nord, Voy. en Scand. et Lap., pl. 15. a-f; Collett, Norske Nordh. Exped., Fiske, 123, pl. 4, fig. 32, 1880; Jordan & Gilbert, Synopsis, 789; Gilbert, Rept. U. S. Fish Comm. 1893 (1896), 455.
? *Gymnelis pictus*, Günther, Cat., iv, 324, 1862, no locality.

2841. GYMNELIS STIGMA (Lay & Bennett).

Head 6; depth 11. D. 90; A. 70. Form, size, and general appearance of *G. viridis*, the dorsal inserted farther back, an eye's diameter behind pectoral. A large, round black ocellus, ringed with white, on dorsal fin above vent; 2 or 3 other ocelli sometimes present; head and nape with small white spots; body with faint dark shades and bands. Otherwise

as in *G. viridis*, from which it may not be distinct; but the above characters appear in our specimens (from near the Pribilof Islands) and in Richardson's figure of *G. unimaculatus*. The white spots on the head were mistaken for "very small scales" in the original description of *G. stigma* from Dr. Collie's notes. This description is, in substance, as follows: No trace of ventral fins; dorsal, caudal, and anal fins united into a transparent ridge; rays of branchial covering distinct; scales very small. Color dilute brown, with void swathes and spots; a purplish spot near beginning of dorsal fin. Snout obtuse; chin with a large gibbosity; teeth small. Length about 5 inches. (Lay & Bennett.) Arctic regions, Greenland to Bering Sea, with the preceding, and apparently equally common. (*στίγμα*, spot.)

Ophidium stigma, Lay & Bennett, Zool. Beechey's Voy., 67, pl. 20, fig. 1, 1839, Kotzebue Sound. (Coll. Dr. Collie.)
Gymnelis viridis var. *unimaculatus*, Richardson, Last Arctic Voyage, 367, 1854, Northumberland Sound. (Coll. Edward Belcher.)
Gymnelis stigma, Günther, Cat., iv, 325, 1862; Jordan & Gilbert, Synopsis, 789, 1883.

943. LYCOCARA, Gill.

Uronectes, Günther, Cat., iv, 325, 1862 (*parrii*); name preoccupied in Crustacea.
Lycocara, Gill, Proc. Ac. Nat. Sci. Phila. 1884, 180 (*parrii*).

Body ensiform, compressed; tail long and tapering; ventrals none; vent not far distant from the head; numerous minute teeth in jaws and on palate; lower jaw the longer; no barbel; scales and gill openings unknown. One species, very imperfectly known, no specimens having been obtained by any recent collector. (*λύκος*, wolf; *κάρα*, head.)

2842. LYCOCARA PARRII (Ross).

Head 4. D. 50; A. 45; P. 37. Head very obtuse, its length, depth, and breadth equal; head broader than the body, flattened and grooved between the eyes, which are lateral and rather large; lower jaw the longer; jaws and palate with minute teeth; greatest depth of body somewhat more than length of head; neck much arched. Dorsal inserted just behind head; pectoral extending beyond vent. Vent not far distant from head. Color uniform. Baffins Bay. (Günther.) (Named for Capt. William Edward Parry, the Arctic explorer.)

Ophidium parrii, Ross, Parry's Third Voyage, App., 109, 1826, Baffins Bay.
Uronectes parrii, Günther, Cat., iv, 326, 1862; Jordan & Gilbert, Synopsis, 789, 1883.

944. MELANOSTIGMA, Günther.

Melanostigma, Günther, Proc. Zool. Soc. Lond. 1881, 21 (*gelatinosum*).

This genus is distinguished from *Bothrocara* by the much more elongate teeth, which in the jaws, as well as on the vomer and palatines, stand in single series. Gill openings much smaller than in related forms, reduced to a small foramen above the base of the pectoral. Skin loose and movable, as in *Liparis*, enveloping the vertical fins; pectorals very small;

ventrals none. Body tapering very rapidly backward; the tail very slender. Deep-sea fishes, of soft substance, allied to *Bothrocara*, but with stronger teeth. (*μέλας*, black; *στίγμα*, spot.)

a. Maxillary reaching beyond front of pupil; color purplish gray, becoming black on the tail. GELATINOSUM, 2843.

aa. Maxillary not reaching beyond vertical from front of pupil; color uniform deep black. PAMMELAS, 2844.

2843. MELANOSTIGMA GELATINOSUM, Günther.

Body enveloped in a loose, delicate skin, as in *Liparis*. Head large, deep, compressed, with obtuse snout. Eye large, 3½ in head, and longer than snout. Cleft of mouth rather oblique, but lower jaw not projecting beyond upper; lips not fleshy; gill opening reduced to a very narrow foramen above base of pectoral fin; origin of dorsal fin and root of pectoral enveloped in loose skin of body; dorsal fin probably commencing above middle of pectoral, low at first, but becoming considerably higher posteriorly; pectorals very narrow, consisting of a few rays only. Upper parts tinged with a purplish-gray; sides marbled with same color, which toward end of tail becomes more intense, almost black; inside of mouth, gill openings, and vent black. Total length of the type specimen 5½ inches; distance of the snout from the gill opening ⅞ inches, from the vent 1⅝ inches. (Günther.) Deep waters of the western Atlantic; originally known from the Straits of Magellan, but since obtained at various localities from Cape Cod to West Indies, in 500 to 1,000 fathoms. The identity of these specimens with the original types from South America may be questionable. (*gelatinosus*, jelly-like.)

Melanostigma gelatinosum, GÜNTHER, Proc. Zool. Soc. London 1881, 21, Tilly Bay, Straits of Magellan, in 24 fathoms (Coll. H. M. S. *Alert*, Dr. Coppinger); GÜNTHER, Challenger Report, XXII, 82, 1887; GOODE & BEAN, Bull. Mus. Comp. Zool., X, No. 5, 209, 1883; JORDAN, Cat. Fish. N. A., 125, 1885; GOODE & BEAN, Oceanic Ichthyology, 314, 1896.

2844. MELANOSTIGMA PAMMELAS, Gilbert.

Head 8 in total length; depth 12½; pectoral narrow, its length 2⅝ in head; eye large, 3½ in head; snout short and broad, 7 in head. Well distinguished from *M. gelatinosum* by the wider, blunter head, the smaller, less oblique mouth, the uniform black coloration, and the arrangement of the teeth in the jaws in 2 series. As in *M. gelatinosum*, the head and body are enveloped in a loose, thin skin, which is thrown into folds in alcoholic specimens, and entirely conceals anterior portion of dorsal and anal fins. On dissection the dorsal is seen to have its origin close behind the head, at a point over middle of pectoral fin; anal beginning immediately behind vent, the rays of both fins enveloped in a gelatinous, subcutaneous tissue. Head broad, with its greatest width equaling its greatest depth; mouth broad, somewhat oblique, with equal jaws, the maxillary reaching vertical from front of pupil; each jaw with teeth in two distinct series in front, in a single series laterally in lower jaw, the outer teeth in front enlarged, almost canine-like. Gill opening a small pore above base of pectoral, its diameter about ½ that of eye. Color intense black on head and abdomen, brownish black elsewhere. Length

of type 4½ inches. Coast of southern Alaska. Three other specimens are at hand from *Albatross* Station 3126 (lat. 36° 49′ 20″ N., long. 122° 12′ 30″ W.; depth 456 fathoms). In the smallest, 2½ inches long, the head and abdomen are jet-black, but the rest of the body is only slightly dusky. (Gilbert.) (πᾶς, all; μέλας, black.)

Melanostigma pammelas, Gilbert, Rept. U. S. Fish Comm. 1893 (1896), 472, pl. 35, coast of southern Alaska, at Albatross Station 3202, lat. 36° 46′ 10″ N., long. 121° 58′ 45″ W., in 382 fathoms.

Family CCVII. DEREPODICHTHYIDÆ.

Deep-sea fishes of slender body, scaleless, and without lateral line, somewhat resembling the *Zoarcidæ*, but with each ventral fin reduced to a slender, unbranched filament, the two very closely approximate, and springing from a common projecting base located far forward, below the eye. Gill opening a narrow, vertical slit. Character otherwise given below. A single species known; apparently intermediate between the *Zoarcidæ* and the *Ophidiidæ*.

945. DEREPODICHTHYS, Gilbert.

Derepodichthys, Gilbert, Rept. U. S. Fish Comm. 1893 (1886), 456 (*alepidotus*).

Body slender; no scales; no lateral line; ventral fins reduced each to a slender, unbranched filament, the two very closely approximate, and springing from a common projecting base, which is located far forward below the eye, as *Ophidion*. Gill opening a narrow, vertical slit, little wider than base of pectorals. Teeth cardiform, curved, few in number, in narrow bands or irregular single series on jaws, vomer, and palatines. (δέρη, throat; πούς, foot; ἰχθύς, fish.)

2845. DEREPODICHTHYS ALEPIDOTUS, Gilbert.

Head 8⅓ in total length; depth of head and body 2⅙ in head; width of head 2⅔ in head; distance from tip of snout to base of ventrals 2⅛ in length of head. Distance from tip of snout to front of dorsal 5 ⅓ in total length, from tip of snout to vent 3⅜ in total. Head and body very long and slender, the former resembling a *Lycodes* in appearance, being moderately compressed, with a flattish occiput and a gentle rounded decurved rostral profile. Mouth slightly oblique, quite at lower side of snout; the lower jaw shorter, fitting within the upper; maxillary and premaxillary entirely concealed within the thick skin of the upper lip, which is directly continuous with that of the forehead, the upper jaw being therefore nonprotractile; angle of mouth under front of pupil, its distance from tip of snout 2⅔ in head. Teeth cardiform, curved, few in number, in narrow bands or irregular single series on jaws, vomer, and palatines. Eye small, not filling the elongate orbit, the diameter of exposed portion of eyeball slightly less than ¾ length of snout, the latter 3¼ in head. A series of large mucous pores on snout and lower part of cheeks; a second series on mandible; no pores on body. Gill slit vertical, not continued forward, its lower end slightly above base of lower pectoral rays; length of slit ¼ length of

head, slightly less than distance between slits. Pectorals long and slender, reaching halfway to vent, 1⅓ in head; dorsal and anal confluent with the caudal, concealed in the thick integument, so that the rays can not be counted. Color in spirits, light brownish, the dorsal and pectorals whitish, the anal with a dark margin which becomes black posteriorly; lips dusky; abdominal region blue black. Coast of British Columbia. A single specimen, 4½ inches long, dredged off Queen Charlotte Island. (Gilbert.) (*ἀλεπιδωτός*, scaleless.)

Derepodichthys alepidotus, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 456, Queen Charlotte Island, at Albatross Station 3342, in 1,588 fathoms.

Family CCVIII. OPHIDIIDÆ.

(THE CUSK EELS.)

Body elongate, compressed, more or less eel-shaped, usually covered with very small scales, which are not imbricated, but placed in oblique series at right angles with each other; head large, lower jaw included; both jaws, and usually vomer and palatines also, with villiform or cardiform teeth; premaxillaries protractile; gill openings very wide, the gill membranes separate, anteriorly narrowly joined to the isthmus behind the ventrals; pseudobranchiæ small. Gills 4, a slit behind the fourth; vent more or less posterior. Vertical fins low, without spines, confluent around the tail; tail isocercal; ventral fins at the throat, each developed as a long, forked barbel. Air bladder and pyloric cæca present. To this Dr. Gill adds the following characters, shared more or less by related families: "Orbito-rostral portion of cranium contracted and shorter than the posterior, the cranial cavity closed in part by the expansion and junction of the parasphenoid and frontals, the supraoccipital horizontal and cariniform posteriorly, the exoccipitals expanded backward and upward behind the supraoccipital, the exoccipital condyles contiguous, and with the hypercoracoid (scapula, Parker) fenestrate (or foraminate) about its center, and the hypercoracoid with its inferior process divergent from the proscapula." Genera 7, species about 25. Carnivorous fishes; found in most warm seas, some of them descending to considerable depths, the group especially well represented in tropical America. (*Ophidiidæ*, group *Ophidiina*, Günther, Cat., IV, 376–380, 1862.)

a. Head scaly, at least above; body covered with scales imbricated in quincunx; snout usually with a spine at tip; opercle with or without spinous tip; air bladder, so far as known, ovate, without posterior foramen. LEPOPHIDIUM, 946.

aa. Head scaleless; scales of body rudimentary, scarcely embedded.

b. Air bladder oblong-ovate, not contracted behind, and without posterior foramen.

c. Opercle ending in a flat point behind, without spine. OPHIDION, 947.

cc. Opercle ending behind in a strong spine concealed in the skin. CHILARA, 948.

bb. Air bladder short, thick, reniform or orbicular, with a large foramen behind.

d. Opercle ending in a flat point, without spine. RISSOLA, 949.

dd. Opercle ending behind in a spine concealed in the skin. OTOPHIDIUM, 950.

946. LEPOPHIDIUM, Gill.

Leptophidium, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 210 (*profundorum*), name preoccupied in Serpents by *Leptophidium*, HALLOWELL, 1860.
Lepophidium, GILL, Amer. Nat., Feb., 1895, 16 (*profundorum*).

Body much elongate, moderately compressed, with back and abdominal regions arched, more compressed and slowly decreasing in height backward to an abruptly rounded point; scales regularly imbricated in quincunx oval, and with striæ radiating backward; head with imbricated scales, extending to forehead; snout high, projecting forward, and obtusely rounded, armed above with a short, nearly concealed spine directed forward and somewhat downward, obsolete in 1 species; mouth moderate, oblique; teeth of jaws villiform, immersed in a mucous membrane, separated by an interval from the longer ones in the outer row, which are pointed and usually movable; vomer and palatines with teeth. Deep waters of America on both coasts. Perhaps a fuller knowledge of the species of this genus will lead to its subdivision. (λέπος, scale; *Ophidium*, from the squamation.)

a. Snout without decurved hook or spine; gill rakers 8; head 5 in length; depth 7; pectorals 10; body marbled, the vertical fins edged with black. MARMORATUM, 2846.

aa. Snout with a decurved hook or spine at tip, sometimes more or less concealed in the skin.

b. Gill rakers 7 to 9 in number.

c. Head large, 3¾ to 4⅓ in length.

d. Body stoutish, the depth 6 in length; scales 125; no black blotch on front of dorsal. EMMELAS, 2847.

dd. Body slender, the depth 9¼ in length; dorsal with a black blotch in front; scales 180. STIGMATISTIUM, 2848.

cc. Head moderate, 6 in length; depth 10; vertical fins black-edged.

e. Anterior teeth in jaws movable; pectoral 11 in body; body without white spots. PROFUNDORUM, 2849.

ee. Anterior teeth in jaws not movable; pectoral 13 to 14 in body; body with whitish spots. CERVINUM, 2850.

bb. Gill rakers 4 in number.

f. Scales moderate, 175 to 200 in lateral line.

g. Body without dark cross bars; dorsal and anal margined with black; air bladder oblong.

h. Head 4⅓ in length; depth 8; pectoral 10½. Pacific species. PRORATES, 2851.

hh. Atlantic species imperfectly described. BREVIBARBE, 2852.

gg. Body with dark cross bars; dorsal spotted with black; anal wholly black; head 5⅓ in length; depth 8⅓. PARDALE, 2853.

ff. Scales minute, about 250 in lateral line; head 4⅔; depth 7⅓ to 8; color nearly plain, the fins dark edged. MICROLEPIS, 2854.

2846. LEPOPHIDIUM MARMORATUM (Goode & Bean).

Head 5; depth 7⅓; eye 4 in head; snout about 5. Body somewhat elongate, stoutish anteriorly, gradually tapering; head thickish; interorbital area broad, convex, its width nearly equal to length of snout, which is blunt, spineless; eye circular, somewhat exceeding length of snout. Maxillary extending to vertical through posterior margin of orbit,

the mandible far beyond, its length equal to that of postorbital portion of head. Teeth on vomer and in jaws in villiform bands, the outer series in the latter slightly enlarged. Pseudobranchiæ present; gill rakers short, 8 below angle of first arch, the longest less than ½ diameter of eye. Branchiostegals 7. Ventrals as long as postorbital part of head. Dorsal origin at distance from snout contained 4½ in total length, with 28 rays in a space equal to length of head, counting from the origin of the fin; anal origin separated from snout by distance 2⅜ in total length; length of pectoral 2 in head, or 10 in total. Scales closely imbricated, ornamented with delicate concentric striæ; lateral line apparently complete, located about ¼ distance from dorsal to ventral outline. Color yellowish gray, marbled along the upper half of head and body with olive brown; dorsal and anal fins with black margins. Gulf Stream, in 213 fathoms. (Goode & Bean.) (*marmoratus*, marbled.)

Leptophidium marmoratum, Goode & Bean, Proc. U. S. Nat. Mus. 1885, 423, lat. 23° 10′ 39″ N., long. 82° 20′ 21″ W., in 213 fathoms (Type, No. 37237, U. S. Nat. Mus. Coll. *Albatross*); Goode & Bean, Oceanic Ichthyology, 348, 1896.

2847. LEPOPHIDIUM EMMELAS (Gilbert).

Head 3¾ to 4; depth 5⅝ to 6; eye 4⅛ in head; snout 4⅛; interorbital width 7; vertebræ 13 + 44 = 57; maxillary 2⅛ to 2¼ in head; ventral filament 2⅜; pectoral 2 in head; scales 8–125–18 or 20 before dorsal. Body deep, compressed. Maxillary reaching slightly beyond orbit. Jaws slender and weak, the teeth in very narrow bands, the outer not enlarged. Rostral ridge very sharp, bearing a flat spine at its base directed upward and backward, terminating in a very slender sharp spine anteriorly. Opercle ending in a weak spinous point behind. Gill rakers short and slender, the longest ½ pupil, 8 or 9 movable ones developed. Skull and all bones of head very thin and papery. Dorsal beginning over base of pectorals, the nape midway between its origin and middle of orbit. Distance from snout to origin of anal 1⅓ in distance from latter to end of tail. Scales large, covering cheeks, opercles and top of head forward to middle of interorbital space. Color brownish, much dusted with minute specks; fins blackish, the vertical fins with an indistinct narrow whitish margin; inside of mouth dusky; the roof of mouth, lining of gill cavity, and peritoneum jet-black. Coast of Lower California. Many specimens, the longest 9 inches, from *Albatross* Stations 3007 and 3008, in 362 and 306 fathoms. (Gilbert.) (ἐν, within; μέλας, black.)

Leptophidium emmelas, Gilbert, Proc. U. S. Nat. Mus. 1890, 110, coast of Lower California. (Coll. *Albatross*.)

2848. LEPOPHIDIUM STIGMATISTIUM (Gilbert).

Head 4⅓ in length; depth 9¼; eye 4⅓ in head; snout 6; interorbital width 6½; maxillary reaching slightly beyond posterior border of eyes, 2⅓ in head. A strong rostral spine. Outer teeth scarcely enlarged, evidently so only in front of upper jaw. Gill rakers long and slender, strongly curved forward at tip, the longest equaling ½ eye; 7 well-developed gill-rakers present. Opercle ending in a rounded process, a broad soft flap

projecting beyond it. Dorsal inserted behind middle of pectorals, the nape equidistant from front of dorsal and base of rostral spine; pectorals $2\frac{2}{3}$ in head; longest ventral filament $2\frac{3}{4}$ in head; scales small, about as in *L. prorates*, 180 transverse series, 28 in front of dorsal, continued forward on top of head to front of pupil; cheeks and opercles scaly. Color dusky olivaceous, lighter below; dorsal with a large black blotch on anterior rays, the margin obscurely dusky; anal broadly margined with jet-black; caudal with median rays black at base, the outer rays and the margin light; lining of gill cavity jet-black; inside of mouth white; peritoneum bright silvery. A single specimen 10 inches long. Coast of Lower California. (Gilbert.) Much resembling *L. prorates*, differing in dentition, in gill rakers, and in color. (*στίγμα*, brand; *ἱστίον*, sail.)

Leptophidium stigmatistium, GILBERT, Proc. U. S. Nat. Mus. 1890, 109, off Lower California, at Albatross Station 2996, in 112 fathoms.

2849. LEPOPHIDIUM PROFUNDORUM (Gill).

Head 6; depth 10. Body very slender; scales regularly arranged in quincunx order, those on head extending to forehead, opercles, and cheeks; snout high, projecting, armed with a concealed spinous hook; teeth villiform, separated by an interval from an outer row of longer, slender, movable teeth; eye longer than snout, $3\frac{1}{2}$ in head; lateral line obsolete behind; vent toward end of first third of length; ventral fins short; gill rakers 8. Light rufous; vertical fins margined with black. Gulf Stream, off the coast of Florida. (Gill.) One specimen known. (*profundorum*, of the depths.)

Leptophidium profundorum, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 211, Gulf Stream, off the Coast of Florida (Coll. Commodore Rodgers); GOODE & BEAN, Oceanic Ichthyology, 347, 1896.

Ophidium profundorum, JORDAN & GILBERT, Synopsis, 793, 1883.

2850. LEPOPHIDIUM CERVINUM (Goode & Bean).

Head about $6\frac{1}{4}$; depth about $10\frac{1}{2}$; eye 4 in head; ventrals 3 in head. Body elongate, slender; head slender, somewhat compressed; interorbital area broad, convex, its width equal to length of snout, and $5\frac{1}{3}$ in head; snout sharp, conical, armed with a short but sharp spine, and somewhat overhanging mouth; eye much exceeding length of snout; maxillary extending nearly to vertical through posterior margin of orbit, $2\frac{2}{3}$ in head; mandible extending behind same vertical, its length equal to that of head without postorbital portion. Jaws, vomer, and palatines with narrow bands of villiform teeth, some of which are noticeably enlarged (not movable). Pseudobranchiæ present. Gill rakers short, 8 below angle of first arch, 4 of which are rudimentary, the longest 5 in diameter of eye. (In *L. profundorum* the gill rakers are slenderer and longer, though about equally numerous on the first arch.) Scales in about 11 rows from the origin of the dorsal to the median line of the body. Dorsal origin far back, at a distance from the snout $4\frac{4}{5}$ in total length; at a distance from the eye equal to the head's length. (In *L. profundorum* this distance is $\frac{2}{3}$ of the head's length and the first ray of the dorsal is nearly over the

middle of the extended pectoral; in *L. cervinum*, over its tip, or nearly so.) Distance of anal origin with snout 3 in total length. Length of pectoral 2 in head's length and 13 to 14 in that of body (10 in *L. marmoratum*, 11 in *L. profundorum*). Scales ornamented with radiating striæ, densely covering all parts of the fish except snout, under surface of head, and the fins; lateral line continued almost to end of tail. Color brownish yellow, with numerous subcircular spots of white, with diameter ½ that of eye, along the upper half of body; vertical fins with narrow black margin. Gulf Stream. (Goode & Bean.) A specimen from off Sand Key Light, Florida, recorded by Mr. Garman. (*cervinus*, deer-like, from the faun-color.)

Leptophidium cervinum, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 422, lat. 40° 1′ N., long. 69° 56′ W., depth 76 fathoms (Type, No. 28764. Coll. *Fish Hawk*); GOODE & BEAN, Oceanic Ichthyology, 346, 1896.

Lepophidium cervinum, GARMAN, Bull. Iowa Lab. Nat. Hist. 1896, 91.

2851. LEPOPHIDIUM PRORATES (Jordan & Bollman).

Head 4⅛ to 4⅜ (4⅜ to 4⅔ in total); depth 7⅛ to 8⅕ (7⅗ to 8½); eye 4⅓ in head; snout 5; maxillary 2¼; interorbital 1⅗ in eye; pectoral 2⅛ in head; inner ventral filament shortest, the longer 2⅜ in head. Body moderately elongate, compressed, considerably stouter than in *L. profundorum*. Mouth large, maxillary reaching about ½ pupil's length beyond posterior border of eye. Outer teeth slightly enlarged, a little movable, those of upper jaw largest. Gill rakers rather long and slender, ¼ length of eye, 4 developed. Tip of snout with a strong spine directed forward and slightly downward; opercle without spine, ending in a flat projection covered by skin. Dorsal beginning over middle of pectorals, longest ray 4 in head. Scales regularly imbricated, but very small, about 225 in a longitudinal series; scales on top of head extending forward to base of ethmoid spine; sides of head covered with small scales; lateral line not reaching end of tail. Air bladder oblong-lanceolate. Color olivaceous, paler below; scales rather profusely dotted with black; a pale shade across opercles; lower jaw, gular region, and anterior branchiostegals dusted; dorsal and anal margined with black, the band on anal the broader; pectorals pale. Specimens of this species were obtained at Panama and at *Albatross* Station 2801, south of Panama. Length of type 10 inches. (*πρωρᾱ́της*, prow-bearing, from the rostral spine.)

Leptophidium prorates, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 172, Panama. (Type, No 41149, U. S. Nat. Mus. Coll. *Albatross*.)

2852. LEPOPHIDIUM BREVIBARBE (Cuvier).

A short decurved spine at tip of snout; teeth strong; occiput and opercles scaly. Vertical fins edged with black. (Kaup.) Air bladder oblong ovate, without contracted portion and without posterior foramen; no single anterior bone replaced by cartilage. (Müller.) West Indies and Brazil; a scarcely known species; apparently close to *L. prorates*, but very insufficiently described. (*brevis*, short; *barba*, beard.)

Ophidion brevibarbe, CUVIER, Règne Animal, Ed. 2, vol. II, 358, 1829, Brazil; MÜLLER, Abhandl. Berl. Acad. 1843, 153, pl. 4, f. 4; KAUP, Apodes, 154, pl. 16, f. 1; GÜNTHER, Cat. Fish., IV, 379, 1862.

2853. LEPOPHIDIUM PARDALE (Gilbert).

Head $5\frac{1}{2}$ in length; depth $8\frac{1}{3}$; eye $3\frac{3}{5}$ in head; snout $4\frac{3}{5}$; interorbital $1\frac{1}{2}$ in eye. Body very slender, with a short head and small mouth; maxillary scarcely reaching vertical from posterior border of orbit, its length $2\frac{1}{5}$ in head; outer teeth very little enlarged, not movable; teeth present on jaws, vomer, and palatines. Gill rakers slender, the longest $\frac{1}{4}$ eye, 4 developed. Tip of snout with a strong, concealed spine, as in *L. prorates*. Opercle ending in a short spine. Nape midway between front of dorsal and front of pupil; dorsal beginning over middle of pectorals, which are $2\frac{1}{4}$ in head; ventral filaments very short; the inner the longer, $4\frac{3}{5}$ in head. Scales very small, about 200 in a longitudinal series, extending forward on top of head to middle of interorbital space; cheeks and opercles scaly. Light olive, a series of 8 black bars downward from back, scarcely reaching lateral line, sometimes continuous with their fellows of the other side, and alternating with smaller black spots on dorsal outline; below the smaller spots a series of round spots nearly as large as eye along middle line of sides; sides and lower parts of head and body dusted with rather coarse black specks; dorsal light, the margin with 10 elongate black blotches, usually longer than the interspaces; caudal dusky at base, its distal half white; anal wholly black; peritoneum and lining of gill cavity white. Lower California. A single specimen, length $7\frac{1}{8}$ inches, from *Albatross* Station 3014, in 29 fathoms. (Gilbert.) (*πάρδαλις*, leopard.)

Leptophidium pardale, GILBERT, Proc. U. S. Nat. Mus. 1890, 108, off Lower California. (Type, No. 44382. Coll. Dr. Gilbert.)

2854. LEPOPHIDIUM MICROLEPIS (Gilbert).

Head $4\frac{2}{5}$ in length; depth $7\frac{1}{3}$ to 8; eye $4\frac{1}{4}$ to 5 in head; snout 5; interorbital width $6\frac{1}{3}$; maxillary extending beyond orbit, $2\frac{1}{4}$ to $2\frac{1}{3}$. Rostral spine very strong, as in *L. prorates*. Outer teeth enlarged, not at all movable, those in upper jaw largest. Four gill rakers developed, the longest $3\frac{1}{2}$ in eye. Opercle ending in a short concealed spinous point. Dorsal inserted in front of middle of pectorals, the distance from nape to front of dorsal usually less than from nape to middle of eye; longest ventral filament $3\frac{1}{8}$ to $3\frac{1}{2}$ in length of head; pectorals $2\frac{1}{4}$ to $2\frac{1}{3}$ in head. Scales exceedingly small, regularly imbricated, in about 250 transverse series, 35 transverse series between nape and dorsal (about 175 transverse rows in *L. prorates*, 25 series between nape and dorsal). Top of head scaly as far as front of eyes. Cheeks and opercles scaly. Color as in *L. prorates*, the lining of peritoneum and gill cavity silvery white, the former with little or no black specking. Closely related to *L. prorates*, differing principally in the much smaller scales. Gulf of California. Many specimens, the longest 14 inches, from *Albatross* Stations 3015 and 3016, in 145 and 76 fathoms. (Gilbert.) (*μικρός*, small; *λεπίς*, scale.)

Leptophidium microlepis, GILBERT, Proc. U. S. Nat. Mus. 1890, 109, Gulf of California. (Coll. Dr. Gilbert.)

947. OPHIDION (Artedi) Linnæus.

(Cusk Eels.)

Ophidion (Artedi) Linnæus, Syst. Nat., Ed. x, 259, 1758 (*barbatum*).

Ophidium, Linnæus, Syst. Nat., Ed. xii, 431, 1766, and of most recent authors; changed spelling.

Body moderately elongate, compressed; scales small, usually not imbricated, but arranged in short, oblique series, often placed at right angles with each other, much as in *Anguilla*. Head naked; teeth villiform, those of the outer series more or less enlarged, none of them movable; teeth on vomer and palatines bluntish, some of them enlarged. Vent well behind pectorals. Opercle without distinct spine; sometimes (*O. barbatum*) a distinct spine at tip of snout. Air bladder oblong-ovate, tapering behind, without foramen. Shore species mostly European. (*Ophidium*, an ancient name, from *ὀφίδιον*, a small snake.)

a. Gill rakers 4.
  b. Head 4⅔ in length; depth 7; fins not dark edged. BEANI, 2855.
  bb. Head 6 in length; inner ray of ventral 1½ in length of outer, which is shorter than head; fins dark-edged. HOLBROOKI, 2856.
aa. Gill rakers 6 or 7; head 5½ to 6 in length; depth 8 to 10; color silvery, unspotted; fins not dark-edged; ventrals nearly as long as head. GRAELLSI, 2857.

2855. OPHIDION BEANI, Jordan & Gilbert.

Head 4⅔ in length; depth about 7. Head small, the profile not very obtuse; snout 4⅔ in head; eye 3¼, more than twice the narrow interorbital space; mouth oblique, the maxillary reaching to posterior border of pupil, 2 in head; lower jaw slightly included; teeth small, in narrow bands in the jaws, the outer series in upper jaw somewhat enlarged; vomerine and palatine teeth small, subequal; head naked; snout spineless; opercle without spine; no evident pseudobranchiæ; gill rakers rather long and strong, 4 below angle of arch; occiput nearly midway between origin of dorsal and front of eye. Air bladder long and slender, occupying nearly the whole length of abdominal cavity, tapering backward. Very light olive, somewhat punctate above, slightly silvery below; fins without trace of dark edging (but being mutilated they may have been dark-edged in life). Gulf of Mexico. Two specimens, 1 of which is in good condition and about 4 inches long, were taken from the stomach of a red snapper at Pensacola. (Named for Dr. Tarleton Hoffman Bean.)

Ophidium grœllsi, Jordan & Gilbert, Proc. U. S. Nat. Mus. 1882, 301; Jordan & Gilbert, Synopsis, 963, 1883; not of Poey.

Ophidion beani. Jordan & Gilbert, Proc. U. S. Nat. Mus. 1883, 43, Snapper Banks off Pensacola (Coll. Jordan & Stearns. Type, No. 30868, U. S. Nat. Mus.); Jordan, Cat. Fishes N. A., 126, 1885.

2856. OPHIDION HOLBROOKI (Putnam).

Head 6 in total length. Inner barbel nearly ⅔ length of the outer; outer barbel equal to distance from center of eye to point of operculum; maxillary reaching to posterior border of eye. Length of eye equal to distance from its posterior margin to ridge of preoperculum. Dorsal and anal

with a black margin. Gill rakers 4; air bladder long, pointed, with a foramen. (Putnam.) Length 6 inches. Gulf of Mexico; recorded from Key West, Florida; not seen by us. (Named for Dr. John Edwards Holbrook, the distinguished ichthyologist of Charleston.)

?*Ophidion josephi*, GIRARD, U. S. and Mex. Bound. Surv., Ichth., 29, 1859, St. Joseph Island, Texas; JORDAN & GILBERT, Synopsis 793, 1883; quite as likely to be *Rissola marginata*.

Ophidium holbrooki, PUTNAM, Proc. Bost. Soc. Nat. Hist. 1874, 342, Key West, Florida; JORDAN & GILBERT, Synopsis, 793, 1883.

2857. OPHIDION GRAELLSI, Poey.

Head $5\frac{3}{4}$ in body; depth 10; eye 3 in head; pectoral $2\frac{1}{5}$; ventral scarcely as long as head. Body elongate, compressed; mouth large, the maxillary reaching to posterior margin of pupil; small teeth on jaws, vomer, and palatines; eye very large, greater than length of snout; interorbital space $\frac{3}{8}$ of eye, a sharp ridge along its middle to tip of snout, where it ends in a sharp spine; opercles unarmed; about 6 gill rakers developed on lower part of gill arch, apparently none above; pseudobranchiæ small, if present. Air bladder, injured in specimen examined, apparently lanceolate; dorsal and anal low, confluent with caudal, which ends in a point; pectorals small, their ends scarcely reaching midway from their base to front of anal; ventrals with 2 filamentous rays, the outer scarcely as long as head, the inner $\frac{1}{2}$ as long. Color in spirits, reddish brown, with silvery reflections on sides; head silvery, upper part of eye black; fins the color of the body, with no dark edgings. Coasts of Cuba; rare. Here described from a specimen from Havana, Cuba, $2\frac{1}{2}$ inches in length, sent by Professor Poey. Poey has also sent a drawing of his original type, a much larger specimen, which he describes as follows:

Head $5\frac{1}{5}$; depth 8; snout rounded; eye large, $4\frac{1}{2}$ in head; maxillary reaching posterior border of eye; teeth small, slender, with a villiform band behind them; teeth on vomer and palatines; scales small; head scaly, except on snout; lateral line high; branchiostegals 7; dorsal beginning over second third of pectoral, joining anal behind; about 100 rays in each fin; vent a little behind first third of length. Yellowish brown, silvery on side of head; no black on fins. Air bladder distinct; no pyloric cæca. Intestine with 2 short turns. Cuba. (Poey.) Air bladder, gill rakers, and ventral not described. Length 230 mm. Rare; not reaching a foot in length. (Named for Mariano de la P. Graëlls, director of the Botanic Garden at Madrid, "comme témoignage de mon estime pour ses travaux scientifiques, et pour la zèle qu'il déploit . . . pour l'acquisition des objets et l'acclimatation des espèces.")

Ophidion graellsi, POEY, Memorias, II, 425, 1860, Havana (Coll. Poey); POEY, Synopsis, 402, 1867.

948. CHILARA, Jordan & Evermann.

Chilara, JORDAN & EVERMANN, Check-List Fish. N. and M. A., 482, 1896 (*taylori*).

This genus contains a single robust species which differs from *Ophidion* only in the presence of a stout concealed spine at tip of opercle; the air bladder

is oblong-ovate, the head naked and the snout without spine. (χιλάρι, the modern Greek name of the species of *Ophidion* and *Rissola*.)

2858. CHILARA TAYLORI (Girard).

Head 6; depth 8; head large, little compressed, naked; top of head with conspicuous mucous pores; dorsal fin beginning over the pectorals; outer ray of ventral little more than ½ length of head, inner about ⅓; air bladder ovate, not contracted; 7 gill rakers below the angle of the arch; pseudobranchiæ developed; no spine on the end of the snout; opercle with a flat spine concealed in its membranes; outer teeth in both jaws considerably enlarged, the upper largest. Color light olive; head and upper parts covered with conspicuous round dark, olive-brown spots; chin dusky; vertical fins edged with black. Length 12 inches. Coast of California, from Monterey to San Diego; not rare in waters of moderate depth. (Named for A. S. Taylor, its discoverer.)

Ophidium taylori, Girard, Pac. R. R. Surv., x, Fishes, 138, 1858, Monterey, California (Type, No. 867. Coll. A. S. Taylor); Jordan & Gilbert, Synopsis, 793, 1883.

949. RISSOLA, Jordan & Evermann.

Rissola, Jordan & Evermann, Check-List Fish. N. and M. A., 483, 1896 (*marginatum*).

This genus contains species agreeing with *Ophidion* in general characters, but with the air bladder short, broad, spherical or kidney-shaped, with a posterior foramen. Species chiefly of the Mediterranean. (Named for Anastase Risso, apothecary at Nice, author of the Ichthyologie de Nice, 1810, and Histoire Naturelle de l'Europe Méridionale, 1826, two of the very best of local faunal works, the foundation of our knowledge of the fishes of the Mediterranean.)

2859. RISSOLA MARGINATA (DeKay).

Head 6½; depth 7½; eye 4 in head; maxillary reaching posterior margin of orbit; air bladder short and broad, with a foramen on the under side; upper ray of ventral about equaling length of head; inner ray ½ length of outer; gill rakers 4; color nearly plain brownish; dorsal and anal fins margined with black. Coast of the United States, from New York south to Pensacola and the coast of Texas; not very common; very similar to the Mediterranean species *Rissola rochii* (Müller), but probably distinct. (*marginatus*, margined.)

Ophidium marginatum, De Kay, N. Y. Fauna: Fish., 315, 1842, New York Harbor; Putnam, Proc. Bost. Soc. Nat. Hist. 1874, 342; Jordan & Gilbert, Synopsis, 792, 1883.

?*Ophidium josephi*, * Girard, U. S. and Mex. Bound. Surv., Zool., 29, 1859, Saint Joseph Island, Texas; Jordan & Gilbert, Synopsis, 793, 1883.

* The scanty description of *Ophidion josephi* agrees fairly with either *Rissola marginata* or *Ophidion holbrooki*, and may be either. The following is the substance of Girard's account:

"Head 6 in length; eye moderate, 4 in head; maxillary extending to opposite its posterior margin; origin of dorsal at some distance behind base of pectorals. Body shorter and pectorals more elongate than in *O. taylori*. Pale olive, sprinkled all over with brownish specks; belly and sides of head plain; vertical fins edged with black."

950. OTOPHIDIUM, Gill.

Otophidium, GILL, in JORDAN, Cat. Fish. N. A., 126, 1885 (*omostigma*).

This genus differs from *Ophidion*, in the form of the air bladder, which is short, thick, and with a large foramen (not examined in *O. galeoides*). The opercle ends in a concealed spine as in *Chilara*. Species American, so far as known. (*οὖς ὠτός*, ear; *Ophidium*.)

a. Gill rakers 4.
  b. Head long, 4⅓ to 4½ in length; depth 5⅓ to 6.
    c. Scapular region with a jet-black spot; pseudobranchiæ little developed; ventrals ⅓ length of head; maxillary 1⅔ in head. OMOSTIGMUM, 2860.
    cc. Scapular region without jet-black spot; pseudobranchiæ well developed; ventrals with the inner ray longest, ⅓ head; body with dark cross bands. INDEFATIGABILE, 2861.
  bb. Head moderate, 5⅓ in body; depth 6; a pale spot before dorsal; pale spots along lateral line; ventral ⅓ head. GALEOIDES, 2862.

2860. OTOPHIDIUM OMOSTIGMUM (Jordan & Gilbert).

Head 4⅓ in length; depth about 6. Body comparatively short, highest at occiput, thence tapering rapidly to tip of tail; upper profile of head very convex; snout blunt; mouth horizontal, the lower jaw included; maxillary not quite reaching posterior border of orbit; teeth in jaws uniform, strongly incurved, in rather broad bands; a single series of small teeth on vomer, those on palatines minute; maxillary 1⅔ in head; eye large, 3 in head, much larger than snout, equaling twice interorbital width; opercle terminating in a strong, compressed spine, the length of which is about ⅔ diameter of pupil; gill rakers very small, 4 below on anterior arch. Longest ventral filament ½ length of head; the shorter ⅔ length of longer. Distance from origin of dorsal to tip of snout 3⅓ in total length; distance from origin of anal to snout 2⅓ in total length. Scales minute, embedded. Pseudobranchiæ probably present (type reexamined by us). Air bladder short, thick, with a large posterior foramen. Color light olive green, silvery on belly, cheeks, and lower side of head; sides above with a few irregular, large, scattered, dark blotches, about 9 of these along base of dorsal fin; an intensely black, round blotch on scapular region, rather larger than pupil; dorsal with black blotches; anal largely black; upper half of eye black, lower half bright silvery. Gulf of Mexico. A single specimen, 3½ inches long, taken from the stomach of a red snapper, at the Snapper Banks off Pensacola. (*ὦμος*, shoulder; *στίγμα*, spot.)

Genypterus omostigma, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 301, Pensacola Snapper Banks (Coll. Jordan & Stearns. Type, 29670, U. S. Nat. Mus.); JORDAN & GILBERT, Synopsis, 963, 1883.

Otophidium omostigma, GOODE & BEAN, Oceanic Ichthyology, 345, fig. 305, 1896.

2861. OTOPHIDIUM INDEFATIGABILE, Jordan & Bollman.

Head 4⅔ (4½ in total); depth 5⅔ (5⅞); eye large, 3 in head; snout 4. Body rather short, compressed, width of nape 2⅓ in head. Mouth large; maxillary reaching to opposite posterior margin of pupil, 1⅓ in head; outer row of teeth of each jaw very slightly enlarged. Interorbital space 2 in eye; interorbital area with a thin crest under the skin, this ending in 2 com-

pressed spines, 1 turning forward, the other backward over front of eye, these spines concealed by the skin. Gill rakers short and thick, less than ½ pupil, 4 developed. Dorsal beginning at end of anterior third of pectorals, longest ray 3½ in head; pectorals 2 in head; inner ventral filament longest, 2 in head. Air bladder short and thick, with a foramen. Scales very small, more or less imbricated on body; head naked. Opercle with a sharp, partly concealed spine. Pseudobranchiæ present. Color pale yellowish brown, silvery on belly and sides of head; back with about 12 irregular dark cross bands, the alternate ones being narrower and broken up into spots, 2 before dorsal; a few scattered spots about as large as pupil on sides, these most distinct about the shoulder; dorsal pale, first rays black, and with 3 or 4 other black blotches on upper part; anal black, margined with white; pectorals pale, axil dusky; caudal and posterior part of anal pale; chin pale. A single specimen obtained at Indefatigable Island, Galapagos Archipelago. Length 4 inches. (*indefatigabilis*, tireless.)

Otophidium indefatigabile, Jordan & Bollman, Proc. U. S. Nat. Mus. 1889, 172, Indefatigable Island, in the Galapagos Archipelago. (Type, No. 44393. Coll. *Albatross*.)

2862. OTOPHIDIUM GALEOIDES (Gilbert).

Head 5⅓ in length; depth 6. D. 125. Maxillary reaching beyond pupil, 2⅓ in head; snout 4⅔; eye 3⅔. Gill rakers short and broad, 4 of them developed. Opercle ending in a sharp concealed spine. Outer teeth little enlarged. Dorsal beginning over middle of pectorals, the nape equidistant between front of dorsal and tip of snout. Caudal very short and bluntly rounded, as in *Chilara taylori*, the rays not projecting beyond dorsal and anal; pectorals 1⅔ in head; ventral filament ½ head. Scales as in *C. taylori*, not at all imbricated, arranged with their long axes frequently at right angles to each other; head naked. Color light olive, without bars, a narrow dusky streak along base of dorsal, and a round light spot at origin of dorsal; a series of small olive-brown spots along lateral line, with a few scattering spots below it but none above; nape and head without spots; vertical fins translucent; dorsal with a large black blotch on tip of anterior rays, the fin behind this narrowly edged with black, which does not surround the caudal; anal with much silvery-white pigment anteriorly on distal portion, becoming dusky behind; pectorals translucent, edged with white below; peritoneum, buccal, and gill cavities white. Closely related to *Otophidium indefatigabile*, differing in color, and in the much shorter head, smaller mouth, less imbricated scales, the more posterior insertion of dorsal, and the absence of spines on head. Air bladder not examined. Pseudobranchiæ present. Gulf of California. One specimen, 5½ inches long, from *Albatross* Station 3025, in 9½ fathoms. (Gilbert.) (γαλῆ, shark; εἶδος, appearance.)

Otophidion galeoides, Gilbert, Proc. U. S. Nat. Mus. 1890, 110, Gulf of California, lat. 31° 21′ 15″ N., long. 113° 59′ W. (Type, No. 44381. Coll. *Albatross*.)

Family CCIX. LYCODAPODIDÆ.

Deep-sea fishes allied to the *Fierasferidæ*, differing chiefly in the normal position of the vent, which is remote from the head, and just before the

anal fin; gill openings large, the membranes united anteriorly only, free from the isthmus, as in *Fierasfer*. Pseudobranchiæ wanting; no scales; no lateral line; no ventral fins. One genus with 4 known species, from the North Pacific.

951. LYCODAPUS, Gilbert.

Lycodapus, GILBERT, Proc. U. S. Nat. Mus. 1890, 107 (*fierasfer*).

Body naked. Ventrals wanting. Vertical fins united around the tail. Gill openings wide, continued forward under the throat; the gill membranes anteriorly narrowly united, loosely joined to the isthmus by a fold of lax skin. Branchiostegals 6. No pseudobranchiæ. Gills 4, a wide slit behind inner arch. Gill rakers developed. Teeth present in jaws and on vomer and palatines, none of them enlarged. Vent remote from the throat. (*Lycodes; ἄπους*, footless.)

a. Body slender, the depth 8 to 11 in length.
b. Head rather large, 4⅜ to 5⅓ in length.
c. Head, body, and fins with very many mucous pores; dorsal rays 70; anal 60. DERMATINUS, 2863.
cc. Head, body, and fins with very few mucous pores; dorsal rays 82; anal 70. FIERASFER, 2864.
bb. Head small, 7⅜ in length; gill openings not extending above base of pectorals. PARVICEPS, 2865.
aa. Body very slender, the depth about 15 in length; dorsal rays about 100. EXTENSUS, 2866.

2863. LYCODAPUS DERMATINUS, Gilbert.

Head 4¾; depth 1⅛ in head; eye 5 in head; snout 4; maxillary 2½. D. 70; A. 60. Very similar to *L. fierasfer*, but the head, body and fins covered with a thick loose skin which contains numerous pores, or openings for the mucous canals. One series of these runs along middle of sides and forms the lateral line; it rises anteriorly above the gill opening, and is continued forward on top of head, the two meeting between eyes; a second series runs between eye and upper lip, and curves around on middle of cheek, running upward to behind eye; one series runs along a fold bordering mandible, 1 along preopercular margin, and 1 on opercle. In *L. fierasfer* a few pores are visible on mandible, and 1 or 2 can frequently be made out on preopercular margins. The skin is very thin and delicate, and the fin rays are very evident through the membrane. The general proportions and the dentition of the type are essentially as in *L. fierasfer*, but the vomerine teeth are long and hooked backward. Mandible heavier than in *L. fierasfer*. Origin of dorsal vertically above axil of pectorals. Length of head and trunk ⅓ total length. Teeth in narrow bands in the jaws, a single series on vomer and palatines. Gill membranes very narrowly joined below and free from the isthmus, as in *L. fierasfer*. Pectorals much longer than in *L. fierasfer*. General color in spirits light brownish yellow, made somewhat dusky by the pigment spots in the skin; body, and especially the fins, darker posteriorly. Aleutian

Islands, in deep water. Only the type known, an example 4¾ inches long. (δέρμα, skin; *dermatinus*, skinny.)

Lycodapus dermatinus, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 471, pl. 35, Aleutian Islands, lat. 37° 54′ 10″ N., long. 123° 30′ W., at Albatross Station 3162, in 552 fathoms.

2864. LYCODAPUS FIERASFER, Gilbert.

Head 5½; depth 10; eye 4½ in head; snout 3⅓; maxillary 2 to 2½. D. 82; A. 70. Body compressed, elongate, tapering rather rapidly backward, the tail not produced to a filament; head flat above, the cheeks deep, vertical, the mouth very oblique, with the lower jaw slightly the longer and nearly entering the upper profile; skull very thin and papery, translucent; jaws weak; gape of mouth wide, the maxillary reaching vertical from behind front of pupil; teeth all small, in a very narrow band in jaws, in a single series on vomer and palatines; interorbital width ⅔ of eye; snout broad, depressed, spatulate, its tip prominent, turned upward, the upper profile thus longitudinally concave; an evident median ridge on snout and interorbital space; gill slits continued forward below to vertical from middle of eye, the membranes united for a distance equaling diameter of pupil; gill rakers short, less than diameter of pupil, strongly toothed, about 10 on horizontal limb of arch; head without conspicuous mucous pores or cavities. Dorsal beginning well forward, its distance from occiput slightly less than that from occiput to nostril. Dorsal and anal rays slender, all articulated, branched only at tips; caudal not distinct, the rays springing from end of tail not projecting beyond the others; origin of anal immediately behind vent, its distance from snout nearly equaling ⅛ total length; pectorals narrow, varying in length, about 2½ in head. Body and fins invested in a rather lax transparent skin, without traces of scales. Color, body translucent, dusted with black specks; abdomen blackish; lips, inside of mouth, lining of gill cavity, and peritoneum jet-black; iris silvery. (Gilbert.) North Pacific. The types, several specimens, the longest 5¼ inches, from *Albatross* Stations 2980, 3010, 3072, off Lower California, in 610 to 1,005 fathoms. Also taken near Unalaska in 109 fathoms. (*Fierasfer*, the pearlfish.)

Lycodapus fierasfer, GILBERT, Proc. U. S. Nat. Mus. 1890, 108, off Lower California, in 610 to 1,005 fathoms (Coll. Dr. Gilbert); JORDAN, Proc. Cal. Ac. Sci. 1896, 234, pl. 23.

2865. LYCODAPUS PARVICEPS, Gilbert.

Head 7⅔; depth 11; eye 4½ in head; snout 3¼; least interorbital width 5; maxillary 2⅙; pectoral 2¾. D. 100; A. about 85 (both counted to middle of caudal); P. 9; no ventrals. Upper profile of head nearly straight, not longitudinally concave as in *L. fierasfer;* head deeper and narrower, the snout less spatulate; skin thicker. A conspicuous series of pores on mandible and along preopercular margin; gill slit very oblique, extending anteriorly as far as vertical from eye, the membranes then narrowly united, free from the isthmus except at extreme front; gill slit superiorly much more restricted than in *L. fierasfer*, not extending above base of pectorals, while in the latter it extends above them for ⅔ diameter of eye. Mouth

oblique, maxillary reaching vertical from middle of eye; jaws even at tip, the mandible slightly included laterally; mandibular teeth in a moderate band anteriorly, the inner series enlarged, narrowing posteriorly to a single row; premaxillary teeth of uniform size, in a narrow band throughout; vomer with 4 canine-like teeth; palatine teeth small, in a single close-set series. Distance from origin of dorsal to occiput slightly less than that from occiput to posterior nostril; head and trunk contained $3\frac{1}{5}$ in tail. Body brownish in spirits, fins whitish, translucent; everywhere dusted with black specks; tail and fins distinctly blackish posteriorly; orbit blackish above; gill cavity silvery, blackish anteriorly; mouth blackish, except anteriorly; peritoneum black, the color not showing through the abdominal wall. Similar to *L. fierasfer*, differing in the much smaller head, longer, slenderer body, the thicker skin with more evident mucous pores, and in the more restricted gill openings. Aleutian Islands, in moderately deep water. Only the type known, a specimen about 5 inches long. (Gilbert.) (*parvus*, small; *-ceps*, head.)

Lycodapus parviceps, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 455, north of Unalaska Island at Albatross Station 3324, in 109 fathoms. (Coll. Dr. Gilbert.)

2866. LYCODAPUS EXTENSUS (Gilbert).

Head $6\frac{2}{4}$; depth $15\frac{1}{8}$; eye $4\frac{1}{4}$ in head; snout $3\frac{2}{5}$; interorbital width $1\frac{1}{8}$ in eye; pectoral $2\frac{2}{5}$ in head. D. 96 (the extreme end of the tail wanting). Gill openings as in *L. fierasfer*, extending well above base of ventrals. Skin thin, the mucous pores inconspicuous, evident on mandible and along margin of preopercle. Upper profile of head longitudinally concave, shaped as in *L. fierasfer*, but slenderer, its depth greater than that of body. Mouth oblique, the maxillary reaching vertical from middle of eye, $2\frac{1}{5}$ in head. Teeth in narrow bands in each jaw, tapering laterally to single series; vomerine teeth more numerous than in *L. parviceps* or *L. fierasfer*, small, not canine-like, in a single series; palatine teeth wanting, as in some individuals of *L. fierasfer*. Occiput midway between front of dorsal and anterior nostril; pectorals slenderer and longer than in *L. fierasfer*. Head and trunk contained $2\frac{3}{4}$ times in tail. Color light brownish, the black peritoneum visible through the skin of the abdomen; mouth and gill cavity largely dusky; a narrow dark-brown streak along base of dorsal and anal, occupying, toward tip of tail, the entire height of both fins. An extremely slender elongate form, with head smaller than *L. fierasfer*, but otherwise resembling that species more than *L. parviceps*. Aleutian Islands, in rather deep water. Only the type, a specimen 4 inches long, known. (Gilbert.) (*extensus*, stretched out.)

Lycodalepis extensus, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 455, north of Unalaska, at Albatross Station 3324, in 109 fathoms. (Coll. Dr. Gilbert.)

Family CCX. FIERASFERIDÆ.

(THE PEARL-FISHES.)

Body elongate, compressed, tapering into a long and slender tail; no scales; teeth cardiform, on jaws, vomer, and palatines; canine teeth often present; no barbels; lower jaw included; vent at the throat; gill mem-

branes somewhat united, free from the isthmus; no pseudobranchiæ; no pyloric cæca; vertical fins very low, confluent, without spines; no ventral fins; pectoral fins present or absent. Small shore fishes of tropical seas, often living in shells of mollusks, echinoderms, etc., being especially often commensal with the pearl oyster and with the larger *Holothuria*. Genera 3; species 12. (*Ophidiidæ*, group *Fierasferina*, Günther, Cat., IV, 381–384, 1862.)

a. Pectoral fins present; no distinct caudal fin; gill membranes connected anteriorly only. FIERASFER, 952.

952. FIERASFER, Cuvier.

Fierasfer, CUVIER, Règne Anim., Ed. 1, II, 239, 1817 (*imberbe=acus*).
Echiodon, THOMPSON, Proc. Zool. Soc. London 1837, 55 (*drummondi*).
Diaphasia, LOWE, Proc. Zool. Soc. London 1843, 92 (*acus*).
Oxybeles, RICHARDSON, Voy. Erebus and Terror, Fishes, 74, 1844–48 (*homei*).
Porobronchus, KAUP, Ann. Mag. Nat. Hist. 1860, 272 (larva of *Fierasfer acus*).
Carapus,* GILL, Proc. Ac. Nat. Sci. Phila. 1864, 152 (after RAFINESQUE, 1810; not type).
Vexillifer, GASCO, Bull. Assoc. Nat. Med. Napoli 1870, 59 (larva of *Fierasfer acus*).
Lefroyia, JONES, Zoologist, IX, 1874, 3838 (*bermudensis*).

Gill membranes little connected, leaving the isthmus bare. No distinct caudal fin; pectoral fins developed. The species of this genus are not well known, and their characters and nomenclatures are uncertain. It is not unlikely that the American species are all reducible to one, *Fierasfer affinis* or *dubius*, but our scanty material will not justify us in taking this view. (*Fierasfer*, the ancient name, from φιερός, sleek and shining.)

a. Vomer with canine teeth; pectoral about ½ length of head.
b. Front teeth of upper jaw enlarged; head 7 to 8 in length; depth 11½ to 15 times in length of body. AFFINIS, 2867.
bb. Front teeth of upper jaw not enlarged; head 6½ in length; depth about 10½ times in length of body. ARENICOLA, 2868.
aa. Vomer with small teeth, scarcely canine-like; pectoral about 2½ in head; head 7 to 8½ in body. BERMUDENSIS, 2869.

2867. FIERASFER AFFINIS† (Günther).

(PEARL-FISH.)

Head 7½; depth of head 15. Maxillary extending slightly beyond orbit; lower teeth larger than the upper, except 2 to 4 front teeth of upper jaw, which are about equal to lower teeth; vomer with 3 to 6 teeth, 2 or 3 of

* The name *Carapus*, Rafinesque, has been substituted for *Fierasfer* by Gill and Poey. This change seems to us not justifiable, as it is certainly not desirable. The name *Carapus* first appears in Rafinesque's Indice d'Iltiologia Siciliana, 57, 1810. No type is mentioned by Rafinesque, but the diagnosis is taken from that of Lacépède's second subgenus under *Gymnotus*, which contains the three species, *carapo*, *fierasfer*, and *longirostratus*. Of these species, *carapo* is the original Linnæan type (Ed. X) of the genus *Gymnotus*. *Carapus* should therefore be regarded as a synonym of *Gymnotus*. The Brazilian name *carapo* evidently suggested the word *Carapus*, although Dr. Gill derives the name from κάρα, head; ἄπους, footless, an ex post facto distinction from *Ophedion*. In a list of Sicilian fishes, on page 37 of Rafinesque's Indice, published somewhat later, the name *Carapus acus* appears for *Fierasfer acus*. This reference of a species of *Fierasfer* to *Gymnotus* or *Carapus* was due to Rafinesque's ignorance of its relations.

† In the Museum of Comparative Zoology is "one valve of a pearl oyster, in which a specimen of *Fierasfer dubius* is beautifully inclosed in a pearly covering, deposited on it by the oyster." (Putnam.)

these canine-like. Pectoral ½ head; vent under base of pectoral. Dorsal fin low, but distinct; anal much more developed than dorsal, its longest rays about in the middle of the fish. Air bladder long, slightly constricted behind. Gill membranes not covering isthmus. Color in spirits, uniform light brown, with a short silvery band along the sides of the abdomen made by confluent spots. (Putnam, description of *F. dubius*.) Panama; especially common among the Pearl Islands, chiefly in shells of pearl oysters. This species should probably stand as *Fierasfer affinis*.

The following notes are from numerous specimens, 3 to 4 inches long, from Pearl Islands, collected by Prof. Bradley, these also being types of *Fierasfer dubius:* Head 6¾ to 7⅛; eye 4½ to 5 in head. Teeth in upper jaw small, acute, in a rather narrow band; sometimes a few in the front of the jaw inconspicuously enlarged; those in lower jaw and on palatines conic, blunt, in somewhat wider bands, the outer series of lower jaw enlarged, canine-like; vomer with a narrowly oblong patch of small, blunt teeth, surrounding a median series of 3 to 6 conspicuously enlarged, retrorsely curved canines, which are usually much the largest teeth in the mouth. Two specimens from *Albatross* Station 3021, Lower California, agree in general with the above account: Head 7⅛; depth 11⅛; eye 4; 2 upper teeth on each side somewhat enlarged, about as large as lateral teeth on mandibles; vomerine canines larger. Professor Putnam refers also to *Fierasfer dubius* specimens from Key Biscayne, Florida (Coll. Theodore Lyman); Tortugas (Coll. Gustav Würdemann); Cape Florida (Coll. Würdemann), and New Providence, Bahama (Coll. F. G. Shaw). These specimens apparently belong rather to *Fierasfer bermudensis*, if that species be different. (*affinis*, related, to *Fierasfer acus*.)

? *Fierasfer affinis*,* GÜNTHER, Cat., IV, 381, 1862, no locality given.

Fierasfer dubius, PUTNAM, Proc. Bost. Soc. Nat. Hist. 1874, 344, Pearl Islands (Coll. Prof. Frank H. Bradley); JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 629; JORDAN & GILBERT, Synopsis, 791, 1883.

2868. FIERASFER ARENICOLA, Jordan & Gilbert.

Head 6½ in length; depth 10½; eye 5 in head; snout 5. Body with nape slightly elevated, thence tapering regularly to the tail. Snout blunt, rounded, protruding; mouth subinferior, nearly horizontal, large, the lower jaw included; gape wide, the maxillary ⅓ length of head, extending beyond vertical from orbit; teeth in upper jaw very small, acute, in a narrow band, none of them enlarged; those in lower jaw and on vomer blunt, conic, in a wide band; those in outer series acute; a few on each side of mandible and 2 or 3 anteriorly on vomer, enlarged, canine-like. Gill openings very wide, the branchiostegal membranes little united, leaving nearly all of isthmus uncovered; the membranes

* *Fierasfer affinis*, Günther, is thus described:

"The length of the head is ⅕ of the total; its greatest width is rather less than ⅓ of its length. Gill openings rather wide, the united gill membranes leaving the greater portion of the isthmus uncovered. Teeth cardiform; a pair in front of the upper jaw, a series on the side of the lower, and several others on the vomer larger than the rest. Dorsal fin low but very distinct. The length of the pectoral nearly ⅔ that of the head. (This species is) similar to *F. acus*, but with a very different dentition." (Günther.) Described from a specimen 8 inches long, from unknown locality. This description, so far as it goes, agrees with *Fierasfer dubius*, but the specimen may not be American.

united as far back only as vertical from end of maxillary; opercle adherent above the upper angle, which is produced in a point extending above the base of pectorals; below the angle the opercular margin runs very obliquely forward. Eye large, greater than interorbital width. Origin of dorsal fin distant from nape by the length of the head, the fin a very inconspicuous fold anteriorly, becoming higher posteriorly, where the rays are evident; anal well developed along entire length, beginning immediately behind vent and running to tail, its rays visible; caudal exceedingly short; pectorals very well developed, more than ½ length of head; vent just in front of base of pectorals. Head and body perfectly translucent; a faint silvery luster on middle of sides anteriorly; a few inconspicuous small light yellowish spots along middle of sides (disappearing in alcohol); tip of tail dusky; upper margin of orbit black. Pacific coast of Mexico. A single specimen, 3¼ inches long, was found buried in the sand at low tide on the beach at Mazatlan. This specimen may be identical with *Fierasfer dubius*, but it is more robust than Putnam's types, with longer head and without enlarged teeth in upper jaw. It may be regarded as distinct, pending investigation. (*arena*, sand; *colo*, I inhabit.)

Fierasfer arenicola, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 363, Mazatlan. (Type, No. 29244. Coll. C. H. Gilbert.)

2869. FIERASFER BERMUDENSIS (Jones).

Head 8½ in length; eye 4, longer than snout; mouth large, the maxillary reaching beyond orbit; pectoral 2½ in head. Teeth small, acute, uniserial, 3 in a line on the vomer; palatine teeth small.* Color pale brownish, a bluish streak crossing the nape between the opercles, 4 pale points on the back. Vertebræ 100. Length 140 mm. West Indies. This description (by Poey) from a specimen taken in the stomach of a holothurian at Havana. Others are recorded from Key West and St. Thomas, the latter from an oyster; not seen by us; doubtfully distinct from *Fierasfer dubius* or *affinis*, but the vomerine teeth said to be smaller.

?*Carapus affinis*, POEY, Synopsis, 402, 1867; not *Fierasfer affinis*, GÜNTHER.

Lefroyia bermudensis, JONES, Zoologist, IX, 1874, 3838, Bermuda.† (Coll. General Lefroy.)

* In another specimen, according to Poey, the teeth are villiform, with an enlarged series outside, the lower teeth largest, the ninth, tenth, and eleventh largest; teeth on vomer small, acute, in a row; palatine teeth bluntish.

† *Fierasfer bermudensis* (Jones) was thus originally described:

"Total length rather more than 4½ inches. Greatest depth at the vertical of the pectorals 3⅛ lines. The length of the head is slightly more than ⅓ of the total length. The greatest width of the head rather less than ⅓ of its length. Body naked, attenuate, compressed. Facial outline rugose. Eye moderate; horizontal diameter of eyecup 1¾ lines; vertical diameter 1½ lines. Gape of the mouth ovoid. Lower jaw shorter and received within the upper. Cardiform teeth of irregular size in both jaws, vomer, and palatines, those of the latter largest. Branchiostegals 7, inflated, united below. Vent thoracic. Pectorals originating at the upper angle of the operculum, 3 lines in extent, and composed of very delicate soft rays. Dorsal indistinct, commencing in a groove about the vertical of the twentieth anal ray, continuous to caudal extreme, where, in conjunction with the anal, it forms a small filamentous tip. Anal prominent, commencing immediately behind the vent in advance of the vertical of the upper angle of the operculum, and extending to the caudal extreme. About its center it is equal in depth to that of the body at same position. Owing to the delicate texture of the fins it is impossible to ascertain certainly the number of rays, but those of the anal exceed 140. Color, when dried out of spirits, golden yellow; the body transparent, showing the vertebræ within; a condition, according to Lefroy, equally observable in life.

"I propose to publish it as *Lefroyia bermudensis*, in compliment to the gallant officer to whom I am indebted for the specimen." (J. Matthew Jones.)

This species is probably identical with the one called *affinis* by Poey.

Family CCXI. BROTULIDÆ.

(The Brotuloid Fishes.)

Body elongated, compressed, regularly tapering behind, the tail generally subtruncate at base of caudal fin, not isocercal; vent submedian; scales cycloid and minute, embedded in the lax skin, which more or less envelops the fins, sometimes wanting; gill openings very large, the membranes mostly free from the isthmus; vertical fins united or contiguous at base of caudal; dorsal fin commencing not far from nape; caudal narrow or pointed; ventral fins small, few-rayed, attached to the humeral arch and more or less in advance of pectoral. Pyloric cæca few (1 or 2), rarely obsolete or in increased number (12); maxillaries generally enlarged behind and produced toward their upper angle. (Gill.) Pseudobranchiæ small or wanting, hypercoracoid with the usual foramen, as in Blennioid fishes. These fishes are closely related to the *Zoarcidæ*. In spite of various external resemblances to the *Gadidæ*, their affinities are rather with the Blennioid forms than with the latter. Genera about 45, species about 100; largely of the depths of the sea; 2 species degenerated into blind cave fishes. We have not had material for any elaborate study of these fishes and follow closely the arrangemment given by Goode & Bean. (*Brotuloidæ*, Gill, Proc. Ac. Nat. Sci. Phila. 1863, 252, and 1884, 175.)

Brotulinæ:
- *a*. Snout and lower jaw each with well-developed barbels; vertical fins united; teeth on vomer and palatines.
  - *b*. Ventrals each reduced to a bifid filament. Brotula, 953.
- *aa*. Snout and lower jaw without barbels.
  - Lucifuginæ:
    - *c*. Species blind, dwelling in fresh-water streams in caves; barbels replaced by cilia.
      - *d*. Palatines with strong teeth; teeth in lower jaw strong. Stygicola, 954.
      - *dd*. Palatines toothless; teeth in jaws villiform. Lucifuga, 955.
    - *cc*. Species marine, the eyes usually well developed.
      - Brosmophycinæ:
        - *e*. Caudal fin differentiated, on a distinct caudal peduncle.
          - *f*. Snout and lower jaw with small cilia; head naked, or nearly so. Brosmophycis, 956.
          - *ff*. Snout and lower jaw without cilia; head more or less scaly. Ogilbia, 957.
        - *ee*. Caudal fin not differentiated, without distinct peduncle.
          - Bythitinæ:
            - *g*. Ventrals inserted on the isthmus, not far from the humeral symphysis.
              - *h*. Pectorals normal, simple; eyes present.
                - *i*. Lateral line present posteriorly, but broken in the middle; palatines with teeth; ventrals a pair of filaments each of 2 closely united rays. Bythites, 958.
                - *ii*. Lateral line obsolete posteriorly.
                  - *j*. Ventrals each of a single ray.
                    - *k*. Lateral line distinct on front of body.
                      - *l*. Preopercle without spines; head scaly (except snout); opercle with a single spine; vent median. Catætyx, 959.

ll. Preopercle with 3 or 4 spines, opercle with a single one; head partially naked.
DICROMITA, 960.

kk. Lateral line obsolescent, almost, or quite invisible; opercle with a feeble spine; head smooth; eyes small.

m. Ventral consisting of a single ray.
BASSOZETUS, 961.

mm. Ventral bifid. MŒBIA, 962.

jj. Ventrals each of a pair of rays.

n. Caudal fin exserted, but confluent with anal and dorsal.

o. Head scaly.

p. Preopercle with small spines at its angle, opercle with 1 spine.
NEOBYTHITES, 963.

pp. Preopercle unarmed.

q. Opercle with 2 spines; ventrals close together.
BENTHOCOMETES, 964.

qq. Opercle with 1 strong spine; ventrals far apart.
BASSOGIGAS, 965.

nn. Caudal not confluent with vertical fins, but without distinct peduncle; teeth on jaws, vomer, and palatines in villiform bands; preopercle unarmed; head scaly.

r. Opercle with a flat spine; snout much produced and dilated; lateral line very indistinct (or absent?).
BARATHRODEMUS, 966.

rr. Opercle a triangular flap, unarmed; lower pectoral rays prolonged, the lowest filamentous. NEMATONUS, 967.

iii. Lateral line represented by 3 rows of pores—dorsal, lateral, and ventral; head with spines.

s. Ventrals of 2 distinct rays; opercular spine moderate, straight. POROGADUS, 968.

ss. Ventrals each of 2 united rays, opercular spines strong, curved. PENOPUS, 969.

hh. Pectorals with the lower rays differentiated.

t. Preopercle armed with 3 spines; opercle armed with 1 spine; lateral line obsolete posteriorly; ventrals bifid.
DICROLENE, 970.

tt. Preopercle unarmed; a single spine on opercle; lateral line absent (?); ventrals each a pair of filaments, closely united throughout. MIXONUS, 971.

APHYONINÆ:

gg. Ventrals inserted on humeral symphysis; lateral line obsolete (in almost every case); ventrals each of a single filament; body naked; notochord persistent.

u. Eye visible through the skin; a few fang-like teeth on vomer and mandible. BARATHRONUS, 972.

uu. Eye not visible; no teeth on maxillary or palatines; teeth on vomer rudimentary, those on mandible small.
APHYONUS, 973.

953. BROTULA, Cuvier.

(Brótulas.)

Brotula, Cuvier, Règne, Anim., Ed. 2, II, 296, 1829 (*barbata*).

Body elongate, compressed, covered with minute smooth scales; eye moderate; mouth medium, with villiform teeth on jaws, vomer, and palatines; lower jaw included; each jaw with 3 barbels on each side. Dorsal fin long and low, the dorsal and anal joined to the caudal. Ventral fins each reduced to a single filament of 1 ray. Eight branchiostegals. Air bladder large, with 2 horns posteriorly. One pyloric cæca. Vertebræ 16 + 39 = 55. Tropical seas, in water of moderate depth. (*Brótula*, Spanish name of *Brotula barbata*.)

2870. BROTULA BARBATA (Bloch & Schneider).

(Brótula.)

Head $4\frac{1}{2}$; depth about 5. D. 123; A. 93; V. 1. Upper jaw the longer. Ventral fin $\frac{1}{2}$ as long as head. Dorsal commencing behind vertical from root of pectoral; vertical fins covered with thick skin. Color nearly uniform brown. Length 12 to 18 inches. West Indies; rare; in water of moderate depth. One specimen obtained by us in the market of Havana. (*barbatus*, bearded.)

Brótula, Parra, Dif. Piezas Hist. Nat., 70, lam. 31, fig. 2, 1780, Havana.
Enchelyopus barbatus, Bloch & Schneider, Syst. Ichth., 52, 1801; after Parra.
Brotula barbata, Cuvier, Règne Anim., Ed. 2, II, 296, 1829; Poey, Memorias, II, 102, lam. 9, fig. 2, 1860; Günther, Cat., IV, 371, 1862.

954. STYGICOLA, Gill.

Stygicola, Gill, Proc. Ac. Nat. Sci. Phila. 1863, 252 (*dentatus*).

This genus differs from *Lucifuga* in the presence of palatine teeth. The teeth in the jaws are larger. As in *Lucifuga*, the single known species inhabits cave streams in Cuba. (στύξ, *Styx*, the river of the lower regions; *colo*, I inhabit.)

2871. STYGICOLA DENTATUS (Poey).

Head $2\frac{2}{3}$; depth $3\frac{3}{5}$. D. 90; A. 70; P. 17; V. 1. Vertebræ 11 + 37 = 48. Eyes usually wanting, occasionally represented by a rudiment; head elevated at the nape, the general form less slender than in *Lucifuga*, the belly more prominent; no scales on the nape; strong teeth, well separated, on the palatines as well as the vomer; teeth in the jaws larger than in *Lucifuga*; posterior with a large apophysis. Color translucent violet, with darker areas on nape and throat. Caves of the province of San Antonio, in southern Cuba. Largest specimen 5 inches long. (Poey.) (*dentatus*, toothed.)

Lucifuga dentatus, Poey, Memorias, II, 102, 1860, Cave of Cajío (Coll. Noda), Cave of Castle La Industria (Coll. Dubrocá), Cave of Ashton (Coll. Fabre); Günther, Cat., IV, 373, 1862.
Stygicola dentata, Gill, Proc. Ac. Nat. Sci. Phila. 1863, 252.

955. LUCIFUGA, Poey.

(Cuban Blindfish.)

Lucifuga, Poey, Memorias, II, 95, 1860 (*subterraneus*).

Body moderately elongate, translucent pinkish, covered with minute scales. Eye rudimentary, covered by the skin; bands of villiform teeth in the jaws and vomer, none on the palatines; nostrils 2 on each side; no barbels; head with small tactile cirri; no spines on head; gills 4; no pseudobranchiæ; gill opening large, extending forward nearly to the symphysis, the gill membranes not united; branchiostegals 7 or 8; vertical fins low, united around the tail; ventrals each reduced to a short thin filament; male with an anal papilla, no pyloric cæca; air bladder large, rounded behind, joined to the base of the skull. Cave streams of Cuba; the eyes having undergone a degeneration similar to that seen in *Amblyopsis*. These fishes have no relation to the blind cave fishes of North America, but are derived from marine types, their ancestors being evidently allies of *Ogilbia* and *Brotula*. It is known that blindfishes are found also in caves of the islands of Jamaica, but no specimens have been seen by naturalists. (*lux*, light; *fugo*, I flee.)

2872. LUCIFUGA SUBTERRANEUS, Poey.

(Pez Ciego.)

Head $2\frac{2}{5}$; depth $3\frac{2}{5}$. Branchiostegals 7. D. 70; A. 70; P. 51; V. 1; C. 9. Vertebræ 11 + 36 = 47. Body elongate, compressed, tapering, pointed; head low at nape, much depressed anteriorly, broad, covered with soft, white, wrinkled skin, with microscopic cirri, having firm and conical tubes; no barbels on lips or chin; skin of head with many pores; scales not ciliate, present on body and top of head and on opercles; lateral line median, marked by a series of microscopic cirri like those on head, these wanting posteriorly; eyes wanting; nasal openings double; mouth large; lower jaw shorter; lips fleshy; maxillary broad at tip, $\frac{2}{5}$ length of head; teeth in jaws very short and sharp, in a band; vomerine teeth larger; no palatine teeth; pharyngeal teeth slender; tongue smooth; gill openings large; males with an anal papilla; fin rays simple, flexible, jointed but not branched; dorsal beginning at a point about $\frac{1}{2}$ nearer tip of snout than tip of caudal; anal smaller, beginning farther back, the 2 fins fully joined to the pointed caudal; pectoral fin short, falcate, nearly $\frac{1}{4}$ head; ventral in front of pectoral, a slender ray not $\frac{1}{2}$ length of maxillary. Color transparent rosy, head reddish, becoming darker in alcohol. No pyloric cæca; intestines short; air bladder large. Described from 12 specimens, the longest about $4\frac{1}{2}$ inches; found in caves of the jurisdiction of San Antonio, in the southern part of Cuba. (Poey.) (*sub*, under; *terra*, earth.)

Lucifuga subterraneus, Poey. Memorias, II, 96, 1860, San Antonio, Cuba (Coll. D. Tranquilino); Sandalio de Noda (Coll. D. Juan Antonio Fabre); first coll. from Cajío Cave, 1831 (Noda); second, Cave at La Industria (Coll. Dubrocá); third, Ashton Cave, San Andreas (Coll. Fabre); fourth, Cave of the Dragon (Coll. Fabre); fifth, Cave at the Castle of Concord (Coll. Layunta).

956. BROSMOPHYCIS, Gill.

Brosmophycis, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 168 (*marginatus*).
Halias, AYRES, Proc. Cal. Ac. Sci. 1861, 52 (*marginatus*); preoccupied.

Body elongate, moderately compressed; head unarmed; snout not long; teeth sharp, curved, in bands on jaws, vomer, and palatines; small cilia above snout and on anterior part of lower jaw. Body covered with thin cycloid scales; scales on head rudimentary or wanting. Caudal fin differentiated, entirely separated from the dorsal and anal; caudal peduncle slender. California. This genus is very close to *Ogilbia*, differing in the ciliated lips. Its species reaches a larger size. (*Brosmius; Phycis.*)

2873. BROSMOPHYCIS MARGINATUS (Ayres).

Head 4½ in body; depth 6⅞. D. 92; A. 70; eye 7½ in head; snout 4½; maxillary 2; pectoral 1⅝; caudal 3; body elongate, moderately compressed; snout blunt; profile of head straight from snout; snout scarcely overhanging mouth; jaws subequal, the teeth conical, sharp, and slightly curved back, in bands on jaws, vomer, and palatines; maxillary reaching ½ the eye's diameter beyond eye; snout and lower jaw thickly covered with small cilia; head naked with the exception of small scales above; 2 large pores at tip of chin, a few large ones around preopercle and preorbital, 1 around gill opening, behind which is a pocket in the skin; about 3 short gill rakers developed below the angle of first arch, with many rough plates, not differentiated from those on the other arches. Dorsal and anal long and low, the rays embedded in the skin; tips of last rays each beyond the base of the caudal about ½ the length of caudal rays; distance of front of dorsal from snout 3⅓ in length of body; origin of anal a little nearer base of caudal than tip of snout; pectoral reaching about half way from its base to front of anal; ventrals developed as long filaments; caudal slender and rounded behind. Color bright reddish brown; fins edged with bright rose-red. Coast of California, in water of moderate depth; rare. Here described from a specimen, 12 inches in length, collected off San Francisco by Mr. W. G. W. Harford. (*marginatus*, edged.)

Brosmius marginatus, AYRES, Proc. Cal. Ac. Nat. Sci., I, 1854, 13, San Francisco (Coll. W. O. Ayres); GIRARD, Pac. R. R. Surv., X, Fishes, 141, 1858.
Brosmophycis marginatus, GILL, Proc. Ac. Nat. Sci. Phila. 1861, 168; GILL, Proc. Ac. Nat. Sci. Phila. 1862, 280.
Halias marginatus, AYRES, Proc. Cal. Ac. Nat. Sci., pt. 2, 1861, 52.
Dinematichthys marginatus, GÜNTHER, Cat. Fishes Brit. Mus., IV, 375; JORDAN & GILBERT, Synopsis, 796, 1883.

957. OGILBIA, Jordan & Evermann.

Ogilbia, JORDAN & EVERMANN, in EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897 (February 9, 1898), 132 (*cayorum*).

Body moderately elongate, covered with minute, smooth, embedded scales; sides of head with similar scales; lateral line inconspicuous; opercle with a very small spine, preopercle unarmed; no strong hook on maxillary; no barbels nor cilia; teeth in jaws in bands, similar teeth on

vomer and palatines; caudal free from the dorsal and anal; lower lip without cirri; dorsal and anal rays covered by the skin, ventrals each reduced to a filament of 2 rays; anal papilla of the male without horny claspers. Small fishes of the tropical shores of America, living in rock pools and shallows among algæ. This genus is closely allied to the East Indian genus *Dinematichthys*, differing in the absence of anal papilla and claspers and in the shorter vertical fins. (Named for J. Douglas Ogilby, the accomplished naturalist of the museum of Sydney, in recognition of his excellent work on the fishes of Australia.)

a. Snout very short, about 7 in head; eye small, 10 or 11 in head; scales small, obscure, snout very short, 6½ in head. VENTRALIS, 2874.

aa. Snout longer, about 4 in head; eye about 8½ in head; scales larger, distinct. CAYORUM, 2875.

2874. OGILBIA VENTRALIS (Gill).

Head 4½ in body; depth 5½. D. 64; A. 50; scales about 100; eye 10 or 11 in head; maxillary 2; pectoral 1⅔; caudal 1⅝. Body elongate, moderately compressed; snout blunt, the profile behind snout nearly straight to occiput; mouth large, the maxillary extending 2 or 3 times the eye's diameter behind eye; teeth small, in bands on jaws, vomer, and palatines; eye very small, nearer snout than posterior end of maxillary; no cilia on snout and chin; body apparently naked to the unaided eye; but body and top of the head covered with small scales, which can be seen by the aid of a lens. Origin of dorsal distant from tip of snout by a space contained 3¼ times in body; front of anal about midway between tip of snout and base of caudal; tips of last dorsal and anal rays reaching about to the middle of caudal rays, but not connected; pectorals scarcely reaching midway between their base and the front of anal; ventrals filamentous; caudal slender and rounded behind. Color in spirits, light brown above, lighter below; fins all colorless; without distinct marking anywhere. Gulf of California; not rare in rock pools; several specimens, 2 to 4 inches in length taken by us at Mazatlan. Here described from a specimen, 2 inches in length, from La Paz Harbor, Lower California. (*ventralis*, pertaining to the belly.)

Brosmophycis ventralis, Gill, Proc. Ac. Sci. Phila. 1863, 253, Cape San Lucas. (Coll. Xantus.)

Dinematichthys ventralis, Jordan, Proc. Cal. Ac. Sci. 1895, 502, pl. 54.

2875. OGILBIA CAYORUM, Evermann & Kendall.

Head 4; depth 4½; eye 8½; snout 4. D. about 68; A. about 50; scales about 14–87–13; maxillary 1⅓; pectoral 1⅓; ventral 1½; caudal 2⅓. Body moderately elongate, compressed; head moderate, snout blunt; mouth large, jaws subequal, maxillary extending beyond vertical of eye a distance nearly equal to length of snout; eye very small, high up, situated in anterior third of head; nostril small, close to eye; teeth small, in bands on jaws, vomer, and palatines; back elevated, strongly arched from snout to origin of dorsal fin, thence descending in a nearly straight line to base of caudal; ventral outline comparatively straight, slightly concave at front of anal. Dorsal and anal long and low, distinct from caudal, the

posterior rays longest, about 3½ in head, base of each scaled; distance from tip of snout to origin of dorsal about 3 in length of body; origin of anal under about twenty-second dorsal ray, equidistant between tip of snout and base of caudal; scales very small, embedded, but showing distinctly under a lens; cheek and opercles partially covered with minute, embedded scales; top of head naked; opercle with a large, flat, flexible spine on level with eye. No barbels, cilia, nor tubercles; 2 large mucous pores at symphysis of lower jaw and 2 on preorbitals near anterior edge on each side; a row of 5 or 6 pores on lower jaw and edge of preopercle. Color uniform pale olivaceous or light brown, finely punctate with minute brown specks. Key West. Only the type known, an example, 2⅛ inches long, seined on a shoal covered with algæ at Key West. (Cayo Hueso, or Boné Key, the original Spanish name for the Island of Key West, whence the name *cayorum*, of the keys.)

Ogilbia cayorum, Evermann & Kendall, Bull. U. S. Fish Comm. 1897 (Feb. 9, 1898), 132, pl. 9, fig. 14, Key West, Florida. (Type, No. 48792. Coll. Evermann & Kendall.)

958. BYTHITES, Reinhardt.

Bythites, Reinhardt, Dansk. Vidensk. Selsk. Afhandl., vii, 1838, 178 (*fuscus*).

Body elongate, covered with minute scales. Head large, thick; mouth large; jaws equal; no barbel; bands of teeth in the jaws and on vomer and palatines. Branchiostegals 8; gill membranes united, free from the isthmus; eyes moderate. Lateral line interrupted. Vertical fins united; ventral fins reduced to simple filaments, each composed of 2 rays closely united. Air bladder large; 2 pyloric cæca. A thick, conical, anal papilla (in the male). Greenland. (*βυθίτης*, an animal of the depths, from *βύθιος*, the deep.)

2876. BYTHITES FUSCUS, Reinhardt.

Head about 4; depth 4½. Body somewhat compressed, lipariform; snout obtuse, naked, with minute cirri. Mandible long, curved, extending far behind vertical from posterior margin of orbit; eye small; scales moderate on body; lateral line complete, but interrupted over vent, the two parts slightly overlapping the same vertical; vertical fins confluent, enveloped in thick skin; pectorals broad, lanceolate, with broad base; ventrals filiform, reaching behind origin of pectoral, as long as pectoral and ⅔ as long as head; a conspicuous anal papilla in the male. The only known specimen, now in the museum at Copenhagen, was obtained in Greenland half a century ago. (Goode & Bean.) (*fuscus*, dusky.)

Bythites fuscus, Reinhardt, Dansk. Vidensk. Selsk. Afh., vii, 1838, 178, Greenland; Günther, Cat., iv, 375, 1863; Jordan & Gilbert, Synopsis, 795, 1883; Goode & Bean, Oceanic Ichthyology, 316, 1896.

959. CATÆTYX, Günther.

Catætyx, Günther, Challenger Report, xxii, 104, 1887 (*messieri*).

Body compressed, elongate, covered with very small and thin scales; lateral line indistinct, interrupted. Head oblong, with somewhat pointed snout, covered with very small scales, only the anterior part of the snout

naked; bones of the head rather firm, but with the muciferous system well developed, the canals having wide openings along the infraorbital, and on the lower limb of the preoperculum; eye rather small; nostrils far apart, the posterior in front of the eye and the anterior at the extremity of the snout; operculum with a spine behind; no other armature on the head; snout not swollen, but the upper jaw slightly overlapping the lower; barbels none; mouth wide; bands of villiform teeth in the jaws, on the vomer, and the palatine bones; a series of larger teeth along the sides of the lower jaw; tail not much attenuated; vertical fins confluent; ventrals close together, reduced to a pair of fine, simple filaments, and inserted somewhat behind the isthmus, below the middle of the operculum. Gills 4, with short, broad gill rakers and well developed laminæ; pseudobranchiæ none; branchiostegals 8; pyloric appendages. Deep seas. Two species known. (*κατα*ί, at the bottom; *τύξις*, *τυγχάνω*, find.)

2877. CATÆTYX RUBRIROSTRIS, Gilbert.

Depth of body below origin of dorsal equals $\frac{1}{3}$ distance from end of snout to vent, 7 in length; head 4; distance from snout to origin of dorsal $3\frac{1}{5}$; from snout to vent $2\frac{2}{3}$; maxillary extending beyond eye, $2\frac{2}{3}$ in head; eye equaling snout, $5\frac{1}{2}$; interorbital width 7; width of snout $3\frac{1}{3}$. Teeth in upper jaw in a narrow band, minute, compressed, narrowly triangular, none of them enlarged; in the lower jaw a still narrower band of similar teeth, the posterior row slightly enlarged and increasing a little in size on sides of jaw, where it is accompanied by a single series only of the smaller teeth; this lateral series is continued backward far beyond premaxillary band; teeth on vomer and palatines similar to those in sides of lower jaw, the former in a V-shaped patch, the latter in a long and very narrow band. Anterior nostril in a short tube at tip of snout, the posterior large, without tube, immediately in front of eye; system of mucous pores well developed but not conspicuous, the pores collapsing on account of the thinness of the skin covering head; large mucous tube below eye, extending around front of snout and opening by slit-like pores along edge of snout and lower margin of infraorbital flap, opening posteriorly by a vertical slit $\frac{1}{2}$ as long as pupil, immediately above end of maxillary; another series of pores along mandible and at edge of expanded limb of preopercle; no other evident pores. Angle of preopercle much expanded, its width equaling diameter of pupil; a sharp, strong spine arising from anterior portion of opercle, the structure of the gill flap apparently like that of *Bassogigas stelliferoides;* a short, sharp spine directed backward immediately behind posterior nostril; no other spines on head; gill openings wide, continuing forward to below posterior margin of orbit, the membranes wholly free from the isthmus; gill rakers short but not very broad, about $\frac{1}{2}$ length of pupil, only 2 or 3 developed immediately in front of angle of arch. Dorsal beginning over or slightly behind middle of pectorals, the distance from its origin to occiput equaling or somewhat exceeding distance of latter from tip of snout; dorsal and anal fully united to caudal. The caudal has a base of appreciable width, bearing about 12 close-set rays, which extend much beyond tips of last dorsal and

anal rays; origin of anal nearer snout than base of caudal; ventrals slender, each consisting of a single ray, inserted very near together, under anterior portion of opercle, their length about equaling that of maxillary; pectorals with about 23 rays, evenly rounded behind, their length $1\frac{2}{3}$ in head. Scales very small, cycloid, regularly imbricated, in about 135 transverse series; nape and belly scaled, as is also the head, excepting snout, mandible, suborbital, and sometimes interorbital areas. Color dusky olive, the ventrals white, the other fins black, at least on distal portions; opercles, gill membranes, sides and top of snout, and posterior portion of abdomen blue black; snout flushed with dark ruby red in life; lining membrane of mouth and gill cavity, and peritoneum jet-black. Closely related to *C. messieri*, differing in the shorter, broader snout, the wider preopercle, the more anterior origin of anal, and apparently in the gill rakers and pores on head. Off coast of California. Four specimens known, the longest $4\frac{1}{2}$ inches in length. (Gilbert.) (*ruber*, red; *rostrum*, snout.)

Catætyx rubrirostris, GILBERT, Proc. U. S. Nat. Mus. 1890, 111, off coast of California, at Albatross Stations 2909, 2925, and 2936, in 205 to 359 fathoms. (Type, No. 44379.)

960. DICROMITA, Goode & Bean.

Dicromita, GOODE & BEAN, Oceanic Ichthyology, 319, 1896 (*agassizii*).

Brotulids resembling in form and general appearance *Catætyx* and *Diplacanthopoma*, having the lateral line obsolete, or interrupted posteriorly; ventrals a pair of simple, fine filaments, and with teeth upon the palatines. It has, however, 3 or 4 small spines upon the preoperculum, as well as a sharp spine upon the upper angle of the operculum; and the lateral line, though indistinct, is traceable for $\frac{1}{2}$ or $\frac{3}{4}$ the length of the body, which, like the upper part of the head, is covered with small, deciduous scales, the opercular region being apparently scaleless, and the bones of the suborbital region almost uncovered, with conspicuous sinuses, which show through the transparent texture of the surface. Head oblong; snout somewhat produced, depressed, and turgid, resembling, though in a less degree, that of *Barathrodemus*. Eye moderate, conspicuous. Mouth wide; teeth villiform, in bands on the jaws and palatines, and very minute upon the vomer, which has a roughened, knob-like enlargement at its angle. Vent premedian. Ventral fins confluent; ventrals rooted very close together, each reduced to a fine, flexible, simple filament, planted somewhat behind the isthmus and below the middle of the operculum. Gills 4, with well-developed laminæ and rather long, slender gill rakers. Branchiostegals 8. Pseudobranchiæ apparently absent. (δίκρος, forked; μίτος, thread.)

2878. DICROMITA AGASSIZII, Goode & Bean.

Body elongate, much compressed, its height about $\frac{1}{5}$ of its total length, its width about $\frac{1}{2}$ its greatest height; head slightly greater than height of body, about twice its own width; mouth very large, the maxillary curved and much dilated at its extremity, reaching far behind the vertical from the posterior margin of orbit; jaws nearly equal, the snout considerably

produced and dilated, its length equal to diameter of eye and ¼ length of head. Teeth very fine, villiform, in bands on jaws and palatines, and also present on vomer, though very small, especially upon the rounded, globular process of the angle. Lateral line very indistinct, interrupted, but extending behind the vent at least ⅓ of the way to tip of tail. Dorsal origin nearly in vertical from the axil of the pectoral; ventrals very slender, villiform, closely approximate at their roots, and less than ½ as long as the head. Color brownish. A specimen was obtained by the *Blake* off Granada, Station XCIII, at a depth of 291 fathoms. The collateral type was obtained by the *Albatross* at Station 2374, in lat. 29° 11′ 30″ N., long. 85° 29′ W., at a depth of 26 fathoms. (Goode & Bean.) (Named for Prof. Alexander Agassiz.)

Dicromita agassizii, GOODE & BEAN, Oceanic Ichthyology, 319, fig. 285, 1896, off Granada, in 291 fathoms (Coll. the *Blake*); Lat. 29° 11′ 30″ N., Long. 85° 29′ W., in 26 fathoms. (Coll. *Albatross*.) (Type in M. C. Z.)

961. BASSOZETUS, Gill.

Bathynectes, GÜNTHER, Ann. and Mag. Nat. Hist., II, 1878, 20 (*compressus*); name preoccupied in Crustacea.

Bassozetus, GILL, Proc. U. S. Nat. Mus., VI, 1883, 59 (*normalis*).

Bathyonus, GOODE & BEAN, Proc. U. S. Nat. Mus., VIII, 1886, 603 (*catena*).

Body compressed, with long tapering tail, covered with deciduous thin scales of moderate size. Bones of the head very soft and cavernous, the upper opercular spine very feeble, ridge-like; no other armature of the head. Head scaly, except the snout, which is obtusely rounded off, with the jaws equal or nearly equal in front. Mouth very wide; bands of villiform teeth in the jaws, on the vomer, and palatine bones. Barbels none. Eye small; anterior nostril about midway between the posterior and the extremity of the snout. Vertical fins confluent; ventrals close together, reduced to a pair of simple filaments, and inserted below the rounded angle of the preoperculum. Gills 4, with short gill laminæ, but with long stiff gill rakers on the first branchial arch. Pseudobranchiæ none. Branchiostegals 8. Pyloric appendages none. (*βάσσων*, for *βαθύς*, deep; *ζητέω*, seek.)

a. Body moderately elongate, the depth 9 to 10 in length; dorsal rays 116; anal 92 to 96.
b. Head 6 in length. NORMALIS, 2879.
bb. Head about 7½ in length. COMPRESSUS, 2880.
aa. Body more elongate, the depth 12½ in length; head 8⅔; head with conspicuous, chain-like rows of pores. CATENA, 2881.
aaa. Body excessively attenuate, the depth more than 16 times in length: dorsal rays 138; anal 115; distance from snout to vent nearly 4 times in body. TÆNIA, 2882.

2879. BASSOZETUS NORMALIS, Gill.

Head 6 in total length; eye 4 in snout. D. 116; A. 96. Body much compressed, its width in the region of vent not more than ⅛ of its height, which at the same point is about ⅛ of the total length; greatest height of body, over the origin of the pectorals, about ⅔ the distance from base of pectorals to vent, the vent being about twice as distant from base of

caudal rays as from snout. Head moderately compressed, flat above; snout obtuse rounded, turgid; lower jaw considerably included. Bones of head not completely ossified, very cavernous in the alcoholic specimen, the head showing many deep sinuosities and depressions. Eye very small, situated about midway between the tip of the snout and the vertical from the posterior end of maxillary. Teeth all small and short, densely set, forming narrow, villiform bands; vomerine band open V-shaped. Dorsal fin beginning far in advance of origin of pectoral and above upper angle of gill opening, rays longest in region of vent; anal beginning immediately behind vent, its rays not quite so long as those of dorsal; pectoral with broad base, short, not exceeding much more than halfway to vertical from vent, its length considerably less than that of postorbital portion of head. Ventral rays very slender, villiform, reaching almost to vent, far beyond pectoral, their length almost equal to that of head. Scales moderate, very deciduous, extending upon cheeks and on top of head almost to tip of snout; no evidence of a lateral line. Color light, the head and abdomen blackish; inside of mouth purplish brown. The *Blake* secured specimens from Station CCIV, in lat. 24° 33′ N., at a depth of 1,920 fathoms, and from Station LXXXIV, off Dominica, in 1,131 fathoms. The *Albatross* also obtained examples (No. 49416, U. S. Nat. Mus.) from Station 2380, in Lat. 28° 02′ 30″ N., Long. 87° 43′ 45″ W., at a depth of 1,430 fathoms; (No. 33306, U. S. Nat. Mus.) from Station 2042, in lat. 39° 33′ N., Long. 68° 26′ 45″ W., at a depth of 1,555 fathoms. (Goode & Bean.) West Indies, Gulf of Mexico, and to lat. 40° N., in region of the Gulf Stream. (*normalis*, normal.)

Bassozetus normalis, Gill, Proc. U. S. Nat. Mus. 1883, 259, Lat. 39° 33′ N., Long. 68° 26′ 45″ W., in 1,555 fathoms (Type, No. 33306. Coll. *Albatross*); Goode & Bean, Oceanic Ichthyology, 322, fig. 287, 1896.

2880. BASSOZETUS COMPRESSUS (Günther).

D. 116; A. 92; P. 23; V. 1. The greatest depth of the body is above the end of the gill cover and about ½ length of trunk; vent twice as distant from extremity of tail as from snout, consequently the tail is more moderately attenuated. Head compressed like the body, and about ⅔ length of trunk; superficial bones form large muciferous cavities which, when full, must give to the head a much more evenly rounded appearance than in the preserved state, when the supporting bony ridges project more or less from under the skin. Snout slightly swollen, but the jaws nearly even in front, the wide mouth slightly ascending forward; maxillary with the form usual in these Gadoid fishes, dilated behind, and extending far behind the eye. Eye very small, ½ length of snout, and $\frac{1}{11}$ that of head, placed high up on the side, and not possessing an orbital fold of integument; interorbital space rather convex and equal in width to 3 diameters of eye. Teeth all very small, short, densely set, and forming villiform bands, the broadest on maxillary bone and quite uncovered on the sides, no labial folds being developed; palatine band broader than the mandibulary, and the vomerine band V-shaped, each arm being bent with the convexity inward. Gill opening and cavity very wide and of an intense black; gill rakers much longer than the laminæ, 15 in number on

the anterior arch, besides some rudimentary ones above. Dorsal fin commencing above upper end of gill opening, with short rays partly hidden in the skin, becoming longer in middle of fin, but remaining of moderate length; anal shorter; pectoral with a rather narrow base, quite free, and composed of feeble rays, its length only ½ that of head; ventral rays very feeble, reaching somewhat beyond the root of pectoral. In the specimens examined only very few of the thin, cycloid scales have been preserved; they are of moderate size, there being about 16 in a transverse series running from the vent to the dorsal fin; the lateral line, if it was developed, can no longer be traced. Blackish, with the fins, head, and abdomen black. Specimens of this very fine and truly bathybial fish were obtained at great depths on the southeast of New Guinea, off the Philippine Islands, and in the mid-Atlantic; the exact localities being 75 miles east-southeast of Raine Island, Station 184, depth 1,400 fathoms; two specimens, 17 and 4½ inches long. Philippine Islands, Station 205, depth 1,050 fathoms; one specimen, 5½ inches long. Mid-Atlantic, Station 107, depth 1,500 fathoms; one specimen, 5½ inches long. The young are extremely similar to the old, but have a larger eye, which is ⅛ of length of head. The specimen from Station 205 (Philippine Islands) has longer ventral filaments, extending nearly to the vent. (Günther.) (*compressus*, compressed.)

Bathynectes compressus, GÜNTHER, Ann. Mag. Nat. Hist., II, 1878, 20, Challenger Station 107, mid-Atlantic, in 1,500 fathoms. (Coll. *Challenger*.)
Bathyonus compressus, GÜNTHER, Challenger Report, XXII, 109, 1887.
Bassozetus compressus, GOODE & BEAN, Oceanic Ichthyology, 322, 1896.

2881. BASSOZETUS CATENA, Goode & Bean.

Head 8⅔; depth 1½ in head or 12½ in body; eye 5; snout 5; interorbital width 5. Body very elongate, much compressed, and tapering into a slender, whip-like tail. Head without spines, very cavernous, not much compressed, higher than body. Interorbital area somewhat convex. The muciferous channel upon the infraorbital ring shows in its course several wide subcircular sinuses, closely approximated; a similar row upon the posterior edge of the preoperculum and continued forward upon the under surface of the mandible; the vertex also has a semicircle of similar sinuses. Maxillary extending beyond vertical through posterior margin of orbit, its length equal to that of postorbital part of head; mandible ⅔ as long as head and equal in length to height of body; jaws, vomer, and palate with bands of villiform teeth, the vomerine band V-shaped. Nostrils in front of middle of eye, separated by a slight interspace, the anterior nearer to its mate than to tip of snout. Branchiostegals 8; pseudobranchiæ absent. Gill rakers long and numerous, the longest slightly exceeding diameter of eye, 15 developed below angle of first arch, besides several rudiments; dorsal origin slightly behind that of pectoral, its distance from tip of snout about 7⅓ in total, rays well developed; in the anterior ⅓ of the fin, in a space equal to length of head, were counted 20 rays, the longest of which is ⅔ as long as head; anal origin under twenty-first dorsal ray, its rays shorter than those of dorsal; pectoral extending to vertical from eighteenth ray of dorsal, ⅔ as long as head; ventrals composed each

of a simple filament, the origin slightly in advance of vertical through pectoral origin, the length ⅔ that of head, not reaching nearly to vent, the distance of which from origin of ventrals is slightly greater than length of head. Color brownish yellow; head and abdomen blackish. Gulf of Mexico, in great depths. Only the type known, 237 mm. long. (*catena*, chain, from the arrangement of the mucous cavities on the head.)

Bassozetus catena, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 603, Lat. 28° 00′ 15″ N., Long. 87° 42′ W., in 1,467 fathoms (Type, No. 37341. Coll. *Albatross*); GÜNTHER, Challenger Report, XXII, 111, 1887; GOODE & BEAN, Oceanic Ichthyology, 323, fig. 286, 1896.

2882. BASSOZETUS TÆNIA (Günther).

D. 138; A. 115; P. 30; V. 1. The greatest depth of the body is below the origin of the dorsal fin and about ⅓ of the length of the trunk, the vent being not quite thrice as distant from the extremity of the tail as from the snout; therefore, the whole of the fish, and especially the tail, is much attenuated. Head not compressed, low and long, forming ⅓ length of trunk. Structure of the bones of the head as in *B. compressus*. Snout rather swollen and broad, the upper jaw but slightly overlapping the lower; maxillary extending far behind the eye, which is very small, ⅓ length of snout, about 1⅛ that of head, and ¼ width of interorbital space. Teeth very small and short, densely set, forming narrow, villiform bands; vomerine bands open Λ-shaped. Gill cavity deep black; gill rakers long and slender, 16 in number, with some rudimentary ones in front and behind. Dorsal fin commencing above upper end of gill opening, with short rays partly hidden in the skin, the rays becoming longer on the anterior third of tail, but remaining of moderate length, the anal rays still shorter; pectoral with a broad base, quite free, and composed of rather feeble rays, its length equal to that of postorbital portion of head; ventral rays very feeble, reaching nearly to the middle of the pectoral. The scales must have been extremely thin and rather small; there were probably about 20 in a transverse series running from the vent to the dorsal fin. The lateral line can not be made out. Light colored (possibly pink in life), with the head and abdomen black. Only 1 specimen known of this eminently bathybial fish, obtained in mid-Atlantic (*Challenger* Station 104) at a depth of 2,500 fathoms. Its total length is 10 inches. (ταινία, ribbon.)

Bathyonus tænia, GÜNTHER, Challenger Report, XXII, 110, 1887, pl. 23, fig. A, mid-Atlantic, Station 104, at a depth of 2,500 fathoms; GOODE & BEAN, Oceanic Ichth., 323, 1896.

962. MŒBIA, Goode & Bean.

Mœbia, GOODE & BEAN, Oceanic Ichthyology, 331, 1896 (*gracilis*).

Brotulids resembling *Bassozetus* in general form, excepting that the tail is prolonged in a very slender filament, the dorsal and anal rays being extremely short posteriorly, but positively confluent with the caudal rays, which are much longer and much exserted; ventrals each bifid, instead of a single ray, as in *Bassozetus*. Head very cavernous, the sinuses large and conspicuous on the infraorbital ring, on the mandible, and the pre-

operculum. A single, short, feeble spine on the shoulder, but none upon the operculum or preoperculum, though certain projections seem to show above the eye, doubtless due to the shrinkage of the integument upon the underlying projections of the bone. Mouth very wide, the extremity of the maxillary much dilated; posterior nostrils very wide and separated from the eye by a small, spinous projection of bone; teeth in narrow bands, that on the vomer V-shaped, with the 2 arms straight. A few large scales in a row starting from the upper angle of the gill opening and terminating over the axle of the pectoral. Gill rakers on outer arch rather numerous, long and slender. Pseudobranchiæ represented by 2 minute globules. Deep sea; 2 species known. ("Named in honor of Prof. Karl Möbius, director of the Royal Zoological Museum in Berlin, who has added much to our knowledge of marine life by his noble work, Die Fauna der Kielerbucht, and by numerous other writings.")

2883. MŒBIA PROMELAS (Gilbert).

Head 2 in trunk; depth 3. Body 3⅛ to 3¼ in tail. Tail produced into a filament, the caudal basis extremely narrow, supporting 5 long slender rays which are firmly bound together. Mouth terminal, large, the maxillary much dilated at tip, reaching well behind the eye, 1⅘ in head; lower jaw included, the tip slightly produced. Teeth in villiform bands on mandible, premaxillary, vomer, and palatines, the band on mandible very narrow, that on vomer with the diverging arms much incurved, the anterior angle rounded. Tongue toothless, some of the basibranchials forming a sharply elevated dentigerous crest. Gill laminæ extremely narrow, the gill rakers of outer arch very long and slender, 1 (with 4 rudiments) above angle, 15 below. Infraorbital chain with 6 mucous sinuses, the mandible with 5, preopercle with 5, and a number on top of head; these are all bridged over with very delicate membrane which is easily ruptured. A row of low, strong spinous points directed posteriorly on the ridge running backward from the eye; no other spines on head, though a number of short spinous points are made evident when the skin is removed; opercular spine rather weak. A distinct membranaceous flap runs along the projecting edge of shoulder girdle, connecting pectorals with upper end of gill flap; pectorals slender, equaling postorbital part of head; ventrals each of a bifid filament, the two branches joined at the base for a very short distance, variable in length, reaching to or nearly to tips of pectorals, usually contained about 1⅓ times in head; dorsal beginning a trifle behind base of pectorals, its distance from occiput equaling distance of latter from front of eye. Scales very small, apparently covering a part of top of head; 3 series of large pores on sides; 1 from upper end of gill slit backward parallel with dorsal outline; a second along middle of sides; the third beginning halfway between base of pectorals and ventral outline, extending backward on belly and along base of anal fin, these lines all somewhat indistinct, and it can not be determined how far they extend backward. Color light brown; head (except occiput), mouth, gill cavity, and abdomen jet-black; fins dusky. This species closely resembles *Mœbia gracilis* (Günther), from New Guinea, dif-

fering in the following respects: Depth 3 in trunk (in *gracilis* 3½); eye 6 in head (in *gracilis* 5½); vomerine patch of teeth with the two arms incurved (U-shaped in *gracilis*); dorsal fin beginning behind the pectoral (over root of pectoral in *gracilis*); an additional series of large scales (lateral line) along middle of sides, and another along ventral outline; trunk 2⅛ in tail (2⅜ in *gracilis*); ventrals shorter, not reaching past tips of pectorals. Five specimens, the largest 9½ inches long, from *Albatross* Station 3010, at a depth of 1,005 fathoms, in the Gulf of California. (Gilbert.) (*πρό*, before; *μέλας*, black.)

Porogadus promelas, GILBERT, Proc. U. S. Nat. Mus. 1891, 547, Gulf of California, in 1,005 fathoms. (Coll. *Albatross.*)

963. NEOBYTHITES, Goode & Bean.

Neobythites, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 600 (*gilli*).

Brotulids having the body elongate, compressed, covered with small scales, and the head also scaled; lateral line incomplete, obsolete posteriorly. Eye moderate; snout moderate, rounded, slightly produced, the lower jaw slightly included; no barbel. Teeth villiform, in narrow bands in jaws and palatines; vomerine teeth in a V-shaped patch; 2 weak spines at angle of preoperculum, and a stronger one at angle of operculum. Gill openings wide, the membranes deeply cleft and not attached to the isthmus; vertical fins united; ventrals reduced each to a bifid ray. Branchiostegals 8. Pseudobranchiæ present, but small. Air bladder present. (*νέος*, new; *Bythites.*)

a. Scales about 88 in longitudinal series; depth 4⅜ in length. GILLII, 2884.
aa. Scales about 123 in longitudinal series; depth 5¾ in length; dorsal rays 101. MARGINATUS, 2885.

2884. NEOBYTHITES GILLII, Goode & Bean.

Body compressed, its height contained 4⅜ times in total length, and less than length of head; interorbital area convex, its width equal to diameter of circular eye, 3⅜ in length of head, and 1½ in length of snout in young. Head compressed, deeper than broad, with wide sinuses, its length contained 4½ times in that of the body; snout obtusely rounded, slightly produced; mouth large, the maxillary extending considerably behind the vertical through posterior margin of eye, expanded posteriorly; mandible still longer, its length about 2¼ times in height of body; interorbital space convex. Teeth in villiform bands in jaws and on palatines; vomerine patch subcircular, with angles extended posteriorly. Gill rakers moderately long and slender, somewhat numerous, the longest about ⅔ diameter of eye, 11 developed and 3 rudiments below the angle. Pseudobranchiæ absent; gill opening wide, the membrane deeply cleft, free from the isthmus behind. A single long, flat spine attached to posterior portion of operculum, high up, extending back to its edge; a small hidden spine at lower angle of preoperculum. Nostrils small, the anterior one in a very short tube, almost upon tip of snout; posterior nostril slightly larger, not tubular, immediately in front of middle of eye. Scales mod-

erate, upon head and body, in 88 vertical rows, 7 rows between dorsal origin and lateral line, which becomes obsolete in its posterior half, 16 or 17 from vent forward to lateral line; dorsal origin behind that of ventral and pectoral, its distance from snout contained 4 times in total length, its rays moderately long; anal origin under eighteenth dorsal ray, its distance from snout contained $2\frac{1}{8}$ times in body length, rays rather slenderer than those in the dorsal; caudal rays 6 or 7 in number, their length contained 9 times in total length, not differentiated from those of the adjacent fins; pectoral origin well forward, its base somewhat concealed by the flap of the operculum, its length about equal to $\frac{2}{3}$ that of head; ventrals each a bifid ray, the inner filament the longer, inserted slightly in advance of the base of the pectoral, not far from humeral symphysis, and reaching nearly to vent, its length nearly equal to height of body; distance from origin of ventral to vent slightly greater than height of body; color light yellow, with silvery reflections, with cloudings of brown above lateral line and numerous black chromatophores; a series of irregular brown blotches above the lateral line, with 1 or 2 much darker, extending upon the dorsal fin. In many specimens the color is uniform yellow, with simply the dark ocelli showing. (Goode & Bean.) Atlantic, in rather deep water, from Gulf Stream to the coast of Brazil. (Named for Dr. Theodore Gill.)

Neobythites gillii, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 601, Lat. 28° 36′ N., Long. 85° 33′ W., in 111 fathoms (Type, No. 37340. Coll. *Albatross*); GÜNTHER, Challenger Report, XXII, 103, 1887; GOODE & BEAN, Oceanic Ichth., 325, fig. 289, 1896.

Neobythites ocellatus, GÜNTHER, Challenger Report, XXII, 103, pl. 21, fig. B, 1887, off Pernambuco, in 350 fathoms.

2885. NEOBYTHITES MARGINATUS, Goode & Bean.

Head $4\frac{2}{3}$ in total length; depth $5\frac{1}{7}$. D. 101; scales 7–123–29. Body compressed, somewhat elongate; interorbital area convex, its width greater than the diameter of the circular eye. Mouth large, the maxillary extending considerably behind vertical through posterior margin of orbit, its length 2 in head; mandible slightly more than $\frac{2}{3}$ height of body. Teeth as in *N. gillii*. Gill rakers slightly longer than $\frac{1}{2}$ the diameter of eye, 7 and 3 rudiments below the angle of the anterior arch. Pseudobranchiæ absent. A long flat spine upon the upper edge of the operculum, extending back nearly to its margin; 2 short, flat spines upon the angle of the preoperculum. Nostrils as in *N. gillii*. Scales small, very closely imbricated, the lateral line obsolete in its posterior half. Distance of dorsal origin from snout 4 times in total length; anal origin under fourteenth dorsal ray, at a distance from the snout $2\frac{2}{3}$ times in total length. Caudal of about 8 or 9 rays, very closely placed, about $10\frac{1}{2}$ times in total length; pectoral placed much as in *Benthocometes*, its length about $2\frac{1}{8}$ times that of the head, extending to vertical through the vent; ventral a bifid ray inserted in advance of base of pectoral, not reaching to the vent, its length considerably less than height of body; distance of ventral origin from vent slightly more than height of body. Color light yellowish brown, an obscure narrow band of darker brown commencing on the snout, inter-

rupted by the eye, and extending backward ⅔ distance to tail; another beginning on the snout, extending over eye and back as far as first described, interrupted posteriorly; dorsal fin milky white at base in its anterior third; above this a blackish band extending whole length of fin; a narrow white margin above. The type is from *Blake* Station LXXIX, off Barbados, in 209 fathoms. (*marginatus*, edged.)

Neobythites marginatus, Goode & Bean, Bull. Mus. Comp. Zool., x, No. 5,162, 1883 off Barbados, in 209 fathoms (Coll. *Blake*); Goode & Bean, Oceanic Ichthyology, 326, fig. 290, 1896.

964. BENTHOCOMETES, Goode & Bean.

Benthocometes, Goode & Bean, Oceanic Ichthyology, 327, 1896 (*robustus*).

Brotulids, similar in appearance and structure to *Neobythites* and *Bassogigas*, distinguished by 2 short flat spines upon the anterior portion of the operculum, placed at some distance from each other, and by the absence of spines on the preoperculum. The lateral line is complete, and extends without interruption to the posterior fourth of the body, where it becomes obsolete. The vomerine teeth are bunched in a circular patch instead of being arranged in triangular form. The head is comparatively short, with the jaws in front nearly equal; the snout not produced, but obtuse, rounded, and almost declivous in its outline. Deep sea. Two species known. (*βένθος*, the depths; *κωμήτης*, inhabitant.)

2886. BENTHOCOMETES ROBUSTUS, Goode & Bean.

Body rather short and deep, its greatest height nearly 4⅜ in total length and about equal to length of head; interorbital area convex, its width greater than diameter of the circular eye, and 1½ times length of snout; head about 4 times diameter of eye; mouth moderate, the maxillary extending to vertical through posterior margin of eye, the mandible a little beyond, its length equal to that of postorbital part of head. Teeth in villiform bands in jaws and on palatines; vomerine teeth bunched in a circular patch. Gill rakers moderate, the longest a little more than twice in diameter of eye, 4 above angle of first arch, 11 below. Pseudobranchiæ rudimentary. Gill opening wide, the membrane deeply cleft behind, free from the isthmus. A pair of short flat spines upon the anterior portion of the operculum. Nostrils small, the anterior as close to the snout as the posterior ones are to the eyes; no apparent cirri. Scales minute; lateral line obsolete on the last fourth of body. Dorsal origin behind that of ventral and pectoral, its distance from snout 3⅜ times in body; height of dorsal fin moderate, the longest ray about 3 times in head; anal origin under eighteenth ray of dorsal, the height of fin about equaling that of dorsal; vertical fins not connate with the caudal, which consists of 12 or 13 very slender rays, its length nearly equal to ½ head; pectoral with a broad base, close to gill opening, its length nearly ⅔ that of head; ventral a single bifid ray, inserted in advance of vertical through base of pectorals, and not far from humeral symphysis, reaching nearly halfway to vent, the distance of which from the origin of the ven-

tral is equal to length of head. Color yellowish brown. The type of this species, a specimen 88 mm. long, was taken by the *Blake* from Station XCIV, off Moro Castle, Cuba, at a depth of from 250 to 400 fathoms. A collateral type specimen (No. 29057) was obtained by the *Fish Hawk* from Station 1043 in Lat. 38° 39′ N., Long. 73° 11′ W., at a depth of 130 fathoms. (Goode & Bean.) West Indies, to lat. 39° N., in Gulf stream. (*robustus*, stout.)

Neobythites robustus, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 161, 1883, off Moro Castle, Cuba, in from 250 to 400 fathoms. (Type in M. C. Z. Coll. *Blake*.)
Benthocometes robustus, GOODE & BEAN, Oceanic Ichthyology, 327, fig. 288, 1896.

965. BASSOGIGAS, Gill.

Bassogigas, GILL MS. in GOODE & BEAN, Oceanic Ichthyology, 328, 1896 (*gillii*).

Brotulids having the body elongate, compressed, covered with a thick, heavy skin, which upon the head covers and obscures all the angles of the skull; scales small, covering body and head completely; lateral line indistinct for the greater part of the course, but apparently extending at least ⅔ of the way from the operculum to the tail; eye moderate; vertical fins completely united; ventrals a pair of bifid filaments inserted behind the humeral symphysis and remote at their bases, short, rather stout; snout without barbels, slightly produced, the lower jaw being barely included; villiform teeth in the jaws, on the vomer and palatines; vomerine patch V-shaped, but with its arms broadly expanded and thicker at the angle, so that it is almost triangular; operculum with a long, sharp spine; preoperculum unarmed; branchiostegals 8; air bladder present; pseudobranchiæ small. Deep sea. (*βάσσων*, for *βαθύς*, deep; *γίγας*, giant.)

a. Dorsal fin with 83 rays; anal 67. GILLII, 2887.
aa. Dorsal fin with 95 rays; anal 82. STELLIFEROIDES, 2888.

2887. BASSOGIGAS GILLII, Goode & Bean.

D. 83; C. 6; A. 67. Head rather short and broad, with snout slightly overlapping the lower jaw; diameter of the eye scarcely ⅓ of the length of the snout and about 1½ that of head; maxillary extending far behind eye, the vertical from the anterior margin of orbit nearly bisecting it, its length ½ that of head, and its posterior margin ending in a broad triangular dilation; teeth normal; anterior and posterior nostrils separated by a space greater than diameter of eye; preoperculum with a square, rounded angle; no armature; operculum with a strong, sharp spine above, the tip of which projects slightly beyond the opercular flap; distance of vent from root of pectoral slightly more than length of head, as far removed from this point as is the anterior nostril; scales moderate, covering the entire head; lateral line somewhat conspicuous, obsolete in its posterior third. Dorsal and anal fins enveloped in thick scaly skin; origin of dorsal in advance of middle of pectoral; pectorals rounded, broad, and very short; less than ½ as long as head and extending about ⅓ distance from origin to vertical from vent; ventrals inserted somewhat behind angle of preoperculum, extending to vertical from axil of pectoral, and about ⅔ of distance from origin to vent; each ventral filament bifid, the inner

part being the longer. Color uniform grayish brown; fins darker. The type of this species was obtained by the *Albatross* from Station 2684, off Cape Henlopen, Delaware, in Lat. 39° 35′ N., Long. 70° 54′ W., at a depth of 1,106 fathoms. (Goode & Bean.) (Named for Dr. Theodore Gill.)

Bassogigas gillii, GOODE & BEAN, Oceanic Ichthyology, 328, fig. 291, 1896, off Cape Henlopen, Delaware, in 1,106 fathoms. (Type, No. 39417. Coll. *Albatross*.)

2888. BASSOGIGAS STELLIFEROIDES (Gilbert).

Head 4 to 4⅓ in length; depth 5 to 5¼. D. 95; A. 82; scales 110. Physiognomy strikingly like that of the Sciænoid genus *Stellifer*. Mouth large, oblique, the lower jaw included, maxillary reaching well beyond orbit, ½ length of head. Teeth uniform, small, in narrow bands, those on vomer in a ⌒-shaped patch; a well-developed band on palatines; tongue smooth, a well-developed dentigerous crest on median line behind it; no barbel at symphysis. Snout short, bluntly rounded, about equaling diameter of orbit, slightly overhanging mouth, 5 in head; interorbital width 4; upper limb of preopercle extending obliquely downward and backward, largely adnate, the angle produced into a free membranaceous flap which entirely conceals the narrow interopercle, and bears no spines. The structure of the gill flap does not appear to have been correctly interpreted. The opercle is strong, but of small extent, forking at its base, 1 branch continued straight backward as a strong spine, the second a narrow flat process downward and somewhat backward, parallel with and little distant from margin of preopercle. Filling the deep notch between these 2 processes, and forming the greater portion of the gill flap, is the thin membranaceous subopercle. Branchiostegal rays 7. Gill rakers long and slender, the longest ⅔ diameter of orbit, 7 above angle, 13 and about 5 rudiments below. Nape midway between front of dorsal and front of eye; dorsal and anal similar, uniform, low, joined to base of caudal, the latter truncate, projecting well beyond them; ventrals inserted under angle of preopercle, each of a single ray forked to the very base, the 2 branches united by membrane for a distance equaling ⅔ orbit, the inner filaments being longest, ⅓ longer than head, and extending well beyond front of anal; pectorals long and narrow, 1½ in head; a narrow membranaceous flap connecting base of pectorals with upper angle of opercular flap. Scales small, well imbricated, entirely investing body and head, including gular membrane and part of gill membranes; lateral line nearly complete, lacking for about ⅛ length of body, running high, parallel with dorsal outline. Color silvery gray, dusted with coarse black specks, darker along dorsal outline; dorsal and anal with a narrow light streak at base, otherwise dusky, becoming black posteriorly, and with a narrow white margin; caudal black, with a broad white terminal bar; pectorals and ventrals white, with few black specks; peritoneum silvery white; mouth white anteriorly, its posterior portion and gill cavity jet-black. Pacific Ocean, off coast of Lower California. Many specimens from *Albatross* Station 2996, in 112 fathoms. Length 7 inches. (*Stellifer*, a genus of *Sciænidæ*; *εἶδος*, resemblance.)

Neobythites stelliferoides, GILBERT, Proc. U. S. Nat. Mus. 1891, 112, off Lower California. (Type, No. 44383. Coll. Dr. Gilbert.)

966. BARATHRODEMUS, Goode & Bean.

Barathrodemus, Goode & Bean, Bull. Mus. Comp. Zool., x, No. 5, 200, 1883 (*manatinus*).

Body brotuliform, much compressed; head compressed; mouth moderate. Head unarmed, except for a short flattened spine at upper angle of opercle. Snout long, projecting far beyond premaxillaries, its tip much swollen; jaws subequal in front. Teeth minute, in villiform bands on jaws, vomer, and palatines. No barbels. Anterior nostrils on the outer angles of the dilated snout, circular, each surrounded by a cluster of mucous tubes. Posterior nostrils above front of eye. Gill openings wide, the membranes not united. Gill rakers rather few. Body and head covered with small, thin, scarcely imbricated scales. Dorsal and anal long. Caudal fin separate, long, and slender. Ventrals close together, far in front of pectorals, each reduced to a single bifid ray. Deep-sea fishes. (*βάραθρον*, a gulf or deep abyss; *δῆμος*, people.)

2889. BARATHRODEMUS MANATINUS, Goode & Bean.

Head about 6 in total length; depth 7½. D. 106; A. 86; C. 2+5+2; P. 18 to 20; V. 1/1; scales about 175. Body much compressed. Dorsal and anal outline approaching at an equal angle the horizontal axis. Scales small, about 175 rows between the branchial opening and the tail, and about 34 rows, counting upward and forward obliquely from the origin of the anal to the dorsal line; lateral line apparently absent. Head considerably compressed, with rounded upper surface, its width contained 2½ times in its length, its greatest height equaling ⅔ its length. Snout slightly longer than the horizontal diameter of the eye, and projecting beyond tip of upper jaw a distance equal to vertical diameter of eye, much dilated and swollen, the anterior pair of nostrils being situated at the most salient angles; snout in general form resembling that of a manatee, whence the specific name. Mouth moderate, its cleft extending to the vertical from the center of the orbit; length of upper jaw equal to twice horizontal diameter of eye, and contained 2½ times in length of head; posterior portion of maxillary considerably expanded; maxillary largely included within a skinny sheath; when the mouth is closed the lower jaw is entirely included within the upper. Vomer and palatine with bands of teeth more than twice as broad as the bands of the intermaxillaries and on the mandible. Eye elliptical in form, its vertical diameter ⅔ of its horizontal, the latter being equal to distance from tip of snout to posterior nostril, and contained 5¼ times in length of head; distance of eye from dorsal outline equal to ½ its horizontal diameter, and to ⅕ height of head in a perpendicular through center of eye; interorbital space rounded, its width equal to horizontal diameter of eye. Dorsal fin inserted in the vertical above insertion of pectoral, at a distance from end of snout equal to that of insertion of pectoral; anal inserted under twenty-first to twenty-third dorsal ray, and at a distance from snout about equal to ⅓ body length; height of dorsal and anal fins about equal to ½ height of body at insertion of anal, their bases extending almost to insertion of caudal; caudal composed of 9 rays, the 5 medial ones almost

equal in length, though the tip of the tail is slightly rounded, about equal to height of body midway between branchial opening and base of tail; ventrals inserted almost under middle of operculum, in length about equal to ½ length of head; pectorals inserted under origin of dorsal, and at a distance behind branchial opening equal to ⅔ vertical diameter of eye, its length equal to greatest height of the body. Color grayish brown; abdominal region black. (Goode & Bean.) Gulf stream, north of the Bermudas, in 647 to 1,395 fathoms. (*Manati*, like the manatee or sea cow.)

Barthrodemus manatinus, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 200, 1883, Lat. 33° 35′ 20″ N., Long. 76° W., in 647 fathoms (Type in M. C. Z. Coll. *Blake*); JORDAN, Cat., 127, 1885; GÜNTHER, Challenger Report, XXII, 100, 1887; GOODE & BEAN, Oceanic Ichthyology, 332, fig. 294, 1896.

967. NEMATONUS, Günther.

Nematonus, GÜNTHER, Challenger Report, XXII, 114, 1887 (*pectoralis*).

Body compressed, with long tapering tail. Bones of head soft, muciferous channels moderately developed, and with the integument very thin or absent on the upper portion and snout. Operculum cartilaginous and flat; a broad process near its upper angle corresponding to the opercular spine in some of the related genera, the head otherwise unarmed, though irregular by reason of the cranial bones. Snout much depressed, broad, rounded; jaws equal in front; mouth very wide; bands of villiform teeth in jaws, on vomer and palatines. Barbel none. Eyes small. Vertical fins confluent; ventrals a pair of bifid filaments close together, on the isthmus, close to the humeral symphysis. Gills 4, with very short laminæ and rather short, incurved, acicular gill rakers on the first arch, and much shorter, less numerous, spatulate ones on the 3 other arches. Pseudobranchiæ rudimentary. No traces of a lateral line, though the body is covered with scales of considerable size, almost as large as the eye, and the cheek with others still larger. *Nematonus* differs from *Porogadus* not only in the absence of spines upon the head, as Günther has indicated, but in the much less ossified opercular apparatus, in the shorter and thicker head, in the absence of the 3 series of pores simulating lateral lines, and in the tendency to prolongation in the lower rays of the pectoral, which increase from the uppermost to the lowermost in *Nematonus*, while *Porogadus* has a lanceolate fin, and also in the extreme exsertion of the caudal rays. (*νῆμα*, thread; *Onus*, the rockling.)

2890. NEMATONUS PECTORALIS, Goode & Bean.

D. 93; A. 73; P. 17; V. 2. Body moderately elongate, much compressed, the tail much shorter and more robust than in *Bassozetus catena*, its height equaling 1⅛ times length of head and ⅓ that of body. Head stoutish, not much compressed, lower than body, its length contained 5⅓ times in the body; snout compressed, broad at its tip, its length exceeding diameter of the circular eye; interorbital area slightly convex, its width slightly exceeding twice diameter of eye, 3 times in head. Maxillary

extending far behind eye, its length less than that of preorbital portion of head; mandible as long as postorbital portion of head; jaws, vomer, and palatines with narrow bands of villiform teeth, normally arranged. Branchiostegals 8. Gill lamellæ very short; gill rakers long and numerous, 18 on first arch below the angle, 5 above, 4 of which are rudimentary. Pseudobranchiæ present, but very rudimentary. Anterior nostrils on the top of the snout and near the median line of the head, near its tip, separated by a space about equal to diameter of eye; posterior nostrils in front of eye. Muciferous pores large, arranged much as in *B. catena*. Dorsal origin in the same vertical with that of pectorals, its distance from tip of snout contained 5 times in total and equaling twice length of maxillary. Rays well developed in anterior third, the longest ⅔ of head; anal origin under twentieth dorsal ray, its rays nearly as long as those of dorsal; pectoral with its penultimate ray produced, extending to thirteenth ray of anal, nearly twice as long as head; ventrals originating in advance of vertical through pectorals, and each a bifid filament; distance of ventral origin from tip of snout equaling length of ventral and about ¾ as long as head; distance of ventral origin from vent considerably greater than length of head; distance from tip of ventral to vent equal to ½ the length of the head. Number of scales in transverse series from vent to dorsal about 23; from the upper angle of the gill opening to the vertical through origin of anal 32. Color brownish yellow; head and abdomen blackish. The type (No. 37342, U. S. Nat. Mus.) was taken at *Albatross* Station 2380, Lat. 28° 02′ 30″ N., Long. 87° 43′ 45″ W. in 1,430 fathoms. It is 183 mm. long to the caudal base, 215 with caudal. Another young specimen, 70 mm. long, was taken at *Blake* Station XCV, off Dominica, in 330 fathoms. (Goode & Bean.) (*pectoralis*, pertaining to the breast.)

Nematonus pectoralis, Goode & Bean, Proc. U. S. Nat. Mus. 1885, 604, Lat. 28° 02′ 30″ N., Long. 87° 43′ 45″ W., in 1,430 fathoms (Type, No. 37342. Coll. *Albatross*); Günther, Challenger Report, xxii, 114, 1887; Goode & Bean, Oceanic Ichth., 333, fig. 295, 1896.

968. POROGADUS, Goode & Bean.

Porogadus, Goode & Bean, Proc. U. S. Nat. Mus. 1885, 682 (*miles*).

Body brotuliform, much compressed; head with numerous spines on interorbital space, 2 pairs on the shoulders, 1 at angle of operculum and a double series on angle of preoperculum; head with numerous mucous pores, as in *Bassozetus;* mouth large; snout moderate, not projecting much beyond the upper jaw; jaws nearly equal in front; teeth in villiform bands in jaws and on vomer and palatines; barbel none; gill openings wide, membranes narrowly united, not attached to the isthmus; gills 4; gill laminæ short; gill rakers moderate, numerous; pseudobranchiæ absent; caudal fin of few rays, on a very narrow base, not prolonged, scarcely differentiated from the vertical fins; dorsal and anal fins well developed; pectorals simple, moderate; each ventral a single bifid ray close to the humeral symphysis; branchiostegals 8; scales small; lateral line apparently triple, or replaced by 3 series of pores—1 close to ventral outline, 1 median, and another along base of dorsal. (πόρος, pore; *Gadus*, the codfish.)

2891. POROGADUS MILES, Goode & Bean.

Head 6½; depth 10; eye 5¾ in head; body much compressed, elongate, tapering to a very slender tail; head long, moderately compressed, subconical, the profile gradually ascending in nearly a straight line from tip of snout to origin of dorsal; interorbital space slightly convex, spiny, its width 4⅔ times in length of head, and slightly greater than diameter of eye; opercles and head generally covered with numerous and strong spines, as described in the generic diagnosis; mouth very large and wide; maxillary extending far behind eye and much expanded at its tip, its length more than ½ that of head; length of mandible equal to greatest height of body; jaws, vomer, and palatines with narrow bands of villiform teeth, none of which is enlarged; gill rakers 15 on anterior arch below the angle, 3 rudimentary ones above. Anterior pair of nostrils nearly on top of snout and somewhat nearer its tip than to eye, separated by a narrow space and placed immediately in front of middle of eye; behind each posterior nostril a strong spine projecting outward and upward; pores of the head arranged much as in *Bassozetus*; scales minute; lateral line not to be clearly made out; 3 rows of minute pores on each side of dorsal, median, and ventral, beginning near head and extending well toward extremity of tail. Dorsal origin slightly behind vertical through pectoral base, its distance from snout nearly 6 times in length of body, its rays moderately long, the longest about as long as snout, and very numerous; anal origin in vertical from twenty-second or twenty-third dorsal ray, its distance from snout 3⅓ times in length of body, its rays about as long as those of dorsal; pectoral imperfect, its length in the type equaling ½ that of head; ventrals a bifid filament, placed close to the humeral symphysis, well in advance of pectoral, its length equal to height of body; distance from origin of ventrals to vent nearly equal to length of head; ventral not reaching vent by a distance equal to length of snout. Color blackish brown. The type (No. 35625, U. S. Nat. Mus.) is 153 mm. in length, from *Albatross* Station 2230, lat. 38° 27′ N., long. 73° 02′ W., at a depth of 1,168 fathoms. (Goode & Bean.) (*miles*, a soldier.)

Porogadus miles, Goode & Bean, Proc. U. S. Nat. Mus. 1885, 602, Lat. 38° 27′ N., Long. 73° 02′ W., in 1,168 fathoms; Goode & Bean, Oceanic Ichthyology, 334, fig. 292, 1896.

969. PENOPUS, Goode & Bean.

Penopus, Goode & Bean, Oceanic Ichthyology, 335, 1896 (*macdonaldi*).

Body stout in front, tapering behind; tail not greatly exceeding the length of the rest of the fish; head scaly, thick, its top surface flat, with depressed and moderately projecting snout; a pair of minute postnasal spines; a strong and much curved spine on the operculum; several weak spines on the angle of the preoperculum, and several at the posterior angle of the suboperculum; mouth moderately large, the lower jaw included; several narrow slit-like pores along the margin of the preorbital and suborbital; 2 minute pores on under surface of mandible near its symphysis, and not far behind them 2 long slit-like pores; the

anterior nostril in a long slit, the posterior larger, oblong in shape, and ½ concealed by a fold of skin; eye small; the teeth appear only in minute asperities, the intermaxillary band much wider in front than behind; mandibulary band narrow throughout; vomerine band very narrow, V-shaped; palatines in a long, broad band; gill openings wide, deeply cleft in front, narrowly joined to the isthmus; branchiostegals 8; no pseudobranchiæ; gill rakers long and slender, not numerous; gill laminæ moderately long, a long slit behind the fourth gill; scales very small; lateral lines 3; caudal fin consisting of few rays, well differentiated from the dorsal and anal; dorsal beginning not far behind head; ventrals slightly in advance of the pectorals and composed of 2 rays, united by membrane, which forms a margin around them; pectoral normal, several of its upper rays simple; vent not much in advance of middle of total length. This genus agrees with *Porogadus* in nearly every respect except in the scarcity of spines on the head and in the structure of the ventrals. *Porogadus* has the ventrals composed of 2 distinct rays which are separated throughout their entire length, but in *Penopus* the 2 rays are inclosed in a membrane which connects them and forms a margin around them. In *Porogadus*, also, the suboperculum has a smooth margin and the opercular spine is weaker than in *Penopus*, and is not curved. Deep seas. (*πήνη*, thread; *πούς*, foot.)

2892. PENOPUS MACDONALDI, Goode & Bean.

D. 137; A. 102. Greatest height of body equaling length of postorbital part of head and about ⅓ of total without caudal; greatest width of body anteriorly about ⅔ of its greatest height; head stout, its greatest width equaling ⅔ of its greatest depth and more than ⅓ of its length; width of interorbital space about ⅕ length of head; eye very small, its length less than ½ width of interorbital space; distance from eye to tip of snout equaling length of intermaxillary; distance of anterior nostril from tip of snout equaling length of eye; distance of posterior nostril from eye slightly less than its distance from tip of snout; maxillary expanded behind and reaching somewhat behind eye, its length equaling that of snout; mandible extending much behind eye, its length equal to postorbital part of head. Dorsal beginning over middlle of pectoral, its rays well developed, those in middle of fin longer than anterior ones; anal beginning under twenty-seventh ray of dorsal, middle rays longest; pectoral nearly ½ length of head and about equal to distance of its tip from vent; ventral about ⅓ distance of its origin from origin of anal. Lateral lines 3, the uppermost beginning at the upper angle of the gill opening, quickly approaching top of body near base of dorsal and merging into dorsal base about middle of tail; median lateral line beginning a little behind head and extending almost to root of caudal, becoming very faint posteriorly; lowermost lateral line with its origin under and not far from base of pectoral, extending along lower side of tail and merging into base of anal fin somewhat beyond middle of length of tail. Color yellowish brown; operculum, opercular flap and branchiostegal membrane, pectoral, and ventral dusky. Only a single specimen, 315 mm. long, known; obtained by the *Albatross* September 18, 1886, at Station 2716, Lat. 38°

29′ 30″ N., Long. 70° 57′ W., in 1,631 fathoms. (Goode & Bean.) (Named for Hon. Marshall McDonald.)

Penopus macdonaldi, Goode & Bean, Oceanic Ichthology, 336, fig. 293, 1896, Lat. 38° 29′ 30″ N., Long. 70° 57′ W., in 1,631 fathoms. (Type, No. 39433.)

970. DICROLENE, Goode & Bean.

Dicrolene, Goode & Bean, Bull. Mus. Comp. Zool., x, No. 5, 202, 1883 (*intronigra*).

Brotulids with body moderately compressed; head somewhat compressed; mouth large; tip of maxillary much dilated; eye large, placed close to dorsal profile. Head with supraorbital spines; several strong spines on preopercle and 1 long spine at upper angle of opercle. Snout short, not projecting beyond the upper jaw; jaws subequal. Teeth in narrow villiform bands in each jaw, on head of vomer, and on palatines. No barbel. Gill openings wide, membranes not united; gills 4; gill laminæ of moderate length; gill rakers rather long, not numerous; pseudobranchiæ absent. Caudal not confluent with dorsal and anal, but without a distinct peduncle. Dorsal and anal fins long; pectoral rays in 2 groups, several of the lower ones being separated and much produced; ventrals a pair of bifid rays, close together on the isthmus. Branchiostegals 8. Body and head covered with small scales; lateral line close to base of dorsal fin, apparently becoming obsolete on posterior third of body. Stomach siphonal; pyloric cæca few and rudimentary; intestine shorter than body. Deep sea; a single species known. (*δικρόος*, forked; *ὠλένη*, limb.)

2893. DICROLENE INTRONIGRA, Goode & Bean.

Head 5; eye large, 4 in head; interorbital width 4. D. 100; A. about 85; C. 6 or 7; V. 1; P. 19+7 or 8; scales 110 to 120. Body moderately compressed, its dorsal and ventral outlines approaching at an equal angle the horizontal axis, and tapering to a narrow point. Head somewhat compressed, with flattish upper surface, which is encroached upon by the upper margin of orbit; a strong spine at posterior upper margin of orbit, pointing backward and upward; a long, sharp spine at upper angle of opercle, its exposed portion 2 in eye; 3 equidistant spines on lower posterior border of preopercle, much weaker than that on opercle. Large muciferous cavities in bones of head; a row of large cavities extending backward from upper angle of orbit, and continuous with those on lateral line. Mouth large, its cleft considerably more than $\frac{1}{2}$ head, the maxillary extending beyond eye and with scales upon its expanded tip. Distance from snout to origin of dorsal fin $\frac{3}{5}$ total length; anal inserted under twenty-fifth or twenty-sixth dorsal ray; height of dorsal and anal fins each about equal to eye; length of caudal fin 2 in distance from snout to dorsal; ventrals about equal to upper jaw; pectorals inserted close to branchial aperture, the 8 lower rays free and much prolonged, the longest and most anterior being about 3 in body, and more than 3 times as long as the contiguous posterior ray of the normally constructed portion of the fin, which is, however, about equal to the last free rays. West Indies, Gulf of

Mexico, and Gulf Stream in various localities, and off coast of Soudan and on the bank d'Arguin, in deep water. (*intro,* within; *niger,* black.)

Dicrolene intronigra, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 202, 1883, Gulf Stream, Lat. 39° 59′ 45″ N., Long. 68° 54′ W. (Coll. *Blake*); GÜNTHER, Challenger Report, XXII, 107, 1887; VAILLANT, Exp. Sci. Travailleur et Talisman, 258, pl. 23, fig. 2, 1888; GOODE & BEAN, Oceanic Ichthyology, 338, fig. 297 A and B, 1896.

971. MIXONUS, Günther.

Mixonus, GÜNTHER, Challenger Report, XXII, 108, 1887 (*laticeps*).

Lower pectoral rays free, not united by membrane with, but inserted on the same base as, the upper part of the fin; they are but slightly stronger than the other rays and prolonged. Body elongate, compressed, covered with small, very thin and deciduous scales. Head slightly compressed, broad and flat above, depressed in front, naked (with the exception of the parts between the mandibles, and, perhaps, of the cheeks). Bones thin, with muciferous system moderately developed; only 1 small spine above on the operculum; preoperculum without spine. Eye small. Vertical fins united, but the narrow caudal projecting beyond the short dorsal and anal rays. Ventrals each reduced to a filament, which consists of 2 rays firmly bound together in their whole length; they are inserted behind the humeral symphysis and close together. Snout broad, rounded, scarcely overlapping the lower jaw. Mouth very wide; villiform teeth in the jaws, on the vomer, and palatine bones. Gill laminæ short; gill rakers long, not very closely set. Pseudobranchiæ none. (*μίξις*, mixture, half; *Onus*, a synonym of *Gaidropsarus*, the rockling.)

2894. MIXONUS LATICEPS (Günther).

Head 2; depth 3; eye 8 in head; snout 4. P. 17. Greatest depth of body below origin of dorsal fin; distance of vent from snout ⅔ its distance from extremity of spinal column. Crown of head remarkably convex, covered with an extremely thin and transparent skin, which, perhaps, in older examples is scaly; interorbital space less convex, and equaling in width the length of snout including the eye; eye small, above middle of length of the maxillary; posterior nostrils wide, open, in front of the eye. Distance of vent from ventrals exceeds length of head; origin of dorsal fin above root of pectorals, its rays of moderate length, but longer than those of anal; pectoral with a rather narrow base, as long as head without snout, its rays feeble, 3 or 4 lower ones a little stouter, detached, and prolonged; ventral filaments not reaching as far backward as pectoral. Gill rakers 10, much longer than the laminæ. Whitish, with the abdomen and gill apparatus black. Mid-Atlantic, in profound depths. One specimen, 5½ inches long, was obtained in mid-Atlantic (*Challenger* Station 104), at the enormous depth of 2,500 fathoms. The second (type of *Sirembo guntheri*) was taken off Cape Verde, in 3,200 meters. (*latus*, broad; *-ceps*, head.)

Bathynectes laticeps, GÜNTHER, Ann. Mag. Nat. Hist. 1878, 20, mid-Atlantic, in 2,500 fathoms. (Coll. *Challenger*.)

Sirembo guntheri, VAILLANT, Exp. Sci. Trav. et Talisman, 268, pl. XXIV, fig. 5, 1889, off the Cape Verde Islands, at a depth of 3,200 meters.

Mixonus laticeps, GÜNTHER, Challenger Report, XXII, 108, pl. 25, fig. 8, 1887; GOODE & BEAN, Oceanic Ichthyology, 339, fig. 296 A, 1896.

972. BARATHRONUS, Goode & Bean.

Barathronus, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 164, 1883 (*bicolor*).

Brotulids having the head stout, body and tail compressed, covered closely by skin; scaleless; vent far behind pectoral, included in a cleft; mouth wide, oblique, the lower jaw projecting; intermaxillary teeth rudimentary; several fang-like teeth on the head of the vomer, none on palatines, a few rather large, recurved, separated teeth in the mandible; nostrils close together and small; eye visible through the skin, partly upon the top of the head, with or without dark pigment in the iris; barbel none; gill rakers very numerous and slender, and rather long; gill laminæ well developed on all the arches; no pseudobranchiæ; head full of mucíferous channels; gill membranes not united, but covered by a fold of skin; ventrals reduced to single simple rays, placed in advance of the pectorals and close to the humeral symphysis; dorsal and anal placed far back; caudal scarcely differentiated, composed of rather numerous, very slender rays upon a somewhat narrow base. (*βάραθρον*, the abyss; *ὄνος*, *Onus*, the rockling.)

2895. BARATHRONUS BICOLOR, Goode & Bean.

Head $5\frac{1}{5}$ in total, its width $\frac{2}{3}$ its length; depth $6\frac{1}{3}$; orbit $4\frac{2}{3}$ in head; interorbital width $4\frac{3}{4}$. D. about 70; A. 57. Body much compressed; eye concealed by the skin; maxillary extending slightly beyond the perpendicular through posterior margin of orbit, almost entirely concealed under the preorbital, and much expanded at tip, where its width is rather greater than that of eye. Intermaxillary very thin, broad, and slightly protractile; vomer very close to intermaxillary symphysis, its head somewhat raised and bearing 3 fang-like teeth (2 of which are off one side and 1 on the other in the type), separated by a moderately wide interspace; mandible with 5 enlarged, separate, recurved teeth upon each side, which increase in size posteriorly, its upper edge, posteriorly, produced above the level of the tooth-bearing surface, and received under the expanded maxillary; longest gill raker about as long as eye. Dorsal origin distant from the snout about $\frac{1}{2}$ total length; dorsal rays well developed, numerous, long, and slender, the longest about 3 times in length of head; anal originating in vertical from fourteenth dorsal ray, equidistant from eye and base of caudal, longest rays about as long as those in the dorsal; pectoral with a fleshy base, its length a little less than height of body. Ventral well in advance of pectoral, close to humeral symphysis, the rays being placed very close together at their origin, the length of the fin contained about 9 times in the total length, about 3 times in the distance from its origin to the vent. Caudal with about 10 rays, its length about 8 times in total length. Color yellowish white, with a broad vertical band of black from origin of ventral nearly to vent; another similar and narrower band above it upon each side. The type, 120 mm. long, from *Blake* Station LXXI, off Guadaloupe, at a depth of 769 fathoms. (Goode & Bean.) (*bicolor*, two-colored.)

Barathronus bicolor, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 164, 1883, off Guadaloupe, in 769 fathoms (Coll. *Blake*); GOODE & BEAN, Oceanic Ichthyology, 341, fig. 298, 1896.

973. APHYONUS, Günther.

Aphyonus, GÜNTHER, Ann. Mag. Nat. Hist. 1878, 22 (*gelatinosus*).

Head, body, and tapering tail strongly compressed, enveloped in a thin, scaleless, loose skin. Vent far behind the pectoral, at nearly the middle of the total length. Snout swollen, projecting beyond the mouth, which is wide. No teeth in the upper jaw; small conical teeth in the lower, pluriserial in front and uniserial on the side. Vomer with a few rudimentary teeth; palatine teeth none. Nostrils close together, small. No externally visible eye. Barbel none. Ventrals reduced to simple filaments, placed close together and near to the humeral symphysis. Gill membranes not united. Four branchial arches, the posterior without gill laminæ, the anterior with very short gill rakers and with rather short gill laminæ. Head covered with a system of wide muciferous channels and sinuses, the dermal bones being almost membranaceous, while the others are in a semicartilaginous condition. Notochord persistent, but with a superficial indication of the vertebral segments, as in some Leptocephaline forms. (Günther.) (*ἀφύη*, anchovy, a small translucent fish; *Onus*, the rockling.)

2896. APHYONUS MOLLIS, Goode & Bean.

Body much compressed, its greatest height 6 in its total length. Head thicker than body, its height slightly greater. Length of head about 4¼ in total, width over ½ its length. Snout 3⅓ in length of head. Eye not externally visible. Diameter of orbit, as seen through the skin, about ¼ length of head. Maxillary extending to vertical through posterior margin of orbit, the mandible somewhat farther back, its length nearly equal to height of body. A few weak teeth on vomer, palatines, and mandible, and very rudimentary ones in maxillary, not visible to the eye, but appreciable to the touch. Gill laminæ on the fourth and rudimentary gill rakers, 8 rudiments and 4 developed below the angle. Dorsal origin almost over posterior edge of operculum, its distance from the snout ¼ of total length, dorsal rays more than 110, well developed, the longest 3 in head; anal origin slightly nearer base of caudal than to the tip of snout, its rays shorter than those in the dorsal; pectoral with a fleshy base, its origin somewhat behind that of the dorsal, its length equal to width of head; ventral origin in advance of that of pectoral, close to humeral symphysis, the fin a single simple ray, whose length equals that of the pectoral, its tip not reaching vent by a space equal to height of head. Skin not loose. Texture of body rather firm, not transparent; whitish. Gulf of Mexico, in deep water. This species is closely allied to *Aphyonus gelatinosus*. (Goode & Bean.) (*mollis*, soft.)

Aphyonus mollis, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 163, 1883, Lat. 24° 36′ N., Long. 84° 5′ W., in 955 fathoms (Coll. *Blake*); GOODE & BEAN, Oceanic Ichthyology, 342, fig. 299, 1896.

Family CCXII. BREGMACEROTIDÆ.

Body stout, with robust caudal portion, truncate or convex behind, almost without procurrent caudal rays above or below; vent before mid-

dle of body; suborbitals moderate; no barbels, spines, nor cirri on head; mouth terminal, with minute teeth on jaws and vomer, none on palatines; ventrals jugular, extremely long, few-rayed, the rays dilated and separate nearly to base. Dorsal fins 2, the first an elongate, slender occipital ray; second dorsal on posterior half of body, of soft rays, depressed medially, so that it forms 2 lobes; no spines in fins. Anal nearly similar to the soft dorsal and similarly depressed in the middle; dorsal and anal depressible in a groove of scales. Hypercoracoid perforate; no pseudobranchiæ; gill openings wide, the membranes free from the isthmus. A single genus with 2 or 3 species found in the open sea, probably near the surface; widely distributed. The presence of the hypercoracoid foramen shows that this family is allied to the *Brotulidæ* rather than to the *Gadidæ*. From the *Brotulidæ* it is mainly distinguished by the development of its dorsal and ventral fins.

974. BREGMACEROS, Thompson.

Bregmaceros, THOMPSON, in Charlesworth's Mag. Nat. Hist., IV, 1840, 184 (*macclellandii*).
Calloptilum, RICHARDSON, Voy. Sulph., Fish., 94, pl. 46, figs. 4–7, 1843 (*mirum*).
Asthenurus, TICKELL, Journ. Asiat. Soc. Bengal 1865, 32 (*atripinnis*).

Characters of the genus included above. (*βρέγμα*, the upper part of the head, the nape; *κέρας*, horn.)

a. Scales in transverse series 14; scales in lateral series 58 to 64. MACCLELLANDII, 2897.

aa. Scales in transverse series 10; in lateral series 65; anterior lobes of dorsal and anal lower than in *B. macclellandii*. ATLANTICUS, 2898.

2897. BREGMACEROS MACCLELLANDII, Thompson.

Head $5\frac{2}{3}$; depth $6\frac{2}{3}$. D. about I, 18–X–22 (16 + X–15); A. about 18, X, 22 (22, X, 20); V. 4 or 5; scales 58–14 (64–14). Body moderately elongate, compressed, the form somewhat as in *Ophidion*, the back not elevated. Head short and small, moderately compressed; bones of head thin, without serrature or spine; eye moderate, 3 in head; interorbital space ridged, about as broad as eye; snout blunt, rather shorter than eye; mouth very oblique, the jaws subequal; maxillary reaching to beyond middle of eye, $2\frac{1}{4}$ in head; lower jaw flattish, curved upward; teeth in both jaws moderate, slender, close set, recurved, apparently in a single series. Tongue conspicuous; no teeth evident on vomer or palatines; branchiostegals 7 or 8; gill membranes separate, free from the isthmus; no evident pseudobranchiæ; gill rakers obsolete; no barbels about jaws. Body with rather large, thin, caducous scales (nearly all of them fallen in the typical specimens so that they can not be counted). Dorsal fin beginning with a single long and very slender spine on occiput, this nearly $\frac{1}{3}$ longer than head. Behind this, for a distance about equal to its length, the rudimentary rays, if present, do not rise above the sheath on each side. Nearly opposite the vent begins the dorsal proper, the distance of its first ray from snout being about $\frac{2}{5}$ length of body; about 12 rays are moderately elevated, about $\frac{3}{4}$ length of head. The others are gradually shorter and more slender, becoming too

small to count, until just before caudal, where the fin becomes conspicuous again, this posterior lobe not ½ so high as the anterior. Anal opposite dorsal and similar to it, the first ray close behind vent; caudal free from dorsal and anal, the caudal peduncle truncate at its base. Ventrals of 3 long rays, with a fourth at the inner base of the third; this fourth is probably a rudiment of 2. The ventrals are jugular in position, the rays very long and filamentous, the longest about ½ the body, reaching to the middle of anal fin. Pectorals inserted high, somewhat shorter than head. Vent slightly behind end of anterior ⅓ of total length. Color brown above, sides and below silvery; back and base of anal closely dotted with dusky; dorsal mostly dusky; caudal pale, dusky at base, with a narrow white cross bar; lower fins pale; the dark marking on front of back assume something of the form of lengthwise streaks. Tropical Pacific; Bay of Bengal; Philippine Islands; coast of China, etc., east to the coast of Central America, living near the surface in the open sea. Here described from the types of *Bregmaceros bathymaster*, two specimens, 1¾ and 2 inches in length, dredged at *Albatross* Station 2804, south of Panama, in 47 fathoms depth. Two others, 4 inches long, found later off the coast of Panama. A recomparison of these latter specimens with Günther's * detailed account of *B. macclellandii* shows no difference whatever, and we regard *B. bathymaster* as identical with the latter. Günther counts the scales 64–14; we find 58–14. In our largest specimens the ventrals reach middle of anal. (Named for Dr. John McClelland, of the Bengal Medical Service, who first studied the fishes of the Ganges.)

Bregmaceros macclellandii, THOMPSON, in Charlesworth's Mag. Nat. Hist., IV, 1840, 184, mouth of the Ganges; GÜNTHER, Cat., IV, 368, 1862.

Calloptilum mirum, RICHARDSON, Voyage Sulphur, Fish., 95, pl. 46, figs. 4–7, 1843.

Asthenurus atripinnis, TICKELL, Journ. Asiat. Soc. Bengal 1865, 32, with plate, Bay of Bengal.

Bregmaceros bathymaster, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, Lat. 8° 13′ 30″ N., Long. 79° 37′ 45″ W., southwest of Panama. (Type, No. 41137. Coll. *Albatross*.)

Bregmaceros atripinnis, DAY, Proc. Zool. Soc. Lond. 1869, 522, Bay of Bengal; types, same specimens described by TICKELL.

2898. BREGMACEROS ATLANTICUS, Goode & Bean.

Head 5⅔; depth 7⅔ in total length. D. I–15, X, 16; A. 15 or 16 + X (7 or 8) + 21 or 22; scales 65–10. Length 46 mm. Body compressed, moderately elongate. Interorbital area convex, its width greater than eye, which is 4 in head; jaws even in front; maxillary reaching to vertical through middle of eye; mandible to vertical through posterior margin of eye; teeth in intermaxillary minute, apparently in a single series; mandibu-

* The following is Dr. Günther's account of *Bregmaceros macclellandii*, taken from specimens from the China Sea:

"B. 7; D. I, 16+X+15; A. 22 | X+20; V. 5 or 6; scales 64–14. Occipital ray very slender, longer than head; dorsal and anal fins depressible in a groove formed by the scales along the bases of these fins; anterior portions of dorsal and anal elevated, connected with the posterior lower portion by a series of very short extremely feeble rays. Vent at end of anterior third of total length. Three outer rays of ventral fins dilated, compressed, simple, much elongate, reaching to or nearly to middle of anal; the second and third rays sometimes united at base. Silvery, minutely dotted with brown."

lary teeth biserial, the inner teeth enlarged. Cephalic appendage reaching nearly to base of first dorsal, its length 4½ in total. Distance of dorsal from snout 2½ in total, that of the anal the same; the dorsal and anal fins received in a groove formed by the scales along their bases; anterior portion of second dorsal and second anal less elevated than in *B. macclellandii.* The differentiations between the developed and undeveloped rays of the anal are so slight that the limits of the so-called anterior and posterior sections of the fin can not be determined. Length of the longest anal ray about 2 in body length. Specimens were obtained by the *Blake* at the following stations: XCIX, off Granada, 90 fathoms; CXIII, off Neris, 305 fathoms; CLXXXV, Lat. 25° 33′ N., Long. 84° 21′ W., 101 fathoms. (Goode & Bean.) This species seems doubtfully distinct from *B. macclellandii.* (*atlanticus*, of the Atlantic.)

Bregmaceros atlanticus, GOODE & BEAN, Bull. Mus. Comp. Zool., XII, No. 5, 165, 1886, West Indies, off Granada and Neris (Coll. *Blake*); GOODE & BEAN, Oceanic Ichthyology, 388, fig. 331, 1896.

Suborder ANACANTHINI

(THE JUGULAR FISHES.)

Vertical fins very long, destitute of true spines; tail isocercal, the posterior vertebræ progressively smaller; ventrals jugular, without spines; hypercoracoid without perforation or foramen; no pseudobranchiæ. The osteological characters of this group, called by him *Gadoidea*, are thus given by Dr. Gill:

"Jugulares with the orbito-rostral portion of the cranium longer than the posterior portion, the cranial cavity widely open in front; the supraoccipital well developed, horizontal and cariniform behind, with the exoccipitals contracted forward and overhung by the supraoccipital, the exoccipital condyles distant and feebly developed, with the hypercoracoid entire, the hypocoracoid with its inferior process convergent toward the proscapula, and the fenestra between the hypercoracoid and hypocoracoid." (Gill, Proc. Ac. Nat. Sci. Phila. 1884, 170.)

A large and important group, chiefly confined to the cold depths of the ocean and the northern seas. From all other typical fishes they are separated by the entire hypercoracoid. (*ἀν-* privative, without; *ἄκανθα*, spine.)

a. Caudal fin present; tail not greatly elongate; body tapering or coniform behind, with many procurrent caudal rays above and below; suborbitals moderate.

b. Frontal bones paired, with a triangular excavated area above, the divergent frontal crests continuous from the forked occipital crest; ribs wide, approximated, channeled below or with inflected sides; no barbels.

MERLUCCIIDÆ, CCXIII.

bb. Frontal bones normal, not forming a triangular excavated area above; ribs normal; chin with a barbel (rarely obsolete). GADIDÆ, CCXIV.

aa. Caudal fin wanting; tail very long, tapering behind; suborbitals very broad.

MACROURIDÆ, CCXV.

Family CCXIII. MERLUCCIIDÆ.

(The Hakes.)

Body moderately elongate, covered with small, smooth, deciduous scales; posterior part of body coniform and with the caudal rays procurrent forward; vent submedian. Head elongate, depressed, pike-like; suborbital bones moderate; mouth terminal, with strong teeth; no barbels; ventrals subjugular; dorsal fins 2, a short anterior and long posterior one, a long anal corresponding to the second dorsal; ribs wide, approximated, and channeled below or with inflected sides; frontal bones paired, excavated, with divergent crests continuous from the forked occipital crest. A single genus, with about 4 species; large cod-like fishes, of voracious habit, inhabiting moderate depths, and distinguished from the *Gadidæ* mainly by the structure of the frontal bones and the ribs. (*Merlucciidæ*, Gill, Proc. Ac. Nat. Sci. Phila. 1884, 772.)

975. MERLUCCIUS, Rafinesque.

(Hakes.)

Merluccius, Rafinesque, Caratteri di Alcuni Nuovi Generi, etc., 26, 1810 (*merluccius*).
Onus, Rafinesque, Indice d'Ittiol. Sicil., 12, 1810 (*riali* = *merluccius*); substitute for *Merluccius*.
Merlangus, Rafinesque, Indice d'Ittiol. Sicil., 30, 1810 (*riali*); substitute for *Onus*.
Merlus, Guichenot, in Gay, Hist. Nat. Chili, Zool., II, 328, 1847 (*gayi*).
Stomodon, Mitchill, Rept. Fish. N. Y. 1814. 7 (*bilinearis*).
Homalopomus, Girard, Proc. Ac. Nat. Sci. Phila. 1856, 132 (*trowbridgei*).
Epicopus, Günther, Cat. Fish. Brit. Mus., II, 248, 1860 (*gayi*).

Body elongate, covered with small, deciduous scales. Head slender, conical, the snout long, depressed; a well-defined, oblong, triangular excavation at the forehead, bounded by the ridges on the separated frontal bones, these ridges converging backward into the low occipital crest; eye rather large; edge of preopercle free; preopercle with a channel behind its crest, crossed by short radiating ridges; mouth large, oblique; maxillary extending to opposite the eye; lower jaw longer; no barbels; jaws with slender teeth, of various sizes, in about 2 series, those of the inner row longer and movable; vomer with similar teeth; palatines toothless. Branchiostegals 7. Gill rakers long; gill membranes not united. Dorsal fins 2, well separated, the first short, the second long, with a deep emargination; anal emarginate, similar to second dorsal; ventral fins well developed, with about 7 rays; vertebræ peculiarly modified, the neural spines well developed and wedged into one another; frontal bone double and the skull otherwise peculiar in several respects. Species several, very similar in appearance; ill-favored fishes of soft flesh and fragile fins, inhabiting water of some depth. Large voracious fishes, little valued as food. (*Merluccius*, the ancient name, meaning sea pike.)

a. Scales moderate, about 110 in lateral line; teeth very strong. D. 10-36; A. 36. MERLUCCIUS, 2899.
aa. Scales small, 135 to 150 in lateral line; teeth moderate. D. 11 to 13-41; A. 41.
b. Ventrals long, about 1⅔ in head. BILINEARIS, 2900.
bb. Ventrals short, about 2½ in head. PRODUCTUS, 2901.

2899. MERLUCCIUS MERLUCCIUS (Linnæus).

(European Hake.)

Head large, $3\frac{1}{3}$; depth $6\frac{1}{2}$. D. 10–36; A. 36; vertebræ 25 + 26; scales 150. Ventrals a little more than $\frac{1}{2}$ head; teeth very long. Dusky above, silvery below; dorsal, caudal, and distal part of pectoral blackish; inside of opercle black; inside of mouth black posteriorly, pale in front; peritoneum black. Coasts of Europe, generally abundant, south to Madeira and Italy, straying to Greenland.* Here described from specimens taken at Genoa. The identity of the Greenland Hake with *M. merluccius*† is perhaps uncertain. (Eu.) (*merluccius*, ancient name; *mare*, sea; *Lucius*, pike.)

Gadus merluccius, Linnæus, Syst. Nat., Ed. x, 254, 1758, Europe; after authors.
Merluccius smiridus, Rafinesque, Caratteri, etc., 26, 1810; Jordan & Gilbert, Synopsis, 809, 1883; Lilljeborg, Sveriges Fiske, ii, 121, 1891.
Gadus ruber, Lacépède, Hist. Nat. Poiss., v, 673, 1803, Scotland; Dieppe; on notes by M. Nöel; young.
Gadus merlus, Risso, Ichth. Nice, 122, 1810, Nice.
Onus riali, Rafinesque, Indice d'Ittiol. Sicil., 26, 1810; substitute for *merluccius*.
Merlucius vulgaris, Fleming, Brit. Anim., 195, 1828; Günther, Cat., iv, 344, 1862.
Merluccius esculentus, Risso, Eur. Mérid., iii, 1826, 220, Nice.
? *Merluccius ambiguus*, Lowe, Proc. Zool. Soc. Lond. 1840, 37, Madeira.
Merluccius sinuatus, Swainson, in Lowe, Proc. Zool. Soc. 1840, 38.
Merlucius lanatus, Gronow, Cat. Fish., Ed. Gray, 130, 1854, Mediterranean.
Epicopus gayi, Günther, Cat., ii, 248, 1860, no locality; not *M. gayi*, Guichenot, which is the Chilian Hake.
Merluccius linnæi, Malm, Götheborgs och Bohusläns Fauna, 489, 1877.

2900. MERLUCCIUS BILINEARIS (Mitchill).

(Silver Hake; New England Hake; Whiting.)

Head $3\frac{2}{3}$; depth $6\frac{1}{2}$. D. 13–41; A. 40; scales 100 to 110. Top of head with W-shaped ridges very conspicuous; eye shorter than snout and less than interorbital width; maxillary reaching posterior border of pupil; teeth not very large, smaller than in the European species, *Merluccius merluccius*. Scales larger than in other species; pectorals and ventrals long, the latter reaching $\frac{3}{4}$ distance to vent, their length about $\frac{3}{5}$ that of head. Grayish, darker above, dull silvery below; axil and edge of pectoral somewhat blackish; inside of opercle dusky silvery; inside of mouth dusky bluish; peritoneum nearly black. Coasts of New England and northward to Straits of Belle Isle; south, in deep water, to the Bahamas; rather common; used as food; breeding in deep water, though often taken near shore, northward. This species resembles the European Hake, *Merluccius merluccius*, but the latter has smaller scales, about 150, and larger teeth. (*bilinearis*, two-lined.)

* The Iceland Hake has been described as *Merluccius argentatus* (Faber). According to Faber, it has large teeth, the mouth white within, and the rays D. 15–43; A. 51; the fins deeply notched. It is perhaps a valid species, and, if so, it doubtless occurs in Greenland. (*argentatus*, silvered.)

Gadus merluccius (*argentatus*), Faber, Fische Islands, 90, 1829, Iceland.
Merluccius argentatus, Günther, Cat., iv, 346, 1862.

† "Dans l'Amérique du Nord, on cite ce poisson de Grœnland, mais l'exactitude de cette indication paraît douteuse." (Collett, Comp. Sci. Hirondelle, 1896, 58.)

Stomodon bilinearis, MITCHILL, Rept. Fish. N. Y., 7, 1814, **New York.**
Gadus albidus, MITCHILL, Journ. Ac. Nat. Sci. Phila., I, 1817, 409, **New York.**
Merlucius albidus, STORER, Hist. Fish. Mass., 363.
Merlucius bilinearis, GOODE & BEAN, Bull. Essex Inst., XI, 9, 1879, JORDAN & GILBERT, Synopsis, 809, 1883; GOODE & BEAN, Oceanic Ichthyology, 386, fig. 330, 1896.

2901. MERLUCCIUS PRODUCTUS (Ayres).

Head $3\frac{3}{4}$; depth 7. D. 11–41; A. 43; V. 7; scales 136. Head with the W-shaped ridges less strongly marked; maxillary reaching center of pupil; eye large; pectorals long and narrow, reaching vent; ventrals much smaller than in *M. bilinearis*, reaching halfway to vent, their length about $\frac{2}{3}$ that of head; caudal somewhat forked. Scales quite small, deciduous. Teeth moderate. Silver gray; head dusted with coarse black dots; inside of mouth and opercle jet-black; peritoneum silvery, with black specks. Length 3 feet. Pacific coast of America, from Santa Catalina Island northward to Puget Sound; everywhere abundant at moderate depths; used as food. (*productus*, drawn out.)

Merlangus productus, AYRES, Proc. Cal. Ac. Nat. Sci. 1855, 64, **San Francisco.**
Homalopomus trowbridgii, GIRARD, Proc. Ac. Nat. Sci. Phila. 1856, 132, **Astoria, Oregon.** (Coll. Lieut. W. P. Trowbridge.)
Gadus productus, GÜNTHER, Cat., IV, 338, 1862.
Merluccius productus, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 247; JORDAN & GILBERT, Synopsis, 809, 1883.

Family CCXIV. GADIDÆ.

(THE CODFISHES.)

Body more or less elongate, the caudal region moderate, coniform behind, and with the caudal rays procurrent above and below; vent submedian; suborbital bones moderate; scales small, cycloid; mouth large, terminal; chin with a barbel, more or less developed. Gill openings very wide; gill membranes separated or somewhat united, commonly free from the isthmus; no spines, the fin rays all articulated. Dorsal fin extending almost the length of the back, forming 1, 2, or 3 fins; anal fin long, single or divided; caudal fin distinct, or confluent with the dorsal and anal; ventral fins jugular, but attached to the pubic bone, each of 1 to 8 branched rays. Gills 4, a slit behind the fourth. No pseudobranchiæ. Edge of preopercle usually covered by skin of head. Pyloric cæca usually numerous, but sometimes few or none. Air bladder generally well developed. Genera about 25, species about 140; an important family, many of its members being highly valued as food. They inhabit chiefly the northern seas, sometimes venturing into the oceanic abysses. One genus (*Lota*) is confined to the fresh waters. (*Gadidæ*, Günther, Cat., IV, 326–369.)

GADINÆ:

a. Anal divided into 2 separate fins; dorsal fin divided into 3.
b. Lower jaw distinctly projecting; barbel small or obsolete; caudal concave behind.
c. Teeth in upper jaw slender, wide, set in 1 or 2 series; caudal forked. BOREOGADUS, 976.
cc. Teeth in upper jaw in a villiform band, the outer somewhat larger; caudal lunate.

d. Subopercle and postclavicle normal, both thin and flat, not enlarged and ivory-like. POLLACHIUS, 977.

dd. Subopercle and postclavicle enlarged, the bone dense and smooth, like ivory. THERAGRA, 978.

bb. Lower jaw included; barbel well developed; caudal not concave behind.

e. Hypocoracoid not swollen and ivory-like; lateral line pale; supraoccipital crest moderate.

f. Transverse processes of vertebræ thickened, swollen, and ivory-like at tip; small codfishes of the Arctic. ELEGINUS, 979.

ff. Transverse processes of vertebræ not swollen at tip.

g. Vent in front of second dorsal; size very small. MICROGADUS, 980.

gg. Vent below second dorsal; typical codfishes of large size. GADUS, 981.

ee. Hypocoracoid much swollen and ivory-like; lateral line black; mouth small, the maxillary not reaching to opposite eye; supraoccipital crest very high. MELANOGRAMMUS, 982.

aa. Anal fin forming a continuous fin or sometimes deeply notched; dorsal not divided into 3 fins.

h. Dorsal fin divided into 2 fins.

i. Anterior dorsal composed of distinct rays, similar to those in second dorsal.

j. Ventral fins rather broad, each of about 6 rays.

MORINÆ:

k. Anal fin with a deep notch.

l. Snout not much depressed, its edge without keel; tail slender. LEPIDION, 983.

ll. Snout flat, depressed, keeled on the edge; tail attenuate. ANTIMORA, 984.

LOTINÆ:

kk. Anal fin not notched; mouth terminal.

m. Vomer toothless.

n. Teeth in jaws unequal, outer series enlarged.

o. Barbel obsolete. URALEPTUS, 985.

oo. Barbel well developed. LOTELLA, 986.

nn. Teeth in jaws all villiform; barbel developed. PHYSICULUS, 987.

mm. Vomer with teeth; head not compressed.

p. Vomer and mandible without canines. Fresh-water species. LOTA, 988.

pp. Vomer and mandible armed with canines. Deep-water species. MOLVA, 989.

PHYCINÆ:

jj. Ventral rays very slender, each of 1 or 2 rays.

q. Ventrals each of 2 or 3 slender rays. UROPHYCIS, 990.

qq. Ventrals each of a single bifid ray. LÆMONEMA, 991.

GAIDROPSARINÆ:

ii. Anterior dorsal formed of a single slender ray, followed by a band of fringes; ventrals each of 5 to 7 rays.

r. Barbels 3; snout with 2 barbels, 1 at each nostril, none at tip; chin with 1 barbel. GAIDROPSARUS, 992.

rr. Barbels 4; snout with 3 barbels, 1 at tip of snout and 1 on each nostril; chin with 1 barbel, head high and compressed; no canines. ENCHELYOPUS, 993.

BROSMINÆ:

hh. Dorsal fin continuous, undivided; ventrals several-rayed; teeth on jaws, vomer, and palatines; mouth large; frontal bone. BROSME, 994.

976. BOREOGADUS, Günther.

Boreogadus, GÜNTHER, Cat. Fish. Brit. Mus., IV, 336, 1862 (*fabricii*).

This genus is closely allied to *Pollachius*, the body more slender, the caudal fin more deeply forked, and the teeth in both jaws slender, sharp, wide set, in 1 or 2 series. Small codfishes of the Arctic. (*βόρειος*, northern; *Gadus*.)

2902. BOREOGADUS SAIDA (Lepechin).

Head $3\frac{1}{2}$; depth $5\frac{1}{2}$; eye 4 in head; snout $3\frac{1}{3}$; interorbital space $4\frac{2}{3}$; gill rakers 9 to 13 + 30 to 32. D. 13–14–20; A. 16–21. Body slender, little compressed; head long, rather pointed, the lower jaw projecting; barbel minute; maxillary reaching middle of pupil; mandible 2 in head; teeth in upper jaw in 1 series, except in front, when the row is double; teeth in lower jaw uniserial; teeth nearly uniform in size, sharp, and wide set; teeth on vomer few, similar to those in jaws. Gill rakers numerous, long and slender, the longest $\frac{1}{2}$ eye; vent slightly before second dorsal; caudal peduncle slender, rounded, its depth scarcely more than $\frac{1}{2}$ eye. Pectorals reaching vent, $1\frac{1}{2}$ in head; ventrals $1\frac{1}{3}$, the second ray exserted for $\frac{2}{3}$ its length; first dorsal highest; front of second dorsal midway between tip of snout and base of caudal. Caudal forked for a distance equal to $\frac{1}{2}$ eye, the tips rounded. Color plain brownish, silvery below, the body with fine black points, most numerous above; dorsals and caudals dusky, the rays blackish distally, their edge narrowly white; anal similarly colored, pale at base; pectorals uniform dusky, pale-edged; ventrals somewhat dusky; peritoneum blackish. Length 6 to 8 inches. Arctic seas of Asia and America, from Greenland to Siberia; generally common in the far North, but rare in Bering Sea and south of Greenland. Here described (by Mr. Norman B. Scofield) from specimens from Davis Straits and Melville Bay, Greenland, the largest $6\frac{3}{4}$ inches long, and from specimens taken by Mr. Scofield at Point Barrow, Port Clarence, and Herschel Island. There is no difference between Greenland and Alaskan specimens. The range of fin rays is D. 12 to 15–12 to 15–18 to 22; A. 15 to 18–20 to 22. Concerning its habits Mr. Scofield observes:

"This fish appears to be quite abundant north of Bering Straits. It was especially brought to our notice by its habit of hiding in small holes in the floating ice, from which it was dislodged by our steamer striking and turning over the blocks of ice. This floating ice was usually in 7 fathoms of water and 1 or 2 miles from the coast. At Herschel Island we took it with the seine in shallow water along the beach. Lucien H. Turner reports it from St Michaels, where he took it through the ice in February, and was told by the natives that it appeared there only in winter. According to Richardson it spawns in Greenland in February, laying its eggs in the seaweeds along the shore under the ice." According to Richardson, in Northumberland Sound, "when hotly pursued by the Beluga or white whale, it has been observed, in its endeavors to escape, to leap by hundreds on the ice." (Eu.) (*saida*, Russian name.)

Gadus saida, LEPECHIN, Nov. Comm. Ac. Sci. Petrop. 1774, 512, White Sea; PALLAS, Zoogr. Rosso-Asiat., III, 199, 1811; GÜNTHER, Cat., IV, 337, 1862; COLLETT, Norske Nord-Havs Exped , 126, 1880; JORDAN & GILBERT, Synopsis, 307, 1883.

Merlangus polaris, SABINE, Supp. Parry's Voyage, CCXI, 1824, Baffins Bay; RICHARDSON, Last Arctic Voyage, 27, 1824.

Gadus fabricii, RICHARDSON, Fauna Bor.-Amer., III, 245, 1836, northern bays of Greenland; after *Gadus æglefinus* of Fabricius.

Gadus agilis, REINHARDT, Danske Vid. Selsk. Afh., VII, 126, 1838, Greenland.

Gadus glacialis, PETERS, Nord Pol. Expd., II, 172, 1874.

Pollachius polaris, GILL, Cat. Fish. East Coast N. A., 218, 1861.

Boreogadus polaris, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 233.

Boreogadus saida, BEAN, Bull. U. S. Nat. Mus., IV, 108, 1879; SCOFIELD, in Jordan & Gilbert, Rept. Fur Seal Invest., 1898.

977. POLLACHIUS, Nilsson.

(POLLACKS.)

Pollachius, NILSSON, in Bonaparte, Catalogo Metodico Pesci Europ., 45, 1846 (*pollachius*).

Body rather elongate, covered with minute scales; mouth moderate or large, the lower jaw projecting; barbel very small or obsolete; villiform teeth on vomer, none on palatines; teeth in jaws equal or the outer slightly enlarged; gill membranes more or less united; subopercle and postclavicle not enlarged and not ivory-like; dorsal fins 3; anal 2; caudal lunate; vent under first dorsal. Large fishes of the northern seas. (Polog or Pollack, the English vernacular name, latinized as *Pollachius*, as though derived from *πολλαχῆ*, many fashioned.)

2903. POLLACHIUS VIRENS (Linnæus).

(POLLACK; COAL-FISH; GREEN COD.)

Head 4; depth 4¼. D. 13-22-20; A. 25-20; scales about 150; vertebræ 54. Body rather elongate, compressed; snout sharp and conic; mouth rather small, oblique; maxillary reaching beyond front of orbit; lower jaw slightly the longer; teeth in the upper jaw nearly equal, the outer series not being especially enlarged; barbel rudimentary or obsolete; gill membranes considerably united, free from isthmus; vent under first dorsal; caudal fin lunate; pectorals short, scarcely reaching anal; ventrals short, their origin in front of base of pectoral a distance about equal to diameter of eye. Greenish brown above; sides and below somewhat silvery; lateral line pale; fins mostly pale; sometimes a black spot in the axil. North Atlantic; common northward on both coasts, south to Cape Cod and France. (Eu.) (*virens*, green.)

Gadus virens, LINNÆUS, Syst. Nat., Ed. X, 253, 1758, Seas of Europe; after *Gadus tripterygius imberbis* of the Fauna Suecica; GÜNTHER, Cat., IV, 339; JORDAN & GILBERT, Synopsis, 807, 1883.

Gadus carbonarius, LINNÆUS, Syst. Nat., Ed. X, 254, 1758, seas of Europe; after *Gadus dorso tripterygius imberbi* of ARTEDI.

Gadus colinus, LACÉPÈDE, Hist. Nat. Poiss., II, 416, 1800, England, etc.; after LE COLIN of Danberton.
Gadus virens, LACÉPÈDE, Hist. Nat. Poiss., II, 417, 1800.
Merlangus purpureus, MITCHILL, Trans. Lit. and Phil. Soc., I, 1815, 370, New York.
Merlangus leptocephalus, DE KAY, New York Fauna: Fishes, 288, pl. 45, fig. 146, 1842, Long Island.
Merlangus purpureus, STORER, Rept. Fish. Mass., 130, 1839.
Pollachius carbonarius, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 233.

978. THERAGRA, Lucas, a new genus.

(ALASKAN POLLACKS.)

Theragra, LUCAS, in Jordan & Gilbert, Rept. Fur Seal Invest. 1896 (1898) (*chalcogrammus*).

This genus is closely allied to *Pollachius*, differing in the following respects: Suboperculum thick, smooth and dense instead of being thin and squamous as in *Pollachius;* the postclavicle is also similar in structure while its proximal portion is subcircular in *Theragra* and rhomboidal in *Pollachius*; this ivory-like character of the suboperculum and postclavicle is so marked that it serves to distinguish these bones at a glance, being entirely different from what is found in the corresponding bones of other gadoids. The Alaskan Pollack farther differs from the Atlantic Pollack in having 19 precaudal vertebræ and 33 caudal, instead of 23 precaudals and 32 caudals; the bodies of the vertebræ are also slightly longer and more deeply sculptured in the Alaskan fish and the spinous process of the anterior dorsals less elevated. The vertebral differences between the 2 genera are merely differences of degree and of specific value only, but the differences between the subopercula and postclavicula are different in kind, distinguishing the Alaskan Pollack not only from the Atlantic Pollack, but from other gadoids. (θήρ, beast; ἄγρα, prey or food; the Alaskan Pollack being a chief food of the fur seal, *Callorhinus*.)

a. Dorsal rays about 13 or 14–17–18 or 19; anal rays 20–20; side with 2 interrupted dark longitudinal bands. CHALCOGRAMMA, 2904.
aa. Dorsal rays 10 or 11–13 to 15–16; anal rays 16 to 19–16 to 19; sides plain dusky; body less elongate, the snout blunter, the fins lower. FUCENSIS, 2905.

2904. THERAGRA CHALCOGRAMMA (Pallas).

(ALASKA POLLACK.)

Head 4; depth 6. D. 12–14–18; A. 20–20. Eye 5 in head; snout $3\frac{1}{6}$; maxillary $2\frac{1}{2}$; snout conic, sharp, rounded in profile; mouth oblique; maxillary reaching middle of pupil; chin with a minute barbel; teeth small, those of the outer row above slightly enlarged; eye large, wider than the flat interorbital space, 4 in head. Gill membranes somewhat united, the posterior outline deeply emarginate; vent under interspace between first and second dorsal; first dorsal higher than the others, the second lowest; ventrals filamentous, reaching nearly to vent; pectorals long, reaching past front of anal, $1\frac{1}{2}$ in head; caudal somewhat concave. Olivaceous above, sides silvery, with 2 interrupted stripes of dark, brassy,

olive along sides, these irregular on their edges, about ½ width of eye, with uneven edges; a trace of a third similar stripe below anteriorly, the stripes very irregular; back mottled. Dorsal plain dark olive; pectoral quite dark; lower fins ashy; caudal ashy olive. Bering Sea and neighboring waters, probably south to Sitka and the Kurils. Our specimens from Unalaska, Robben Reef, Komandorski and Pribilof islands and Bristol Bay. Excessively common throughout Bering Sea, swimming near the surface, and furnishing the greater part of the food of the fur seal. This animal rarely catches the true codfish, which swims nearer the bottom. Length 3 feet. (*χαλκός*, brass; *γραμμή*, line.)

Gadus chalcogrammus, PALLAS, Zoogr. Rosso-Asiat., III, 198, 1811, Kamchatka; GÜNTHER, Cat., IV, 340, 1862; JORDAN & GILBERT, Synopsis, 807, 1883.

Gadus periscopus, COPE, Proc. Am. Philos. Soc. Phila. 1873, 30, Unalaska (Coll. George Davidson).

Pollachius chalcogrammus, JORDAN, Cat. Fish. N. A. (130) 918, 1885.

Theragra chalcogramma, JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

2905. THERAGRA FUCENSIS (Jordan & Gilbert).

(WALL-EYED POLLACK; PUGET SOUND POLLACK.)

Head 3¾ in body; depth 5½. D. 10–13–16 to 11–15–16; A. 16–19 to 19–19; eye 4½ in head; maxillary 2⅔; pectoral 1⅔; longest caudal ray 2. Body elongate, not greatly compressed; mouth large, the maxillary reaching to below middle of eye; jaws with minute, sharp, curved teeth, the outer series enlarged; teeth on vomer, palatines toothless; lower jaw projecting, a very small barbel under its tip; interorbital space wide, very slightly and evenly convex, wider than the diameter of eye; nostrils much nearer eye than tip of snout, the posterior much the larger; head almost entirely covered with small scales; gill rakers numerous, the longest as long as pupil, about 5+27 in number. Distance of origin of first dorsal from snout 3⅛ in body; first rays of first dorsal reaching far past the ends of last rays where fin is depressed; first rays of other dorsals and anals scarcely reaching the base of last rays; caudal slightly forked or subtruncate when spread, the lobes subequal; end of pectoral reaching to front of anal; ventrals inserted in front of base of pectoral in distance a little more than diameter of eye, ending in a filamentous point. Color nearly plain sooty, with no distinct lateral bands, and with generally only a trace of a pale lateral streak along the side; on the head some diffuse dark spots; fins all dusky. The band of teeth in the premaxillary is wider than in *Theragra chalcogramma*, and the band is widened at the anterior end; the body is shorter; eye smaller; color darker; fins not so high; caudal not so deeply forked. Pacific coast, from Vancouver Island to Monterey, abundant in Puget Sound; probably northward to Kadiak, replacing *T. chalcogramma* to the southward. This form may intergrade with *Theragra chalcogramma*, though the original types seem well separated. Little is known of its range to the northward. Scofield and Seale took a specimen in Chignik Bay in northern Alaska, which

seems as near *T. fucensis* as *T. chalcogramma*. Its rays * are D. 11–16–17; A. 18–17; ventrals reaching $\frac{3}{5}$ distance to vent; interorbital space wider than eye; coloration dark. But its body is as slender as in *T. chalcogramma*. (*fucencis*, from the straits of Juan de Fuca.)

Pollachius chalcogrammus fucencis, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1893, 315, Puget Sound at Tacoma. (Type, No. 44455. Coll. David H. Hume.)

979. ELEGINUS, Fischer.

Eleginus, FISCHER, Mem. Soc. Nat. Moscow, v, 4, 2d Ed., 252–257, 1813 (*navaga*); not *Eleginus* of later authors.

Tilesia, SWAINSON, Nat. Hist. Class'n Fishes, II, 300, 1839 (*gracilis*); name preoccupied.

Pleurogadus, BEAN, in JORDAN, Cat. Fish. N. A., 130, 1885 (*gracilis*); substitute for *Tilesia* preoccupied.

This genus differs from the other codfishes in the structure of the transverse processes of the vertebræ, which are club-shaped, narrow at base, but expanding distally into a rounded hollow bulb at their tips. Skeleton otherwise essentially as in *Microgadus*, the skull similar. Small codfishes of the Arctic seas. (ἐλεγῖνος, a social fish mentioned by Aristotle.)

2906. ELEGINUS NAVAGA (Kölreuter).

(WACHNA COD.)

Head $3\frac{1}{3}$ in length of body; depth 6; eye $5\frac{2}{3}$ in head; snout 3; interorbital space $4\frac{1}{2}$; gill rakers 20 or 21; barbel small, equal to pupil; dorsal 13–18–18; anal 22–20; scales small, 157 transverse rows above lateral line from gill opening to first rudimentary caudal rays. Body slender and rounded with a rather long head; snout viewed from above rounded, but running to a rather sharp point when viewed from the side; lower jaw included, the fleshy snout projecting beyond the maxillary, its length slightly greater than that of the snout; tip of maxillary on a vertical with the front of the pupil; articulation of mandible with quadrate bone on a vertical running midway between pupil and posterior edge of eye; teeth

* The following is the count of fin rays in 13 specimens of *Theragra* of the two species:

Dorsal.	Anal.	Locality.
T. chalcogramma:		
13—15—20	19—20	Kamchatka.
13—15—20	19—21	Unalaska.
14—19—23	24—22	Pribilof Islands.
14—16—21	21—23	Do.
14—17—18	23—21	St. Paul Island.
13—15—19	21—22	Do.
14—17—19	21—20	Do.
14—17—18	22—20	Kamchatka.
11—16—17	19—17	Chignik Bay.
T. fucensis:		
10—15—17	18—16	Puget Sound.
11—15—16	19—18	Do.
10—14—16	16—19	Do.
12—13—17	19—19	Do.

all slender and curved backward, those in upper jaw in several irregular rows, the outer row regular and with slightly larger teeth; teeth in lower jaw in a single row except in front where they are in a double row; teeth on vomer few and about the size of the smaller teeth in the upper jaw; gill rakers moderate, the longest not quite equal to diameter of pupil; caudal peduncle compressed, its depth equal to diameter of eye; vent under front of second dorsal; pectoral fin not reaching vent, its length 1¾ times in head; ventrals reaching halfway to vent, the second ray moderately produced; first dorsal highest; distance between second and third dorsals twice distance between first and second; caudal fin very slightly concave; third ray of second dorsal midway between tip of snout and base of middle caudal rays. Color somewhat mottled, grayish brown above, light silvery below; the 3 dorsals and caudal dusky and edged with white; pectorals uniform dusky; ventrals but slightly dusted with black; anal with a few punctulations at their anterior ends; peritoneum pale. Arctic shores of Asia and North America, south to Bering Sea, locally abundant. It reaches the length of about a foot. Here described (by Norman B. Scofield) from numerous specimens, the largest 11 inches long, taken at Port Clarence by Scofield and Seale, and at Petropaulski by the *Albatross* (Fur Seal Invest. of 1896). The range of the fin rays is D. 12 to 15—18 to 21—18 to 21; A. 20 to 23—20 to 23. Mr. Scofield has prepared a skeleton of this species for comparison with that of *Microgradus proximus* from San Francisco. There is very little difference in the skulls. There is no difference in the neural spines of the vertebræ. The transverse processes of the vertebræ in *Microgadus proximus* are flattened and plate-like, while in *Eleginus navaga* they are club-shaped, narrow at base where they leave the centrum, but expanding into a rounded hollow bulb at the distal end. This character defines the genus *Eleginus*. (*navaga*, a Russian name.)

Gadus navaga, KÖLREUTER, Nov. Comm. Ac. Petrop., XIV, 1770, 484, pl. 12, coast of northern Russia; PALLAS, Zoogr. Rosso-Asiat., III, 196, 1811.

Gadus gracilis, TILESIUS, Mém. Ac. Imp. Petersb., II, 1810, 354, Kamchatka; JORDAN & GILBERT, Synopsis, 804, 1883.

Gadus wachna, PALLAS, Zoogr. Rosso-Asiat., III, 182, 1811, Kamchatka.

Tilesia gracilis, SWAINSON, Nat. Hist. Fish., II, 300, 1839; BEAN, Proc. U. S. Nat. Mus. 1881, 243.

Pleurogadus gracilis, BEAN, in JORDAN, Cat. Fish. N. A., 130, 1885.

Eleginus navaga, GILL, Proc. U. S. Nat. Mus. 1890, 303.

980. MICROGADUS, Gill.

(TOMCODS.)

Microgadus, GILL, Proc. Ac. Nat. Sci. Phila. 1865, 69 (*proximus*).

Very small codfishes allied to *Gadus*, but with the vent placed before the second dorsal and with a different structure of the cranium. The following is Professor Gill's account of the skull of *Microgadus proximus*, the italicised portions indicating the differences from *Gadus:*

The cranium is proportionally broader toward the front and less flattened, while the brain case is flattened below, *decidedly swollen* on each

side of a depressed *sphenoidal groove*, and has an ovate cardiform shape; the *paraoccipital* or epiotic is not produced into an angle behind, but is obtusely rounded, and its posterior or *outwardly descending ridge blunt;* the opisthotic is well developed, oblong, and with its reentering angle *high up*, and, on a line with it, the surface is divided into 2 parts—a *narrow* and a flattened one, and a lower expanded one, much swollen; the alisphenoid or prootic is *oblong*, acutely emarginate in front, swollen from the region of the high anterior sinus, and above a little produced forward; the great *frontal* is a little longer than broad, with supraoccipital crest *continued forward* on the bone, and near the front expanded upward, and with the *expanded portion* behind dividing into narrow *lateral wings;* the lateral testiform ridges of the frontal are continued forward and *curved outward* toward the antero-lateral angles; the anterior frontals are *mostly covered in front* by the great frontal, and are much *developed* in the direction of the antero-lateral angles, the inferior expanded axillary portion being very narrow; the nasal has a rounded ridge in front, continued well below, and its posterior crest is *laminar* and trenchant.

Species American; valued as food. (*μικρός*, small; *γαδος*, *Gadus.*)

a. Second anal with 21 or 22 rays; snout rather long; body semitranslucent; first anal and ventrals pale; body scarcely blotched with blackish. • PROXIMUS, 2907.

aa. Second anal with 16 to 20 rays; snout shorter; body opaque; first anal and ventrals dusky; body blotched above with blackish. TOMCOD, 2908.

2907. MICROGADUS PROXIMUS (Girard).

(CALIFORNIA TOMCOD.)

Head 3½ in body; depth 5. D. 14–18–18 to 21; A. 21 or 22–21 or 22; V. 6–7; eye 5 in head; maxillary 2½; pectoral 2; highest dorsal spine 2; middle caudal rays equal to snout. Head long, convex above, somewhat compressed, with vertical sides; eye moderate; mouth rather large; maxillary reaching to below pupil; barbel small; teeth in each jaw in a band, the outer row a little enlarged. Gill membranes a little connected, free from the isthmus. First dorsal highest, somewhat falcate; first anal longer and higher than second; pectorals moderate, reaching anal; ventrals filamentous, scarcely reaching anal; caudal slightly emarginate or subtruncate when fin is spread. Lateral line very distinct, wavy, high anteriorly, slightly interrupted posteriorly. Vent below first dorsal. Color olivaceous above, pale, or slightly translucent white below; dorsal fins dusky, paler at base; first anal and ventrals uncolored; second anal dusted with dark points. Monterey to Unalaska; abundant; a food-fish of considerable importance, the flesh delicate but without much flavor. Here described from a specimen, 8 inches in length, from Alaska, *Albatross* Station 3213. It reaches the length of about a foot. (*proximus*, near, to *Microgadus tomcod.*)

Gadus proximus, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 141, San Francisco; GIRARD, U. S. Pac. R. R. Surv., X, Fishes, 142, 1858; JORDAN & GILBERT, Synopsis, 805, 1883.
Morrhua californica, AYRES, Proc. Cal. Ac. Nat. Sci. 1854, 9, San Francisco.
Gadus californicus, GÜNTHER, Cat., IV, 332, 1862.
Microgadus proximus, GILL, Proc. Ac. Nat. Sci. Phila. 1865, 69.

2908. MICROGADUS TOMCOD (Walbaum).

(Tomcod; Frostfish.)

Head 4 in body; depth 5. D. 13 to 15–15 to 19–16 to 18; A. 17 to 21–16 to 20; eye 5 in head; maxillary $2\frac{2}{3}$; pectoral $1\frac{3}{5}$; middle caudal rays $2\frac{1}{4}$; first dorsal rays $1\frac{2}{3}$. Snout rounded, less produced than in *Microgadus proximus;* mouth short; maxillary $2\frac{1}{2}$ in head, reaching pupil; eye large, $3\frac{2}{5}$ in head; barbel small; pectorals reaching vent; ventrals filamentous, not reaching vent; vent under interval between first and second dorsals. Color olive brown, distinctly blotched and spotted with darker, lighter on the belly; more opaque than in *M. proximus;* back and sides profusely punctulate; dorsals and caudal blotched with darker; anals coarsely punctulate anteriorly, colorless posteriorly; ventrals and pectorals dusky. Virginia to Labrador; very common northward, and valued as a food-fish. Here described from a specimen, 9 inches in length, from Boston, Massachusetts. Length 1 foot. (*tomcod,* a vernacular name.)

Tomcod, Schöpf, Schrift. Naturf. Freunde, viii, 140, 1780, **New York.**
Gadus tomcod, Walbaum, Artedi Piscium, iii, 133, 1792, after Schöpf; Jordan & Gilbert, Synopsis, 806, 1883.
Gadus frost, Walbaum, Artedi Piscium, iii, 134, 1792, **North America**; after Frost-fish of Pennant.
Gadus tomcodus, Mitchill, Trans. Lit. and Phil. Soc., i, 1815, 368, **New York**; Günther, Cat., iv, 331, 1862.
Gadus pruinosus, Mitchill, Trans. Lit. and Phil. Soc., i, 1815, 368, **New York.**
Gadus tomcodus fuscus, Mitchill, Trans. Lit. and Philos. Soc., i, 1815, 369, **New York.**
Gadus tomcodus luteus, Mitchill, Trans. Lit. and Philos. Soc., i, 1815, 369, **New York.**
Gadus tomcodus mixtus, Mitchill, Trans. Lit. and Philos. Soc., i, 1815, 369, **New York.**
Gadus polymorphus, Mitchill, Trans. Lit. and Philos. Soc., i, 1815, 369, **New York.**
Morrhua americana, Storer, Rept. Fish. Mass., 120, 1839, **coast of Massachusetts.**

981. GADUS (Artedi) Linnæus.

(Codfishes.)

Gadus, Linnæus, Syst. Nat., Ed. x, 251, 1758 (*morhua*); after Artedi.
Morrhua, Oken, Isis 1817, 1182 (*morrhua;* on *les Morrhues* of Cuvier).
Cepphus, Swainson, Nat. Hist. Class'n Fishes, ii, 300, 1839 (*macrocephalus*).

Body moderately elongate, compressed and tapering behind. Scales very small; lateral line present, pale. Head narrowed anteriorly; mouth moderate, the maxillary reaching past front of eye; chin with a barbel; teeth in jaws cardiform, subequal; vomer with teeth; none on the palatines; cranium without the expanded crests seen in *Melanogrammus;* no part of the skeleton expanded and ivory-like. Dorsal fins 3, well separated; anal fins 2; ventral fins well developed, of about 7 rays. Species of the Northern Seas; highly valued as food. (*Gadus,* the Latin name, akin to the English word cod.)

a. Eye moderate, about $\frac{1}{3}$ snout in adult; axil without dusky spot.
b. Air bladder large. Atlantic codfish. CALLARIAS, 2909.
bb. Air bladder small. Pacific codfish. MACROCEPHALUS, 2910.
aa. Eye large, more than $\frac{1}{2}$ length of snout; axil with a dusky spot; caudal peduncle slender. OGAC, 2911.

2909. GADUS CALLARIAS,* Linnæus.

(Common Codfish.)

Head 3½ to 4½; depth about 4. D. 14–21–19; A. 20–18. Head large, but varying much in size; maxillary about reaching middle of orbit; occipital keel not greatly developed; teeth strong, cardiform, in narrow bands, those of the outer row in the upper jaw and of the inner row in the lower jaw somewhat enlarged. Eye moderate, about ⅓ length of snout. First dorsal little elevated, its height about ½ length of head; vent under front of second dorsal; caudal slightly emarginate; pectorals ⅓ length of head. Greenish or brownish, subject to many variations, sometimes yellowish or reddish; back and sides with numerous rounded brownish spots; lateral line pale; fins dark. North Atlantic, south to Virginia, and France; one of the most important of food-fishes. (Eu.) (*Callarias*, an old name of the codfish.)

Gadus callarias, Linnæus, Syst. Nat., Ed. x, 252, 1758, young examples, Baltic Sea and oceans of Europe, after *Gadus*, etc., *cauda integra* of the Fauna Suecica; Cuvier, Règne Animal, Ed. 2, vol. ii, 332, 1829; Jordan & Gilbert, Synopsis, 804, 1883.

Gadus morhua, Linnæus, Syst. Nat., Ed. x, 252, 1758, seas of Europe, after *Gadus*, etc., *cauda subæquali* of the Fauna Suecica; Richardson, Fauna Bor.-Amer., 242, 1836.

Gadus barbatus, Linnæus, Syst. Nat., Ed. x, i, 252, 1758.

Gadus vertagus, Walbaum, Artedi Pisc., iii, 143, 1792; after *Jägershen*, Klein, Hist. Nat. Pisc., v, 7, pl. 2, fig. 1, 1749.

?Gadus heteroglossus, Walbaum, *l. c.*, 144; after *Hornbogen* of Klein.

Gadus arenosus, Mitchill, Trans. Lit. and Philos. Soc., i, 1815, 368, New York.

Gadus rupestris, Mitchill, Trans. Lit. and Philos. Soc., i, 1815, 368, New York.

? Gadus nanus, Faber, Fische Islands, 113, Iceland.

Morrhua americana, Storer, Hist. Fish Mass., 343, 1867.

Gadus morrhua, Günther, Cat., iv, 328, 1862; Goode & Bean, Oceanic Ichthyology, 354.

2910. GADUS MACROCEPHALUS, Tilesius.

(Alaska Codfish.)

Head 3 in body; depth 4⅘. D. 13–18–16. A. 21–17; eye 6 in head; maxillary 2½; highest dorsal ray 3; pectoral 2½; middle caudal rays 4. Head large, the snout blunt; mouth large, the maxillary reaching to below front of pupil, snout projecting beyond mouth, lower jaw included; teeth strong, cardiform, in narrow bands on jaws and vomer; interorbital wide, 1½ times wider than diameter of eye, very slightly convex. Gill rakers moderate, about equal to pupil in length, 3 + 17 in number. Pectoral reaching to below end of first dorsal; ends of first dorsal rays reaching second dorsal when fin is depressed; ventrals inserted in front of base of pectorals in distance equal to diameter of eye; veins under front of second dorsal; caudal subtruncate. Color brownish, lighter below, back and sides with numerous brownish spots; fins, with the exception of first anal and ventrals, dusky. This species is very abundant in Bering Sea,

* We retain the name *Gadus callarias*, Linnæus for the codfish, instead of the commonly used name *Gadus morhua*, applied by Linnæus to the same species, because the name *Gadus callarias* stands first on the page on which it occurs. To accord priority to the name standing first is essential to fixity, and not the less so if the competing names are of the same actual date, published by the same author. It is not justice nor elegance, but fixity, which the rules of nomenclature aim to secure.

on both shores, in 15 to 130 fathoms, forming an important article of commerce. Its range southward extends to the offshore banks of Oregon. In external respects we recognize no distinction between this species and the common eastern codfish, except that the head seems larger. Here described from a specimen 20 inches long, taken in the Straits of Fuca by the *Albatross*. Concerning this species Dr. Gilbert observes:

It has been frequently pointed out, and is well known to fishermen that the Pacific codfish has a smaller air bladder or sound than the Atlantic cod. Pending an examination of this question, which we are not now in a position to make, we propose to recognize the Pacific fish as a distinct species. (*μακρός*, long; *κεφαλή*, head.)

Gadus macrocephalus, Tilesius, Mém. Acad. Sci. St. Petersb., II, 1810, 360, Kamchatka; Günther, Cat., IV, 330, 1862.
Gadus pygmæus, Pallas, Zoogr. Rosso-Asiat., III, 1811, Kamchatka.
Gadus auratus, Cope, Proc. Am. Philos. Soc., 1873, 30, Unalaska.

2911. GADUS OGAC, Richardson.

(Greenland Codfish.)

Head 3½. D. 14 or 15–18 to 20–17 to 20; A. 20 to 22–18 to 19; V–6. This species resembles the common cod (*Gadus callarias*), but differs from it as follows: It has a more slender caudal peduncle, larger eye, greater interorbital width, longer barbel, more advanced position of ventral fins, and a longer pectoral fin. Color dark, blackish brown above, lighter below, with yellowish marblings; the tip of the dorsal, anal, and caudal fins black; ventral and pectorals dark brown or black, a dusky spot on the axil; barbel black. Coast of Greenland; not seen by us. The above notes from specimens collected at Godhavn, Greenland, examined by Ensign Dresel. (*ogac*, a native name.)

Gadus ogac, Richardson, Fauna Bor.-Amer., 246, 1836, Greenland; Reinhardt, Vid. Selsk. Naturvid. Math. Afh. 1838; Dresel, Proc. U. S. Nat. Mus. 1884, 246.
Gadus ogat, Kröyer, Voy. Scand. et Lap., pl. 19.

982. MELANOGRAMMUS, Gill.

(Haddocks.)

Melanogrammus, Gill, Proc. Ac. Nat. Sci. Phila. 1862, 280 (*æglefinus*).
Ægleſinus, Malm, Götheborgs och Bohusläns Fauna, 481, 1877 (*æglefinus*).

This genus is distinguished from *Gadus* by its smaller mouth, the produced first dorsal fin, black lateral line, and especially by the great enlargement of the hypocoracoid, which is dense and ivory-like. The lateral line is always black, and the supraoccipital and other crests on the head are largely developed. Food fishes of large size. (*μέλας*, black: *γραμμή*, line.)

2912. MELANOGRAMMUS ÆGLEFINUS, Linnæus.

(Haddock.)

Head 3¾; depth 4½. D. 15–24–21; A. 23–21. Snout long and narrow, overlapping the small mouth; maxillary barely reaching front of orbit;

teeth subequal, large, in a cardiform band in upper jaw; in a single series on lower jaw and on vomer; occiput carinated; a ridge extending backward from each orbit; eye very large, $\frac{2}{3}$ length of snout, 4 in head. Anterior rays of first dorsal elevated, $\frac{2}{3}$ length of head, the fin pointed, higher than second and third dorsals; caudal lunate; vent below front of second dorsal. The skull in this species is more depressed than in *Gadus callarias*, broader, and thinner in texture; occipital crest exceedingly high, much higher than in *Gadus*, the wing-like projections at its base anteriorly spreading widely, raised above the surface of the skull. Dark gray above, whitish below; lateral line black; a large dark blotch above the pectorals; dorsals and caudal dusky. North Atlantic, on both coasts, south to France and North Carolina; in deeper water to Cape Hatteras; an important food-fish, reaching a considerable size. (Eu.) (*æglefinus*, an old name of the haddock, from the French Aiglefin or Aigrefin, according to Bellon; perhaps from *aigre faim*, extremely hungry, voracious.)

Gadus æglefinus, LINNÆUS, Syst. Nat., Ed. x, 251, 1758, seas of Europe, after *Gadus*, etc., *cauda biloba*, of the Fauna Suecica; JORDAN & GILBERT, Synopsis, 803, 1883.
Morrhua æglefinus, FLEMING, British Animals, 191, 1828.
Morrhua punctatus, FLEMING, British Animals, 192, 1828.
Melanogrammus æglefinus, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 280; *ibid.* 1863, 237; GOODE & BEAN, Oceanic Ichthyology, 354, 1896.
Æglefinus linnæi, MALM, Götheborgs och Bohusläns Fauna, 481, 1877.

983. LEPIDION, Swainson

Lepidion,* SWAINSON, Nat. Hist. Class'n Anim., I, 318, 1838, and II, 300, 1839 (*lepidion*).
Haloporphyrus, GÜNTHER, Cat. Fish. Brit. Mus., IV, 358, 1862 (*lepidion*).

Body elongate, covered with small scales; head not greatly depressed, higher than broad; the snout subconical, obtusely rounded; tail tapering behind; jaws with bands of villiform teeth; a roundish patch of teeth on vomer; no teeth on palatines; chin with a barbel; branchiostegals 7. Caudal fin separate; 2 dorsal fins and 1 anal; the first dorsal short; ventrals narrow, of 6 rays. Deep waters. The American species distinguished from the *Lepidion lepidion* (Risso), of the Mediterranean, by its non-filamentous first dorsal. (*λεπίδιον*, diminutive of *λεπίς*, scale:—small-scaled.)

2913. LEPIDION VERECUNDUM, Jordan & Cramer.

Head $3\frac{1}{3}$; depth $4\frac{1}{4}$. D. VIII–40; A. 37; V. apparently 4 (some rays broken on each side); scales about 75, not to be exactly counted. Body robust, compressed, tapering from the large head to the very slender, attenuate tail, which is not so broad as pupil; head large, not greatly compressed, not keeled above, its sides scaly; lower jaw with some scales; interorbital space depressed, $5\frac{1}{4}$ in head; eye very large (in young), $2\frac{2}{3}$ in head; snout short, depressed, not pointed, and with lateral keel, $5\frac{1}{4}$ in head; preorbital very narrow; mouth rather large, oblique, the maxillary reaching to below front of pupil, $2\frac{2}{3}$ in head; lower jaw slightly longer, its tip with a stiffish pointed projection representing the barbel; teeth small, in bands, a few

* *Lepidion* is sufficiently distinct from *Lepidia*, Savigny, 1817.

on vomer. No spines on snout or opercles. Gill membranes somewhat united, free from isthmus. Gill rakers slender, rather long, 10 to 12 on lower part of arch. Scales very small, mostly lost posteriorly (in our specimen) and not to be exactly counted; lateral line not evident. First dorsal rather low and long, none of its rays produced, the longest about ½ head; ventrals filamentous, ½ head; pectorals about ½ head; caudal 2½ head; anal deeply notched behind the middle, its posterior lobe highest. Color uniform purplish black, the fins paler. One young individual, 2¼ inches long, from *Albatross* Station 2993, off the Revillagigedo Islands. (*verecundus*, modest.)

Lepidion verecundum, JORDAN & CRAMER, Proc. U. S. Nat. Mus. 1896, 456, Revillagigedo Islands, at Albatross Station 2993. (Coll. *Albatross.*)

984. ANTIMORA, Günther.

Antimora, GÜNTHER, Ann. Mag. Nat. Hist. 1876, 2 (*rostrata*).

This group differs from *Lepidion* in the form of the snout, the backward position of the vent, the imperfect division of the anal, in which latter respect it approaches *Mora*. In *Lepidion* the snout is subconical, obtusely rounded; in *Antimora* it forms a flat, triangular lamina, sharply keeled at the sides, resembling the snout of *Macrourus*. Body elongate, compressed, tapering into a slender tail. Scales very small. Head entirely scaly, even to the gill membranes. Snout depressed, thin and flat, projecting beyond the mouth; mouth rather large; chin with a barbel; jaws with bands of villiform teeth; a small roundish patch of teeth on vomer, none on palatines. Dorsal fins 2, the first short, its anterior ray produced into a long filament; anal fin deeply notched, almost separated into 2 fins; ventral fins with 6 rays, 1 of them filamentous; caudal truncate. Branchi ostegals 7. Deep-water fishes. (*ἀντί*, opposite; *Mora*, a related genus.)

a. Head rather small, 4½ in length; scales 115. VIOLA, 2914.
aa. Head rather large, about 3½ in length; scales 130. MICROLEPIS, 2915.

2914. ANTIMORA VIOLA (Goode & Bean).

Head 4½ in body; depth 5. D. 4–53; A. 40; V. 6; scales 11–115–27. Snout broad, pointed at tip, much depressed, forming a roof-like projection above mouth; a conspicuous keel extending backward from tip of snout along the suborbital to the posterior margin of the eye. Mouth U-shaped, wholly inferior; maxillary nearly reaching posterior margin of orbit; interorbital space flat, as wide as the large eye, the orbital ridges somewhat elevated; barbel about ½ diameter of orbit. First dorsal with its first ray much produced, longer than head; anal fin deeply notched near its middle. Caudal peduncle as long as eye, its depth more than ½ its length; longest ray of ventrals reaching about halfway to vent; pectoral 1½ in head. Color deep violet or blue black; inside of mouth and opercles blue black. Banks of Newfoundland and southward, in deep water. (Goode & Bean.) (*viola*, violet.)

Haloporphyrus viola, GOODE & BEAN, Proc. U. S. Nat. Mus., 1, 1878, 256, La Have Bank, 400 to 500 fathoms; JORDAN & GILBERT, Synopsis, 800, 1883.
Antimora viola, GOODE & BEAN, Oceanic Ichthyology, 372, fig. 324, 1896.

2915. ANTIMORA MICROLEPIS, Bean.

Head about 4 in total length with caudal; depth $5\frac{2}{3}$ without caudal; eye 4 in head, nearly equal to snout. D. 4 or 5-51; A. 41; barbel very slender, 2 in eye; gill rakers short, slender, 4 + 11. Maxillary reaching to nearly below posterior edge of eye; longest ray of first dorsal about $\frac{1}{2}$ as long as head; anal deeply emarginate, beginning under twentieth ray of second dorsal; second ventral ray $1\frac{1}{6}$ in head. Scales very small, about 9 rows between origin of second dorsal and lateral line, and about 130 in lateral line. Color olivaceous, deeper on opercles and branchiostegal membranes and on inside of mouth. Off Queen Charlotte Islands. Several specimens taken by the *Albatross* at different stations in Bering Sea, at depths of 350 and 351 fathoms, and off the coasts of the Queen Charlotte Islands and California, at depths of 1,588 and 455 fathoms. One large specimen, from off Bogoslof Island, has the filamentous ray of first dorsal $\frac{7}{8}$ length of head, and the eye is shorter than the snout. (*μικρός*, small; *λεπίς*, scale.)

Antimora microlepis, Bean, Proc. U. S. Nat. Mus. 1890, 38, off Cape St. James, Queen Charlotte Island, at Albatross Station 2860, in 876 fathoms (Type, No. 45361); Gilbert, Rept. U. S. Fish Comm. 1893 (1896), 456 and 473; Jordan & Gilbert, Rept. Fur Seal Invest., 1898.

985. URALEPTUS, Costa.

Uraleptus, Costa, Archiv fur Naturgesh. 1858, 87 (*maraldi*).
Gadella, Lowe, Proc. Zool. Soc. Lond. 1843, 91 (*gracilis*).

Body elongate, compressed, and tapering posteriorly, covered with small scales. A separate caudal; 2 dorsal fins and 1 anal; ventral fins narrow, with flat base, composed of 6 rays. Upper and lower jaw with an outer series of strong curved teeth; vomerine and palatine teeth none; chin without barbel. Branchiostegals 7. Deep waters of the Atlantic. (*οὐρά*, tail; *λεπτός*, slender.)

2916. URALEPTUS MARALDI (Risso).

Head 4; depth 6. Head rather thick, its greatest width equal to its height, which is somewhat more than $\frac{1}{2}$ its length; cleft of mouth oblique, wide, the maxillary extending to below posterior margin of orbit; lower jaw received within the upper, but both nearly equal in length anteriorly, each armed with a series of rather large, curved, widely set teeth; another series of small teeth within the outer in the upper jaw. Snout rather broad, obtusely rounder, scarcely longer than eye, which is $4\frac{1}{2}$ in head; interorbital space emarginate on each side of upper part of orbit, its width somewhat more than diameter of eye. Nape broad, scarcely elevated, with a spine on each side pointing outwards and covered by skin. Operculum small, with a slender horizontal spine posteriorly, the part below the spine being emarginate; gill membranes united below the throat by a rather narrow cutaneous bridge, not attached to the isthmus; gill openings wide; gills 4, a slit behind the fourth; pseudobranchiæ glandular. Trunk rather low; tail tapering into a very narrow band;

first dorsal fin commencing behind vertical from base of pectoral, somewhat higher than long, and not higher than second; second dorsal commencing immediately behind the first, its rays increasing somewhat in length posteriorly, one of the longest being ½ as long as head, the whole fin naked; caudal fin slender, slightly rounded, entirely free from dorsal and anal, and nearly ½ as long as head; anal fin commencing at some distance behind the vent, which is situated below the origin of the first dorsal, very similar to the second dorsal; pectoral inserted somewhat below middle of body, its length equaling distance between front margin of eye and end of operculum; ventrals narrow, slender, the outer ray produced into a filament shorter than the pectoral. Scales extending over the whole head, the chin and the thin lips being naked. (Günther.) Tropical Atlantic. This form, originally described from Nice, has since been found at Madeira by Johnson, and at Naples and Catania by Giglioli. The *Blake* obtained a poor specimen, apparently of this form, at station LXXXI, off the Island of Nevis, in the West Indies. (Goode & Bean.) (Eu.) (A personal name for one of "quelques hommes que les talens, le mérite, la gloire ou l'amitié m'ont désignés.")

Gadus maraldi, Risso, Ichth. Nice, 123, pl. 6, fig. 13, 1810, Nice.
Merlucius attenuatus, Cocco.
Gadella gracilis, Lowe, Proc. Zool. Soc. Lond. 1843, 91, Madeira. (Type in University of Cambridge.)
Merlucius maraldi, Risso, Eur. Mérid., III, 220, 1826.
Uraleptus maraldi, Günther, Cat., IV, 349, 1862; Günther, Challenger Report, XXII, 87, 1887; Goode & Bean, Oceanic Ichthyology, 367, fig. 320, 1896.

986. LOTELLA, Kaup.

Lotella, Kaup, Archiv fur Naturgesch. 1858, 88 (*schlegeli*).

This genus differs from *Physiculus* chiefly in the presence in both jaws of an outer row of large teeth. Deep sea. (Name, a diminutive of *Lota*.)

2917. LOTELLA MAXILLARIS, Bean.

Head about 4⅓; depth 5. D. 5–55; A. 44; V. 10; scales about 7 or 8–115–14 or 15. Snout short; eye 3 in head; maxillary reaching vertical through anterior margin of pupil, its length equaling that of postorbital part of head. Teeth in narrow bands in jaws, the outer series enlarged; vomer and palate apparently without teeth. Vent situated about under eighth ray of second dorsal; distance of first dorsal from tip of snout 4 times in total length including caudal; ventrals extending to about vertical from origin of second dorsal, not reaching nearly to vent; longest ray of first dorsal a little more than ½ as long as head; none of the rays of second dorsal or of anal as long as first ray of first dorsal; longest ray of second dorsal not much exceeding ½ of height of body; longest ray of anal about ½ length of ventral; origin of anal about under tenth ray of second dorsal; ventrals situated about under beginning of posterior third of head, their length ⅓ that of second dorsal base; origin of pectoral somewhat in advance of that of first dorsal, the fin imperfect, but its length probably slightly exceeding that of ventral; caudal

rounded. Color very light brown; the margins of the dorsal and anal, in their posterior portions, blackish. (Goode & Bean.) Gulf stream, Lat. 40° N., in 396 fathoms. (*maxillaris*, pertaining to the upper jaw.)

Lotella maxillaris, BEAN, Proc. U. S. Nat. Mus. 1884, 241, Lat. 39° 55′ N., Long. 70° 28′ W. in 396 fathoms (Type, No. 29832. Coll. *Fish Hawk*); GOODE & BEAN, Oceanic Ichthyology, 368, 1896.

987. PHYSICULUS, Kaup.

Physiculus, KAUP, Archiv fur Naturgesch. 1858, 88 (*dalwigkii*).

Body elongate, covered with small scales; head entirely scaly; snout broad, obtusely rounded, projecting beyond the mouth; mouth of moderate size; chin with a barbel; jaws with bands of villiform teeth; vomer and palatines toothless. Dorsals 2; anal fin single, not notched; ventral fin with 5 rays, the outer ray filamentous; caudal rounded, slender, free; branchiostegals 7. (φυκίς, an ancient name of some fish living in the *Fucus*, φῦκος, probably a species of *Gobius*.)

a. Scales moderate, about 62 in a longitudinal series; gill rakers few; dorsal rays 10–49; anal 54; ventral reaching fourth anal ray. FULVUS, 2918.

aa. Scales very small, about 100 in a longitudinal series.

b. Gill rakers few, about 11 below arch; head 4 in length; depth 5; ventrals filamentous at tip, as long as head, reaching tenth ray of anal. NEMATOPUS, 2919.

bb. Gill rakers undescribed, probably few; head 4 in length; depth 5; ventrals shorter than head, reaching front of anal. KAUPI, 2920.

bbb. Gill rakers very numerous, 7+18; head 3¼ in length; depth 4⅔; ventrals reaching seventh anal ray. RASTRELLIGER, 2921.

2918. PHYSICULUS FULVUS, Bean.

Head about 4; depth 4⅔. D. 10–49; A. 54; V. 7; scales 6–61 to 62–16. Head broad and depressed; snout short; eye 3½ in head; the length of the upper jaw 2½ in head, about equal to space between ventrals and anal origin; maxillary not quite reaching vertical through hind margin of eye; barbel 6 in head. Teeth in narrow bands in jaw; no outer series of enlarged teeth, but a few in the middle of the bands in each jaw are slightly larger than the others; all of the teeth, however, inconspicuous; vomer and palate smooth. Vent situated about under third ray of first dorsal; distance of first dorsal from tip of snout equaling 3 times length of its base, its longest ray twice length of snout, and slightly exceeding length of longest of second dorsal; length of second dorsal base 3 times length of pectoral, which is nearly 5½ times in total without caudal. Origin of anal about in a vertical let fall from base of fifth ray of first dorsal; distance of ventral from tip of snout about 5¾ times in body. Tip of ventral when extended backward reaching base of fourth anal ray; length of middle caudal ray 3 in head. Lateral line very indistinct, situated rather high, following pretty closely the contour of back. Gill rakers moderately short and not numerous. General color a light yellowish brown; under surface of head, the abdomen, margins of dorsal and anal fins, lips, and axil of pectoral very dark brown; a dark brown blotch on the subopercular; inside of mouth and gill membranes white. (Bean.)

Caribbean Sea, north to 40° in region of Gulf Stream, reaching a depth of 955 fathoms. (*fulvus*, brownish yellow.)

Physiculus fulvus, BEAN, Proc. U. S. Nat. Mus. 1884, 240, Lat. 40° 1′ N., Long. 69° 56′ W. in 79 fathoms (Type, No. 28766. Coll. *Fish Hawk*); GOODE & BEAN, Oceanic Ichthyology, 366, 1896.

2919. PHYSICULUS NEMATOPUS, Gilbert.

Head 4 in length; depth 5. D. 7 to 9–56 to 61; A. 59 to 64; scales 90 to 105. Length of caudal peduncle to base of median caudal rays 5¼ in head. Snout very broadly rounded, its width twice its length, which is 4⅜ in head; eye 3¾; interorbital 4; maxillary 2, reaching slightly beyond vertical from posterior margin of orbit. None of the teeth enlarged; palate smooth. Branchiostegal membranes more narrowly joined than in the *P. rastrelliger*, but wholly free from isthmus. Gill rakers short and slender, 11 movable ones on horizontal limb of arch. Origin of first dorsal over base of pectorals, its distance from tip of snout 3⅜ in length; base of first dorsal equaling snout and ½ eye, its highest ray 2¼ in head; free portion of caudal peduncle ⅓ diameter of orbit; notch of dorsal and anal fins not conspicuous, the posterior dorsal rays little longer than those which precede, 2½ in head; caudal 2⅛; pectorals 1½; ventrals with broad base and 7 rays, the outer 2 filamentous, the second the longest, reaching base of tenth to twelfth anal rays, and as long as head; distance between bases of ventrals equals interorbital width; scales small, regularly imbricated, becoming minute on snout, which they completely invest, as well as mandible and gular membranes; lateral line present on anterior half of body only, 8 scales above it anteriorly. Color light olive brown, sprinkled with dark specks, the sides of head and trunk with silvery luster; snout, mandible, and gular membrane dusky; abdominal area, branchiostegal membranes, base of ventrals, axillary blotch, and front of anal, purplish black; posterior edge of gill membranes and opercular flap white; dorsals dusky, with an inconspicuous darker margin, which becomes more marked posteriorly; anal darker, margined with black; caudal blackish; pectorals and filamentous portion of ventral white. Inside of mouth and gill cavity white; peritoneum silvery, rendered black on sides by clusters of spots. Coast of southern California. Many specimens, the largest 7 inches long, from *Albatross* Stations 2997, 3011, 3015, and 3016, in 71 to 221 fathoms. (Gilbert.) (*νῆμα*, thread; *πούς*, foot.)

Physiculus nematopus, GILBERT, Proc. U. S. Nat. Mus. 1890, 114, coast of southern California. (Types, No. 46486 and 46555. Coll. Dr. Gilbert.)

2920. PHYSICULUS KAUPI, Poey.

Head 3⅓; depth 4. D. 10–60; A. 60; P. 30; V. 8; C. 17; scales 12 to 15—over 100. Body and head short, swollen; tail regularly narrowed; vent below base of pectoral; eye high, equal to snout, 4½ in head; nostrils with valves; snout blunt; lower jaw the shorter; maxillary reaching slightly beyond eye; each jaw with a band of cardiform teeth, none on palate and tongue; maxillary sloping under skin of cheek; barbel a little longer than eye; opercular bones covered with skin, without spines; no spines at

nape; gill membranes somewhat united, free from isthmus; gills 4, a slit behind fourth; no pseudobranchiæ; lateral line parallel with the back to beyond middle of body, then turning down suddenly, continuing to base of caudal. Scales small, cycloid, not easily counted; head scaly, even to the lips; vertical fins with small scales; ventral filamentous, equaling $\frac{2}{3}$ length of head, all the rays except the first short; ventral with 2 filamentous rays, which reach to front of anal and are about $1\frac{1}{2}$ in head; first dorsal as high as long, beginning behind base of pectoral, its longest rays about $2\frac{1}{2}$ in head; soft dorsal and anal low, free from the small rounded caudal; pectoral falcate, $1\frac{1}{2}$ in head. Color yellowish brown, bluish on belly; second dorsal and anal edged with darker brown. Type, 1 specimen, 250 mm. long. (Poey.) Deep waters of the Atlantic.

This species has constantly (as far as is shown in our specimens) a broader base to the ventral fins than *Physiculus dalwigkii*, and they are formed of 7 rays, of which the largest may or may not reach the anal fin; the fin rays vary within proportionate limits; they are, D. 9 or 10—60 to 66; A. 60 to 70; there are 13 scales between the anterior dorsal and lateral line; the caudal peduncle is shorter and less slender than in the Madieran form, but otherwise the species are so similar as to scarcely deserve specific separation. Poey obtained a specimen at Cuba, and Melliss 2 at St. Helena. These differ in no respect from 5 examples, 11 to 16 inches in length, found by the *Challenger* off Inosima in 345 fathoms. (Günther, Challenger Report, XXII, 88, pl. XVII, fig. A, 1887.) (Named for Dr. J. J. Kaup, author of a work on the Apodal fishes.)

Physiculus kaupi, POEY, Repertorio, I, 186, 1865, Matanzas. (Coll. Don Cirilo Dulzaides.)
? Physiculus japonicus, HILGENDORF, Sitz. Naturf. Freunde, Berlin, 1879, 80, Japan.

2921. PHYSICULUS RASTRELLIGER, Gilbert.

Head $3\frac{1}{2}$ in length; depth $4\frac{3}{4}$. D. 8 or 9–53 to 61; A. 57; scales 100 to 110. Length of caudal peduncle to base of median caudal rays, $2\frac{1}{3}$ in head. Snout short and broadly rounded, $4\frac{1}{4}$ in head; eye $3\frac{3}{5}$; interorbital width $4\frac{1}{8}$ to $4\frac{3}{5}$; maxillary $2\frac{1}{4}$, extending to vertical from posterior margin of pupil. Teeth in rather broad bands, none of them enlarged; width of patch on premaxillaries $\frac{1}{2}$ pupil; vomer and palatines toothless. Branchiostegal membranes broadly united, joined to the isthmus anteriorly, the width of the free fold more than $\frac{1}{2}$ pupil. Gill rakers numerous, slender, moderately long, the longest $\frac{1}{3}$ diameter of orbit, about 7 above angle, 17 to 19 below, the anterior ones short but movable. Origin of first dorsal slightly in advance of base of pectorals, its distance from tip of snout $3\frac{2}{7}$ in length; base of first dorsal equaling length of snout, its longest ray $2\frac{2}{5}$ in head; free portion of caudal peduncle equaling diameter of eye; second dorsal notched, the median rays $\frac{2}{3}$ the height of the highest anterior rays, the posterior highest, equaling first dorsal and longest caudal rays; anal similar to soft dorsal, but lower; ventrals under middle of opercle, the distance between their bases little less than interorbital width, equaling distance from vent to anal fin; ventrals with 7 rays, the outer 2 produced, the second the longest, reaching base of seventh or eighth anal ray; pectorals with broad base, covered with lax membrane,

containing 26 to 28 rays, their length 1⅓ to 1⅔ in head. Scales small, comparatively little reduced on top of head, a broad ring encircling snout in front of eyes naked, a very narrow patch of scales between this laterally and premaxillaries; scales in 100 to 110 transverse rows, 8 or 9 between lateral line and front of dorsal; lateral line wanting on posterior part of body, in the latter part of its course present on occasional scales only. Color uniform grayish olive on sides, each scale, or at least its marginal ½, closely covered with minute dark specks; gular and branchiostegal membranes, ventral region, and axil of pectorals blue black; basal portion of vertical fins light bluish, margined with blackish; pectorals dusky; ventrals blue black at base, the distal portion white; lining membrane of mouth and gill cavity white; peritoneum silvery, but in places so filled with black specks as to appear black. Coast of southern California. Many specimens, the longest 8 inches, from *Albatross* Stations 3045 and 2987, in 184 and 171 fathoms. (Gilbert.) (*rastrelliger*, bearing small gill rakers; *rastrum*, rake; *gero*, I bear.)

Physiculus rastrelliger, GILBERT, Proc. U. S. Nat. Mus. 1890, 113, coast of southern California. (Type No. 48266. Coll. *Albatross*.)

988. LOTA (Cuvier) Oken.

(BURBOTS.)

Les Lottes, CUVIER, Règne Anim., Ed. 1, vol. 2, 215, 1817 (*lota*).
Lota, OKEN, Isis 1817, 1182 (*lota*).

Body long and low, compressed behind. Head small, depressed, rather broad; anterior nostrils each with a small barbel; chin with a long barbel; snout and lower parts of head naked; mouth moderate, the lower jaw included; each jaw with broad bands of equal, villiform teeth; vomer with a broad, crescent-shaped band of similar teeth; no teeth on palatines. Gill openings wide, the membrane somewhat connected, free from the isthmus. Scales very small, embedded; vertical fins scaly. Dorsal fins 2, the first short, the second long, similar to the anal; caudal rounded, its outer rays procurrent; ventrals of several rays. One or 2 species, living in fresh waters of northern regions. (*Lota*, the ancient name used by Rondelet, in French, *la Lotte*.)

2922. LOTA MACULOSA (Le Sueur).

(BURBOT; LAKE LAWYER; LING.)

Head 4⅓ in body; depth 5½. D. 13–76; A. 68; ventral 7; eye 7 in head; pectoral 1⅓ in head; maxillary 2⅔; middle caudal rays 2⅔. Body elongate, not much compressed anteriorly; head slightly depressed; mouth large, the maxillary reaching to posterior margin of eye; teeth villiform, in bands on jaws and vomer; barbel longer than the small eye; interorbital broad, nearly twice diameter of eye; gill rakers very short, about 3 + 6 in number; anterior nostrils with barbels; body covered with small embedded scales; pectorals scarcely reaching to below front of dorsal; ending of ventrals filamentous; caudal rounded; vertebræ 21 + 38 = 59; cæca 30. Dark olive, thickly marbled and reticulated with blackish;

yellowish or dusky beneath; young often sharply marked, the adult becoming dull grayish; vertical fins with dusky margins. Length 2 feet. Lakes and sluggish streams. New England and Great Lake region, north to the Arctic seas and west to the headwaters of the Missouri, the Frazer River basin, and Bering Straits; abundant northward; rare in the Ohio River and the Upper Mississippi; a rather coarse and tasteless fish, seldom used as food. Here described from a specimen, 18 inches long, from Lake Michigan at Michigan City, Indiana. The American Burbot is very close to the common species of northern Europe and Asia, *Lota lota* (Linnæus) = *Lota vulgaris*, Cuvier = *Lota communis*, Rapp, and may prove wholly identical with the latter. In *Lota lota* the pectorals reach beyond front of dorsal, being 1⅓ in head. (*maculosus*, spotted.)

Gadus maculosus, Le Sueur, Jour. Ac. Nat. Sci. Phila., I, 1817, 83, Lake Erie.
Molva maculosa, Le Sueur, Mém. Mus., V, 1819, pl. 16.
Lota maculosa, De Kay, New York Fauna: Fishes, 284, pl. 52, fig. 168, 1842.
Gadus compressus, Le Sueur, Jour. Ac. Nat. Sci. Phila., I, 1817, 84, Connecticut River.
Lota compressa, De Kay, New York, Fauna: Fishes, 285, pl. 78, figs. 244, 245, 1842.
Gadus lacustris, Mitchill, Amer. Monthly Mag., II, 1818, 244, Sebago Pond, Maine (Coll. Henry A. S. Dearborn).
Molva huntia, Le Sueur, Mém. Mus., V, 1819, 161, Connecticut River.
Lota inornata, De Kay, New York Fauna: Fishes, 283, pl. 45, fig. 145, 1842, Hudson River, Lansingburgh, N. Y.
Lota brosmiana, Storer, Boston Journ. Nat. Hist., IV, 1839, pl. 5, fig. 1, New Hampshire.

989. MOLVA, Fleming.

(Lings.)

Molva, Fleming, British Animals, 192, 1828 (*vulgaris*).
Molva, Nilsson, Skandinav. Fauna, IV, 573, 1832 (*molva*).

Body elongate, covered with very small scales. Chin with a barbel; lower jaw included; bands of teeth on jaws and vomer; lower jaw with large canines which are arrow-shaped and movable; vomer with a curved series of canines mixed with small teeth, these mostly fixed; no teeth on palatines. Gill membranes broadly united. Two dorsal fins, both well developed; 1 anal fin; ventrals with several rays. Northern seas. (An old name of the salt-water ling.)

2923. MOLVA MOLVA (Linnæus).

Head 5; depth 7 or 8. D. 13 to 16-63 to 70; A. 57 to 66 (vertebræ 27 + 37 = 64). Upper jaw the longer, the maxillary reaching to below middle of orbit. Teeth cardiform in the jaws, with an inner row of rather widely separated and larger ones on mandible; a semicircular band on vomer, among which a few larger ones are interspersed. First dorsal inserted over the latter half of pectoral, its greatest height ⅔ that of body below it; pectoral about ½ as long as head; anal insertion in vertical over seventh or eighth ray of second dorsal. Barbel longer than eye, which is about equal to width of interorbital space. Scales small, covering head and fins. Color black gray, lighter on the sides and beneath; vertical fins edged with white; a dark blotch at the posterior end of the first dorsal, and a more distinct one on the end of the second dorsal. Arctic parts of the

Atlantic, south in deep water. This fish, the "ling" of Europe, is found from Spitzbergen to the Gulf of Gascony, where specimens have been taken very exceptionally at Arcachon and San Juan de Luz. It is very rare, however, south of the British Channel, and most abundant along the coast of northern Europe, especially in the German Ocean and off Norway. It is rare about Iceland, Greenland, and the Faroe Islands, and has never been found in the Baltic. It is said to have been found in the deep water off Newfoundland, but we have been unable to find the specific record. Collett states that on the Norwegian coast young examples rarely occur in less depth than 100 fathoms, and according to Lilljeborg the largest are caught in from 80 to 150 fathoms. (Goode & Bean.) (*molva*, an ancient name.)

Gadus molva, LINNÆUS, Syst. Nat., Ed. x, 254, 1858, seas of Europe; after *Gadus dorso dipterygia*, ARTEDI.

Molva vulgaris, FLEMING, British Animals, 192, 1828; GÜNTHER, Cat., IV, 361, 1862; GOODE & BEAN, Oceanic Ichthyology, 364, fig. 317, 1896.

Gadus raptor, NILSSON, Prodromus, 46, Sweden.

Molva linnæi, MALM, Götheborgs och Bohusläns Fauna, 491, 1877.

990. UROPHYCIS, Gill.

(CODLINGS.)

Phycis, BLOCH & SCHNEIDER, Syst. Ichth., 56, 1801 (*tinca* = *blennioides*); not *Phycis*, Fabricius, 1798, a genus of Lepidoptera.

Phycis, RAFINESQUE, Amer. Monthly Mag. 1818, 243 (*marginata*).

Urophycis, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 240 (*regius*).

Emphycus, JORDAN & EVERMANN, new subgenus (*tenuis*).

Body rather elongate; head subconic; mouth rather large, the maxillary reaching to below eye; lower jaw included; chin with a small barbel; jaws and vomer with broad bands of subequal, pointed teeth; palatines toothless. Dorsal fins 2, the first sometimes produced at tip; second dorsal long, similar to the anal; ventrals wide apart, filamentous, each of 3 slender rays, closely jointed, appearing like one beftd filament. Gill membranes somewhat connected, narrowly joined to the isthmus. (*οὐρά*, tail; *Phycis*.)

UROPHYCIS:

a. First dorsal fin not elevated, none of its rays filamentous.

b. Scales moderate, 90 to 95 in a longitudinal series.

c. Dorsal rays 8–43; anal 45; sides with some pale spots. REGIUS, 2924.

cc. Dorsal rays 10–66; anal rays 57; barbel minute. CIRRATUS, 2925.

bb. Scales small, 120 to 155 in a longitudinal series.

d. Dorsal rays 13–57; anal about 50; scales 120; sides with some pale spots. FLORIDANUS, 2926.

dd. Dorsal rays 10–62; anal about 53; scales 155. EARLLI, 2927.

EMPHYCUS (*ἔν-φύκος*, in the seaweed):

aa. First dorsal fin elevated, 1 or more of its rays filamentous.

e. Scales about 140; dorsal rays 9–57; anal 48; ventrals reaching vent. TENIUS, 2928.

ee. Scales about 110; dorsal rays 9–57; anal 50; ventrals reaching beyond vent, not longer than head. CHUSS, 2929.

eee. Scales about 90; dorsal rays 9–56; anal 56; second dorsal filamentous; ventrals very long, nearly 3 times length of head. CHESTERI, 2930.

Subgenus UROPHYCIS.

2924. UROPHYCIS REGIUS (Walbaum).

Head $4\frac{1}{3}$ in body; depth 5. D. 8–43; A. 40; scales about 90; eye $4\frac{1}{3}$ in head; maxillary 2; pectoral $1\frac{2}{3}$; caudal $1\frac{2}{3}$. Body rather elongate, compressed; mouth large, the maxillary reaching slightly past posterior margin of eye; lower jaw included; cardiform teeth on jaws and vomer; interorbital flattish, about equal in width to the diameter of eye; gill rakers short, 3 + 12 in number; origin of dorsal over base of pectorals; pectorals slender, barely reaching to front of anal; ventrals filamentous, composed of 2 rays each with the inner ray the larger, inserted in front of base of pectoral in distance equal to $1\frac{1}{2}$ diameter of eye, their ends reaching beyond front of anal; front of anal nearer snout than base of caudal, by nearly a head's length; caudal subtruncate. Pale brownisht inged with yellowish, the lateral line dark brown, interrupted by white spots; inside of mouth white; first dorsal largely black, this color surrounded by white; second dorsal olivaceous, with irregular round dark spots; caudal, anal, and pectorals dusky; ventrals and lower edge of pectorals white; 2 vertical series of round dark spots on the sides of the head. North Atlantic, south to Cape Fear; ranging from shallow water to a depth of 167 fathoms. Here described from a specimen, 8 inches in length, from Charleston, South Carolina. The species is said to exhibit electric powers in life. (*regius*, royal.)

Blennius, sp., SCHÖPF, Schrift. Naturf. Freunde, Berlin, VIII, 1780, 142, New York.
Blennius regius, WALBAUM, Artedi, Pisc, III, 186, 1792; after SCHÖPF.
Enchelyopus regalis, BLOCH & SCHNEIDER, Syst. Ichth., 53, 1801, after SCHÖPF.
Gadus punctatus, MITCHILL, Trans. Lit. and Phil. Soc. N. Y., I, 1815, 372, New York.
Urophycis regius, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 240.
Phycis regius, GOODE & BEAN, Oceanic Ichthyology, 357, 1896.
Phycis regalis, GÜNTHER, Cat., IV, 355, 1862.
Phycis punctatus, DE KAY, N. Y. Fauna: Fishes, 292, 1842.

2925. UROPHYCIS CIRRATUS, Goode & Bean.

Head 4; depth 5. D. 10–66; A. 57; scales 6-93-20. Body moderately stout; eye large, about 4 in head; interorbital space 2 in eye. Maxillary not reaching posterior margin of orbit in large specimens, but in smaller ones it extends fully to that vertical; mandible extending far beyond posterior margin of eye, its length about equal to postorbital part of head; barbel minute in all examples examined, its length usually about $\frac{1}{5}$ that of eye. Teeth in villiform bands in both jaws, the intermaxillary bands being wider than those of mandible; vomerines in a narrow villiform band. Gill rakers 2 + 12, the largest club-shaped at end, the longest 4 in eye. Gill membranes attached to isthmus, but with a narrow, free posterior border. Length of pectorals about $\frac{1}{2}$ distance from ventral to anal origin, reaching to about the twenty-sixth row of scales; ventral reaching in some specimens slightly beyond origin of anal; in 1 individual almost to middle of anal fin; none of the dorsal rays filamentous, the longest from $2\frac{1}{2}$ to 3 times in head; base of first dorsal about equal to length of eye in most specimens; in smaller examples somewhat greater, about 3 in head;

vent under sixteenth ray of second dorsal. Color light brown; lower parts minutely dotted; dorsals with narrow dark margins; caudal with a broad dark margin; anal with a narrow dark margin in its posterior third; roof of mouth and interior of gill cavity dark brown. Deep water of the Gulf of Mexico. (Goode & Bean.) (*cirratus*, bearing cirri.)

Phycis cirratus, GOODE & BEAN, Oceanic Ichthyology, 358, 1896, Gulf of Mexico at Lat. 29° 03′ 15″ N., Long. 88° 16′ W. (Type, No. 39059. Coll. *Albatross*.)

2926. UROPHYCIS FLORIDANUS, Bean & Dresel.

Head 4 in body; depth 6. D. 13–57; A. 49; scales about 120; eye 6 in head; maxillary 2; height of first dorsal 2½; middle caudal rays 2. Body rather elongate, compressed, head subconic; mouth large, the maxillary reaching to below posterior margin of orbit; upper jaw and snout somewhat projecting beyond lower; small cardiform teeth, in narrow bands on jaws and vomer; barbel very slender, small; interorbital space wide, slightly convex, nearly twice as wide as eye; gill rakers small, slender, 2+11 in number. Origin of dorsal a little behind the vertical from base of pectoral; first dorsal high, slightly falcate; second dorsal a little higher in its anterior end, higher than anal; origin of anal about midway between tip of snout and base of caudal; pectoral slender, reaching an eye's diameter beyond front of second dorsal; ventrals inserted twice diameter of eye in front of pectorals, 2-rayed, the inner ray the longer, not reaching to vent in larger examples, reaching to front of anal in small ones; caudal long and rounded. Color in spirits, reddish brown, light below, a small black spot above eye, a vertical series of 3 or 4 behind eye, and 2 on opercle, these spots less than ½ pupil, distinct and clear cut; a dark streak from preorbital across cheek to edge of opercle, lateral line black, interrupted at short intervals by white spots; fins dusky, with the exception of pectorals and ventrals, dark toward the ends of the rays. Gulf of Mexico, in rather shallow water, coming to shore in abundance about Pensacola in cold weather. Here described from a specimen, 7½ inches in length, from Pensacola, Florida. (*floridanus*, from Florida.)

Phycis floridanus, BEAN & DRESEL, Proc. Biol. Soc. Wash. 1884, 100, Pensacola, Florida (Coll. Silas Stearns); JORDAN, Cat. Fish. N. A., 129, 1885.

2927. UROPHYCIS EARLLI, Bean.

Head 3⅓ in body; depth 5. D. 10–60; A. 53; scales 155; eye 6 in head; maxillary 2. Body moderately elongate, not much compressed anteriorly; mouth large, the maxillary reaching to below or very slightly past posterior margin of eye; snout and upper jaw projecting beyond lower jaw; teeth strong, cardiform in a narrow band on vomer and lower jaw, in a rather wide band in upper; interorbital wide, convex, about 1½ times eye; gill rakers short and blunt, about 2+9. Origin of dorsal slightly behind the vertical from base of pectoral; origin of anal about midway between snout and base of caudal; ventrals 2-rayed, the inner a little the longer, not reaching to vent. Color brown, with some light spots on the second dorsal fin and on the sides; anal and both dorsals margined with brown.

Atlantic coast of United States, southward in water of moderate depth; not common. Here described from a specimen, 17 inches in length, from Charleston, South Carolina. (Named for R. Edward Earll, then assistant to the United States Fish Commission.)

Phycis earlli, BEAN, Proc. U. S. Nat. Mus., III, 1880, 69, Charleston, S. C. (Coll. R. E. Earll. Type, Nos. 25207, 25208, and 25209); JORDAN & GILBERT, Synopsis, 798, 1883.

Subgenus **EMPHYCUS**, Jordan & Evermann.

2928. UROPHYCIS TENUIS (Mitchill).

(CODLING; WHITE HAKE; SQUIRREL-HAKE.)

Head 4¼; depth 5½. D. 9-57; A. 48; scales 138. Snout longer than eye, narrower and more pointed than in *P. chuss*. Eye large, usually wider than interorbital space; maxillary reaching beyond pupil. Filamentous dorsal ray about ⅔ length of head; ventral fins about reaching vent. Scales very small. Brownish, lighter and yellowish below; fins very dark. Banks of Newfoundland to Cape Hatteras; abundant northward in rather deep water, reaching a depth of 304 fathoms. The species resembles *Phycis chuss*, differing chiefly in the smaller scales. (*tenuis*, slender.)

Gadus tenuis, MITCHILL, Trans. Lit. and Phil. Soc. N. Y. 1815, 372, New York.
Phycis dekayi, KAUP, Archiv Natur. 1858, 89, North America.
?Phycis rostratus, GÜNTHER, Cat. Fish. Brit. Mus., IV, 353, 1862, no locality; D. 9–59 to 62; A. 49 to 50; scales ca. 150; ventrals immaculate, reaching front of anal.
Phycis tenuis, DE KAY, N. Y. Fauna: Fishes, 293, 1842; GILL, Proc. Ac. Nat. Sci. Phila, 1863, 238; JORDAN & GILBERT, Synopsis, 799, 1883; GOODE & BEAN, Oceanic Ichthyology, 359, fig. 312, 1896.

2929. UROPHYCIS CHUSS (Walbaum).

(CODLING; SQUIRREL-HAKE.)

Head 4½; depth 5. D. 9-57; A. 50; scales 110. Body rather slender; head depressed; eye large, about equal to interorbital width; maxillary reaching posterior margin of pupil; filamentous dorsal ray about ⅔ length of body, when perfect; pectorals ⅔ length of head; ventral fins extending beyond the vent; scales comparatively large. Brownish above, sides lighter and tinged with yellowish; thickly punctulate with darker; below pale; inside of mouth white; vertical fins somewhat dusky; anal fin margined with pale; lateral line not dark. Atlantic coast, from Gulf of St. Lawrence to Virginia; common northward; reaching a depth of 300 fathoms. (*chuss*, a vernacular name now obsolete, apparently derived from cusk.)

Chuss, SCHÖPF, Schrift. Naturf. Freunde, Berlin, VIII, 1780, 143, New York.
Blennius chuss, WALBAUM, Artedi Pisc., 186, 1792; after SCHÖPF.
Enchelyopus americanus, BLOCH & SCHNEIDER, Syst. Ichth., 53, 1801; after SCHÖPF.
Gadus longipes, MITCHILL, Trans. Lit. and Phil. Soc. N. Y., 1, 372, pl. 1, fig. 4, 1815, New York.
Phycis marginatus, RAFINESQUE, Amer. Month. Mag., Jan., 1818, 205, Point Judith, Rhode Island. D. 10-60; A. 40; ventral reaching anal; tail black-edged.
Phycis americanus, STORER, Rept. Fish. Mass., 138, 1839; GÜNTHER, Cat., IV, 353, 1862.
Phycis chuss, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 237; JORDAN & GILBERT, Synopsis, 799, 1883; GOODE & BEAN, Oceanic Ichthyology, 359, fig. 311, 1896.

2930. UROPHYCIS CHESTERI, Goode & Bean.

Head $4\frac{1}{3}$; depth 5; orbit $3\frac{1}{3}$ in head; maxillary 2; barbel about 3 in orbit. D. 9 or 10–55 to 57; A. 56; C. 5, 18 to 21, 5; P. 17 or 18; V. 3; scales 7–90 or 91–28. Vent situated under the twelfth ray of second dorsal, and equidistant from tip of snout and end of second dorsal; distance of dorsal fin from snout equal to twice length of mandible; third ray of first dorsal extremely elongate, extending to a point (thirty-third ray of second dorsal) $\frac{3}{5}$ of distance from snout to tip of caudal, its length more than twice that of head, and more than 4 times as long as the rays immediately preceding and following it; anal fin inserted immediately behind vent, its distance from root of ventrals equal to that of dorsal from snout; as in other species of the genus, ventral of 3 rays, the first 2 much prolonged, the first contained 3 times in length of body, the second almost 3 times as long as head, reaching to fortieth anal ray or $\frac{3}{4}$ of distance from snout to tip of caudal, the third shorter than diameter of orbit; pectoral 4 times as long as operculum. Scales large and thin, easily wrinkling with the folding of the thick, loose skin, particularly in the median line of sides of body. Lateral line much broken on posterior half of body. (Goode & Bean.) Atlantic coast of United States, in 100 to 500 fathoms, with *Macrourus bairdi*, the most abundant fish on the continental slope, swarming everywhere below the 100-fathom line. (Named for Capt. Hubbard C. Chester.)

Phycis chesteri, Goode & Bean, Proc. U. S. Nat. Mus. 1878, 256, off Cape Ann, in 140 fathoms (Coll. Captain Chester); Jordan & Gilbert, Synopsis, 800, 1883; Goode & Bean, Oceanic Ichthyology, 360, fig. 313, 1896.

991. LÆMONEMA, Günther.

Læmonema, Günther, Cat. Fish. Brit. Mus., IV, 356, 1862 (*yarrellii*).

Body of moderate length, covered with small scales; fins naked. A separate caudal; 2 dorsal fins and 1 anal, the anterior dorsal composed of 5 rays; ventrals reduced to a single long ray, bifid at its end. Bands of villiform teeth in the jaws; a small group of vomerine teeth; none on the palatine bones. Chin with a barbel. Branchiostegals 7. Deep sea. (λαιμός, throat; νῆμα, thread.)

a. Scales 13–140–31; barbel $\frac{1}{3}$ eye; dorsal and anal with narrow black edgings. BARBATULUM, 2931.

aa. Scales 16–160–38; barbel $\frac{2}{3}$ eye; a large, triangular, black blotch on tail and adjacent parts of vertical fins. MELANURUM, 2932.

2931. LÆMONEMA BARBATULUM, Goode & Bean.

Head $4\frac{2}{3}$; depth $4\frac{1}{2}$; orbit 3 in head; upper jaw more than 2; barbel about 2 in eye. D. 5–63; A. 59; P. 19; V. 2; scales 13–140–31. Vent situated under sixth or seventh ray of second dorsal. Distance of first dorsal from snout 4 in body; base of first dorsal $\frac{1}{2}$ as long as middle caudal rays, that of second slightly more than 3 times length of head; first dorsal composed of 5 rays, the first of which is elongate, 3 times as long as middle caudal rays, extending to base of twenty-fourth ray of second dorsal; anal fin inserted at a distance from tip of snout equal to twice length of head,

its distance from insertion of ventrals being equal to length of head; length of ventrals equal to that of pectorals, their tip not extending to vent. Scales small, very thin, deciduous, crowded anteriorly; lateral line not well defined on posterior part of body. Color similar to that of the various species of *Phycis;* the dorsal and anal fins with narrow black margins. The length of the first dorsal ray is very variable, being shorter in younger individuals. This species differs from *L. yarrellii* by its much smaller scales, and from *L. robustum* by the greater number of rays in the dorsal and anal fins, and its much shorter ventrals. (Goode & Bean.) Gulf Stream, reaching a depth of 312 fathoms. (*barbatulus,* having small barbels.)

Læmonema barbatula, Goode & Bean, Bull. Mus. Com. Zool., x, 204, 1883, Gulf Stream, Lat. 32° 43′ N., Long. 77° 20′ W., in 230 fathoms, and Lat. 28° 35′ N., Long. 73° 13′ W.; Goode & Bean, Oceanic Ichthyology, 362, figs. 315 and 315A, 1896.

2932. LÆMONEMA MELANURUM, Goode & Bean.

Head about $4\frac{2}{3}$; depth $4\frac{2}{3}$; eye 3 in head; snout 4; interorbital width 6. D. 6–57; A. 55; P. 25; V. 2; Br. 7; scales 16–160–38. Maxillary extending to below middle of eye; intermaxillary nearly $\frac{1}{2}$ length of head; mandible slightly more than twice length of snout. Teeth in intermaxillary and mandible in villiform bands; vomerine teeth in a small circular patch on middle of head of bone. Barbel about as long as snout. Distance of first dorsal from tip of snout about 4 in snout; length of first ray of dorsal equaling that of head without snout; last ray of dorsal scarcely more than $\frac{1}{4}$ as long as first; ventral consisting of a single bifid ray, its distance from tip of snout equal to length of head, its length nearly equal to that of dorsal or the pectoral when extended, not reaching vent by a distance equal to length of snout; pectoral equaling that of longest dorsal ray, and also equaling head without snout; second dorsal higher anteriorly, and posteriorly much higher than in middle; longest anterior ray $\frac{1}{2}$ length of ventral; longest posterior ray $\frac{1}{2}$ length of head. Vent under eighth ray of second dorsal. Gill rakers 5 + 15, the longest $\frac{1}{4}$ as long as snout. Color very light brown, the dorsals and anal with a narrow dark margin; a conspicuous, large, triangular, dark blotch on last rays of dorsal and anal, and a dark blotch occupying almost the whole of caudal, leaving a margin of whitish around it. (Goode & Bean.) Caribbean Sea, north to New York; reaching a depth of 1,467 fathoms. (*μέλας,* black; *οὐρά,* tail.)

Læmonema melanurum, Goode & Bean, Oceanic Ichthyology, 363, fig. 316, 1896, Gulf Stream, Lat. 30° 44′ N., Long. 79° 26′ W., in 440 fathoms. (Type, No. 38270. Coll. *Albatross.*)

992. GAIDROPSARUS, Rafinesque.

(Three-bearded Rocklings.)

Gaidropsarus, Rafinesque, Indice d'Ittiol. Siciliana, 1810 (*mustellaris = mediterraneus*); description from a rough figure of Rondelet.

Les Mustèles, Cuvier, Règne Anim., Ed. I, vol. 2, 215, 1817 (*tricirrhatus = mediterraneus*).

?*Mustela,* Oken, Isis. 1817 (for *les Mustèles;* not *Mustela,* a genus of mammals).

Onos, Risso, Hist. Eur. Mérid., III, 214, 1826 (*mustella* = *mediterraneus*).
Mustela, Stark, Elem. Nat. Hist., I, 425, 1828 (after *les Mustèles*).
Motella, Cuvier, Règne Anim. Ed. 2, vol. II, 334, 1829 (*vulgaris* = *tricirratus*).
Onus, Günther, corrected spelling.

Body rather elongate, covered with minute scales; head not compressed, the upper jaw the longer; snout with 2 conspicuous barbels, the chin with 1; teeth on jaws and vomer in bands, palatines toothless; dorsals 2, the anterior of a single long ray followed by a series of short fringe-like rays concealed in a groove; second dorsal and anal long, similar to each other; caudal rounded or lanceolate; ventral rays 5 to 7. Small fishes of the northern seas, descending to deep water. We here regard the 5-bearded Rocklings (*Ciliata*, Couch, 1832) = *Couchia*, Thompson, 1856 = *Molvella*, Kaup, 1858, as a distinct genus, distinguished by the 5 barbels at the tip of the snout. (*γαϊδραψάρα*, a modern Greek name used by Rondelet for a species of this group.)

The name *γαϊδροψάρον* is now applied in Athens to the Pollack-like fish, ***Micromesistius poutassou*** (Risso).

According to Prof. Horace A. Hoffman "the name *γαϊδουροψάρον* is modern, meaning donkey fish. *Γαΐδουρος* = *γάδαρος* = ass, donkey. The ancients called a certain fish *ὄνος*, ass. Dorio, in Athenæus, VII, 99, says some persons call the *ὄνος* (i. e., the fish *ὄνος*) *γάδος*. Epicharmus, in his Marriage of Hebe, says: 'Wide-gaping *χάνναι* and monstrous-bellied *ὄνοι*.' (See Aristotle 599b 33, 601a 1, 620b 29, frag. 307, 1530a.) According to Aristotle the *ὄνος* has a mouth opening wide (literally, breaking back), like the *γαλεοί*. It leads a solitary life, is the only fish which has its heart in its belly, has stones in its brain like millstones in form, and is the only fish which lies torpid in the warmest days under the reign of the dog star, Sirius, the other fishes going into this torpid state in the wintriest days. The *ὄνος*, *βάτος*, *ψῆττα*, and *ῥίνη* bury themselves in the sand, and after they make themselves invisible they wave the things in their mouths which fishermen call little rods or little wands (*ῥαβδία*). (Hoffman & Jordan, Fishes of Athens, Proc. Ac. Nat. Sci. Phila. 1887, 146.)

a. First ray of first dorsal long, as long as head; head small, 5¾ in length; teeth rather feeble, uniform. D. 59; A. 45; P. 25. Color uniform brick red. ENSIS, 2933.

aa. First ray of first dorsal short, about as long as snout.

b. Pectoral rays 22 to 24; upper jaw without cirri or rudimentary barbels along the premaxillary; maxillary reaching posterior border of eye; head 5⅛ in length. D. 56; A. 45. Color reddish. ARGENTATUS, 2934.

bb. Pectoral rays 16; upper jaw with short cirri or barbels along the premaxillary; maxillary reaching far beyond eye. D. 50; A. 42. Head 4 in length. Color brownish. SEPTENTRIONALIS, 2935.

2933. GAIDROPSARUS ENSIS (Reinhardt).

Head 5¾; depth 4½. D. 59; A. 44 to 46; P. 22 to 27; V. 8. Body unusually deep, being greatest at the vent; head small; eye rather large, nearly as long as snout, equaling interorbital area, and in anterior half of head; posterior margin of orbit nearly equidistant between tip of snout and posterior margin of operculum. Mouth normal; supramaxillary end-

ing under posterior margin of pupil. Teeth in a narrow band in each jaw, some of those at least in outer row of upper jaw slightly enlarged and brownish colored; teeth of vomer forming a short curved band in 2 rows. Nasal barbel about equaling diameter of eye. Chin barbel small and not much exceeding ½ diameter of eye. Foremost ray of first dorsal springing from back above opercular margin; second dorsal fin low in front, but rising rapidly to seventh or eighth ray, behind which it is nearly uniform for a long distance, and the highest at posterior portion; anal fin much lower than second dorsal; caudal slightly emarginate, almost truncate behind, its median rays about ⅔ as long as head; pectorals nearly ¾ as long as head, produced toward the upper angles, the third ray being longest; ventral fins with their bases mostly in advance of pectorals, the longest ray filamentous and nearly equaling pectoral. Lateral line obsolescent. (Goode & Bean.) Atlantic coast of North America, from Greenland to Cape Hatteras; in deep waters, reaching a depth in the Gulf Stream of 1,081 fathoms. (*ensis*, sword.)

Motella ensis, REINHARDT, Dansk. Vidensk. Selsk. Afhandl., VII, 15, 1838, Greenland.
Onos rufus, GILL, Proc. U. S. Nat. Mus. 1883, 259, Gulf Stream; GILL, Proc. Ac. Nat. Sci. Phila. 1884, 172; JORDAN, Cat. Fish. N. A., 128, 1885.
Onos ensis, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 241; GILL, Cat. Fish. E. Coast U. S., 18, 1873; JORDAN & GILBERT, Synopsis, 797, 1883; JORDAN, Cat. Fish. N. A., 128, 1885; GOODE & BEAN, Oceanic Ichthyology, 381, fig. 327, 1896.

2934. GAIDROPSARUS ARGENTATUS (Reinhardt).

Head 5; depth 5¼. D. 54 to 59; A. 45 or 46. Body elongate; head small; teeth in several rows, 1 row more enlarged than the others; maxillary reaching posterior border of eye; eye large, 5½ in head; interorbital space scarcely exceeding the eye. Lateral line with about 27 enlarged pores along its entire length. First ray of first dorsal short, little longer than snout; vent near middle of length. Reddish gray, changing to bluish on the head and abdomen; tips of dorsal, anal, and caudal red, also the barbels and first ray of first dorsal; cavity of mouth pale. Coasts of Greenland (Collett), south to Faroë and Bear Islands; not seen by us. There can be no doubt that *Motella argentata* is the young of the species later called *Motella reinhardti*. (Eu.) (*argentatus*, silvered.)

Motella argentata, REINHARDT, Dansk. Vidensk. Selsk. Afh., VII, 128, 1838; Greenland; young.
Motella reinhardti, KRÖYER MS., 1852; COLLETT, Forh. Vid. Selsk. Chr., No. 14, 83, 1878, Greenland.
Couchia argentata, GÜNTHER, Cat., IV, 365, 1862.
Ciliata argentata, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 241; GILL, Cat. Fish. E. Coast U. S., 18, 1873.
Onos reinhardti, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 241; GILL, Cat. Fish. E. Coast U. S., 18, 1873; COLLETT, Norske Nord-Havs Exp., 131, 1880; JORDAN & GILBERT, Synopsis, 797, 1883; JORDAN, Cat. Fish. N. A., 128, 1885; GOODE & BEAN, Oceanic Ichthyology, 383, 1896.

2935. GAIDROPSARUS SEPTENTRIONALIS (Collett).

Head 4; depth 5⅔. D. 50; A. 42; P. 16. Three barbels, 2 at the nostrils, 1 at the chin, besides a row of about 8 shorter rudimentary barbels along the edge of the upper lip; eye small, ⅓ length of snout; cleft of mouth

extending far beyond eye, its length nearly equal to that of postorbital part of head; teeth rather small, unequal; outer teeth of upper jaw and some of the inner teeth of lower enlarged; first ray of first dorsal short, about as long as snout; vent midway between tip of snout and last anal ray; lateral line with about 20 large pores. Grayish brown, paler below; cavity of mouth white. Coast of Norway; 1 specimen known from Greenland. (Collett.) (Eu.) (*septentrionalis*, northern.)

Motella septentrionalis, COLLETT, Ann. Mag. Nat. Hist. 1874, 15, 82, Lofoten, Norway.
Onos septentrionalis, COLLETT, Norske Nord-Havs Exped., 139, 1880; JORDAN, Cat. Fish. N. A., 128, 1885.

993. ENCHELYOPUS, Bloch & Schneider.

(FOUR-BEARDED ROCKLINGS)

Enchelyopus, BLOCH & SCHNEIDER, Syst. Ichth., 50, 1801 (*cimbrius;* the first species mentioned and the one left as type after elimination of the genera, defined prior to *Rhinonemus*).
Rhinonēmus, GILL, Proc. Ac. Nat. Sci. Phila. 1883, 241 (*cimbrius*).

Barbels 4, 1 at each nostril, 1 at tip of snout, and 1 at the chin; head high and compressed anteriorly; teeth in narrow bands, some of them enlarged; otherwise essentially as in *Gaidropsarus*. North Atlantic. (*ἐγχελυωπός*, resembling an eel; *"facie anguillaris."*)

2936. ENCHELYOPUS CIMBRIUS (Linnæus).

(FOUR-BEARDED ROCKLING.)

Head 5½; depth 9. D. 45 to 50; A. 41 or 42; V. 5. Body slender, tapering from the shoulders back; caudal peduncle narrow, 4 in head; snout moderate, blunt, rounded, not depressed, a little shorter than the eye; eye large, subcircular, 4 in head; interorbital space narrow, equal to vertical diameter of eye, 6 in head; teeth villiform, those in the upper jaw unequal, small, with about 8 enlarged in front, those of the lower jaw long and slender, of equal length, a few somewhat enlarged in front; maxillary reaching beyond posterior border of eye, a barbel at each nostril, 1 on tip of snout and 1 on chin, stitch-like; lateral line with about 35 enlarged pores along its entire length; first (free) ray of dorsal nearly as long as head; ventral ½ head; caudal acute. Light olivaceous (salmon-red); first dorsal ray and posterior end of dorsal and anal abruptly black, as is lower half of caudal; pectorals and ventrals pale; sides of head somewhat silvery; cavity of mouth dark bluish. North Atlantic, on both coasts, south in deep water to the Gulf Stream; common in Massachusetts Bay; our specimens from Woods Hole; the young ("mackerel midges") silvery, unlike the adult in appearance. (Eu.) (*cimbrius*, welsh.)

Gadus cimbrius, LINNÆUS, Syst. Nat., Ed. 12, I, 440, 1766, Atlantic Ocean; Scania (Coll. Dr. Strussenfelt).
Motella caudacuta, STORER, Proc. Bost. Soc. Nat. Hist., III, 1848, 5, Cape Cod, Provincetown, Mass. (Coll. Herman M. Smith); STORER, Amer. Ac. Sci., 411, 1867; STORER, Hist. Fish. Mass., 183, 1867.
Rhinonemus caudacuta, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 241; GOODE & BEAN, Amer. Journ. Sci. and Arts 1877, 476; JORDAN, Cat. Fish. N. A., 128, 1885.
Motella cimbria, NILSSON, Prod. Ich. Scand., 48, 1832; BELL, Can. Nat. and Geol., IV, 209, 1859.

Onos cimbrius, GOODE & BEAN, Proc. U. S. Nat. Mus. 1878, 349; GOODE & BEAN, Bull. Essex Inst., XI, 1879; JORDAN & GILBERT, Synopsis, 797, 1883; GOODE & BEAN, Bull. Mus. Comp. Zool., X, No. 5, 217, 1883.

Rhinonemus cimbrius, JORDAN, Cat. Fish N. A., 128, 1885; GOODE & BEAN, Oceanic Ichthyology, 384, fig. 328, 1896.

Enchelyopus cimbricus (misprint for *cimbrius*), BLOCH & SCHNEIDER, Syst. Ich., 50, pl. 9, 1801.

994. BROSME (Cuvier) Oken.

(CUSKS.)

Les Brosmes, CUVIER, Règne Animal, Ed. I, vol. 2, 216, 1817 (*brosme*).

Brosme, OKEN, Isis, 1817, 1182; after CUVIER.

Brosmius, CUVIER, Règne Animal Ed. 2, vol. II, 334, 1829 (*brosme*).

Body moderately elongate, covered with very small scales. Mouth rather large, with teeth in the jaws, vomer, and palatines, some of those on the vomer and palatines enlarged; chin with a barbel; branchiostegals 7. Dorsal fin single, continuous, not elevated, not notched; anal fin similar, but shorter; caudal fin rounded; ventral fin several-rayed. Northern seas. (From the Danish vernacular name, *brosme*.)

2937. BROSME BROSME (Müller).

(CUSK.)

D. 98; A. 71; P. 24; V. 5. Body cylindrical, posteriorly compressed; head flattened above. Mouth large, oblique, maxillary reaching beyond orbit; lower jaw included; several rows of sharp teeth on jaws, vomer, and palatines; barbel about 5 in head; interorbital greater than the diameter of eye. Origin of dorsal above anterior half of pectoral; pectoral round, $2\frac{1}{2}$ in head; caudal rounded behind. Brownish above, the sides yellowish, sometimes mottled with brown; young uniform dark slate color, or with transverse yellow bands; vertical fins bordered with blackish, and with a white edge. (Storer.) North Atlantic, south to Cape Cod and Denmark; rare southward on our coasts. (*brosme*, a Danish name.)

Gadus brosme, MÜLLER, Prodr. Zool. Dan., 41, 1776, Denmark; FABRICIUS Fauna Grœnlandica, 140, 1780.

Gadus lubb, EUPHRASEN, Vet. Akad. Handl. 1794, 223, tab. 8.

Gadus torsk, BONNATERRE, Encycl. Meth., 51, 1788, Söndmöre, Norway; after Strom.

Brosmius vulgaris, FLEMING, British Anim., 194, 1828.

Brosmius flavescens, LE SUEUR, Mém. Mus., V, 1819, 158, Banks of Newfoundland; chin with 2 barbels; lower jaw longest.

Brosmius flavescens, GÜNTHER, Cat., IV, 369, 1862; STORER, Hist. Fish. Mass., 368, 1867.

Enchelyopus brosme, BLOCH & SCHNEIDER, Syst. Ichth., 51, 1801.

Brosmius brosme, GÜNTHER Cat., IV, 369, 1862; JORDAN & GILBERT, Synopsis, 802, 1883; GOODE & BEAN, Oceanic Ichthyology, 385, fig. 329, 1896.

Blennius torsk, LACÉPÈDE, Hist. Nat. Poiss., II, 508, 1800.

Family CCXV. MACROURIDÆ.

(THE GRENADIERS.)

Body elongate, tapering into a very long compressed tail, which ends in a point; scales moderate, usually keeled or spinous, sometimes smooth. Suborbital bones enlarged, sometimes cavernous. Teeth villiform or cardi-

form, in bands, on the jaws only; tip of lower jaw with a barbel; premaxillary protractile. Dorsal fins 2, the first short and high, of stiff, spine-like branched rays; the second dorsal very long, usually of very low feeble rays, continued to the end of the tail; anal fin similar to the second dorsal, but usually much higher; no caudal fin; ventrals small, subjugular, each of about 8 rays. Branchiostegals 6 or 7. Lateral line present. Gills $3\frac{1}{2}$ or 4, a slit behind the fourth. Gill rakers small; gill membranes free or narrowly united to the isthmus, usually more or less connected; pseudobranchiæ wanting or rudimentary; pyloric cæca numerous; air bladder present. Genera 18; species about 50, chiefly of the northern seas, all in deep water. They differ from the codfishes chiefly in the elongate and degenerate condition of the posterior part of the body. Dr. Gill succinctly defines the group as "Gadoidea with an elongated tail tapering backward and destitute of a caudal fin, postpectoral anus, enlarged suborbital bones, inferior mouth, subbrachial ventrals, a distinct anterior dorsal, and a long second dorsal and anal converging on end of tail." We here follow Goode & Bean in the general arrangement of the genera of *Macrouridæ*. Some of these can, however, be only provisionally adopted, as the characters of dentition, form of mouth, and character of the second dorsal spine or ray, are subject to much intergradation. These characters seem much more distinct on paper than they are in fact. Still, most of the genera here adopted will ultimately prove valid. (*Macruridæ*, Günther, Cat., IV, 390–398, 1862.)

a. First branchial arch free, without fold of membrane across it; mouth large; second dorsal well developed, higher than the anal.

BATHYGADINÆ:

b. Gills $3\frac{1}{2}$; snout short and blunt, the jaws even in front; teeth in villiform bands, sometimes obsolete; bones of head soft and cavernous; scales smooth; first dorsal low, its spine not produced. BATHYGADUS, 995.

TRACHYRINCHINÆ:

bb. Gills 4; snout rather long.

c. Teeth in upper jaw in 2 series, the outer enlarged, those in lower jaw in 1 series; mouth subterminal; barbel obsolete; nape without scaleless fossæ; vomer with teeth; bones of head soft and cavernous; tail very long, flagelliform; anal fin with an elevated anterior lobe. STEINDACHNERIA, 996.

cc. Teeth in both jaws in villiform bands; barbel developed; mouth inferior; a naked fossa on each side of nape; a row of armed scales along base of dorsal anteriorly; opercle very small; anal not elevated in front. TRACHYRINCUS, 997.

MACROURINÆ:

aa. First branchial arch with a fold of membrane across its terminal portion; gills 4, a slit behind the fourth; barbel well developed.

d. Teeth not all in villiform bands, those of lower jaw in 1 series; mouth rather large, with more or less of lateral cleft.

e. Upper jaw without villiform band behind the enlarged anterior teeth, the inner teeth, if present chiefly uniserial, not in villiform bands.

f. Dorsal fins widely separated, the interspace greater than base of first.

g. First dorsal with the spine not serrate, its insertion over pectoral or nearly so; pectoral placed high, opposite upper angle of gill cleft; scales small, bristly; bones of head cavernous; ventrals short and weak. MALACOCEPHALUS, 998.

gg. First dorsal with its spine more or less strongly serrate; pectoral inserted below upper angle of gill cleft.

h. Scales nearly smooth, with weak ridges which are not spinigerous. MOSELEYA, 999.

hh. Scales rough, with strong ridges. NEMATONURUS, 1000.

ff. Dorsal fins near together, the interspace less than base of first; scales rough.

i. Dorsal spine weak, unarmed or very nearly so; pectorals moderate. ALBATROSSIA, 1001.

ii. Dorsal spine very strongly serrate; pectorals very long. BOGOSLOVIUS, 1002.

ee. Upper jaw with a distinct villiform band behind the outer series of enlarged teeth; dorsal spine serrate; dorsal fins not widely separated. CHALINURA, 1003.

dd. Teeth in villiform bands above and below, the outer scarcely enlarged and not separated from the rest; the lower band sometimes becoming a single series laterally; scales rough.

j. Mouth wide, with considerable lateral cleft.

k. Dorsal spine finely barbed; skull rather firm; dorsals moderately separated. CORYPHÆNOIDES, 1004.

kk. Dorsal spine entirely smooth; bones of skull very thin and papery; dorsals well separated. HYMENOCEPHALUS, 1005.

jj. Mouth inferior, small, with little lateral cleft; a more or less distinct ridge across the suborbital region.

l. Scales spinous, very rough.

m. Scales distinct, regularly imbricated.

n. Long dorsal spine serrate in front; mouth subinferior, below the short snout. MACROURUS, 1006.

nn. Long dorsal spine smooth; mouth wholly inferior, below the long sturgeon-like snout. CŒLORHYNCHUS, 1007.

mm. Scales indistinct, scarcely imbricated; the whole body rough-villous; dorsal spine smooth. TRACHONURUS, 1008.

ll. Scales all thin and smooth, dorsal fin slightly serrulate. LIONURUS, 1009.

995. BATHYGADUS, Günther.

Bathygadus, GÜNTHER, Ann. Mag. Nat. Hist. 1878, 23 (*cottoides*).

Head large, fleshy, without prominent ridges, spiny armature or external depressions; nape elevated, hump-like. Snout broad, obtuse, not produced; mouth terminal very large, with small villiform teeth or none; suborbital ridge very low, not joined to the angle of the preoperculum. Maxillary entirely received within a groove under the prefrontal and suborbital bones, its tips narrowed and blade-like; premaxillaries protractile downward, separated anteriorly, rib-shaped, compressed vertically, very broad and without true teeth; provided posteriorly with a short flange, which is received under the maxillary; mandible received within the intermaxillary bones, without true teeth, but with minute asperities, similar to those in the upper jaw; vomer and palatines toothless. Barbel sometimes present. No pseudobranchiæ. Gill rakers numerous, moderate, lanceolate, with minute denticulations along their inner edge. Branchiostegal membrane free from the isthmus, deeply cleft. Branchiostegals 7, very stiff. Gill opening very wide; gills $3\frac{1}{2}$; anterior gill arch free. Operculum with a blunt, spine-like prominence at its angle. Ventrals below the pectorals, many-rayed, the anterior rays produced; dorsal consisting for the most part of branched rays, higher than the anal, the

first dorsal low, without differentiated spine. Scales cycloid, unarmed; lateral line strongly arched over the pectoral. Deep seas. This genus differs from *Macrourus* and its allies in the structure of both the first and last gill arches. It is perhaps the most primitive of the family and as such is nearest allied to the *Gadidæ*. (*βαθύς*, deep; *Gadus*, codfish.)

a. Pectoral and ventral fins moderate, not much, if any, longer than head.
b. Jaws without teeth; pectoral fin broad, of 25 rays; depth 5⅔ in length.
ARCUATUS, 2938.
bb. Jaws with small teeth; pectoral fin narrow; depth 6 to 6½ in length.
c. Eye moderate, 5 in head; pectoral rays 14, the fin ½ as long as head.
FAVOSUS, 2939.
cc. Eye very large, 2¾ in head; pectoral as long as head without snout.
MACROPS, 2940.
aa. Pectoral and ventral fins much produced, much longer than head, each reaching about halfway to tip of caudal; depth 7½ in length; pectoral rays 13.
LONGIFILIS, 2941.

2938. BATHYGADUS ARCUATUS, Goode & Bean.

Head 5 in total length; depth 5⅔; eye 4½ in head; snout 4½. D. II, 9 or 10–135; A. 120; P. 25; V. 8; scales 8–140–13 or 14 (counting backward from vent to lateral line), 22 counting forward. Body shaped much as in *Chalinura simula*, but the nape still more convex; back gibbous, the dorsal outline rising rapidly from interorbital region to origin of first dorsal, whence it descends gradually to end of tail. Scales moderate, cycloid, subovate, without armature, those of abdominal region and those above pectorals the largest; lateral line strongly arched over the pectorals, length of the arched portion contained about 3¼ times in straight portion, greatest height of arch about ¼ its chord; scales covering all parts of head except jaws and chin. Interorbital area flat, its width 6 in head; postorbital portion of head about 2½ times diameter of eye; operculum terminating in a flat obtuse spine, its length, including the flap, about equal to diameter of eye; preoperculum entire, with a prominent ridge in advance of its posterior edge; snout very broad, obtuse, the intermaxillaries extending beyond it, its width at nostrils equal to about twice length of eye; posterior extremities of intermaxillary processes elevated, producing a decided hump upon top of snout; ridge formed by prefrontal and suborbital bones terminating very slightly behind posterior margin of orbit, and not connected with angle of preoperculum. Nostrils immediately in front of lower part of eye, not tubular, the anterior one very small, porelike, only about ⅓ as large as posterior one; distance of anterior nostril from tip of snout about ⅔ length of eye. Length of barbel 6⅔ in length of body, and equal to length of head without snout, more than 3 times as long as eye. No true teeth, the intermaxillaries and mandible being broad plates, covered with minute asperities; a naked space at the symphysis of intermaxillaries; distance of first dorsal from snout nearly 3⅓ times length of its base, the first spine minute, the second (in the type) somewhat mutilated, its length nearly 3 in length of head, not stouter than the branched rays, and entirely smooth; second dorsal fin separated from first by a very short interspace, equal to about ⅓ of length of eye, its rays long, subequal, the first slightly the longest, its length equal to that of base of

first dorsal; anal much lower than dorsal, the longest rays being in front, its third ray about ½ as long as first ray of second dorsal; this fin inserted under the seventh ray of second dorsal; about 3 of the terminal anal rays might be considered caudal rays; pectoral inserted slightly in advance of ventral, which is in about the same vertical with the origin of the first dorsal, second ray of pectoral slightly produced; length of pectoral equal to that of head without snout; ventral insertion distant from tip of snout a distance equal to that of first dorsal from snout, the first and second rays filamentous, the latter slightly the longer, and extending to the fifteenth or eighteenth ray of anal fin. Color brown; vertical fins bluish or black; peritoneum black; inside of gill covers and roof of mouth bluish. (Goode & Bean.) West Indies and Gulf of Mexico. Three specimens known; the type from near Martinique. (*arcuatus*, arched.)

Bathygadus arcuatus, Goode & Bean, Bull. Mus. Comp. Zool., xii, No. 5, 158, 1883, off Martinique, in 334 fathoms (Coll. *Blake*); Goode & Bean, Oceanic Ichthyology, 421, 1896.

2939. BATHYGADUS FAVOSUS, Goode & Bean.

Head 5½ in total length; depth about 6; eye 5 in head; snout about 4. D. II, 9-125; A. 110; V. 9; P. 14; B. 7; scales 10-135-16. Body heavy, stout, the profile descending gradually and in a slight curve from first dorsal to snout. Scales small, deciduous, cycloid, without armature; interorbital area slightly convex, its greatest width about 3 in head; the postorbital part of head 2⅔ times as long as eye; snout broad, oblique, its width at the nostrils a little more than that of interorbital area; nostrils close to and in front of middle of eye, the posterior somewhat the larger; no barbel. Teeth in both jaws in villiform bands, a naked space at symphysis of intermaxillaries; intermaxillary bands more than twice as wide as those of mandible; vomer and palatines toothless. Gill rakers 20 + 25, the longest on anterior arch slightly more than ½ eye; pseudobranchiæ present, very rudimentary in some individuals, in others wanting or present only on one side; first dorsal distant from snout a distance slightly more than length of head, length of its base about equal to width of snout at nostrils, the fin consisting of 2 spines, the first minute, and 9 branched rays; length of longest dorsal spine, which is armed, 2 in head; second dorsal beginning immediately behind first, the membrane being continuous; anterior rays longest, apparently about ¼ length of head; anal lower than second dorsal, its distance from snout about equal to ⅓ of total length; pectoral inserted under anterior rays of first dorsal and very slightly in advance of origin of ventral, its length more than ½ that of head; distance of ventral from snout 5 times in total length; this fin inserted nearly under base of pectoral; the first ray somewhat produced, its tip reaching to fourth ray of anal fin. Color bluish brown, darkest upon head and abdomen. West Indies. The type specimen, 350 mm. in length, was obtained by the *Blake* from Station LXXX, off Martinique, at a depth of 472 fathoms. (Goode & Bean.) (*favosus*, like honeycomb.)

Bathygadus favosus, Goode & Bean, Bull. Mus. Comp. Zool., xii, No. 5, 160, 1883, off Martinique in 472 fathoms (Coll. *Blake*); Goode & Bean, Oceanic Ichthyology, 420, fig. 352, 1896.

2940. BATHYGADUS MACROPS, Goode & Bean.

Head 5¼ in total length; depth 6½; eye 2⅔ in head; snout 5. D. II, 8—about 125; V. 8. Body somewhat compressed; scales small, deciduous, about 25 rows in an oblique line from the vent to the dorsal fin, 24 from the upper angle of operculum to the vertical through origin of the anal; interorbital area nearly flat, its width 4 in head; postorbital part of head somewhat longer than diameter of eye; snout broad, obtuse; nostrils close to eye, the posterior nearly twice as large as anterior one; maxillary extending to vertical through posterior margin of orbit, its length equal to that of head without its postorbital portion; length of mandible 3 times that of snout; intermaxillaries and mandible provided with narrow bands of villiform teeth, those of the mandible much shorter. A minute barbel, about ⅓ as long as snout. Vomer and palate toothless. Gill rakers lanceolate, elongate, 7 + 26, the longest 7 in head; pseudobranchiæ absent; distance of first dorsal from snout nearly 5 times in total length, second or longest ray in the typical specimen twice length of snout; second dorsal almost continuous with the first, its anterior rays the longest, about 4 times in length of head; anal inserted under fourteenth ray of second dorsal, its rays all very short; in a distance equal to length of head, counting back from insertion, there are 33 rays; pectoral inserted under first branched ray of first dorsal, its length in the most nearly perfect specimens equaling length of head without snout; ventral origin very slightly behind origin of pectoral under third branched ray of dorsal, reaching nearly to vent when laid back, its length equaling 3 times that of the snout. Branchiostegals 7. Color yellowish gray, lighter below. (Goode & Bean.) In deep waters of the Gulf of Mexico off the United States coast, in 321 to 347 fathoms. (*μακρός*, large; *ὤψ*, eye.)

Bathygadus macrops, Goode & Bean, Proc. U. S. Nat. Mus. 1885, 598, Gulf of Mexico, Lat. 28° 34′ N., Long. 86° 48½ W., in 335 fathoms (Type, No. 37339. Coll. *Albatross*); Günther, Challenger Report, xxii, 156, 1887; Goode & Bean, Oceanic Ichthyology, 423, 1896.

2941. BATHYGADUS LONGIFILIS, Goode & Bean.

Head about 5¾ in total length; depth 7½; eye 4 in head; snout 4. D. II, 8 or 9—about 140; P. 13; V. 8; scales about 142. Body more compressed than in *B. macrops*; scales small, cycloid, deciduous, about 25 rows from the vent upward and forward to the dorsal fin, interorbital area flattened, its greatest width 3⅜ times in total length of head; postorbital portion of head twice as long as eye; snout and nostrils normal; maxillary reaching somewhat beyond posterior margin of orbit, its length twice in distance from snout to origin of first dorsal; length of mandible 2½ times in snout; barbel slender, long, its length equal to 1½ times orbital diameter. Teeth in narrow villiform bands in each jaw, none on vomer or palatine bones; gill rakers very long and slender, numerous, 17 + 35, the longest nearly 6 in head; pseudobranchiæ absent; first dorsal of 2 stout spines, the first minute, the second elongate, and 8 or 9 branched rays, its distance from snout 5½ in total; second or longest simple ray nearly 8 times length of

snout, and reaching to or beyond the thirtieth ray of the second dorsal; second dorsal almost continuous with the first, its anterior rays longest and not diminishing rapidly in size toward tail; anal inserted under ninth ray of second dorsal, its rays much shorter than those of dorsal, and situated about same distance apart; pectorals inserted under anterior portion of first dorsal, first ray much produced, extending more than halfway from its insertion to tip of tail; ventral origin slightly behind origin of pectoral, under third branched ray of dorsal, its first ray much enlarged, extending more than halfway from its insertion to tip of caudal, its length 2⅓ times in total length; branchiostegals 7. Color yellowish gray, abdomen bluish. This form is closely allied to *B. multifilis*, described by Günther from off the Philippines (Challenger Report, XXII, 155, pl. 42, fig. B, 1887), which, however, appears to have a smaller eye, less elongate filaments, and ventrals inserted in advance of the first dorsal, while the anal appears to be further back, under the twelfth or thirteenth ray of second dorsal. Both species are provided with long, slender barbels; in other respects they are closer to *B. cottoides*, the typical species, than to *B. macrops*. (Goode & Bean.) Deep waters of the Gulf of Mexico, in 525 to 739 fathoms. (*longus*, long; *filum*, thread.)

Bathygadus longifilis, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 599, Gulf of Mexico, Lat. 28° 47′ 30″ N., Long. 87° 27′ W., in 724 fathoms (Type, No. 37338. Coll. *Albatross*); GÜNTHER, Challenger Report, XXII, 157, 1887; ALCOCK, Ann. Mag. Nat. Hist. 1890, 302; ALCOCK, *l. c.* 1891, 123; GOODE & BEAN, Oceanic Ichthyology, 422, 1896.
Hymenocephalus longifilis, VAILLANT, Exp. Sci. Trav. et Tails, 218, pl. 23, fig. 1, 1888.

996. STEINDACHNERIA, GOODE & BEAN.

Steindachneria, GOODE & BEAN, in AGASSIZ, Three Cruises of the *Blake*, II, 26, 1888 (no type; short diagnosis*); not *Steindachneria*, EIGENMANN, Nematognathi, Occasional Papers, I, Cal. Ac. Sci. 1890, 100 and 202, a genus of Siluroid fishes.
Steindachneria, GOODE & BEAN, Oceanic Ichthyology, 419, 1896 (*argentea*).
Steindachnerella,† EIGENMANN, American Naturalist, February, 1897, 159 (*argentea*).

Body compressed, with tapering tail. Mouth large, terminal. Dorsal fins continuous, both elevated anteriorly; anal divided, the anterior portion elevated, the posterior low. Teeth in each jaw biserial, the outer much enlarged, vomerine teeth present. Bones of head soft and cavernous. Eye large. Gill membranes connected anteriorly, free from the isthmus. Gill rakers slender, rather numerous; vent in anterior third of length. No pseudobranchiæ. Branchiostegals 7. No barbel. Pectorals and ventrals both below first dorsal. Scales thin, cycloid, deciduous. Deep seas. ("This remarkable genus is named in honor of Dr. Franz Steindachner, Custos of the Imperial Zoological Museum of Vienna," one of the ablest naturalists of the century.)

* "*Steindachneria*, a Macruroid with a high differentiated first anal spine." (Goode & Bean.)
† As the original diagnosis of the Macrourid genus *Steindachneria*, although very short, is correct and sufficient for identification, the name in question should be retained for it rather than *Steindachnerella*, and the Silurid genus *Steindachneria*, Eigenmann should receive a new name.

2942. STEINDACHNERIA ARGENTEA, Goode & Bean.

Head 5⅓ in total; depth 7½, at anal origin 8; eye 3⅓ in head; snout about 5½; interorbital width 5½; maxillary 2; premaxillary 2; mandible 1¾; gill rakers 4 or 5+19; D. VIII, 123+; A. 10+113; P. 15; V. 8. Head and body compressed; tail tapering to a very fine point. Scales small, deciduous, cycloid, 6 rows between lateral line and origin of soft dorsal. Nostrils nearer eye than end of snout, the anterior nostril nearly circular, the posterior much longer and slightly concave; no barbel. Maxillary dilated at the extremity and somewhat produced downward into an obtuse point, reaching nearly to a vertical at posterior margin of orbit, and concealed by the preorbital; premaxillaries slightly protractile, much attenuated posteriorly; mandible reaching slightly behind eye. Premaxillary and mandibular teeth biserial, those of the outer series enlarged and rather widely set, some of the enlarged teeth slightly sagittate at tip; vomerine teeth well developed; upper pharyngeal teeth in 2 broad, well-developed patches. Gill rakers slender, the longest about 2 in eye. Distance from snout to first dorsal about ⅓ total length, the first spine elongate, filiform, and reaching fourteenth ray of second dorsal; base of first dorsal about 1 in head; longest ray of second dorsal about 2½ in head, the rays diminishing in size rapidly, the last minute; origin of anal under sixth ray of second dorsal, not far behind the vent, the anterior elevated portion consisting of 10 rays, all of which except the first are divided, the second ray longest, twice length of eye, the tenth ray only about ⅓ length of second, and separated by a small membrane from rest of fin which consists of very minute rays. Vent under fourth ray of second dorsal. Origin of ventrals under base of pectorals and about under third spine of first dorsal; first ventral ray filamentous, reaching origin of anal; pectoral reaching to below fifteenth ray of second dorsal. Gulf of Mexico. Only the type known. Length 233 mm. (*argenteus*, silvery.)

Steindachneria argentea, Goode & Bean, Oceanic Ichthyology, 419, fig. 351, 1896, off delta of Mississipi River, Lat. 39° 14′ 30″ N., Long. 88° 09′ 30″ W., in 68 fathoms. (Type, No. 37350. Coll. *Albatross*.)

997. TRACHYRINCUS, Giorna.

Trachyrincus, Giorna, Mem. Accad. Imp. Turin, xvi, 1803, 178 (no type mentioned).
Lepidoleprus, Risso, Ichth. Nice, 197, 1810 (*trachyrincus*).
Oxycephas, Rafinesque, Caratteri, 31, 1810 (*scabrus* = *trachyrincus*).
Lepidosoma, Swainson, Nat. Hist. Class'n Fish., ii, 261, 1839 (*trachyrhynchus*).
Trachyrhynchus, Günther, Challenger Report, xxii, 152, 1887; corrected spelling.

Snout produced in a long depressed process which is sharply pointed in front, with a sharp lateral edge, which is continued in a straight line across the suborbital region. Mouth inferior, horseshoe-shaped, placed like the mouth of a sturgeon. Teeth in both jaws in villiform bands; chin with a barbel; a scaleless fossa on each side of nape. Second dorsal well developed. Scales moderate, spinigerous; a series of larger scales, each armed with a projecting ridge, along each side of base of dorsal and anal anteriorly. Opercle small. Gill membranes scarcely united; gills 4; first gill arch free, with short, styliform gill rakers. Deep seas. This

genus and its allies differ from *Macrourus* in the important character of the structure of the first gill arch. (τραχύς, rough; ῥύγχος, snout; hence properly, but not originally, spelled *Trachyrrhynchus*.)

2943. TRACHYRINCUS HELOLEPIS, Gilbert.

Head 3½ in total; depth 7; eye large, 4 in head, = interorbital width; snout 2½, its greatest width 1⅔ in its length. D. 11. Snout depressed, flat, narrowly triangular, tapering to a sharp point, its lateral ridges continuous backward over suborbital chain and across cheek. Interorbital space wide and flat. Ethmoidal ridge not prominent. Mouth wholly inferior, U-shaped, overpassed by the snout by a distance contained 3¼ in head. Barbel slender, short, less than ⅓ diameter of orbit. Teeth finely villiform, in very broad bands in each jaw, none of them enlarged. Maxillary reaching to or almost to vertical from hinder margin of orbit, 3⅓ in head. Opercle very small, triangular, its length behind preopercular margin scarcely more than ½ diameter of orbit; outer gill arch not adnate to the opercle, its lower limb with 17 short gill rakers, which are not tubercular. Distance of dorsal fin from nape 3⅓ in head, the 2 dorsal fins closely approximated; second dorsal ray not spine-like, soft and flexible, and not longer than the succeeding rays, its length ⅔ the diameter of orbit. Vent located immediately in front of origin of anal fin, its distance from ventrals 1½ in head. Ventrals short, inserted well in advance of base of pectorals, the outer ray little produced, its length 1⅓ in diameter of orbit. Scales all with their margins embedded, and therefore appearing non-imbricated, the central portion of each projecting, tubercle-like, and bearing a single strong central spine, with sometimes 2 or 3 smaller ones; belly and breast sometimes covered with much smaller scales similarly armed; no naked area between bases of ventrals; enlarged plates along bases of dorsals and anal bearing each a strong compressed backwardly-curved spine, usually without distinct serrations; from the base of the central spine radiate lines of short spinous points; dorsal series of plates continued forward to the nape, the predorsal portion of the included groove covered with scales; ventral series scarcely extending beyond vent, but extending farther posteriorly than do the dorsal plates; scales on top of head with a median serrated ridge; temporal fossæ small but evident, naked. Color apparently dark brown; gill cavity and peritoneum black. Pacific Ocean, off the coast of Central America, in deep water. Only the type known, a specimen 18 inches long. (ἥλος, tubercle; λεπίς, scale.)

Trachyrhynchus helolepis, Gilbert, Proc. U. S. Nat. Mus. 1891, 562, Pacific Coast of Central America in deep water. (Type, No. 48205.)

998. MALACOCEPHALUS, Günther.

Malacocephalus, Günther, Cat. Fish. Brit. Mus., iv, 396, 1862 (*lævis*).

Intermaxillary teeth biserial, mandibulary teeth uniserial. Mouth lateral; snouth short, obtuse. Head without prominent ridges, with wide muciferous cavities. Dorsal fin over origin of pectorals, its longest spine

smooth; dorsal fins widely separated. Pectorals short, placed high, opposite upper angle of gill cleft. Scales small, bristly. Origin of lateral line at upper angle of gill cleft. (*μαλακός*, soft; *κεφαλή*, head.)

2944. MALACOCEPHALUS OCCIDENTALIS, Goode & Bean.

Eye 2¼ in head; barbel slightly longer than eye; snout 4 in head; interorbital space 4. Agreeing with Günther's description of *M. lævis*, but differing in the position of the vent, the ventrals, and the anal fin, the last commencing at a distance behind the vent equal to length of snout; distance of vent from origin of ventrals less than its distance from origin of anal; ventrals originate under middle of first dorsal; origin of pectorals under that of first dorsal, the pectorals as long as head without postorbital flap; ventrals reaching to or slightly beyond origin of anal. Gill rakers rudimentary, $x+11$. Second dorsal spine nearly equal to length of head; first branched dorsal ray about as long as head. Atlantic Ocean, off Cape Hatteras, and Caribbean Sea. Length 8½ inches; a doubtful species, perhaps identical with *M. lævis*. (*occidentalis*, western.)

Malacocephalus occidentalis, Goode & Bean, Proc. U. S. Nat. Mus. 1885, 597, off Cape Hatteras, at Albatross Station 2310, Lat. 35° 44′ N., Long. 79° 51′ W., in 132 fathoms. (Type, No. 37336.)

999. MOSELEYA, Goode & Bean.

Moseleya, Goode & Bean, Oceanic Ichthyology, 417, 1896 (*longifilis*).

This genus is near *Nematonurus*, having the mouth small, the upper teeth in 1 or 2 series, the dorsal spine weakly serrate, and the dorsal fins well separated. The chief difference lies in the scales, which are feebly ridged and nearly or quite smooth. The typical species, *M. longifilis* (Günther), is from off the coast of Japan. ("Named in honor of Prof. Henry N. Moseley, F. R. S., of Oxford University, whose contributions to natural history while naturalist of H. M. S. *Challenger* we desire to commemorate.")

2945. MOSELEYA CYCLOLEPIS (Gilbert).

Dorsal II–8 or 9; ventral 12; eye 4⅓ in head; snout 3⅔; maxillary 2⅔. Head smooth, compressed, without conspicuous ridges; median and lateral rostral ridges terminating in slightly projecting points, the median process, a short portion of the median ridge, and the edge of the membrane connecting median with lateral processes, with spinous scales and points. Snout projecting beyond the premaxillaries for ⅔ its length. Eye small, less than snout, very slightly exceeding interorbital space; mouth small, wholly inferior, maxillary reaching vertical from posterior margin of pupil. Premaxillary teeth in 2 series, the outer similar to those in mandible, not enlarged or canine-like, the inner series smaller, directed obliquely backward; a single series of teeth in mandible, not widening into a patch at symphysis. Barbel thick at base, ⅔ length of snout. Preopercle incurved above the angle, the lower limb expanded, the marginal region striate. First dorsal inserted behind axil of pectoral (second spine broken in both specimens examined), the basal portion smooth, a single sharp

barb showing that the spine is serrate; base of first dorsal equals length of snout; interspace between dorsals exceeding length of first dorsal base by $\frac{1}{8}$ to $\frac{2}{5}$ length of latter. Vent immediately in advance of origin of anal, under middle of interspace between dorsals; dorsal low and inconspicuous and the anal higher, as usual in this group; pectorals very slender, $1\frac{9}{10}$ in length of head; outer ventral ray filamentous, reaching third or fourth anal ray. Scales mostly lost, the few remaining on head either entirely smooth or bearing a single median keel with 1 or 2 low spinous points; those on body without spines, either entirely smooth or showing traces of a low median keel; 6 scales in an oblique series between lateral line and middle of base of dorsal. Color dark brown, the anterior portion of back and sides with small scattered black spots; opercles, lower side of head including gill membranes and ventral area black, as are also the mouth and gill cavity and the peritoneum. A species with the general appearance, including the protruding snout, the inferior mouth and comparatively weak dentition of *Nematonurus armatus* and *N. affinis*, but with the dorsals less widely separated, the vent anterior in position, and the scales unarmed, as in *Moseleya longifilis*. (Gilbert.) Coast of British Columbia. Two specimens, the longest 150 mm., from Station 3342, off Queen Charlotte Islands, depth 1,588 fathoms. (*κύκλος*, circle; *λεπίς*, scale.)

Nematonurus cyclolepis, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 458, off Queen Charlotte Islands, at Albatross Station 3342, in 1,588 fathoms.

1000. NEMATONURUS, Günther.

Nematonurus, GÜNTHER, Challenger Report, Deep-Sea Fishes, XXII, 124, 150, 1887 (*armatus*).

Body rather robust, covered with rough, strongly-ridged scales. Head short; mouth small or moderate, more or less inferior; teeth in upper jaw rather strong, in 1 series or nearly so; lower teeth uniserial; mucous cavities small; pectoral fin inserted low, below upper angle of gill cleft; ventrals well developed, the outer ray filamentous; long ray of dorsal serrated; space between dorsals long, much greater than length of first dorsal. Deep seas. A well-marked genus, distinguished by its rough, firm scales and the wide space between dorsals. (*νῆμα*, thread; *οὐρά*, tail.)

a. Depth $6\frac{1}{2}$ in length; scales without distinct median keel. GOODEI, 2946.

aa. Depth $5\frac{2}{3}$ in length; scales with the median keel prominent; suborbital narrow, with well-marked mucous partitions. SUBORBITALIS, 2947.

2946. NEMATONURUS GOODEI* (Günther).

Head $5\frac{2}{3}$; depth $6\frac{1}{2}$; eye 5 in head; snout $4\frac{1}{4}$; interorbital width $4\frac{1}{4}$; postorbital part of head $8\frac{1}{2}$; first dorsal II, 8 or 9; second dorsal 105; A. 110; P. 20; V. 10; scales 7–150–18, small, strong, free portions covered by series of small vitreous spines arranged in about 6 rows; no specialization of the central row, though the median spine at margin of scale projects

* By some inadvertence this species is recorded by Goode & Bean as a *Hymenocephalus* (Oceanic Ichth., 407). On p. 408 it is said to be a *Nematonurus*. It has obviously no affinity with *Hymenocephalus*, and is, in fact, an ally of *Nematonurus armatus*.

most strongly and is longest. Width of interorbital area a little greater than horizontal diameter of orbit and length of operculum; snout triangular, depressed, its tip in axis of body nearly on a level with lower margin of eye, its lower surface forming an angle with the body axis, about equal to that formed with same by its upper profile; superior ridge pronounced anteriorly, but ending in advance of concavity in interorbital space; lateral ridges prominent, continuing posteriorly to eye, with strong angular projections in front of nostrils; no ridges continued from supraorbital region; nostrils rather close to eye; barbel shorter than eye; tip of lower jaw under anterior nostril; cleft of mouth under posterior margin of orbit; under surface of head naked, with the exception of a few minute, spiny tubercles on under surface of mandible; suborbital ridge very slightly developed; the intermaxillary a long bone, nearly as long as the maxillary; mouth large; teeth on intermaxillary in a double series, those of the outer series much larger than the inner; teeth in mandible uniserial. Dorsal spine strongly serrated; distance of first dorsal from snout equal to nearly 4 times length of its base, its distance from anterior margin of orbit equal to length of head; first spine minute, second strongly serrated, nearly $\frac{2}{3}$ length of head, when laid down is far from reaching origin of second dorsal; when the fin is erect its superior margin is nearly at right angles to plane of back and slightly convex; distance between dorsals twice length of base of first, the second beginning in the perpendicular from fifth ray of anal; anal about 3 times as high as second dorsal; vent under thirtieth scale of lateral line directly in advance of the anal and at a distance from ventral considerably greater than length of that fin; distance of pectoral from snout slightly more than length of head, its length less than that of dorsal spine, slightly more than $\frac{1}{2}$ its distance from the snout, its insertion (upper axil) in middle line of body; insertion of ventral under that of pectoral, slightly in advance of that of dorsal, its first ray not greatly prolonged, about $\frac{1}{2}$ length of distance of fin from snout; branchiostegal membrane narrowly attached to the isthmus, leaving no free margin behind; gill rakers very small tubercles, only 10 below angle on first arch. Color dark reddish brown, spines upon the scales with a metallic luster; young with 3 stellate bosses upon snout, 1 at tip, 1 at some distance upon each side. Length of specimen described 322 millimeters. (Goode & Bean.) Gulf Stream, from Cape Cod to Havana; generally abundant. (Named for George Brown Goode.)

Macrurus asper, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 196, 1883, Gulf Stream south of New England, Lat. 41° 24′ 25″ N., Long. 65° 35′ 30″ W., in 1,242 fathoms; name preoccupied by *Macrurus asper*, GÜNTHER; JORDAN, Cat., 131, 1885.

Macrurus goodei, GÜNTHER, Challenger Report, XXII, 136, 1887; substitute for *Macrurus asper*.

Hymenocephalus goodei, GOODE & BEAN, Oceanic Ichthyology, 407, fig. 340, 1896.

2947. NEMATONURUS SUBORBITALIS (Gill & Townsend).

Head $5\frac{1}{3}$; depth $5\frac{4}{5}$; eye 5 in head; snout $4\frac{2}{4}$; maxillary $2\frac{3}{4}$. D. 12–85; A. 102; P. 19; V. 11. Mouth wholly inferior; scales closely adherent and rather large, mostly short and roundish, with considerable exposed sur-

faces, having radiating ridges beset with weak spines; head a little more than $\frac{1}{5}$ of the entire length; snout projecting but little; median and lateral tubercles faintly developed; infraorbital narrow, divided into 2 well marked areas, an upper wider, distinguised by the glassy tubercular scales, and the narrow lower, almost skinny and scaleless; the ridge independently, is little marked; teeth biserial in the upper jaw, robust in the outer row, very weak in the inner; uniserial in lower jaw and scarcely incurved; dorsal spine strongly serrate, $1\frac{1}{2}$ in head; pectoral $1\frac{7}{8}$ in head; ventrals $1\frac{3}{4}$ in head, with short filaments, reaching vent; interspace between dorsals $\frac{1}{2}$ greater than base of first. Bering Sea. Only the type, 20 inches long, known, the above description taken from it by us. (*suborbitalis*, pertaining to the region below the eye.)

Macrurus (*Nematonurus*) *suborbitalis*, GILL & TOWNSEND, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17, 1897), 234, Bering Sea, southwest of Pribilof Islands, Albatross Station 3603, in 1,771 fathoms. (Type, No. 48773, U. S. Nat. Mus. Coll. *Albatross*.)

1001. ALBATROSSIA, Jordan & Evermann, new genus.

Albatrossia, JORDAN & EVERMANN, new genus (*pectoralis*).

This genus has the form and appearance of *Chalinura*, with the dentition of *Nematonurus*, and the dorsal spines of *Malacocephalus* and *Optonurus;* teeth in the upper jaw strong, in an irregular double series, the outer enlarged; the inner series growing double with age; lower teeth uniserial or nearly so; scales small, rather firm, rough; dorsal spine weak, smooth or very slightly serrate; dorsal fins close together; ventrals well developed; pectorals moderate. Size large. (Named for the good ship *Albatross*, in remembrance of her splendid contributions to our knowledge of the life of the deep seas.)

2948. ALBATROSSIA PECTORALIS (Gilbert).

Head 6 in total; depth $1\frac{2}{3}$ in head; eye $4\frac{1}{2}$ to 5 in head, $1\frac{1}{4}$ in snout. D. X–128; A. 121; V. 7; P. 17; mouth wide, lateral, the short snout projecting beyond premaxillaries for a distance about equaling $\frac{1}{2}$ diameter of orbit; suborbital ridge and lateral ridge on snout inconspicuous; a strong median ridge on snout and a pair of parallel ridges forward from above nostrils; maxillary reaching well behind vertical from posterior margin of orbit, $2\frac{1}{3}$ in head; teeth in 2 somewhat irregular series in front of premaxillaries, the outer series enlarged, the inner directed obliquely inward, the two series merging into one laterally; mandible with a single row, similar to inner series of upper jaw; barbel short, $\frac{3}{7}$ to $\frac{2}{7}$ diameter of orbit; angle of preopercle bluntly rounded, not produced; outer gill arch adnate, as usual in *Macrourus*, 7 short tubercular gill rakers present on its free portion; first dorsal spine slender and weak, with 1 or 2 small retrorse prickles near its middle; distance between dorsals equal to $\frac{2}{3}$ base of first; vent immediately in front of anal origin, its distance from base of ventrals slightly more than $\frac{1}{2}$ head; pectorals long and narrow, reaching vertical from ninth or tenth ray of second dorsal, more than $\frac{1}{2}$ length of head; outer ventral ray produced into a long slender filament, reaching $\frac{5}{6}$ the

distance from its base to front of anal; scales rather small, 10 or 11 in a series between lateral line and origin of second dorsal or middle of first dorsal; scales on sides very thin and flexible, readily deciduous, each furnished with low diverging ridges, usually 3 in number, bearing few minute spinules, and projecting but little beyond the margins of the scales; entire head, including snout and mandibles, invested with much smaller scales irregularly imbricated, those on the opercles marked similarly to those on sides, the others usually each with a single median ridge terminating in a spinous point; no naked spots or pits on head or between ventral fins; a small narrow area behind and below axil of pectorals. Color light grayish, darker on belly and head; mouth, gill cavity, and peritoneum black; lateral line black; dorsals and ventrals dusky; anal lighter, edged with blackish; pectorals black. Bering Sea to Oregon. Specimens have been taken at *Albatross* Stations 3071, 3074, and 3075, in depths of 685 to 877 fathoms, off the coast of Oregon, and from near Bogoslof Island in Bering Sea in 664 fathoms. It is a large, firm-fleshed species, easily recognized. (*pectoralis*, pertaining to the pectoral.)

Macrurus (*Malacocephalus*) *pectoralis*, GILBERT, Proc. U. S. Nat. Mus. 1891, 563, off the coast of Oregon. (Coll. Dr. Gilbert.)

Macrurus (*Nematonurus*) *magnus*,* GILL & TOWNSEND, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17, 1897), 234, Bering Sea, southwest of Pribilof Islands. (Types, No. 48770 and 48771, U. S. Nat. Mus. Coll. *Albatross*.)

Albatrossia pectoralis, JORDAN & GILBERT, Report Fur Seal Invest., 1898.

1002. BOGOSLOVIUS, Jordan & Evermann, new genus.

Bogoslovius, JORDAN & EVERMANN, new genus (*clarki*).

This genus is close to *Chalinura*, from which it is distinguished by its dentition, having the teeth in the upper jaw in 2 series, the outer slender and sharp, slightly arrow-shaped; those of the inner small, close set, replacing the villiform band of *Chalinura*. Scales excessively rough; ventral filament produced; dorsal spine filamentous, sharply serrate; dorsal fins close together; pectorals inserted below upper angle of gill opening. Deep seas. (Named for the volcanic island, St. John Bogoslof, in Bering Sea, near which the typical species was dredged.)

a. Ventrals much longer than head, reaching far beyond front of anal. CLARKI, 2949.
aa. Ventrals shorter than head, scarcely reaching front of anal. FIRMISQUAMIS, 2950.

* We have examined the type and cotypes of *Macrurus* (*Nematonurus*) *magnus*, Gill & Townsend, and find them to agree fully with *Albatrossia pectoralis* (Gilbert). The type may be redescribed as follows: Head $5\frac{1}{4}$; depth $7\frac{1}{2}$; eye $4\frac{3}{4}$ in head; snout $4\frac{2}{3}$ to $4\frac{1}{2}$; maxillary $2\frac{1}{2}$; pectoral 2 in head; ventral with short filament, $2\frac{1}{2}$ in head. Mouth large, with lateral cleft. Dorsals well separated, the interspace not $\frac{1}{2}$ base of first dorsal; long dorsal spine smooth, or with 1 or 2 roughnesses near its tip, its length $3\frac{2}{3}$ in head; second dorsal low; pectoral inserted low, below angle of opercle. Scales moderately large, readily deciduous, decidedly oblong or long, with a small exposed surface which is beset with about 5 radiating ridges with conspicuous spinigerous ridges on dorsal surface, but not armed at tip; head regularly conical; snout rather long, projecting $\frac{1}{3}$ its length beyond mandible; tubercles feebly developed, plain, and continuous from 3 parallel ridges; infraorbital flat, with the crest rather nearer the orbit than its lower margin; its entire surface scaly; teeth in the upper jaw biserial or triserial in front, the outer series strongly hooked, the inner series considerably smaller and well separated from the outer series; an irregular series between in the type specimen; teeth in lower jaw uniserial or irregularly biserial. Three specimens, the largest (type of *M. magnus*) 43 inches long.

2949. BOGOSLOVIUS CLARKI, Jordan & Gilbert.

Eye $4\frac{2}{3}$ in head; maxillary $2\frac{1}{2}$. D. II, 12– ; P. 19; V. 10. Snout short, slightly exceeding diameter of eye, $3\frac{9}{10}$ in head; median and nasal ridges very little projecting anteriorly, without radiating spines; tip of snout very little projecting beyond the mouth, for a distance not exceeding $\frac{1}{8}$ the interspace between ends of median and nasal ridges. Suborbital ridge inconspicuous, scarcely extending beyond the eye; mucous pores on head prominent. Mouth large, oblique, the lower jaw included, the maxillary nearly reaching vertical from posterior edge of orbit. Outer premaxillary teeth slender, sharp, unequal, rather distant, not very strong, slightly widened and arrow-shaped near tip, becoming very small toward angle of mouth; within this, and well separated from it, a close-set series of short teeth directed inward. Mandibular teeth slender, unequal, in a single series corresponding to outer series in upper jaw, slightly widening at symphysis, which is not prominent. Barbel very short, less than $\frac{1}{2}$ diameter of pupil. Eye of moderate size, equaling distance from tip of snout to middle of anterior nostril, $1\frac{1}{10}$ in interorbital width. Preopercle broadly rounded, the angle little produced backward, leaving a strip of interopercle exposed along its entire length. Gill membranes joined to the isthmus, with a narrow free edge. Gill rakers very short and thick, 3 + 12 in number, including rudiments. Dorsal beginning above base of pectorals, the second spine long, filamentous at tip, $1\frac{2}{3}$ in head, its anterior margin sharply serrate, except in basal third; base of first dorsal $2\frac{1}{4}$ in head; interspace between dorsals very short, usually less than diameter of pupil. Pectorals very long and slender, equaling or exceeding length of head behind snout; insertion of pectorals below upper angle of gill opening. Outer ventral ray excessively produced, twice or more than twice length of head in uninjured adults, reaching base of fiftieth anal ray or beyond. Vent immediately before anal origin. Scales in a strip along the back firm and very rough, none others preserved in our specimens; scales with 3 to 5 sharp, radiating ridges, each ridge with several sharply projecting spines, the posterior of which project beyond the margin of the scale. Color very light gray, the vertical fins blackish posteriorly; mouth and gill cavity and peritoneum jet-black. Bering Sea. Known from 4 specimens, 24 to 41 cm. long, from *Albatross* Station 3634, off Bogoslof Island, in 664 fathoms. (Named for George Archibald Clark, secretary of the Fur Seal Commission for 1896 and 1897, in recognition of his researches on the mammalia of Bering Sea.)

Bogoslovius clarki, Jordan & Gilbert, Report Fur Seal Invest., 1898, Bering Sea off Bogoslof Island, in 664 fathoms.

2950. BOGOSLOVIUS FIRMISQUAMIS (Gill & Townsend).

Head 5 in total; depth $6\frac{1}{2}$; eye $4\frac{2}{3}$ in head; snout $3\frac{2}{3}$; second dorsal spine $1\frac{2}{3}$ in head; pectoral 2; ventral $1\frac{1}{2}$; maxillary $2\frac{1}{3}$. D. II, 10-126; A. 105; P. 20; V. 8. Scales firmly affixed, oblong or rather short, and with considerable exposed surfaces, which have subequal radiating ridges beset with numerous acute spinelets, the ridges varying from 3 to 8 in number; head regularly convex in profile: rostral tubercles obsolete and infraor-

bital ridge rounded; barbel greater than pupil; teeth biserial or partly triserial above; second dorsal spine with short retrorse serræ, the lower fifth smooth; base of first dorsal 3⅓ in head; interspace between dorsal fins ½ base of first dorsal, greater than diameter of pupil. This species is distinguishable from most American *Macrouri* by the very firm scales, and from *B. clarki* by the much shorter ventral. Bering Sea. Only the type, 31 inches long, known. (*firmus*, firm; *squama*, scale.)

Macrurus firmisquamis, Gill & Townsend, Proc. Biol. Soc. Wash., xi, 1897 (Sept. 17, 1897), 234, Bering Sea, southwest of Pribilof Islands. (Type, No. 48772, U. S. Nat. Mus. Coll. *Albatross.*)

1003. CHALINURA, Goode & Bean.

Chalinura, Goode & Bean, Bull. Mus. Comp. Zool., x, No. 5, 198, 1883 (*simula*).
Chalinurus,* Günther, Challenger Report, xxii, 124, 144, 1887; change in spelling.

Scales cycloid, fluted longitudinally, with slightly radiating striæ. Snout long, broad, truncate, not much produced. Mouth lateral, subterminal, very large. Head without prominent ridges, except the subocular ones and those upon the snout. Suborbital ridge not reaching angle of preopercle. Teeth in the upper jaw in a villiform band, with an outer series much enlarged, those of the lower jaw uniserial, large. No teeth on vomer or palatines; small pseudobranchiæ present. Gill rakers spiny, strong, depressible, in double series on anterior arch. Ventrals below the pectorals; chin with a barbel. Dorsal spine serrate; soft dorsal much lower than anal. Deep sea fishes. Species numerous. This genus is allied to *Macrourus*, differing in the dentition; the genus *Optonurus*, with dorsal spine unarmed, is very close to *Chalinura*. (χαλινός, a strap or thong; οὐρά, tail.)

a. Snout long, longer than eye, which is 5 in head; pectoral 1⅔ in head; dorsal spine 1½ in head; scales 130. SERRULA, 2951.

aa. Snout moderate, about as long as eye, which is 4 in head; dorsal, pectoral, and ventral produced, the pectoral 1⅓ in head, the dorsal spine and ventral filament each about as long as head. FILIFERA, 2952.

aaa. Snout very short, as long as eye, which is 5 in head; ventrals very long. SIMULA, 2953.

2951. CHALINURA SERRULA, Bean.

Head 5½ in total length. D. II, 9–76 (?); scales 7 or 8–130–17; Br. 6. Cheeks and opercles scaly; snout with a median serrated keel on the nose; diameter of eye less than length of snout, 5 in head; maxillary reaching vertical from posterior margin of eye, its length 2½ in head; mandible about 2 in head, a row of 5 pores on its under surface and 6 pores on the edge of the suborbitals; branchiostegal membrane narrowly free from the isthmus, the first gill opening restricted as in *Macrourus;* gill rakers small tubercles, 11 below the angle of the first arch, and only 1 or 2 above the angle; length of pectoral equals postorbital part of head; ventrals

* Goode & Bean rightly protest against the wanton "action of the English ichthyologists in changing the form of the generic name" *Chalinura*. *Chalinura* is perfectly correct, and should be used even if it were not so, as it is the original form, the only reason for changing it being that other generic names in the group end in *urus*.

about as long as head; longest dorsal spine strongly serrated and nearly equaling length of head without snout; dorsals separated by an interspace $\frac{2}{3}$ as long as head. Color brown; head, abdomen, and inside of mouth purple, the purple areas less marked in the type specimen, which is 12½ inches long. Coast of British Columbia, east of Prince of Wales Island, in 1,569 fathoms. (*serrula*, a fine saw.)

Chalinura serrula, Bean, Proc. U. S. Nat. Mus. 1890, 37, east of Prince of Wales Island, in 1,569 fathoms. (Coll. *Albatross*.)

2952. CHALINURA FILIFERA, Gilbert.

D. II, 12 to 14; P. 20 to 22; V. 9 or 10; eye 4 in head. Snout short, slightly exceeding diameter of eye, $3\frac{9}{10}$ in head, median ridge and nasal ridges terminating each in a much projecting point, furnishing each with a short rosette of radiating spines and ridges, outline between these points concave; tip of snout projecting beyond premaxillaries for a distance equaling that which separates the central rosette from 1 of the lateral ones; infraorbital ridges inconspicuous, not reaching angle of preopercle behind or bony portion in front. Mouth large, slightly oblique, with extensive lateral cleft, the maxillary reaching vertical from posterior margin of pupil, $2\frac{3}{5}$ in head, equaling distance from tip of snout to middle of eye. Outer series of teeth in premaxillary strong, succeeding from a narrow band of smaller cardiform teeth; mandibular teeth similar to inner band of upper jaw, the band becoming slightly wider at the prominent symphysis. Barbel short, $\frac{1}{2}$ to $\frac{2}{3}$ length of snout. Eye large, the diameter of orbit slightly less than interorbital width on snout. Angle of preopercle produced backward, concealing all but the extreme posterior angle of interopercle, the margin appearing serrulate when divested of skin; gill membranes joined to isthmus, with a posterior free margin; gill rakers very short and heavy, 1 + 11. Dorsal beginning vertically above base of pectorals, the second spine extremely long and slender, smooth basally; the terminal half rather strongly toothed, becoming very slender toward tip and terminating in a long membranaceous filament. (In 1 specimen it exceeds length of head, in the others it equals $\frac{5}{6}$ that length.) Length of base of first dorsal equaling $\frac{1}{5}$ length of head; interspace between dorsals short, $\frac{2}{3}$ to $\frac{3}{4}$ length of snout. Pectorals very long and slender, equaling the head without the snout; outer ventral rays very long and filamentous, equaling length of head; vent immediately in advance of anal origin. Scales rather thin, those on back and sides with above 5 diverging ridges, each of which bears a number of short rigid spinules directed very obliquely backward, the posterior projecting but little beyond the margin of the scale; 8 or 9 scales in an oblique series between the middle of first dorsal and the lateral line. Dark brown; the fins, gill membranes, lips, nostrils, and underside of snout black; anterior part of mouth and lining of gill cavity purple; peritoneum blackish brown. Related to *C. serrula*, Bean, from the same region and depth, differing in the larger eye, shorter mental barbel, longer snout, longer pectoral fins, shorter interspace between dorsals and the longer dorsal fin. Coast of British Colum-

bia; known from 3 specimens, 520 to 550 mm. long. (Gilbert.) (*filum*, thread; *fero*, I bear.)

Chalinura filifera, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 458, off Queen Charlotte Islands, at Albatross Station 3342, in 1,588 fathoms.

2953. CHALINURA SIMULA, Goode & Bean.

Head 5; depth 6; orbit 6 in head; snout 3; interorbital width greater than eye; postorbital part of head 3 times as long as eye; opercle 2 in upper jaw. D. II, 9–113; A. 118; P. 20; V. 9; Br. 6; scales 8–150–17 to 19. Body shaped much as in *Coryphænoides*, but rather stout; back more gibbous in profile, the dorsal outline rising quite rapidly from the interorbital region to origin of first dorsal, thence descending almost in a straight line to end of tail. Preopercle emarginate on its posterior limb. Snout broad, obtuse, scarcely projecting beyond the mouth, its width nearly as great at tip as its own length; median ridge very prominent, gibbous in outline when viewed laterally; lateral ridges starting almost at right angles with the median, and continued upon sides of head; no supraorbital ridges. Nostrils in front of middle of eye, and nearer its anterior margin than to tip of snout; barbel longer than eye; teeth in upper jaw in a broad villiform band, the outer series very much enlarged; lower jaw with teeth in a single series. Scales rather small, but with indications, particularly on the head, of radiating striæ. Origin of first dorsal from snout 4½ in its base, or from anterior margin of orbit 1 in head; first dorsal spine very short, second rather stout, 1½ in head, and with a simple serration anteriorly, the serræ closely appressed to the spine; second dorsal separated from the first by a distance equal to length of upper jaw; anal high, its average rays about 3 times as long as those of dorsal, inserted slightly behind perpendicular from last ray of first dorsal; pectoral inserted over base of ventral; origin of ventral from snout less than its longest ray, which is produced in a filament extending to base of eighteenth anal ray. (Goode & Bean.) West Indies and Gulf Stream, in deep water. (*simulus*, pug-nosed.)

Chalinura simula, GOODE & BEAN, Bull. Mus. Comp. Zool., X, No. 5, 199, 1883, Gulf Stream, at Blake Station 308, Lat. 41° 25′ 45″ N., Long. 65° 35′ 30″ W., in 1,242 fathoms; JORDAN, Cat., 132, 1885; GOODE & BEAN, Oceanic Ichthyology, 412, fig. 345, 1896.
Macrurus simulus, GÜNTHER, Challenger Report, XXII, 148, 1887.

1004. CORYPHÆNOIDES, Gunner.

Coryphænoides, GUNNER, Trondhj. Selsk. Skrift., III, 50, 1765 (*rupestris*).
Branchiostegus, RAFINESQUE, Analyse de la Nature 1810, 86 (substitute for *Coryphænoides*).

Snout short, obtuse, high, obliquely truncated, soft to the touch, except its bony center; mouth broad, terminal, its cleft lateral; head without prominent ridges, the membrane bones of the side of the head soft and papery; teeth villiform in both jaws, those in the outer series of upper jaws somewhat enlarged. Scales spinous, second or elongate dorsal ray finely serrated in front. Lower jaw with a barbel at tip. Deep Sea.

Close to *Macrourus*, differing in the larger terminal mouth. (*κορύφαινα, Coryphæna; εἶδος*, resemblance.)

a. Head 4 in length; gill rakers 4+15=19. RUPESTRIS, 2954.
aa. Head 6 in length; gill rakers 3+11=14. CARAPINUS, 2955.

2954. CORYPHÆNOIDES RUPESTRIS, Gunner.

D. 10; P. 19; V. 7; gill rakers 4+15=19. Head short, rather compressed; snout short, obliquely truncated in front; cleft of mouth wide, lateral, extending to beyond the center of eye; intermaxillary not much shorter than maxillary. Teeth in villiform bands in each jaw; barbel very small. Interorbital space convex, its width being considerably more than diameter of eye, which, in a specimen 3 feet long, is equal to the length of the snout and ¼ of that of the head. Scales equally rough over the whole of their surface, all the spinelets being directed backward; 7 or 8 scales in a transverse series between the dorsal fin and the lateral line; head entirely covered with small scales. Anterior dorsal spine armed with numerous small closely set barbs; outer ventral ray produced into a long filament. Distance between the vent and isthmus ⅔ the length of the head. The gill membrane entirely free from the isthmus behind. Intermaxillary continues beyond its vertical process and extending almost as far back as the maxillary, these 2 bones being about equal in length; last third of intermaxillary toothless; intermaxillary teeth in a very narrow band, which is uniform in width, the outer teeth only slightly enlarged; mandible with villiform teeth in a broad bunch-like band at the symphysis and becoming uniserial behind. Eye nearly circular. Snout projecting slightly. Gill rakers longer and less tubercular in character than in *Macrourus berglax* and *M. acrolepis*. The suborbital ridge feebly developed and very abruptly curved upward and narrowed in front of the eye where it joins the nasal ridge. In *M. berglax* and *M. acrolepis* the suborbital ridge is very strong and is continued almost in a straight line toward the nasal ridge. (Goode & Bean.) Arctic seas and the north Atlantic, on both coasts south to the banks of Newfoundland and Norway, in deep water. (Eu.) (*rupestris*, living about rocks.)

Coryphænoides rupestris, GUNNER, Trondhjem Selsk. Skrift., III, 50, pl. 3, fig. 1, 1765, Norway; COLLETT, Norges Fiske, 131; JORDAN & GILBERT, Synopsis, 812, 1883; GOODE & BEAN, Oceanic Ichthyology, 402, 1896.
Lepidoleprus norvegicus, NILSSON, Prodr. Ichth. Scand., 51, 1832, Norway.
Coryphænoides norvegicus, GÜNTHER, Cat., IV, 396, 1862.
Macrourus stromii, REINHARDT, Dansk. Vidensk. Afhandl., VII, 129, 1828: GAIMARD, Voy. Skand., Poiss., pl. 11.
Macrurus rupestris, GÜNTHER, Challenger Report, XXII, 138, 1887.

2955. CORYPHÆNOIDES CARAPINUS, Goode & Bean.

Head 6. D. II, 8–100; A. 117; V. 10; eye 4 in head. Snout acute, projecting beyond the mouth, its tip at a distance from the mouth equal to or greater than diameter of eye. Bones of head very soft and flexible, its surface very irregular, there being a very prominent subocular ridge, a prominent ridge extending from tip of snout to middle of interorbital space, and a curved ridge extending from upper anterior margin of orbit

over cavity containing nostrils to a prominent point at side of and slightly posterior to tip of snout; barbel $\frac{2}{3}$ as long as eye. Interorbital space almost twice diameter of eye, equal to length of upper jaw; preoperculum crenulate; upper jaw extending to vertical through posterior margin of pupil, its length equaling $\frac{1}{2}$ that of head without snout; mandible extending behind vertical through posterior margin of orbit, its length 3 times in distance from tip of snout to origin of first dorsal. Teeth in villiform bands on intermaxillary and mandible, the mandibulary series uniserial in about the second half of its length. First ray of dorsal very short, second compressed anteriorly and serrated, with slender teeth closely appressed and bent upward, its length equaling length of head and greater than height of body; this fin seated upon a hump-like elevation of the back, its base as long as snout; second dorsal beginning over tenth or twelfth anal ray, and at a distance from end of first dorsal equal to length of head without snout; vent located not far behind vertical from end of first dorsal. Scales 22 to 24 in a transverse series (the position of the lateral line can not be determined, but there appear to be 4 above it); scales oval, membranaceous, showing several parallel ridges composed of small spines. Gill membrane very deeply cleft and attached to the isthmus; gill rakers short and stout, about 11 below the angle on the first arch. (Goode & Bean.) Gulf Stream, in deep water. (*carapinus*, formed as in *Carapus*.)

Coryphænoides carapinus, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 195, 1883, Gulf Stream, Lat. 41° 24′ 45″ N., Long. 65° 35′ 30″ W., in 1,242 fathoms (Type in M. C. Z. Coll. *Blake*); GÜNTHER, Challenger Report, Deep-Sea Fishes, XXII, 139, 1887; GOODE & BEAN, Oceanic Ichthyology, 404, fig. 339, 1896.

1005. HYMENOCEPHALUS, Giglioli.

Hymenocephalus, GIGLIOLI, Pelagos, Genoa, 228, 1884 (*italicus*).
Mystaconurus, GÜNTHER, Challenger Report, Deep-Sea Fishes, XXII, 124, 1887 (*italicus*).

This genus is closely allied to *Coryphænoides*, differing in the smooth dorsal spine, and the membranaceous skull. First dorsal broad, placed far forward over base of pectoral; second dorsal and anal origins nearly opposite, and separated by a considerable space from the vertical from the end of first dorsal; vent far from ventrals. Head large, naked, soft, and cavernous; snout abrupt, perpendicular, or parabolic; mouth lateral, wide. Eye very large, orbital margin forming part of profile of head. Barbel long. Pectoral rather narrow (10 to 16 rays). Scales thin, deciduous, with fine short spines. Under parts in advance of ventral wholly or partly naked. Deep seas. Remarkable for the papery structure of the bones of the head. (ὑμήν, membrane; κεφαλή, head.)

2956. HYMENOCEPHALUS CAVERNOSUS (Goode & Bean).

Head about 6 in total length; depth 7. D. II, 10–133; A. 27 rays, in a space equal to length of head. Body stoutish, the bones of head very soft and cavernous, spongy, in many places without muscular covering; interorbital area doubly concave, with a spinous medial ridge, its greatest

width about 2⅔ in length of head; postorbital portion of head about ½ its length, 1⅓ as long as eye, which is circular, its diameter contained 2⅔ times in length of head. Snout broad, very obtuse, its width at nostril nearly equal to interorbital width, its length 4⅔ times in that of the head; nostrils normal. Teeth in each jaw in villiform bands, very small; a naked space at the symphysis of intermaxillaries; vomer and palatine toothless. Gill-rakers very short, minute, and rather numerous, about 18 below angle of anterior arch. Pseudobranchiæ absent. Barbel ⅔ as long as eye. First dorsal composed of 2 spines, the first minute, inserted at a distance from the snout equal to length of head, the second as long as head without snout, and 10 branched rays, its base equal to diameter of eye; second dorsal almost rudimentary, its rays remarkably short, about 133 in number, its distance from first dorsal ½ length of head; anal much higher than second dorsal, its distance from snout contained about 3½ times in total length; anterior anal rays longest, in length about ¾ diameter of eye; pectoral inserted under first branched ray of first dorsal, its length equal to twice that of eye and about ⅔ that of head. Scales (on type) mostly wanting, except a few on breast and nape, these being rough with small points, dentate behind. Ventral slightly behind the pectoral, its first ray filamentous, reaching to the base of the tenth anal ray, consisting of 11 rays. Color gray, with silvery tints on sides; abdomen and lips dark. (Goode & Bean.) Gulf of Mexico, in deep water. One young individual known. Length 162 mm. (*cavernosus*, cavernous.)

Bathygadus cavernosus, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 598, Gulf of Mexico, at Albatross Station 2398, Lat. 28° 45′ N., Long. 86° 26′ W., in 227 fathoms (Type, No. 37337. Coll. *Albatross*); GÜNTHER, Challenger Report, XXII, 156, 1887.

Hymenocephalus cavernosus, GOODE & BEAN, Oceanic Ichthyology, 408, fig. 341, 1896.

1006. MACROURUS, Bloch.

Macrourus, BLOCH, Ichth., V, 152, 1787 (*rupestris* = *berglax*).

Macruroplus, BLEEKER, Versl. Med. Akad. Welenth. Amsterd., VIII, 1874, 369 (*serratus*).

Macrurus, GÜNTHER, Cat., IV, 392, 1862; corrected spelling.

Snout broadly conical, high, projecting beyond mouth; mouth moderate, its cleft horizontal, U-shaped, entirely inferior; teeth in both jaws in villiform bands, those of the outer series not enlarged; head with roughened bony ridges, one of which, on the suborbital and preorbital, simulates the suborbital stay of the Cottoids; eyes very large; scales imbricate, very rough, keeled. Dorsal spine long, serrated on the anterior edge. Deep water fishes. (μακρός, long; οὐρά, tail, hence correctly written *Macrurus*, but *Macrourus* is the original name as given by Bloch.)

a. Top of head with 4 to 6 distinct ridges; depth 6 to 7 in length; 5 scales between lateral line and dorsal.

b. Anal rays 148; scales each with a strong ridge. BERGLAX, 2957.

bb. Anal rays 121; scales each with 3 to 5 spinules, otherwise almost unarmed; ridges on top of head very rough. HOLOTRACHYS, 2958.

aa. Top of snout with indistinct ridges or with none.

c. Pectoral fin moderate, 1½ to 2 in head.

d. Body rather elongate, the depth 7 to 8 in length; bones of head rather firm; dorsal spine strongly serrated.

e. Head short, 6⅓ in head; pectoral more than ⅓ head; snout with bony ridges above. BAIRDII, 2959.

ee. Head 5⅓ in length; ventrals 5 in body; pectoral 1⅓ in head. LEPTURUS, 2960.

eee. Head longer, 4¾ in length; pectoral ⅓ as long as head; eye as long as snout, 4 in head; ventral 8 in body. ACROLEPIS, 2961.

dd. Body rather robust, the depth 5⅓ in length; head without ridges above; scales spinous, not ridged; dorsal spine 1⅔ in head. STELGIDOLEPIS, 2962.

cc. Pectoral fin elongate, about as long as head; head elevated, not ridged above, the bones soft; eye large; second dorsal spine rough, nearly as long as head; scales each with 7 to 9 ridges. CINEREUS, 2963.

2957. MACROURUS BERGLAX, Lacépède.

D. 12–124; A. 148; P. 18 or 19; V. 8. Short snout, subtrihedral, pointed in front, much shorter than the large eye, which is ⅓ or ⅖ length of head in the adult. Intermaxillary very short, ½ length of maxillary, and not continued beyond its expanded vertical process. Eye oblong. Whole under surface of head below suborbital and nasal ridge naked; axil of pectoral naked; space between ventrals scaled; body scales each with a single strong median keel, made up of 5 to 8 spines directed backward; some scales, particularly of head, have also 2 lateral keels; 6 longitudinal series of scales between first dorsal fin and lateral line; first dorsal spine indistinctly denticulated toward the point; length of pectoral nearly or quite ⅓ length of head; longest spine of dorsal very finely serrated along its anterior margin, the serrations becoming obsolete near its base. Vent situated behind origin of second dorsal fin. Gill rakers very small, tubercular, 9 to 11 on the first arch; gill membranes broadly joined, free from the isthmus behind. This form, originally discovered on the coast of Norway, has been found abundantly as far south as Georges Bank, where the halibut fishermen catch it, or some closely allied form, on their trawls. The first specimen seen by American naturalists was picked up floating at the surface off the mouth of New York Harbor. The *Albatross* obtained it from Station 2528, in Lat. 41° 47′ N., Long. 65° 37′ 30″ W., at a depth of 677 fathoms. Günther knew it from Finmark and Greenland, as well as from New England. He calls attention to remarkable individual variations in the specimens examined by him. (*berglax*, Norwegian name, from *berg*, cliff; *lax*, salmon.)

Macrourus berglax, LACÉPÈDE, Hist. Nat. Poiss., III, 170, 1800, Greenland, Söndmöre; JORDAN, Cat. Fish. N. A., 131, 1885.

Macrourus fabricii, SUNDEVALL, Vet. Akad. Handl. 1840, 6; COLLETT, Norges Fiske, 128, 1875; LILLJEBORG, Sverig. og. Norges Fiske, 242; GOODE & BEAN, Cat. Fish. Essex Co. and Mass. Bay, 7, 1879; GÜNTHER, Challenger Report, XXII, 130, 1887.

Macrourus rupestris, GÜNTHER, Cat. Fish. Brit. Mus., IV, 390, 1862 (not of Gunner).

Macrurus berglax, GOODE & BEAN, Oceanic Ichthyology, 391, fig. 334, 1896.

2958. MACROURUS HOLOTRACHYS, Günther.

Head 4¾ in length; depth 6¾. D. 12–115 to 125; A. 121; P. 20 or 21; V. 5; eye large, round, as long as snout, 2⅖ in head, much wider than interorbital space. Snout triangular, each point with a tubercle, covered with

strong spines, this border continued as a strong ridge below eye, extending across opercle, this crest covered with coarse, spinous tubercles; mouth rather small, the maxillary reaching middle of eye; teeth very small, close set. Head with salient ridges above, covered with spinous scales; 1 ridge above eye, toward upper angle of gill opening, another ridge along the vertex, nearly parallel with this above it, besides a short temporal ridge; vent far back, under seventh ray of second dorsal. First dorsal not far behind eye, the long ray slightly serrulate; ventrals with a short filament. Scales each with a median crest of 3 to 5 spinules, otherwise almost unarmed. Five scales between lateral line and dorsal. (Collett.) Depths of the Atlantic. Known from 2 specimens, the type 9 inches long, from the mouth of Rio de la Plata, in 600 fathoms; the second, above described, about a foot long, from the banks of Newfoundland, in 1,267 fathoms. (ὅλος, wholly; τραχύς, rough.)

Macrurus holotrachys, Günther, Ann. Mag. Nat. Hist. II, 1878, 24 mouth of Rio de la Plata in 600 fathoms; Günther, Challenger Report, XXII, pl. 28, fig. B, 1887; Collett, Compagnes Scient. de l'Hirondelle, 1896, 83, pl. 2, fig. 6; Goode & Bean, Oceanic Ichthyology, 396, 1896.

2959. MACROURUS BAIRDII, Goode & Bean.

(Common Rat-tail.)

Head 6½ in total length; depth 8; greatest width 13. D. II, 11–137; A. 120; P. 15; V. 7; scales 6–152–19 or 20. Body much compressed posteriorly, tapering from first dorsal to tip of tail; scales irregularly polygonal, the free portions covered with transparent vitreous spines, arranged in from 10 to 12 irregular longitudinal rows. On head and upper part of body, in advance of the first dorsal, the median row of spines most prominent, and presenting the appearance of a low median keel. Lateral line nearly straight, formed by a smooth groove, which replaces 2 or 3 median rows of spines of each scale; greatest height at posterior margin of orbit greater than width at same point, 1⅔ times in length of head; width of interorbital area equal to length of snout and length of maxillary; length of postorbital region about equal to horizontal diameter of orbit; length of operculum about ½ length of mandible. Snout sharp, a front view presenting 4 ridges radiating from tip at right angles to each other, the lower one being merely a fold in the skin of the under surface of the head, horizontal ridges continued into the ridges upon the suborbitals; ridge extending backward from tip of snout upon top of head lost in the interorbital space; branches of the horizontal ridges continued upon upper margins of orbits, and there disappearing. Nostrils immediately in front of orbit, the posterior pair much the longer. Mouth situated entirely on lower side of head; symphysis of lower jaw in vertical from anterior margin of orbit, and articulations of mandibles in vertical from posterior margin of orbit; width of cleft of mouth equal to distance between symphysis of maxillaries and line connecting their articulations; upper jaw protractile vertically. Teeth conical, somewhat recurved, of nearly uniform size, arranged in villiform bands; palate smooth. Distance of first dorsal from snout about 4 times the length of its base, and from anterior margin of

orbit equal to length of head; first spine very short, not much longer than the teeth of the second spine; second spine in length twice horizontal diameter of orbit, stout, its anterior margin armed from base to tip with 15 teeth pointing upward, the uppermost slender; its length to tip of filament almost equal to distance from origin of second dorsal, this tip when laid back reaching almost to second dorsal; rays decreasing regularly in length so that, when the fin is upright, its shape approximates that of a right-angled triangle, the hypothenuse of which is the second dorsal spine, and its perpendicular side a line touching the tips of the rays; length of base of second dorsal less than that of the anal, its origin over the thirteenth scale of lateral line. Length of longest ray less than length of barbel; all rays very feeble; membrane scarcely perceptible; distance of anal from snout $3\frac{4}{5}$ times in its length at base, its origin under eighteenth scale of lateral line; length of first ray $\frac{1}{2}$ the length of tenth, and 3 times the length of last ray, the length of rays increasing to a point beneath anterior part of first dorsal, and thence gradually decreasing to tip of tail; distance of pectoral from snout 4 times width of interorbital area, its length twice length of mandible; insertion above the middle of depth of body, on a level with center of orbit, its third ray longest, its tip reaching to vertical from base of fourth ray; insertion of ventral behind pectoral and almost under that of first dorsal, its distance from snout slightly exceeding twice its length; tip of ventral filament reaching base of third anal ray. Ground color, light brownish gray; under parts silvery; belly darker, bluish; under surface of snout pink, as is also the first dorsal, except spines; spines of dorsal, ventral, and anterior anal rays blackish; throat, branchiostegal membrane, and isthmus rich deep violet; sclerotic coat green; eyes very dark blue. This species was the first deep-sea fish obtained by the Fish Commission or described by an American ichthyologist. It ranges in depths from 9 to 1,255 fathoms. This species is distinguished by Günther from his *Macrourus æqualis*, which it closely resembles, (1) by its longer snout, which is nearly equal to the diameter of the eye, and (2) by the smaller number of ventral rays (7). (Goode & Bean.) West Indies to Massachusetts Bay, usually in great depths; excessively abundant on the continental slope, with *Phycis chesteri*, far outnumbering all other deep-sea fishes in the region. (Named for Spencer Fullerton Baird.)

Macrourus bairdii, Goode & Bean, Amer. Journ. Sci. and Arts 1877, 471, Massachusetts Bay; Goode, Proc. U. S. Nat. Mus. 1880, 337, 475; Günther, Challenger Report, xxii, 135, pl. 22, fig. B, 1887; Goode & Bean, Oceanic Ichthyology, 393, fig. 335, 1896.

2960. MACROURUS LEPTURUS, Gill & Townsend.

Head $5\frac{1}{4}$; depth 8; eye $4\frac{2}{3}$ in head; snout 4; maxillary $2\frac{3}{4}$. D. XIV–122; A. 116; P. 20; V. 8. Scales deciduous and moderate, oblong or oval with reduced exposed surfaces, those on the back or above the lateral line with a few, 3 to 5, ridges beset with spines, but those below mostly unarmed; head regularly conical; snout moderately extended; median tubercle very projecting, the lateral well developed, connected with the median by a well-defined ridge; infraorbital vertical, with the ridge linear and near

the orbit; teeth cardiform in both jaws; the lower teeth beset the outer slope of the jaw. Ventral as long as head; pectoral $1\frac{2}{3}$ in head; dorsal spine serrate, $1\frac{1}{3}$ in head. Apparently close to *M. acrolepis*, but probably with shorter head, longer ventrals, and longer dorsal spine, the eye also larger. Length 22 to 26 inches. Bering Sea. Only 2 specimens known. (λεπτός, slender; οὐρά, tail.)

Macrurus lepturus, GILL & TOWNSEND, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17, 1897), 233, Bering Sea, southwest of Pribilof Islands, Albatross Station 3604, in 1,401 fathoms. (Type, No. 48767, U. S. Nat. Mus. Coll. *Albatross*.)

Macrurus dorsalis,* GILL & TOWNSEND, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17, 1897), 233, Bering Sea, southwest of Pribilof Islands, Albatross Station 3604, in 1,401 fathoms. (Type, No. 48768, U. S. Nat. Mus. Coll. *Albatross*.)

2961. MACROURUS ACROLEPIS, Bean.

Head about $4\frac{1}{2}$; depth at ventrals 7; eye $3\frac{1}{2}$ in head; snout 4; maxillary $2\frac{4}{5}$; mandible $2\frac{3}{5}$; pectoral about 2; ventral about $1\frac{2}{3}$. D. XI–111 +; A. 94 +; P. 20. Form of *M. berglax*, width of head $\frac{3}{4}$ its height; interorbital width $\frac{3}{4}$ eye; snout moderate, pointed. Origin of first dorsal from snout a distance 3 times length of upper jaw; base of first dorsal $3\frac{1}{2}$ in head, or about 3 times distance between dorsals; first dorsal spine very short, the second about $1\frac{2}{3}$ in head, serrate in front. Distance of anal from snout $2\frac{1}{2}$ times its length; distance of pectoral from snout slightly greater than head; distance of ventral origin from snout $\frac{1}{5}$ its length. Length 2 feet or more. Coasts of Vancouver Island, Washington and Oregon, in deep water, in 345 to 786 fathoms; common. A small specimen taken by us off Bogoslof Island. Our specimens have 11 rays in the first dorsal, not II, 11 or 13, as given by Bean. (ἄκρος, sharp; λεπίς, scale.)

Macrurus acrolepis, BEAN, Proc. U. S. Nat. Mus. 1883, 362, Straits of Juan de Fuca, near Neah Bay, Washington (Coll. James G. Swan, from the stomach of a fur seal); JORDAN, Cat. Fish. N. A., 131, 1885; GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 457.

Macrourus acrolepis, JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

2962. MACROURUS STELGIDOLEPIS, Gilbert.

Head $4\frac{3}{5}$; depth $5\frac{1}{2}$; eye small, $3\frac{1}{2}$ to 4 in head; snout $4\frac{1}{2}$. D. II, 10 or 11; A. 130; scales 155; 5 or 6 scales between lateral line and base of first dorsal. Body deep, the lower profile rapidly rising along anterior portion of base of anal, the tail thus abruptly becoming slender. Head short and deep; snout heavy, little produced, acute at extreme tip; infraorbital ridge not prominent on sides of head or snout, not continued backward on preopercle. A pair of narrow, transverse naked strips on upper surface of snout near tip, separated on each side by a single scale from the naked

*The following is Gill & Townsend's description of *Macrourus dorsalis*:

"Dorsal 15–120; anal 122; pectoral 21; ventral 9. Scales deciduous and rather small, diversiform, with small exposed surfaces; near the dorsal they have about 5 radiating spinigerous ridges, but below the lateral line these ridges are fewer and unarmed; snout short, projecting a considerable length beyond the eye and a little beyond the supramaxillary; median tubercle very prominent; connecting ridge well defined; infraorbital nearly vertical, with the ridge linear and near the orbit; teeth cardiform."

To this we add the following, from our examination of their type: Head $5\frac{2}{3}$; depth 7; eye $4\frac{2}{3}$; snout 4; interorbital width slightly greater than eye; maxillary $2\frac{2}{3}$; ventral fin $\frac{1}{5}$ longer than head, $4\frac{1}{2}$ in body; pectoral $1\frac{1}{2}$ in head. Dorsal spine strongly serrate, its length equal to that of head.

nostril fossa; a double series of scales intervenes between the nostrils and the orbit; lower side of snout wholly naked anteriorly, partly scaled laterally. Mouth large, overhung by premaxillaries for a distance about ½ diameter of orbit; premaxillaries in advance of nostrils; maxillaries reaching vertical from posterior margin of pupil, 2⅔ in head; snout about equaling interorbital width; barbel long, ⅝ orbit. Teeth in cardiform bands of equal width in both jaws, narrowed laterally, but not to a single series; anterior series in upper jaw enlarged, in lower jaw all the teeth of equal size. Preopercle broadly rounded, the angle but moderately produced, a narrow strip of the interopercle visible for its entire length; outer gill arch partially joined to cover, as usual; gill rakers obsolete; gill membranes united, forming a wide free fold across isthmus posteriorly. Scales without ridges, their exposed surfaces thickly beset with spines which are usually without definite arrangement; the marginal spine longest, thence decreasing in length to the base, about 40 present on each scale on middle of sides; scales on head crowded, the spines shorter and not directed backward as on the body; a rosette of short spines on tip of snout; no naked area between ventrals; mandible and gill membranes partly scaled; no considerable naked area in axil of pectorals. Dorsal inserted over base of pectorals, the length of its base slightly less than ½ the interspace between base of dorsals; second dorsal spine rather short and fragile, furnished anteriorly with a series of retrorse spinules, its length slightly exceeding ½ that of head, its tip not reaching origin of second dorsal; origin of anal fin well in advance of second dorsal; the vent unusually far forward, its distance from base of ventrals 2 to 2½ in its distance from anal fin; ventrals less widely separated than in *M. scaphopsis*, the outer ray produced, extending beyond front of anal; ventrals with 10 rays; pectorals with 22 to 24 rays; longest pectoral ray equals ½ head. Color very dark brownish, lighter on tail; lower side of head, breast, and abdominal region, including front of anal and base of pectorals, blue black; roof of mouth, valvular flap of membrane behind bands of teeth, gill membranes, and upper posterior portion of opercular lining, black; mouth and gill cavity otherwise white; peritoneum bright silvery, with little black specking; fins dusky. (Gilbert.) Coast of southern California. Two specimens, the longest 12 inches in length, from *Albatross* Station 2960, in 267 fathoms. (στελγίς, a scraper; λεπίς, scale.)

Macrurus stelgidolepis, Gilbert, Proc. U. S. Nat. Mus. 1890, 116, coast of southern California, at Albatross Station 2960, in 267 fathoms.

2963. MACROURUS CINEREUS, Gilbert.

(Pop-eye.)

D. II, 10 or 11; ventral 9; 7 scales between lateral line and first dorsal. Eye 3¾ to 4 in head; snout about 4, high and blunt, but little overlapping the mouth, terminating in a pointed prolongation of the median ridge, which bears at its tip a bony tubercle furnished with radiating ridges; nasal ridges terminating in shorter and smaller, but similar, tubercles, the outline between them concave; tip of snout overpassing the premaxillaries for ⅜ its length; eye very large and protuberant; mouth of

moderate size, the maxillary reaching vertical from hinder margin of orbit, equaling length of snout and ½ of eye. Teeth finely villiform, in each jaw, the outer series not at all enlarged, the mandibular band narrow. Barbel short and slender, its length less than ½ diameter of pupil; interorbital width ⅔ diameter of orbit, equaling length of snout; preopercle greatly expanded, much overlapping the interorbital below, leaving exposed only the extreme posterior angle. Gill membranes narrowly joined, with a posterior fold, free from the isthmus; gill rakers short, compressed, almost tubercular, 2 + 12. Origin of dorsal well behind base of pectorals; second dorsal spine long and filamentous, strongly spinous except on extreme base and tip; length of spine ⅝ to ⅔ head; base of first dorsal equaling diameter of orbit; interspace between dorsals ⅜ to ⅔ base of first dorsal; pectoral long and slender, equaling length of head behind anterior nostril opening, about as long as the filamentous outer ventral ray; vent immediately in front of anal origin. Scales on sides well imbricated, each with 7 to 9 parallel ridges which bear short sharp spines directed very obliquely backward; 7 scales between lateral line and base of first dorsal. Color uniform light grayish on body and fins, with the exception of the blackish pectorals and ventrals; sides of head silvery; mouth, gill cavity, and peritoneum brownish or purplish black; gill membranes and gular membrane dusky. (Gilbert.) Bering Sea; excessively abundant in the depths, where it outnumbers all other fishes. Numerous specimens from north of Unalaska Island, at *Albatross* Stations 3307 and 3329, in 1,033 and 399 fathoms; and the North Pacific, south of Unimak Island, *Albatross* Station 3340, in 695 fathoms. Our many specimens from near Bogoslof Island, in 664 fathoms. (*cinereus*, ashy gray.)

Macrourus cinereus, GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 457, near Unalaska and Unimak Islands, in 399 to 1,033 fathoms; JORDAN & GILBERT, Report Fur Seal Invest., 1898.

1007. CŒLORHYNCHUS, Giorna.

Cœlorhynchus, GIORNA, Mém. Ac. Sci. Turin, XVI, 178, 1803 ("*Cœlorhynche la ville*").
Krohnius, COCCO, Lettera al Sig. Augusto Krohn, Pesci del Mare de Messina, 1, 1844 (*filamentosus*; larva).
Paramacrurus, BLEEKER, Versl. Med. Ak. Wetensk. Amsterd. 1874, 103 (*australis*).
Oxymacrurus, BLEEKER, Versl. Med. Ak. Wetensk. Amsterd. 1874, 103 (*japonicus*).

This genus agrees with *Macrourus* in all essential respects, except that the small mouth is wholly below the long-pointed, sturgeon-like snout. Dorsal spine smooth in typical species, those with serrate spine having been lately separated under the generic name *Cœlocephalus*. (Gilbert & Cramer, Proc. U. S. Nat. Mus. 1896, 422) (*acipenserinus*). Species numerous. (κοῖλος, hollow; ῥύγχος, snout.)

a. Head large, 3⅓ in length; depth 7; eye ⅓ length of the long snout, 4 in head; dorsal spine moderate. OCCA, 2964.
aa. Head short, 4⅓ to 5 in length.
b. Body rather elongate, the depth 8 in length; eye as long as snout, 3 in head. CARMINATUS, 2965.
bb. Body less elongate, the depth 6⅓ to 6¾ in length.
c. Dorsal spine long; anal rays about 110; scales 124. CARIBBÆUS, 2966.
cc. Dorsal spine very short; anal rays 95; scales 98. SCAPHOPSIS, 2967.

2964. CŒLORHYNCHUS OCCA (Goode & Bean).

Head 3¼ in total length; depth 7 in total length; snout exceedingly elongate, nearly twice as long as diameter of eye; a black flap between nostrils; angle of mouth nearly reaching vertical from posterior margin of the orbit; ridge of head very strong and continuous from snout to angle of preopercle, having, also, strong supraocular and occipital ridges; eye nearly round, its horizontal diameter 4 in head and equal to interorbital space; ventral originating under middle of first dorsal, and extending to fourth ray of anal; distance from ventral origin to vent 3½ in length of head; second spine of dorsal weak and smooth, its length equal to postorbital part of head, its base slightly less than distance between first and second dorsals; squamation excessively rough, each scale bearing about 5 large spines besides many smaller ones, the median spine of the large series being much the largest; 5 rows of scales between origin of dorsal and lateral line, 19 from vent forward to lateral line and 12 backward; barbel ¼ as long as snout. This species has scales similar to those of *Macrourus berglax*, there being a strong median keel formed by series of spines, of which the last is the largest; surface of each scale also with about 4 or more lateral ridges formed by series of short spines. In a much larger example (U. S. Nat. Mus. No. 37334), measuring 18 inches in length, the lateral series of keels have greatly increased in number, the individual spines having become more prominent, so that the median keel has become less conspicuous than in the type. In the larger specimen referred to, the nakedness of the under surface of the head is even more pronounced than in the smaller, in which the under surface of the head beneath the suborbital and nasal ridge is almost entirely naked. The intermaxillary has a very short bone similar in structure and dentition to that of *Macrourus berglax*, that is to say, the intermaxillary teeth are in a rather broad villiform band, and the outer teeth are not enlarged; mandibulary teeth in a similar broad villiform band; mouth entirely inferior and small. Gill membranes attached across the isthmus, very little emarginate, and not deeply cleft; in the large example the gill membrane is attached to the isthmus and not deeply cleft, but there is a very narrow free margin behind. The gill rakers are very short, tubercular, and few in number, certainly not more numerous than in *M. berglax;* in the large example only 8 little tubercles can be seen on the first gill arch. Second spine of the dorsal in the type specimen is smooth, with the exception of 2 weak spines near its tip, but in the large example there is no trace of serrations on the dorsal spine. (Goode & Bean.) Length 450 mm. Gulf of Mexico and West Indies, in deep water. (*occa*, a harrow, from the rough scales.)

Macrurus occa, Goode & Bean, Proc. U. S. Nat. Mus. 1885, 595, Gulf of Mexico, Lat. 28° 34′ N., Long. 86° 48′ W., in 335 fathoms. (Type, No. 37334. Coll. *Albatross.*)
Cœlorhynchus occa, Goode & Bean, Oceanic Ichthyology, 400, figs. 332, 333, and 337, 1896.

2965. CŒLORHYNCHUS CARMINATUS (Goode).

Head about 5 in total length; depth 8; eye about 5 in head, equaling interorbital width; snout equaling eye or postorbital part of head; length of opercle about 2 in snout. Body less elongate than in *M. bairdii*. Snout

long, sharp, depressed, triangular, the lower surface more nearly parallel with the axis of body than in *M. bairdii;* lateral ridges more pronounced, continued in a straight line under eye and upon preopercle; strong horizontal ridges running from supraorbital margins to gill openings, parallel with subocular ridges; nostrils immediately in front of orbit; barbel very short. Teeth small, conical, somewhat recurved, arranged in villiform bands. Origin of first dorsal to snout 4⅓ times its base, its distance from anterior margin of orbit much less than length of head; first spine very short, hardly perceptible above the skin; second spine about 2 in head, slender and unarmed, when laid back its tip reaching to or beyond origin of second dorsal, the spines decreasing in length very gradually, the sixth being nearly as long as second, so that the fin is not so triangular as in *M. bairdii*; second dorsal beginning in a perpendicular from seventh anal ray; anal much higher than in *M. bairdii,* nearly equal to ½ interorbital width, its origin under eighteenth scale of lateral line, its longest rays as long as interorbital width; distance of pectoral from snout equaling twice its own length, which about equals longest dorsal spine; origin of pectoral below middle of depth of body and below level of middle of orbit, its tip not reaching origin of anal; insertion of ventrals behind pectoral, slightly in advance of first dorsal, its distance from snout greater than twice its length, the long filament not reaching anal. Color silvery gray. Length 250 mm. This species is extremely close to the common Mediterranean species, *C. cælorhynchus* (Risso), but the spines on the scales are a little larger. West Indies, Gulf of Mexico, and in the Gulf Stream in deep water; abundant; taken at many stations by the *Albatross,* the *Blake,* the *Fish Hawk,* and the *Challenger,* in 115 to 464 fathoms. (*carminatus,* from *carmen,* a wool card.)

Macrurus carminatus, Goode, Proc. U. S. Nat. Mus., III, 1880, 346 and 475, Gulf Stream off Rhode Island, Lat. 40° 02′ 54″ N., Long. 70° 23′ 40″, at Fish Hawk Station 871, in 115 fathoms (Type, No. 26007); Goode & Bean, Bull. Mus. Comp. Zool., X, No. 5, 196, 1883.

Macrurus (Cœlorhynchus) carminatus, Günther, Challenger Report, Deep-Sea Fishes, XXII, 129, pl. 5, fig. 13, 1887.

Cœlorhynchus carminatus, Goode & Bean, Oceanic Ichthyology, 398, fig. 336, 1896.

2966. CŒLORHYNCHUS CARIBBÆUS (Goode & Bean).

Head 4½ in total length; depth 6½ in total length. 1 D. II, 8; 2 D. at least 110; A. 110+; scales 6–124–15 or 16. Body normal in shape; scales moderate, strong, densely covered with minute spines, without enlarged median keel; interorbital area flat, its greatest width about 5 times in length of head; postorbital portion of head about 3 in head, and just as long as eye, which is oval, and 1⅔ as long as its vertical diameter. Snout long, thin, diaphanous, with acuminate point, its general form resembling that of *C. carminatus.* The nostrils close to the orbit, the posterior one much the larger. Teeth in each jaw in villiform bands, minute. Barbel slender and short, its length ⅓ that of eye. Maxillary extending to vertical through middle of pupil; upper jaw about 3 in head; mandible 2½; intermaxillary short. Outer series of teeth on intermaxillary and mandible not enlarged, the teeth not becoming uniserial. Gill membranes narrowly attached to the isthmus; gill rakers minute, tubercular, about 10 on

first arch. Suborbital ridge very strong, continued almost in a straight line by the lateral ridge of the snout; under surface of head, except chin and branchiostegal region, densely covered with small, spiny tubercles; a naked space on underside of snout, occupying almost entire distance from front of mouth to tip of snout, widest anteriorly, the greatest width 5 in snout; intermaxillary protractile in a vertical direction; mouth distinctly inferior. Origin of second dorsal over seventh anal ray, about an eye's diameter behind first dorsal; length of anal rays about 4 in head; origin of pectoral in front of first dorsal, its length 2 in head, its tip reaching fifth anal ray. Color silvery gray, with yellowish and lavender tints. Length 290 mm. Caribbean Sea north to the Gulf of Mexico, in deep water. (*caribbæus*, of the Caribbean Sea.)

Macrurus caribbæus, Goode & Bean, Proc. U. S. Nat. Mus. 1885, 594, Gulf of Mexico, at Albatross Station 2377, Lat. 29° 07′ 30″ N., Long. 88° 08′ W., in 210 fathoms (Type, No. 37333); Günther, Challenger Report, Deep-Sea Fishes, xxii, 124, note 3, 1887.
Cœlorhynchus caribbæus, Goode & Bean, Oceanic Ichthyology, 401, fig. 338, 1896.

2967. CŒLORHYNCHUS SCAPHOPSIS (Gilbert).

Depth 6¾ in total length; head 4⅙; scales 98. D. II, 8; A. ca. 95. Snout flattened, acute, the conspicuous infraorbital ridge forming a strong ridge along its sides, the two meeting at tip in a salient point; an evident keel extending from tip of snout to middle of interorbital area; supraorbital ridge dividing anteriorly, 1 branch running down in front of nostril, the other separating nostril fossa from orbit; between the ridges the head is covered with a soft, yielding integument, which is semitranslucent. Lower side of snout wholly naked below, and with a large naked area above on each side of tip; snout projecting beyond mouth for a distance equaling length of maxillary. Mouth of moderate size, the maxillary reaching vertical from posterior margin of pupil, 3⅓ in head. Teeth villiform, in a broad band in upper jaw, in a narrower band below, not reduced to a single series laterally in either jaw, and none of the teeth enlarged. Eye large, elliptical, equaling length of snout, 3¼ in head; interorbital width 4⅓. Barbel short, about ½ pupil; preopercular angle greatly produced backward, wholly concealing the interopercle, the strong infraorbital ridge failing to reach preopercular margin by only ⅛ diameter of pupil. Structure of gills as usual in this genus, the gill rakers obsolete; gill membranes broadly united, joined to isthmus, across which they form posteriorly a very narrow free fold. Besides the ridges already described on head, there are a pair on occiput, a pair from upper posterior margin of orbit to upper angle of gill opening, and a median ridge on nape reaching about halfway from occiput to dorsal. These ridges, as well as the interorbital space and the area between the occipital ridges, covered with scales compressed to a knife-like edge, which is provided with a single series of backward-directed spines; scales on infraorbital and rostral ridges bearing stellate spines or are similar to those on temporal region, sides of head, and body generally; scales on body large; 3 longitudinal series between lateral line and middle of first dorsal; each scale provided with a ridge bearing about 6 backward-directed spines, and from 2 to 4

pairs of lateral ridges also bearing spines, the lateral ridges sometimes extending the whole width of scales, sometimes confined to their basal portion; marginal spines longest; axil of pectoral naked, its base anteriorly with small cycloid scales; a naked, much depressed, elliptical area between bases of ventrals in all specimens; second dorsal spine smooth, weak, little exceeding length of soft rays, equaling length of snout and orbit; base of first dorsal $1\frac{1}{2}$ in interspace between dorsals, which is $2\frac{2}{3}$ in head; distance from front of anal to snout equaling $\frac{1}{3}$ total length; ventrals with the outer ray produced, about reaching front of anal; pectorals reaching beyond anal $\frac{1}{2}$ length of head; ventrals with 7 rays; pectorals with 15 to 17 rays. Color light olive brown, dusted with coarse black specks; axil of pectorals, belly, ventrals, and branchiostegal membranes blue black; lower side of head dusky; mouth anteriorly, including tongue and $\frac{1}{2}$ of palate, white, its posterior part and most of lining of gill cavity jet-black; inner lining of cheeks abruptly white; lower part of iris silvery; peritoneum silvery, with coarse dusky specks; vertical fins dusky, the anterior portion of anal black. Coast of southern California. Many specimens, the longest 12 inches long, from *Albatross* Station 3015, in 145 fathoms. (Gilbert.) (*σκάφη*, spade; *ὄψις*, face.)

Macrurus (Cœlorhynchus) scaphopsis, GILBERT, Proc. U. S. Nat. Mus. 1890, 115, Albatross Station 3015, coast of southern California, in 145 fathoms.

1008. TRACHONURUS, Günther.

Trachonurus, GÜNTHER, Challenger Report, Deep-Sea Fishes, XXII, 124, 1887 (*villosus*).

Scales not imbricated, separated by furrows, and densely covered with sharp spinules, so that the animal seems villous to the touch; dorsal spine smooth; dorsal much lower than anal; teeth in both jaws in villiform bands; snout obtuse, the mouth subinferior; suborbital ridge little developed. This genus is distinguished from *Cœlorhynchus* by the indistinct squamation. (*τραχύς*, rough; *οὐρά*, tail.)

2968. TRACHONURUS SULCATUS (Goode & Bean).

Head $7\frac{1}{2}$ in total length; depth about $9\frac{1}{2}$; eye $3\frac{2}{3}$ in head; snout 4 to $4\frac{1}{2}$. D. II, 8 or 9, the second of numerous low rays; A. 120; V. 7; P. 13; scales 7–175 or more—33. Barbel $2\frac{1}{2}$ to 2 in eye. Body elongate, rapidly contracted behind the abdomen; the tail long and whip-like. Scales moderate, strongly armed, each with 8 to 10 spinelets, irregularly placed, less numerous in the young, which feel bristly to the touch, separated by wide deep furrows; armature of head similar to that of body, but the scales upon snout, cheeks, and chin have very feeble spines. Interorbital area nearly flat, its length equaling diameter of eye or about 3 in head; postorbital part of head as long as eye; snout short, obtuse, scarcely overhanging the mouth; nostrils somewhat above level of middle of eye, the anterior one nearly upon the dorsal outline. Upper jaw with 2 series of teeth in villiform bands, the outer series slightly enlarged; teeth of lower jaw in a single series; maxillary reaching to vertical through hind margin of pupil in adult, shorter in younger individuals; length of upper jaw,

including maxillary, 3 in head; mandible 2 in depth of body; barbel 2 to 2½ in eye. Gill rakers very small, tubular, almost rudimentary, about 10 below angle of first arch; attachment of membrane to first arch very extensive, but free from isthmus; no pseudobranchiæ. First dorsal comparatively low, the first spine rudimentary, the second elongate and smooth; insertion of first dorsal immediately over or somewhat behind base of pectoral, its distance from snout 1⅓ in head, its base equal to snout, its longest spine, when laid down, reaching behind origin of second dorsal, or 1½ to 2 in head; second dorsal very low, its distance from first 3 to 4 in head, 32 rays in a distance equal to length of head; 22 in same distance of anal; anal much higher than second dorsal, yet very low, its longest ray equal to eye; distance of anal origin from snout 4⅜ in total length, or nearly under origin of second dorsal; pectoral inserted under or somewhat in front of origin of first dorsal, its length about 2 in head; ventral inserted behind vertical from end of base of first dorsal, extending to origin of anal, its length about equaling eye; vent about midway between origin of ventrals and anal. Color brown; abdomen and lower parts of head blackish in the young. West Indies and Gulf of Mexico, in deep water; taken both by the *Albatross* and the *Blake*. (*sulcatus*, furrowed.)

Coryphænoides sulcatus, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 596, Gulf of Mexico, at Albatross Station 2394, Lat. 28° 38′ 30″ N., Long. 87° 02′ W., in 420 fathoms (Type, No. 37335); GOODE & BEAN, Oceanic Ichthyology, 403, 1896.

Macrurus (Malacocephalus) sulcatus, GÜNTHER, Challenger Report, Deep-Sea Fishes, XXII, 169, 1887.

Trachonurus sulcatus, GOODE & BEAN, Oceanic Ichthyology, 410, fig. 343, 1896.

1009. LIONURUS, Günther.

Lionurus, GÜNTHER, Challenger Report, XXII, Deep-Sea Fishes, 124, 1887 (*filicauda*).

This genus is close to *Macrourus*, differing in the smooth, flaccid scales, and soft, cavernous skull, characters associated with its extreme bathybial degradation. (λεῖος, smooth; οὐρά, tail.)

a. Barbel minute, not ½ pupil; eye small, 5 in head; tail very slender.
FILICAUDA, 2969.

aa. Barbel moderate, 1⅔ in eye; eye 3¾ in head. LIOLEPIS, 2970.

2969. LIONURUS FILICAUDA (Günther).

D. 11; P. 20; V. 9; cæca 7. Snout considerably projecting beyond the mouth, pointed in the middle, twice as long as eye, which is unusually small, only ⅓ as wide as interorbital space. Mouth rather wide, extending beyond the center of the eye. Upper teeth villiform, in a very narrow band, those of mandible very small, biserial. Barbel minute. Preoperculum with the angle produced backward, broadly rounded and crenulated on the margin. Terminal portion of the tail prolonged into a long filament, more slender than in any of the other species. Bones of head soft. Scales of moderate size, thin, cycloid, and deciduous, 6 or 7 in a transverse series between the first dorsal spine and the lateral line; snout

and inferior half of the infraorbital region naked. Second dorsal spine slender, with the barbs in front very inconspicuous and sometimes entirely absent; distance between dorsal fins less than length of head; outer ventral ray produced into a small filament. Distance between vent and isthmus less than length of head. Head and trunk whitish, tail brownish, lower part of head and gill openings black. (Günther.) This species is clearly one of those in this family which extends to the greatest depths. The decrease in the size of the eye, the very soft bones, the comcomitant want of firmness in the structure of the scales, and the tail, which tapers into a very fine filament, indicate its abyssal abode. The scales are nearly all gone in all the specimens obtained. The species appears to be abundant in individuals, and has, like a true deep-sea fish, a wide distribution. (Günther.) Antarctic Ocean and deep seas off both coasts of South America. (*filum*, thread; *cauda*, tail.)

Coryphænoides (*Lionurus*) *filicauda*, GÜNTHER, Ann. and Mag. Nat. Hist., XX, 1878, 27, Deep seas on both sides of South America, in 1,375 to 2,650 fathoms.
Macrurus filicauda, GÜNTHER, Challenger Report, Deep-Sea Fishes, 141, pl. 34, fig. B, 1887.
Lionurus filicauda, GOODE & BEAN, Oceanic Ichthyology, 409, fig. 342, 1896.

2970. LIONURUS LIOLEPIS, Gilbert.

Head $4\frac{2}{3}$ in length; depth $6\frac{1}{5}$; maxillary nearly reaching vertical from posterior margin of orbit, $2\frac{4}{7}$ in head; eye $3\frac{3}{4}$; interorbital space concave, equaling snout, $4\frac{1}{8}$; barbel $\frac{2}{3}$ eye. D. II, 10. A. 120. Snout short and high, with well-marked lateral ridge, the extreme tip flattened; the median ethmoidal ridge is prominent, and the supraocular ridge is continued forward on the snout, meeting the lateral ridge in a projecting point. Top of snout wholly naked mesially, a narrow band of scales around each margin and in front. Lower side of head, including under side of snout, mandibles, gill membranes, and most of interopercles, naked; a very small patch of scales on posterior part of interopercle. Mouth moderate, the snout overhanging the premaxillaries for a distance equaling $\frac{1}{2}$ maxillary. Teeth in rather narrow cardiform bands in each jaw, not, however, forming single series laterally; the outer series in upper jaw only is enlarged; angle of preopercle little produced, not concealing the interopercle; infraorbital ridge not continued on to it. Gill membranes forming posteriorly a rather wide free fold across isthmus; outer gill arch joined to gill cover as usual in this genus. Scales small, everywhere cycloid, very deciduous, lost in most specimens; no spines developed, but occasionally can be seen traces of a median ridge and a pair of lateral ridges; about 6 or 7 series of scales between lateral line and base of first dorsal. Origin of first dorsal over or in advance of base of pectorals; base of first dorsal $1\frac{1}{8}$ to $1\frac{2}{3}$ in interval between dorsals; second dorsal spine usually smooth, occasionally with from 1 to 3 weak prickles near the middle; length of spine $1\frac{2}{3}$ in head. Origin of anal slightly behind first dorsal, the vent midway between base of ventrals and anal; ventrals short, the outer ray slightly produced, with from 10 to 12 rays; pectorals with 20 or 21 rays. Color very dark brown; snout, opercles, lower side of head, and abdominal region black or blue black; mouth and gill cavity

black; peritoneum dusky silvery. Coast of southern California. Many specimens taken at *Albatross* Station 2980, in 603 fathoms. (λεῖος, smooth; λεπίς, scale.) (Gilbert.)

Macrurus (*Lionurus*) *liolepis*, GILBERT, Proc. U. S. Nat. Mus. 1890, 117, coast of southern California, at Albatross Station 2980, in 603 fathoms.

Remotely related to the Scombriform fishes, and perhaps derived from the same ancestral stock as the *Trichiuridæ*, is the singular

Suborder TÆNIOSOMI.

(THE RIBBON-FISHES.)

This group is thus defined by Dr. Gill:

"Scapular arch subnormal, post-temporal undivided and closely applied to the back of the cranium, between the epiotic and pterotic, or upon the parietal; hypercoracoid perforate at or near the margin; cranium with the epiotics enlarged, encroaching backward and juxtaposed behind, intervening between the exoccipitals and supraoccipital; prootic and opisthotic represented chiefly by the enlarged prootic; suborbital chain imperfect; the copular bones separated by intervening cartilaginous elements; the hypopharyngeals styliform and parallel with the branchial arches; epipharyngeals in full number (4 pairs), and mostly compressed; the dorsal fin composed of inarticulate rays or spines, separable into lateral halves, and the ventrals (when present) subbrachial. A myodome may be present or absent, none being developed in the *Regalecidæ*, but 1 being distinct and supplemented by a dichost in the *Trachyteridæ*." (Gill.)

"The ribbon-fishes," says Günther, "are true deep-sea fishes, met with in all parts of the oceans, generally found when floating dead on the surface or thrown ashore by the waves. Their body is like a band, specimens of from 15 to 20 feet long being from 10 to 12 inches deep and about an inch or two broad at their thickest part. The eye is large and lateral; the mouth small, armed with very feeble teeth; the head deep and short. A high dorsal fin runs along the whole length of the back, and is supported by extremely numerous rays, its foremost portion, on the head, is detached from the rest of the fin, and composed of very elongate flexible spines. The anal fin is absent. The caudal fin (if preserved, which is rarely the case in adults) has an extra-axial position, being directed upward like a fan. The ventrals are thoracic, either compressed of several rays or reduced to a single long filament. The coloration is generally silvery, with rosy fins. When these fishes reach the surface of the water the expansion of the gases within their bodies has so loosened all the parts of their muscular and bony system that they can be lifted out of the water with difficulty only, and nearly always portions of the body and fins are broken and lost. The bones contain very little bony matter, and are very porous, thin, and light. At what depth ribbon fishes live is not known; probably the depths vary for different species; but although none has yet been obtained by means of the deep-sea dredge, they must be abundant at the bottom of all oceans, as dead fishes or fragments of them are frequently obtained.

Some writers have supposed from the great length and narrow shape of these fishes that they have been mistaken for 'sea serpents,' but as these monsters of the sea are always represented by those who have had the good fortune of meeting with them as remarkably active, it is not likely that harmless ribbon-fishes, which are either dying or dead, have been the objects described as 'sea serpents.'" (*ταινία*, ribbon; *σῶμα*, body.)

FAMILIES OF TÆNIOSOMI.

a. Ventral fins reduced each to a single long filament, thickened at the tip; anterior rays of dorsal produced; mouth small; caudal fin short or wanting. REGALECIDÆ, CCXVI.

aa. Ventral fins normally developed or else wanting.

b. Caudal fin short, fan-shaped, inserted at an angle with axis of body; the tail not much produced beyond it. TRACHYPTERIDÆ, CCXVII.

bb. Caudal fin short, the tail beyond it ending in a long filament, longer than rest of body. STYLEPHORIDÆ, CCXVIII.

Family CCXVI. REGALECIDÆ.

(OAR-FISHES.)

Body very elongated and compressed, the head oblong, the opercular apparatus well developed (the operculum extended backward, the subopercular obliquely behind it, and the interoperculum extended upward below the 2), the preorbital chain oblique and widest at the second bone; ventrals represented by single elongate rays, the cranium with the myodome atrophied and the dichost suppressed, the supraoccipital pushed forward by the extensive development of the epiotics which encroach forward on the roof as well as back and sides of the cranium, and with short ribs. (Gill.) Superficial characters are the very long dorsal, extending the whole length of the back and with the rays at the nape much produced; pectorals very short; caudal fin short or wanting; anal very low; head small; mouth very short; no air bladder; pyloric cæca numerous. One genus, with 2 or more species. Very large, surface-swimming fishes of the open seas; the great size, undulating motion and projecting mane causing them frequently to be taken for sea serpents. (*Regalecidæ*, Gill, Standard Nat. Hist., III, 1885; GILL, Amer. Nat. 1890, 482.)

1010. REGALECUS, Brünnich.

(OAR-FISHES.)

Regalecus, BRÜNNICH, Nya Sammlung, III, 414, 1788 (*glesne*).
Gymnetrus, BLOCH & SCHNEIDER, Syst. Ichth., 487, 1801 (*remipes*).
Xypterus, RAFINESQUE, Indice, 59, 1810 (*imperati*).

Characters of the genus included above. "It is not certain that there is more than 1 species of *Regalecus*, although, as the synonymy which follows clearly shows, various names have been suggested in connection with the comparatively few individuals which, during the past century and a half, have been captured in the North Atlantic. There appears to be consider-

able possibility of individual variation in proportions of height to length, and in the number of rays in the dorsal fin, but it is a fact well known to ichthyologists that constancy is not to be expected in forms in which the number of vertebræ and fin rays has been extended far beyond the normal average. It should also be said that most of the individuals studied have been in very imperfect condition, and also that in many instances the observations have been made by untrained observers, so that it seems doubtful whether there is really more than 1 species to be assigned to the Atlantic fauna. At all events, Günther, Collett, Lütken, and Day agree in the idea that it is impossible to discriminate between the forms already described, and we follow their lead in considering them all, for the present, as a single species. It is not impossible, of course, that, should better material be obtained, it may be desirable to separate the group into more subspecies, but until this shall be done discrimination leads to confusion rather than to definite knowledge. The fishes belonging to the genus *Regalecus* are very remarkable, not only on account of their peculiar appearance and structure, but because of their enormous size. They have been known to attain the length of 20 feet, and it is more than probable that they grow very much longer, and that many of the creatures popularly identified with the "sea serpent" are only large individuals of this type. Indeed, it seems quite safe to assign to this group all the so-called "sea serpents" which have been described as swimming rapidly near the surface, with a horse-like head raised above the water, surmounted by a mane-like crest of red or brown. The individual which came ashore at Hungry Bay, in Bermuda, in 1860, and which was about 17 feet long, was described by the people who saw it before its capture as being very much larger, and as having a head of an immense horse with a flaming red mane." (Goode & Bean.) (*rex*, king; *halec*, herring. The species have long been known as "king of the herrings," as have those of *Trachypterus*.)

2971. REGALECUS GLESNE (Ascanius).

(Oar-Fish; Sea Serpent.)

Head 16 to 20; depth 12 to 24; eye 4 to 6 in head; snout short, truncated. D. 275 to 400; P. 11 to 14; V. I. Body very elongate. Cleft of mouth vertical, the upper jaw very protractile; jaws minute or absent. Anterior 8 to 15 rays of dorsal forming an elevated crest, sometimes in 2 parts, the posterior rays with membranaceous tips; each ventral ray with a lobate membranaceous tip; skin with numerous bony tubercles; lateral line placed low. Color silvery gray, with a few spots or streaks of darker hue, most numerous anteriorly. Günther (Challenger Report, XXII, 73 to 76) has in the most painstaking manner brought together a list of the specimens taken in the North Atlantic, as far as they are known to science. He mentions 14 known upon the Scandinavian coasts from 1740 to 1852; 19 on the British coasts from 1759 to 1884; 1 in the Mediterranean (he states, however, that about ½ a dozen specimens have been observed in the Mediterranean); 1 in the Bermudas; 3 at the Cape of Good Hope; 1 in the Indian Ocean, and 5 off the coast of New Zealand. He calls attention to

the fact that of those observed on the British and Scandinavian coasts 4 were observed in the month of January, 5 in February, 8 in March, 2 in April, 1 in May, 1 in June, 1 in July, 2 in August, 1 in September, and 1 in October. He also calls attention to the fact that by far the greater proportion of their capture, in the Northern Hemisphere at least, is in the stormy season. This agrees with what we know of the capture of *Trachypterus*, which likewise seems to be brought to the surface only by great commotions of the ocean. The popular name of *Regalecus* is oarfish, in allusion to the blade-like expansion of the extremities of the 2 ventral fins. *Regalecus* is also called in the books the "king of the herrings." Strangely enough, no representative of this genus has been found on the coast of North America. Günther is of the opinion that the distribution of this fish in the depths of the sea is the same as that of *Trachypterus*. The similarity in their geographical distribution is quite remarkable. (Goode & Bean.) (Eu.) (*glesne*, from "Glesnæs," a farm at Glesvær, near Bergen, where the type of the species was taken.)

Spada marina, IMPERATO, Hist. Nat., 679, 687, 1599, **Naples.**
Regalecus glesne, ASCANIUS, Icones Rerum Nat., II, pl. 11, about 1788, **Glesvær, Norway.**
Ophidium glesne, ASCANIUS, Nya Saml. Vid. Selsk. Skrivt., III, 419, 1788.
Regalecus remipes, BRÜNNICH, Nya Saml. Vid. Selsk. Skrivt., III, 1788, 414, taf. B., figs. 4, 5; WALBAUM, Artedi Piscium, III, 647, tab. 3, fig. 4, 1792.
Cepola gladius, WALBAUM, Artedi Piscium, III, 617, 1792.
Gymnetrus hawkenii, BLOCH, Ichthyol., XII, 88, 425, 1792.
Gymnetrus grillii, LINDROTH, Vet. Akad. Handl. 1798, 291, pl. 8.
Gymnetrus ascanii, SHAW, Gen. Zool., IV, 197, 1803; after Ascanius.
Xypterus imperati, RAFINESQUE, Indice, 59, 1810; after Ferrante Imperato.
Gymnetrus longiradiatus, RISSO, Eur. Mérid., III, 296, 1826, **Nice.**
Gymnetrus telium, CUVIER & VALENCIENNES, Hist. Nat. Poiss., X, 361, pl. 299, 1834, **Nice.**
Regalecus banksii, CUVIER & VALENCIENNES, Hist. Nat. Poiss., X, 365, 1834, **Filey Bay, Yorkshire.**
Gymnetrus capensis, CUVIER & VALENCIENNES, Hist. Nat. Poiss., X, 376, 1834, **Cape of Good Hope.**
Regalecus glesne, ASCANIUS, Icones Rerum Naturalium, 1806, pl. 11; LACÉPÈDE, Hist. Nat. Poiss., II, 214, 215, 1800; GOODE & BEAN, Oceanic Ichthyology, 480, fig. 395, 1896.
Gymnetrus remipes, BLOCH & SCHNEIDER, Syst. Ichth., 482, tab. 88, 1801; YARRELL, Brit. Fishes, Ed. 2, I, 223, and Ed. 3, II, 301.
Gymnetrus glesne, CUVIER & VALENCIENNES, Hist. Nat. Poiss., X, 366.
Gymnetrus gladius, CUVIER & VALENCIENNES, Hist. Nat. Poiss., X, 352, pl. 298, 1835.
Regalecus gladius, GÜNTHER, Cat., III, 308, 1861.

Family CCXVII. TRACHYPTERIDÆ.

(THE KING OF THE HERRINGS.)

Body moderately elongate, strongly compressed, naked, the skin smooth or prickly. Lateral line present. Head short; the mouth rather small, terminal, with feeble teeth; premaxillaries protractile; opercles unarmed; opercular apparatus abbreviated (the operculum extended downward, the suboperculum below it, and the interoperculum contracted backward and bounded behind by the operculum and suboperculum); the cranium with a myodome and dichost, the supraoccipital continued behind into a prominence; the epiotics confined to the sides and back of the cranium, and without ribs. Eye large, lateral; branchiostegals 6; gill membranes

separate, free from the isthmus; gills 4, a slit behind the fourth. Pseudobranchiæ well-developed, in a pouch formed by a fold of the mucous membrane. Dorsal fin single, extending from the head to the tail, its rays all technically spinous, being neither articulated nor branched, but all very soft, flexible, and fragile; anal fin wanting; pectorals short; ventrals thoracic, the rays elongate, less than I, 5 in number, usually atrophied in the adult; caudal fin either rudimentary or else divided into 2 parts, the upper and larger fan-shaped, directed obliquely upward from the slender tip of the tail. Bones very soft, the muscles little coherent. Pyloric cæca very numerous. Vertebræ in large number. Deep-sea fishes, often of large size, found in most warm seas. Their extreme fragility renders them rare in collections, and the species are little known. One genus; species about 12. The ribbon-fishes are well known in the Eastern Atlantic and the Mediterranean, and have even been found as far west as Madeira [and Cuba]. Some few representatives have been found on the west coast of South America, and 1 or 2 examples have been taken in New Zealand. They are generally admitted to be true deep-sea fishes, which live at very great depths, and are only found when floating dead on the surface or washed ashore by the waves. Almost nothing is known of their habits except through Nilsson's observations in the Far North. This naturalist, as well as Olafsen, appears to have had the opportunity of observing them in life. They say that they approach the shore at flood tide on sandy shelving bottoms, and are often left by the retreating waves. Nilsson's opinion is that its habits resemble those of the flat fishes, and that they move with one side turned obliquely upward, the other toward the ground; and he says that they have been seen on the bottom in 2 or 3 fathoms of water, where the fisherman hook them up with the implements employed to raise dead seals, and that they are slow swimmers. This is not necessarily the case, however, for the removal of pressure and the rough treatment by which they were probably washed upon the shore would be demoralizing, to say the least. *Trichiurus*, a fish similar in form, is a very strong, swift swimmer, and so is *Regalecus*. Whether or not the habits of *Trachypterus arcticus*, on which these observations were made, are a safe guide in regard to the other forms is a matter of some doubt, but it is certain that they live far from the surface, except near the Arctic Circle, and that they only come ashore accidentally. They have never been taken by the deep-sea dredge or trawlnet, and, indeed, perfect specimens are very rare, the bodies being very soft and brittle, the bones and fin rays exceedingly fragile. A considerable number of species have been described, but in most instances each was based upon 1 or 2 specimens. It is probable that future studies may be as fruitful as that of Emery, who, by means of a series of 23 specimens, succeeded in uniting at least 3 of the Mediterranean species, which for half a century or more had been regarded as distinct. The common species of the Eastern Atlantic, *Trachypterus atlanticus*, is not rare, 1 or more specimens, according to Günther, being secured along the coast of northern Europe after almost every severe gale. We desire to quote the recommendation of Dr. Günther, and to strongly urge upon any one who may be so fortu-

nate as to secure 1 of these fishes, that no attempt should be made to keep it entire, but that it should be cut into short lengths and preserved in the strongest spirits, each piece wrapped separately in muslin. (Goode & Bean.)

1011. TRACHYPTERUS, Gouan.

(King of the Herrings.)

Trachypterus, Gouan, Hist. Poiss., 104, 153, 1770 (*trachypterus*).
Bogmarus, Bloch & Schneider, Syst. Ichth., 518, 1801 (*islandicus* = *arcticus*).

Body elongate, compressed, ribbon-shaped, the dorsal fin extending the entire length of the back. Anal absent; each ventral well developed, if present, but sometimes absent. Caudal present and placed for the most part above the longitudinal axis of the body. No air bladder. Pyloric appendages numerous. Ventrals appearing to be absent in some individuals, but Day calls attention to the fact that most of the specimens of *T. arcticus* taken along the coast of Great Britain had no ventrals. In the very young, as has been shown by Emery, the fin rays commence to grow when it is about 6 mm. long, and continue to lengthen until it is about 24 mm. long, after which a partial shortening takes place. Ventrals very elongate in the young, and the caudal rays much longer than in the grown fish. Young individuals (from 2 to 4 inches) are not rarely met with near the surface; they possess the most extraordinary development of fin rays observed in the whole class of fishes, some of them being several times larger than the body, and provided with lappet-like dilatations. There is no doubt that fishes with such delicate appendages are bred and live in depths where the water is absolutely quiet, as a sojourn in the disturbed water of the surface would deprive them at once of organs which must be of some utility for their preservation. (Goode & Bean.) (τραχύς, rough; πτερόν, fin.)

a. Color bright metallic silvery, a jet-black blotch at base of dorsal; 3 dark spots on side, 2 smaller ones on belly; anterior profile, snout, and tip of mandible, jet black; caudal and ventral fins carmine red in life. REX-SALMONORUM, 2972.

aa. Color shining leaden gray; no black. TRACHYURUS, 2973.

2972. TRACHYPTERUS REX-SALMONORUM, Jordan & Gilbert.

(King of the Salmon.)

Head 8⅓; cross depth at nape 8. D. V–170; C. 8; V. 6; P. 11. Body long and slender, closely compressed and ribbon-shaped, as usual in the genus. Head short, deeper than long, the anterior profile steep and nearly straight to the base of the nuchal crest; dorsal fin beginning on the top of nuchal crest, which is directly over the second third of the diameter of eye; height of crest slightly more than diameter of eye, the latter greater than length of snout, and 3 in head. Mouth oblique; maxillary rugose and very broad, its width ½ its length; length of lower jaw greater than length of snout, 2⅔ in head, its angle under the front of the orbit. Opercular bones rugose, entirely covering the gills. Premaxillary covered with minute and feeble teeth, in addition to which in this specimen are 3 canines, 2 on one side and 1 on the other, directed very obliquely

backward. On the side having 2 canines, 1 is placed directly behind the other; lower jaw with 3 strong canines on one side and 2 strong and 1 weak canine on the other, all directed obliquely backward and inward. Dorsal fins slightly connected at base; the filamentous rays of the first dorsal not quite twice the length of head; ventrals inserted just below axil of pectorals, filamentous, about ½ longer than head; pectorals ⅕ longer than eye; caudal rays simple to near tip, where is sometimes a single fork, the longest filamentous rays about 3 times length of head; dorsal fin much lower than body, longest rays of second dorsal nearly ⅔ length of head; a series of spinules along base of dorsal, 1 pair for each ray. Lateral line well developed, with a series of small inconspicuous plates, each of which has a minute central prickle. Lower part of the body thickly beset with small spinous tubercles; rest of the skin naked; rays of all the fins accompanied by a series of small prickles. Coloration everywhere bright metallic silvery, an oblong jet-black blotch a little longer than eye lying close along base of dorsal and beginning 1½ diameters of eye behind eye; 3 larger spots, dusky but not black, lying behind this along side between lateral line and dorsal fin; 2 smaller dusky spots on belly, the one just behind base of ventrals, the other under the second of the 4 spots of back; these spots, except the first one mentioned, are all diffuse and a little less than twice the diameter of eye in length and about twice as long as deep; anterior profile below crest, including front of snout and tip of mandible, jet-black; caudal and ventral fins carmine red in life; other fins unmarked. Length 17 inches. This species bears some resemblance to *Trachypterus altivelis* described by Kner from Valparaiso. The latter species has, however, the nuchal crest much lower and farther back, the first dorsal and the ventrals much lower, the second dorsal fin higher, the skin rougher, the 4 black spots different in size and position from those found in our specimen, and the caudal rays divided near the base. It is probable that the 3 specimens of *Trachypterus* mentioned in the Synopsis of the Fishes of North America, p. 619, and referred with doubt to *Trachypterus altivelis*, really belong to the present species. Four specimens known; 1 from Santa Cruz, California, taken by Dr. C. L. Anderson; 2 from the Straits of Fuca, taken by Mr. J. G. Swan, and the type, obtained by a fisherman (Mr. Knox) in the open sea outside the bay of San Francisco. According to Mr. Swan the species is known by the Makah Indians west of the Straits of Fuca as "king of the salmon," and its destruction is believed to have a baneful influence on the salmon fishing. "When the king of the salmon is killed the salmon will cease to run." (*rex*, king; *salmonorum*, of the salmon.)

?Trachypterus altivelis, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 52; JORDAN & GILBERT, Synopsis, 618, 1883; specimen from Santa Cruz; not of KNER.

Trachypterus rex-salmonorum, JORDAN & GILBERT, Proc. Cal. Ac. Sci. 1894, 145, pl. 9, open sea outside Bay of San Francisco. (Type, No. 1382, L. S. Jr. Univ. Coll. Mr. Knox.)

2973. TRACHYPTERUS TRACHYURUS, Poey.

D. 82; P. 15; V. 6. Eye 2½ in head, high, as long as snout. Mouth almost vertical. Bones of head thin as paper. Lateral line a little concave on the middle of trunk. No scales; pectorals small; ventrals behind

pectorals, very long, reaching past vent, which is at second third of length, including caudal; dorsal almost as high as body, without plume in front. Vertebræ 36 + 18. Shining leaden gray, a silvery band produced by the vertebral column showing through. Cuba. (Poey); not seen by us. (*τραχύς*, rough; *οὐρά*, tail.)

Trachypterus trachyurus, POEY, Memorias, II, 420, 1861, Cuba.

Family CCXVIII. STYLEPHORIDÆ.

Body elongate, compressed, ribbon-shaped; the dorsal extending from head nearly to end of tail; tail terminating in an exceedingly long, cord-like appendage, about twice as long as head. Anal absent; ventrals absent; caudal erected upward, having its rays connected by a rather firm membrane. Snout produced; mouth small, toothless; maxillary bones small, short, hidden behind premaxillaries; mandible long, extending far behind the eye. Eye large, turned forward; suborbital very large, covering nearly the whole of cheek and extending backward behind eye. Opercles small. Gill openings wide; gills 4. Vent premedian. Branchiostegals 4. (Goode & Bean.) This family is based on a single specimen obtained in the West Indies in 1790 and preserved in the British Museum. The relations of the fish are uncertain, and it may not belong to the *Tæniosomi*. Its nearest relations are, however, apparently with *Trachyterus*.

(*Stylephoridæ*, SWAINSON, Nat. Hist. Class'n Fishes, II, 47, 1839.)

1012. STYLEPHORUS, Shaw.

Stylephorus, SHAW, Trans. Linn. Soc. Lond., I, 1791, 90 (*chordatus*).

Characters of the genus included above. (*στῦλος*, a style or projecting part; *φορέω*, to bear.)

2974. STYLEPHORUS CHORDATUS, Shaw.

Head 6; depth 5. D. 110; C. 6; P. 13; B. 4. Snout produced, subcylindrical; mouth small and toothless; maxillary bones small, short, and hidden behind the intermaxillaries; mandible long, extending far behind eye; eyes large, close together, directed forward toward snout; suborbital very large, covering nearly the whole of cheek, and extending backward behind eye; opercles small; gill openings very wide; gills 4. Vent situated before middle of total length; pectorals pointed, directed upward, about ½ as long as head; dorsal extending from head nearly to end of tail; caudal directed upward, and having its rays connected by a rather firm membrane, the tail terminating in a narrow band-like appendage about twice as long as body. Color uniform silvery. (Günther.) This remarkable form is known only from a single specimen, 11 inches long, with the caudal appendage 22 inches in length, which was taken in the Atlantic, between Cuba and Martinique, about the year 1790, and is now in the British Museum. It is undoubtedly an inhabitant of great depths. (Goode & Bean.) (*chordatus*, with a chord; from *χόρδη*, string.)

Stylephorus chordatus, SHAW, Trans. Linn. Soc. London, I, 1791, 90, pl. 6, between Cuba and Martinique; SHAW, Zool., IV, 87; SHAW, Naturalists' Miscellany, VIII, pl. 274; BLAINVILLE, Journ. Phys., LXXXVII, 60, pl. 1, fig. 1; CUVIER & VALENCIENNES, Hist. Nat. Poiss., X, 381; GÜNTHER, Cat., III, 306, 1861; GOODE & BEAN, Oceanic Ichthyology, 482, pl. 66, figs. 393 and 394, 1896.

Suborder HETEROSOMATA.

(The Flatfishes.)

"Cranium posteriorly normal; anteriorly with twisted vertex, to allow 2 orbits on the same side, or 1 vertical and 1 lateral; basis cranii not quite simple. Dorsal fin long, of jointed rays; superior pharyngeals 4, the third longest, much extended forward, the inferior separate." (Cope.) This suborder includes the 2 families, *Pleuronectidæ* and *Soleidæ*. Its nearest relationship is probably with the *Gadidæ*, although the developed pseudobranchiæ and the thoracic ventral fins, indicate an early differentiation from the anacanthine fishes. In the very young fishes the 2 sides of the body are alike and the eyes are 1 on each side, with normal cranium. (ἕτερος, different; σῶμα, body.) (*Anacanthini pleuronectoidei*, Günther, Cat., IV, 399, 504.)

Families of Heterosomata.

a. Preopercular margin more or less distinct, not hidden by the skin and scales of the head; eyes large, well separated; mouth moderate or large; teeth present. Pleuronectidæ, CCXIX.

aa. Preopercular margin adnate, hidden by the skin and scales of the head; eyes small, close together; mouth very small, much twisted; teeth rudimentary or wanting. Soleidæ, CCXX.

Family CCXIX. PLEURONECTIDÆ.*

(The Flounders.)

Body strongly compressed, oval or elliptical in outline; head unsymmetrical, the cranium twisted, both eyes being on the same side of the body, which is horizontal in life, the eyed side being uppermost and colored, the blind side lowermost and usually plain. In the very young fish the bones of the head are symmetrical, 1 eye on each side, and the body is vertical in the water. In most species the cranium becomes twisted, bringing the upper eye over with it. Eyes large, well separated. Mouth small or large, the dentition various, the teeth always present; premaxillaries protractile; no supplemental maxillary bone; pseudobranchiæ present. Gills 4, a slit behind the fourth; lower pharyngeals separate; no air bladder; preopercle with its margin usually distinct, not wholly adnate or hidden by the skin of the head; vent not far behind head, the viscera confined to the anterior part of the body. Scales various, rarely absent, usually small. Lateral line usually present, extending on the caudal fin, sometimes duplicated or wanting. Dorsal fin long, continuous, of soft rays only, beginning on the head; anal similar, shorter; caudal various, sometimes coalescent with dorsal and anal; pectorals inserted rather high, rarely wanting; ventrals under the pectorals, usually of several soft rays, one of them sometimes wanting. Fishes mostly carnivorous, inhabiting sandy bottoms in all seas, some species ascending rivers. Many of them are important food-fishes. Genera about 55; species

* For complete synonymy and descriptions of the American species of this family of fishes see "A review of the flounders and soles (*Pleuronectidæ*) of America and Europe," by David Starr Jordan and David Kop Goss, in Report United States Fish Comm. for 1886, 225-342, pls. 1 to 9, first published in 1889.

nearly 500. The group "*Bibroniidæ*" recently recognized by some of the Italian ichthyologists as a separate family ("*Bibronidi*"), is composed entirely of larval forms in the early stages of their development. In this condition the eyes are symmetrical and the body translucent. Several generic names have been given to these peculiar forms (*Peloria, Bibronia, Coccolus, Charybdia, Bascanius, Delothyris*), but, of course, these genera can have no permanent place in the system. *Peloria* has been shown by Dr. Emery to be the young of *Platophrys*. The others seem to belong to the *Cynoglossinæ* or to some allied group, but we are not yet certain as to the correct identification of any of them. We recognize among the *Pleuronectidæ* 6 subfamilies—*Hippoglossinæ, Psettinæ, Samarinæ, Pleuronectinæ, Oncopterinæ*, and *Pelecanichthyinæ*. These subfamilies are natural groups and are in most cases easily distinguished, although some few aberrant genera exist, which serve as links joining one group to another. Thus *Isopsetta* of the *Pleuronectinæ* is certainly a near ally of *Psettichthys*, which is as certainly a genuine member of the *Hippoglossinæ*. The *Hippoglossinæ* and the *Pleuronectinæ* are largely arctic in their distribution, few of the former group and none of the latter extending into the Tropics. The *Oncopterinæ* seem to take the place of the *Pleuronectinæ* in antarctic waters, but the species of this group are few in number. The *Psettinæ* and the soles are, on the other hand, essentially warm-water fishes, their representatives in the north being comparatively few. The *Samarinæ* are few in number and belong to the East Indian fauna, and the single species of *Pelecanichthyinæ* belongs to the bassalian fauna of the Pacific. As the tropical *Hippoglossinæ* and all the *Psettinæ* are sinistral species, the eyes and color being on the left side of the body, it follows that the tropical flounders are nearly all left-sided species, while those of arctic and antarctic waters are chiefly dextral species, the eyes and color on the right. The *Hippoglossinæ* are the most generalized of the flatfishes. From the northern representatives of this group, the allies of *Hippoglossoides*, the *Pleuronectinæ*, are certainly descended. The *Psettinæ* are apparently derived from ancestors of the type of *Paralichthys*. The soles show closest affinities with the *Psettinæ*, from ancestors of which group they have become degraded. Very remarkable is the relation between the number of vertebræ and the geographical distribution of the various species. It has been already noticed by Dr. Gill, Dr. Günther and others that in some groups of fishes northern representatives have the number of vertebræ increased. In no group is this more striking than in the flounders, as the following table, showing the numbers of the vertebræ in various species, will clearly show. The numbers inclosed in brackets are copied from Dr. Günther; the others represent our own count of specimens.

Numbers of vertebræ in flounders.

I.—HIPPOGLOSSINÆ.

Hippoglossus hippoglossus	16 + 34 = 50
Atheresthes stomias	12 + 37 = 49
Hippoglossoides platessoides	13 + 32 = 45
Lyopsetta exilis	11 + 34 = 45
Eopsetta jordani	11 + 32 = 43

Psettichthys melanostictus	11 + 29 = 40
Paralichthys oblongus	11 + 30 = 41
Paralichthys dentatus	10 + 30 = 40
Paralichthys lethostigmus	10 + 27 = 37
Paralichthys albiguttus	10 + 27 = 37
Paralichthys californicus	10 + 25 = 35
Xystreurys liolepis	12 + 25 = 37
Ancylopsetta quadrocellata	9 + 26 = 35

II.—Pleuronectinæ.

Glyptocephalus zachirus	13 + 52 = 65
Glyptocephalus cynoglossus	[58]
Microstomus pacificus	12 + 40 = 52
Microstomus kitt	[13 + 35 = 48]
Parophrys vetulus	11 + 33 = 44
Pleuronectes platessa	[14 + 29 = 43]
Isopsetta isolepis	10 + 32 = 42
Lepidopsetta bilineata	11 + 29 = 40
Limanda limanda	[40]
Liopsetta glacialis	13 + 27 = 40
Pleuronichthys decurrens	14 + 26 = 40
Pleuronichthys verticalis	13 + 25 = 38
Flesus glaber	11 + 26 = 37
Flesus flesus	[12 + 24 = 36]
Pseudopleuronectes americanus	10 + 26 = 36
Hypsopsetta guttulata	11 + 24 = 35
Platichthys stellatus	12 + 23 = 35

III.—Psettinæ.

Monolene sessilicauda	[43]
Lepidorhombus whiff-iagonis	[11 + 30 = 41]
Citharichthys sordidus	11 + 29 = 40
Platophrys lunatus	9 + 30 = 39
Arnoglossus laterna	10 + 28 = 38
Arnoglossus grohmanni	10 + 28 = 38
Zeugopterus punctatus	[12 + 25 = 37]
Platophrys ocellatus	10 + 27 = 37
Lophopsetta maculata	11 + 25 = 36
Bothus rhombus	12 + 24 = 36
Syacium papillosum	11 + 25 = 36
Citharichthys arctifrons	10 + 26 = 36
Syacium micrurum	10 + 25 = 35
Phrynorhombus regius	10 + 25 = 35
Citharichthys spilopterus	10 + 24 = 34
Citharichthys macrops	10 + 24 = 34
Etropus microstomus	10 + 24 = 34
Etropus crossotus	10 + 24 = 34
Azevia panamensis	33
Psetta maxima	12 + 19 = 31

The subdivision of the flounders into genera leaves room for considerable variety of opinion. Most of the species are well defined and easily recognized, but they do not fall readily into generic groups unless we regard almost every well-marked species as the type of a distinct genus. A natural result of an attempt at sharply defining the genera is to reach what seems an extreme degree of generic subdivision. On the other hand, attempts to unite these smaller groups to form larger ones often leave these larger ones at once unnatural and ill-defined.

It will probably appear to some that the process of generic subdivision has been in this paper carried too far. It is possible that this is true, but the arrangement which we have adopted seems to bring out the relations of the different forms better than can be done by a more conservative view of the genera. (*Pleuronectidæ*, Günther, Cat., IV, 1862.)

SUBFAMILIES OF PLEURONECTIDÆ.

A. Ventral fins symmetrical, similar in position and in form of base, the ventral of the colored side not extended along the ridge of the abdomen.
  a. Mouth nearly symmetrical, the dentition nearly equally developed on both sides, the gape usually but not always wide. (Halibut tribe.) HIPPOGLOSSINÆ, I.
  aa. Mouth unsymmetrical, the jaws on the eyed side with nearly straight outline, the bones on the blind side strongly curved; teeth chiefly on the blind side.
    b. Eyes and color on the right side (with occasional exceptions). (Flounder tribe.) PLEURONECTINÆ, II.

AA. Ventral fins unsymmetrical, dissimilar in position and usually also in form, the ventral fin of the eyed side being extended along the ridge of the abdomen. Eyes and color on the left side. (Turbot tribe.) PSETTINÆ, III.

ANALYSIS OF GENERA.

I. HIPPOGLOSSINÆ.

(HALIBUT TRIBE.)

Large-mouthed flounders with the ventral fins symmetrical.—Mouth symmetrical, the jaws and the dentition nearly equally developed on both sides; gape usually wide, the maxillary more than $\frac{1}{3}$ length of head. Lower pharyngeals narrow, usually with but 1 or 2 rows of sharp teeth; teeth in jaws usually acute. Eyes large; edge of preopercle free. Pectoral and ventral fins well developed, *the ventral fins similar in position* and in form of base, the ventral fin of the eyed side not being attached along the ridge of the abdomen. Septum of gill cavity without foramen.

a. Vertebræ and fin rays much increased in number (the vertebræ about 50; dorsal rays about 100, anal rays about 85); body comparatively elongate; caudal fin lunate; lateral line simple; anal spine mostly obsolete. Dextral species, arctic in distribution. (Genera allied to *Hippoglossus*.)
  b. Large teeth in both jaw arrow-shaped, biserial, some of them depressible; upper eye with vertical range; gill rakers short; scales deciduous, ciliated; lateral line without arch; flesh soft. Vertebræ (in *A. stomias*) 12 + 37 = 49. ATHERESTHES, 1013.
  bb. Large teeth not arrow-shaped, biserial above, uniserial below; scales very small, cycloid; gill rakers long and slender; eyes strictly lateral.
    c. Lateral line without anterior arch; lower pharyngeal teeth uniserial. REINHARDTIUS, 1014.
    cc. Lateral line with an anterior arch; lower pharyngeal teeth biserial; vertebræ (in *H. hippoglossus*) 16 + 34 = 50. HIPPOGLOSSUS, 1015.

aa. Vertebræ and fin rays in moderate number (vertebræ less than 46; dorsal rays fewer than 95; anal rays fewer than 75); caudal fin double truncate or rounded, the median rays longest.
  d. Lateral line without distinct anterior arch; vertebræ 40 to 46; body normally dextral;* caudal peduncle distinct; scales ciliated; anal spine usually strong. Species of subarctic distribution. (Genera allied to *Hippoglossoides*.)

* Frequently sinistral in *Hippoglossoides elassodon*.

e. Lateral line simple, without accessory dorsal branch; teeth sharp, those of lower jaw uniserial; dorsal beginning above eye.

f. Teeth in the upper jaw biserial.

g. Scales comparatively large, thin, and deciduous; lateral line 70; body slender, the flesh soft; vertebræ (in *L. exilis*) 11 + 34 = 45. LYOPSETTA, 1016.

gg. Scales small and adherent; lateral line 96; body robust, the flesh firm; vertebræ (in *E. jordani*) 11 + 32 = 43. EOPSETTA, 1017.

ff. Teeth in the upper jaw uniserial; scales small and flesh firm; vertebræ (in *H. platessoides*) 13 + 32 = 45. HIPPOGLOSSOIDES, 1018.

ee. Lateral line with an accessory dorsal branch; scales small, firm, ctenoid; dorsal fin beginning before the eye; teeth sharp, unequal, some of them canine-like; mouth not large; lower pharyngeal teeth sharp, uniserial; vertebræ (in *P. melanostictus*) 11 + 29 = 40. PSETTICHTHYS, 1019.

dd. Lateral line with an arch in front; no accessory branch; vertebræ in smaller number (35 to 41); anal spine usually obsolete; body normally sinistral. (Species chiefly of the temperate or subtropical seas, none of them Arctic and none European.) (Genera allied to *Paralichthys*.)

h. Dorsal fin beginning above the pupil; teeth rather small; no canines; body indifferently dextral or sinistral (in some species at least).

i. Scales ctenoid.

j. Teeth in upper jaw in 2 series; gill rakers broad. VERASPER, 1020.

jj. Teeth all uniserial; gill rakers slender. HIPPOGLOSSINA, 1021.

ii. Scales cycloid; teeth uniserial; gill rakers short and thick.

k. Teeth small, pointed, equal. LIOGLOSSINA, 1022.

kk. Teeth unequal, blunt, conical; caudal fin subsessile, the caudal peduncle extremely short; skin of shoulder girdle with patches of cup-shaped scales; vertebræ (in *X. liolepis*) 12 + 25 = 37. XYSTREURYS, 1023.

hh. Dorsal fin beginning in advance of eye; teeth sharp, uniserial or smooth.

l. Scales weakly ciliated; caudal fin with a distinct peduncle; mouth large; teeth unequal, some of the anterior canine like; gill rakers rather long and slender; no dorsal lobe nor produced ventral rays; vertebræ 35 to 41. PARALICHTHYS, 1024.

ll. Scales very strongly ctenoid on both sides of body; mouth smallish, with small, sharp teeth; anterior rays of dorsal more or less exserted, thus forming a more or less distinct lobe; gill membranes considerably united; gill rakers short and broad; caudal peduncle short; left ventral produced; vertebræ (in *A. quadrocellata*) 9 + 26 = 35.

m. Lateral line with its tubes much branched, covering parts of contiguous scales; dorsal lobe low; left ventral much produced. RAMULARIA, 1025.

mm. Lateral line with its tubes simple, not branched.

n. Body broad, ovate, the depth more than ½ length; dorsal lobe and left ventral moderately produced. ANCYLOPSETTA, 1026.

nn. Body elliptical, the depth not more than ½ length; dorsal lobe and left ventral greatly produced. NOTOSEMA, 1027.

lll. Scales entirely smooth; caudal peduncle short; mouth small; gill rakers short and thick; dorsal with an anterior lobe; left ventral elongate. GASTROPSETTA, 1028

II.—PLEURONECTINÆ.

(Flounder Tribe.)

Mouth small, unsymmetrical, the jaws on the eyed side with nearly straight outline, the bones on the blind side strongly curved; dentition chiefly developed on the blind side; eyes large; edge of preopercle not hidden by the scales; pectoral fins well developed; vertical fins well separated; ventral fins nearly or quite symmetrical, that of the eyed side not prolonged along the ridge of the abdomen; anal spine usually strong (obsolete in *Microstomus* and *Embassichthys*). Body dextral (except frequently in *Platichthys stellatus*). Species arctic or subarctic in distribution.

a. Vertebræ in moderate number, from 10 + 26 = 36 to 11 + 33 = 44; dorsal rays 65 to 80; anal rays 45 to 60.

b. Teeth small, acute, in several series; lateral line nearly straight, with an accessory dorsal branch; lower pharyngeals narrow, with small biserial teeth; scales cycloid.

c. Lips thick, each with several longitudinal folds; dorsal fin beginning on the blind side; vertebræ 38 to 40. Pleuronichthys, 1029.

cc. Lips simple; dorsal fin beginning on the median line; vertebræ (in *H. guttulata*) 11 + 24 = 35. Hypsopsetta, 1030.

bb. Teeth chiefly uniserial, all more or less blunt, conical or incisor-like.

d. Lateral line with an accessory dorsal branch.

e. Lateral line without distinct arch in front.

f. Teeth compressed, incisor-like, close set.

g. Scales closely imbricated, mostly cycloid; upper eye on median line; vertebræ (in *P. vetulus*) 11 + 33 = 44. Parophrys, 1031.

gg. Scales scarcely imbricated, all very strongly ctenoid; eyes both lateral. Inopsetta, 1032.

ff. Teeth conical, separated, not incisor-like; scales closely imbricated, all strongly ctenoid; mouth comparatively large (approaching that of *Psettichthys*); vertebræ (in *I. isolepis*) 10 + 32 = 42. Isopsetta, 1033.

ee. Lateral line with a distinct arch in front; scales imbricated, rough-ctenoid; vertebræ (in *L. bilineata*) 11 + 29 = 40. Lepidopsetta, 1034.

dd. Lateral line without accessory dorsal branch.

h. Lateral line with a distinct arch in front; scales imbricated, rough-ctenoid; vertebræ (in *L. limanda*) about 40. Limanda, 1035.

hh. Lateral line without distinct arch in front.

i. Scales regularly imbricate, all (on eyed side) ctenoid in both sexes; no stellate tubercles on head nor on bases of dorsal and anal fins; teeth, incisor-like, close set; lower pharyngeals very narrow, each with 2 rows of separate, conical teeth; fin rays scaly. Pseudopleuronectes, 1036.

ii. Scales imperfectly imbricated, or else not all ctenoid.

j. Scales chiefly cycloid in both sexes; lower pharyngeals small and narrow, separate, each with about 1 row of small, bluntish teeth; teeth incisor-like, close set, forming a cutting edge; no stellate scales at base of dorsal and anal. Pleuronectes, 1037.

jj. Scales rough-ctenoid in the male, more or less cycloid in the female (fin rays scaly in the male, naked in the female); lower pharyngeals very large, more or less united in the adult, their surface somewhat concave, the teeth in 5 or 6 rows, large, blunt, close set; teeth in jaws incisor-like; fin rays of dorsal and anal without tubercles at base. Liopsetta, 1038.

jjj. Scales all in both sexes and on both sides of the body represented by coarse scattered stellate tubercles; similar tubercles between bases of dorsal and anal rays; lateral line without scales; lower pharyngeals broad, each with 3 rows of blunt, coarse teeth; teeth incisor-like. PLATICHTHYS, 1039.

aa. Vertebræ in increased number (varying from 13 + 35 = 48 to 13 + 52 = 65); dorsal rays 90 to 120; anal rays 70 to 100; teeth broad, incisor-like; scales small, all cycloid. (Genera allied to *Glyptocephalus.*)

k. Left side of skull normal; anal spine obsolete; vertebræ 48 to 52.

l. Body elongate, the depth 2½ to 3 in length; vertebræ 48 to 52. MICROSTOMUS, 1040.

ll. Body stouter, the depth 2 to 2½ in length; vertebræ more numerous, about 63. EMBASSICHTHYS, 1041.

kk. Left side of skull with large mucous cavities; anal spine strong; vertebræ 58 to 65. GLYPTOCEPHALUS, 1042.

III.—PSETTINÆ.

(TURBOT TRIBE.)

Large-mouthed flounders, with the ventral fins unsymmetrical.—Mouth symmetrical, the dentition nearly equally developed on both sides; gape usually wide (narrow in *Platophrys, Etropus,* etc.), the maxillary commonly more than ⅓ length of head; lower pharyngeals narrow, each with one or more rows or a narrow band of small, sharp teeth; teeth in jaws acute; eyes not minute; pectorals and ventrals usually well developed; edge of preopercle free; ventral fins dissimilar in form or in position, that of the left or eyed side inserted on the ridge of the abdomen, its base extended along this ridge, its rays more or less wide apart; caudal fin rounded or subtruncate; no accessory lateral line; anal spine usually weak or obsolete; a pelvic spine sometimes developed; vertebræ in moderate or small number, 31 to 45. Body sinistral. Species chiefly tropical or subtropical in distribution.

a. Pectoral fin of both sides present; septum of gill cavity below gill arches without foramen; a deep emargination near the isthmus; ventral fins free from anal.

b. Vomer with teeth; lateral line with a strong arch in front; teeth subequal, in villiform bands; body broadly ovate; caudal fin subsessile; interorbital area broad; scales small, cycloid; gill rakers long and slender; anterior dorsal rays produced; vertebræ 36. LOPHOPSETTA, 1043.

bb. Vomer toothless; ventral fins free from anal; caudal fin subsessile.

c. Lateral line with a distinct arch in front; teeth small, uniserial, or imperfectly biserial.

d. Interorbital space more or less broad, deeply concave, at least in the males; form broad ovate; gill rakers short and thick.

e. Scales small, ctenoid, adherent, 75 to 100 or more; anterior rays of dorsal not elevated; pectoral of left side usually filamentous in the male; vertebræ (in *P. lunatus*) 9 + 30 = 39. PLATOPHRYS, 1044.

ee. Scales moderate, 60 to 70; anterior rays of dorsal greatly produced; no lateral line on blind side. PERISSIAS, 1045.

dd. Interorbital space a narrow ridge; dorsal not elevated in front.

f. Gill rakers obsolete; interorbital area armed with a spine; scales rough. ENGYOPHRYS, 1046.

ff. Gill rakers slender; right ventral elongate; scales ctenoid. TRICHOPSETTA, 1047.

cc. Lateral line without arch in front.

g. Teeth in upper jaw biserial, in the lower uniserial, the front teeth of upper jaw enlarged; vertebræ 35 or 36; gill rakers short; interorbital space broad in the male. SYACIUM, 1048.

gg. Teeth in each jaw uniserial; interorbital space very narrow, the ridges coalescing between the eyes.

h. Mouth not very small, the maxillary more than ⅓ length of head.

i. Gill rakers very short and thick, tubercle-like.

j. Scales cycloid, small, and firm. CYCLOPSETTA, 1049.

jj. Scales small, firm, ctenoid. AZEVIA, 1050.

ii. Gill rakers slender, of moderate length; scales thin, deciduous, ciliated; vertebræ 34 to 40. CITHARICHTHYS, 1051.

hh. Mouth very small, the teeth subequal, the maxillary less than ⅓ length of head; scales thin; teeth uniserial; vertebræ 9 + 25 = 34. ETROPUS, 1052.

aa. Pectoral fin of blind side wanting; eyes very close together; caudal fin subsessile; teeth small, uniserial; mouth moderate; lateral line of eyed side arched, that of right side nearly straight; dorsal fin beginning on snout, its anterior rays not exserted, its rays all simple and very numerous; gill rakers few and feeble; scales small; body thin, very elongate; vertebræ (in *M. sessilicauda*) 43; (deep-sea flounders). MONOLENE, 1053.

1013. ATHERESTHES, Jordan & Gilbert.

Atheresthes, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 51 (*stomias*).

Eyes and color on the right side. Body very long and slender, closely compressed, tapering into a long and slender caudal peduncle; head elongate, narrow; mouth extremely large, oblique; the long and narrow maxillary extending beyond the eye; each jaw with 2 irregular series of sharp, unequal, arrow-shaped teeth, some of them long and wide set, and others short and close set, sharp; the long teeth freely depressible. Gill rakers numerous, long, slender, and stiff, strongly dentate within. Scales rather large, thin and readily deciduous, slightly ciliated, those on the blind side similar, smooth; lateral line without arch. Fins low and fragile; dorsal commencing over the eye, its anterior rays low, the posterior rays somewhat forked; no anal spine; pectorals and ventrals small, both of the latter lateral; caudal lunate. The single species which constitutes this genus is one of the most remarkable of the flounders. Of all the group, it approaches in form and general characters most nearly to the Gadoid fishes, from ancestors of which we may presume the flounders to be descended, although Dr. Gill has suggested the possibility of their descent from Trachypteroid fishes. (*ἀθήρ*, the beard or spike of an ear of corn; *ἐσθίω*, to eat; from the arrow-shaped teeth.)

2975. ATHERESTHES STOMIAS (Jordan & Gilbert).

(THE ARROW-TOOTHED HALIBUT.)

Head about 3⅔ in length; depth 3½; eye large, 4⅔ in head. D. 103; A. 86; scales 135; vertebræ 12 + 37 = 49. Head long, the snout protruding, somewhat truncate at tip; mouth excessively large; the maxillary more than ½ length of head, and reaching behind eye; premaxillary in front above the level of the lower eye; teeth in upper jaw anteriorly in a single

series, long, slender, and wide set, much smaller and closer set behind; on sides of jaw the teeth are very small and in 2 distinct series, the inner of which corresponds to the single series in front, the teeth thus gradually increasing in size forward; teeth in inner series of lower jaw very sharp and slender, longer than the upper teeth, wide set, alternating with shorter, depressed teeth; outside of these larger teeth is a series of fixed small teeth; all of the long teeth in both jaws depressible and conspicuously arrow-shaped toward their tips; inner series of small teeth in upper jaw also arrow-shaped, depressible; interorbital space scaly, ridged, not a third width of eye. Gill rakers long and strong, about 4 + 13 in number, the longest more than ½ diameter of eye. Upper eye with its range entirely vertical. Scales extremely thin, irregular in size, not evenly imbricated; lateral line very prominent. Dorsal fin beginning just behind the middle of the eye; caudal peduncle nearly as long as the pectoral fin, about ⅔ length of head. Plain olive brown, the margins of the scales darker; blind side dusted with black points. Length 2 feet. Bering Sea to San Francisco, common northward; not rare in deep water off San Francisco, and is brought in in considerable numbers from the sweep-nets (*parranzelle*) used in Drakes Bay. At Unalaska it occurs commonly in shallow water. In the north the flesh is firmer and the coloration more pronounced. Dr. Gilbert dredged it in abundance on both sides of the peninsula of Alaska and in Bristol Bay, in 32 to 406 fathoms. Mr. Scofield found it abundant in Chignik Bay, and it was taken by us in 1897 at Unga and Karluk. (στομίας, large mouthed.)

Platysomatichthys stomias, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 51, 301, San Francisco. (Coll. Jordan & Gilbert.)

Atheresthes stomias, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 57, 454; BEAN, Proc. U. S. Nat. Mus. 1881, 242; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 66; JORDAN & GILBERT, Synopsis, 820, 1883; BEAN, Proc. U. S. Nat. Mus. 1883, 354; JORDAN, Nat. Hist. Aquat. Anim., 188, pl. 53, 1884; JORDAN & GOSS, Review Flounders and Soles, 236, pl. 1, 1889; GILBERT, Rept. U. S. Fish Comm. 1893 (1896), 459.

1014. REINHARDTIUS, Gill.

Reinhardtius, GILL, Cat. Fishes East Coast N. A., 50, 1861 (*hippoglossoides;* no description).

Platysomatichthys, BLEEKER, Comptes Rendus, Ac. Sci. Amsterdam, XIII, 1862, 426 (*pinguis = hippoglossoides*).

Reinhardtius, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 218 (*hippoglossoides*).

Eyes and color on right side. Body more or less elongate, compressed; head long and large; mouth large; maxillary reaching beyond eye; jaws with strong, unequal teeth, the upper with 2 series in front, these converging behind; lower jaw with a single series of strong, distant teeth; no teeth on vomer or palatines. Gill rakers few, short, stout, and rough. Fins rather low; caudal fin lunate. Lower pharyngeal teeth in 1 row. Scales small, cycloid; lateral line without anterior curve. One species known, an arctic fish, in some degree intermediate between the true halibut and *Atheresthes*. (Named for Prof. Johann Reinhardt, of the University of Copenhagen, an able investigator of the fishes of Greenland.)

2976. REINHARDTIUS HIPPOGLOSSOIDES (Walbaum).

(GREENLAND HALIBUT.)

Head $3\frac{4}{5}$ in length; depth nearly 3. D. 100; A. 75; scales 160; orbit 8 in head; snout about $3\frac{1}{2}$, more than twice as long as orbit; eyes even in front; interorbital space flat, scaly, wider than the orbit; lower jaw prominent; length of maxillary $2\frac{1}{2}$ in head; teeth conical, pointed; upper jaw with 2 series, convergent posteriorly, those of the outer series gradually smaller posteriorly; a pair of strong canine teeth anteriorly in the inner series, the other teeth of this series being very small; lower jaw with a series of strong, distant teeth. Gill rakers short, thick, and strongly dentate. Fins naked. Longest dorsal rays $\frac{1}{3}$ length of head; no anal spine; dorsal and anal rays all simple, the dorsal beginning over posterior third of the eye. Scales very small, not ciliated. Yellowish brown. Reaching a very large size. Arctic parts of the Atlantic, south to Finland and the Grand Banks; not very common. (Eu.) (*ἱππόγλωσσος*, halibut; *εἶδος*, resemblance.)

Pleuronectes cynoglossus, FABRICIUS, Fauna Grœnlandica, 163, 1780, Greenland; not of LINNÆUS.

Pleuronectes hippoglossoides, WALBAUM, Artedi Piscium, 115, 1792; based on FABRICIUS.

Pleuronectes pinguis, FABRICIUS, Zoologiske Bidrag., 43, 1824, Greenland.

Hippoglossus grœnlandicus, GÜNTHER, Cat., IV, 404, 1862, Greenland.

Reinhardtius hippoglossoides, GILL, Cat. Fishes East Coast N. A., 50, 1861; GILL, Proc. Ac. Nat. Sci. Phila. 1864, 218.

Platysomatichthys hippoglossoides, GOODE & BEAN, Bull. Essex Inst., II, 7, 1879; COLLETT, Norske Nord-Havs Exped., 142, 1880; JORDAN & GILBERT, Synopsis, 819, 1883; GOODE, Nat. Hist. Aquat. Anim., 197, pl. 56, 1884; JORDAN & GOSS, Review Flounders and Soles, 237, pl. II, 1889; and of late American writers generally.

Hippoglossus pinguis, REINHARDT, Kgl. Dansk. Vidensk. Selsk., 116, 1838.

Platysomatichthys pinguis, BLEEKER, *l. c.*, 426, 1862.

1015. HIPPOGLOSSUS, Cuvier.

(HALIBUT.)

Hippoglossus, CUVIER, Règne Animal, Ed. 1, II, 221, 1817 (*hippoglossus*).

Eyes and color on the right side. Form oblong, not strongly compressed. Mouth wide, oblique; teeth in the upper jaw in 2 series, those below in 1; anterior teeth in upper jaw, and lateral teeth in lower, strong; no teeth on vomer or palatines; lower pharyngeal teeth in 2 rows. Dorsal fin beginning above the eye, its middle rays elevated, the posterior rays of dorsal and anal bifid; caudal fin lunate; ventral fins both lateral. Scales very small, cycloid; lateral line with a strong curve in front. Gill rakers few, short, compressed, wide set. Vertebræ 16 + 34. Largest of the flounders. This genus contains but 1 species, the well-known halibut; abundant on both coasts of the North Atlantic and of the North Pacific. (*Hippoglossus*, the ancient name of the halibut, from *ἵππος*, horse; *γλῶσσα*, tongue.)

2977. HIPPOGLOSSUS HIPPOGLOSSUS (Linnæus).

(HALIBUT.)

Head $3\frac{3}{4}$; depth about 3. D. 105; A. 78; scales 150 or more. Body comparatively elongate, not strongly compressed, deep mesially, thence rapidly tapering each way; head broad; eyes large, separated by a very broad

flattish area; lower eye slightly advanced; mouth large, the maxillary reaching middle of orbit. Nearly uniform dark brown; blind side white. One of our most important food-fishes, reaching a weight sometimes of 400 pounds. Found in all northern seas, southward in deep water to France, Sandy Hook, and occasionally to the Farallones off San Francisco; abundant throughout the North Atlantic as also the North Pacific and Bering Sea, in water of moderate depth; taken with hook and line on all cod banks.

Pleuronectes hippoglossus, LINNÆUS, Systema Naturæ, Ed. x, 269, 1758, European Ocean.
Hippoglossus vulgaris, FLEMING, British Animals, 197, 1828; GÜNTHER, Cat., iv, 403, 1862; DAY, Fishes Great Britain, ii, 5, pl. 44; STORER, Fish. Mass., 145, 1839; DE KAY, New York Fauna: Fishes, pl. 49, f. 157, 294, 1842; STORER, Synopsis Fish. N. A., 475, 1847; LOCKINGTON, Rep. Com. Fisheries California, 39, 1878–79; LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 71; BEAN, Proc. U. S. Nat, Mus. 1879, 63; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 454; GOODE, Proc. U. S. Nat. Mus. 1880, 471; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 66; BEAN, Proc. U. S. Nat. Mus. 1881, 242; JORDAN & GILBERT, Synopsis, 819, 1883; BEAN, Cat. Col. Fish. U. S. Nat. Mus. 1883, 20; DRESEL, Proc. U. S. Nat. Mus. 1884, 244; GOODE, Nat. Hist. Aquatic Anim., 189, pl. 54, 1884; and of American writers generally.
Hippoglossus maximus, GOTTSCHE, Archiv fur Naturgesch. 1835, 164, no locality.
Hippoglossus gigas, SWAINSON, Nat. Hist. Class'n Anim., ii, 302, 1839, no locality.
Hippoglossus pontius, BONAPARTE, Catalogo Metodico, 47, 1846, Black Sea; after PALLAS.
Hippoglossus americanus, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 220.
Hippoglossus hippoglossus, JORDAN, Cat. Fish. N. A., 133, 1885; JORDAN & GOSS, Review Flounders and Soles, 237, pl. 3, 1889.

1016. LYOPSETTA, Jordan & Goss.

Lyopsetta, JORDAN & GOSS, in JORDAN, Cat. Fish. N. A., 135, 1885 (*exilis*).

Teeth sharp, those of the lower jaw uniserial, the upper jaw biserial; lateral line simple (without accessory dorsal branch) and without distinct anterior arch. Scales comparatively large, thin, ciliated, and deciduous; body dextral; anal spine usually strong; vertebræ about 45; body slender, the flesh soft; dorsal fin beginning above eye. This genus contains but a single species, a small, soft-bodied flounder, of the waters of the North Pacific. In its technical characters *Lyopsetta* is very close to *Hippoglossoides*, but the species has the soft flesh of *Atheresthes*. (λύω, to loosen; ψῆττα, flounder.)

2978. LYOPSETTA EXILIS (Jordan & Gilbert).

Head 4; depth 3¼. D. 78; A. 62; V. 6; scales 16–71–18. Body slender, compressed, the flesh soft; caudal peduncle slender; mouth not large, very oblique, the gape curved; lower jaw scarcely projecting, with a knob at symphysis; maxillary rather narrow, reaching middle of pupil, 2⅔ in length of head; teeth small, slender, close set, nearly uniform; above in 2 series, below in 1. Eyes large, separated by a sharp, scaly ridge; lower eye advanced. Scales comparatively large, thin and deciduous, ctenoid, but not so rough as in the other species, those on blind side similar, less rough. Lateral line prominent, rising anteriorly, without trace of arch. Fins low, fragile; anal preceded by a spine; caudal fin long, rather pointed; pectorals small, the right pectoral little more

than ⅓ length of head. Dorsal beginning immediately in front of pupil; anal higher than dorsal. Gill rakers short, slender, toothed, 9 below angle, the longest about ½ diameter of orbit. Pale olivaceous brown, with dark points, forming edgings on each scale; bronze spots sometimes present; fins mostly dusky; dorsal and anal edged anteriorly with yellowish; ventrals largely yellow. Length 12 inches. North Pacific, in rather deep water; San Francisco to Puget Sound. This small flounder is brought in in large quantities by the sweep nets off San Francisco. It is of little value as a food-fish. (*exilis*, slender.)

Hippoglossoides exilis, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 154, off San Francisco (Type, No. 27121. Coll. Jordan & Gilbert); JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 454; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 67; JORDAN & GILBERT, Synopsis, 827, 1883.

Lyopsetta exilis, JORDAN & GOSS, Review Flounders and Soles, 238, 1889.

1017. EOPSETTA, Jordan & Goss.

Eopsetta, JORDAN & GOSS, in JORDAN, Cat. Fish. N. A., 135, 1885 (*jordani*).

Teeth sharp, those of the lower jaw uniserial, the upper biserial; scales small, ciliated, and adherent; lateral line without accessory dorsal branch and without distinct anterior arch; anal spine usually strong; body normally dextral, robust, the flesh firm; dorsal fin beginning above eye; vertebræ about 43. This genus contains but a single species, a large flounder which is abundant on the coast of California. It is very close to the genus *Hippoglossoides*. (ἑως, morning; πῆττα, flounder.)

2979. EOPSETTA JORDANI (Lockington).

(CALIFORNIA "SOLE.")

Head 3⅛; depth 2½. D. 94; A. 72; scales 96. Body broadly elliptical. Dorsal and ventral outline equally and regularly curved. Mouth oblique, the jaws about even, the symphyseal knob but little projecting; gape curved; maxillary broad, reaching to behind pupil, 2⅔ in head; teeth in 2 series in the upper jaw, the inner series small and distant from the outer, which is considerably enlarged in front; lower jaw with a single series similar to the outer series in the upper jaw, but larger. Gill rakers roughish, strong, about 15 below angle, the longest about ½ as long as eye. Lower pharyngeals rather narrow, each with a single row of sharp teeth. Eyes large; interorbital space a narrow, blunt, scaly ridge. Dorsal beginning over anterior margin of pupil, the rays all simple; caudal fin with the middle rays slightly produced; anal preceded by a spine; pectoral ½ length of head. Scales of colored side small, firm, strongly ciliated, nearly uniform over head and body; lower jaw and snout scaleless; scales on blind side smooth. Olive brown, nearly uniform; membrane of dorsal and anal fins clouded with darker. Length 20 inches. Pacific Coast of the United States from Puget Sound to Point Concepcion. One of the commonest flatfishes of the California coast, being found in abundance in shallow water from Monterey northward. It is a good food-fish, and large numbers are dried each year by the Chinese. (Named for David Starr Jordan.)

Hippoglossoides jordani, LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 73, San Francisco (Coll. W. N. Lockington); LOCKINGTON, Rep. Com. Fisheries California 1878–79, 40; LOCKINGTON, Scientific Press Supplement, April, 1879, 120; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 454; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 67; JORDAN & GILBERT, Synopsis, 826, 1883; JORDAN, Nat. Hist. Aquat. Anim., 187, 1884.
Eopsetta jordani, JORDAN & GOSS, Review Flounders and Soles, 239, 1889.

1018. HIPPOGLOSSOIDES, Gottsche.

Hippoglossoides, GOTTSCHE, Archiv fur Naturgesch. 1835, 164 ("*limanda*"=*platessoides*).
Citharus, REINHARDT, Kong. Dansk. Vid. Selsk. 1838, 116 (*platessoides*); not *Citharus* BLEEKER, 1862.
Drepanopsetta, GILL, Cat. Fish. East Coast N. A., 50, 1861 (*platessoides*).
Pomatopsetta, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 217 ("*dentata*"=*platessoides*).

Eyes and color on the right side (except sometimes in *H. elassodon*). Body oblong, moderately compressed; mouth rather large, with 1 row of sharp teeth on each jaw; no teeth on vomer or palatines; gill rakers rather long and slender; scales ctenoid; lateral line nearly straight, simple; dorsal fin low in front, beginning over or before the eye; ventrals both lateral; caudal double truncate, produced behind. This genus, as here restricted, contains 3 closely related species, 2 of the North Pacific, 1 of the North Atlantic. All are essentially arctic species, inhabiting shallow waters in the regions where they are most abundant. (*ἱππόγλωσσος, Hippoglossus; εἶδος*, resemblance.)

a. Dorsal rays about 88; anal about 70; gill rakers $x+10$; interorbital space with an obtuse, prominent, rather broad ridge. PLATESSOIDES, 2980.
aa. Dorsal rays about 82; anal about 61; gill rakers $x+12$ to 14; interorbital space with a narrow, nearly naked ridge. ELASSODON, 2981.
aaa. Dorsal rays 72 to 76; anal 56 to 60; gill rakers $x+12$; interorbital space moderate, with 2 rows of scales.
b. Depth $2\frac{1}{3}$ in length; D. 76; A. 60; pectoral $\frac{1}{3}$ length of head. ROBUSTUS, 2982.
bb. Depth $2\frac{2}{3}$ in length; D. 72; A. 56; pectoral $\frac{2}{3}$ in length of head. HAMILTONI, 2983.

2980. HIPPOGLOSSOIDES PLATESSOIDES (Fabricius).

(SAND-DAB.)

Head $3\frac{3}{4}$; depth $2\frac{1}{3}$. D. 88 (80 to 93); A. 70 (64 to 75); scales 90 (pores). Body ovate; mouth moderate, oblique; maxillary narrow, reaching to below pupil, $2\frac{2}{3}$ in length of head; teeth rather small, conical, larger anteriorly, in 1 row in each jaw, those in the lower largest. Eyes rather large, the upper longer than snout, $4\frac{1}{3}$ in head; lower jaw included, but with a projecting knob at the chin; snout thick, scaly; interorbital space narrow, with a raised obtuse ridge entirely covered with rough scales in about 6 series; mandible with a series of scales; gill rakers rather short and robust, not toothed, about 10 below angle, the longest less than $\frac{1}{3}$ length of eye; fins with small, rough scales; a strong preanal spine; pectoral not quite $\frac{1}{3}$ length of head. Reddish brown, nearly plain. The identity of the American and European representatives of this species (*platessoides* and *limandoides*) is now conceded by all writers. A little difference is recognizable between arctic and subarctic examples, the

former having a somewhat greater number of fin rays. Thus Greenland specimens, according to Collett, have D. 88, A. 69; specimens from Finmark have D. 92, A. 72; these representing the var. *platessoides*. Specimens from England (var. *limandoides*) have D. 80, A. 66, while those from intermediate localities present in general fin formulæ likewise intermediate, showing that no sharp division is possible. This is a rather common food-fish of the deep waters northward, on both sides of the ocean. North Atlantic, south to Cape Cod, and the coasts of England and Scandinavia. (Eu.) (*platessa*, the plaice; *εἶδος*, resemblance.)

Pleuronectes linguatula, MÜLLER, Zool. Dan. Prodromus, 45, 1776; not of LINNÆUS.
Pleuronectes platessoides, FABRICIUS, Fauna Grœnlandica, 164, 1780, Greenland.
Pleuronectes limandoides, BLOCH, Ausl. Fische, III, 24 tab. 186, 1787, Europe, and of various copyists.
Pleuronectes limandanus, PARNELL, Edinburgh New Phil. Journ. 1835, 210.
Citharus platessoides, REINHARDT, Kongl. Dansk. Vid. Selsk., 116, 1838.
Drepanopsetta platessoides, GILL, Cat. Fish. East Coast N. A., 50, 1861.
Hippoglossoides platessoides, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 217; COLLETT, Norske Nord-Havs. Exped., 144, 1880; GOODE, Proc. U. S. Nat. Mus. 1880, 471; JORDAN & GILBERT, Synopsis, 826, 1883; STEARNS, Proc. U. S. Nat. Mus. 1883, 125; GOODE, Nat. Hist. Aquatic Anim., 197, pl. 55, 1884; JORDAN & GOSS, Review Flounders and Soles, 240, pl. 4, 1889; GOODE & BEAN, Ocean Ichthyology, 438, 1896, and of recent American writers generally.
Hippoglossoides limandoides, GÜNTHER, Cat., IV, 405, 1862; DAY, Fishes Great Britain and Ireland, II, 9, pl. 45, 1884.
Hippoglossoides limanda, GOTTSCHE, Archiv fur Naturgesch. 1835, 168; not *Pl. limanda*, LINNÆUS.
Platessa dentata, STORER, Rept. Fish. Mass., 143, 1839; DE KAY, N. Y. Fauna: Fishes, 298, 1842; STORER, Synopsis, 476, 1846.
Hippoglossoides dentatus, GÜNTHER, Cat., IV, 406, 1862; GÜNTHER, Challenger Report, XXII, Fishes, 3, 1887.
Pomatopsetta dentata, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 217.

2981. HIPPOGLOSSOIDES ELASSODON, Jordan & Gilbert.

Head 3½; depth 2½; eye 4 in head. D. 77 to 87; A. 59 to 67; V. 6; scales 45–100–40. Body oblong-elliptical; caudal peduncle about as long as deep; upper profile of head continuous with the outline of back; depression over eye slight; mouth rather large, the gape curved, considerably wider on the blind side; lower jaw projecting, with a symphyseal knob; maxillary narrow, reaching beyond middle of pupil, 2½ in head; teeth small, close set, nearly uniform, in a single row. Gill rakers slender, smooth, 14 to 16 below arch, the longest nearly ½ diameter of orbit. Eyes large, separated by a narrow, knife-like ridge, which is naked, or with a single series of scales. Scales small, firm, rough, those on tail roughest, those on blind side similar, mostly smooth anteriorly. Lateral line rising anteriorly, but without arch; dorsal beginning immediately in front of pupil; anal preceded by a spine; caudal long; pectoral of eyed side ½ length of head; ventral reaching past front of anal; pectoral and ventral of eyed side with prickle-like scales. Brownish, nearly uniform, sometimes spotted with darker; fins grayish, irregularly blotched with dusky. Body sometimes sinistral. Length 18 inches. Bering Sea south to Cape Fattery; a rather abundant shore fish in Puget Sound, and it

seems to be still more common northward, being, in Alaska, a food-fish of some importance. Abundant north and south of the Aleutian Islands and in Bristol Bay. Our specimens from Kamchatka agree in all respects; D. 77 to 84; A. 60 or 61. Pectoral not quite $\frac{1}{2}$ head. Interorbital ridge sharp, with 1 series of scales; gill rakers $x+14$. (ἐλασσόω, to diminish; ὀδούς, tooth.)

Hippoglossoides elassodon, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 278, Seattle; Tacoma (Type, No. 27263. Coll. D. S. Jordan); JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 454; BEAN, Proc. U. S. Nat. Mus. 1881, 242; JORDAN & GILBERT, Synopsis, 826, 1883; BEAN, Proc. U. S. Nat. Mus. 1883, 20; JORDAN, Nat. Hist. Aquat. Anim., 188. pl. 52, 1884; JORDAN & GOSS, Review Flounders and Soles, 241, pl. 5, 1889; JORDAN & GILBERT, Rept. Fur Seal Invest., 1898.

2982. HIPPOGLOSSOIDES ROBUSTUS, Gill & Townsend.

Head $3\frac{3}{5}$; depth $2\frac{1}{5}$; eye $5\frac{2}{3}$ in head. D. 76; A. 60; scales 95 (pores). Interorbital space a broad, somewhat elevated ridge with 2 rows of scales. Body rather high, its greatest height nearly equaling $\frac{1}{3}$ the length from snout to base of caudal; profile decurved above the eye; body thick; scales on head separate and rarely touch each other. Gill rakers long, $x+11$. Maxillary $2\frac{1}{2}$ in head, directed upward anteriorly; teeth of the single row mostly separated from each other by intervals equal to width of teeth, curved inward, and uniform on the sides; toward front 4 or 5 enlarged, preceded by 2 smaller, leaving the middle toothless; in the lower jaw of nearly uniform size and inclining backward. Pectoral $\frac{1}{4}$ head; ventrals reaching first or second anal ray. Scales on body ciliated or weakly ctenoid, those on cheek smoother; no ctenoid scales on blind side. Caudal shorter than in *H. hamiltoni*, $1\frac{3}{5}$ in head. No exserted nasal tubes. Color plain brown. Bering Sea. Only the type known, $12\frac{1}{2}$ inches long, from which we have taken the above description. (*robustus*, robust.)

Hippoglossoides robustus, GILL & TOWNSEND, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17, 1897), 234, Bering Sea, Lat. 56° 14′ N., Long. 164° 08′ W., Albatross Station 3541, in 49 fathoms. (Type, No. 48766, U. S. Nat. Mus. Coll. *Albatross*.)

2983. HIPPOGLOSSOIDES HAMILTONI, Jordan & Gilbert, new species.

Head $3\frac{1}{5}$ in length; depth $2\frac{2}{5}$; longest diameter of upper eye $3\frac{1}{2}$ in head; snout (measured from upper eye) 5 in head; maxillary of colored side $2\frac{1}{3}$, of blind side $2\frac{1}{6}$, in head; depth of caudal peduncle equaling its length, $3\frac{1}{3}$ in head. D. 72; A. 56; P. 11; pores in lateral line 91. Upper profile of head continuing the dorsal curve without interruption, there being a slight depression above the eye and an increased convexity on the snout; mandible very heavy, projecting anteriorly, so that its symphyseal profile completes the curve of the snout; a very short prominence at symphysis directed vertically downward; gape strongly curved and the mouth narrowed anteriorly, so that the maxillary and premaxillary are almost wholly concealed along the middle of their length by the overarching prefrontal; teeth acute, in a single series in each jaw, all except the anterior teeth in each jaw short; at the symphysis of lower jaw the teeth are

longer and directed inward, while in the anterior end of each premaxillary the teeth are still more enlarged, and the series on each side describes a strong curve with its convex side toward the median line; maxillary reaching vertical from slightly behind middle of lower eye; nostril tubes conspicuous, the anterior in closest proximity to the upper lip, which it entirely overhangs; posterior nostril tube wider and slightly shorter; eyes of nearly equal size, and opposite, separated by a wider ridge than in *H. elassodon*, the ridge bearing in its narrowest portion 2 well-defined rows of strongly spinous scales; a conspicuous series of pores joining lateral line with upper margin of upper eye, and another encircling the lower eye below and behind; a third series along mandible and preopercle; 1 large pore above posterior nostril; gill rakers slender, unarmed, 2 above the angle, 11 or 12 below it, the longest $2\frac{3}{4}$ in eye; dorsal fin beginning above front of pupil, the longest ray $2\frac{5}{6}$ in head; anal preceded by a strong spine, its height equaling that of dorsal; pectoral very long and slender, $\frac{3}{5}$ length of head, that of blind side shorter, $\frac{1}{2}$ length of head; ventrals reaching to base of fourth or fifth anal ray; caudal long, evenly rounded behind, the middle rays not longer than those adjacent, their length equaling distance from tip of snout to preopercular margin; scales on colored side strongly ctenoid except in a strip along middle of sides anteriorly; elsewhere each scale provided with 2 to 4 long spines; on blind side they are smooth except on nape and caudal peduncle; cheeks, opercles, and interorbital space covered with larger, rougher scales than those on sides; mandible and snout naked; a single series surrounding each eye anteriorly, and 1 on maxillary or colored side; blind side of head with maxillary naked; cheeks covered with minute smooth thin scales, the opercles with a few scattered spinous scales, the preopercle naked. Color nearly uniform brownish, without distinctive markings on body or fins. One specimen, 17 cm. long, from *Albatross* Station 3641, off Dalnoi Point, Kamchatka; depth 16 fathoms. Allied to *Hippoglossoides elassodon*, from which it differs in the fewer fin rays and scales, the wider interorbital space, the longer caudal and pectoral fins and the much smaller symphyseal knob. The nasal tubes are larger, the scales rougher, and the anterior part of lateral line more arched. Its relations with *H. robustus* are much nearer but the species are apparently distinct. (Named for Gerald Edwin H. Barrett-Hamilton, of Dublin, member of the British Commission of Fur Seal Investigation, 1896 and 1897, who made valuable collections of Kamchatkan fishes.)

Hippoglossoides hamiltoni, JORDAN & GILBERT, Rept. Fur Seal Invest., 1898, Dalnoi Point, Kamchatka. (Coll. *Albatross*.)

1019. PSETTICHTHYS, Girard.

Psettichthys, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 140 (*melanostictus*).

Body dextral; teeth uniserial, sharp, unequal, some of them canine-like; mouth moderate, the lower pharyngeal teeth sharp, uniserial; scales small, ctenoid, ciliated, and firm; lateral line with an accessory dorsal branch and without distinct anterior arch; anal spine strong; dorsal fin

beginning before the eye; vertebræ about 40; flesh firm. This genus contains but 1 species, found on the coast of California. It is nearly related to *Hippoglossoides*, but possesses the peculiar accessory dorsal branch to the lateral line, characteristic of so many of the Pacific coast flounders. (*ψῆττα*, the turbot; *ἰχθύς*, fish.)

2984. PSETTICHTHYS MELANOSTICTUS, Girard.

Head 4; depth $2\frac{1}{3}$. D. 85; A. 60; scales 112. Body not very deep, elliptical; mouth rather small, the maxillary extending to below pupil, $2\frac{3}{7}$ in head; teeth large, in a single series in each jaw, those in lower jaw largest; a few large canines in front of each jaw. Eyes very small, separated by a broad, flat, scaly space, without ridge; lower eye slightly in advance of upper; gill rakers rather stout, weak, hooked at tip, 14 below the angle; scales very small, ctenoid on colored side; lateral line nearly straight, with a long accessory dorsal branch; dorsal commencing in advance of upper eye, the anterior rays elevated, slender and exserted, the longest about $\frac{1}{3}$ length of head; first ray of dorsal nearly free from its membrane; pectoral fin short, $2\frac{1}{3}$ in head; anal fin preceded by a spine; caudal large, strongly convex; lower pharyngeals very narrow, each with 1 row of sharp, recurved teeth. Grayish brown, finely speckled with darker on body and fins. Pacific coast of North America, from Sitka south to Monterey. This is one of the commoner flounders of the Pacific coast, being everywhere known by the name of "Sole." It lives near the shore, and reaches a length of about 20 inches. In color this species is quite unlike the species of *Hippoglossoides*, but in most other respects the two groups are closely allied. (*μέλας*, black; *στικτός*, spotted.)

Psettichthys melanostictus, Girard, Proc. Ac. Nat. Sci. Phila. 1854, 140, San Francisco; Astoria, Oregon; Girard, U. S. Pac. R. R. Surv., x, Fishes, 154, 1858; Günther, Cat., iv, 420, 1862; Lockington, Rep. Com. Fisheries Cal. 1878–79, 40; Lockington, Proc. U. S. Nat. Mus. 1879, 76; Jordan & Gilbert, Proc. U. S. Nat. Mus. 1880, 453; Jordan & Gilbert, Proc. U. S. Nat. Mus. 1881, 67; Jordan, Nat. Hist. Acquatic Animals, 186, pl. 51, 1884; Jordan & Goss, Review Flounders and Soles, 241, pl. 6, 1889.

Hippoglossoides melanostictus, Jordan & Gilbert, Synopsis, 828, 1883.

1020. VERASPER, Jordan & Gilbert, new genus.

Verasper, Jordan & Gilbert, Report Fur Seal Invest., 1898 MS. (*moseri*).

This genus is allied to *Xystreurys* and *Hippoglossina*, having few short gill rakers like the former and strongly ctenoid scales like the latter. It differs strongly from all its congeners in having the premaxillary teeth in 2 series, teeth uniformly small, without canines. Body dextral; dorsal inserted above the front of pupil; lateral line strongly arched above the root of the pectoral, without recurrent dorsal branch; scales firm, extremely spinous; gill rakers short, thick, and triangular, few in number; none of the fin rays notably produced or exserted. Japan and Kuril Islands; 2 species known, the following and *V. variegatus* (Schlegel), a common food fish of Japan, the 2 very closely related. (*verus*, true; *asper*, rough, the word being suggested by *Veratrum*.)

2985. VERASPER MOSERI, Jordan & Gilbert, new species.

Head 3⅓ in length to base of caudal; depth 2. D. 82; A. 58; pectoral 12; pores in lateral line 84. Depth of caudal peduncle 4 in greatest depth of body; length of caudal peduncle, measured axially, 1⅞ in its depth. Head much depressed, with rather wide, flat interorbital space, resembling in appearance *Psettichthys melanostictus*, its thickness at interorbital space equaling distance between pupils of upper and lower eyes. Mouth small, very oblique, the gape strongly arched, the broad maxillary reaching a vertical behind middle of pupil, 2⅘ in head; mandible narrowing toward tip, with very rudimentary symphyseal knob. Teeth in upper jaw in 2 distinct series throughout, those of the outer series increasing slightly in size toward front of jaw, but none of them canine-like; mandibular teeth in 1 row, except at symphysis, where a few teeth form a short outer series. Nasal openings of eyed side approximated in front of middle of interorbital space, the anterior with a short tube, the posterior with a raised rim. Eyes small, their anterior margins opposite, the diameter of lower eye equaling distance from tip of snout to posterior nostril, 6⅓ in head. Interorbital space rather broad and flat, not ridge-like, its total width equaling ½ diameter of orbit. Gill rakers short, broad, triangular, minutely toothed on inner margin, ⅓ diameter of eye; 7 present on horizontal limb of outer arch. Lateral line with a short high anterior arch, the cord of which is ⅕ the straight portion; height of arch ⅓ its length; behind the arch lateral line descending in a gentle curve to middle of sides, the scales very rough, each possessing several long, sharp spines diverging from median portion of posterior margin; anterior and posterior portions of dorsal and anal fins naked, the rays of the middle portion each with a series of strongly ctenoid scales; caudal densely scaled to tip; pectorals and ventrals naked; head covered with strongly spinous scales, excepting snout, maxillary, and mandible; on blind side of head the snout, jaws, preopercle, subopercle, lower half of opercle, and all but a central strip on interopercle, scaleless; on blind side the scales are rough on head, ventral area, and along bases of ventral fins, largely smooth elsewhere. Dorsal beginning above front of pupil, the rays increasing in length to the forty-fifth, which is 2⅔ in head; longest anal ray (the seventeenth) 2⅓ in head. Caudal broadly rounded, 1⅔ in head; pectoral short and broad, 2⅔ in head; ventrals of nearly equal length, reaching origin of anal, 3⅓ in head; no anal spine. Color in spirits, centers of the scales light gray, the margins dark brown; fins light or dusky, the vertical fins with conspicuous black bars, parallel with the rays, these most evident on under side where the pigment seems principally to occur, and are seen through the fin more faintly on the colored side; lining of cheeks and gill cover of colored side dusky; peritoneum gray. Kuril Islands; 1 male 28 cm. long, from Shana Bay, Iturup Island; also taken at Hakodate. (Named for Jefferson Franklin Moser, U. S. N., Lieutenant-Commander, in charge of the U. S. Fish Commission Steamer *Albatross*, and a member of the United States Fur Seal Commission for 1896.)

Verasper moseri, JORDAN & GILBERT, Rept. Fur Seal Invest., 1898 MS., Shana Bay, Iturup Island, Kuril Group. (Type No. 48797. Coll. *Albatross*, Capt. J. F. Moser.)

1021. HIPPOGLOSSINA, Steindachner.

Hippoglossina, STEINDACHNER, Ichth. Beiträge, V, 13, 1876 (*macrops*).

Teeth rather small, uniserial, no canines; lateral line with a strong arch in front, and with no accessory dorsal branch; anal spine obsolete; body indifferently dextral or sinistral (in some species at least). Scales ctenoid; dorsal fin beginning above pupil; gill rakers rather long and slender. This genus is intermediate between *Hippoglossoides* and *Paralichthys*, agreeing with the former in the insertion of the dorsal and in general appearance, and with the latter in the direction of the lateral line. Several species are now known. Some of them are dextral, and perhaps all of them are normally so, or perhaps, as in the case of *Xystreurys liolepis*, all are indifferently dextral or sinistral. (A diminutive of *Hippoglossus*, the halibut.)

a. Mouth large, the maxillary extending to opposite posterior margin of eye, 2 in head; gill rakers numerous, 4 + 13; dorsal rays about 68; anal 53. STOMATA, 2986.

aa. Mouth moderate, the maxillary extending to opposite middle of pupil, about 2½ in head.

b. Dorsal rays about 66; anal 52; depth of body 2⅓ in length. MACROPS, 2987.

bb. Dorsal rays about 62; anal 48; depth of body 2⅔ in length; gill rakers 2 + 8 or 9. BOLLMANI, 2988.

2986. HIPPOGLOSSINA STOMATA, Eigenmann & Eigenmann.

Head 2¾ to 3 in length; depth 2½ to 2⅔. D. 67 to 70; A. 52 to 54; scales 80. Sinistral. Eye (not orbit) large, 5 in head; lower orbit slightly in advance of upper; interorbital a narrow ridge. Form, elongate elliptical, the profile depressed over the eye. Mouth large, maxillary extending to posterior margin of eye, as long as or longer than pectoral, 2 in head; lower jaw about 1⅔ in head. Teeth small, uniserial; anterior nares of each side with long dermal flaps. Scales of left side all ctenoid, those of right side cycloid on anterior half or two-thirds of body; middle third of interorbital naked, anterior and posterior thirds scaled. Gill rakers 4 + 13 or 14. Dorsal beginning over middle of eye, anterior rays with but 1 or 2 scales, rest scaled to near tip, all but last 8 rays simple; anal similar to dorsal, with a strong procumbent spine; highest dorsal and anal rays about 3⅓ in head; pectoral of colored side about 2 in head, that of blind side shorter; caudal double truncate, 5 to 5½ in length. Brown, strongly tinged in life with robin's-egg blue; numerous spots of light blue and light and dark brown; 5 pairs of large, dark-brown ocelli along dorsal and ventral parts of eyed side, the alternate ones longer and more conspicuous; fins colored like body, profusely mottled with light and dark; sinistral pectoral barred; a dark-brown spot above and below on caudal peduncle just in front of caudal, showing conspicuously on blind side. The eggs are probably pelagic; they are transparent, and measure 1.2 mm. in diameter; the single oil globule measures 0.16 mm. Coast of southern California; 2 specimens obtained in deep water off San Diego, November 7, 1889, both females, 1 with ripe eggs. (Eigenmann & Eigenmann.) (στοματός, large mouthed.)

Hyppoglossina stomata, EIGENMANN & EIGENMANN, Proc. Cal. Ac. Sci. 1890, 22, San Diego. (Coll. C. H. Eigenmann.)

2987. HIPPOGLOSSINA MACROPS, Steindachner.

Head $2\frac{3}{4}$; depth $2\frac{1}{3}$. D. 66; A. 52; scales 75 to 80; upper orbit $3\frac{1}{2}$ in head. Body elliptical, deeper than in related species; mouth moderate, the maxillary reaching to middle of eye; teeth small, sharp, uniserial; lower eye slightly in front of upper; eyes separated by a naked narrow ridge; nostrils close together, the anterior ending in a tube; horizontal limb of preopercle somewhat concave, the vertical convex. Dorsal beginning over middle of eye; pectoral of left side $\frac{1}{2}$ head, much longer than maxillary, which is $2\frac{3}{5}$ in head; interorbital space a narrow ridge; scales of left side all strongly ctenoid, those on blind side ciliated only on posterior third of body; no anal spine. Color brownish, with obscure darker blotches. Body sinistral (in the only specimen known). (Steindachner.) Pacific coast of Mexico. One specimen from Mazatlan; not seen by us. (*μακρός*, large; *ὤψ*, eye.)

Hippoglossina macrops, Steindachner, Ichth. Beitr., v, 13, pl. 3, 1876, Mazatlan; Jordan & Goss, Review Flounders and Soles, 242, 1889.

2988. HIPPOGLOSSINA BOLLMANI, Gilbert.

Head 3 ($3\frac{3}{4}$ to $3\frac{1}{7}$) in length; depth $2\frac{3}{5}$ to $2\frac{3}{4}$ ($3\frac{1}{4}$ to $3\frac{1}{2}$); snout 5 in head. D. 60 to 63; A. 47 to 49; scales along lateral line 70 to 75. Body regularly elongate, elliptical; dorsal and ventral outlines equally curved; orbital rim entering anterior profile, which is equally curved before and behind eyes; greatest depth of body above pectorals. Mouth rather large, the maxillary reaching about to middle of pupil, $2\frac{1}{3}$ to $2\frac{3}{5}$ in head. Teeth equally developed on both sides, small and equal, uniserial. Premaxillary spine prominent. Interorbital space a narrow, sharp, naked ridge; eyes large, the lower slightly in advance of upper, $3\frac{3}{4}$ to 4 in head. Gill rakers moderately long and slender, the longest 3 in length of ventral of eyed side; 2 + 8 or 9 developed, the last 2 much shorter. Scales small, firm, strongly ctenoid, those below pectoral much reduced, about 40 in a cross series; arch of lateral line strongly marked, $2\frac{3}{4}$ to $2\frac{1}{1}$ in straight part. Dorsal beginning above middle of pupil of upper eye, its anterior rays low, its longest rays $2\frac{1}{7}$ in head; a strong antrorse spine before anal; pectoral of eyed side 2 in head, that of blind side $2\frac{1}{2}$ to $2\frac{3}{5}$ in head; ventrals subequal, each 6-rayed, 4 in head, extending more than $\frac{1}{2}$ their length beyond anus; each is lateral, but that of eyed side nearest ridge of abdomen, and a little behind its fellow; last ray of left ventral joined to abdomen alongside of anal spine; caudal acute, its peduncle long. Color grayish brown, a row of 6 round, bluish spots, smaller than pupil, along base of dorsal, 4 similar spots along base of anal, and a few indistinct smaller ones on rest of body and head; body with 6 large black spots somewhat smaller than eye, these regularly 4 below dorsal and 2 above anal, the first of dorsal above arch of lateral line, the second above anterior third of straight part, the third at base of last rays and almost forming a cross bar with the 1 at base of anal rays. Dorsal, anal, and caudal dusky, with small whitish spots; a pale spot at base of last 4 dorsal and anal rays; a small black spot at base of outer caudal rays on peduncle; pectorals and ventrals dusky, but not spotted; right side immaculate. Length

7 inches. Pacific coast of Colombia. Numerous specimens were dredged at *Albatross* Station 2805, at a depth of 51½ fathoms. This species differs from *Hippoglossina stomata* in the gill rakers, which are shorter and fewer in number, and in the larger scales on sides. Scales in 16 rows between lateral line and back, instead of 21 or 22, as in *H. stomata*. Gill rakers somewhat shorter, 8 or 9 on anterior limb, 2 on upper limb. In *H. macrops* the gill rakers are slender, close set, 13 or 14 on anterior limb, 4 on vertical limb. In other respects of color, fin rays, and squamation agreeing perfectly with *H. stomata*. (Named for Charles Harvey Bollman.)

Hippoglossina macrops, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 175; not of STEINDACHNER.

Hippoglossina bollmani, GILBERT, Proc. U. S. Nat. Mus. 1890, 122, Albatross Station 2805, southwest of Panama, in 51½ fathoms. (Type, No. 41143.)

1022. LIOGLOSSINA, Gilbert.

Lioglossina, GILBERT, Proc. U. S. Nat. Mus. 1890, 122 (*tetrophthalmus*).

This genus is allied to *Hippoglossina*, but its scales are all cycloid, the teeth are small, pointed, uniserial, and uniform, and the gill rakers short and thick. (λειῶς, smooth; πλῶσσα, tongue; for *Hippoglossina*.)

2989. LIOGLOSSINA TETROPHTHALMA, Gilbert.

Head large, 3¼ in length in a specimen 1 foot long. D. 76 to 83; A. 58 to 62; lateral line (pores) 97. Body of moderate height, the profile distinctly angulated above upper pupil, the snout projecting; length of caudal peduncle ½ its depth, its outlines diverging backward; depth of body 2⅛ in length; snout projecting beyond profile, bluntly rounded, the lower jaw included. Mouth large, the maxillary reaching nearly to vertical from posterior border of lower eye, 2⅓ in head; a blunt projecting process anteriorly from head of maxillary. Teeth small, pointed, in a single close-set series in each jaw, none of them enlarged; vomer toothless; lower eye slightly in advance of upper; vertical from front of upper falling midway between front of orbit and front of pupil of lower eye; vertical diameter of upper orbit but little more than ⅓ its longitudinal diameter, which is contained 3½ in head; interorbital space a blunt high ridge, entirely scaleless, its width ⅔ diameter of orbit. Anterior nostril of blind side with a very long flap, that of eyed side shorter; a well-marked cutaneous flap on lower eye above pupil. Gill rakers very large, broad, and strong, well toothed on inner edges, longest equaling diameter of pupil, the number on outer gill arch 10 or 11. First dorsal ray over anterior margin of pupil of upper eye, the fin not high, its highest ray 3 in head; anal similar; caudal sharply double truncate, the median rays produced; ventrals rounded, equal, barely reaching front of anal; no spine before anal fin; pectorals moderate, with 9 or 10 developed rays, ½ length of head; ventral 6. Scales rather small, growing distinctly larger posteriorly, everywhere smooth; head scaled, except snout, interorbital area, mandible, and part of maxillary, the latter with a patch of scales on posterior end of its expanded portion; on blind side an area around nostrils, and the greater part of exposed portion of preorbital, scaleless; fin rays of vertical fins,

all with bands of fine scales, those on caudal especially broad; lateral line with a broad arch in front, the cord of which is $3\frac{2}{3}$ in straight portion. Color dusky brownish, with 2 conspicuous pairs of round black spots narrowly edged with gray, the anterior pair about $\frac{1}{2}$ size of orbit, the posterior larger than pupil; the anterior pair under beginning of posterior third of dorsal, and about halfway between lateral line and dorsal and anal margins, respectively; the posterior pair nearer outline of body and about under the tenth before the last dorsal ray; vertical fins obscurely blotched with darker; ventral of eyed side with conspicuous black blotch margined with white, occupying the distal portion of its inner 2 rays; pectoral unmarked; membrane of gill cavity and peritoneum white. Two specimens, each about 12 inches long, from the Gulf of California, taken in 29 and 76 fathoms, at *Albatross* Stations 3014 and 3016. (Gilbert.) (*τετρα-*, four; *ὀφθαλμός*, eye, or eye-like spot.)

Lioglossina tetrophthalmus, GILBERT, Proc. U. S. Nat. Mus. 1890, 122, Gulf of California. (Coll. Dr. Gilbert.)

1023. XYSTREURYS, Jordan & Gilbert.

Xystreurys, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 34 (*liolepis*).

Body broad, covered with small smooth scales. Teeth rather small, uniserial and bluntly conical, unequal; no canines; caudal fin subsessile, the caudal peduncle extremely short; skin of shoulder girdle with patches of cup-shaped scales; lateral line with a strong anterior arch, no accessory branch; vertebræ about 37; gill rakers short and thick. This genus is very close to *Hippoglossina*, differing chiefly in the subsessile caudal fin, the smooth scales, and the peculiar, short, thick gill rakers. The typical species, like some other Parific coast flounders, is almost indifferently dextral or sinistral. (*ξύστρον*, raker; *εὐρύς*, wide, from the broad gill rakers.)

2990. XYSTREURYS LIOLEPIS, Jordan & Gilbert.

Head $3\frac{1}{5}$; depth $1\frac{5}{6}$. D. 80; A. 62; scales 123. Vertebræ $12+25=37$. Body elliptical ovate, broad and compressed, its curves regular; the profile continuous with curve of back; mouth small, very oblique, the lower jaw included; maxillary reaching about to pupil, $2\frac{2}{3}$ in head; eyes rather large, $4\frac{1}{2}$ in head, separated by a very narrow, blunt scaly ridge; teeth small, conical, blunt, in a single row; those in lower jaw subequal, close set; those in upper jaw more distant, decreasing in size backward; teeth $\frac{12+13}{14+15}$. Gill rakers $2+7$, very short, broad, and strong, minutely serrate on inner margin, about 7 below angle, the longest scarcely $\frac{1}{4}$ as long as the eye. Scales small, oblong, cycloid, the smaller accessory scales extremely numerous; lateral line without dorsal branch, with a broad curve above pectorals; branchial arches and skin of the shoulder girdle with small, cup-shaped, tubercular scales. Dorsal rather high, firm, low in front, beginning just in advance of middle of pupil, highest near the middle of the body; caudal peduncle very short and deep, its depth 4 times its length. Pectoral of eyed side falcate, usually much longer than head,

its length varying considerably. Caudal fin somewhat double truncate, with rounded angles, the middle rays being produced. Anterior nostril of blind side with a long flap. Color olive brown, mottled with darker, sometimes with very distinct round black blotches; vertical fins blotched with dark; pectoral of colored side with oblique bars. Length 15 inches. Southern California, rather common from Point Concepcion southward to San Diego. It is a very variable species, the coloration and the length of the pectoral fins having a wide range of variation. The body is indifferently dextral or sinistral. (*λεῖος*, smooth; *λεπίς*, scale.)

Xystreurys liolepis, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 34, Santa Barbara; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 454; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 66; JORDAN & GOSS, Review of Flounders and Soles, 243, 1889.
Paralichthys liolepis, JORDAN & GILBERT, Synopsis, 825, 1883.

1024. PARALICHTHYS, Girard.

(BASTARD HALIBUTS.)

Paralichthys, GIRARD, U. S. Pac. R. R. Surv., x, 146, 1858 (*maculosus* = *californicus*).
Pseudorhombus, BLEEKER, Comptes Rendus, Acad. Sci. Amsterd., XIII, 1862, 5, Notice sur quelques genres de la famille des Pleuronectidæ (*polyspilos*).
Uropsetta, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 330 (*californicus* = *maculosus*).
Chænopsetta, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 218 (*ocellaris* = *dentatus*).

Eyes and color normally on the left side. Body oblong; mouth large, oblique; each jaw with a single row of usually slender and sharp teeth, which are more or less enlarged anteriorly; no teeth on vomer or palatines. Gill rakers slender. Scales small, weakly ctenoid or ciliated; lateral line simple, with a strong curve anteriorly. Dorsal fine beginning before the eye, its anterior rays not produced; both ventrals lateral; caudal fin double truncate, or double concave, its middle rays produced; no anal spine. Species numerous, found in all warm seas. This genus, as now restricted, contains a considerable number of species, inhabiting both coasts of America and the eastern and southern coasts of Asia. As indicated by the reduced number of vertebræ, the species range further southward than do those of the type of *Hippoglossoides*. (*παράλληλος*, parallel; *ἰχθύς*, fish.)

a. Gill rakers in large number, about 9 + 20.
b. Gill rakers as long as eye and very slender. D. 72; A. 55; depth 2⅔ in length. CALIFORNICUS, 2991.
bb. Gill rakers shorter, about ⅔ length of eye. D. 80; A. 61; depth 2¼ in length. ÆSTUARIUS, 2992.
aa. Gill rakers in moderate number (5 + 11 to 6 + 21), rather long and slender.
c. Dorsal rays 70 to 75; anal rays 54 to 60.
d. Head small, lateral line 4½ in length; depth 2½; interorbital space rather broad and flatish, ⅔ diameter of eye; eyes small, 5⅔ in head; gill rakers rather short, 4 + 15, the longest about ⅔ eye. BRASILIENSIS, 2993.
dd. Head rather large, 3⅓ in length; depth 2 to 2¼; eyes small.
e. Gill rakers 5 to 6 + 15 to 18; eyes wide apart. ADSPERSUS, 2994.
ee. Gill rakers 5 + 11; eyes close together. WOOLMANI, 2995.
cc. Dorsal rays 85 to 93; anal rays 67 to 73; gill rakers 5 + 15 or 16, long and slender, the longest ⅔ length of eye; body ovate, the depth about 2⅔ in length; head about 3⅔. DENTATUS, 2996.

aaa. Gill rakers few, shortish, wide set, the number 2 + 8 to 3 + 10.

f. Body ovate, more or less compressed and opaque; depth about 2⅓ in length; no distinct, definitely placed ocelli; scales cycloid.

g. Dorsal rays in large number (85 to 93, as in *P. dentatus*); anal rays 65 to 73; pores of the lateral line about 100; accessory scales few; gill rakers 2 + 10, lanceolate, dentate, wide set, and much shorter than the eye. LETHOSTIGMUS, 2997.

gg. Dorsal rays in moderate number (70 to 80); anal rays 54 to 61.

h. Scales very small, about 120 in lateral line; depth of body about ⅓ length; head 3⅔ in length; gill rakers roughly toothed, 3 + 9 in number. SQUAMILENTUS, 2998.

hh. Scales moderate, 90 to 100 pores in the lateral line; interorbital width about equal to length of eye; dorsal rays 75 to 81; anal rays 59 to 61; gill rakers 2 or 3 + 9 or 10. Coloration, grayish brown with numerous (more or less distinct) whitish blotches, which are rarely obsolete; vertebræ 10 + 27 = 37. ALBIGUTTUS, 2999.

ff. Body oblong, strongly compressed, semitranslucent; scales weakly ciliated; about 93 pores in lateral line. Coloration, light grayish, thickly mottled with darker; 4 large horizontally oblong, black ocelli, each surrounded by pinkish area; 1 just behind middle of the body, below the dorsal, 1 opposite this, above anal, and 2 similar smaller spots below last rays of dorsal and above last of anal; vertebræ 11 + 30 = 41. OBLONGUS, 3000.

2991. PARALICHTHYS CALIFORNICUS (Ayres).

(BASTARD HALIBUT; MONTEREY HALIBUT.)

Head 3¾ to 4¼; depth 2⅔. D. 70; A. 55; scales 100. Vertebræ 10 + 25 = 35. Body rather long and thickish; caudal peduncle long; head small; eye small, little wider than the broad, flattish interorbital space; maxillary as long as pectoral, ⅓ length of head, reaching beyond eye; teeth slender, sharp, rather long, the canines moderate. Scales small, finely ciliate, each scale surrounded by narrow accessory scales; scales on blind side similar; fins with ctenoid scales. Dorsal low, beginning over front of upper eye just past pectoral, pointed, reaching curve of lateral line, 2¼ in head, that of blind side shorter and rounded behind; arch of lateral line 3½ or 4 in straight part. Gill rakers very long and slender, numerous, as long as eye, about 9 + 20; lower pharyngeals narow, with small slender teeth. Anal spine small, concealed. Grayish brown, uniform, or mottled with blackish and pale, the head sometimes sprinkled with black dots; young brownish, with bluish spots. Coast of California, Tomales Bay to Cerros Island. This large flounder is one of the common food-fishes of the Pacific coast, where it takes the place occupied on the Atlantic side by *Paralichthys dentatus*. It reaches a length of 3 feet and a weight of 60 pounds. From its resemblance to the halibut, it usually goes by the name of bastard halibut. It is readily distinguished from the Atlantic members of the same genus by its fewer fin rays and by its more numerous gill rakers. As was first shown by Mr. Lockington, the small fish called *Paralichthys maculosus*, is simply the young of the larger fish, then called *Uropsetta californica*. Unlike other species of the genus, *Paralichthys californicus* is almost as frequently dextral as sinistral. (*californicus*, Californian.)

Pleuronectes maculosus, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 155, young, San Diego.

Paralichthys maculosus, GIRARD, U. S. Pac. R. R. Surv., x, Fishes, 147, 1858, not *Rhombus maculosus*, CUVIER, also a species of *Paralichthys;* GÜNTHER, Cat., IV, 431, 1862; GILL, Proc. Ac. Nat. Sci. Phila. 1864, 197; LOCKINGTON, Rep. Com. Fisheries California 1878–79, 41; LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 79; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 454; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 66; JORDAN, Nat. His. Aquat. Anim., 182, 1884.

Hippoglossus californicus, AYRES, Proc. Cal. Ac. Nat. Sci. 1859, 29, and 1860, fig. 10, adult, San Francisco.

Pseudorhombus californicus, GÜNTHER, Cat., IV, 426, 1862.

Uropsetta californica, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 330; GILL, Proc. Ac. Nat. Sci. Phila. 1864, 198.

Paralichthys californicus, JORDAN & GILBERT, Synopsis, 821, 1883; JORDAN & GOSS, Review Flounders and Soles, 245, 1889.

2992. PARALICHTHYS ÆSTUARIUS, Gilbert & Scofield.

Head $3\frac{3}{8}$; depth $2\frac{1}{4}$; eye $5\frac{1}{2}$; interorbital space flat, 12 in head, $\frac{1}{2}$ diameter of eye; maxillary 2 in head, equal to pectoral fin; gill rakers 9 + 20, the longest $\frac{3}{8}$ length of eye. D. 72 to 83; A. 58 to 64. (In 7 specimens examined the rays are: Dorsal 72, 79, 81, 81, 82, 83, 83; anal 58, 60, 60, 62, 63, 63, 64.) Vertebræ 10 + 28; scales weakly ciliated, with small accessory scales, 105 in the lateral line; length of the arch contained 4 times in straight part of lateral line, 2 in head; height of arch $4\frac{1}{2}$ in head. Four of the 7 specimens are sinistral. Color pale chocolate brown. Specimens small, 6 to 9 inches in length. Taken at Shoal Point, at mouth of the Colorado River, Mexico, by the United States Fish Commission steamer *Albatross*. This species is distinguished from other members of the genus by its numerous fin rays and many gill rakers. It is nearest related to *Paralichthys californicus*. (*æstuarius*, pertaining to the river mouth.)

Paralichthys æstuarius, GILBERT & SCOFIELD, Proc. U. S. Nat. Mus. 1897, 499, pl. xxix, Gulf of California, at mouth of Colorado River, Sonora. (Type, No. 48128. Coll. C. H. Gilbert.)

2993. PARALICHTHYS BRASILIENSIS (Ranzani).

Head $4\frac{1}{3}$; depth $2\frac{1}{3}$. D. 70 to 75; A. 54 to 60; scales not very small, about 100 in course of lateral line; interorbital space rather broad and flattish, $\frac{3}{5}$ diameter of eye; eyes small, $5\frac{3}{5}$ in head; gill rakers rather short, 4 + 15, the longest about $\frac{3}{5}$ eye; pectoral $1\frac{1}{3}$ in head; curve of lateral line high and short, 4 in straight part, its height $1\frac{1}{4}$ in its length; mouth moderate, the maxillary $2\frac{1}{3}$ in head; teeth rather few, the anterior canines large. Color dark brown, more or less mottled and spotted with paler. South America; said to range northward to Guatemala. Here described from numerous specimens from Rio Janeiro and from Maldonado, in the Museum of Comparative Zoology. The locality "Guatemala" given by Günther seems to be somewhat doubtful, and the species may not occur in West Indian waters at all. (*brasiliensis*, living in Brazil.)

Hippoglossus brasiliensis, RANZANI, Nov. Spec. Pisc., 10, tab. 3, 1840, Brazil.

Platessa orbignyana, VALENCIENNES, D'Orbigny Voy. S. Amer. Mérid. Poiss., pt. 5, pl. 16, fig. 1, 1847.

Rhombus aramaca, CASTELNAU, Anim. nouv. ou rares, Poiss., 78, pl. 40, fig. 3; not of CUVIER.

Pseudorhombus vorax, GÜNTHER, Cat., IV, 429, 1862, South America.

Pseudorhombus brasiliensis, GÜNTHER, Fishes Centr. Amer., 473, 1869.

Paralichthys brasiliensis, JORDAN & GOSS, Review Flounders and Soles, 246, 1889.

2994. PARALICHTHYS ADSPERSUS (Steindachner).

Head $3\frac{1}{2}$; depth $2\frac{1}{4}$. D. 75; A. 58; scales 106; eye 6 in head; interorbital $\frac{3}{4}$ vertical diameter of eye; maxillary $2\frac{1}{3}$; mandible $1\frac{5}{6}$; pectoral 2; caudal $1\frac{3}{8}$. Body moderately elongate and compressed; mouth large, the maxillary reaching a little past eye; teeth large, sharp, and slightly recurved, larger in front of jaws; snout very slightly produced; interorbital moderately wide, its posterior half with scales; anterior nostril with a flap which reaches to middle of posterior nostril; gill rakers 3 to 6 + 15 to 17, hardly as long as eye. Snout and mandible naked; end of maxillary and rest of head with scales; the rays of all the fins with small scales; the membrane naked; each scale on body with a row of accessory scales around its posterior edge; scales cycloid, the accessory scales giving the fish a rough feeling; curve of lateral line nearly 5 in the straight part, pectoral reaching slightly past curve of lateral line, its tip pointed; pectoral of blind side shorter, not reaching to end of curve, its tip blunt; origin of dorsal over anterior edge of upper eye, bending slightly toward the blind side; caudal double lunate. Color brownish gray, thickly mottled with many larger and smaller spots, points, and rings; side with 3 or 4 larger spots of irregular form and ocellated with paler.

Specimens taken by Dr. Jordan at Mazatlan are described as follows: "Head $3\frac{1}{2}$; depth about 2 in length of body. D. 73 (70 to 76); A. 57 (53 to 60); P. 12; V. 6. Scales on lateral line about 106 + 8 with 35 dorsally and 36 ventrally. Flesh firm. Body oblong, moderately compressed; mouth large, oblique, the mandible very heavy, slightly projecting; 4 canine teeth on each side of lower jaw in adult specimens, 8 in young, the 2 anterior teeth long; anterior teeth of upper jaw strong, but smaller than those in the lower jaw; the lateral teeth very small and close set. Eye small, shorter than snout, about 7 (6 to 8) in length of head; interorbital area smooth, flattish, $\frac{3}{5}$ width of eye. Scales cycloid, small anteriorly and larger posteriorly; lateral line strongly arched anteriorly, arch about $3\frac{1}{3}$ in straight part. Gill rakers of medium length, broad, retrorse-serrate on inner side, longest about $\frac{2}{3}$ length of eye, from 4 + 13 to 5 + 14 in number, counted in 8 specimens; pectoral fin about as long as mandible, slightly more than $\frac{1}{2}$ length of head. Dorsal low, anterior origin opposite anterior margin of eye; caudal barely double concave; caudal peduncle very strong; anal spine obsolete; ventral fins small, inserted symmetrically; fins all scaly. Color: Large specimens are dark brown, with blotches on fins; small specimens are covered with pearly white and very dark brown blotches; the brown blotches almost circular, larger and with less definite outlines near the center of the body, very dark and distinct on caudal. Seven specimens were taken by the Hopkins Expedition in the estuary at Mazatlan, where they reach a length of 44 cm. Several specimens were also taken at La Paz. These specimens seem to be identical with *Paralichthys adspersus*. The original types have on an average more gill rakers than we find in our Mazatlan specimens, but this character is subject to variation, and no other distinction appears. In one of Dr. Steindachner's types from Callao (No. 11,417, Mus. Comp. Zool.) we find the gill rakers longer, 6 + 17; depth $2\frac{1}{4}$ in length; D. 67; A. 51; scales 120;

arch of lateral line barely twice as long as high, nearly 5 in straight part; maxillary $2\frac{1}{6}$ in head. Mr. Garman has kindly examined for us 6 other specimens, with the following results:

"*Paralichthys adspersus* from Callao, has gill rakers—

"'$\frac{7}{17}$ as long as eye;

"'$\frac{5}{15}$ about $\frac{2}{3}$ as long as the eye.

"'$\frac{7}{18}$ nearly as long as the eye.

"'$\frac{3}{14}$ about $\frac{2}{3}$ as long as the eye.

"'$\frac{5}{15}$ about $\frac{2}{3}$ as long as the eye.

"'$\frac{6}{17}$ near $\frac{3}{4}$ as long as eye.'" *

We are now disposed to regard these Mazatlan specimens as identical with *Paralichthys adspersus*, the range of variation in the number of gill rakers in the latter probably including the former. Pacific coast of tropical America, from Gulf of California to the coast of Peru; everywhere abundant and very variable. (*adspersus*, covered with spots.)

Pseudorhombus adspersus, STEINDACHNER, Ichthyol. Notizen, v, 9, pl. 2, 1867, Chinchas Islands.

Paralichthys adspersus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 370; JORDAN & GILBERT, Bull. U. S. Fish. Comm. 1882, 108 and 111; JORDAN, Cat. Fish N. A., 133, 1885; JORDAN & GOSS, Review Flounders and Soles, 246, 1889; JORDAN & WILLIAMS, Proc. Cal. Ac. Sci. 1895, 503.

2995. PARALICHTHYS WOOLMANI, Jordan & Williams.

Head $3\frac{1}{2}$; depth about 2; gill rakers 5+11. D. 74; A. 57; P. 12; V. 6; scales 100. Flesh firm; body oblong; mouth large, mandible heavy, not projecting; about 8 teeth on each side of lower jaw, the anterior ones long and slender; teeth in upper jaw smaller than those in lower jaw, the lateral teeth very small and close set. Eye small, $5\frac{1}{2}$ in length of head; interorbital area moderately prominent, narrow, about $\frac{2}{3}$ length of eye. Scales cycloid, small anteriorly and increasing in size posteriorly, covering head and fins; lateral line greatly arched anteriorly, arch about $3\frac{1}{2}$ times in length of straight portion. Gill rakers slender, the longest about $\frac{1}{2}$ in length of eye. Pectoral and ventral fins small; pectoral about $\frac{1}{3}$ in length of head; origin of dorsal opposite anterior margin of eye; caudal ending in an obtuse angle, not double concave; caudal peduncle strong; anal spine obsolete. Body and fins blotched with deep brown and pearly white and specked with very dark brown, blotches more definite on median fins and especially on caudal where there are 3 indefinite lines of blotches crossing the fin. Galapagos Islands. One specimen taken by the *Albatross* in 1890, which was at first identified as *Paralichthys adspersus*, from which species it differs but little except in the number of gill rakers. (Named for Mr. Albert Jefferson Woolman, of Duluth, Minnesota, in recognition of his work on the fishes of Mexico and Florida.)

Paralichthys woolmani, JORDAN & WILLIAMS, Proc. U. S. Nat. Mus. 1896, 457, Galapagos Islands. (Type, No. 47575. Coll. *Albatross*.)

* Garman, in lit., May 3, 1895.

2996. PARALICHTHYS DENTATUS (Linnæus).

(Summer Flounder.)

Head $3\frac{1}{2}$ to 4; depth $2\frac{2}{3}$; eye 6 in head; maxillary 2; pectoral $2\frac{1}{3}$; ventral $3\frac{1}{2}$; caudal peduncle 4; caudal $1\frac{1}{4}$. D. 86 to 91; A. 65 to 71; lateral line 108 (tubes). Curve of lateral line $3\frac{2}{3}$ to $4\frac{1}{3}$ in straight portion; body ovate; maxillary about $\frac{1}{2}$ head, reaching past posterior margin of eye; mouth large, oblique, the gape curved; canines large, conical, wide set; gill rakers comparatively long and slender, longest $\frac{2}{3}$ eye, 5 + 15 to 6 + 18 in number; interorbital area a rather flattish ridge, in the adult about equal to vertical diameter of eye, narrower in the young, forming a bony ridge; scales cycloid, each with numerous small accessory scales; vertebræ 11 + 30 = 41. Color in life, light olive brown; adults with very numerous small white spots on body and vertical fins; sometimes a series of larger white spots along bases of dorsal and anal fins; about 14 ocellated dark spots on sides, these sometimes little conspicuous, but always present; a series of 4 or 5 along base of dorsal, and 3 or 4 along base of anal, those of the 2 series opposite, and forming pairs; 2 pairs of smaller less distinct spots midway between these basal series and lateral line anteriorly, with a small one on lateral line in the center between them; a large distinct spot on lateral line behind middle of straight portion; fins without the round dark blotches. Atlantic coast of United States, from Cape Cod to Florida; the common flounder of the coasts of the Northern States, its range apparently not extending much south of Charleston. Of the species found in that region it is the most important from a commercial point of view. It reaches a length of about 3 feet and a weight of about 15 pounds. It has been confounded by nearly all writers with the more southern species now called *P. lethostigmus*, from which it is best distinguished by its much greater number of gill rakers and by its mottled coloration. On account of this confusion it is impossible wholly to disentangle its synonymy from that of *P. lethostigmus*. So far as the proper nomenclature of the two is concerned, this confusion makes little difference. There is no doubt that this is the original *Pleuronectes dentatus* of Linnæus, as the original Linnæan type is still preserved in London. This has been examined by Dr. Bean and its identity with the present species fully established. It seems also certain that this is the *Platessa ocellaris* of De Kay, who properly distinguishes his *ocellaris* from his *oblonga*, the latter being *P. lethostigmus*. A little doubt must be attached to the *P. melanogaster* of Mitchill, very scantily described from a doubled (black-bellied) example of this species or of *P. lethostigmus*. As the former species is much more common about New York than the latter it is probable that Mitchill's fish belonged to it. We have also received a doubled example from New York corresponding exactly to Mitchill's description. We may therefore regard the name *melanogaster* as a synonym of *dentatus*. The differences in the gill rakers of these species were first noticed by Jordan & Gilbert in 1883. These authors erroneously referred all these synonyms to the species with the few gill rakers and described the present one as new under the name *Paralichthys ophryas*. The discovery of the Linnæan type of *Pleuronectes dentatus* has rendered a reconsideration of this matter

necessary, and it is evident that to the "*P. ophryas*" belong also the prior names *dentatus, melanogaster,* and *ocellaris.* (*dentatus,* toothed.)

Pleuronectes dentatus, LINNÆUS, Syst. Nat., Ed. XII, 1, 458, 1766, and of numerous copyists; MITCHILL, Trans. Lit. and Phil. Soc. N. Y. 1815, 390.

Pleuronectes melanogaster, MITCHILL, Trans. Lit. and Phil. Soc. N. Y. 1815, 390, New York; doubled example.

Platessa ocellaris, DE KAY; N. Y. Fauna: Fishes, 300, pl. 47, fig. 152, 1842, New York.

Paralichthys ophryas, JORDAN & GILBERT, Synopsis, 822, 1883, Charleston.

Platessa dentata, STORER, Rept. Fish. Mass., 143, 1839.

Pseudorhombus dentatus, GOODE & BEAN, Proc. U. S. Nat. Mus. 1879, 123.

Paralichthys dentatus, GOODE, Nat. Hist. Aquat. Anim., 178, 1884, detailed account; includes *P. lethostigma*; JORDAN, Cat. Fish. N. A., 134, 1885; JORDAN & GOSS, Review Flounders and Soles, 246, 1889.

Pseudorhombus ocellaris, GÜNTHER, Cat., IV, 430, 1862; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1878, 370.

Chænopsetta ocellaris, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 218.

Paralichthys ocellaris, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 617.

2997. PARALICHTHYS LETHOSTIGMUS, Jordan & Gilbert.

(SOUTHERN FLOUNDER.)

Head $3\frac{3}{8}$; depth $2\frac{1}{4}$. D. 85 to 92; A. 65 to 73; pores about 100. Body ovate, more or less compressed and opaque; no distinct, definitely placed ocelli; scales cycloid. Mouth wide, oblique, the mandible very heavy and much projecting; 8 to 10 teeth on each side of the lower jaw, the 2 anterior teeth very long; anterior teeth of upper jaw strong, but smaller than those in the lower jaw; the lateral teeth very small and close set; eyes small, shorter than the snout, about 6 in head; interorbital space in adult broad, flattish, and scaly, as wide as length of eye. Accessory scales few; gill rakers 2 + 10, lanceolate, dentate, wide set, and much shorter than the eye; caudal peduncle rather long; length of arch of lateral line nearly $\frac{1}{3}$ that of straight part. Color dusky olive, darker than in *P. dentatus,* and with very few darker mottlings or spots. This species is the common large flounder of the South Atlantic and Gulf coasts of the United States, ranging as far north as New York. It very closely resembles *Paralichthys dentatus,* with which it has been repeatedly confounded. It is, however, sharply distinguished by the character of the gill rakers. It is also always darker in color, and almost uniform, while *P. dentatus* is usually profusely spotted. Its only tenable name is the recent one, *Paralichthys lethostigmus.* South Atlantic and Gulf coasts of United States, north to New York. (λήθη, forgetfulness; στίγμα, spot, from the absence of spots.)

Platessa oblonga, DE KAY, New York Fauna: Fishes, 299, pl. 48, fig. 156, 1842, New York, not *Pleuronectes oblongus,* MITCHILL; STORER, Syn. Fish. N. A., 477, 1846.

Paralichthys lethostigma, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1884, 237, Jacksonville, Florida; JORDAN & GOSS, Review Flounders and Soles, 247, 1889.

Pseudorhombus oblongus, GÜNTHER, Cat., IV, 426, 1862.

Chænopsetta dentata, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 218.

Pseudorhombus dentatus, GOODE, Proc. U. S. Nat. Mus. 1879, 110; GOODE & BEAN, Proc. U. S. Nat. Mus. 1879, 123.

Paralichthys dentatus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 302; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 617; BEAN, Cat. Coll. Fish. Proc. U. S. Nat. Mus. 1883, 45; JORDAN & GILBERT, Synopsis, 822, 1883.

2998. PARALICHTHYS SQUAMILENTUS, Jordan & Gilbert.

Head $3\frac{2}{3}$; depth 2. D. 78; A. 59; scales 123 (pores). Body deep, strongly compressed; caudal peduncle very short; profile angulated at front of upper eye. Head wide, the eyes large, wide apart. Mouth very large, oblique, the broad maxillary reaching well beyond pupil, its length more than $\frac{1}{2}$ the head. Lower jaw projecting; mandible with a sharp compressed knob at symphysis; teeth few, unequal, in a single row, about 8 in each jaw canine-like, the 2 in front of lower jaw longest; lateral teeth of upper jaw minute. Interorbital space a narrow scaleless bony ridge, slightly concave anteriorly, scarcely $\frac{1}{2}$ diameter of pupil. Scales very small, smooth, adherent; curve of lateral line $4\frac{1}{3}$ in straight part; snout, jaws, and preopercle naked. Gill rakers short, 3+9 in number, triangular, roughly toothed, little higher than wide, the longest nearly $\frac{1}{2}$ eye. Dorsal beginning over front of eye, the anterior rays $4\frac{1}{2}$ in head; pectoral short, shorter than maxillary; anal spine weak; caudal double rounded. Brownish; body and fins spotted with darker; caudal mottled with white; pectorals banded, with dark spots. South Atlantic and Gulf coasts of United States. This species is very close to *Paralichthys albiguttus*, from which it differs chiefly in the small scales. It seems to be rather rare. Besides the original types from Pensacola, another referred to the same species is in the National Museum from Charleston. (*squamilentus*, scaly.)

Paralichthys squamilentus, Jordan & Gilbert, Proc. U. S. Nat. Mus. 1882, 303, Pensacola (Type, No. 30862); Jordan & Gilbert, Synopsis, 823, 1883; Bean, Cat. Coll. Fish, U. S. Nat. Mus. 1883, 45; Jordan & Goss, Review Flounders and Soles, 248, 1889.

2999. PARALICHTHYS ALBIGUTTUS, Jordan & Gilbert.

(Gulf Flounder.)

Head $3\frac{3}{4}$; depth $2\frac{1}{3}$. D. 72 to 80; A. 59 to 61; scales 9 to 100 (pores); eye 6 or 7 in head; maxillary $1\frac{7}{8}$; pectoral $2\frac{1}{5}$; ventral 3; caudal $1\frac{1}{3}$; curve of lateral line 3 in straight part. Body moderately elongate-elliptical; mouth large, the maxillary reaching past eye; jaws subequal; teeth strong, slender, and curved, about 7 on side of lower jaw, 4 or 5 moderate canines in front of upper jaw, the lateral teeth being minute, close set; interorbital space $\frac{3}{8}$ length of eye, the upper ridge rather prominent behind upper eye, scaled posteriorly; mandibles naked; a small patch of scales on maxillary; gill rakers broad and toothed behind, the longest $2\frac{1}{2}$ in eye, 3+10 in number. Fins low; anterior rays of dorsal not elevated nor exserted, the longest rays behind the middle, $2\frac{3}{4}$ in head; pectoral not reaching to end of curve; caudal double lunate. Scales moderate, cycloid, covered with epidermis which bears small flaps about the borders of many of the scales. Dark olive, mottled with dusky, and marked by numerous more or less distinct pale spots, which are sometimes obsolete; three dark spots, bordered with white, sometimes present, particularly in the young, 1 on lateral line posteriorly and 1 above and below anterior end of straight part of lateral line. Vertebræ 10+27=37. South Atlantic and Gulf coasts of the United States. This species is common on the South Atlantic and Gulf coasts. It has the few gill rakers of *P. lethostigmus*, the mottled coloration of *P. dentatus*, while from each it is distinguished by its smaller number

of dorsal and anal rays. In the number of its vertebræ it agrees with *P. lethostigmus*. It seems to reach a smaller size than either of these species. Here described from a specimen, 16 inches in length, collected at Cedar Key, Florida. (*albus*, white; *gutta*, spot.)

Pseudorhombus dentatus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1878, 370; not of LINNÆUS.

Paralichthys albigutta, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 302, Pensacola (Type, No. 30818. Coll. Dr. Jordan); JORDAN & GILBERT, Synopsis, 823, 1883; JORDAN & SWAIN, Proc. U. S. Nat. Mus. 1884, 233; JORDAN & GOSS, Review Flounders and Soles, 248, 1889.

3000. PARALICHTHYS OBLONGUS (Mitchill).

(FOUR-SPOTTED FLOUNDER.)

Head 4; depth $2\frac{1}{4}$. D. 72; A. 60; scales 93. Body comparatively elongate, strongly compressed. Eyes large, nearly 4 in head, separated by a prominent, narrow, sharp ridge. Upper jaw with very numerous small, close-set teeth laterally, and 4 or 5 canines in front; the lateral teeth abruptly smaller than the anterior; each side of lower jaw with 7 to 10 teeth. Chin prominent. Maxillary narrow, reaching past middle of pupil, $2\frac{1}{4}$ in length of head. Gape curved; gill rakers short and toothed behind, 2 + 8. Scales weakly ctenoid or cycloid. Dorsal low, beginning over front of eye, some of the anterior rays exserted, but not elongate, the longest rays behind middle of fin, not quite $\frac{1}{2}$ head; caudal $1\frac{1}{4}$ in head; pectoral $1\frac{2}{3}$; anal spine obsolete. Grayish, thickly mottled with darker and somewhat translucent; 4 large, horizontally oblong, black ocelli, each surrounded by a pinkish area, 1 just behind middle of the body below the dorsal, 1 opposite this above anal, 2 similar smaller spots below last rays of dorsal and above last of anal. Coasts of New England and New York. This species is rather common on the coast of Cape Cod and the neighboring islands, but it has been rarely noticed elsewhere. The limits of its range are not yet definitely known. It is a very strongly marked species. Its translucency of coloration indicates that it lives in deeper water than the other species of the genus. Here described from specimen from Woods Hole.

Another specimen in our collection from Woods Hole, Massachusetts, referred to this species, shows the following characters: Brownish, somewhat mottled, without traces of ocelli (possibly faded); fins similar. Body rather elongate, slenderer than in other species and more compressed; mouth rather large, oblique, the lower jaw not projecting, the maxillary $2\frac{1}{3}$ in head, reaching to opposite posterior border of pupil; about 12 teeth on each side of lower jaw, the anterior rather long, about equal to anterior teeth of upper jaw; lateral teeth of upper jaw becoming gradually smaller posteriorly, much larger, less numerous, and more widely set than in other species of this genus. Eyes large, longer than snout, 4 to $4\frac{1}{2}$ in head, separated by a narrow, elevated, bony ridge, narrower than pupil, anteriorly scaleless, and curved behind the upper eye posteriorly. Scales moderate, cycloid, rather thin; curve of lateral line $4\frac{3}{4}$ in straight part. Gill rakers 2 + 8 in number, rather long and slender, about $4\frac{1}{2}$ in maxillary. Dorsal beginning above middle of eye, its anterior rays not longer than

others, the middle rays a little longer than longest of anal, which are about $\frac{1}{2}$ head; caudal as long as head; anal spine obsolete; ventrals small; pectoral $1\frac{3}{4}$ in head. Head $4\frac{1}{6}$; depth $2\frac{3}{5}$. D. 77; A. 63; scales 90. Length about 14 inches. (*oblongus*, oblong.)

Pleuronectes oblongus, MITCHILL, Trans. Lit. and Phil. Soc., I, 1815, 391, New York.
Platessæ quadrocellata, STORER, Proc. Boston Soc. Nat. Hist. 1847, 242; STORER, Hist. Fish. Mass., 397, pl. 31, fig. 3, Provincetown.
Chænopsetta oblonga, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 218.
Paralichthys oblongus, GOODE, Proc. U. S. Nat. Mus. 1880, 472; JORDAN & GILBERT, Synopsis, 824, 1883; JORDAN & GOSS, Review Flounders and Soles, 249, pl. 8, 1889; GOODE & BEAN, Oceanic Ichthyology, 436, 1896.

1025. RAMULARIA, Jordan & Evermann.

Ramularia, JORDAN & EVERMANN, new genus (*dendriticus*).

This genus is close to *Ancylopsetta*, differing mainly in the structure of the lateral line, the tubes of which are borne by series of smaller, concealed cycloid scales, the free edges of which are notched to the opening of the pore; these scales are concealed in the skin, and from the pores proceed backward membranaceous tubes which ramify over the bases of contiguous scales. Dorsal scarcely elevated in front; left ventral much produced. Body broad ovate, sinistral, with very rough scales. Gill rakers few, very broad. (*ramulus*, a branchlet, from the tubes of the lateral line.)

3001. RAMULARIA DENDRITICA (Gilbert).

Head $3\frac{5}{8}$ in length; depth $1\frac{3}{4}$. D. 84; A. 63; scales 100; 36 scales in a series upward and backward from lateral line. Body very broad, its depth $1\frac{3}{4}$ in length, the two outlines equally curved; profile not very strongly angulated in front of upper eye. Lower eye slightly in advance of upper; interorbital space a rather broad, convex, scaly ridge, about $\frac{1}{3}$ upper eye, which is contained about 5 times in head and is equal to snout. A blunt spine on snout on head of maxillary. Nostril openings very broad, without tube, the anterior with a narrow flap. Mouth moderate, very oblique, the gape curved, the maxillary reaching slightly beyond vertical from middle of lower eye, 3 in head. Teeth in a single, rather close-set series in each jaw, strong, conical, directed very obliquely inward, becoming gradually larger toward front of jaw, but not canine-like. Gill rakers very short, barely movable, as broad as long, strongly toothed, 6 on anterior limb. Dorsal beginning over middle of upper eye, the anterior rays partly free toward tips, but little, if any, elevated above those that follow, the first $2\frac{2}{3}$ in head; dorsal highest in its posterior third, the longest ray $2\frac{2}{5}$ in head; anal similar, the rays of posterior third of each fin slightly forked at tip; caudal peduncle deep and short, its depth about $\frac{1}{2}$ head, its length $\frac{1}{2}$ its depth; caudal rounded, almost double truncate; ventrals with narrow bases, the left one slightly in advance of the right; fin greatly produced, reaching far beyond front of anal, a trifle shorter than head; left pectoral $1\frac{3}{5}$ in head. Scales very strongly ctenoid, the edge spinous, the entire exposed portion rough; width of anterior arch of lateral line $3\frac{1}{2}$ in straight portion; tubes of lateral line borne by a series of smaller con-

cealed cycloid scales, the free edges of which are notched to the opening of the pore; these scales entirely covered by the integument, and from the pores there proceed backward membranaceous tubes, ramifying over the bases of contiguous scales; this is true also of lateral line of blind side; eyed side entirely scaled except snout and mandible. Vertical fins covered with thick skin, each ray accompanied by 1 or 2 series of ctenoid scales; left ventral also scaled. Color olive brown, with 3 large black ocellated spots larger than orbit, the posterior one on lateral line in front of caudal peduncle, the 2 anterior under middle of dorsal, halfway between lateral line and dorsal and anal outlines, respectively; each spot with a light center; distal portion of vertical fins more or less brown on right side. (Gilbert.) Gulf of California. A single specimen, 13 inches long, from *Albatross* Station 3022, in 11 fathoms. (*dendriticus*, like a tree, branched; *δένδρον*, tree.)·

Ancylopsetta dendritica, GILBERT, Proc. U. S. Nat. Mus. 1890, 121, Gulf of California at Albatross Station 3022, in 11 fathoms.

1026. ANCYLOPSETTA, Gill.

Ancylossetta, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 224 (*quadrocellata*).

Body sinistral, broadly ovate, the depth more than ½ length; mouth moderate; teeth uniserial, unequal, some of the anterior enlarged; caudal fin with a very short peduncle; scales very strongly ctenoid on both sides of the body; anterior rays of dorsal notably exserted, the rays of the anterior part of the fin elongate, thus forming a distinct lobe; gill membranes considerably united; gill rakers short and broad, with rough teeth; left ventral produced; vertebræ about 35. This genus is very close to *Paralichthys*, differing in the subsessile caudal fin, the short gill rakers, the rough scales, and in the prolongation of the anterior rays of the dorsal fin. (*ἄγκυλος*, hook; *ψῆττα*, turbot.)

3002. ANCYLOPSETTA QUADROCELLATA, Gill.

Head 3¾ to 3⅘; depth 1⅘. D. 70 to 76; A. 57 to 59; pores in lateral line 83 to 90; vertical series of scales 70; fourth or fifth dorsal ray longest, nearly ⅔ length of head. Caudal 1⅓ in head; ventral of colored side 1⅔. Body oval, compressed, very deep; an abrupt angle above eye; mouth very small, the maxillary reaching to below middle of orbit, 2½ in length of head; teeth comparatively small, about 14 on each side of lower jaw; no strongly differentiated canines in either jaw. Eyes moderate, separated by a very narrow, sharp, scaly ridge; gill rakers very short, thick, few in number, 2 + 6 or 7, the longest less than ½ diameter of pupil; scales rather small, very strongly ctenoid, those on blind side also rough; curve of lateral line rather low; tubes of lateral line simple; dorsal beginning in front of pupil, its anterior rays long and filiform, much exserted; caudal short and rounded, 1⅘ in head; ventral fin of colored side rather long, as long as pectoral, ½ length of head; anal spine wanting. Brownish olive, with 4 large, oblong, ocellated spots, the first above the arch of the lateral line; the 3 posterior forming an isosceles triangle, the hindmost

being on the lateral line; the ocellated spots are frequently furnished with a bright white center, and the sides and vertical fins have often a few scattered white spots; a small, indistinct, dark spot on middle of each eighth or tenth ray of dorsal and anal. Vertebræ 9 + 26 = 35. South Atlantic and Gulf coasts of the United States; not rare; a very handsome species. (*quadrocellatus*, having 4 ocelli.)

Ancylopsetta quadrocellata, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 224; not *Platessa quadrocellata*, STORER; JORDAN & GOSS, Review Flounders and Soles, 250, 1889.

Paralichthys ommatus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 616, Charleston; JORDAN & GILBERT, Synopsis, 824, 1883; JORDAN & SWAIN, Proc. U. S. Nat. Mus. 1884, 234; JORDAN & GOSS, Review Flounders and Soles, 250, 1889.

Pseudorhombus quadrocellatus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1878, 370.

1027. NOTOSEMA, Goode & Bean.

Notosema, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 192, 1883 (*dilecta*).

Body sinistral, elliptical in form, the caudal fin pedunculate. Mouth moderate, beneath the central axis of the body. Eyes large, close together, the upper one nearly encroaching upon the profile, the lower slightly in advance of the upper. Teeth in a single series in the jaws, about equally developed on each side, largest in front, absent on vomer and palatines. Pectoral fins somewhat unequal, that upon the blind side $\frac{2}{3}$ as large as the other; dorsal fin commencing slightly behind anterior margin of upper eye, the first 8 rays separated into a distinct subdivision of the fin, several of them being prolonged; caudal rounded, sinistral; ventral much elongated. Scales small, ctenoid on colored side of body; lateral line prominent, strongly arched, alike on both sides, the tubes simple. Gill rakers moderately numerous, rather stout, subtriangular, pectinate posteriorly. Pseudobranchiæ well developed. Vertebræ 35. This genus is scarcely distinct from *Ancylopsetta*, the body more elongate, the dorsal and ventral rays more produced. (*νῶτος*, back; *σῆμα*, banner.)

3003. NOTOSEMA DILECTUM (Goode & Bean).

Head $3\frac{1}{3}$; depth 2. D. 68; V. 6. A. 54 to 56; scales 48 (pores) on straight part of lateral line; width of interorbital area almost imperceptible; mandible reaching to middle of pupil of lower eye, its length 2 in head; upper jaw $2\frac{1}{3}$ times length of head. Origin of dorsal over anterior margin of eye, second and third rays the longest, which are 2 in greatest depth of body; anal beginning close to vent, its posterior rays longest; caudal pedunculate, double truncate; pectoral of eyed side subtriangular, its length $5\frac{1}{2}$ in length of body; ventral of eyed side much produced, its length more than 3 times that of its mate. Color dark brown, speckled with darker, 3 large subcircular ocellated spots nearly as large as eye, with white center, dark iris, narrow, light margin, and a brown encircling outline, these arranged in an isosceles triangle, the apex on the lateral line, the others before it and distant from the lateral line a distance equal to their own diameter; blind side white; fins blotched with dark brown. (Goode & Bean.) Gulf Stream. Known from the original

types obtained in the deep waters (75 fathoms) of the Gulf Stream, off the Carolina coast. (*dilectus*, delightful.)

Notosema dilecta, GOODE & BEAN, Bull. Mus. Comp. Zool., x, No. 5, 193, 1883, Gulf Stream off the coast of South Carolina; GOODE & BEAN, Oceanic Ichthyology, 437, 1896.

Paralichthys stigmatias, GOODE, Nat. Hist. Aquat. Anim., 182, 1884; by inadvertance for *dilectus*.

Ancylopsetta dilecta, JORDAN, Cat. Fish. N. A., 134, 1885; JORDAN & GOSS, Review Flounders and Soles, 250, 1889.

1028. GASTROPSETTA, B. A. Bean.

Gastropsetta, BARTON A. BEAN, Proc. U. S. Nat. Mus. 1894, 633 (*frontalis*).

Body oblong-ovate, highly arched in front, covered with small, cycloid, embedded scales; lateral line arched in front, deflected downward on caudal peduncle. Teeth small, in a single series in each jaw. Dorsal fin beginning in advance of eye, its anterior rays produced, not connected by the irregular and broadly fringed membrane. Gill rakers very short, almost as broad as long, few in number. Ventral of eyed side produced, ending in a long filamentous ray in the young. This genus is closely allied to *Ancylopsetta*, from which it differs in form of body, and especially in the entirely smooth scales, singularly branched and produced anterior dorsal rays, and very short and broad gill rakers. (γαστήρ, belly; ψῆττα, turbot or flounder.)

3004. GASTROPSETTA FRONTALIS, B. A. Bean.

Head $4\frac{3}{4}$; depth $2\frac{1}{2}$; middle caudal rays $2\frac{4}{5}$; eye large, $3\frac{3}{4}$ in head. D. 60; A. 48; V. 6; P. I, 10. Mouth of moderate size, maxillary $2\frac{1}{2}$ in head, the jaws curved; interorbital ridge prominent, very narrow. Dorsal beginning in front of eye on snout, its anterior rays singularly branched, the third and fourth longest, almost equaling length of head; anal fin beginning at vent, which is situated on blind side, its anterior rays scarcely produced; ventral of colored side much produced; middle caudal rays long. Color in spirits, light brown; 3 black spots on body, 2 along back, and 1 near anal base; fins with dusky blotches; several vertical stripes across eyes. A smaller specimen from *Albatross* Station 2317 has D. 62; A. 52; V. 6; P. I, 11. Gill rakers short, broad laminæ, 2 + 7. Teeth weak, uniserial. Anterior rays of dorsal greatly produced, the third $1\frac{1}{2}$ times as long as head. Ventral of eyed side very long, ending in a thread-like filament. Color as in the preceding. An example from *Albatross* Station 2373 near Apalachicola, is 224 mm. long; its depth 90 mm. D. 60; A. 49; P. I, 10; V. 6; C. 15. Vent situated in a deep notch, which forms the front margin of abdomen, and not on side, as in other specimens. Color darker than that of the Key West examples, being dark reddish brown; body spotted and fins blotched as in the preceding. Two specimens obtained by the *Albatross*, January 15, 1885, at Station 2317, Lat. 24° 25′ 45″ N., Long. 81° 46′ 45″ W., near Key West, Florida, in 45 fathoms of water, the type 8 inches long, the other one 6 inches. (B. A. Bean.) (*frontalis*, pertaining to the forehead.)

Gastropsetta frontalis, BARTON A. BEAN, Proc. U. S. Nat. Mus. 1894, 633, Key West. (Type, No. 37668, U. S. Nat. Mus. Coll. *Albatross*.)

1029. PLEURONICHTHYS, Girard.

Pleuronichthys, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 139 (*cœnosus*).
Heteroprosopon, BLEEKER, Comptes Rendus Acad. Amsterdam, XIII, 1862, 8 (*cornutus*).
Parophrys, GÜNTHER, Cat. Fishes, IV, 454, 1862; not of GIRARD.

Eyes and color on the right side. Body deep; head short, with very short, blunt snout; mouth small, with several series of slender, acute teeth, which are most developed on the blind side, and are often wanting in 1 or both jaws on the colored side; no teeth on vomer or palatines; lips thick, with several lengthwise folds within which is a series of short fringes. Lower pharyngeals narrow, each with a double row of very small teeth. Gill rakers wide set, very short and weak. Lateral line nearly straight, with a dorsal branch in our species. Scales small, cycloid, non-imbricate, embedded. Dorsal fin anteriorly twisted from the dorsal ridge toward the blind side; anal fin preceded by a spine; caudal fin convex behind. Intestinal canal elongate. Herbivorous species, feeding chiefly on algæ. Pacific Ocean. This well-marked genus contains 3 American species, which are very closely related to each other. The Asiatic species, *Platessa cornuta*, Schlegel, of the coasts of China and Japan, is also a member of this group, having an accessory branch to the lateral line as in the American species. This species bears some resemblance to *Pl. verticalis*. The species of *Pleuronichthys* spawn in the spring, and live in comparatively deep water. (*πλεῦρον*, side; *ἰχθύς*, fish.)

a. Dorsal fin beginning on the level of the lower lip, its first 9 rays on the blind side. DECURRENS, 3005.
aa. Dorsal fin beginning on level of upper lip, its first 5 rays being on the blind side.
b. Interorbital ridge posteriorly with a strong spine directed backward, some tubercles on interorbital ridge. VERTICALIS, 3006.
bb. Interorbital ridge prominent, but without spines and conspicuous tubercles. CŒNOSUS, 3007.

3005. PLEURONICHTHYS DECURRENS, Jordan & Gilbert.

Head $3\frac{1}{2}$; depth $1\frac{3}{4}$. D. 72; A. 40; scales 80; eye 3 in head; maxillary $4\frac{1}{2}$; pectoral $1\frac{3}{8}$; highest dorsal rays $1\frac{1}{2}$; anal rays $1\frac{3}{8}$; caudal 1. Body short and wide; mouth very small, the maxillary reaching nearly to pupil; teeth villiform, in moderate bands on blind side, a narrow band on eyed side of lower jaw; eyes very large, the upper edge of upper eye even with profile; snout extremely short; a blunt tubercle in front of upper eye, another at each end of the narrow interorbital ridge, the posterior largest, but usually not spine-like; 2 or 3 above the latter behind the upper eye; some prominences above the opercle; gill opening short, not extending above upper edge of pectoral. Dorsal beginning very low, on level of end of maxillary, its first 9 rays on the blind side; anal spine well developed, the origin of anal a little behind vertical from base of pectoral; pectoral of eyed side a little larger than its mate, both rounded behind; ventral of blind side shorter than that of eyed side, and placed slightly before it, caudal well rounded. Scales cycloid, embedded, a space between them anteriorly; lateral line without arch, slightly curved. Color brownish, usually much mottled with chocolate and grayish, often finely spotted with brownish on body and fins; all fins darker than body;

dorsal, anal, caudal, and ventrals narrowly edged with white; pectoral uniformly blackish. Pacific coast of United States, south to Monterey. This species is rather scarce along the California coast, being taken chiefly in deep water. It reaches a larger size than either *P. verticalis* or *P. cœnosus*. Here described from a specimen from San Francisco market, 8 inches in length. (*decurrens*, running down.)

Pleuronichthys cœnosus, LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 97; not *Pleuronichthys cœnosus*, GIRARD.

Pleuronichthys quadrituberculatus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 50, not of PALLAS; JORDAN, Nat. Hist. Aquat. Anim., 189, 1884.

Pleuronichythys decurrens, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 453, San Francisco; Monterey Bay (Coll. Jordan & Gilbert); JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 69; JORDAN & GILBERT, Synopsis, 829, 1883; JORDAN & GOSS, Review Flounders and Soles, 282, 1889.

3006. PLEURONICHTHYS VERTICALIS, Jordan & Gilbert.

Head 4 in body; depth 2. D. 65; A. 45; scales about 80; vetebræ 13 + 25 = 38. Form broad ovate, the outlines regular; head small, somewhat constricted behind the upper eye; eyes large, but smaller than in *P. decurrens*. Interorbital ridge narrow; a small tubercle or prominence in front of upper eye; a large one in front of upper edge of lower eye; another larger and sharper at interior edge of the interocular space; another at the posterior edge of interocular spine ridge; this latter developed into a long, sharp, triangular spine, which is nearly as long as the pupil, and is directed backward; a prominent tubercle at posterior lower angle of upper eye; upper edge of operele somewhat uneven, but no other tubercles present. Mouth small, as in other species; the lips thick, with lengthwise plicæ. Teeth in a broad band on the left (blind) side of each jaw; no teeth on the right side in either jaw. Gill rakers very small, weak, and flexible, about 10 in number. Scales essentially as in other species, small, cycloid, embedded, scarcely imbricated; lateral line nearly straight, with an accessory branch which extends to the middle of the dorsal fin. Dorsal fin beginning on blind side at level of premaxillary, there being but about 4 of its rays on left side of median line; vertical fins less elevated than in other species, the longest rays of dorsal about $\frac{1}{2}$ length of head; anal fin preceded by a spine; caudal peduncle short and deep; caudal fin elongate, rounded behind; pectorals short, nearly equal; ventrals moderate, reaching anal spine. Color dark olive brown, with round grayish spots, the body and fins mottled with blackish. This species agrees in habits and general characters with *Pleuronichthys decurrens*. Coast of California, in rather deep water. The above description from the original type. (*verticalis*, pertaining to the vertex.)

Pleuronichthys verticalis, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 49, San Francisco (Coll. Jordan & Gilbert); JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 169; JORDAN & GILBERT, Synopsis, 829, 1883; JORDAN, Nat. Hist. Aquat. Anim., 189, 1884; JORDAN & GOSS, Review Flounders and Soles, 282, 1889.

3007. PLEURONICHTHYS CŒNOSUS, Girard.

Head $3\frac{1}{4}$; depth 2. D. 68; A. 49; scales 61; eye 3 in head; pectoral $1\frac{1}{2}$; dorsal and anal rays $1\frac{1}{2}$; caudal a little longer than head. Body ovate; snout scarcely produced; mouth small, maxillary reaching past front of

lower eye; 3 or 4 rows of teeth on blind side of jaws, 1 on eyed side of lower; eyes very large; interorbital a high, narrow ridge, somewhat angulated behind, but with no conspicuous spine or tubercle; snout very short, about ½ eye; gill opening not extending above upper edge of pectoral. Scales cycloid, embedded, some distance apart anteriorly, their edges not in contact; lateral line nearly straight, with a long dorsal branch which reaches past middle of body. Dorsal and anal high; origin of dorsal on blind side on a level with premaxillary, its first 5 rays on blind side; origin of anal under base of pectoral; pectoral of eyed side a little larger than its mate; caudal well rounded. Color dark brown, usually mottled, the colors variable; our specimens from Puget Sound, very dark, the fins colored like body, with light and dark spots; a conspicuous black spot on lateral line on middle of sides. Pacific coast, from Sitka to San Diego. This species is comparatively common in rather deep water and about rocks, being most abundant about Puget Sound. Its apparent abundance as compared with the other species of the genus is doubtless due to its inhabiting shallower waters than they. It is quite variable in form. The above description from a specimen, 6 inches long, from Seattle. (*cœnosus*, muddy.)

Pleuronichthys cœnosus, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 139, San Francisco; GIRARD, U. S. Pac. R. R. Surv., x, Fishes, 151, 1858; LOCKINGTON, Rep. Com. Fisheries California, 1878–79, 45; LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 97; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 50; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 453; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 68; JORDAN & GILBERT, Synopsis, 830, 1883; JORDAN, Nat. Hist. Aquat. Anim., 189, 1884; JORDAN & GOSS, Review Flounders and Soles, 282, 1889; JORDAN, Proc. Cal. Ac. Sci. 1895, 852.

Parophrys cœnosa, GÜNTHER, Cat., IV, 456, 1862.

1030. HYPSOPSETTA, Gill.

(DIAMOND FLOUNDERS.)

Hypsopsetta, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 195 (*guttulatus*).

Eyes and color on the right side; body broad, ovate, rhomboid; mouth very small; teeth slender, equal, acute, in several series; lips thick, not plicate; lateral line nearly straight, with an accessory dorsal branch; scales small, smooth; dorsal fin beginning on the dorsal ridge, not turned to the blind side at its insertion; anal spine present; caudal fin convex; gill rakers little developed. This genus consists of a single species, abundant on the coast of California. It is very close to *Pleuronichthys*, from which it differs only in a few characters of comparatively minor importance. Its range is in shallower and warmer water than that of the species of *Pleuronichthys*, and, in accordance with this fact, its flesh is firmer and its number of vertebræ fewer than in the latter genus. (*ὕψι*, deep; *ψῆττα*, flounder.)

3008. HYSOPSETTA GUTTULATA (Girard).

(DIAMOND FLOUNDER.)

Head 3⅘; depth 1⅔. D. 68; A. 50; scales 95. Body very deep, somewhat angulated near middle of back and belly; eyes moderate, separated by a flattish, raised area; head without spines or tubercles; scales of opercular

region little developed; those of blind side reduced; no teeth on right side of either jaw; accessory lateral line long, ½ length of body; anal spine small; pectorals about ½ length of head; ventrals rather short; caudal peduncle much deeper than long; caudal large, nearly as long as head. Brown, with numerous pale-bluish blotches in life, these disappearing in spirits; blind side white, with a strong tinge of yellow along profile of head; fins plain, sometimes with black specks. Coast of California and southward, Cape Mendocino to Magdalena Bay. This species is one of the most abundant in the shore waters of the California coast. It is a food-fish of fair quality. (*guttulatus*, with small spots.)

Pleuronichthys guttulatus, GIRARD, Proc. Ac. Nat. Sci. Phila. 1856, 137, Tomales Bay, California (Coll. E. Samuels); GIRARD, Journ. Boston Soc. Nat. Hist. 1857, pl. 25, figs. 1–4; GIRARD, U. S. Pac. R. R. Surv., X, Fishes, 152, 1858; LOCKINGTON, Rep. Com. Fisheries California, 1878–79, 44; LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 94.

Parophrys ayresi, GÜNTHER, Cat., IV, 1862, 457, San Francisco. (Coll. Dr. W. O. Ayres.)

Pleuronectes guttulatus, GÜNTHER, Cat., IV, 445, 1862.

Hypsopsetta guttulata, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 195; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 453; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 68; JORDAN & GILBERT, Synopsis, 830, 1883; JORDAN, Nat. Hist. Aquat. Anim., 185, 1884; JORDAN & GOSS, Review of Flounders and Soles, 283, 1889.

1031. PAROPHRYS, Girard.

Parophrys, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 139 (*vetulus*).

Body rather elongate, covered with small, cycloid scales; scales of the head roughish. Head rather pointed; mouth small, the teeth uniserial, all more or less blunt, compressed, incisor-like, close set. Lateral line with an accessory dorsal branch; upper eye on median line of top of head. A single species, on the Pacific coast of America. The narrow interorbital space and the vertical range of the upper eye give it a peculiar physiognomy, but in most regards it is not very different from some of the species of *Pleuronectes*. (παρά, near; ὀφρύς, eyebrow, from the narrow interbital.)

3009. PAROPHRYS VETULUS, Girard.

Head 3½; depth 2½; eye 4½ in head. D. 74 to 86; A. 54 to 68; scales 105 (tubes). Body elongate-elliptical; snout very prominent, much protruding, forming an abrupt angle with the descending profile; depth of head opposite middle of upper eye about equaling distance from middle of orbit to snout; eyes large, separated by a very narrow, high ridge, the upper with vertical range; mouth very small; maxillary not reaching pupil; teeth trenchant, small, and rather narrow, widened at tip, about 45 teeth on left side of lower jaw; few teeth on right side of lower jaw. Accessory lateral line long. Pectoral about ½ length of head; caudal truncate, 1½ in head; fin rays entirely scaleless; scales on body all cycloid, those on cheeks often slightly ciliated. Uniform light olive brown; the young somewhat spotted with blackish. Pacific coast of North America, Sitka to Santa Barbara. This small flounder lives in waters of moderate depth. It is, next to *Platichthys stellatus*, probably the most abundant of the flounders of the California coast. (*vetulus*, an old man.)

Parophrys vetulus, GIRARD, Proc. Ac. Nat. Sci. Phila. 1854, 140, California; GÜNTHER, Cat., IV, 455, 1862; LOCKINGTON, Rep. Com. Fish. Cal. 1878–79, 45; LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 100; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 453; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 68; JORDAN, Nat. Hist. Aquat. Anim., 185, 1884; JORDAN & GOSS, Review Flounders and Soles, 284, 1889.

Pleuronectes digrammus, GÜNTHER, Cat., IV, 445, 1862, Victoria. (Coll. Earl Russell.)

Parophrys hubbardi, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 281, San Francisco.

Pleuronectes vetulus, JORDAN & GILBERT, Synopsis, 831, 1883.

1032. INOPSETTA, Jordan & Goss.

Inopsetta, JORDAN & GOSS, in JORDAN, Cat. Fish. N. A., 136, 1885 (*ischyrus*).

This genus resembles *Parophrys*, differing chiefly in having the scales less imbricated, all strongly ctenoid, and having the eyes both lateral, the snout much less acute than in *Parophrys*. A single species, closely allied to *Platichthys stellatus*, but separated from it by the curious character common to many of our Pacific coast flounders, of having an accessory branch to the lateral line. (ἴς, strength; ψῆττα, flounder.)

3010. INOPSETTA ISCHYRA (Jordan & Gilbert).

Head 3¼; depth 2. D. 70 to 76; A. 52 to 57; V. 6; scales 85. Body oblong, robust; caudal peduncle rather long; snout projecting, forming an angle with the profile; mouth oblique, the chin projecting; teeth $\frac{5+25}{10+22?}$ narrowly incisor-like, bluntish, in a single, rather close-set series; maxillary reaching past front of orbit, 5⅔ in head; eyes large; interorbital space rather broad, scaly, continuous with a ridge above opercle; head mostly covered with scales like those of the body, but smaller and rougher; gill rakers feeble; lower pharyngeals each with 2 rows of coarse, blunt teeth; scales thick and firm, adherent, not closely imbricated, those in front well apart; all the scales strongly ctenoid; blind side with similar scales, almost as strongly ctenoid; vertical fins mostly scaly; lateral line conspicuous, its scales less rough than the others; a distinct short accessory lateral line on both sides, extending to about the tenth dorsal ray, less than ⅙ head; a series of pores around lower eye behind; dorsal beginning over pupil, its anterior rays low, its highest rays nearly ½ length of head; caudal large, double truncate; pectoral of right side about ½ head. Light olive-brown, vaguely clouded with light and dark; fins reddish brown; a few roundish dusky blotches on dorsal and anal; pectoral and caudal tipped with dusky; blind side white, immaculate, or with small, round rusty spots; left side of head sometimes rusty tinged. Puget Sound. This species is known only from 4 specimens taken by Dr. Jordan at Seattle in 1880. It is a large, rough flounder, with firm, white flesh. (ἰσχυρός, robust.)

Parophrys ischyrus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 276 and 453, Puget Sound (Coll. Dr. Jordan); JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 67; JORDAN, Nat. Hist. Aquat. Anim., 185, 1884.

Pleuronectes ischyrus, JORDAN & GILBERT, Synopsis, 832, 1883.

Inopsetta ischyra, JORDAN, Cat. Fish. N. A., 136, 1885; JORDAN & GOSS, Review Flounders and Soles, 284, 1889.

1033. ISOPSETTA, Lockington.

Isopsetta, LOCKINGTON MS., in JORDAN & GILBERT, Synopsis, 832, 1883 (*isolepis*).

Body much compressed, elliptical in form; mouth rather large; the teeth chiefly uniserial, all more or less blunt, separated, not incisor-like; scales closely imbricated, all strongly ctenoid; lower pharyngeals each with a double row of bluntish teeth. A single species found on the coast of California. *Isopsetta* approaches in many respects very close to the large-mouthed flounders of the type of *Hippoglossoides*, and it may fairly be said to be intermediate between *Psettichthys* and *Lepidopsetta*. Its affinities on the whole are nearest the latter, but the close relation of the *Hippoglossinæ* and *Pleuronectinæ* is clearly shown. (ἴσος, equal; ψῆττα, flounder.)

3011. ISOPSETTA ISOLEPIS (Lockington).

Head 4; depth 2¼. D. 88; A. 65; scales 88; vertebræ 10 + 32 = 42. Body elliptical, much compressed, moderately deep, the curvature very regular; head moderate, strongly compressed, the profile little depressed above the eye; eyes rather large; interorbital space broad, flattish, with several series of scales. Scales on cheeks similar to those on body, rather large, ctenoid, and closely imbricated. Mouth comparatively large, maxillary reaching pupil, 3⅜ in head; teeth not large, about $\frac{11+14}{9+24}$, conical, close set, in 1 somewhat irregular series, or partly in 2 series, those on colored side small; lower pharyngeals each with a double row of bluntish teeth. On the blind side the scales are more or less ctenoid, sometimes smooth; those on the cheeks weakly ctenoid; most of the opercle, the preopercle, interopercle, and subopercle on blind side naked; lateral line with a very slight arch in front, the depth of which is less than ⅓ the length; accessory branch nearly as long as head; fins rather low, mostly covered with ctenoid scales. Color brownish, mottled and blotched with darker. This small flounder is rather common off the coast of California, where it reaches a length of about 15 inches. It much resembles *Psettichthys melanostictus*, but its small mouth and blunt dentition indicate a real affinity with the small-mouthed flounders, among which it is here placed. Its nearest relative among our species is doubtless *Lepidopsetta bilineata*. Puget Sound to Point Concepcion, in rather deep water; not rare. (ἴσος, equal; λεπίς, scale.)

Lepidopsetta umbrosa, LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 106; not of GIRARD.
Lepidopsetta isolepis, LOCKINGTON, Proc. U. S. Nat. Mus. 1880, 325, San Francisco.
Parophrys isolepis, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 453 and 1881, 67; JORDAN & GILBERT, Synopsis, 832, 1883; JORDAN, Nat. Hist. Aquat. Anim., 186, 1884.
Isopsetta isolepis, JORDAN, Cat. Fish. N. A., 136, 1885; JORDAN & GOSS, Review Flounders and Soles, 285, 1889.

1034. LEPIDOPSETTA, Gill.

Lepidopsetta, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 195 (*umbrosus*).

Body robust; mouth small. Teeth stout, conical, little compressed, bluntish, in 1 series, rather irregularly placed. Lateral line with a distinct arch in front and accessory dorsal branch; scales imbricated,

rough ctenoid, smooth in the very young. A single species, abundant on the Pacific coast of North America. It is close to *Inopsetta*, from which it is separated by the arch of the lateral line, and still closer to *Limanda*, from which the accessory branch of the lateral line alone separates it. (*λεπίς*, scale; *ψῆττα*, flounder.)

3012. LEPIDOPSETTA BILINEATA (Ayres).

Head $3\frac{3}{8}$; depth $2\frac{1}{8}$. D. 80; A. 60; teeth $\frac{27+7}{25+10}$; scales 85. Vertebræ 11 + 29 = 40. Body broadly ovate, thickish; mouth moderate, turned toward the left side; teeth stout, conical, little compressed, bluntish, in 1 series, rather irregularly placed. Lower pharyngeals broad, with 2 rows of blunt teeth. Gill rakers few, very short, thick and weak, without teeth. Snout projecting; eyes large, separated by a prominent ridge, which, like the cheeks and upper portion of opercle, is covered with rough stellate scales; lower eye advanced; opercle, subopercle, and interopercle of left side scaly; preopercle naked. Scales rather small, mostly ctenoid, not closely imbricated, those on the blind side smooth; scales on cheeks and other parts of head very rough; scales of body smoother and less closely imbricated anteriorly, the degree of roughness variable, northern specimens (var. *umbrosus*) being roughest. Lateral line moderately arched anteriorly, with an accessory dorsal branch, which is less than $\frac{1}{2}$ length of head; height of arch less than $\frac{1}{3}$ its length. Dorsal beginning over eye, its anterior rays low; caudal convex; anal preceded by a spine; a concealed spine behind ventrals; rays of dorsal and anal all simple; dorsal and anal somewhat scaly; caudal $\frac{3}{7}$ length of head; pectoral $\frac{1}{3}$ head. Lower pharyngeals broad, each with 2 rows of blunt teeth. Yellowish brown, with numerous round pale blotches. Pacific coast of North America, Bering Strait to Monterey. This species is one of the commonest of the flounders of the Pacific coast, its abundance apparently increasing toward the northward. In Bering Sea it far outnumbers all other flounders. We have specimens from Bering Island, Medni Island, Unalaska, St. Paul, St. George, and Chignik Bay. It reaches a weight of 5 or 6 pounds and is an inhabitant of shallow waters. Specimens from Puget Sound and northward are rougher than southern specimens and constitute a slight geographical variety, for which the name *Lepidopsetta bilineata umbrosa* may be used. This is the same as *P. perarcuatus* of Cope. (*bilineatus*, two-lined.)

Platessa bilineata, AYRES, Proc. Ac. Nat. Sci. Cal. 1855, 40, San Francisco.

Platichthys umbrosus, GIRARD, Proc. Ac. Nat. Sci. Phila. 1856, 136, Puget Sound.

Pleuronectes perarcuatus, COPE, Proc. Ac. Nat. Sci. Phila. 1873, 30, Unalaska.

Pleuronectes umbrosus, GÜNTHER, Cat., IV, 454, 1862.

Pleuronectes bilineatus, GÜNTHER, Cat., IV, 444, 1862; JORDAN & GILBERT, Synopsis, 833, 1883.

Lepidopsetta bilineata, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 195; LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 103; LOCKINGTON, Rep. Com. Fisheries California, 1878–79, 46; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 453; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 68; BEAN, Proc. U. S. Nat. Mus. 1881, 241; BEAN, Cat. Coll. Fish. U. S. Nat. Mus. 1883, 19; BEAN, Proc. U. S. Nat. Mus. 1883, 353; JORDAN, Nat. Hist. Aquat. Anim., pl. 50, 184, 1884; JORDAN & GOSS, Review Flounders and Soles, 286, 1889.

1035. LIMANDA, Gottsche.

(Mud Dabs.)

Limanda, Gottsche, Archiv fur Naturgsch. 1835, 100 (*limanda*).
Myzopsetta, Gill, Proc. Ac. Nat. Sci. Phila. 1864, 217 (*ferruginea*).

Teeth chiefly uniserial; lateral line with a distinct arch in front, and without accessory dorsal branch; scales imbricated, rough ctenoid; vertebræ about 40. This genus is closely allied to *Pseudopleuronectes*, from which it differs only in the presence of an arch on the anterior part of the lateral line. (*Limanda*, an old name of the European Dab, *Limanda limanda*, from *limus*, mud.)

a. Head comparatively large, 3¼ to 4⅓ in length.
b. Dorsal rays 85; anal rays 62.
c. Scales rather small, 90 to 100 in lateral line; scales of right side ctenoid, closely imbricated, those of blind side mostly smooth; teeth conical, close set, forming a continuous series, about 11 + 30 in lower jaw; snout abruptly projecting, forming in front of upper eye a sharp angle with the descending profile. FERRUGINEA, 3013.
cc. Scales larger, wide apart, about 80 in lateral line; scales of blind side more or less rough. ASPERA, 3014.
bb. Dorsal rays 60 to 70; anal 47 to 53; scales small, 86 to 95; snout long, protruding; scales of blind side smooth. PROBOSCIDEA, 3015.
aa. Head very short, 5⅓ in length; snout very short; interorbital space very narrow. D. 64; A. 63; scales 88. BEANII, 3016.

3013. LIMANDA FERRUGINEA (Storer).

(Rusty Dab.)

Head 4 in length; depth 2⅓. D. 85; A. 62; scales 100. Body ovate-elliptical, strongly compressed; teeth small, conical, close set, in a single series on each side in each jaw, about 11 + 30 in the lower jaw; snout projecting, forming a strong angle above upper eye, with the descending profile; gill rakers of moderate length, very weak, not toothed; eyes moderate, 4½ in head, the lower slightly in advance of upper, separated by a high, very narrow ridge, which is scaled posteriorly, and is continued backward as an inconspicuous but rough ridge to the beginning of the lateral line; scales imbricate, nearly uniform, those on right side rough ctenoid, those on left side nearly or quite smooth; scales on body rougher than on cheeks; caudal peduncle short, higher than long; dorsal inserted over middle of eye, its middle rays highest; pectoral less than ⅔ length of head; caudal fin rounded; anal spine present; lateral line simple, with a rather low arch in front, the depth of which is barely ⅔ the length; a concealed spine behind ventrals; ventral of colored side partly lateral, the other wholly so; anal spine strong. Brownish olive, with numerous, irregular, reddish spots; fins similarly marked; left side with caudal fin, caudal peduncle and margins of dorsal and anal fins lemon yellow. Atlantic coast of North America, Labrador to New York. This species is rather common northward on our Atlantic coast. It is allied to the European Dab, but has smaller scales and a more prominent snout. Our specimens are from the east coast of Massachusetts. (*ferrugineus*, rusty red.)

Platessa ferruginea, D. H. STORER, Rept. Fish. Mass., 141, pl. 2, 1839, Cape Ann; DE KAY New York Fauna: Fishes, 297, pl. 48, fig. 155, 1842; STORER, Syn. Fish. N. A., 476, 1846.
Platessa rostrata, H. R. STORER, Bost. Journ. Nat. Hist., VI, 1850, 268, Labrador.
Pleuronectes ferrugineus, GÜNTHER, Cat., IV, 447, 1862; JORDAN & GILBERT, Synopsis, 834, 1882.
Myzopsetta ferruginea, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 217.
Limanda ferruginea, GOODE, Proc. U. S. Nat. Mus. 1880, 472; GOODE, Hist. Aquat. Anim., pl. 49, 1884; GOODE & BEAN, Oceanic Ichthyology, 427, 1896; JORDAN & GOSS, Review Flounders and Soles, 287, 1889.
Limanda rostrata, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 217.

3014. LIMANDA ASPERA (Pallas).

(ALASKA DAB.)

Head 3½; depth 2. D. 69; A. 53; scales about 80. Form of *Lepidopsetta bilineata*. Teeth small, almost conical, on both sides of the mouth; interorbital space narrow, scaly; opercle and preopercle naked below; gill rakers very feeble; pharyngeals not very broad, their teeth bluntish, not paved; scales small, wide apart, partly embedded, each one with 1 to 4 spinules, which are almost erect; anterior scales with 3 to 4 of these spinules; posterior mostly with 1; scales of blind side smoother; only middle rays of dorsal and anal scaly; no accessory lateral line; anal spine present; twentieth anal ray and thirty-seventh dorsal ray longest; caudal double truncate. Brown, nearly plain, the blind side with tinges of lemon yellow. Bering Sea, generally common, south to Vancouver Island and to the Okhotsk Sea. We have specimens from Petropaulski and Robben Reef, Bristol Bay, and Herendeen Bay. It is especially abundant in Bristol Bay, and, according to Dr. Gilbert, it is an excellent food-fish. Dr. Bean has also collected it in various localities in Alaska. Its scales are larger and rougher than in *L. ferruginea* which, in many respects, it resembles. A specimen from the island of Saghalien is in the museum at Cambridge. The above description is from examples taken by Dr. Bean. (*asper*, rough.)

Pleuronectes asper, PALLAS, Zoogr. Rosso-Asiat., III, 425, 1811, east coast of Siberia; GÜNTHER, Cat., IV, 454, 1862; STEINDACHNER, Pleuronectiden, etc., aus Decastris Bay, 1870–1875; JORDAN & GILBERT, Synopsis, 835, 1883.
Limanda aspera, BEAN, Proc. U.S. Nat. Mus. 1881, 242; BEAN, Cat. Coll. Fish, U.S. Nat. Mus. 1883, 20; BEAN, Proc. U. S. Nat. Mus. 1883, 354; BEAN, Hist. Aquat. Anim., 184, pl. 48, 1884; JORDAN & GOSS, Review Flounders and Soles, 288, 1889.

3015. LIMANDA PROBOSCIDEA, Gilbert.

Depth 2¼ to 2½ in length; head large, 3 to 3⅓ in length in a specimen 7 inches long. D. 63 to 67; A. 47 to 49; scales 86 to 95. Resembling *L. ferruginea*, but having fewer rays in dorsal and anal, larger scales and longer snout. Profile sharply angulated above front of upper eye, the snout convexly protruding; form varying from very slender to broadly elliptical, the 2 outlines equally curved; caudal peduncle short, widening backward, its least depth twice its length; mouth oblique, maxillary reaching beyond front of lower eye, 4 in head; teeth narrow, little compressed, in a single series on both sides of the jaw, extending farther back on the blind side; eyes on right side; lower eye well in advance of upper, the diameter of

upper eye 5½ to 6 in head, 1½ in snout; vertical from front of upper eye, falling midway between front of orbit and front of pupil of lower eye; interorbital space a very narrow, sharp ridge, naked in females, with a single series of ctenoid scales in males; gill rakers short, about equal to diameter of pupil, 13 or 14 in number, 9 or 10 on lower limb; scales loosely imbricated, ctenoid in males on colored side, smooth in females; blind side of both sexes smooth; head scaled on eyed side in males; the opercle, subopercle, interopercle, and preopercle mostly naked in females; head on blind side naked; rays of vertical fins with a single series of ctenoid scales; dorsal fin beginning slightly behind front of upper eye, the first 3 rays usually higher and with membranes more deeply incised than in those which follow; highest portions of both dorsal and anal fins behind the middle of the body; these fins about equal, their longest rays equal to the snout and eye; caudal ⅘ head; pectorals short, ⅓ in head; ventrals reaching beyond front of anal, 3⅓ in head; the usual small antrorse spine in front of anal fin. Color light grayish or brownish, thickly covered with small whitish spots; entire left side with margins of dorsal, caudal, and anal fins bright lemon yellow (as in *ferruginea*); vertical fins grayish, with an occasional dark-brown ray. Specimens described 7½ inches long. Bering Sea; several specimens from *Albatross* Stations 3239 and 3240, in Bristol Bay, in 11½ to 14½ fathoms; 1 young individual from Herendeen Bay. (Gilbert.) (*proboscideus*, having a long snout or proboscis.)

Limanda proboscidea, GILBERT, Report U. S. Fish Comm. 1893 (1896), 460, pl. 33, Bristol Bay and Herendeen Bay. (Coll. *Albatross*.)

3016. LIMANDA BEANII, Goode.

Head 5½; depth 2⅗. D. 64; A. 63; scales 88. Body elliptical, with angular outlines, strongly compressed; head very short; snout abbreviated; mouth small, subvertical; teeth small, apparently in two rows, chiefly on the blind side of lower jaw; eyes large, as long as mandible; interorbital space very narrow. Dorsal fin beginning about pupil, its rays long, wide apart, exserted; right ventral near the median line; caudal broad, fan-shaped. Lateral line with an abrupt curve, the length of which is twice its height and about equal to length of head, its scales highly specialized; lateral line on colored side less developed; scales small, strongly ctenoid on the right side; larger and cycloid on the blind side. Grayish, mottled with darker; a conspicuous black blotch on the outer rays of caudal on each side. (Goode.) Deep water off the coasts of New England; not common. (Named for Dr. Tarleton Hoffman Bean.)

Limanda beanii, GOODE, Proc. U. S. Nat. Mus. 1880 (Feb. 16, 1881), 473, southern coast New England, Fish Hawk Stations, 875, 876; GOODE & BEAN, Oceanic Ichthyology, 428, pl. 102, figs. 355a and 355b, 1896.

Pleuronectes beani, JORDAN & GILBERT, Synopsis, 835, 1883; JORDAN & GOSS, Review Flounders and Soles, 288, 1889.

1036. PSEUDOPLEURONECTES, Bleeker.
(WINTER FLOUNDERS.)

Pseudopleuronectes, BLEEKER, Comptes Rendus Acad. Amst., Pleuron., 7, 1862 (*planus*).

Body oblong, with firm flesh; the scales firm, regularly imbricated, strongly ctenoid on eyed side in both sexes; fin rays scaly; mouth small;

teeth uniserial, incisor-like, close set, all more or less blunt; lower pharyngeals very narrow, each with 2 rows of separate, conical teeth. This genus is distinguished from *Pleuronectes* chiefly by the well-imbricated ctenoid scales, and from *Limanda*, which it more closely resembles, by the want of arch to the lateral line. Besides the typical species, we refer to this genus a second from the North Pacific. (*ψεῦδος*, false; *Pleuronectes*.)

a. Dorsal rays 65; anal rays 48; scales 83; vertical fins nearly plain.
AMERICANUS, 3017.

aa. Dorsal rays 58; anal rays 38; scales 70; vertical fins with black bars.
PINNIFASCIATUS, 3018.

3017. PSEUDOPLEURONECTES AMERICANUS (Walbaum).

(Common Flatfish; Winter Flounder.)

Head 4 in length; depth 2¼. D. 65; A. 48; scales 83. Body elliptical; an angle above eye. Head covered above with imbricated, strongly ctenoid scales, similar to those on the body; blind side of head nearly naked; interorbital space rather broad, strongly convex, its width ½ eye, entirely scaled; teeth compressed, incisor-like, widened toward tips, close set, forming a continuous cutting edge; some of teeth often emarginate, sometimes movable; right side of each jaw toothless. Highest dorsal rays less than length of pectorals, and more than ½ length of head; anal spines present. Dark rusty brown, spotted or nearly plain; young olive brown, more or less spotted and blotched with reddish. Atlantic coast of North America, from Labrador to Chesapeake Bay. This small flounder is one of the most abundant of the group on our Atlantic coast. It reaches a length of about 15 inches and a weight of less than 2 pounds. It is a very good food-fish and sells readily in the markets. Along the south coast of Massachusetts this species is more abundant than any other of the flatfishes. The specimens examined by us are from Labrador, Cape Breton, Anticosti, Grand Menan, Boston, Provincetown, Woods Hole, New Bedford, and Somers Point, New Jersey.

Flounder, Schöpf, Schrift. Gesellschaft Naturforscher Freunde, VIII, 1788, 148, New York.
Pleuronectes americanus, Walbaum, Artedi Piscium, 113, 1792, based on the *Flounder* of Schöpf; Bloch & Schneider, Syst. Ichth., 150, 1801; Günther, Cat., IV, 443, 1862; Jordan & Gilbert, Synopsis, 837, 1883; Stearns, Proc. U. S. Nat. Mus. 1883, 125.
Pleuronectes planus, Mitchill, Trans. Lit. & Philos. Soc. N. Y., I, 1815, 387, New York.
Platessa pusilla, De Kay, New York Fauna: Fishes, 296, pl. 47, fig. 153, 1842, New York; Storer, Synopsis, 477, 1846.
Platessa plana, Storer, Rept. Fishes Mass., 140, 1839; De Kay, New York Fauna: Fishes, 295, pl. 49, fig. 158, 1842; Storer, Synopsis, 476, 1846.
Pseudopleuronectes planus, Bleeker, Comptes Rendus Amsterd., XIII, 1862, 7.
Pseudopleuronectes americanus, Gill, Proc. Ac. Nat. Sci. Phila. 1864, 216; Goode, Nat. Hist. Aquat. Anim., 182, pl. 44, 1884; Jordan & Goss, Review Flounders and Soles, 289, 1889.

3018. PSEUDOPLÆURONECTES PINNIFASCIATUS (Kner).

Head 3¼ in body; depth 2⅛. D. 58; A. 38; scales 70; eye 5⅜ in head; snout 5; highest anal ray 2; pectoral 2; caudal 4½ in body. Body subelliptical, the snout rather pointed and not forming an angle above eye; mouth rather small, maxillary reaching scarcely to the middle of the lower eye; interorbital space rather broad, ½ width of eye; a rather prominent

rugose ridge above opercle, with a smaller similar ridge behind it; both sides of jaws with teeth, those on blind side stronger; origin of dorsal over middle of upper eye. Color brown, with vague dusky spots; 6 or 7 blackish vertical bars on dorsal and anal; similar lengthwise blotches on caudal. Okhotsk Sea, east to Kamchatka. (Steindachner.) Not seen by us. From the excellent figure we conclude that it belongs to *Pseudopleuronectes*, although its pharyngeals have not been described. It seems to us nearer to *P. americanus* than to *Liopsetta glacialis*. (*pinna*, fin; *fasciatus*, banded.)

Pleuronectes pinnifasciatus, KNER, in STEINDACHNER, Ueber einige Pleuronectiden, etc., aus Decastris Bay, 2, pl. 1, fig. 1, 1870, Decastris Bay, mouth of Amur River; JORDAN & GOSS, Review Flounders and Soles, 290, 1889.

1037. PLEURONECTES (Artedi) Linnæus.

(PLAICE.)

Pleuronectes, ARTEDI, Genera, etc., in part, 16, 1738.

Pleuronectes, LINNÆUS, Syst. Nat., Ed. X, 268, 1758 (*platessa*); included all known *Pleuronectidæ*.

Platessa, CUVIER, Règne Animal, Ed. 1, II, 220, 1817 (*platessa*).

Pleuronectes, SWAINSON, Nat. Hist. Class'n Anim., II, 302, 1839 (*platessa*).

Pleuronectes, BLEEKER, Comptes Rendus Acad. Amsterd., XIII, 1862 (*platessa*); and of most recent authors.

Body oblong, with firm flesh. Mouth small, teeth uniserial, incisor-like, compressed, forming a continuous cutting edge. Lateral line straightish, without arch or accessory dorsal branch. Scales imperfectly imbricated, chiefly cycloid in both sexes; lower pharyngeals small and narrow, separate, each with 1 or 2 rows of small bluntish teeth. No stellate scales along bases of dorsal and anal. Species mostly European; valued as food. (πλεῦρον, side; νέκτης, swimmer.)

3019. PLEURONECTES QUADRITUBERCULATUS, Pallas.

Head 3⅔; depth 2. D. 68; A. 50; scales 78. Mouth very small, with small, incisor-like teeth, rounded at tip. Eyes separated by a narrow ridge; about 5 small, prominent, conical, obtuse, bony tubercles in a row above the opercle, continuous with the direction of the lateral line, which is straight, without accessory dorsal branch; tubercle above opercle largest. Scales small, cycloid in all specimens examined. Anal spine present. Grayish, mottled with paler and with round black spots; fins very dark. Bering Sea on both coasts, south to Kadiak; not common. Our specimens from Avatcha Bay, Bristol Bay, Herendeen Bay, Chernofsky Harbor, Grantley Harbor, Chignik Bay, and Robben Island. The above description from a small specimen (No. 28025, U. S. Nat. Mus.) collected by Mr. W. J. Fisher at Kadiak. The species proves, as suspected by Jordan & Goss, to be a true *Pleuronectes*, having the lower pharyngeals narrow, separate, with 2 rows of bluntish teeth. (*quadrituberculatus*, having four tubercles.)

Pleuronectes quadrituberculatus, PALLAS, Zoogr. Rosso-Asiat., III, 423, 1811, sea between Kamchatka and Alaska; BEAN, Proc. U. S. Nat. Mus. 1881, 241; JORDAN & GILBERT, Synopsis, 836, 1883.

Pleuronectus pallasii, STEINDACHNER, Ichth. Beitr., VIII, 45, 1879, Kamchatka.

Parophrys quadrituberculatus, GÜNTHER, Cat., IV, 456, 1862.

Platessa quadrituberculata, JORDAN & GOSS, Review Flounders and Soles, 292, 1889.

1038. LIOPSETTA, Gill.

(Eel-back Flounders.)

Liopsetta, Gill, Proc. Ac. Nat. Sci. Phila. 1864, 217 (*glaber*); females.
Euchalarodus, Gill, Proc. Ac. Nat. Sci. Phila. 1864, 222 (*putnami*); males.

Teeth chiefly uniserial, incisor-like; scales imperfectly imbricated, rough ctenoid in the male, more or less cycloid in the female (fin rays scaly in the male, naked in the female); lower pharyngeals very large, more or less united in the adult, their surface somewhat concave, with teeth in 5 or 6 rows, large, blunt, close set; lateral line without arch or dorsal branch. This genus comprises several species of small flounders of the Arctic seas. The genus is distinguished by the large, half-united pharyngeals, as also by the peculiar squamation, the scales in the males being very rough, in the females smooth. This difference has given rise to the nominal genus *Euchalarodus*, based on the males, while *Liopsetta* was based on the smoother females, which were erroneously supposed to be scaleless. (λεῖος, smooth; ψῆττα, flounder.)

a. Dorsal rays 55 or 56; anal 40 to 42.
b. Pectoral fin short, ⅓ length of head in males, shorter in females. GLACIALIS, 3020.
bb. Pectoral fin long, 1⅓ in head in males, nearly 2 in females. PUTNAMI, 3021.
aa. Dorsal rays 59 to 62; anal 45 or 46; pectoral 1⅔ in head in males. OBSCURA, 3022.

3020. LIOPSETTA GLACIALIS (Pallas).

(Arctic Flounder.)

Head 4; depth 2½. D. 56; A. 42. Form of *Liopsetta putnami*. A roughened ridge above the cheeks and opercles on the eyed side. Eyes separated by a narrow, smooth, bony ridge. Scales minute, embedded, nonimbricate, ctenoid in the males, smooth in the females; scales on blind side similar, less developed; scales of lateral line a little larger. Teeth colored, incisor-like, forming an even edge, mostly on blind side. An anal spine; pectorals short. Dark brown, the fins spotted. Arctic shores of Alaska and Siberia, south in Bering Sea to Petropaulski, St. Michaels, and Bristol Bay. Our specimens from Port Clarence, Petropaulski, Bristol Bay, mouth of Nushagak River, and Kotzebue Sound; the description from specimens from the last-named locality taken by Dr. Bean. It is said to be abundant in the Arctic Ocean and as far south as Bristol Bay. "Although small, its great abundance and fine flavor make it important as an article of food." The male is the rough fish described by Pallas as *P. cicatricosus*. The smoother female is Dr. Günther's *Pleuronectes franklinii*, the sexual differences being much as in *Liopsetta putnami*. *Liopsetta dvinensis* of the northern coasts of Russia may be the same species. (*glacialis*, icy.)

Pleuronectes glacialis, Pallas, Itin., III, App., 706, mouth of River Obi; Bloch & Schneider, Syst. Ichth., 150, 1801; Pallas, Zoogr. Rosso-Asiat., III, 424, 1811; Richardson, Fauna Bor.-Amer., Fish., 258, 1836; De Kay, N. Y. Fauna: Fishes, 302, 1842; Storer, Synopsis Fish. N. A., 479, 1846; Bean, Proc. U. S. Nat. Mus. 1881, 241; Jordan & Gilbert, Synopsis, 837, 1883; Bean, Cat. Coll. Fish. U. S. Nat. Mus., 20, 1883; Bean, Nat. Hist. Aquat. Anim., 184, pl. 47, 1884.

Pleuronectes cicatricosus, Pallas, Zoogr. Rosso-Asiat., III, 424, 1811, male, sea between Kamchatka and Alaska.

? *Platassa dvinensis*, LILLJEBORG, Veb. Ak. Handl. 1850, 360, tab. 20, mouth of River Dwina.
Pleuronectes franklinii, GÜNTHER, Cat. Fish., IV, 442. 1862, Arctic seas of America, female; BEAN, Proc. U. S. Nat. Mus. 1881, 241.
Liopsetta glacialis, JORDAN & GOSS, Review Flounders and Soles, 295, pl. 17, 1889.

3021. LIOPSETTA PUTNAMI (Gill).

(EEL-BACK FLOUNDER.)

Head 3½; depth 2. D. 55; A. 40; scales 70 (pores). Body oblong, ovate. Eyes rather small, separated by a naked elevated ridge. Jaws sometimes each with 2 distinct rows of teeth, the interrupted outer series of truncate, close set, thickish, incisor-like teeth, which are sometimes movable; the inner row of similar teeth more widely set and rather distant from the outer row (and often or generally wanting); about 20 teeth in outer row in lower jaw; right side of each jaw toothless; interorbital ridge continuous, with a broad, naked, smoothish, tuberculose ridge, which joins the lateral line. Scales small, distant, nonimbricate, smooth in the female, and more or less ctenoid in the male, those on blind side smaller. Fins moderate, somewhat scaly; anterior rays of dorsal low; pectoral a little more than ½ head; bases of vertical fins not tuberculate; anal spine present; lower pharyngeals separate, broad, with coarse teeth. Grayish brown, mottled with darker brown; fins with blackish spots. Length 10 inches. Atlantic coast of North America, from Cape Cod northward to Labrador and beyond; occasionally found in abundance. This species is rather common along the coast of northern Massachusetts and northward to Labrador. Specimens are frequently found in the markets, mixed with those of *Pseudopleuronectes americanus*. The numerous specimens in our possession were found in the markets of Indianapolis, having been sent thither from Boston. The remarkable sexual differences in the species have been fully discussed by Dr. Bean (Proc. U. S. Nat. Mus. 1878, 345), the form formerly called *Euchalarodus putnami* being the male, and that called *Pleuronectes glaber* being the female of the same species. These conclusions of Dr. Bean are fully corroborated by our series of specimens in which both sexes are fully represented.

Although *Liopsetta putnami* is abundant where found, its ascertained range is somewhat limited. The specimens in the United States National Museum represent localities from Salem, Massachusetts, to Belfast, Maine. In the Museum of Comparative Zoology the localities represented are Providence, Boston, Salem, Grand Manan, and Labrador. (Named for Prof. Frederic Ward Putnam.)

Platessa glabra, STORER, Proc. Boston Soc. Nat. Hist. 1843, 130, female, **Massachusetts**; STORER, Syn. Fish. N. A., 477, 1846; STORER, Hist. Fish. Mass., 199, pl. 31, fig. 1, 1867; PUTNAM, Bull. Essex Inst., VI, 1874, 12; not *Platessa glabra* of RATHKE, 1837, a species of *Flesus*.

Euchalarodus putnami, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 216–221, **Salem, Massachusetts** (Coll. F. W. Putnam), male; GILL, Report U. S. Fish Comm. 1873, 794; GOODE & BEAN, Amer. Journ. Sci. and Arts, XIV, 1877.

Liopsetta glabra, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 217.

Pleuronectes glaber, GILL, Report U. S. Fish Comm. 1873, 794; GOODE & BEAN, Amer. Journ. Sci. and Arts, XIV, 1877, 476; XVII, 1879, 40; GOODE & BEAN, Proc. U. S. Nat. Mus. 1878, 347; JORDAN & GILBERT, Synopsis, 836, 1883; GOODE, Nat. Hist. Aquat. Anim., 183, pl. 45, 1884.

Liopsetta putnami, JORDAN & GOSS, Review Flounders and Soles, 294, pl. 16, 1889.

3022. LIOPSETTA OBSCURA (Herzenstein).

To this species we refer 2 males from Shana Bay, Iturup Island. The scales on the colored side are everywhere strongly ctenoid and imbricated, while in Herzenstein's types (supposed to be females) they were cycloid. In our specimens the head is somewhat smaller, $3\frac{3}{10}$ in length instead of $3\frac{1}{10}$ to $3\frac{5}{9}$; the depth is greater, $2\frac{1}{5}$ in length instead of $2\frac{3}{7}$ to $2\frac{3}{8}$; the interorbital space is covered with very fine scales, not naked; the curve of the lateral line seems more marked, its cord contained 5 instead of 6 times in the straight portion. All of the fins are higher than in the female types, the pectoral of colored side being $1\frac{3}{5}$ head, the caudal $1\frac{1}{6}$, the ventral $\frac{1}{2}$ head, and the highest dorsal ray $1\frac{2}{3}$. Some of these differences may well be sexual. The lower pharyngeals are short and broad, the two closely appressed but united in our specimens, 27 and 29 cm. long. The teeth are large and very blunt, like cobble stones, and are arranged in 1 row along the outer edge; a row of larger teeth along the inner edge, and a short row along the posterior edge of the triangle. The arrangement is very similar is that found in *L. glacialis*, but here a few small teeth, without definite arrangement, are interposed in the middle of the bone, between the third series described. Dorsal 59 and 62; anal 45 and 46; tubes in the lateral line 79. Color on eyed side uniform dark brown on body and fins, the extreme tips of the fin rays white; on blind side yellowish white, with a few irregular scattered dark spots; dorsal and anal yellowish at base, becoming more or less mottled with dusky on distal half, the fins marked with broad dark bars parallel with the rays, about 7 on the anal fin, 10 or 11 on the dorsal; caudal light on basal half more or less blotched with darker, becoming black posteriorly. With this species we identify also a number of young individuals, 9 to 15 cm. long, from the same locality (Iturup Island). They are probably young females, but the viscera are in such condition as to prevent positive determination. The scales are perfectly smooth, but in other respects they agree perfectly with the adult males, except in their more varied coloration; head and body brownish, profusely spotted in coarser or finer pattern with light gray; also with a few scattered black spots edged with gray; markings on the fins as described for adults. In 7 specimens the dorsal contains 60, 62, 62, 62, 64, 65, and 66 rays; anal 45, 45, 45, 46, 47, 47, 48. Sea of Okhotsk. Our specimens from Shana Bay, Iturup Island, one of the Kurils; originally described from Mantchuria. (*obscurus*, dark.)

Pleuronectes obscurus, Herzenstein, Mélanges Biologiques, 127, 1890, Mantchuria.
Liopsetta obscura, Jordan & Gilbert, Rep. Fur Seal Invest., 1898.

1039. PLATICHTHYS, Girard.

(Starry Flounders.)

Platichthys, Girard, Proc. Ac. Nat. Sci. Phila. 1854, 136 (*rugosus*=*stellatus*).

Body very robust, broad, not greatly compressed. Mouth small; teeth chiefly uniserial, incisor-like. Scales all in both sexes and on both sides of body reduced to coarse scattered stellate tubercles, which are not

imbricated; similar tubercles between bases of dorsal and anal rays; lateral line without scales, with no anterior arch or accessory lateral line; lower pharyngeals broad, each with 3 rows of blunt coarse teeth. A single species, the largest of the small-mouthed flounders, and distinguished from related forms chiefly by the development of coarse stellate tubercles instead of scales. (*πλατύς*, flat; *ἰχθύς*, fish.)

3023. PLATICHTHYS STELLATUS (Pallas).

(Great Flounder.)

Head $3\frac{3}{5}$; depth 2. D. 58; A. 42. Vertebræ 34. Body broad and short, the snout forming a slight angle with the profile; lower jaw projecting; interocular space rather broad, with very rough scales; large rough scales at base of dorsal and anal rays and on sides of head; similar but smaller scales scattered over the body; lateral line smooth; fins without scales; a cluster of bony prominences above opercle. Teeth incisor-like, truncate, rather broad, $\frac{10+15}{12+16}$. Lower pharyngeals broad, with coarse paved teeth. Dark brown or nearly black, with lighter markings; fins reddish brown; dorsal and anal with 4 or 5 vertical black bands; caudal with 3 or 4 black longitudinal bands. Pacific coast of America, from Point Concepcion to the Arctic Ocean and south to the Amur River. This is one of the largest of the American flounders, reaching a weight of 15 to 20 pounds. Of the small-mouthed flounders it is much the largest species known. It is an excellent food-fish, and from its size and abundance it is one of the most important of the group in the region where it is found, constituting half the total catch of flounders on our Pacific coast, and it is equally abundant in Bering Sea. It lives in shallow water and sometimes ascends the larger rivers. It is one of the most widely distributed of all the flounders, its range extending from San Luis Obispo, where it was obtained by Jordan & Gilbert, to the mouth of the Anderson and Colville rivers on the Arctic coast, where it was observed by Dr. Bean, and to Port Clarence, where Mr. Scofield obtained specimens. We have also specimens from Petropaulski, Bering, Medni, and Robben islands and from Bristol Bay. A specimen from the island of Saghalien in Asia is in the museum at Cambridge. (*stellatus*, starry.)

Pleuronectes stellatus, Pallas, Zoographia Rosso-Asiatica, III, 416, 1811, Kamchatka, Aleutian and Kuril Islands; Günther, Cat., IV, 443, 1862; Steindachner, Pleur. von Decastris Bay, 1870, 1; Jordan & Gilbert, Proc. U. S. Nat. Mus. 1880, 453; Jordan & Gilbert, Proc. U. S. Nat. Mus. 1881, 68; Bean, Proc. U. S. Nat. Mus. 1881, 420; Jordan & Gilbert, Synopsis 835, 1883; Bean, Proc. U. S. Nat. Mus. 1883, 353; Bean, Cat. Coll. Fish. U. S. Nat. Mus. 1883, 20; Jordan, Nat. Hist. Aquat. Anim., 184, pl. 46, 1884.

Platichthys rugosus, Girard, Proc. Ac. Nat. Sci. Phila. 1854, 139, 155, San Francisco; Presidio; Petaluma; Girard, U. S. Pac. R. R. Surv., X, Fishes, 148, 1858.

Platessa stellata, De Kay, N. Y. Fauna: Fishes, 301, 1842; Storer, Synopsis, 478, 1846.

Platichthys stellatus, Lockington, Rep. Com. Fish. Cal. 1878–79, 43; Lockington, Proc. U. S. Nat. Mus. 1879, 91; Jordan & Goss, Review Flounders and Soles, 296, 1889.

1040. MICROSTOMUS,* Gottsche.

(Smear Dabs.)

Microstomus, Gottsche, Archiv fur Naturgesch. 1835, 150 (*latidens*); not *Microstoma*, Risso, 1826.

Cynicoglossus, Bonaparte, Fauna Italica, 1837, fasc., xix (*cynoglossus*, Nilsson, not of L.).

Cynoglossa, Bonaparte, Catalogo Metodico Pesci Europei, 48, 1846 (*microcephalus*); not *Cynoglossus*, Hamilton, 1822.

Brachyprosopon, Bleeker, Comptes Rendus Acad. Sci. Amsterd., xiii, Pleuron., 7, 1862 (*microcephalus*).

Cynicoglossus, Jordan & Gilbert, Synopsis, 460, 1883 (*microcephalus*).

Body elongate, compressed; mouth very small; teeth broad, incisor-like, on blind side only; scales small, all cycloid; vertebræ numerous (48 to 52); dorsal rays 90 to 100; anal rays 70 to 85; anal spine obsolete; left side of skull normal, without mucous cavities; ventral fins with 5 rays each. Arctic seas. This genus is widely separated from *Pleuronectes*

* We here retain the generic name *Microstomus*, although in accordance with recent usage of most ornithologists and ichthyologists, it should be suppressed, as identical with *Microstoma*. The two words are from the same root, and differ only in the termination. But is not this difference enough? The code of nomenclature of the American Ornithologists' Union very properly declares that "a name is only a name, and has no necessary meaning," and therefore no necessarily correct spelling, except the spelling selected by the writer from whom it dates its origin. As a result of this, the original spelling of each generic name is (undoubted misprints aside) the orthography to be adopted, regardless of all questions as to the correct etymology of the word. As a necessary sequence, it seems to us that all generic names, not actually preoccupied by names spelled in the same way, should be tenable. There is no other certain boundary line between names tenable and names untenable. We therefore regard all generic names as available unless used in zoology earlier and in exactly the same orthography. Among American genera of fishes we may therefore use the following, notwithstanding their earlier analogies:

Microstomus for	*Cynicoglossus* notwithstanding the prior	*Microstoma.*
Heterodontus	*Cestracion*	*Heterodon.*
Lucania	———	*Lucanus.*
Thymallus	*Choregon*	*Thymalus.*
Nebris	———	*Nebria.*
Xiphidion	*Xiphister*	*Xiphidium.*
Amitra	*Monomitra*	*Amitrus.*
Scytalina	*Scytaliscus*	*Scytalinus.*
Lagochila	*Quassilabia*	*Lagocheilus.*
Auchenopterus	*Cremnobates*	*Auchenipterus.*
Lyopsetta	———	*Liopsetta.*
Leucos	*Myloleucus*	*Leucus.*
Pterophryne	*Pterophrynoides*	*Pterophrynus.*
Scaphirhynchus	*Scaphirhynchops*	*Scaphorhynchus.*
Lepidion	*Haloporphyrus*	*Lipidia.*
Gramma	———	*Grammia.*
Stenotomus	———	*Stenotoma.*

If *Microstomus* be discarded, the next name in order of date is *Cynicoglossus*.

The following is Bonaparte's definition of *Cynicoglossus* as quoted by Gill (Proc. Ac. Nat. Sci. Phila. 1864, 222):

"Secundo è *Cynicoglossus* nob. che come il *Pl. cynoglossus* L. ha la linea laterale retta, la bocca piccola, i denti come quello di sopra [*Platessa*] ma la mascelle iguale, con labbra turgide, e l' ano senza spina."

Later, in his Catalogo Metodico dei Pesci Europei, Bonaparte changes this name from *Cynicoglossus* to *Cynoglossa*, giving the sole species as *Cynoglossa microcephala*, and quoting as its synonym "*Pleuronectes cynoglossus*, N. Nilss.", showing that his identification of the Linnæan species coincided with that of Nilsson, who at first used the name "*Pleuronectes cynoglossus*" for the present species instead of the species of *Glyptocephalus*. In Bonaparte's Catalogo, *Glyptocephalus*, Gottsche, is regarded by Bonaparte as synonymous with *Platessa*.

It is thus evident, as Dr. Gill has suggested, that Bonaparte meant to refer to the *Pleuronectes microcephalus* instead of *Pl. cynoglossus*, he "having followed Nilsson in his erroneous identification" of the latter with the former. In further evidence of this we have the fact that *Cynicoglossus microcephalus* (*kitt*) has no anal spine, while such a spine is present in the species of *Glyptocephalus*. We would be, therefore, justified in the use of *Cynicoglossus* instead of the later *Brachyprosopon*, if *Microstomus* should be regarded as ineligible on account of the prior name *Microstoma*. (Jordan & Goss.)

and its allies by its greatly increased number of vertebræ, a character accompanied by a similar increase in the number of fin rays. It is close to *Glyptocephalus*, but the lack of the cavernous structure of the bones of the head, a structure peculiar to the species of that genus, sufficiently distinguishes it. (*μικρός*, small; *στόμα*, mouth.)

a. Dorsal rays 85 to 93; anal rays 70 to 76; head very small, about 5 in length; eye 4 in head. KITT, 3024.

aa. Dorsal rays 102; anal ray 85; head 4½ in length; eye 3 in head. PACIFICUS, 3025.

3024. MICROSTOMUS KITT (Walbaum).

(SMEAR DAB.)

Head 5¼ in length; depth 2¼. D. 85 to 93; A. 70 to 76; scales 130; caudal 1⅓ in head; pectoral 1¾. Body moderately elongate; mouth small, the maxillary not reaching to front of lower eye; teeth on blind side conical, rather compressed and blunted, 11 to 13 on either jaw; eyes close together, the lower slightly in advance; gill rakers short, not numerous. Origin of dorsal above middle of upper eye, its rays larger in the posterior half of body; pectorals about equal in size; no spine before anal; caudal rounded; head, except snout, entirely scaled; scales cycloid; lateral line with a small curve; vertebræ 13 + 35 = 48. Color dull yellowish, blotched, and with dark spots, especially over the chest and along the base of anal fin; dark blotches and spots on anal, caudal, and ventral fins; dark base to pectoral, which has also some cloudy markings. (Day.) Seas of the north of Europe in rather deep water, south to Cornwall. Recorded by Steindachner (as *Pleuronectes gilli*), from the sea between Iceland and Greenland. This small flounder is rather common in the waters of northern Europe. It reaches the length of a foot or more, and is said to be excellent as food. Like its congener, *Microstomus pacificus*, this species is very slimy in life. *Pleuronectes gilli*, as described by Dr. Steindachner, seems to differ from *Microstomus kitt* only in the larger head, which is but 4⅔ in the length to base of caudal. It is probably not specifically distinct from the latter. Only a single specimen, 10½ inches long, is known. (Eu.) (The specific name "*kitt*," given by Walbaum on the authority of Jago's description, should be adopted for this species. According to Day, the species is still called "*kitt*" on the coast of Cornwall.)

Rhombus lævis cornubiensis, JAGO, in Ray, "Syn. Pisc., 162, tab. 1, fig. 1, 1713."

The Smear Dab, PENNANT, British Zoology, III, 230, pl. 41, 1776.

Pleuronectes kitt, WALBAUM, Artedi Piscium, III, 120, 1792, after RAY; the description in part confused with that of *Lepidorhombus*.

Pleuronectes lævis, SHAW. Gen'l Zool., IV, 299, 1803.

Pleuronectes quenseli, HÖLBÖLL, Bohusläns Fiske, IV, 59, 1821, Bohusläns, Sweden.

Pleuronectes quadridens, FABRICIUS, Kongl. Dansk. Vid. Selsk. Afhandl., I, 39, 1824, Iceland.

Microstomus latidens, GOTTSCHE, Archiv fur Naturgsch. 1835, 150, Zealand.

Pleuronectes gilli, STEINDACHNER, Ichth. Notizen, VII, 40, 1868, Polar Sea north of Iceland.

Pleuronectes microcephalus, DONOVAN, British Fishes, II, pl. 42, 1802; GÜNTHER, Cat., IV, 447; STEINDACHNER, Ichth. Beitr., VIII, 47; DAY, Fishes Great Britain, II, 28, pl. 102; COLLETT, Norges Fiske, 145, and of recent European writers generally.

Pleuronectes microstomus, FABER, Isis, 886, 1828.

Platessa microcephala, FLEMING, British Anim., 198, 1828, and of numerous writers.

Cynoglossa microcephala, BONAPARTE, Catalogo Metodico Pesci Eur., 48, 1845.

Microstomus kitt, JORDAN & GOSS, Review of Flounders and Soles, 1886, 298.

3025. MICROSTOMUS PACIFICUS (Lockington).

(SLIPPERY SOLE.)

Head $4\frac{1}{4}$ in body; depth $2\frac{5}{6}$. D. 102; A. 85; scales 140; eye $3\frac{1}{2}$ in head; maxillary 5; pectoral $1\frac{3}{5}$; greatest height of dorsal $2\frac{1}{3}$; anal $2\frac{1}{3}$; caudal $1\frac{2}{5}$; vertebræ $12+40=52$. Body elongate, elliptical; mouth small, the maxillary reaching just past front of lower eye; teeth long and broad, forming a continuous cutting edge, on blind side only, about 10 teeth on lower jaw; eyes very large, nearly twice as long as snout, the upper even with profile above; interorbital a narrow scaly ridge; gill opening adnate to shoulder girdle above pectoral; gill rakers short, 8 below angle, 5 or 6 very small scarcely developed ones above; scales small, cycloid, not closely imbricated, lateral line nearly straight. Origin of dorsal slightly behind middle of upper eye, caudal truncate or slightly rounded. Color olive brown, blotched on body and fins with darker, all fins blackish toward the ends of the rays. Pacific coast of North America, Monterey to Unalaska, in rather deep water, 15 to 50 fathoms; common. Here described from a specimen, about 14 inches in length, from *Albatross* Station 2927, off the coast of California. This small flounder abounds in deep water about San Francisco, but comes near the shore farther north. It is exceedingly slimy when first taken. The large individuals are considered excellent as food; the smaller are thrown away. It rarely reaches the weight of a pound.

Glyptocephalus pacificus, LOCKINGTON, Rep. Cal. Com. Fisheries, 1878–79, 43, off Point Reyes, California; LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 86; JORDAN, Nat. Hist. Aquat. Anim., 188, 1884.

Cynicoglossus pacificus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 453; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 68; JORDAN & GILBERT, Synopsis, 838, 1883.

Microstomus pacificus, JORDAN & GOSS, Review Flounders and Soles, 299, 1889.

1041. EMBASSICHTHYS, Jordan & Evermann.

Embassichthys, JORDAN & EVERMANN, Check-List Fishes, 506, 1896 (*bathybius*).

This genus is a deep sea representative of *Microstomus*, from which it differs in the increased number of vertebræ (63 instead of 48 to 52). Its fin rays are correspondingly increased, the body is deeper than in *Microstomus*, and it has teeth on both sides of the jaws, as in *Glyptocephalus*. (ἐν, in; βάσσος, for βαθύς deep; ἰχθύς, fish; a fish in the depths.)

3026. EMBASSICHTHYS BATHYBIUS (Gilbert).

Head 4 to $4\frac{2}{3}$ in length; depth 2 to $2\frac{1}{3}$. D. 111 to 117; A. 96 to 98; vertebræ $14+49=63$. Body oval, very thin and deep, the greatest depth at anterior third of body; upper profile very abruptly angulated opposite hinder margin of upper pupil, the anterior half of head conspicuously protruding beyond general outline. Caudal nearly sessile, the peduncle very short. Mouth small, maxillary about $\frac{1}{3}$ length of head in specimens 1 foot long. Teeth broad incisors, slightly notched at tip, nearly equally developed on blind and colored sides, 21 on blind side of lower jaw, 16 on

colored side. As in other members of this group, the lower jaw is the longer, the upper teeth included. Interorbital space wholly scaled, with a very high, rather sharp ∼-shaped ridge. Eyes very large, the upper entering largely into the upper profile, the lower much in advance; front margin of upper orbit on vertical of front of lower pupil; diameter of upper eye $2\frac{3}{4}$ to $2\frac{4}{5}$ in head. Anterior nostrils of both sides in rather long tubes, the posterior margins produced to form short flaps. Preopercular margins adnate, as usual, concealed by scales. No conspicuous mucous excavations on blind side. Gill rakers weak and rather short, 10 or 11 on anterior of arch. Scales very small, cycloid, in about 165 cross rows, the tubes of lateral line much fewer, not regularly arranged; over 50 longitudinal rows above lateral line. Dorsal beginning over posterior edge of pupil; fins low, the highest dorsal rays behind middle of body, $\frac{2}{5}$ length of head; caudal rounded, $1\frac{1}{2}$ in head; pectorals 2 in head; ventrals small, each with 5 rays, as in *Microstomus pacificus*. (*Glyptocephalus cynoglossus* and *zachirus* have 6 rays in each ventral.) Color of eyed side warm brown, darker toward margins, becoming black on vertical fins; everywhere on body and fins coarsely blotched with light blue, the marks so arranged on upper and lower thirds of sides as to form 5 broad bars of bluish, alternating with those of the ground color, and corresponding above and below; lips and branchiostegal membranes black; blind side dusky brownish. This well-marked species differs from the species of *Microstomus* in its much greater depth and bright coloration, and in having teeth well developed on both sides of jaws, as in the species of *Glyptocephalus*. Two specimens from the Santa Barbara Channel, in deep water. (Gilbert.) (*βαθύς*, deep; *βίος*, life.)

Cynicoglossus bathybius, Gilbert, Proc. U. S. Nat. Mus. 1890, 123, Santa Barbara Channel, at Albatross Station 2980, Lat. 33° 49′ 45″ N., Long. 119° 24′ 30″ W., in 603 fathoms. (Type in U. S. N. M. Coll. Gilbert.)

1042. GLYPTOCEPHALUS, Gottsche.

(Flukes.)

Glyptocephalus, Gottsche, Archiv fur Naturgsch. 1835, 156 (type *saxicola* = *cynoglossus*, L.).

Eyes and color on the right side. Body extremely elongate, more than twice as long as deep, much compressed. Head very small and short, its blind side with many excavations and mucous cavities in the skull, mandible, and preopercle. Mouth very small; teeth moderate, incisor-like, broad, equal, close set, in a single series; no teeth on vomer or palatines. Gill rakers short, weak. Lower pharyngeals narrow, with 1 or 2 rows of conical teeth. Lateral line nearly straight, simple; scales very small, smooth; dorsal and anal very long, there being more than 90 rays in the dorsal and more than 80 in the anal; caudal fin rounded; anal spine present; ventral rays 6. Vertebræ in increased number, 58 to 65. Northern seas, in deep water. This genus is one of the most strongly marked in the family, being distinguished from most of the genera by the greatly increased number of vertebræ, and from all of them by the remarkable cavernous

structure of the bones of the head. Two species known. (*γλυπτός*, sculptured; *κεφαλή*, head.)

a. Pectoral fins very short, not falcate, that of right side about ½ length of head; vertebræ 58. CYNOGLOSSUS, 3027.

aa. Pectoral fins of colored side falcate, longer than the head; vertebræ 65. ZACHIRUS, 3028.

3027. GLYPTOCEPHALUS CYNOGLOSSUS (Linnæus).

(CRAIG FLUKE; POLE FLOUNDER.)

Head 5 to 5¼ in body; depth 2⅓ to 3. D. 101 to 112; A. 87 to 100; scales 125; V. 6; highest dorsal and anal rays 2 in head; pectoral a little more than 2; vertebræ 58. Body oblong, fusiform; head small, ovate; the profile slightly decurved; mouth very small, with the cleft oblique; teeth on blind side close set, with incisoral edges, $\frac{1}{2}\frac{7}{5}$; on the eyed side, distant, obtusely conic, $\frac{6}{7}$; eyes moderate, the lower advanced, close together, 3 in head; scales regularly imbricated, lateral line straight; pectoral short, falcate; origin of dorsal above middle of upper eye; anal spine present; caudal convex or angulated behind; pectoral fins very short, not falcate, that of right side about ½ length of head; upper jaw with about 30 teeth; opercle adnate to the shoulder girdle for a short distance only. Color grayish brown; fins with dark spots; tip of pectoral dusky above. North Atlantic, on both coasts, chiefly in deep water, south to Cape Cod and France. This species is found in rather deep water on sandy bottoms. It reaches a length of 12 to 18 inches. This flounder has been taken in great numbers with the beam trawl in deep water off our New England coast. It is pronounced by the United States Fish Commission to be not inferior as a food-fish to the European sole. (Eu.) (*cynoglossus*, a sole; *κύων*, dog; *γλῶσσα*, tongue.)

Pleuronectes, etc., *Corpore oblongo glabro*, GRONOW, Museum Ichthyol., I, IV, 39, etc., Belgium.

Pleuronectes cynoglossus, LINNÆUS, Syst. Nat., Ed. X, 269, 1758, after GRONOW; GÜNTHER, Cat., IV, 449, 1862; DAY, Fishes Great Britain, II, 30, pl. 103; LILLJEBORG, Sveriges och Norges Fiske, II, 386, 1891; and of European writers generally.

Platessa pola, CUVIER, Règne Animal, Ed. II, Vol. 2, 339, 1829, after la Pole of Duhamel.

Pleuronectes saxicola, FABER, Tidsskr. f. Naturv., 5 B., 244, 1828, Denmark.

Pleuronectes nigromanus, NILSSON, Prodr. Ichth. Scand., 55, 1832.

Platessa elongata, YARRELL, Hist. Brit. Fish., 619, 1859, young.

Glyptocephalus acadianus, GILL, Proc. Ac. Nat. Sci. Phila. 1873, 360, Nova Scotia. (Type, No. 12685.)

Glyptocephalus cynoglossus, GILL, Proc. Ac. Nat. Sci. Phila. 1873, 360; GOODE & BEAN, Proc. U. S. Nat. Mus. 1878, 21; GOODE, Proc. U. S. Nat. Mus. 1880, 337; GOODE, Proc. U. S. Nat. Mus. 1880, 475; COLLETT, Norske Nord-Havs Exped. 1880, 150; GOODE & BEAN, Bull. Mus. Comp. Zool., X, No. 5, 195, 1883; JORDAN & GILBERT, Synopsis, 838, 1883; GOODE, Nat. Hist. Aquat. Anim., 198, 1884; JORDAN & GOSS, Review Flounders and Soles, 300, pl. 19, 1889.

Solea cynoglossa, RAFINESQUE, Indice di Ittiologia Siciliana, 53, 1810; based on the Sole or *Cynoglossum* of RONDELET.

Glyptocephalus saxicola, GOTTSCHE, Archiv fur Naturgesch. 1835, 156.

Platessa saxicola, KRÖYER, Danmark's Fiske, 338, 1843.

Pleuronectes elongatus, GÜNTHER, Cat., IV, 450, 1862.

Gyptocephalus elongatus, GILL, Proc. Ac. Nat. Sci. Phila. 1873, 362.

3028. GLYPTOCEPHALUS ZACHIRUS, Lockington.

(Long-finned Sole.)

Head $5\frac{1}{3}$ to $5\frac{1}{2}$; greatest width of body $3\frac{1}{3}$ to $3\frac{1}{2}$; eye $3\frac{1}{3}$ in head; snout 8. D. 94 to 106; A. 79 to 89; P. 11 to 13; V. 6; vertebræ 13 + 52 = 65. Body elongate-ovate, anterior portion of the oval shorter than posterior; snout declivous, almost vertical, its tip level with upper margin of lower eye, its curve uniting without sensible depression with that of nape; dorsal outline rising with a regular gentle curve from snout to about twenty-second dorsal ray, thence declining very gradually and regularly with but slight curvature to caudal peduncle; abdominal outline almost straight from knob of mandible to ventral; from thence to end of anal curved in same manner as dorsal outline; peduncle of tail expanded toward caudal, its least width about $\frac{1}{4}$ of greatest depth of body; greatest distance from anal to lateral line less than length of head. Eyes large, elliptical, the lower in advance of the upper about $\frac{1}{3}$ length of pupil, and scarcely reaching dorsal profile anteriorly. Interocular space very narrow, about $\frac{1}{8}$ of longitudinal diameter of eye, smooth; not raised above the eye in a fresh fish; a slight ridge rising at its posterior part, forming lower posterior margin of upper eye, and dying out on cheek. Nostrils of right side level with upper margin of lower eye; anterior nostril with a short tube, the posterior with a raised margin, and vertical with the front margin of the lower orbit; posterior nostril of blind side in advance of eye; anterior nostril nearly as on colored side; nostrils small and inconspicuous. Gape of mouth very small on colored side, considerably larger on blind side; on the colored side the cleft is nearer vertical than horizontal; posterior end of maxillary reaching very little behind anterior margin of orbit of lower eye, and the symphysis of intermaxillaries about level with upper edge of orbit; mandible projecting in the closed mouth, short, not passing a vertical from front margin of pupil, with a prominent knob below the symphysis, and a smaller one at its posterior extremity. Teeth on both sides of jaws throughout the full length of the gape, in a single row, broad, but thick, forming a blunt, continuous edge, about 34 in lower jaw and rather fewer in the upper, in an individual $11\frac{3}{16}$ inches long; in an example $14\frac{5}{8}$ inches long there were 14 teeth on the colored side and 26 on the blind side of the mandible, the latter the larger; in the intermaxillaries, 13 on the colored side and 23 on the blind side; each lower pharyngeal with a double row of teeth, the inner larger than the outer; the 4 anterior teeth of outer row conspicuously larger than those following; about 12 teeth in each inner row; upper pharyngeals each with a close-set row of 6 or 7 blunt conical teeth. Branchiostegals 7; gill rakers few, flexible, very short. Dorsal commencing between front of orbit and pupil, considerably behind nostrils, long and low, forming a continuous arch of slightly greater curvature than dorsal outline, the longest rays in central portion, and ending opposite anal at about $\frac{2}{3}$ of width of caudal peduncle from origin of caudal; anal with a horizontal spine, the first ray rather distant from the visible portion of the spine, and nearly length of ventral behind pectoral base, similar to the dorsal; almost all the rays of dorsal and anal directly backward; caudal convex on posterior margin, rather narrow, the rays once bifurcate,

sometimes bifurcate again near the tips; pectoral of colored side exceedingly long and lanceolate, about $\frac{1}{4}$ of total length of fish; first 5 rays simple, the others once bifurcate; fourth ray longest, fifth nearly equal, sixth a little longer than third, thence diminishing rapidly. Usual proportion of the first 4 rays 3–8–10–12; pectoral of blind side lanceolate, rather more than $\frac{1}{3}$ of length of that of colored side, and formed of the same number of rays, first 4 simple, the others once forked; fourth and fifth rays longest; ventrals inserted so that their hinder axil is vertical with, or a little posterior to, anterior axil of pectoral, their tips reaching to first anal ray; 4 posterior rays once bifurcate. Lateral line almost straight, rising very slightly anteriorly, formed of a double row of tubes, about 138 in number, excluding those upon caudal; a row of similar pores commencing at ridge under upper eye, and continuing around lower eye almost to its front margin; scales small, smooth, uniform over the body, and extending over the head to snout, on which they are smaller; intermaxillaries and mandible scaleless; scales on blind side similar; caudal scaly on both sides; no scales on the other fins. Color uniform brownish or cinereous; fins darker; the color formed by minute dark spots on the scales; membrane between the fin rays closely set with dark points; blind side whitish, the ground tint clouded with numerous black points. Deep waters of the Northern Pacific, from San Francisco northward; found throughout Bering Sea in 35 to 350 fathoms. This species is a thin, dry flounder, reaching a length of something over a foot. It is taken in the sweep nets in deep water about San Francisco. It is readily known by its long pectoral fin. (ζα-, an intensive particle; χείρ, hand, from the long pectoral.)

Glyptocephalus zachirus, Lockington, Proc. U. S. Nat. Mus. 1879, 88, San Francisco; Lockington, Rep. Com. Fisheries California 1878–79, 42; Jordan & Gilbert, Proc. U. S. Nat. Mus. 1880, 453; Jordan & Gilbert, Proc. U. S. Nat. Mus. 1881, 68; Jordan & Gilbert, Synopsis, 838, 1883; Jordan, Nat. Hist. Aquat. Anim., 188, 1884; Jordan & Goss, Review Flounders and Soles, 301, 1889; Gilbert, Rept. U. S. Fish. Comm. 1893 (1896), 460.

1043. LOPHOPSETTA, Gill.

(Window Panes.)

Lophopsetta, Gill, Proc. Ac. Nat. Sci. Phila. 1862, 216 (*maculatus*).

Eyes and color on the left side. Body broadly ovate, strongly compressed, pellucid; mouth large, oblique, the maxillary reaching to beyond eye; teeth subequal, in narrow bands, or in single series; a small patch of teeth on the vomer. Scales small, cycloid, imbricate, the skin without bony tubercles. Lateral line strongly arched in front, without accessory branch. Dorsal fin beginning on the snout, its anterior rays exserted; anal fin not preceded by a spine; ventral of left side free from the anal, inserted nearly on the ridge of the abdomen, its base broad, the rays well separated; pectoral and ventral fins moderate. One species. Very close to the European genus *Bothus*, Rafinesque (= *Scophthalmus*, Rafinesque, = *Rhombus*, Cuvier, = *Passer*, Valanciennes), from which it differs in the more numerous gill rakers, pellucid body, and produced dorsal rays, all characters of minor importance. The European Turbot (*Psetta*, Swainson), is

also closely related, but the typical species, *Psetta maxima*, is a large robust fish, scaleless and beset with bony tubercles. (λόφος, crest; ψῆττα, turbot.)

3029. LOPHOPSETTA MACULATA (Mitchill).

(Window Pane.)

Head $3\frac{1}{2}$; depth $1\frac{3}{5}$. D. 65; A, 52; scales 85; eye 4 in head; pectoral $1\frac{1}{3}$; highest dorsal rays $1\frac{7}{8}$; highest anal rays $1\frac{1}{5}$; interorbital space $\frac{1}{2}$ eye. Body broadly rhomboid, strongly compressed, translucent in life; mouth large, the maxillary reaching nearly to posterior margin of eye, maxillary of eyed side with a bony tubercle on its anterior end; jaws subequal, the lower with a sharp knob at symphysis; teeth in each jaw in 1 series laterally, in a very narrow band in front; interorbital space rather broad, slightly concave, its posterior third or fourth with scales; gill rakers short and slender, about 8 + 25; maxillary, mandibles, snout, and the greater part of interorbital naked; scales on head and body cycloid, loosely imbricated, those on the blind side a little smaller. Anterior rays of dorsal produced, their ends branched and free, the first on tip of snout, the rays at the beginning of posterior third of fin the highest; origin of anal directly under angle of preopercle; base of ventrals long, that of the eyed side extending along ridge of body from notch in isthmus to front of anal, base of ventral on blind side shorter; pectoral reaching past curve on eyed side, its mate much smaller; caudal rather long. Color light olive brown, almost translucent, everywhere marbled with paler, and with many small, irregular, sharply defined black spots; dorsal, anal, and caudal with larger, round, blended spots of dark brown; pectoral with brown, interrupted cross lines. This small flounder much resembles the European Brill (*Bothus rhombus*), but is smaller, thinner, and more translucent in body. Its weight rarely exceeds a pound or two, and its value as a food-fish is but slight; nevertheless, it is a near ally of the European Turbot (*Psetta maxima*), and in its technical characters it very closely agrees with the latter species. Atlantic coast of United States, from Casco Bay to South Carolina; common. (*maculatus*, spotted.)

Pleuronectes maculatus, Mitchill, Rept. in part, Fish. N. Y., 9, 1814, New York; De Kay, New York Fauna: Fishes, 301, pl. 47, fig. 151, 1842; Storer, Synopsis, 479, 1846; Storer, Hist. Fish. Mass., 204, 1867; Jordan & Goss, Review of Flounders and Soles, 258, 1889.

Pleuronectes aquosus, Mitchill, Trans. Lit. and Phil. Soc. N. Y., I, 1815, 389, pl. 2, fig. 3, New York.

Lophopsetta maculata, Gill, Proc. Ac. Nat. Sci. Phila. 1862, 216; *ibid*, 1864, 220; Jordan & Gilbert, Proc. U. S. Nat. Mus. 1878, 371.

Bothus maculatus, Jordan & Gilbert, Synopsis, 815, 1883.

Rhombus aquosus, Günther, Cat., IV, 411, 1862.

1044. PLATOPHRYS, Swainson.

Solea, Rafinesque, Indice di Ittiologia Siciliana, 52, 1810 (*rhomboide*); not of Quensel, 1806.

Platophrys, Swainson, Nat. Hist. Class'n Fishes, II, 302, 1839 (*ocellatus*).

Peloria, Cocco, Intorno ad Alcuni Pesci del mar di Messina, Giorn. del Gabin., 1844, 21–30, Lettere di Messina (*heckeli*, a larval form of *P. podas*); not *Pelorus* of Montfort, 1808.

? Coccolus,* BONAPARTE, in COCCO, Alcuni Pesci Messina, 21, 1844 (*annectens;* larval form—probably of *P. podas*, with the right eye in transit to the left side).

Bothus, BONAPARTE; Catologo Metodico, 49, 1846 (*podas*); not of RAFINESQUE.

Rhomboidichthys, BLEEKER, Act. Soc. Sci. Indo-Nederl. Maaad. and Makassar, 67, 1357–58 (*myriaster*).

Platophrys, BLEEKER, Comptes Rendus Acad. Sci. Amsterd., XIII, 1862, Pleuron., 5 (*ocellatus*).

Eyes and color on the left side. Body ovate, strongly compressed; mouth of the large type, but comparatively small; the maxillary $\frac{1}{3}$ or less of the length of the head; teeth small, subequal, in 1 or 2 series; no teeth on vomer or palatines. Interorbital space broad and concave, broadest in adult males. Gill rakers moderate. Dorsal fin beginning in front of eye, all its rays simple; ventral of colored side on ridge of abdomen; caudal convex behind; pectoral of left side usually with 1 or more filamentous rays, longest in the male. Scales very small, ctenoid, adherent; lateral line with a strong arch in front. Coloration usually variegated.

This well-marked genus is widely diffused in the warm seas. The sexual differences are greater than usual among flounders, and the different sexes have often been taken for different species. As a rule, in the males, the pectoral fin on the left side is much prolonged, the interorbital area is much widened and very concave, and there are some tubercles about the snout and lower eye. The young fishes, as is usually the case, resemble the adult females. Lately, Dr. Emery has shown that the larval flounder, known as *Peloria heckeli*, is in all probability the young of *Pleuronectes podas*. The generic name, *Coccolus*, based on forms slightly more mature than those called *Peloria*, probably belongs here also. We have seen no larval forms so young as those which have been described as *Peloria heckeli*. We have, however, examined small transparent flounders, one with the eyes quite symmetrical, taken in the Gulf Stream, and another with the eyes on the left side, taken at Key West. Both these may be larvæ of *Platophrys ocellatus*. The figures published by Emery seem to make it almost certain that the corresponding European forms belong to *P. podas*, although some doubt as to this is expressed by Facciolà. The species of *Platophrys* are widely distributed through the warm seas, no tropical waters being wholly without them. All the species of *Platophrys* are extremely closely related and can be distinguished with difficulty. On the other hand, the variations due to differences of age and sex are greater than in any other of our genera. The following analysis of the species of *Platophrys* is very unsatisfactory. There are certainly 3 species (*podas*, the European species, *maculifer*, and *lunatus*) which are known to be distinct in their adult state. The young forms of *maculifer* and *lunatus* are not well known, nor is it known how they differ from *ocellatus*, *spinosus*, and other species which presumably reach a smaller size. Only a thorough study of the species, in all stages of development in their native waters, can give us the characters by which the species can be really discriminated. (πλατύς, broad; ὀφρύς, eyebrow.)

* "Parvus mole et pleuronectiformis, medius inter Pleuronectidas et Bibroniinos hic piscis videtur! Attamen dum illi oculos unilaterales habeant, iste vero bilaterales; in hoc novo genere oculi, alter a latere, altere in vertice vix ad appositum latus convenus positi sunt." (Bonaparte: quoted by Facciolà, Su di Alcuni Rari Pleuronettidi.)

a. Anal rays, at least anteriorly, each with a spinule at base (these formed by a slight widening of the tip of the interhæmal spines, each being covered by a little rough scale); front of dorsal with similar projections.

b. Color brown, with pale rounded spots; fins dotted with brown; a faint dark spot at first ⅓ of lateral line; snout with horny points; mouth small, the maxillary reaching front of eye. SPINOSUS, 3030

aa. Anal rays without spinules at their base.

c. Anterior profile of head convex before the interorbital area, the very short snout scarcely forming a reentrant angle at its base; form elliptic-ovate, the outlines more regular than in *P. lunatus.*

d. Dorsal rays 85 to 95.

e. Scales not very small, about 75 pores in lateral line; no blue markings, at least in the young.

f. Mouth small, the maxillary 3 in head; no spines about the snout; eye 3⅓ in length; interorbital width 3 in head (in the type); pectoral short; curve of lateral line 6 times in straight part. Color dark brown, with numerous stellate white spots, the most distinct of them with darker edgings; these generally scattered over the body; but some of them on sides of body gathered together in little rings; these spots blue rather than white in life. CONSTELLATUS, 3031.

ff. Mouth smaller, the maxillary 3¾ in head. Color light grayish, tinged with reddish, with small round spots of darker gray, and with lighter rings inclosing spaces of ground color. OCELLATUS, 3032.

ee. Scales smaller, 90 to 95 pores in lateral line. Color of adult, reddish gray, the body everywhere covered with rings formed of round, sky-blue spots, which are not confluent and not edged with black; besides these, very few detached spots or other blue markings. MACULIFER, 3033.

dd. Dorsal rays 105; anal rays 80; pectoral short; interorbital space 2⅜ in head; depth 1⅔ in length; scales 91; body deep. Color (specimen 4¾ inches long) grayish, much spotted and mottled with whitish, no blue in young example. ELLIPTICUS, 3034.

cc. Anterior profile of head strongly concave before interorbital area, the projecting snout leaving a marked reentrant angle above it.

g. Mouth not very small; maxillary 3 in head. Color dark olive, with many rings, curved spots, and small round dots of sky blue edged with darker on body, these largest near middle of sides, where some are as large as eye; 3 obscure dark blotches on straight part of lateral line. LUNATUS, 3035.

gg. Mouth small; maxillary 3⅝ in head. Color highly variegated with different shades of gray, the pale blotches rounded, very irregular in size and position; no blue spots. LEOPARDINUS, 3036.

3030. PLATOPHRYS SPINOSUS (Poey).

Depth 1⅓. D. about 74; A. about 57; scales about 80. Anal rays, at least anteriorly, each with a spinule at base, these formed by a slight widening of the tips of the interhæmal spines, each being covered by a little rough scale; front of dorsal with similar projections. Snout with horny points; mouth small, the maxillary reaching front of eye. Eyes very wide apart, 2⅜ in head, the interorbital space 1⅕ in head; pectoral fin short; curve of lateral line 5 in straight part. Color brown, covered with pale rounded spots; fins dotted with brown; a faint dark spot at first third of lateral line. Described from specimens from Cuba, probably the types, 4¼ inches

long, which have been partly dried before being placed in alcohol. Cuba. The original description of this species is a very scanty one. In all respects, unless it be the color, it agrees with the European species, *Platophrys podas*. We have found 2 small specimens sent by Professor Poey to the Museum of Comparative Zoology, which may be the types of this species. They are 4¼ inches long, and have been partly dried in the sun. A result of this has been to increase the prominence of the interhæmal spines. Whether these be the original types or not, the species is an extremely doubtful one. The eyes are farther apart in these specimens than in any of *Platophrys ocellatus*, which we have examined. They agree in this respect with Agassiz's figure of *Rhombus ocellatus*. (*spinosus*, spinous.)

Rhomboidichthys spinosus, POEY, Synopsis, 409, 1868, Cuba; POEY, Enumeratio, 139, 1875. *Platophrys spinosus*, JORDAN & GOSS, Review Flounders and Soles, 266, 1889.

3031. PLATOPHRYS CONSTELLATUS, Jordan.

Head 4; depth 1½; eye 3⅛ in head; interorbital width 3. D. 89; A, 65; scales 75. Body elliptic-ovate, the outlines more regular than in *P. lunatus;* anterior profile of head convex before the interorbital area, the very short snout scarcely forming a reentrant angle at its base; anal rays without spinules at their base; mouth small, the maxillary 3 in head; no spines about the snout; pectoral short; curve of lateral line 6 times in straight part. Color dark brown, with numerous stellate white spots, the most distinct of them with darker edgings; these generally scattered over the body, but some of them on sides of body are gathered together in little rings (perhaps these spots are blue rather than white in life); fins mottled with dark brown, the pectoral finely barred. Specimens examined 3½ inches long. Galapagos Archipelago. Originally described from 3 specimens, the largest 3½ inches long, numbered 11146 on the register of the Museum of Comparative Zoology. They are from James Island, in the Galapagos. The species is closely related to *P. ocellatus* and others, but in color, at least, it is different, and its habitat is remote; locally common. (*constellatus*, with star-like spots.)

Platophrys constellatus, JORDAN, in JORDAN & GOSS, Review Flounders and Soles, 266, 1889, James Island, Galapagos Archipelago. (Types in M.C. Z.)

3032. PLATOPHRYS OCELLATUS (Agassiz).

Head 4 in length; depth 1⅓; eye (lower) 3⅜ in head; snout 5. D. 85; A. 64; scales 75 (pores); vertebræ 37. Body ovate, deep anteriorly, the profile descending steeply, rendered abruptly concave in front of interorbital space by the conspicuously projecting short snout. Mouth very small and oblique, the maxillary reaching vertical from front of lower eye, 3¾ in head; tip of lower jaw entering the profile. Teeth fine, conical, in 2 series in the upper jaw, 1 in the lower, those of the outer row in upper jaw larger and more widely separated than those of the inner series. Snout very short, equaling interorbital width. Interorbital space narrow, deeply concave, closely scaled. Eyes large, the lower in advance of upper. Gill rakers obsolete, 7 rudiments on horizontal branch of anterior arch. Scales moderate, not extending on the fins, those on colored side

ctenoid, those on blind side smooth; arch of lateral line short and high, its base contained 4½ to 5 times in the straight portion. Dorsal fin beginning opposite anterior nostril, the rays nearly uniform in length, the longest about ½ head; pectoral of colored side 4¾ in length; ventral of colored side beginning under middle of lower eye, with 6 rays; the right ventral with 5 rays. Color in life, light grayish with reddish tinge, covered with small round spots of darker gray and with lighter rings inclosing spaces of the ground color; vertical fins similarly colored, with a small black spot near base of each ninth or tenth ray; 2 black spots on median line of body divide the length into nearly equal thirds; some other small black spots scattered over colored side. Western Atlantic, from Long Island to Rio Janeiro, on sandy shores. Here described from Key West specimens, types of *P. nebularis*. This species is very common at Key West in clear, shallow water on sandy bottom. The largest of the numerous specimens taken is 3 inches in length. A specimen similar to these has been taken by Dr. Bean on the south coast of Long Island. This seems to be the same as the Cuban species called *Rhomboidichthys ocellatus* by Poey, and some of the specimens sent by Poey to the Museum of Comparative Zoology are apparently identical with the types of *P. nebularis*. In the Museum of Comparative Zoology we have compared specimens of the real *Platophrys ocellatus* (No. 11423, Rio Janeiro, Agassiz) with a representative specimen of *P. nebularis* (No. 26147, from the Tortugas, Florida), and are unable to find any differences. We adopt, therefore, the name *Platophrys ocellatus* for all, and regard it as one of the widely distributed flounders, like *Etropus crossotus* and *Citharichthys spilopterus*. (*ocellatus*, with eye-like spots.)

Rhombus ocellatus, AGASSIZ, Spix, Pisc. Brasil., 85, pl. 46, 1829, Brazil.
Rhombus bahianus, CASTELNAU, Anim. nouv. rares Amérique du Sud, 1855, Bahia.
Platophrys nebularis, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1884, 31, 143, Key West (Type, 34972. Coll. Dr. Jordan); GOODE & BEAN, Oceanic Ichthyology, 441, 1896.
Platophrys ocellatus, SWAINSON, Nat. Hist. Class'n Fishes, II, 302, 1839; JORDAN & GOSS, Review Flounders and Soles, 266, 1889.
Rhomboidichthys ocellatus, GÜNTHER, Cat., IV, 433, 1862; POEY, Synopsis, 408, 1868.

3033. PLATOPHRYS MACULIFER (Poey).

Head 4; depth 1⅝. D. 90 to 95; A. 70; scales 90 to 95. Body elliptical, ovate. Mouth small, oblique, the maxillary 3⅔ in head; teeth in each jaw in 2 irregular series; filamentous rays of pectorals reaching very nearly to last rays of dorsal; arch of lateral line short and high, its length 1⅓ times its height and 2⅔ in head; snout very short, 4 in head; interorbital area 3¾ in head. Color of adult reddish gray, the body everywhere covered with rings formed of round, sky-blue spots, which are not confluent and are not edged with black; besides these, very few detached spots or other blue markings; head with similar blue spots, but no rings; area inclosed in the blue rings not different from the ground color; caudal with blue spots, other fins with none; dorsal and anal mottled; a large, diffuse, dusky spot at front of straight part of lateral line; 1 better defined on middle of lateral line; a faint one farther back; pectorals grayish, with dark bars. Cuba. We identify specimens taken by Dr.

Jordan at Havana with this species. In the Museum of Comparative Zoology are other specimens similar to these, sent to Cambridge by Poey. In several respects these specimens agree fairly with Poey's *P. ellipticus*, but that species is said to have 104 dorsal rays. (*macula*, spot; *fero*, I bear.)

? *Pleuronectes maculiferus*, POEY, Memorias, II, 316, 1860, Cienfuegos. (Coll. Poey.)
? *Rhomboidichthys maculiferus*, POEY, Synopsis, 408, 1868; POEY, Enumeratio, 139, 1875.
Platophrys maculifer, JORDAN & GOSS, Review Flounders and Soles, 267, 1889.
Platophrys ellipticus, JORDAN, Proc. U. S. Nat. Mus. 1886, 51; not of POEY.

3034. PLATOPHRYS ELLIPTICUS (Poey).

Depth $1\frac{3}{4}$. D. 105; A. 80; scales 91. Body elliptical, ovate; anterior profile of head convex before the interorbital area; pectoral short; interorbital space $2\frac{3}{4}$ in head; body deep. Color (specimen $4\frac{3}{4}$ inches long) grayish, much spotted and mottled with whitish; no blue (in young example). Cuba. Poey describes his *P. ellipticus* as having 104 dorsal rays. In none of our other species does the number of these rays reach 100. Among the specimens sent by Poey to the museum at Cambridge is 1, described above, $4\frac{3}{4}$ inches long, which has 105 dorsal rays. We have therefore assumed that the species to which this specimen belongs is the real *P. ellipticus*, and that the one heretofore called *P. ellipticus* is Poey's *P. maculifer*. Both these assumptions are open to considerable doubt. (***ellipticus***, elliptical.)

? *Pleuronectes ellipticus*, POEY, Memorias, II, 315, 1860, Cuba. (Coll. Poey.)
? *Rhomboidichthys ellipticus*, GÜNTHER, Cat., IV, 434, 1862; POEY, Synopsis, 408, 1868; POEY, Enumeratio, 139, 1875.
Platophrys ellipticus, JORDAN & GOSS, Review Flounders and Soles, 267, 1889.

3035. PLATOPHRYS LUNATUS (Linnæus).

(PEACOCK FLOUNDER.)

Head $3\frac{3}{5}$ in length; depth 2. D. 93; A. 70; scales 90; lower eye 6 in head; maxillary $2\frac{3}{5}$; interorbital $2\frac{3}{5}$; highest dorsal rays $2\frac{3}{5}$; highest anal rays $2\frac{1}{2}$; caudal $1\frac{1}{4}$; base of ventral of eyed side $3\frac{1}{4}$. Vertebræ $9+30=39$. Body elliptical, ovate, strongly compressed; anterior profile concave, the snout projecting, leaving a reentrant angle above it; mouth moderate, the maxillary reaching to middle of pupil of lower eye; jaws subequal, the lower with a well-developed knob at symphysis, teeth small, in an irregular double series in each jaw; anterior end of maxillary with a large blunt spine, pointing outward and forward, a smaller one behind it on upper edge of maxillary, pointing upward and backward; interorbital very wide and deeply concave; orbital rim, below on upper orbit, above on lower, broken up into blunt papillæ; gill rakers short and thick, 9 developed on lower part of arch, none on upper. Anterior part of interorbital, snout, maxillary, and mandible, naked; scales all cycloid; the rays of dorsal and anal with scales, a few on ventral of eyed side; arch of lateral line 5 in straight part. Pectoral of eyed side filamentous, reaching to base of caudal, its mate of opposite side shorter, about $1\frac{3}{4}$ in head; origin of dorsal over snout; ventral of eyed side with a long base, extending from

angle at isthmus, along ridge of body, slightly past front of anal; base of ventral of blind side ½ the length of that of its mate; caudal with the middle rays produced, double convex. Color dark olive, with many rings, curved spots, and small round dots of sky blue edged with darker on body, these largest near middle of sides, where some are as large as the eye; 3 obscure dark blotches on straight part of lateral line; head and vertical fins with sharply defined blue spots, which are mostly round; spots on opercles larger and curved; pectorals with dark bars. West Indies, north to Florida; common. Here described from a specimen from Green Turtle Cay, Florida, 14 inches in length. This handsome and curiously colored species is not rare in the waters of the West Indies. The specimens examined by us are from Cuba, Sombrero, St. Thomas, and other localities in the West Indies. The original figure of this species published by Catesby is a very good one and leaves no room for doubt as to the species intended. The figure of Bloch, called *Pleuronectes argus*, is also fairly accurate, and can refer to no other species. This species reaches a length of some 18 inches, and is the largest in size of the American species of *Platophrys*. We have never seen any young examples which certainly belong to it, and till its development is traced some of the species known from small examples only must be doubtful. (*lunatus*, crescent-shaped, from the spots.)

Solea lunata et punctata (the Sole), CATESBY, Nat. Hist. Carolina, tab. 27, 1725, Bahamas.
Pleuronectes lunatus, LINNÆUS, Syst. Nat., Ed. x, 269, 1758, Bahamas; based on CATESBY; and of the various copyists.
Pleuronectes argus, BLOCH, Ichthyol., tab. 48, 1783, Martinique; after Plumier.
?*Pleuronectes surinamensis*, BLOCH & SCHNEIDER, Syst. Ichth., 156, 1801, Surinam; "*satis parva et glabra;*" fins scaly; mouth small; lateral line arched in front; D. 96; A. 55.
Rhomboidichthys lunatus, GÜNTHER, Cat., IV, 433, 1862; POEY, Synopsis, 408, 1868.
Rhomboidichthys lunulatus, POEY, Enumeratio, 138, 1875.
Platophrys lunatus, JORDAN, Proc. U. S. Nat. Mus. 1886, 51; JORDAN & GOSS, Review Flounders and Soles, 267, 1889.

3036. PLATOPHRYS LEOPARDINUS (Günther).

Head 3¾ in length; depth 1¾; eye (lower) 3½ in head. D. 86 to 88; A. 64 (62 to 66); scales about 80. Mouth very small, the maxillary 3⅞ in head; teeth very small, biserial above. Interorbital space concave, rather broad, its width 3½ in head. Eyes large, the lower considerably before the upper. Lateral line with a short sharp curve anteriorly. Gill rakers very small. Anterior rays of dorsal not elevated; left pectoral not produced, little longer than right, 1¼ in head. Coloration highly variegated with different shades of gray, the pale blotches rounded, very irregular in size and position; no distinct black spots along the lateral line; a large whitish cloud between the eyes; blind side pale, scaled like the eyed side. Gulf of California. This species is known only from the original type from unknown locality, and from a single specimen, 2⅜ inches long, in the United States National Museum, taken by Mr. H. F. Emeric, at Guaymas, Sinaloa. From this the above description was taken. (*leopardinus*, leopard-like.)

Rhomboidichthys leopardinus, GÜNTHER, Cat. Fish., IV, 434, 1862, locality unknown.
Platophrys leopardinus, JORDAN, Proc. U. S. Nat. Mus. 1884, 260, specimen from Guaymas; JORDAN & GOSS, Review of Flounders and Soles, 268, 1889.

1045. PERISSIAS, Jordan & Evermann.

Perissias, JORDAN & EVERMANN, new genus (*tæneopterus*).

This genus differs from *Platophrys* in the larger scales, narrower interorbital, and especially in the greatly produced ribbon-like lobe at the front of the dorsal. From *Engyprosopon* it differs in the short thick gill rakers and in the produced dorsal rays. The lateral line is wanting on the blind side. Deep sea. (*περισσός*, strange.)

3037. PERISSIAS TÆNIOPTERUS (Gilbert).

Head 3⅔; depth 2½. D. 86 to 88; A. 67 to 70; scales 60 to 65, the arch with 15 pores; 20 scales in a series running upward and backward from lateral line. Body elongate; caudal fin subsessile, the last anal and dorsal rays inserted near rudimentary caudal rays; height of caudal peduncle 4 in height of body; upper profile descending very obliquely anteriorly, a slight reentrant angle in front of lower eye; in males the profile slightly angulated in front of upper orbit, below which it ascends more steeply; lower eye much in advance of upper; in females eyes close together, the vertical from middle of lower eye passing through front of upper orbit; diameter of upper orbit 3½ in head; in males the lower eye may be entirely in advance of upper; in females 3 inches long, and in very young males the interorbital space is a narrow, concave, scaleless groove, less than diameter of pupil, running into a deep pit behind lower eye; in males 2 inches long the interorbital space has already widened, and in specimens 3½ inches long is as wide as longitudinal diameter of orbit; it is traversed by an oblique ridge running upward and backward from front of lower eye, separating the anterior scaleless portion from the deep scalely pit behind; supraorbital ridge of lower eye serrated, forming a strong series of spines, less marked in females; anterior rim of upper orbit similarly but less strongly marked; a strong double spine on maxillary in front of nostrils; a spine near end of maxillary in males; mouth small, maxillary not reaching front of pupil, equaling diameter of orbit; teeth small, in a single close-set series in each jaw, equally developed on both sides, with enlarged canines; gill rakers very short and weak, 8 on horizontal limb; anterior nostrils with very short flaps; dorsal beginning above front of lower eye; in all specimens, females as well as males, the first 2 rays detached from the rest of fin, the second ray produced into a flat, ribbon-shaped filament about as long as head; dorsal and anal rays all unbranched; median caudal rays forked; no anal spine; ventral of colored side on ridge of abdomen, the 2 anterior rays in males connected by membrane at base only, produced into flat filaments as long as head, extending far beyond front of anal; pectoral of left side well developed, but small, slightly more than ½ head; that of blind side little developed, about ½ diameter of orbit; scales of left side strongly ctenoid, absent on interorbital space, snout, maxillary, and mandible; lateral line with strong curve anteriorly, the cord of which is contained 5 times in straight portion. Along lateral line are occasional broad cutaneous flaps, colored blue in life; scales of blind side cycloid, the tubes of lateral line obsolete, the course of lateral

line indistinctly indicated by pits at bases of scales and occasional pore-like markings; median rays of dorsal and anal on left side with series of ctenoid scales, otherwise scaleless; caudal rays with double series on both sides. Color on left side olive brown, with many small irregular spots of light gray, with darker border; 3 or 4 dark blotches along lateral line; along dorsal and ventral outlines about 5 pairs of light spots, broadly ocellated with blackish; males with a bright blue spot on anterior profile at base of each of first 10 or 12 dorsal rays and 1 on end of snout; blind side in males with a broad oblique bar covering about ¼ of sides, bluish black in life, dark brown in spirits; from its upper anterior part a number of narrow parallel streaks run forward toward head, much as in *Engyophrys sancti-laurentii;* filamentous rays of dorsal and ventral white; fins all speckled; a small black spot at base of median caudal rays. Differing from all known species of *Platophrys* in the ribbon-shaped prolongations of second dorsal ray and first and second ventral rays of eyed side, and in the obsolete lateral line of blind side. Several specimens from the Gulf of California and the western coast of Lower California, in 40 fathoms. (Gilbert.) (*ταινία*, ribbon; *πτερόν*, fin.)

Platophrys tæniopterus, Gilbert, Proc. U. S. Nat. Mus. 1890, 118, Gulf of California, north of La Paz, at Albatross Station 2998, Lat. 24° 51′ N., Long. 110° 39′ W., in 40 fathoms. (Type, No. 43095. Coll. Gilbert.)

1046. ENGYOPHRYS, Jordan & Bollman.

Engyophrys, Jordan & Bollman, Proc. U. S. Nat. Mus. 1889, 176, (*sancti-laurentii*).

This genus is allied to *Platophrys*, Swainson, but differs from it in having the interorbital space very narrow and armed with a spine, and the scales of moderate size and ctenoid. Gill rakers obsolete. No anal spine. Gill membranes entirely separate. It is still nearer the genus *Engyprosopon*, Günther, but in that group the interorbital space is broader and the gill rakers are developed and slender. (*ἐγγύς*, near together; *ὀφρύς*, eyebrow.)

3038. ENGYOPHRYS SANCTI-LAURENTII, Jordan & Bollman.

Head 2⅖ to 2⅐ (3 to 3½); depth 1⅜ to 2 (2 to 2⅛). D. 78 to 85; A. 68 to 72; scales 60 to 68, along lateral line. Body broadly ovate, much compressed, the greatest depth over pectorals; dorsal and ventral outlines equally curved; profile scarcely concave before eyes. Mouth very small, oblique, the maxillary reaching opposite pupil of lower eye, 4 to 4½ in head. Teeth present on blind side, well developed, close set and even, none on vomer. Snout short, 4½ to 5 in head. Interorbital space a very narrow, sharp, scaleless ridge, the ridge forking above pupil, leaving a very narrow concavity anteriorly; lower ridge armed with a strong spine, turned backward, inserted just above pupil of lower eye. Anterior orbital rim of upper eye rather high, entering profile. Eyes large, lower in advance of upper, 3¾ to 4 in head. Gill rakers almost obsolete, represented by 5 or 6 small fleshy papillæ. Scales moderately small, ctenoid, and not very firmly attached; small scales on rays of dorsal and

anal fins; arch of lateral line short and small, but abrupt, 4 to 5 times in straight part. Dorsal beginning on blind side just behind posterior nostril and in front of eye; pectoral of colored side 2 in head, that of blind side 2½ in head; ventrals of colored side slightly longest, 3 in head; that of colored side with 6 rays, of blind side with 5 or 6 rays. Color of left or eyed side, blackish brown, with scattered white and black spots, the latter most prominent along base of dorsal and anal fin; 3 large, black, nonocellated blotches on straight part of lateral line, the first at beginning, second at middle, and third on peduncle; fins dusky; dorsal and anal with scattered white and black spots; caudal with 5 black spots arranged in a curved series; blind side with 5 or 6 curved parallel dusky bands as wide as eye, the first beginning on interopercle and curving across cheek to along base of dorsal; second beginning at throat and curving along posterior margin of preopercle, and extending on back, parallel with the first from vent; third curving around in front of pectorals, across posterior part of opercle, and extending to base of dorsal fin behind the middle; rest behind pectorals. All of these bands fade out behind middle of body, so that the posterior portion is immaculate. In young examples these bands are very faint or obsolete. Coast of Colombia, southwest of Panama. Numerous specimens, the largest about 4½ inches long, were dredged at *Albatross* Station 2795, at a depth of 33 fathoms, and at *Albatross* Station 2805 at a depth of 51½ fathoms. This peculiar species is distinguished from the species of *Platophrys* and *Engyprosopon* by its very narrow interorbital ridge, from the species of *Arnoglossus* by the form of the body, the short gill rakers, etc., and from all related species by the peculiar coloration of the blind side. (Named for St. Lawrence, in allusion to the gridiron-like markings of the blind side.)

Engyophrys sancti-laurentii, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 176, Pacific Ocean, off coast of Colombia, at Albatross Station 2805, Lat. 7° 56′ N., Long. 79° 41′ 30″ W., and Station 2795, Lat. 7° 57′ N., Long. 78° 55′ W. (Type, No. 41155.)

1047. TRICHOPSETTA, Gill.

Trichopsetta, GILL, Proc. U. S. Nat. Mus. 1888, 603 (*ventralis*).

Body ovate, covered with rather large, ctenoid adherent scales; mouth moderate, the chin prominent; vomer toothless; teeth small, somewhat enlarged and hooked in front, uniserial; maxillaries obliquely truncated behind; interorbital area a narrow ridge, with a median groove in front; none of the dorsal rays produced; ventrals free from the anal; caudal fin subsessile; both pectoral fins present; right ventral much produced, the left on the ridge of the abdomen; lateral line with a strong arch in front. (θρίξ, hair; ψῆττα, turbot, from the prolonged ventral.)

3039. TRICHOPSETTA VENTRALIS (Goode & Bean).

Head 4 in body; depth 2⅓. D. 93; A. 73; pectoral 11 (eyed side), 7 or 8 (blind side); scales 19–66–23; eye 3⅔ in head; maxillary scarcely 2; interorbital very narrow, scaleless, its width 8 in eye; scales strongly ctenoid; dorsal beginning upon snout upon the blind side, in advance of eyes, its

highest rays equaling length of mandible; origin of anal under base of pectoral, its longest ray equaling or slightly exceeding ½ the distance of its anterior ray from snout; caudal equal to length of head without snout; pectorals inserted considerably below origin of lateral line, close to gill opening, that of the eyed side 6 in length of body; that of the blind side almost as long as head. Color light brownish gray; a dark blotch as long as eye on the anterior rays of the anal, a few obscure on different parts, of lighter hue at the junction of the curved and straight portion of the lateral line. (Goode & Bean.) Deep waters of the Gulf of Mexico. (*ventralis*, pertaining to the ventrals.)

Citharichthys ventralis, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 592, deep waters of Gulf of Mexico. (Coll. *Albatross.*)
Arnoglossus ? ventralis, JORDAN & GOSS, Review Flounders and Soles, 262, 1889.
Trichopsetta ventralis, GOODE & BEAN, Oceanic Ichthyology, 440, pl. 109, fig. 372, 1896.

1048. SYACIUM, Ranzani.

Syacium, RANZANI, Novis Speciebus Piscium, Diss. Sec., 20, 1840 (*micrurum*).
Hemirhombus, BLEEKER, Comptes Rendus Acad. Sci. Amsterd., XIII, Pleuron., 4, 1862 (*guineënsis*).
Aramaca, JORDAN & GOSS, in JORDAN, Cat. Fish. N. A., 133, 1885 (*pœtula*).

Body elliptic-ovate, much compressed; interorbital space broad in the males and more or less concave, narrowed in the female; mouth moderate, the gape curved; teeth in the upper jaw biserial, in the lower uniserial; the front teeth of the upper jaw enlarged; vomer toothless; scales rather large, ciliate; lateral line without arch in front; pectoral fins on both sides present; septum of gill cavity below gill arches without foramen; a deep emargination near the isthmus; gill rakers short and thick; dorsal low, its anterior rays not elevated; pectorals both present; caudal subsessile; no anal spine; pectorals produced in the males; ventral fins short, that of colored side on ridge of abdomen. This genus contains a considerable number of species, mostly American and African, which form a transition from *Platophrys* to *Citharichthys*. They fall readily into 2 groups distinguished by the width of the interorbital space. As this width is dependent on age, and as it is subject to various intergradations, the group *Aramaca* founded on it can not be admitted as a distinct genus. (συάκιον, diminutive of σῦαξ, a kind of pulse, the application unexplained.)

a. Snout and orbits without spines or spinous processes.
b. Scales rather large, 50 to 57 in the lateral line; interorbital space broad. Color nearly plain brown, with darker dots or mottlings, no ring-like spots or ocelli; fins mottled; left pectoral barred; blind side sometimes wholly or partly dusky, especially in northern specimens. PAPILLOSUM, 3040.
bb. Scales rather small, 58 to 70 in the lateral line.
c. Scales 65 to 70. Color dark brown, with many rings and spots of light gray and blackish, some of the dark rings with a black central spot; a diffuse dusky blotch on lateral line above pectoral, and 1 near base of caudal peduncle; fins with numerous inky spots and dark markings; blind side pale. MICRURUM, 3041.
cc. Scales 58 to 60.

d. Interorbital space in male broader than eye. Color light brown, with grayish and light bluish dots, some darker areas, and a few round brown spots ocellated with lighter; interorbital space with a vertical brown bar bordered by lighter; fins mottled and spotted.
LATIFRONS, 3042.

dd. Interorbital space not broader than pupil. Color light olive brown, nearly uniform, the vertical fins with elongate dark spots.
OVALE, 3043.

3040. SYACIUM PAPILLOSUM (Linnæus).

Head $3\frac{3}{8}$ in length; depth $2\frac{1}{3}$. D. 82; A. 63 to 70; scales 53; eye 5 in head; maxillary $2\frac{3}{8}$; pectoral of eyed side $1\frac{1}{6}$; caudal $1\frac{2}{3}$. Body elliptic-ovate, the anterior profile regularly decurved, forming an angle above the snout; mouth rather large, arched; maxillary extending to below middle of eye, its posterior end concave; teeth in upper jaw in 2 series, some of the outer forming small canines; lower teeth in 1 row; eye large, 4 in head; lower eye in advance of upper, especially in the adult; interorbital space broad, concave, greater than the long diameter of the eye in the males, about equal to the vertical diameter in the females; accessory scales very numerous; mandible, maxillary, and interorbital with scales; gill rakers short, scarcely as long as pupil, about 2 + 8; dorsal rather low, beginning slightly in front of lower eye, the first 3 or 4 rays on blind side, the anterior rays produced beyond the membrane; ventrals with moderate base, that of eyed side on ridge of body, that of blind side slightly in advance of its mate; anal beginning a little in advance of pectoral; pectoral of eyed side pointed behind, the upper rays filamentous (at least in the male); caudal double truncate. Vertebræ 10 + 26 = 36. Color nearly plain brown, with darker dots or mottlings, no ring-like spots or ocelli; fins mottled; left pectoral barred; blind side sometimes wholly or partly dusky, especially in northern specimens. Charleston to Rio Janeiro, in rather deep water. Here described from an adult specimen from Charleston, a foot in length. Of the species found in the deep waters about Pensacola, and called by Dr. Bean *Hemirhombus pœtulus*, we have numerous specimens. Lately we have received from Mr. Charles C. Leslie, of Charleston, a specimen which shows its presence also in Carolina waters. It has not yet been recorded from Cuba, but in the Museum of Comparative Zoology is a specimen (26104) taken by Mr. Samuel Garman, at Kingston, Saint Vincent. But its range extends much farther to the southward, for among the collections made by Professor Agassiz, at Rio Janeiro, there are many specimens (11375, 4666), the largest about a foot long. These seem to be completely identical with Florida examples, differing only in having the blind side pale, it being usually partly blackish in northern samples. These Brazilian specimens agree very closely with the figure of *Rhombus soleæformis*, except that Agassiz has represented that species as having a dusky blotch at the shoulder. No such marking is apparent in any of our specimens. The coloration and the breadth of the interorbital both render it unlikely that Agassiz's *solcæformis* could have been *micrurum*. The *Aramaca* of Marcgrave, which is the sole basis of *Pleuronectes papillosus*. *Pleuronectes macrolepidotus*, and *Rhombus aramaca*, can not well be any known species other than the present one. According

to Marcgrave's rude figure and his description, this species has the form of a sole, the eyes wide apart, the left pectoral produced, the mouth very large, the body oblong, and the coloration stone-like (sand color) on the left side and white on the eyed side. *Syacium micrurum* is not colored in that way, and its eyes are not noticeably far apart. We therefore adopt for this species the oldest name, *Syacium papillosum*. (*papillosus*, having papillæ.)

Aramaca, MARCGRAVE, Hist. Brasil., 181, 1648, Brazil.

Pleuronectes papillosus, LINNÆUS, Syst. Nat., X, 271, 1758, Brazil; based on MARCGRAVE.

? *Pleuronectes macrolepidotus*, BLOCH, Ausländische Fishe, VI, 25, tab. 190, 1787; apparently based on MARCGRAVE.

Pleuronectes aramaca, DONNDORF, Beiträge zur Ausgabe des Linnæischen Natursystems, XIII, 386, 1798; after MARCGRAVE.

Rhombus aramaca, CUVIER, Règne Animal, Ed. 2, II, 341, 1829; after MARCGRAVE.

Rhombus soleæformis, AGASSIZ, Spix, Pisc. Brasil., 86, tab. 47, 1829, Atlantic Ocean.

Hippoglossus internedius, RANZANI, Novis Speciebus Piscium Dissertatio Secundo, 1840, 14, pl. 4, Brazil.

Hemirhombus soleæformis, GÜNTHER, Cat. Fish., IV, 423, 1862.

Hemirhombus pœtulus, BEAN MS., JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 304, Pensacola (Coll. Silas Stearns); GOODE & BEAN, Proc. U. S. Nat. Mus. 1882, 414; BEAN, Cat. Coll. Fish U. S. Nat. Mus. 1883, 45.

Citharichthys aramaca, JORDAN & GILBERT, Synopsis, 816, 1883.

Citharichthys pœtulus, JORDAN & GILBERT, Synopsis, 964, 1883; JORDAN, Proc. U. S. Nat. Mus. 1884, 38; GOODE & BEAN, Oceanic Ichthyology, 448, pl. 109, fig. 373, 1896.

Aramaca papillosa, JORDAN, Proc. U. S. Nat. Mus. 1886, 602; synonymy confused with *S. micrurum*.

Aramaca soleæformis, JORDAN, Proc. U. S. Nat. Mus. 1886, 602.

Syacium papillosum, JORDAN & GOSS, Review Flounders and Soles, 268, 1889.

3041. SYACIUM MICRURUM, Ranzani.

Head $3\frac{4}{5}$ in length; depth $2\frac{2}{3}$. D. 87 to 92; A. 54 to 68; scales 65 to 70 (pores); eye 4 in head; maxillary $2\frac{1}{2}$ to 3. Form regularly elliptical, the profile evenly convex to end of snout; eyes large, nearly even in front, the male with the interorbital space deeply concave, its width $\frac{2}{3}$ the vertical depth of the eye (or more in Brazilian specimens); female with interorbital area much narrower, with a more or less perfect median groove, its width about equal to depth of pupil; mouth small, the maxillary reaching to below middle of eye; teeth small, slender, in 2 rows above, in 1 row below, the outer series in upper jaw somewhat enlarged, but hardly canine-like; gill rakers very short and thick, about 1 + 7 in number. Scales small, firm, moderately ctenoid; pectoral $1\frac{1}{3}$ in head in the female, reaching nearly to base of caudal in the male; vertebræ 9 + 24 = 33. Color dark brown, with many rings and spots of light gray and blackish, some of the dark rings with a black central spot; a diffuse dusky blotch on lateral line above pectoral, and 1 near base of caudal peduncle; fins with numerous inky spots and dark markings; blind side pale. West Indian fauna, Key West to Rio Janeiro; rather common. We have found in the Museum of Comparative Zoology specimens purporting to be the types of *Hemirhombus ocellatus*, Poey (No. 11144; Poey's number, 88). These are female examples, and they differ from the types of *Hemirhombus æthalion*, obtained in Cuba by Dr. Jordan, only in their greater size. Numerous

specimens (11373) from Rio Janeiro belong to the same species. Among these are males, which have the interorbital space much broader than in the types of *ocellatus* and *œthalion*. Besides these specimens we have examined others from Hayti, Cuba, and Key West, and there can be no reasonable doubt of their identity, and that all are identical with Günther's *Hemirhombus aramaca*. This fish is described and fairly well figured by Ranzani under the name of *Syacium micrurum*. It is the type of his genus *Syacium*, a generic name which, strangely enough, has received no notice from subsequent authors until lately. (*μικρός*, small; *οὐρά*, tail.)

Syacium micrurum, RANZANI, Nov. Spec. Pisc. Diss. Sec., 20, pl. 5, 1840, Brazil; JORDAN & GOSS, Review Flounders and Soles, 269, 1889.

Hippoglossus ocellatus, POEY, Memorias, II, 314, 1860, Cuba.

Hemirhombus aramaca, GÜNTHER, Cat., IV, 42, 1862, Cuba; Jamaica; not *Rhombus aramaca*, CUVIER.

Citharichthys œthalion, JORDAN, Proc. U. S. Nat. Mus. 1886, 52, Havana. (Type, No. 37748. Coll. D. S. Jordan.)

Hemirhombus ocellatus, POEY, Synopsis, 407, 1868; POEY, Enumeratio, 138, 1875.

Citharichthys ocellatus, JORDAN & GILBERT, Synopsis, 964, 1883; JORDAN, Proc. U. S. Nat. Mus. 1884, 143.

Hemirhombus œthalion, JORDAN, Proc. U. S. Nat. Mus. 1886, 602.

3042. SYACIUM LATIFRONS (Jordan & Gilbert).

Head 4; depth 2½. D. 92; A. 72; scales 60. Body elliptical, the dorsal and ventral outlines equally arched; mouth placed low, below axis of body; snout with an abrupt constriction in front of upper orbit, the outline then again convex; eyes on left side, distant, the lower in advance of the upper; a vertical line from anterior margin of upper orbit passing through middle of lower; distance of upper eye from dorsal outline equaling ⅔ its vertical diameter; interorbital space concave, very wide, its width 1½ times diameter of orbit in a specimen 8 inches long, much narrower in the young; a ridge from upper angle in lower eye runs upward and backward to join a ridge from upper orbit. Nostrils on a level with upper margin of lower eye, the anterior with a flap, distant from the posterior, which is circular; length of snout to front of lower eye 4⅓ to 5 in head; mouth very oblique, the gape convex upward and backward; maxillary ⅔ length of head, reaching to middle of lower pupil, very narrow and covered with small scales; teeth small, the upper jaw with 2 series, the front teeth of the outer series somewhat enlarged; lower jaw with a single series; vomer and palatines toothless; gill rakers short and broad, the longest about ½ vertical diameter of pupil; about 7 on anterior limb of arch; pseudobranchiæ present; preopercle with posterior margin nearly vertical, only the lower third free, the upper ⅔ grown fast to opercle and scaled over; the lower margin running very obliquely downward and forward, the angle thus an obtuse one; dorsal fin commencing on the snout in front of upper eye, the first 4 or 5 rays exserted and turned over to the blind side; the highest rays are behind the middle of the fin and are about ⅔ length of head; anal fin similar to dorsal, its origin under base of pectorals; caudal short, about ⅔ length of head, the middle rays the longest, the outer rays slightly prolonged; ventrals un-

symmetrical, that of colored side on the ridge of the abdomen, the other inserted in front of it; pectoral of colored side long, the rays very slender, the two upper prolonged and filamentous, the upper (in adults) more than $\frac{1}{3}$ total length; pectoral of blind side more than $\frac{2}{3}$ of length of head; scales ciliated, somewhat irregular, of moderate size, with small scales intermixed; snout naked, head and body otherwise scaly; scales on interorbital region very small; a series of small scales on basal half of each dorsal and anal ray; base of caudal thickly scaled, a series of small scales running nearly to tip of each ray, lateral line slightly rising anteriorly, but without distinct curve. Color light brown, with grayish and light bluish dots, some darker areas and a few round brown spots ocellated with lighter; interorbital space with a vertical brown bar bordered by lighter; fins mottled and spotted. This species is known only from the original types, taken by Professor Gilbert at Panama. The several variations in this species have not been studied. The species differs from *Syacium ovale* chiefly in the much broader interorbital space. We should regard this as unquestionably the adult male of *S. ovale* were it not that in making large collections of the latter species at Mazatlan we found not one referable to *S. latifrons*. (*latus*, broad; *frons*, forehead.)

Citharichthys latifrons, JORDAN & GILBERT, Bull. U. S. Fish Comm. 1881, 334, Panama. (Coll. C. H. Gilbert.)

Syacium latifrons, JORDAN & GOSS, Review Flounders and Soles, 271, 1889.

3043. SYACIUM OVALE (Günther).

Head $3\frac{3}{5}$ in length; depth $2\frac{1}{10}$. D. 86; A. 69; scales 58; eye $4\frac{1}{2}$ in head; maxillary $2\frac{3}{5}$; pectoral $1\frac{1}{2}$; caudal $1\frac{1}{6}$. Body elliptic-ovate, body outline from snout to caudal peduncle uniform, the snout not produced; mouth moderate; maxillary concave behind, reaching to middle of pupil of lower eye; lower jaw slightly included; teeth biserial in upper jaw, the inner series small and sharp, the outer much larger, irregular, uniserial in lower jaw; the lower eye slightly in advance of the upper; interorbital space narrow, as broad as pupil, concave; gill rakers as long as pupil, $2 + 8$ in number. Scales strongly ctenoid; scales on mandible, maxillary, and a few in front of interorbital, the middle of which is naked; lateral line not curved. Dorsal beginning slightly in front of upper eye on blind side, the anterior rays produced a little beyond membrane; base of ventral of blind side wider than that of eyed side; caudal double lunate. Color light olive brown, nearly uniform, the vertical fins with elongate dark spots; caudal with large, irregular black spots. Pacific coast of tropical America; common at Mazatlan and Panama. Here described from specimens 6 or 7 inches in length, collected at Mazatlan, Mexico, by the Hopkins expedition to Sinaloa. None of these shows the broad interorbital area of *Syacium latifrons*. (*ovalis*, oval.)

Hemirhombus ovalis, GÜNTHER, Proc. Zool. Soc. Lond. 1864, 154, Panama; GÜNTHER, Fish. Centr. Amer., 472, pl. 80, fig. 1, 1869; JORDAN & GILBERT, Bull. U. S. Fish Comm. 1882, 108–111.

Citharichthys ovalis, JORDAN, Proc. U. S. Nat. Mus. 1885, 391.

Syacium ovale, JORDAN & GOSS, Review Flounders and Soles, 271, 1889.

1049. CYCLOPSETTA, Gill.

Cyclopsetta, GILL, Proc. U. S. Nat. Mus., XI, 1888, 601 (*fimbriata*).

Mouth very large; jaws squarely truncated behind; teeth uniserial, those of the upper jaw moderate, of lower jaw enlarged and largest at sides; dorsal and anal almost symmetrical, dorsal commencing in front of eye on snout, scarcely deflected on blind side; caudal slightly peduncular and convex; pectorals subequal and with a subtruncate free margin; ventrals nearly equal, the left on the preanal ridge, the right lateral, each with the inner ray connected by membrane to the body; interbranchial membrane imperforate; gill rakers tubercular and surmounted by blunt denticles. This genus differs from *Azevia* only in the smooth scales. (*κύκλος*, circle; *ψῆττα*, flounder, from the cycloid scales.)

a. Dorsal rays 91 to 95; anal 73 to 75; scales 90 to 95. Color nearly plain, the fins blotched. QUERNA, 3044.

aa. Dorsal rays 80 to 82; anal 62; dorsal and anal with dark ocelli.

b. Scales small, about 90; pectoral fin uncolored; anterior dorsal rays scarcely produced. CHITTENDENI, 3045.

bb. Scales larger, about 70; pectoral fin with black ocellus; anterior rays of dorsal somewhat produced. FIMBRIATA, 3046.

3044. CYCLOPSETTA QUERNA (Jordan & Bollman).

Head $3\frac{1}{3}$ to $3\frac{2}{5}$; depth $2\frac{1}{3}$. D. 91 to 95; A. 73 to 75; scales along lateral line 90 to 95. Body shaped as in *Azevia panamensis*. Mouth large, maxillary $1\frac{2}{3}$ in head. Teeth as in *A. panamensis*, in single series, rather long and slender, the anterior somewhat more enlarged. Snout 5 in head, its tip hooked over the lower jaw so that the outer canines project. Interorbital space rather narrow, slightly concave, with a few small scales, its width a little less than pupil, $\frac{1}{4}$ diameter of eye. Eyes moderate, $5\frac{1}{2}$ in head, the upper somewhat in advance. Gill rakers short and broad, as in *A. panamensis*, each with 3 or 4 strong teeth. Scales small, cycloid on both sides, those below pectorals more reduced than in *A. panamensis*, about 65 in a cross series; anterior part of lateral line bent slightly upward, this portion about $3\frac{1}{2}$ in straight part. Dorsal beginning above and between the nostrils, the anterior rays short, but with free tips; longest ray $2\frac{1}{2}$ in head; pectoral of eye side $1\frac{1}{3}$ to 2 in head, of blind side $2\frac{1}{3}$ to $2\frac{1}{2}$; ventrals subequal, each 6-rayed, $2\frac{2}{3}$ in head, extending $\frac{1}{3}$ their length beyond vent. Color plain brown, unspotted; fins dusky, thickly punctulate; young with 2 large oval indistinct dark spots on dorsal and anal; 3 on caudal, of which the middle is much larger. Distinguished from *A. panamensis* (Steindachner) by having much smaller cycloid scales on eyed side and by its plain coloration. Coast of Colombia. Numerous specimens, the largest about 8 inches in length, were dredged in 7 fathoms at *Albatross* Station 2800 and in 16 fathoms at Station 2802. (*quernus*, oaken, i. e., tanned.)

Azevia querna, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 174, Pacific Ocean off coast of Colombia, at Albatross Station 2802, Lat. 8° 38′ N., Long. 79° 31′ 30″ W. (Type, No. 41159.)

3045. CYCLOPSETTA CHITTENDENI, B. A. Bean.

Head 3½ in body; depth 2½. D. 82; A. 62; scales 90; eye 5 in head. Mouth widely cleft, oblique, the jaws curved; cleft of mouth less than 2 in head. Teeth of each jaw in a single series, those of lower jaw strong and sharp, curved inward and backward, those of upper jaw not so large, and very irregular in size. Ventral fins well developed, that of eyed side being on abdominal ridge, and about ¾ as long as pectoral; pectorals ½ as long as head, their length equaling a little more than ⅓ of body depth, posterior margin oblique; gill rakers very short, tubercular, almost as broad as long, 3 or 4 + 8 in number. Color brown; fins lighter, marked with blackish; 3 small faint blotches of black on first half of dorsal fin, and 3 rather distinct blotches on second half, last blotch extending to caudal peduncle; anal fin with 3 black blotches situated as and similar to those of dorsal fin; ventral of eyed side blackish, that of blind side pale; caudal fin with 3 black spots at its extremity; pectoral fin of colored side blackish; quite a large blotch of black on body under this fin. This species is distinguished from *Cyclopsetta fimbriata* by its shorter head, smaller and closely adhering scales, larger teeth, the little-produced anterior dorsal rays and by the oblique posterior margin of the pectorals. In *C. fimbriata* the scales are rather large and deciduous, the teeth small, the anterior rays of the dorsal considerably produced, and the posterior margin of the pectoral is subtruncate. A single specimen collected by Dr. John F. Chittenden, of the Victoria Institute, Port of Spain, Trinidad Island, and named in his honor. It is 7¾ inches in length. (B. A. Bean.)

Cyclopsetta chittendeni, B. A. Bean, Proc. U. S. Nat. Mus. 1894, 635, Trinidad. (Type, No. 44100. Coll. Dr. Chittenden.)

3046. CYCLOPSETTA FIMBRIATA (Goode & Bean).

Head 3½ in length; depth nearly 2. D. 80; A. 60 or 61; pectoral 10; ventral 6; scales 25–70–31; maxillary 2 in head; caudal 4⅘ in total length; pectoral 5½. Mouth very large, the upper jaw strongly curved, lower jaw included; teeth uniserial in each jaw, some of the anterior ones in the upper jaw being much larger than those following, while those in the lower jaw are still larger than these, some of the teeth in each jaw depressed; upper eye placed at a distance from profile equal to ½ its own diameter, which is a little less than 5 in head; eyes in the same vertical; interorbital ridge low, 4 in eye; gill rakers very short, tubercular, about 9 on lower part of angle. Scales cycloid; curve of lateral line slight, curve 3⅓ in straight part. Dorsal beginning on snout in advance of nostrils, first ray higher than second, highest rays behind middle of fin; origin of anal under base of pectoral, its highest rays behind middle of fin, higher than highest dorsal rays; ventral of eyed side on ridge of abdomen; middle caudal rays produced. Color grayish brown; dorsal and anal fins each with 2 round dark blotches upon their posterior halves, which are slightly larger than eye; a similar dark blotch upon middle of caudal, sometimes with smaller blotches irregularly placed near its outer margin; pectoral with a very narrow dark band near its base, whole of outer half marked with a dark blotch, reticulated and mottled with

lighter; intervening portion pearly white with dark specks upon the rays; blind side cream colored. Deep waters of the Gulf of Mexico. (Goode & Bean.) (*fimbriatus*, fringed; from the produced dorsal rays.)

Hemirhombus fimbriatus, GOODE & BEAN, Proc. U. S. Nat. Mus. 1885, 591, deep waters of the Gulf of Mexico, between Mississippi Delta and Cedar Keys. (Type, No. 37330. Coll. *Albatross.*)

Arnoglossus ? fimbriatus, JORDAN & GOSS, Review Flounders and Soles, 262, 1889.

Cyclopsetta fimbriata, GOODE & BEAN, Oceanic Ichthyology, 451, fig. 368, 1896.

1050. AZEVIA, Jordan.

Azevia, JORDAN, in JORDAN & GOSS. Review Flounders and Soles, 271, 1889 (*panamensis*).

Body elliptical, compressed, covered with small, firm, ctenoid scales; mouth large; teeth in both jaws uniserial; vomer without teeth; gill rakers very short and thick, tubercle-like; interorbital space very narrow in both sexes, the ridges coalescing between the eyes; lateral line without arch in front; ventrals free from the anal; septum of gill cavity below gill arches, without foramen; a deep emargination near isthmus. None of the fins especially modified or with elongate rays. This genus differs from *Citharichthys* in its tubercular gill rakers, as also in its small, firm scales, and other characters of minor importance. (*Azevia*, a Portuguese name for the sole, used at Lisbon, according to Brito-Capello. It probably corresponds to the Cuban name *Acedía.*)

3047. AZEVIA PANAMENSIS (Steindachner).

Head $3\frac{3}{8}$ in length; depth $2\frac{1}{4}$. D. 95; A. 73 to 78; scales 73 to 78; eye 5 in head; maxillary 2; pectoral $1\frac{3}{8}$; caudal $\frac{1}{4}$. Body rather elongate; anterior profile evenly convex; mouth large, the maxillary reaching to posterior border of eye, the upper jaw somewhat hooked over the lower; about 3 teeth in upper jaw enlarged and hooked, canines in lower jaw long and sharp; eyes about even in head; interorbital space very narrow, less than diameter of pupil, a ridge along its middle; gill rakers divided into many sharp points around its edge, very short, as wide as long, about 4 + 9 in number. Scales on posterior part of interorbital, maxillary, and mandible; tip of snout, the greater part of interorbital, and tip of lower jaw naked; scales all strongly ctenoid; lateral line not curved anteriorly. Origin of dorsal at the vertical between tip of snout and front of eyes, scarcely on blind side, the anterior rays somewhat produced beyond membrane, the fin rather low; origin of anal below angle of opercle; pectorals short, that of eyed side pointed, its mate of the opposite side broadly rounded behind; caudal double lunate. Here described from a specimen collected by the Hopkins Expedition to Sinaloa, at Mazatlan, Mexico, about 11 inches in length. We have also examined specimens from Panama, in the museum at Cambridge, a part of the series of Dr. Steindachner's original types. Pacific coast of Central America; common at Mazatlan and Panama. (*panamensis*, from Panama.)

Citharichthys panamensis, STEINDACHNER, Ichth. Beitr., III, 62. 1875, Panama; JORDAN & GILBERT, Bull. U. S. Fish Comm. 1882, 108 and 111; GILBERT, Bull. U. S. Fish Comm. 1882, 112.

Azevia panamensis, JORDAN & GOSS, Review Flounders and Soles, 272, 1889; JORDAN, Proc. Cal. Ac. Sci. 1895, 503.

1051. CITHARICHTHYS,* Bleeker.

(Whiffs.)

Citharichthys, Bleeker, Comptes Rendus Acad. Sci. Amsterd., xiii, Pleuronectoidei, 6 1862 (*cayennensis* = *spilopterus*).
Orthopsetta, Gill, Proc. Ac. Nat. Sci. Phila. 1862, 330 (*sordidus*).
Metoponops, Gill, Proc. Ac. Nat. Sci. Phila. 1864, 198 (*cooperi* = *sordidus*).

Eyes and color on the left side. Body oblong; mouth of the large type, but comparatively small, with 1 series of small, sharp teeth in each jaw; no teeth on vomer or palatines. Gill rakers moderate, slender. Dorsal fin beginning just in front of eye; all the fin rays simple; ventrals of colored side on the ridge of the abdomen; no anal spine; caudal fin convex or double truncate behind; none of the fins produced. Scales thin, deciduous, slightly ctenoid. Lateral line nearly straight, simple. Lower pharyngeals separate, each with a single row of teeth. Vertebræ 30 to 40. This genus includes small flounders of weak organization, especially characteristic of the sandy shores of tropical America. The subgenus *Orthopsetta* includes species of more northern range and somewhat different form, and especially noteworthy as having an increased number of vertebræ. The two groups intergrade so perfectly that no sharp line of division can be drawn between them. (*Citharus*, an allied genus; ἰχθύς, fish—a fish which lies on its κίθαρος, or ribs; that is, on its side.)

Orthopsetta (ὀρθός, straight; ψῆττα, flounder):

a. Vertebræ 37 to 40; interorbital ridge sharply elevated; the head not closely compressed; eyes large; species of the North Pacific.
b. Interocular space concave, scaly, at least behind.
c. Gill rakers $x + 16$ to 18.
d. Scales 65 to 70; dorsal rays 95; anal 77; depth $2\frac{1}{2}$. SORDIDUS, 3048.
dd. Scales 46 to 50; dorsal rays 83 to 87; anal 67 to 70; depth $2\frac{2}{3}$ in length. FRAGILIS, 3049.
cc. Gill rakers $x + 10$ or 11; dorsal rays about 84; anal 65; scales 50; depth $2\frac{1}{4}$ in length. XANTHOSTIGMUS, 3050.
bb. Interocular space a sharp, naked ridge; dorsal rays 85 to 90; anal 68 to 72; scales 55 to 60; head $3\frac{1}{4}$ in length; depth $2\frac{1}{3}$. STIGMÆUS, 3051.

Citharichthys:

aa. Vertebræ 33 to 36; interorbital ridge low and narrow, the head closely compressed. Species of the Atlantic or the Tropics.
e. Eyes large, 3 to $4\frac{1}{2}$ in head.
f. Head large, 3 to $3\frac{1}{2}$ in length.
g. Interorbital space very narrow, 5 in eye; snout with a spine; pectoral of eyed side elongate, $\frac{1}{3}$ longer than head; maxillary $2\frac{1}{4}$ in head. D. 91; A. 73; scales 48. DINOCEROS, 3052.
gg. Interorbital space very broad, 2 in eye; snout without spine; pectoral of eyed side shorter than head; maxillary $2\frac{1}{2}$ in head. D. 78; A. 62; scales 43. PLATOPHRYS, 3053.
ff. Head smaller, about 4 in length.
h. Body comparatively elongate, the depth about $2\frac{1}{3}$ in length; mouth very small; the maxillary $3\frac{1}{2}$ in head; dorsal rays 83; anal 67; scales 40; eye 4 in head. ARCTIFRONS, 3054.

* "As the name *Citharichthys* was introduced a short time before that of *Orthopsetta*, proposed for the *Psettichthys sordidus*, and was framed for a species related to that type, that name must be adopted if the *O. sordida* is not regarded as generically distinct." (Gill.)

hh. Body comparatively broad, the depth about ½ the length; mouth larger.

i. Snout with a strong, sharp spine on eyed side, above upper lip; eyes large, 3 in head; greatest depth of body over the pectorals; interorbital space with a wide ridge, about ½ diameter of eye. D. 74; A. 60; scales 40. UNICORNIS, 3055.

ii. Snout without distinct spine; eyes moderate, 3½ to 4½ in head; greatest depth of body under middle of dorsal; interorbital space a narrow, scaly ridge with a slight median groove; maxillary 2½ in head; teeth small, those in front slightly enlarged; body not very thin; gill rakers moderate, 6 + 13.

j. Dorsal rays 68; anal 52; scales smaller, the lateral line with about 53 pores; sides with whitish blotches. UHLERI, 3056.

jj. Dorsal rays 80; anal 56; scales large, 41 in lateral line; sides and fins with dark blotches. MACROPS, 3057.

ee. Eyes quite small, 5 to 6 in head; snout short, forming an angle with the profile; mouth moderate, oblique, the maxillary 2⅓ to 2⅔ in head; teeth small, the anterior somewhat enlarged; dorsal rays about 80; anal rays 60; body and fins speckled.

k. Scales not very large, 45 to 48 in lateral line; gill rakers long and slender, longer than pupil. SPILOPTERUS, 3058.

kk. Scales large, 40 to 46 in lateral line; gill rakers short, not longer than pupil. GILBERTI, 3059.

Subgenus ORTHOPSETTA, Gill.

3048. CITHARICHTHYS SORDIDUS (Girard).

(SOFT FLOUNDER.)

Head 3⅔; depth 2⅓. D. 95; A. 77; scales 65 to 70. Form elliptical; interocular space concave, scaly, a conspicuous sharp ridge above the lower eye; mouth not large, the maxillary about 3 in length of head; teeth anteriorly subequal, growing much smaller behind. Gill rakers about 7 + 16. Lower pharyngeals narrow, each with 1 row of slender teeth. Scales rather large, thin, and membranaceous, readily deciduous, their edges slightly ciliate; accessory scales numerous. Eye large, much longer than snout, 3½ in head; depth of caudal peduncle less than ⅛ head; pectorals long, nearly ⅔ length of head. Vertebræ 11 + 29 = 40. Dull olive brownish of varying shade, the males with dull orange spots and blotches; each scale with a darker edge; dorsal and anal fins in the male blackish, with dull orange blotches, and edged anteriorly with yellowish; female paler, the fins nearly plain. Pacific coast of North America, in water of moderate depth; British Columbia to Lower California. This small flounder is one of the commonest species on the Pacific coast, being found in water of 10 fathoms or more depth in all localities from the Mexican boundary to British Columbia. Although much larger in size than any other species of the genus, it rarely exceeds 2 pounds in weight. In its deciduous scales and soft flesh it much resembles *Lyopsetta exilis* and *Atheresthes stomias*, 2 species which are often taken in company with it. Of all the species of *Citharichthys*, this one has the most extended range to the northward. (*sordidus*, sordid, from its dull coloration.)

Psettichthys sordidus, GIRARD, Proc. Ac. Nat. Sci. Phila., VII, 1854, 142, San Francisco; Tomales Bay; GIRARD, U. S. Pac. R. R. Surv., X, Fishes, 155, 1858.

Metoponops cooperi, GILL, Proc. Ac. Nat. Sci. Phila. 1864, 198, Santa Barbara; shrivelled specimen. (Type, No. 9407.)

Orthopsetta sordida, GILL, Proc. Ac. Nat. Sci. Phila. 1862, 330.

Citharichthys sordidus, LOCKINGTON, Rep. Com. Fisheries of California, 1878-79, 42; LOCKINGTON, Proc. U. S. Nat. Mus. 1879, 83; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1880, 453; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 67; JORDAN & GILBERT, Synopsis, 817, 1883; BEAN, Proc. U. S. Nat. Mus. 1883, 353; JORDAN & GOSS, Review Flounders and Soles, 274, 1889.

3049. CITHARICHTHYS FRAGILIS, Gilbert.

Head $3\frac{2}{5}$ to $3\frac{3}{5}$ in length; depth $2\frac{3}{5}$ to $2\frac{3}{4}$ (in specimens 5 inches long.) D. 83 to 87; A. 67 to 70; scales 46 to 50. Vertebræ 10 + 27. Body elongate, posteriorly sharply wedge-shaped, tapering to base of caudal; anterior profile very conspicuously angulated above front of upper eye, the snout strongly projecting, its anterior profile nearly vertical; depth of caudal peduncle $2\frac{1}{5}$ in head. Anterior nostril with a short tube and flap, the latter nearly obsolete on blind side. Mouth more oblique than in *C. sordidus;* maxillary reaching vertical from front of pupil, $2\frac{2}{3}$ to $2\frac{3}{5}$ in head. Teeth in a single series, close set, those anteriorily somewhat enlarged, but none of them canine-like. Eyes large, the vertical from front margin of upper eye falling through front of lower pupil; longest diameter of upper orbit $2\frac{2}{3}$ in head; interorbital space narrow, concave, scaled, the lower ridge strongest, its width about $\frac{1}{4}$ diameter of orbit. Symphyseal knob sharp. Gill rakers long, slender, close set (as in *C. sordidus*), 18 on anterior limb of arch, the longest $\frac{1}{3}$ orbit. Scales large, deciduous, somewhat irregularly arranged, 12 or 13 series above lateral line; scales smooth on blind side, minutely spinous on eyed side; lateral line without anterior arch. Dorsal beginning slightly in advance of eye, the longest ray $\frac{1}{2}$ head; pectorals long and narrow, with 11 rays on colored side. Color dusky olivaceous, with occasional slaty-blue spots. This species is closely related to *C. sordidus*, from which it differs in the fewer vertebræ and fin rays and in the larger scales. Many specimens from the Gulf of California in from 18 to 76 fathoms, at *Albatross* Stations 3011, 3016 to 3018, and 3033. (Gilbert.) (*fragilis*, fragile.)

Citharichthys fragilis, GILBERT, Proc. U. S. Nat. Mus. 1890, 120, Gulf of California, east coast of Lower California. (Type, No. 44409. Coll. Dr. Gilbert.)

3050. CITHARICHTHYS XANTHOSTIGMUS, Gilbert.

D. 81 to 86; A. 63 to 67; scales 50. Vertebræ 11 + 26. Body deep, varying from $2\frac{1}{3}$ (in young, 3 inches long) to $2\frac{1}{10}$ (7 inches long) in length of body. Profile angulated above front of upper eye, the snout convexly projecting. Depth of caudal peduncle $\frac{1}{3}$ head. Lower eye in advance, the vertical from front of the upper passing through front of lower pupil. Mouth rather small, the outline somewhat curved, the maxillary reaching the vertical from front of lower pupil, 3 to $3\frac{1}{4}$ in head; mandible with a sharp downward-directed point at symphysis. Teeth in a single close-set series in each jaw, growing slightly larger anteriorly, but without canines.

Anterior nostril with a short tube, and a narrow flap arising from its inner edge. Interorbital width 3¾ to 5 in orbit, slightly concave, the lower ridge much stronger and higher than the upper, scaled posteriorly. Eye large, the orbit 3⅛ to 3⅝ in head. Gill rakers rather long and slender, coarsely dentate on inner margin, distant, 10 or 11 on anterior limb of arch. Scales large, in regular series, appearing cycloid, but the edges very minutely spinous; lateral line gently rising on anterior ⅓, but without curve; fifty vertical series of scales, with as many pores in lateral line; 13 to 15 horizontal series above lateral line. Dorsal beginning immediately behind posterior nostril of blind side, ending so as to leave caudal peduncle free for a distance equaling ½ diameter of eye; ventrals long, reaching beyond origin of anal; pectoral very long and slender, normally with 9 rays, the longest ray on colored side longer than head, about ⅓ length of body. Color light olive brown, irregularly flecked with slaty, and with numerous bright yellow spots broadly ocellated with brownish black, a series of these usually on lateral line, and 2 others halfway between it and the dorsal and ventral outlines, respectively, those of the latter series forming pairs; fins not conspicuously marked, the pectorals sometimes with faint broad dusky cross bars. Both coasts of Lower California. In external appearance the species closely resembles *C. sordidus*, to which, however, it is not closely related, differing in number of scales, fin rays, and vertebræ, and in the size and number of gill rakers. (Gilbert.) Many specimens, from *Albatross* Stations 3039, 3043, and 3044, in 47 to 74 fathoms. (ξανθός, yellow; στίγμα, spot.)

Citharichthys xanthostigma, GILBERT, Proc. U. S. Nat. Mus. 1890, 120, Gulf of California, west coast of Lower California, and Magdalena Bay. (Type, No. 44408. Coll. Dr. Gilbert.)

3051. CITHARICHTHYS STIGMÆUS, Jordan & Gilbert.

Head 3¾ in length without caudal; depth 2⅓; dorsal 87; anal 68; scales 54 (pores). Body moderately deep, the 2 profiles regularly and equally arched; snout short, gibbous, projecting a little beyond the outline; caudal peduncle very short, not high, its length (from end of last vertebra to vertical from last anal ray) about ⅔ its height, which is ⅔ length of head; caudal fin appearing sessile. Mouth moderate, very oblique, the maxillary reaching beyond front of pupil, 2¾ in head; teeth in a single series, subequal in the two jaws, rather long, very slender and numerous, decreasing toward angle of mouth; about 40 teeth in the upper jaw, and 30 in the lower on blind side. Eyes large, close together, separated by a narrow, sharp, scaleless ridge; the upper eye largest, slightly behind the lower, with considerable vertical range; diameter of upper eye 3⅛ in head. Snout and lower jaw scaleless; end of maxillary and rest of head scaled. Gill rakers moderate, not strong, about 9 on anterior limb. Dorsal fin beginning on the vertical from front of upper eye, the first 3 rays being somewhat turned to blind side; the fin low, the highest at beginning of its posterior third, the longest ray nearly ½ length of head; anal spine present, very small; caudal rounded, about equalling length of head; pectoral of colored side 1⅔ in head, of blind side, 2⅕. Scales

moderate, those forming the lateral line persistent, the others deciduous, those on colored side with ciliated margins, on blind side smooth; lateral line without anterior curve, the scales crowded and smaller anteriorly. Color in spirits uniform olivaceous, the scales dark edged; lips and some of the membrane bones of head margined with blackish; fins dusky, each seventh (to tenth) ray of vertical fins with a very small but conspicuous black spot on its middle. The above description is from the original type from Santa Barbara. Numerous specimens dredged by the *Albatross* in 9 to 41 fathoms off the coast of California show the following characters: Gill rakers $x+9$. Specimens 5 mm. long show white spots each with a black half ring on the outer side symmetrically arranged along bases of dorsal and anal; 4 distinct pairs of these, 2 unpaired ones more anteriorly along dorsal base, and a few fainter ones midway between these rows and the lateral line and alternating with them; there are some other scattered light spots. The abdomen is covered by a broad black streak; this, however, is wanting in specimens larger and smaller. Coast of California; rare; in rather deep water. The original type of this species is a young example, taken near Santa Barbara by Capt. Andrea Larco. In the Museum of Comparative Zoology are other specimens collected by Mr. Cary at San Francisco. These have 72 anal rays, while the original type had but 68. A few other specimens have been since obtained. Some of these are full of spawn at a length of 5 inches. (*στιγμαῖος*, speckled.)

Citharichthys stigmæus, Jordan & Gilbert, Proc. U. S. Nat. Mus. 1882, 410, 411, Santa Barbara (Coll. A. Larco. Type, 31099 U. S. Nat. Mus.); Jordan & Gilbert, Synopsis, 965, 1883; Jordan & Goss, Review Flounders and Soles, 274, 1889; Gilbert, Rept. U. S. Fish. Comm. 1893 (1896) 473.

Subgenus CITHARICHTHYS.

3052. CITHARICHTHYS DINOCEROS, Goode & Bean.

Head $3\frac{1}{3}$ in length; depth $2\frac{1}{3}$. D. 91; A. 73; scales 14–48–16; eye $3\frac{1}{2}$ in head; maxillary a little less than 2; greatest height of dorsal 2; pectoral $2\frac{1}{3}$ in body; caudal $5\frac{1}{2}$. Teeth uniserial in both jaws, those in the front much the largest; a strong spine upon the snout overhanging the upper lip, above this a second shorter spine; interorbital very narrow, its width less than 5 in eye, ridge rather prominent, narrow, sharp. Scales thin, deciduous, cycloid, large; lateral line slightly curved over the pectoral. Dorsal beginning on snout, in advance of eye, upon the blind side, its highest rays behind the middle; origin of anal under base of pectoral; third and fourth pectoral rays upon the eyed side elongate, the fin $\frac{3}{5}$ longer than its mate of the opposite side; caudal subsessile, pointed. Color grayish brown above, white below. Vertebræ 33 to 36. West Indies, in deep water. The type specimen, 92 mm. long to base of caudal, was taken by the *Blake*, off Guadeloupe; others were taken off St. Lucie and Barbados, from 310 to 955 fathoms. (Goode & Bean.) (*δεινός*, terrible; *κέρας*, horn.)

Citharichthys dinoceros, Goode & Bean, Bull. Mus. Comp. Zool., xii, No. 5, 157, 1886, off Martinique, St. Lucie, and Barbados; Jordan & Goss, Review Flounders and Soles, 275, 1889; Goode & Bean, Oceanic Ichthyology, 447, 1896.

3053. CITHARICHTHYS PLATOPHRYS, Gilbert.

Head 3; depth 2. D. 78; A. 62; scales 43. Body ovate; caudal fin subsessile, the free portion of caudal peduncle about ½ as long as diameter of pupil, its depth ⅛ length of head. Mouth very oblique; maxillary 2½ in head, reaching vertical from middle of lower eye. Teeth slender, close set, in a single series in each jaw, those in front of upper jaw largest, but not canine-like. Eyes large, the lower much in advance of the upper, their horizontal diameter 3¾ in head. Interorbital space very wide for the genus, concave, divided by an oblique ridge running backward from middle of upper orbit; interorbital width 8⅔ in head, nearly ½ as wide as eye. Distance from tip of snout to front of lower eye ⅔ diameter of eye, from tip of snout to upper eye ⅓ head. Gill rakers short and very slender, less than diameter of pupil, 9 present on horizontal limb of outer arch. Scales large, those on blind side very weakly ctenoid. Dorsal beginning behind nostril on blind side of snout, its longest ray 2⅓ in head; pectoral of eyed side long and narrow, 4 in length, containing 11 rays, that of blind side but ½ its length; ventrals short; caudal rounded, 1⅔ in head. Color in spirits, uniform light brownish (olivaceous in life), without distinctive marks; fins somewhat dusky; ventral of eyed side jet-black, that of blind side blackish on distal portion of inner rays. (Gilbert.) One specimen known, from *Albatross* Station 2799, southwest of Panama. (*πλατύς*, broad; *ὀφρύς*, eyebrow.)

Citharichthys platophrys, Gilbert, Proc. U. S. Nat. Mus. 1890, 454, Albatross Station 2799, southwest of Panama. (Coll. *Albatross*.)

3054. CITHARICHTHYS ARCTIFRONS, Goode.

Head 4 in body; depth 2⅔. D. 82; A. 67; pectorals 9 or 10, 7; scales 8–40–8; eye 4 in head; maxillary 3½; caudal 1. Body comparatively elongate; mouth small; teeth small, the anterior scarcely enlarged; interorbital space narrow, sharp, scaleless; scales cycloid, deciduous; small scales on the rays of the ventral fins; lateral line sharply defined, straight. Dorsal beginning above front of upper eye, its highest ray about 3 times the distance from snout to first ray; origin of anal under base of pectoral; caudal subsessile, triangular; rays of vertical fins all exserted; pectoral inserted low, that of eyed side twice the length of the other. Color dirty light brown. Deep waters of the Gulf Stream. (*arctus*, contracted; *frons*, forehead.)

Citharichthys arctifrons, Goode, Proc. U. S. Nat. Mus. 1880, 341, 472, Gulf Stream off southern coast of New England; Goode & Bean, Bull. Mus. Comp. Zool., Vol. x, No. 5, xix, 194, 1883; Jordan & Gilbert, Synopsis, 818, 1883; Jordan & Goss, Review Flounders and Soles, 275, 1889; Goode & Bean, Oceanic Ichthyology, 442, fig. 366, 1896.

3055. CITHARICHTHYS UNICORNIS, Goode.

Head 4 in length; depth a little less than length. D. 74; A. 60; P. 4 (right), 10 (left); scales 12–40–12; eye 3 in head; maxillary scarcely 2; highest dorsal ray 2. Body deep, its greatest height over the pectorals;

scales thin, deciduous; eye equal to snout or interorbital space; interorbital with a strong ridge; teeth minute, close set, in a single series, stronger on the blind side; a strong, sharp spine on the snout at the anterior termination of the ridge at lower margin of upper eye; caudal pointed, triangular, subsessile; pectoral of left side twice as long as the eye, not ½ longer than right pectoral. Dorsal beginning at side of preorbital spine, its anterior rays being slightly upon the blind side; anal equal to dorsal in height. Ashy gray, with dark lateral line; eyes black. (Goode.) Deep waters of the Gulf Stream. (*unicornis*, having one horn.)

Citharichthys unicornis, GOODE, Proc. U. S. Nat. Mus. 1880, 342, Gulf Stream off southeast of New England; JORDAN & GILBERT, Synopsis, 818, 1883; JORDAN & GOSS, Review Flounders and Soles, 275, 1889; GOODE & BEAN, Oceanic Ichthyology, 444, fig. 369, A & B, 1896.

3056. CITHARICHTHYS UHLERI, Jordan.

D. 68; A. 52; scales 53 (pores). Body comparatively broad, regularly oval, without angle; greatest depth of body under middle of dorsal; eyes moderate, 4½ in head, close together, the orbital ridges coalescent, the lower larger. Teeth small, uniserial; maxillary 2⅕ in head; gill rakers short and very slender, $x + 12$. Color dark brown, with whitish blotches, the fins mottled. Hayti. A single specimen in the Museum of Comparative Zoology, 4½ inches in length. The species is close to *Citharichthys macrops*, but its fin rays and scales are considerably more numerous than in the latter. (Named for Mr. Philip Reese Uhler, the well-known entomologist, its discoverer.)

Citharichthys uhleri, JORDAN & GOSS, Review Flounders and Soles, 275, 1889, Hayti. (Coll. P. R. Uhler. Type in Mus. Comp. Zool.)

3057. CITHARICTHYS MACROPS, Dresel.

Head 4 in body; depth scarcely 2. D. 80; A. 56; scales 14–41–16; lower eye 4 in head; maxillary 2⅓; highest dorsal rays a little over 2; pectoral of eyed side 1⅔; caudal 4 in body; vertebræ $9 + 25 = 34$. Body suboval; upper profile very convex, descending in a sharp curve from nape to front of upper eye, and forming an abrupt angle with the short, blunt snout; mouth moderate, very oblique and curved; maxillary reaching to below middle of eye; teeth minute, uniserial, slightly larger on blind side; interorbital narrow, with a scaleless ridge, which curves upward and backward to upper angle of gill opening; upper eye very close to profile, its anterior margin on the same vertical line with lower; snout shorter than eye; gill rakers about ½ the length of eye, $6 + 13$ in number. Scales large, not ciliated, no accessory scales; origin of dorsal on blind side near tip of snout, anterior rays exserted, the first ray as long as eye, the fin highest at its middle portion; origin of anal under base of pectoral, its highest part a little higher than dorsal; caudal pointed; pectoral of blind side somewhat shorter than that of eyed side. Color in spirits, light-olive brown; body with some 20 dark-brown spots, the largest as large as eye, 4 of these arranged at equal intervals along the lateral line, the second near the middle the most prominent; dorsal and anal fins with a series of round, brown spots, 1 at the middle of every sixth or seventh ray, besides small irregular spots and mottlings;

caudal spotted and mottled with dark brown, and with 2 round, brown spots, 1 above the other at the base of the fin. (Dresel.) South Atlantic and Gulf coasts of the United States; rather common; a well-marked species. We have examined several specimens dredged in the harbor of Beaufort, N. C., by Prof. Oliver P. Jenkins. (*μακρός*, large; *ὤψ*, eye.)

Citharichthys macrops, DRESEL, Proc. U. S. Nat. Mus. 1884, 539; Pensacola (Type No. 21500); JORDAN, Proc. U. S. Nat. Mus. 1886, 29; JORDAN & GOSS, Review Flounders and Soles, 275, 1889.

3058. CITHARICHTHYS SPILOPTERUS, Günther.

Head 3½ in body; depth 2⅕. D. 75 to 80; A. 58 to 61; scales 45 to 48; eye 6 in head; maxillary 2½; pectoral 2⅓; highest dorsal and anal rays 2; caudal 1½. Body moderately elongate, much compressed; snout short, forming an angle with the profile; jaws strongly curved, the upper somewhat hooked over the lower; lower jaw slightly included; maxillary reaching to posterior margin of lower orbit; teeth small, in a single row, the anterior a little enlarged; interorbital area a low, narrow ridge, which is divided only anteriorly; gill rakers short and rather slender, about 3 in eye, 4 + 12 in number; scales cycloid. Origin of dorsal above anterior edge of upper eye, very slightly on blind side, its highest rays in its posterior half; origin of anal slightly behind base of pectoral; pectoral of eyed side very slightly shorter than that of eyed side; vertebræ 34. Color olive brownish, somewhat translucent, with darker dots and blotches; a series of distant obscure blotches along bases of dorsal and anal. Atlantic coast of tropical America north to New Jersey; very common on sandy shores; not found in the Pacific, all west coast references belonging to *C. gilberti*. Here described from a specimen from Havana, 6 inches in length. This little flounder is almost everywhere abundant on the sandy shores of the warmer parts of the Western Atlantic, in shallow water. Careful comparison of specimens from South Carolina, Cuba, and Brazil shows no tangible difference, and we are compelled to regard all as forming a single species. It rarely exceeds 5 or 6 inches in length. It usually comes into the markets mixed with other shore fishes, and it nowhere receives any notice as a food-fish. This species is common in the markets of Havana, and it is evidently the original of Poey's *Hemirhombus fuscus*, although in Poey's description there seems to be some confusion, because the teeth are said to be biserial above, and 60 scales are counted in the lateral line. A specimen from Poey in the museum at Cambridge is labeled "*Hemirhombus fuscus*, type. Collector's number, 87." This belongs to *C. spilopterus*, and it has 48 scales in the lateral line. Bleeker's *C. guatemalensis* agrees in all respects with *C. spilopterus*. We are unable to find any description of *C. cayennensis*, if, indeed, the species has ever been described. Specimens of *C. spilopterus* are in the museum at Cambridge from Cuba, Pará, Sambaia, Pernambuco, Camaru, Rio das Velhas, Rio Janeiro, and San Matheo. (*σπίλος*, spot; *πτερόν*, fin.)

Citharichthys spilopterus, GÜNTHER, Cat., IV, 421, 1862, New Orleans, San Domingo, Jamaica; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 618; JORDAN & GILBERT, Synopsis, 817, 1883; JORDAN, Proc. U. S. Nat. Mus. 1886, 53; JORDAN & GOSS, Review Flounders and Soles, 276, 1889.

Citharichthys cayennensis, BLEEKER, Comptes Rendus Acad. Sci. Amsterd., XIII, 1862, 6, Cayenne; name only.

Citharichthys guatemalensis, BLEEKER, Neder. Tydschr. Dierk. 1864, 73, Guatemala; GÜNTHER, Fish. Centr. Amer., 472, 1869.

Hemirhombus fuscus, POEY, Synopsis, 406, 1868, Havana; POEY, Enumeratio, 138, 1875.

3059. CITHARICHTHYS GILBERTI, Jenkins & Evermann.

Head $3\frac{1}{2}$ to $3\frac{2}{3}$; depth of head 4; depth of body $1\frac{9}{10}$ to $2\frac{1}{6}$. D. 77 to 82; A. 57 to 61; scales 18–40 to 46–19. Body comparatively broad, formed as in *C. spilopterus*, the two profiles about equally arched; snout slightly longer than longest diameter of eye, and without a distinct spine. Eyes on left side, equal in size, small, 5 to $5\frac{2}{3}$ in head; interorbital space narrow, $1\frac{2}{3}$ in eye, low, slightly grooved, and scaled on posterior portion only. Maxillary $2\frac{2}{3}$ in head, reaching barely to posterior border of eye; upper jaw projecting. Teeth small, in a single series; gill rakers 4 + 13, short and slender, not longer than pupil, with a rather broad base, narrowing to a slender stalk. Dorsal fin beginning in front of upper eye, the first 3 rays growing from the blind side, the distance of origin from snout 7 in head; fin rays all simple, $2\frac{3}{10}$ in head; pectorals nearly equal, the one on colored side being slightly longer, $1\frac{9}{10}$ in head; rays on colored side 9; on blind side 8; ventrals $2\frac{3}{10}$ in head; caudal rounded, caudal peduncle short, its depth 8 in the body, equaling height of anal; scales large, ciliated, pretty uniform, those toward head and margins of disk becoming smaller; lateral line gradually descending along the course of about 16 scales, from which point it is straight. Color light brown, with about 15 irregular dark blotches of various sizes, the largest being a pair on the latter third of the disk, 1 on each side of lateral line, as great in diameter as length of ventral fin. Specimens from fresh waters (*C. sumichrasti*) are much darker in color; gray, everywhere closely peppered with dark specks; pectoral and caudal mottled. Pacific coast of tropical America; very abundant in sandy bays from Guaymas to Panama, ascending all the streams. This species very closely resembles *C. spilopterus*, representing the latter on the Pacific coast, and it has been frequently recorded under the name *C. spilopterus*. *C. gilberti* differs mainly in the shorter gill rakers and in the slightly larger scales. Fresh-water specimens (as the type of *C. sumichrasti* from Rio Zanatenco, Chiapas, and numerous examples collected by us in Rio Presidio, near Mazatlan) differ considerably in color, being much darker, but there is no other difference. ("This species is dedicated to Prof. Charles H. Gilbert, whose collection and notes on fishes from Mazatlan, containing undescribed species, this among them, were destroyed by fire in 1883.")

Citharichthys gilberti, JENKINS & EVERMANN, Proc. U. S. Nat. Mus. 1888, 157, Guaymas, Mexico (Type, No. 39627. Coll. Jenkins & Evermann); JORDAN, Proc. Cal. Ac. Sci. 1895, 503.

Citharichthys sumichrasti, JORDAN & GOSS, Review Flounders and Soles, 276, 1889, Rio Zanatenco, Chiapas. (Coll. Prof. Francis E. Sumichrast. Type, 25299, M. C. Z.)

Citharichthys spilopterus, GÜNTHER, Fish. Centr. Amer., 471, pl. 80, fig. 2, 1869; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 382; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 630; JORDAN & GILBERT, Bull. U. S. Fish Comm. 1882, 108–111; not of GÜNTHER, 1862.

1052. ETROPUS, Jordan & Gilbert.

Etropus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 364 (*crossotus*).

Eyes and color on left side. Body regularly oval, deep, and compressed. Head small; mouth very small, the teeth close set, slender, and pointed, somewhat incurved, mostly on the blind side; no teeth on vomer. Eyes small, separated by a narrow, scaleless ridge; margin of preopercle free. Ventrals free from anal, that of colored side inserted on ridge of abdomen, its base rather long. Dorsal fin beginning above eye; caudal double truncate; anal without spine. Scales thin, deciduous, ctenoid on left side, cycloid on blind side. Lateral line simple, nearly straight. Size small. This genus is very close to *Citharichthys*, from which it differs only in the very small size of the mouth, and in the correspondingly weak dentition. The 3 or 4 known species are similar in appearance to the species of *Citharichthys*, and they inhabit the same waters. Another genus extremely close to *Etropus* and *Citharichthys* is *Thysanopsetta*, a South American genus. The teeth in *Thysanopsetta* are, however, arranged in a band. The larval forms are translucent and symmetrical, as in *Platophrys*, *Monolene*, *Arnoglossus*, etc. (ἦτρον, abdomen; πούς, foot; in allusion to the insertion of the ventrals, common to all the *Psettinæ*, but not found in other small-mouthed species.)

a. Snout not acute; dorsal rays 75 to 85.

b. Body comparatively elongate, the depth rather less than ¼ length.

c. Dorsal rays 81; anal 58; head 4¼ in length; eye 3⅔ in head; maxillary 4. MICROSTOMUS, 3060.

cc. Dorsal rays 77; anal 61; head 4 in length; eye 3½ in head; maxillary 4⅓. RIMOSUS, 3061.

bb. Body very deep, the depth more than ⅓ length; eye 3⅔ in head; maxillary 4; head 4½; depth 1⅞ to 2; D. 76 to 85; A. 56 to 67; scales 42 to 48; cirri on subopercle of blind side very numerous, white; olive ground, with darker blotches; fins sanded. CROSSOTUS, 3062.

3060. ETROPUS MICROSTOMUS (Gill).

"D. 81; A. 58; caudal 4, 6, 5, 3; pectoral 10; ventral 6. The height of the body enters about 2⅜ times (0.36–0.37) in the extreme length; that of the caudal peduncle about 11 times. The head forms a fifth of the length, is rather abbreviated, scarcely sinuous above the eyes, blunt at the snout, which scarcely exceeds ⅓ of the head's length and the rostral area is rhombic, and not higher than long. The eyes are even; the longitudinal diameter contained about 3⅔ times (0.05½) in the head's length. The mouth is rather small, the length of the upper only equaling ¼ of the length, and that of the lower ⅖ of the head's length. The teeth are very small, and close together; larger in front. The dorsal commences above the front of the orbit, and is highest and convergent near the fortieth ray, which equals about $\frac{1}{10}$ of the total length; the anal is highest at about the twenty-fifth ray, and is as high or even higher than the dorsal. The caudal is rounded behind, and forms about ⅙ of the length. The pectoral fins are unequally developed, that of the dark side being prolonged, and contained only 6⅔ times in the total length, while that of the white side only equals $\frac{1}{10}$ of the same; the rays are all simple.

The ventral fins are also unequally developed, the right being on the abdominal ridge at its origin, rather in advance of the opercular margin and with its longest rays contained about 14 times in the total length; stretched backward, it extends to the second anal ray; the fin on the white side is more advanced, wider, and its rays longer, contained less than 12 times in the length and extends backward to nearly the third anal ray. The scales are large, angular behind, covered with smaller ones, especially near the point of junction of contiguous ones, where alone they are developed on the blind side; the scales of the eyed side are mostly minutely ciliated behind, unarmed, however, near the lateral line, the scales of which last are quadrate and mostly covered; the scales of the blind side are less angular behind and unarmed. The lateral line runs through about 42 scales, while of longitudinal rows there are 10 above and 14 below the lateral line. The color is uniform reddish brown. A single specimen, little more than 3 inches long, was first obtained by Professor Baird, at Beesleys Point." (Gill.)

This species has not been certainly recognized by recent writers, unless, as supposed by Jordan & Goss, it is identical with *Etropus rimosus*. It is in any event certainly an *Etropus*. In the Museum of Comparative Zoology are numerous young specimens collected at Somers Point, New Jersey, by Dr. Stimpson. These seem to belong to the genus *Etropus*. The teeth are equal; the scales are 44, and the depth of the body is $2\frac{1}{5}$ in its length. The eye is 4 in head, the dorsal rays 75 to 80, and the anal rays 56 or 57. The color is light brown, mottled and spotted with darker. These probably represent the *Citharichthys microstomus* of Gill, collected in the same neighborhood by the same naturalist. We are unable to distinguish them from *Etropus rimosus*. (*μικρός*, small; *στόμα*, mouth.)

Citharichthys microstomus, Gill, Proc. Ac. Nat. Sci. Phila. 1864, 223. Beesleys Point, New Jersey (Coll. Prof. S. F. Baird); Goode & Bean, Oceanic Ichthyology, 446, 1896; Jordan, Proc. U. S. Nat. Mus. 1890, 332.

Etropus microstomus, Jordan & Goss, Review Flounders and Soles, 278, 1889.

3061. ETROPUS RIMOSUS, Goode & Bean.

Head 4 in body; depth 2 to $2\frac{1}{4}$. D. 77; A. 61; scales 12–41–14; eye $3\frac{1}{2}$ in head; maxillary $4\frac{1}{3}$; snout 8; caudal 1. Body somewhat elongate, pear-shaped; mouth very small, its cleft less than $\frac{1}{2}$ the orbit, its angle below anterior margin of lower eye; teeth well developed on blind side on each jaw, also on eyed side of lower jaw in front; eyes placed in the same vertical; upper eye close to dorsal profile, and separated from its mate by a space less than $\frac{1}{8}$ its diameter; interorbital ridge low; nostrils in a line with the interorbital ridge, each in a short tube, the posterior the larger, the anterior midway between tip of snout and front of lower orbit; head entirely scaled; accessory scales numerous. Dorsal commencing on blind side at anterior margin of eye, the highest rays somewhat behind middle of fin, its length 7 times in total length; origin of anal under base of pectoral, its highest rays equal to those of dorsal; pectoral of eyed side longest, equal to head without snout; caudal fin rounded. Color gray, hoary above, with a few irregularly placed indistinct brownish blotches, none of them larger than eye; white below. West coast of Florida;

type, 100 mm. long, collected by the *Albatross* at Station 2408, depth 21 fathoms, between Pensacola and Cedar Keys, Florida. (Goode & Bean.) On reexamining our specimens of *Etropus*, we find that those obtained by Jordan & Evermann from Pensacola differ from the others in the greater elongation of the body and in the somewhat grayer coloration. These correspond fairly to the description and figure of *Etropus rimosus*. All other specimens from the United States coast collected by Dr. Jordan and his associates are, in our opinion, referable to *Etropus crossotus*. The original description of *Citharichthys microstomus*, Gill, fits this species better than any other known. The fish in question is much too elongate for *Etropus crossotus* (depth $2\frac{2}{3}$ in total length), and the mouth is too small for any of the known species of *Citharichthys* (maxillary 4 in head; mandible $2\frac{1}{2}$). We have little doubt of the identity of *Etropus rimosus* and *microstomus*, but leave the matter for further investigation. The separation of *E. rimosus* from *E. crossotus* is not beyond question. (*rimosus*, frosted.)

Etropus rimosus, Goode & Bean, Proc. U. S. Nat. Mus. 1885, 593, coast of Florida, between Pensacola and Cedar Keys, dredged at the depth of 21 fathoms; Goode & Bean, Oceanic Ichthyology, 455, pl. 104. figs. 360, 361, 1896. (The latter figure an excellent representation of the symmetrical, translucent larval form, before the right eye has crossed the forehead.)

Etropus crossotus, Jordan & Evermann, Proc. U. S. Nat. Mus. 1886, 476; not of Jordan & Gilbert.

3062. ETROPUS CROSSOTUS, Jordan & Gilbert.

Head $4\frac{4}{5}$ in length; depth $1\frac{2}{3}$ to 2. D. 76 to 85; A. 56 to 67; V. 6; scales 42 to 48; vertebræ $9+25=34$. Body oval, strongly compressed, with the dorsal and ventral curves nearly equal; both outlines strongly arched anteriorly, the body much deeper in adult specimens. Head very small; snout short; mouth very small, its cleft not so long as diameter of orbit. Teeth conical, pointed, close set, strongly incurved, in a single series, those in upper jaw on blind side only, those in lower jaw on both sides. Eyes large, the lower in advance of the upper, the two separated by a very narrow scaleless ridge, which extends backward above preopercle; edge of opercle on blind side, with a row of conspicuous white cilia. Upper nostril turned somewhat to blind side; anterior nostril on left side, with a very slender cirrus. Dorsal fin commencing over front of upper eye, its middle rays highest, the anterior not elevated; anal fin not preceded by a spine, its middle rays highest; caudal fin very sharply double-truncate, as long as head; pectorals short, that of left side the longer, about $\frac{3}{4}$ length of head; ventral of colored side on ridge of abdomen, the membrane of its last rays nearly reaching base of first ray of anal; ventral of blind side longer than the other, $\frac{1}{2}$ length of head, inserted farther forward than that of colored side. Vent lateral, with a well-developed anal papilla. Scales thin, large, ctenoid on colored side, smooth on blind side, those on the middle part of the body larger; head entirely scaly, except snout and interorbital ridge; rays of vertical fins with scales on the basal half, on colored side; lateral line developed equally on both sides, nearly straight. Color olive brown, with some darker blotches most distinct in the larger specimens; vertical fins finely mottled and streaked with black and gray; pectoral and ventral on left

side spotted. Tropical America on both coasts, north to Cerros Island and North Carolina, south to Panama and Rio Janeiro; the type a single specimen, about 5 inches long, taken with a seine in the Astillero at Mazatlan. This little fish seems to be abundant in all warm and sandy shores of tropical America. It is the smallest and feeblest of all our flounders, and has therefore been generally overlooked by collectors. In the Museum of Comparative Zoology are specimens of this species from Rio Janeiro, Santos, Victoria, Para, and Sambaia, in Brazil. The largest of these is 6 inches in length. Head 5 in length; depth $1\frac{2}{3}$; scales 44; D. 85; A. 67. We have specimens from Charleston, Cedar Keys, New Orleans, Galveston, Beaufort, North Carolina, Mazatlan, Panama, and from several localities along both sides of the coast of Lower California. These vary in form, color, and squamation, but we are unable to point out specific distinctions among them. (*κροσσωτός*, fringed, from the cirri of the subopercle.)

Etropus crossotus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 364, Mazatlan; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 305; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 618; JORDAN & GILBERT, Bull. U. S. Fish Comm. 1882, 108-111; JORDAN & GILBERT, Synopsis, 839, 1883; BEAN, Cat. Fish. Intern. Exh. 1883, 44; JORDAN & SWAIN, Proc. U. S. Nat. Mus. 1884, 234; JORDAN & GOSS, Review Flounders and Soles, 278, 1889.

Etropus microstomus, JORDAN, Proc. U. S. Nat. Mus. 1886, 29; not *Citharichthys microstomus*, GILL.

1053. MONOLENE, Goode.

Monolene, GOODE, Proc. U. S. Nat. Mus. 1880, 338 (*sessilicauda*).

Thyris, GOODE, Proc. U. S. Nat. Mus. 1880, 344 (*pellucidus*; larval form); name preoccupied.

Delothyris, GOODE, Proc. U. S. Nat. Mus. 1883, 110 (*pellucidus*); substitute for *Thyris*.

Body thin, elongate; eyes on the left side, very close together, near the profile; mouth moderate, the length of the maxillary less than $\frac{1}{3}$ that of the head; teeth minute, in a single series, nearly equal on both sides; no teeth on vomer or palatines. Scales rather large, ctenoid on colored side, cycloid on blind side. Lateral line well marked, that of colored side strongly and angularly curved anteriorly, that of blind side nearly straight. Pectoral of blind side wholly absent; dorsal beginning on the snout, its rays all simple. Caudal fin sessile, almost confluent with dorsal and anal. Ventral fins normal, that of the left side on the ridge of the abdomen; gill rakers few, feeble. Vertebræ 43. Deep-sea fishes, closely allied to *Trichopsetta* and *Arnoglossus*, but with the right pectoral obsolete. The translucent larva of *Monolene* is similar to the larva of *Platophrys*. It was at first described as a distinct genus (*Thyris* = *Delothyris*) by Dr. Goode before its true character was recognized. The following are the characters ascribed to the larval genus *Delothyris*:*

* The following are the characters of the species, *Delothyris pellucidus*, Goode:
Colorless, translucent; 3 conspicuous, dusky, longitudinal lines on left side, the middle one faintest. Two streaks on right side; eyes black. Body thin, pellucid, divided into 3 longitudinal tracks by depressions at the bases of the rows of interspinous processes. Scales small, thin, caducous. Head very small; eyes small, protruding, their diameter equal to the interorbital space and $\frac{1}{3}$ the length of the snout; mouth small, formed as in the soles, the upper jaw somewhat hook-shaped. Dorsal fin beginning in advance of the eye, of long, flexible, simple rays, the tips of which are much exserted. Pectorals inserted far below lateral line, that on blind side as long as orbit, the other as long as snout; ventrals reaching past front of anal. Head 5; depth 3. D. 100; A. 80; P. 12 (left), 4 (right). Length 3 inches. (Goode.) Gulf Stream, off the coast of Rhode Island.

Body elongate, soft, and translucent. Head very short; mouth small, toothless. Eyes sinistral, close together, the lower slightly advanced. Pectoral of blind side smallest; ventrals crowded together on median keel of body, their bases prolonged on this keel. Rays simple; dorsal beginning on the snout; caudal subsessile, almost confluent with dorsal and anal. Scales very thin, easily detached, probably cycloid. Lateral line well marked, straight. (*μόνος*, one; *ὠλένη*, arm.)

a. Dorsal rays 99 to 103; anal rays 79 to 81; scales 92. SESSILICAUDA, 3063.
aa. Dorsal rays 124; anal rays 100; scales 105. ATRIMANA, 3064.

3063. MONOLENE SESSILICAUDA, Goode.

Head 5 in body; depth $2\frac{2}{3}$. D. 99 to 103; A. 79 to 84; scales 23–92–25; eye 4 in head; highest dorsal ray 2; highest anal rays slightly more than 3; pectoral $1\frac{1}{3}$; caudal nearly 1. Body moderately elongate; maxillary extending slightly past front of lower eye, with uniserial, subequal teeth; lower eye in advance of upper; interorbital space very small, less than $\frac{1}{6}$ the diameter of eye; head everywhere closely scaled; scales ctenoid; lateral line strongly curved over anterior $\frac{2}{3}$ of pectoral, the curve with 2 angles, 72 scales along straight portion; lateral line of blind side nearly straight. Origin of dorsal over anterior edge of lower eye, longest rays in the posterior fourth of the fin; origin of anal under base of pectoral; pectoral present only on eyed side. Color on left side ashy brown, with numerous more or less distinct darker brown spots; on blind side white, pectoral blackish with traces of lighter transverse bands. Specimens from shallow water near Key West (Coll. Prof. C. C. Nutting), according to Mr. Garman, are much more brightly colored. These are "grayish brown, with numerous spots of darker to blackish over head and body, the spots being $\frac{1}{2}$ as large as the eye or smaller, arranged for the greater part in broad transverse bands as wide as the interspaces, of which bands the first and foremost passes from the nape to the opercle, the second lies immediately behind the pectoral, the third just in front, and the fourth just behind the middle of total length, and the fifth, more indistinct, crosses near the ends of the dorsal and anal; the caudal is crossed by 2 rather indefinite narrow streaks; the pectoral is white at its base and bears 3 or 4 narrow curved transverse bands of white, separating 3 or 4 similar bands of black, which with the white are more distinct in the lower half of the fin." Specimens from 150 fathoms or more have markings similar but less distinct. D. 104; A. 84; V. 6; P. 11; scales 22–93–24. Deep waters of the Gulf Stream, Cape Cod to Key West. (*sessilis*, sessile; *cauda*, tail.)

Monolene sessilicauda, Goode, Proc. U. S. Nat. Mus. 1880, 337, 338, deep sea south of New England; Goode, Proc. U. S. Nat. Mus. 1880, 472; Jordan & Gilbert, Synopsis, 841, 1883; Jordan & Goss, Review Flounders and Soles, 280, 1889; Goode & Bean, Oceanic Ichthyology, 452, figs. 357 A & B, 1896; Garman, Bull. Iowa Lab. Nat. Hist. 1896. 91.

Thyris pellucidus, Goode, Proc. U. S. Nat. Mus. 1880, 344, Gulf Stream off the coast of Rhode Island.

Delothyris pellucidus, Goode, Proc. U. S. Nat. Mus. 1883, 109.

3064. MONOLENE ATRIMANA, Goode & Bean.

Head 4¼ in length of body; depth about 3. D. 124; A. 100; scales 30–105–32; eye 2⅔ in head; maxillary 3; highest dorsal ray 2; left ventral 3½; pectoral 4¼ in body; caudal 6. Body rather elongate; snout slightly produced. Mouth oblique, the maxillary extending to a little behind front of lower eye, teeth uniserial, well developed on both sides; lower eye in advance of upper; interorbital a very narrow ridge, about 9 in eye; nostrils in very short tubes in the same line with the interorbital ridge, the posterior one is slightly less distant from lower eye than the anterior one is from the snout; head everywhere scaly; lateral line strongly arched over anterior third of pectoral. Origin of dorsal on blind side above front of lower eye, longest rays in posterior fourth of fin; highest rays of anal a little higher than dorsal rays; pectoral on eyed side only; caudal sessile, rounded. Color light brownish gray, right ventral pale, other fins dusky; pectoral and eyelids black. West Indies. The type was taken by the *Blake* in 288 fathoms, off Barbados; its length is 114 mm. (Goode & Bean.) (*ater*, black; *manus*, hand.)

Monolene atrimana, Goode & Bean, Bull. Mus. Comp. Zool., xii, 155, 1886, deep waters off Barbados; Jordan & Goss, Review Flounders and Soles, 280, 1889; Goode & Bean, Oceanic Ichthyology, 455, fig. 358, 1896.

Family CCXX. SOLEIDÆ.

(The Soles.)

Body oblong or elongate, usually scaly; mouth very small, much twisted toward the eyed side; the teeth in villiform bands, very small or obsolete; eyes small, close together, with or without a bony ridge between them; edge of preopercle adnate, concealed by the skin and scales; gill openings narrow, the gill membranes adnate to the shoulder girdle above; pectoral fins small or wanting; ventral fins small, 1 or both sometimes wanting. Small fishes living on sandy bottoms, similar to the *Pleuronectidæ* in structure, but much degraded, the fins and teeth having lost many of their distinctive qualities. The vertebræ are usually in increased numbers.*

* The following are the numbers of vertebræ in several species of *Soleidæ:*

I.—Achirinæ.

Achirus fasciatus	8 + 20 = 28
Achirus inscriptus	9 + 19 = 28

II.—Soleinæ.

Synaptura zebra	[8 + 41 = 49]
Solea solea	9 + 40 = 49
Solea kleini	10 + 37 = 47
Solea aurantiaca	[46]
Quenselia ocellata	9 + 28 = 37
Microchirus luteus	8 + 29 = 37
Monochirus hispidus	9 + 25 = 34

III.—Cynoglossinæ.

Symphurus atricaudus	10 + 42 = 52
Symphurus nigrescens	9 + 40 = 49
Symphurus plagiusa	9 + 38 = 47

They are numerous in the warm seas, and those of sufficient size are valued as food. Genera about 12; species 150. The North American species belong to 2 subfamilies very different one from the other. The soles are naturally divisible into 3 subfamilies, each quite distinct from the others, and possibly independently descended or degraded from normal *Pleuronectidæ*. The *Achirinæ*, or American soles, are apparently allied to the *Psettinæ*, and as in the latter, the ventral fin of the eyed side extends along the ridge of the abdomen. The *Soleinæ*, or European soles, show in the insertion of ventral and in other respects a strong resemblance to the *Pleuronectinæ*. The more aberrant *Cynoglossinæ*, or tongue fishes, are perhaps degraded *Soleinæ*, but the eyes are sinistral, as in the *Psettinæ*. In the *Soleinæ* and *Achirinæ* the eyes are dextral, as in the *Pleuronectinæ*.

ACHIRINÆ:

I. Soles with the eyes on the right side and separated by a distinct bony ridge; the ventral with long base confluent with the anal. Body oblong or ovate, with the color on the right side; eyes moderate or small, the upper eye usually more or less in advance of the lower; mouth small, more or less twisted toward the blind side; teeth little developed, in villiform bands; edge of opercle adnate, usually concealed by the scales; gill openings more or less narrowed, the gill membranes adnate to the shoulder girdle above; blind side of head usually with fringes; pectoral fins small, sometimes wanting; ventral fins developed, one or both of them sometimes obsolete; scales usually ctenoid, rarely wanting; lateral line straight, usually single; right ventral with extended base, confluent with the anal fin.

a. Gill openings of moderate extent, confluent below; vertical fins well separated; body ovate in outline, the depth nearly ½ the length; pectoral fins rudimentary or wanting; lateral line straight; scales well developed, ctenoid, those on the head more or less enlarged, those of the blind side of the head with fringes; vertebræ about 28. ACHIRUS, 1054.

aa. Gill openings very small, separate, each reduced to a small slit below angle of opercle; right ventral beginning at the chin; pectoral fins minute or wanting; lateral line straight; snout dilated, the dorsal beginning upon it.

b. Scales present, ctenoid; caudal somewhat confluent with dorsal.

c. Left ventral rudimentary, with 2 rays. APIONICHTHYS, 1055.

bb. Scales none; caudal free from dorsal and anal. GYMNACHIRUS, 1056.

CYNOGLOSSINÆ:

II. Soles with the eyes on the left side, not separated by a bony ridge. Body elongate, more or less lanceolate in outline, with the color on the left side; eyes small, very close together, with no distinct interorbital ridge between them; mouth small, twisted toward the blind side; teeth little developed, in villiform bands; gill openings narrow, the gill membranes adnate to the shoulder girdle above, joined together and free from the isthmus below; pectoral fins wanting (in the adult); ventral fins small, that of the blind side often wanting; vertical fins more or less confluent; scales ctenoid; lateral lines sometimes wanting, sometimes duplicated.

d. Ventral fin of eyed side only present, free from the anal; no pectoral fins; no lateral line; head without fringes. SYMPHURUS, 1057.

1054. ACHIRUS, Lacépède.

(AMERICAN SOLES.)

Achirus, LACÉPÈDE, Hist. Nat. Poiss., IV, 659, 1803 (*fasciatus*, etc.).

Achirus, CUVIER, Règne Animal, Ed. 2, II, 343, 1829 (restriction to *fasciatus*, etc.).

Trinectes, RAFINESQUE, Atlantic Journal and Friend of Knowledge, I, 1832 (*scabra*).

Grammichthys, KAUP, Archiv fur Naturgsch. 1858, 94 (*lineatus; fasciatus*); *Achirus* being restricted to *Pardachirus barbatus*, etc.

Monochirus, KAUP, Archiv fur Naturgsch. 1858, 94 (*maculipinnis*); not of RAFINESQUE. 1814, a genus of *Soleinæ*.

? Aseraggodes, KAUP, Archiv fur Naturgsch. 1858, 103 (*guttulata*).

Baiostoma, BEAN, Proc. U. S. Nat. Mus. 1882, 413 (*brachiale*).

Bæostoma, JORDAN & GILBERT, Synopsis, 965, 1883; amended orthography.

Eyes and color on the right side. Body oblong, bluntly rounded anteriorly. Head small; eyes small, close together, the upper eye in advance of the lower, the two separated by a bony ridge; mouth small, somewhat turned toward the colored side; nasal flaps present, the nostril of the blind side fringed; lip of the colored side fringed; teeth very small, on blind side only; gill openings rather narrow, but confluent below, not reduced to a slit; the branchiostegal region scaled. Head closely scaled everywhere, the scales on the colored side similar to those on the body, those of the nape and chin much enlarged; scales on the blind side anteriorly with their pectinations more or less produced, forming cirri; scales of both sides extremely rough, extending on the fins. Lateral line straight, simple; edge of preopercle covered by the scales. Dorsal beginning on the snout, low in front and thickly scaled, its rays divided; anal fin similar, without spine; caudal fin free, convex; caudal peduncle very short and deep; pectoral fin of left side wanting, that of right side small or obsolete; ventral rays 3 or 4, the ventral fin of the colored side long, connected with the anal by a membrane. This strongly marked genus contains numerous species, all very closely related, and nearly all American. It has been united by Dr. Günther with *Solea*, but for no good reason, as the number of vertebræ is very much fewer than in the European soles, and the right ventral fin is decurrent along the abdomen and united with the anal in the American soles, while it is short and wholly free in all the European forms. The 2 groups belong in fact to distinct subfamilies. It is also worth noticing that the name *Achirus* is prior in date to that of *Solea*. The species with rudimentary pectoral fins have been set apart by Dr. Bean to form the genus *Baiostoma*, but the very slight development of these organs in some of the species and the evidently very close relationship of them all lead us to regard *Baiostoma* as a subgenus only. If we follow Kaup in restricting the name *Achirus* to the Asiatic group called *Pardachirus*, the present genus would receive the name of *Trinectes*. It seems to us, however, that both Lacépède and Cuvier regarded the species called by us *fasciatus* as the type of their genus *Achirus*. (ἄχειρ, without hands; without pectoral fins.)

BAIOSTOMA (βαιός, small; στόμα, mouth):

a. Pectoral fins small, present at least on the right side.

b. Pectoral fin present on both sides, that of the left side rudimentary, of a single ray; that of the eyed side with about 3 rays.

c. Dorsal rays 60 to 67; anal rays about 48; scales 80; depth 1⅔ in length. Color brownish, irregularly spotted with darker, and with about 10 black vertical lines crossing the lateral line. ACHIRUS, 3065.

cc. Dorsal rays 53 to 57; scales 75 to 80; depth 1⅔ in length; scales not very rough, those of colored side with scattered, hair-like appendages, some black, others pale. Color olivaceous; head, body, dorsal, and anal fins covered with a network of dark lines; traces of about 8 dark cross streaks sometimes present. INSCRIPTUS, 3066.

bb. Pectoral of right side only present.

d. Dorsal rays 65 or 66; anal rays 48 to 51.

e. Pectoral well developed, with about 6 rays; scales of eyed side without hair-like filaments; scales of lateral line 77 to 80; chin little prominent. KLUNZINGERI, 3067.

dd. Dorsal rays 50 to 58; anal rays 38 to 48.

f. Pectoral fin of 4 to 6 rays, considerably longer than eye; body with 8 to 10 narrow, vertical dark bars, these sometimes obsolete with age.

g. Vertical fins all with round, dark spots, these usually especially distinct on the caudal fin; some of the scales of eyed side with black, hair-like appendages; pectoral fin with 5 or 6 rays, about 3 in head, its length equal to that from outer edge of 1 eye to outer edge of another. Head $3\frac{1}{4}$; depth $1\frac{1}{3}$. Body with 8 narrow, vertical cross streaks. LINEATUS, 3068.

gg. Vertical fins dark, without distinct markings. Body broad, ovate, the depth about $1\frac{1}{2}$ in length; pectoral fin with 4 rays; scales of right side with numerous black, hair-like appendages. D. 56; A. 42; scales 70. MAZATLANUS, 3069.

ff. Pectoral fins of 1 to 3 rays, about as long as eye.

h. Body with 6 narrow, dark bands, these sometimes obsolete. Body rather narrowly ovate, its depth $1\frac{5}{8}$ in length. D. 58; A. 44; scales 85. FONSECENSIS, 3070.

hh. Body with about 10 black cross lines; depth $1\frac{5}{8}$ in length. D. 61; A. 44; scales 60; pectoral of a single ray. FISCHERI, 3071.

hhh. Body with very numerous (20 to 40) black cross bands, which are as broad as the interspaces.

j. Blind side of snout with few fringes. Depth $1\frac{1}{2}$ in length. D. 55; A. 48; scales 80. Body covered by many blackish, wavy bands; caudal with black spots. SCUTUM, 3072.

ACHIRUS:

aa. Pectoral fins wholly wanting.

k. Dorsal rays 46; anal rays 33; right lower lip with serrated fringes; nostril in a fringed tube; depth $1\frac{1}{2}$ in length; head 3. Color brown, head and body with numerous large, rounded or kidney-shaped white spots, edged with dark brown; scales 70. FIMBRIATUS, 3073.

kk. Dorsal rays 50 to 55; anal rays 37 to 46; right lower lip fringed; left nostril with some fringes; depth $1\frac{1}{3}$ in length; head 4; none of the scales of eyed side with hair-like appendages. Color dusky olive, more or less mottled with about 8 dark vertical stripes, these varying very much in width and number; caudal spotted. FASCIATUS, 3074.

kkk. Dorsal rays 59 or 60; anal rays 41 to 45; snout and chin without evident fringe or barbel; right lower lip fringed; head 4 in length; depth $1\frac{3}{4}$; scales 64. Body with 12 black cross bands with narrower ones between; caudal spotted. PANAMENSIS, 3075.

Subgenus BAIOSTOMA, Bean.

3065. ACHIRUS ACHIRUS (Linnæus).

D. 60 to 67; A. 48; P. right 3, left 1; scales 80. Pectorals rudimentary on both sides; right ventral fin composed of 5 rays, which are continuous with the anal. Scales on the nape and on chin twice as large as those on body; snout with a few fringes on blind side; right lower lip fringed. Height of body $1\frac{5}{8}$ in total length (without caudal); width of interorbital space nearly equal to, or rather more than diameter of eye; upper eye

slightly in advance of lower; longest dorsal rays are in posterior fifth of the fin, $\frac{2}{3}$ of length of head; caudal rounded, rather longer than head. Brownish, irregularly spotted with darker, and with about 10 black vertical lines crossing the lateral line. Coasts of Surinam. (Günther.) Not seen by us.

We know this species only from Dr. Günther's description. *Pleuronectes achirus*, Linnæus, is based on a description by Gronow of some *Achirus* from Surinam. Gronow's fish agrees with the present species in having 60 dorsal rays and 48 anal rays, in being brown, with transverse black bands, with dark spots on the fins, as well as in coming from Surinam. But Gronow explicitly denies the presence of pectorals, and the present species has rudimentary pectoral fins on both sides. Probably these were overlooked by Gronow, and as no other species found in the same region has so large a number of rays, we feel justified in the use of the name *Achirus achirus* for this species. (ἀ-, without; χείρ, hand.)

Pleuronectes oculis dextris, corpore glabro, pinnis pectoralibus nullis, Gronow, Museum, I, No. 42, Surinam.
Pleuronectes achirus, Linnæus, Syst. Nat., Ed. X, 268, 1758, Surinam; based on Gronow.
Solea gronovii, Günther, Cat., IV, 472, 1862, Surinam.
Achirus gronovii, Jordan, Proc. U. S. Nat. Mus. 1886, 602.
Achirus achirus, Jordan & Goss, Review Flounders and Soles, 311, 1889.

3066. ACHIRUS INSCRIPTUS, Gosse.

Head $3\frac{3}{4}$ in body; depth $1\frac{1}{4}$. D. 53 to 57; A. 40; scales 75 to 80; interorbital width less than eye; upper eye in advance of lower. Pectoral fin present on each side, that of the left side rudimentary, of a single ray; that of the eyed side with about 3; left ventral with 1 or 2 small rays, in some specimens entirely absent; right ventral joined to anal. Scales smaller and less rough than usual in this genus, those of nape scarcely enlarged on eyed side, those of blind side much fringed; scales of colored side with scattered hair-like appendages, some black, others pale. Color olivaceous; head, body, dorsal, and anal fins covered with a network of dark lines; traces of about 8 dark cross streaks sometimes present; caudal fin yellowish, nearly plain, or with a few dark dots or reticulations, its base dusky. Vertebræ $8+20=28$. West Indies north to Key West. Known to us from numerous specimens taken by Dr. Jordan at Key West, and from specimens from Hayti, in the museum at Cambridge. These specimens belong undoubtedly to the species called *reticulatus* by Poey, and this is apparently not different from the *inscriptus* of Gosse, as the agreement with the latter is even closer than with the former description. (*inscriptus*, written on.)

Achirus inscriptus, Gosse, Nat. Sojourn Jamaica, 52, pl. 1, fig. 4, 1851, Jamaica; Jordan, Proc. U. S. Nat. Mus. 1884, 143; Jordan & Goss, Review Flounders and Soles, 311, 1889.
Monochir reticulatus, Poey, Memorias, II, 317, 1861, Cuba; Poey, Synopsis, 409; Poey, Enumeratio, 139.
Solea reticulata, Günther, Cat., IV, 472, 1862.
Solea inscripta, Günther, Cat., IV, 473, 1862.
Bœostoma reticulatum, Bean & Dresel, Proc. U. S. Nat. Mus. 1884, 152.

3067. ACHIRUS KLUNZINGERI (Steindachner).

Head $3\frac{3}{4}$ in body; depth $1\frac{5}{7}$. D. 65; A. 51; scales 37–80–42; eye $3\frac{3}{5}$ in head; height of dorsal and anal $1\frac{1}{2}$; caudal 1. Body moderately broad; eyes in the same vertical line; interorbital as wide as length of eye; angle of mouth reaching a little past front of lower eye; right under lip fringed; scales near upper profile of head enlarged, all scales strongly ctenoid; scales of eyed side without hair-like filaments. Pectoral of right side only present, with about 6 rays; caudal round behind. Color brownish, with 9 or 10 narrow blackish cross lines; small rounded blackish spots on the membranes of each of the vertical fins, much as in *A. lineatus*. (Steindachner.) Pacific coast of tropical America; Panama to Guayaquil. (Named for Dr. C. B. Klunzinger, Professor of Zoology at Stuttgart, author of Memoirs on the Fishes of the Red Sea.)

Solea klunzingeri, STEINDACHNER, Zur Fische des Cauca und der Flüsse bei Guayaquil, 44, 1879, Guayaquil.

Achirus klunzingeri, JORDAN, Proc. U. S. Nat. Mus. 1885, 391; JORDAN & GOSS, Review Flounders and Soles, 312, 1889.

3068. ACHIRUS LINEATUS (Linnæus).

Head $3\frac{1}{2}$; depth about $1\frac{1}{2}$. D. 49 to 58; A. 38 to 44; scales 75 to 85. Pectoral fin of right side only developed, of 4 to 6 rays, considerably longer than eye. Body with 8 to 10 narrow vertical dark bars, these sometimes obsolete with age; vertical fins all with round dark spots, these usually especially distinct on the caudal fin; some of the scales of eyed side with black, hair-like appendages; pectoral fin with 5 or 6 rays, about 3 in head, its length equal to that from outer edge of one eye to outer edge of another. Color brown, the young spotted with whitish, the adult sometimes with darker; body with about 8 narrow vertical cross streaks of blackish. West Indies and Brazil, Florida Keys to Uruguay; common and variable.

We have placed the Florida species, *comifer* and *brachialis*, in the synonymy of *lineatus*. They differ from the latter only in the slightly smaller number of the scales and fin rays. The following table shows our count of a number of specimens from different localities:

Locality.	Species.	D.	A.	Scales.
Key West	comifer	50	35	55 to 67
Pensacola	brachialis	51	37	75 to 77
Cienfuegos	lineatus	54	43	85
Rio Janeiro	maculipinnis	57	42	85
Do	maculipinnis	54	44	72
Rio Grande do Sul	maculipinnis	49	38	70
Coary	maculipinnis	53	40	68
Manacapuru	maculipinnis	55	42	75

It is evident from this table that neither the fin rays nor the scales form characters by which the subspecies can be absolutely distinguished. It is evident also, from the examination of large series of specimens, that the

coloration is subject to very great variations—as great as in *Achirus fasciatus*. In some of these the caudal is dark and immaculate, in others pale and usually profusely spotted. In some the ground color is nearly plain blackish, in others it is pale, usually with narrow dark cross bands, but sometimes closely spotted everywhere. The specimens examined by us are from Pensacola and Egmont Key (*brachialis*), Key West (*comifer*), Cienfuegos (Cuba, Poey), Coary, Teffy, Tapajos, Porto Alegre, Pernambuco, Cannarivieras, Manacapuru, Porto do Moz, Rio Grande do Sul, Rio Janeiro, San Matheo, Rosario, Itabapuana, Obidos, Xingu, Gurupa, Jutaby, Curaçao, Para, Bahia, Santarem, Iça, Fonteboa, San Paolo, Rio Trompetas, Sambaia, Manes, Javary, and Tabatinga. The species would appear to be one of the commonest in Brazil. (*lineatus*, striped.)

a. Var. *lineatus*.

Passer lineis transversis notatus, SLOANE, Jamaica, 2, 77, pl. 246, fig. 2, 1725, Jamaica.
Pleuronectes fuscus subrotundus glaber, BROWNE, Jamaica, 445, 1756, Jamaica.
Pleuronectes lineatus, LINNÆUS, Syst. Nat., Ed. x, 268, 1758, Jamaica; based on BROWNE and SLOANE; not of Ed. xii, which is *Achirus fasciatus*.
Monochir maculipinnis, AGASSIZ, Spix, Pisc. Brasil., 88, pl. 49, 1829, Brazil; POEY, Synopsis, 409, 1868.
Monochir lineatus, QUOY & GAIMARD, Voy. Uranie, Zool., 238, 1824.
Achirus lineatus, D'ORBIGNY, Voyage Amér. Mérid., Poiss., pl. 16, fig. 2, 1847; JORDAN & GOSS, Review Flounders and Soles, 312, 1889.
Solea maculipinnis, GÜNTHER, Cat., iv, 473, 1862; KNER, Novara Fische, iii, 289, 1866.
Achirus maculipinnis, JORDAN, Proc. U. S. Nat. Mus. 1886, 602.

b. Var. *brachialis*.

Baiostoma brachialis, BEAN, Proc. U. S. Nat. Mus. 1882, 413, Appalachicola Bay and South Florida. (Types, Nos. 26605 and 30463. Coll. Silas Stearns.)
Bœostoma brachiale, JORDAN & GILBERT, Synopsis, 965, 1883.
Achirus brachialis, JORDAN, Proc. U. S. Nat. Mus. 1884, 149.

c. Var. *comifer*.

Achirus comifer, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1884, 31, Key West. (Coll. Dr. Jordan.)

3069. ACHIRUS MAZATLANUS (Steindachner).

(MEXICAN SOLE; LENGUADO DE RIO; TEIPALCATE.)

Head $3\frac{1}{3}$ in body; depth $1\frac{1}{2}$. D. 56; A. 42; scales 70; eye $7\frac{1}{2}$ in head; dorsal and anal rays $3\frac{1}{2}$ in depth of body; caudal 3. Body broad, oval; eyes small, the upper in advance of the lower; interorbital about $\frac{1}{2}$ the diameter of eye; nostril in a tube, placed just above middle of mouth; pectoral developed on eyed side only, with about 4 rays; origin of dorsal on tip of snout; greatest height of dorsal and anal behind their middle; scales of right side with numerous black hair-like appendages. Color brownish, with 8 or 9 narrow vertical black bars; fins dark, without distinct markings. West coast of Mexico, entering all streams; common and variable. Many specimens from Mazatlan and Rio Presidio examined by us, as also a specimen from Chiapas. (Name from Mazatlan,* the river of the deer.)

* "With eternal sun above thee,
'T is not strange the tall deer loved thee,
That he gave his name, Mazatl,
To thy river, Mazatlan!"

Solea mazatlana, STEINDACHNER, Ichth. Notizen, IX, 23, 1869, Mazatlan; JORDAN & GILBERT, Bull. U. S. Fish Comm. 1882, 108.

Solea pilosa, PETERS, Berliner Monatsber. 1869, 709, Mazatlan.

Achirus mazatlanus, JORDAN, Proc. U. S. Nat. Mus. 1885, 391; JORDAN & GOSS, Review Flounders and Soles, 313, 1889; JORDAN, Proc. Cal. Ac. Sci. 1895, 505.

3070. ACHIRUS FONSECENSIS (Günther).

Head $3\frac{1}{3}$; depth $1\frac{5}{6}$. D. 58; A. 44; P. 2; scales about 85. No trace of a pectoral on the left side, that on the right not much longer than the eye; right ventral fin composed of 5 rays, which are continuous with the anal. Scales on the nape twice or thrice as large as those on the body. The upper part of the snout slightly overlaps the lower jaw. The left anterior part of the head with numerous tentacles; the right lower lip with very distinct slender fringes; nostril on the right side in a wide and short tube. The width of the interorbital space is less than the diameter of the eye; the upper eye is in advance of the lower. The rays of the vertical fins are branched; the longest dorsal rays are $\frac{2}{3}$ of the length of the head. Caudal rounded, as long as the head. Brownish olive, with 6 pairs of deep brown vertical lines extending on the dorsal and anal fins. Pacific Coast of Tropical America. (Günther.) Described from 1 specimen $4\frac{1}{4}$ inches long, from Gulf of Fonseca; 2 others since taken by us in Rio Presidio, near Mazatlan. (Name from Fonseca, the type locality.)

Solea fonsecensis, GÜNTHER, Cat., IV, 475, 1862, Gulf of Fonseca. (Coll. Sir John Richardson.)

Achirus fonsecensis, JORDAN & GOSS, Review Flounders and Soles, 314, 1889; JORDAN, Proc. Cal. Ac. Sci. 1895, 505.

3071. ACHIRUS FISCHERI (Steindachner).

(PEGE OJA.)

Head $3\frac{1}{2}$ in body; depth $1\frac{8}{9}$; caudal $3\frac{1}{4}$. D. 61; A. 44; P. right. 1; V. 5; scales 60 to 62. Pectoral wanting on left side, rudimentary on right, of a single ray scarcely longer than eye; right ventral connected with the anal. Scales on neck and lower portion of head $1\frac{1}{2}$ to 2 times as large as those on body; right side of lower lip fringed; upper jaw not projecting forward over lower jaw; left side of head with dermal flaps only around corner of mouth and on lower jaw. Eyes small, the upper a little further forward than the other, and 2 in snout; breadth of forehead equaling diameter of eye. Dorsal rays increasing gradually in length to the forty-eighth, which is about $\frac{2}{3}$ length of head. Scales strongly ctenoid, the teeth considerably largest at the middle; rays of all the fins, except of pectorals, scaled to their tips, the membranes less fully scaled; only the anterior third of the caudal membranes scaled, and between the last dorsal and anal rays the scales extend slightly upon the fins. A few black thread-like appendages on right side of body between scales. Color of right side dark gray; 2 or 3 blackish cross lines on head, about 10 on body, between them numberless spots of similar color; spots on fins, especially those on caudal, a little larger; a few large dark spots on body, irregular

and poorly defined; blind side reddish yellow. Total length about 10 cm. Rio Mamone, near Panama; known to us only from Steindachner's description and figure. (Steindachner.) Not seen by us. (Named for W. Fischer.)

Solea fischeri, STEINDACHNER, Beiträge Kenntniss Fluss-Fische Sudamer., I, 13, 1879, Rio Mamone, near Panama.
Achirus fischeri, JORDAN, Proc. Ac. Nat. Sci. Phila. 1887, 391.

3072. ACHIRUS SCUTUM (Günther).

Head $3\frac{3}{8}$; depth $1\frac{1}{2}$. D. 55; A. 48; P. 3; scales 80. No trace of a pectoral on left side; right pectoral quite rudimentary, scarcely longer than the eye; right ventral composed of 5 rays, which are continuous with anal. Scales on nape nearly twice as large as those on the body; snout with scarcely any fringes on the blind side, right lower lip fringed. Width of interorbital space less than horizontal diameter of orbit; upper eye slightly in advance of lower. Longest dorsal rays in posterior third of fin, $\frac{2}{3}$ length of head; caudal rounded, longer than head. Grayish; head, body, and fins with numerous blackish, irregular, waving, sometimes bifurcate, transverse bands, which are broader than the interspaces; caudal with rounded deep black spots; the left side uniform white. Pacific coast of Central America. (Günther.) Not seen by us. (*scutum*, a shield)

Solea scutum, GÜNTHER, Cat., IV, 474, 1862, Gulf of Fonseca, Panama.
Achirus scutum, JORDAN & GOSS, Review Flounders and Soles, 314, 1889.

Subgenus ACHIRUS.

3073. ACHIRUS FIMBRIATUS (Günther).

Head 3; depth $1\frac{1}{2}$. D. 46; A. 33; scales 70. Pectorals none; right ventral of 5 rays, which are continuous with the anal. Scales on nape 4 times, those on the chin twice, as large as on the body. Upper part of the snout slightly bent downward over the mandible and forming a short hook; right lower lip broadly fringed, each fringe being serrated; nostril in a short, wide, fringed tube. No tentacles on left side of head. Width of interorbital space equaling diameter of circular small orbit; upper eye slightly in advance of lower. Longest dorsal rays $\frac{2}{3}$ of length of head; rays of vertical fins branched; caudal rounded, its length being $\frac{1}{4}$ of the total. Brown; head and body with numerous large, rounded, or kidney-shaped white spots, edged with dark brown. Gulf of Fonseca, Central America. (Günther.) Known from 1 specimen, $3\frac{1}{6}$ inches long. (*fimbriatus*, fringed.)

Solea fimbriata, GÜNTHER, Cat., IV, 477, 1862, Gulf of Fonseca. (Coll. Sir John Richardson.)
Achirus fimbriatus, JORDAN & GOSS, Review Flounders and Soles, 315, 1889.

3074. ACHIRUS FASCIATUS, Lacépède.

(AMERICAN SOLE; HOG CHOKER.)

Head 4 in body; depth $1\frac{4}{5}$. D. 50 to 55; A. 37 to 46; scales 66 to 75; eye 7 in head; height of dorsal and anal nearly 2; caudal $1\frac{1}{5}$. Body broad, regularly elliptical; mouth moderate, reaching just past front of lower

eye; right lower lip fringed; eyes very small, the upper one in advance of the lower; nostril ending in a wide tube, nearer lower eye than tip of snout; interorbital space with scales, more than ½ eye; head and body scaled with strongly ctenoid scales, none of them with hair-like appendages; lateral line nearly straight; gill opening short, about twice as long as maxillary. Origin of dorsal on tip of snout; last few rays of dorsal and anal rapidly decreasing, giving the fins a truncate appearance posteriorly; pectorals wholly wanting; caudal rounded. Color dusky olive, more or less mottled, and with about 8 dark, vertical stripes, these varying very much in width and in number; vertical fins with the membrane of every second or third pair of rays blackish, besides dark cloudings at base of fin; caudal with numerous longitudinally oblong spots; blind side often with round, dark spots, especially in northern specimens, usually immaculate in southern ones (var. *browni*). Vertebræ 8+20=28. South Atlantic and Gulf coast, from Cape Ann to Brazos Santiago, ascending sandy streams in shallow water. The species is the best known of the American soles, and it is common along our coast, ascending the rivers for a considerable distance above tide water. It seldom exceeds 5 or 6 inches in length, and is of but little value as food on account of its small size. Here described from a specimen, 4 inches long, from Beaufort, North Carolina. This species has not yet been recorded from the West Indies. The form found along the Gulf coast has been described as a distinct species under the name *Solea browni*. The differences are not very evident. We have compared a number of specimens from Boston (*fasciatus*) with others from Pensacola, and find the following differences, none of which is constant: In the Gulf variety (*browni*) the blind side is always immaculate, while in almost all Atlantic examples (*fasciatus*) the blind side is profusely covered with round, dark spots. In 1 specimen, however (11360, Boston), the blind side is immaculate. The darker cross streaks on the eyed side are usually broader and more numerous in southern specimens, and the scales on the blind side of the head rougher. There are no constant differences either in the fin rays or in the scales. We have examined specimens of this species from Boston, Chestertown, Tarrytown, New York, Port Monmouth, Havre de Grace, Potomac River, Neuse River, Beaufort, Charleston, Pensacola, Mobile, and Galveston. In 1 large specimen from Pensacola (11482, M. C. Z.) there is a rudiment of a pectoral fin on the eyed side. It consists of a single ray ⅔ as long as the eye. (*fasciatus*, banded.)

Pleuronectes lineatus. Linnæus, Syst. Nat., Ed. xii, 458, 1766, on a specimen from Charleston, received from Dr. Garden; not *Pleuronectes lineatus* of Ed. x.

Achirus fasciatus, Lacépède, Hist. Nat., Poiss., iv, 659, 662, 1803, Charleston; excl. syn., description based entirely on the Linnæan account of the fish sent by Garden; Jordan & Goss, Review Flounders and Soles, 315, 1889.

Pleuronectes mollis, Mitchill, Trans. Lit. and Phil. Soc. N. Y., i, 1815, 388, pl. 2, fig. 4, New York.

Pleuronectes apoda, Mitchill, Amer. Monthly Mag. and Crit. Rev., Feb., 1818, 244, Straits of Bahama; perhaps *A. lineatus*.

Trinectes scabra, Rafinesque, Atlantic Journal and Friend of Knowledge, i, 1832, Pennsylvania, in fresh water.

Solea browni, Günther, Cat., iv, 477, 1862, New Orleans; Texas.

Achirus lineatus, CUVIER, Règne Animal, Ed. 2. II, 343, 1829; GILL, Cat. Fishes East Coast N. Am., in Rept. U. S. Fish Comm., 1871–72, 794; JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1878, 368; GOODE, l. c., 1879, 110; GOODE & BEAN, l. c., 1879, 123; BEAN, l. c., 1880, 77; JORDAN & GILBERT, l. c., 1882, 618; BEAN, l. c., 1883, 365.
Grammichthys lineatus, KAUP, Archiv fur Naturgesch. 1858, 101.
Achirus mollis, STORER, Synopsis, 228, 1846; STORER, Hist. Fish. Mass., 206, pl. 32, 1867; DE KAY, New York Fauna: Fishes, 303, pl. 49, fig. 159, 1842.
Achirus achirus mollis, JORDAN, Cat. Fish. N. A., 137, 1885.
Solea achirus, GÜNTHER, Cat., IV, 476, 1862; not *Pleuronectes achirus* L.
Achirus achirus, JORDAN, Proc. U. S. Nat. Mus. 1885, 19; JORDAN, Cat. Fish. N. A., 137, 1885.
Achirus lineatus, var. *browni*, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 305.

3075. ACHIRUS PANAMENSIS (Steindachner).

Head 4 in body; depth $1\frac{3}{4}$. D. 59; A. 45; scales 63 to 65; highest dorsal and anal spines 2 in head; caudal 1. Body broad, elliptical; angle of mouth below middle of lower eye; edge of lower lip, on the eyed side, fringed; eyes small, the upper in advance of the lower; interorbital scaled, scarcely as wide as diameter of eye; scales ctenoid; pectorals wholly wanting; origin of dorsal on end of snout; highest rays of anal and dorsal behind their middle; ventral rays short; middle rays of caudal the longest, fin sharply rounded behind. Color brown; about 12 dark cross bands on head and body; between these faint, paler cross bands, which form spots on dorsal and anal; caudal similarly spotted, the spots forming obscure cross bands. (Steindachner.) Pacific coast of tropical America, Panama.

Solea panamensis, STEINDACHNER, Ichthyol. Beiträge, V, 10, taf. II, 1876, Panama.
Achirus panamensis, JORDAN & GOSS, Review Flounders and Soles, 316, 1889.

1055. APIONICHTHYS, Kaup.

Apionichthys, KAUP, Archiv fur Naturgesch. 1858, 104 (*dumerili*).
Soleotalpa, GÜNTHER, Cat., IV, 489, 1862 (*unicolor*).

Gill openings very small, separate, each reduced to a slight slit below angle of opercle; right ventral beginning at the chin, confluent with the anal; pectoral fins wanting or very small; lateral line present, straight; eyes small; snout dilated, the dorsal beginning upon it. Scales present, ctenoid; caudal fin somewhat confluent with dorsal. Left ventral rudimentary, with 2 rays. West Indies and Brazil. This genus is closely related to *Achiropsis*, Steindachner, of the rivers of Brazil, but in the latter genus the ventrals are both well developed. (*ἀ-*, not; *πίων*, fat; *ἰχθύς*, fish.)

3076. APIONICHTHYS UNICOLOR (Günther).

D. 76; A. 57; V. right 5, left 2; scales 92. Body very flat and thin, its height being contained $2\frac{3}{4}$ times in the total length (without caudal), the length of the head $4\frac{1}{8}$ times. The upper part of the snout is dilated, bent downward like an aquiline nose, the end covering the symphysis of the mandibles; the cleft of the mouth is curved, the lower eye being immediately above its angle. The eyes are mere points, rather distant from

each other. The gill opening is reduced to a very small slit, the gill membrane being attached to the sides of the throat. The dorsal fin commences on the extremity of the snout and terminates at the root of the caudal, its rays are simple, and each is accompanied by a series of very small ctenoid scales; the longest rays are not quite ½ as long as the head, and occupy the middle and the third quarter of the fin. Caudal quite free, as long as the head, somewhat pointed. The right ventral appears as a mere continuation of the anal; the left is reduced to 2 minute rays near the vent. The scales on both sides are ctenoid, those on the neck and on the chin being twice the size of those on the body. Color uniform brownish gray. Coast of Surinam and Brazil. The above description from Günther, taken from the type of *Soleotalpa unicolor*. A specimen (No. 4677, M. C. Z.), from Obydos, Brazil, examined by us, differs in coloration, being pale brown, the body and fins profusely covered with round, dark spots of varying sizes, the largest as wide as from eye to eye. Head 4⅓; depth 2⅔. D. 78; A. 56; scales 100; V. 2. Eyes reduced to points, the upper in advance of lower, near middle of length of head; gill openings small, subequal; right ventral beginning at the chin, continuous with anal; dorsal and anal slightly connected with caudal. Steindachner gives D. 72; A. 53; scales 95. Color brownish, mottled with darker spots. Probably Günther's specimen is faded. (*unicolor*, one-colored.)

Apionichthys dumerili, KAUP, Archiv fur Naturgesch. 1858, 104, no locality; no description.

Soleotalpa unicolor, GÜNTHER, Cat., IV, 489, 1862, West Indies. (Coll. Scrivener.)

Apionichthys nebulosus, PETERS, Berliner Monatsberichte 1869, 709, Surinam.

? Apionichthys bleekeri, HORST, Nederl. Tydschr. Dierk., IV, 30, 1878, locality unknown; specimen in Mus. Utrecht. (Description not seen by us.)

Apionichthys unicolor, JORDAN, Proc. U. S. Nat. Mus. 1886, 603.

Apionichthys dumerili, BLEEKER, Nederl. Tydschr. voor Dierkunde, II, 1865, 305; STEINDACHNER, Ichth. Beitr., VIII, 48, 1878.

1056. GYMNACHIRUS, Kaup.

Gymnachirus, KAUP, Uebersicht der Soleinæ, Archiv fur Naturgesch. 1858, 101 (*nudus*).

This genus differs from *Achirus* in the absence of scales; the dorsal and anal are free from the caudal. Brazil. (*γυμνός*, naked; *Achirus*.)

3077. GYMNACHIRUS FASCIATUS, Günther.

Head 4½; depth 1½. D. 68; A. 50; pectoral of right side present, very small, of 2 rays, ⅓ length of eye; jaws hidden in thick skin; lips and left side of head covered with fringes. Gill opening not extending upward as far as pectoral; vertical fins in thick skin. Olive, with 14 brown cross bands as broad as the interspaces, all extending on dorsal and anal, the first across snout, the second and third across eye; caudal with 3 brown bands. (Günther.) Locality unknown, probably Surinam or Brazil; a related species (*G. nudus*, Kaup; no pectoral fins. D. 51; A. 42), being described from Bahia. (*fasciatus*, banded.)

Gymnachirus fasciatus, GÜNTHER, Cat., IV, 488, 1862, locality unknown; JORDAN & GOSS. Review Flounders and Soles, 317, 1889.

1057. SYMPHURUS,* Rafinesque.

(Tongue-Fishes.)

Symphurus, Rafinesque, Indice d'Ittiologia Siciliana, 52, 1810 (*nigrescens*).
Bibronia, Cocco, Alcuni Pesci del mare di Messina, 15, 1844 (*ligulata;* larval form).
Plagusia, Cuvier, Règne Animal, Ed. 2, II, 344, 1829 (based on *Plagusia* of Brown); name preoccupied in Crustaceans, Latreille, 1806.
Plagiusa, Bonaparte, Catalogo Metodico, 51, 1846 (*lactea*); substitute for *Plagusia* preoccupied.
Aphoristia, Kaup, Archiv fur Naturgesch. 1858, 106 (*ornata*).
Glossichthys, Gill, Cat. Fish. E. Coast N. A., 51, 1861 (*plagiusa*).
Ammopleurops, Günther, Cat., IV, 490, 1862 (*lacteus = nigrescens*).
? Bascanius, Schiödte, Naturhist. Tydsskr., V, 269, 1867 (*tœdifer;* larval form).
Acedia, Jordan, in Jordan & Goss, Review Flounders and Soles, 321, 1889 (*nebulosus*).

Body elongate, more or less lanceolate in outline, with the eyes and color on the left side; eyes small, very close together, with no distinct interorbital ridge between them; mouth small, twisted toward the blind side; teeth little developed, in villiform bands; edge of preopercle covered by the scales; gill openings narrow, the gill membranes adnate to the shoulder girdle above, joined together and free from the isthmus below; pectoral fins wanting (in the adult); vertical fins more or less confluent; scales ctenoid; lateral line wanting. Ventral fin of eyed side only present, free from the anal; head without fringes. (*σὺν*, together; *φύω*, to grow; *οὐρά*, tail; from the united vertical fins.)

Symphurus:

a. Scales not minute, ctenoid, 65 to 105 in number; dorsal rays 86 to 100; anal rays 70 to 87.

b. Scales rather large, about 65; head 4⅓; depth 4⅓; color, clouded brown. piger, 3078.

bb. Scales small, moderately ctenoid, 75 to 105 in a longitudinal series.

c. Dorsal and anal pale anteriorly, becoming more or less abruptly black posteriorly.

d. Caudal fin abruptly pale, at least at tip.

e. Body elongate, depth 4½ in length; head 5½. D. 96 to 100; A. 86 or 87; scales 88 to 90. Color, grayish, speckled with brown; dorsal and anal fins black on last tenth, the caudal abruptly pale; tips of fin rays vermilion. marginatus, 3079.

ee. Body deeper, the depth 3⅓ to 3⅝ in length.

f. Color, light brown, irregularly barred and marbled with darker; dorsal and anal with 3 to 6 inky blotches posteriorly. D. 92 to 95; A. 75 to 78. atramentatus, 3080.

---

* We follow Jordan & Goss in using the name *Symphurus* instead of *Aphoristia*, as the so-called *Ammopleurops lacteus* is a genuine member of the latter genus, and as it seems to be evident that the latter species is the original of *Symphurus nigrescens* of Rafinesque. The following is Rafinesque's description: "III. Gen. *Symphurus*. Ala caudale acuta, e riunita all' ale dorsali, ed anali, occhj alla sinistra. *Osserv.* Si dovranno ragguagliare in questo genere due specie del genere *Achirus* di Lacepede, cioè gli *A. bilineatus*, e *A. ornatus*. Sp. no. 44. *Symphurus nigrescens*. Nerastro senza fascie, allungato, una sola linea laterale da ogni lato."

This single lateral line assumed to distinguish *Ammopleurops* from *Aphoristia* is not a real lateral line, but a depression along the median line produced by the junction of the muscles. The species of *Symphurus* are somewhat numerous and very closely allied. With the exception of the European *Symphurus nigrescens*, all of them are American. The development of the species is imperfectly known. According to Giglioli, the larvæ called *Bibronia*, may belong to this genus, and so possibly may *Charybdia*. The name *Plagusia* belongs properly to the present genus rather than to the type of *Plagusia bilineata*, to which it has been restricted by Kaup and Günther. It is, however, preoccupied in crustaceans, and in any case, both *Plagusia* and the substitute name *Plagiusa* are antedated by the name *Symphurus*.

ff. Color, light olive, with numerous roundish brownish black spots much larger than eyes, dorsal and anal black, with narrow white margin. D. 94; A. 77; scales 95.
FASCIOLARIS, 3081.

dd. Caudal fin black, as is a large part of dorsal and anal, the black either continuous or in the form of large spots. Color brownish, often mottled, usually with more or less distinct darker cross bands and with longitudinal streaks along the rows of scales, sometimes nearly plain brown.

g. Scales quite small, 98 to 105.

h. Body decidedly elongate, the depth about $4\frac{2}{3}$ in length. D. 97; A. 82; scales 98. ELONGATUS, 3082.

hh. Body less elongate, the depth $3\frac{1}{3}$ in length; head $5\frac{1}{3}$; longitudinal streaks very distinct. D. 100; A. 80; scales 105. ATRICAUDUS, 3083.

gg. Scales rather larger, 75 to 90.

i. Body rather elongate, the depth $3\frac{2}{3}$ to $4\frac{1}{3}$; dorsal rays 80 to 90; anal 80 to 85; opercular flap large; body with 3 or 4 dark cross bands. LEEI, 3084.

ii. Body less elongate, the depth $3\frac{1}{10}$ to $3\frac{2}{3}$ in length; the head $5\frac{1}{3}$ to $5\frac{2}{3}$. D. 90 to 95; A. 75 to 80.
PLAGUSIA, 3085.

cc. Dorsal and anal pale throughout, or more or less mottled or spotted with darker, the caudal similarly colored, not distinctly black. Body not very elongate, the depth 3 to $3\frac{1}{3}$ in length. (Probably all varieties of *S. plagiusa*.)

j. Body with dark cross bands more or less distinct; the fins mottled or speckled; upper eye slightly in advance of lower.

k. Dorsal rays 86 to 95; anal rays 75 to 80; head 5 in length; depth $3\frac{1}{4}$; scales 85 to 93; cross bands more distinct than in related species. PLAGIUSA, 3086.

kk. Dorsal rays 78 to 85; anal rays 70 to 72; head 5 in length; depth $3\frac{1}{3}$; scales 80 to 90. Color, light brown, with darker cross bars, which become obsolete with age. PUSILLUS, 3087.

jj. Body uniform grayish, without cross bands; last part of dorsal and anal with 3 or 4 oblong black blotches, each somewhat larger than the eye; upper eye directly above lower; head $5\frac{1}{3}$ in length. Scales 85; D. 92; A. 75. DIOMEDEANUS, 3088.

ddd. Caudal and posterior part of dorsal and anal not black, scarcely darker than anterior part; scales 92; D. 93; A. 73. WILLIAMSI, 3089.

ACEDIA (Spanish name of *Symphurus plagusia* at Havana):

aa. Scales very small, ctenoid, each with a median dark streak, which simulates a keel, but is not a ridge; snout and jaws naked; fin rays in increased number.

l. Head $5\frac{2}{3}$; depth $4\frac{2}{3}$. D. 119; A. 107; scales 120. Grayish, everywhere mottled with brown. NEBULOSUS, 3090.

Subgenus SYMPHURUS.

3078. SYMPHURUS PIGER (Goode & Bean).

Head $4\frac{1}{3}$ in total length; depth $3\frac{1}{7}$. D. 90; A. 69 to 75; ventral 4; scales 65–34 (transverse); eye 6 in head; snout $4\frac{1}{2}$; mouth oblique, curved, its angle below middle of lower eye; teeth feeble, closely placed, a little stronger on colored side; nostril tubular, a little nearer eye than tip of snout; eyes moderate in size, very close together, the upper very slightly in advance, its distance from the dorsal outline equal to its diameter; scales large, ctenoid, deciduous. Dorsal begining over middle of upper eye; longest dorsal and anal rays 3 in depth of body; pectorals obsolete.

Color grayish and brownish, with a submetallic luster upon the scales when examined separately; the denticulations of the scales dark and prominent, giving a clouded general aspect; some of the smaller specimens with large, irregular, brownish blotches above, and a dark subcircular blotch near the root of the tail, its diameter twice eye; colorless below. (Goode & Bean.) West Indies and Gulf of Mexico, in deep water; a well-defined species. (*piger*, sluggish.)

Aphoristia pigra, GOODE & BEAN, Bull. Mus. Comp. Zool., XII, 5, 154, 1886, St. Kitts, in about 250 fathoms (Coll. *Blake*); GOODE & BEAN, Oceanic Ichthyology, 460, 1896.
Symphurus piger, JORDAN & GOSS, Review Flounders and Soles, 326, 1889.

3079. SYMPHURUS MARGINATUS (Goode & Bean).

Head 5½ in total length; depth 4½. D. 96 to 100; A. 86 or 87; ventral 4; scales 88 to 90; eye 4½ in head; snout 4½. Body slender, lanceolate; mouth moderate, oblique, curved, its angle below front of pupil of upper eye; dentition feeble; eyes moderate, close together, the upper very slightly in advance; nostril in a long slender tube, midway between lower eye and tip of snout; scales moderate, strongly and sharply denticulate, not keeled; origin of dorsal above posterior margin of upper eye; anal scarcely so high as dorsal; median caudal rays short. Color in life, reddish gray, much speckled with brown; belly bluish gray; bases and membranes covering fin rays dark brown; caudal abruptly pale; tips of dorsal and anal rays and some of the membrane covering caudal rays vermilion. West Indies, in deep water. Described from a specimen, 102 mm. in length, collected by the *Blake* at Station CLXXXI, in 321 fathoms. (Goode & Bean.) (*marginatus*, edged.)

Aphoristia marginata, GOODE & BEAN, Bull. Mus. Comp. Zool., XII, No. 5, 153, 1886, off St. Vincent (Coll. *Blake*); GOODE & BEAN, Oceanic Ichthyology, 459, fig. 376, 1896.
Symphurus marginatus, JORDAN & GOSS, Review Flounders and Soles, 323, 1889.

3080. SYMPHURUS ATRAMENTATUS, Jordan & Bollman.

Head 4⅜ to 5; depth 3⅛ to 3⅜. D. 92 to 95; A. 75 to 78; scales 95 to 100, 38 in a cross series. Body more elongate than in *S. atricaudus*. Eyes larger, the upper in advance of lower, vertical diameter of each 3½ to 4 in head. Cleft of mouth somewhat more curved than in *S. atricaudus*, otherwise similar. Scales larger than in *S. atricaudus;* spines on posterior margin not so strong. Ventral fins (measured from angle of gill opening) 2⅜ to 3 in head. Color light brown, irregularly barred and marbled with darker; several irregular grayish bars most distinct on posterior parts, a distinct narrow, dark bar behind gill opening; anterior part of dorsal and anal fin pale, posterior dark; anterior part with 4 to 7 dusky oblique areas, posterior part with 3 to 6 roundish inky-black spots; caudal black, narrowly tipped with white; each scale with a narrow dark edge. Length about 4½ inches. Pacific Ocean off Colombia, in water of moderate depth; common. Related to *Symphurus atricaudus* (Jordan & Gilbert), but distinguished by having 3 to 6 black oblong blotches on posterior part of dorsal

and anal; the general coloration darker; the scales and eyes larger. (*atramentatus*, inked.)

Symphurus atramentatus, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 177, off coast of Colombia, at Albatross Station 2795, Lat. 7° 57′ N., Long. 78° 55′ W., in 33 fathoms. (Type, 41157, U. S. Nat. Mus. Coll. *Albatross.*)

3081. SYMPHURUS FASCIOLARIS, Gilbert.

Depth 3⅔ in length; head 5⅓. D. 94; A. 77; scales 95. Eye small, 7 in head; cleft of mouth reaching to below middle of lower eye. Color light olive, with numerous roundish brownish-black spots much larger than eye, the largest arranged in 5 vertical dusky cross bars, the spots being connected by a darker ground color; a vertical dusky streak through eye; a wide dusky cross bar, bounded by darker lines on cheeks; dorsal and anal posteriorly black, with narrow white margin; caudal jet-black, with white edge; ventral white. Gulf of California, where several specimens were dredged by the *Albatross*, in shallow water. (Gilbert.) (*fasciolaris*, with narrow bands.)

Symphurus fasciolaris, GILBERT, Proc. U. S. Nat. Mus. 1891, 566, Gulf of California. (Coll. Dr. Gilbert.)

3082. SYMPHURUS ELONGATUS (Günther).

Head 5⅓ in body; depth 4⅔. D. 97; A. 82; scales 98 to 105; eye 10 or 11 in head; gape of mouth 3½; caudal 2½. Body extremely elongate; mouth strongly curved, reaching past lower eye; eyes in contact, the upper in advance; opercle vertical behind, devided into 2 convex flaps by a concave portion, its upper end hardly reaching axis of body; scales not keeled, ctenoid. Pectorals obsolete; dorsal beginning above eye; rays of dorsal and anal short, subequal, the fins confluent with the caudal, which ends in a sharp point; ventral of blind side obsolete, that of eyed side on the body ridge, separated from the anal. Color brownish, often mottled, usually with more or less distinct darker cross bands, and with longitudinal streaks along the rows of scales, sometimes nearly plain brown; caudal fin black, as is a large part of the dorsal and anal, the black either continuous or in the form of large spots. Pacific coast of Central America; not rare. Here described from a specimen, 6 inches long, from *Albatross* Station 2804, in Panama Bay, in 47 fathoms. (*elongatus*, elongate.)

Aphoristia ornata, var. *elongata*, GÜNTHER, Fishes Centr. Amer., 473, 1869, Panama.
Aphoristia elongata, JORDAN & GILBERT, Bull. U. S. Fish. Comm. 1882, 111.
Symphurus elongatus, JORDAN & GOSS, Review Flounders and Soles, 323, 1885.

3083. SYMPHURUS ATRICAUDUS (Jordan & Gilbert).

(SAN DIEGO SOLE.)

Head 5⅓; depth 3⅓. D. 100; A. 80; scales 105. Body oblong-lanceolate, anteriorly somewhat blunt, regularly narrowed behind and ending in a point; the snout rather abruptly truncate; eyes and color on the left side. Eyes very small, nearly even behind, the upper eye the larger and extending farther forward. A single nostril in front of interorbital space, and

apparently a single smaller one below it. Mouth moderate, extending to opposite eye, somewhat turned toward eyed side; lips large, not fringed, the upper with a small black papilla in advance of lower eye, this apparently normal, but it may be a detached piece of skin, hardened by the alcohol; upper jaw scarcely produced, not forming a hook. Teeth small, on blind side only, the edge of the jaw on eyed side forming a smooth ridge. Gill openings narrow, not extending up to level of mouth. Scales very small, ctenoid, pretty regular over the body, much smaller on the head, the rows of scales rendered very distinct by black dots, the stripes converging toward the snout; scales on the 2 sides of the body similar; no lateral line on either side; about 105 scales (100 to 110) in a longitudinal series from the head to the tail, 45 to 50 in cross series. Dorsal fin beginning on head, continuous with anal around the tail; ventral fin of colored side only present, nearly on ridge of abdomen, and separated from the anal by an interval $\frac{1}{2}$ longer than cleft of mouth; rays of middle parts of dorsal and anal fins with a fleshy border at base on blind side. Coloration brownish olive, with vertical dark half bars, irregular in size and position, some of them coming down from the back and others up from the belly, these posteriorly nearly meeting, but anteriorly alternating; streaks of dark points along the rows of scales, these forming very distinct longitudinal streaks; posterior part of dorsal and anal broadly edged with black; right side plain white. San Diego to Cape San Lucas, in sandy bays; common in the bay of San Diego, in which locality the numerous specimens before us were taken. A small specimen, $1\frac{1}{2}$ inches long, with light spots on the colored side and a pale ocellation on the black of the tail, taken by Mr. Lyman Belding near Cape San Lucas, probably belongs to the same species. (*ater*, black; *cauda*, tail.)

Aphoristia atricauda, Jordan & Gilbert, Proc. U. S. Nat. Mus. 1880, 23, San Diego; Jordan & Gilbert, Synopsis, 842, 1883; Jordan & Gilbert, Proc. U. S. Nat. Mus. 1882, 380; Jordan, Proc. U. S. Nat. Mus. 1886, 54.

Symphurus atricauda, Jordan & Goss, Review Flounders and Soles, 324, 1889.

3084. SYMPHURUS LEEI, Jordan & Bollman.

Head 4 to $4\frac{1}{2}$ ($4\frac{1}{3}$ to $4\frac{2}{3}$); depth $3\frac{4}{5}$ to 4 ($4\frac{1}{5}$ to $4\frac{1}{3}$). D. 95 to 100; A. 80 to 85; scales 80 to 90, 35 to 38 in a cross series; ventrals $3\frac{1}{3}$ to $3\frac{1}{2}$ in head. Body more elongate than in *S. atricaudus* or *S. atramentatus*, approaching that of *S. elongatus;* outline of under part of head more oblique than in the other Pacific coast species. Eyes larger than in the preceding species, the upper in advance of lower, their vertical diameter 5 to $5\frac{1}{2}$ in head; cleft of mouth extending slightly farther back than in *S. atricaudus* or *atramentatus*, but not beyond eye as in *S. elongatus;* maxillary reaching posterior border of eye, $3\frac{2}{3}$ to 4 in head; snout $5\frac{1}{3}$ to $5\frac{2}{3}$ in head. Scales comparatively large, not so firmly embedded as in *S. atricaudus* or *atramentatus*, those on opercles rather large. Opercular flap larger than in other Pacific species. Color light brown, speckled with darker, and with 3 or 4 broad black cross bands, width of median bands $2\frac{1}{2}$ to 3 in head, the posterior band widest; caudal and the posterior $\frac{2}{3}$ of the dorsal and anal

black; no black spots on dorsal; scales thickly punctulate, but with no distinct darker edgings. Related to *Symphurus atricaudus* (Jordan & Gilbert), but the body with 4 wide black cross bands, and the form more elongate. Bay of Panama. Many specimens of this species were obtained at *Albatross* Station 2804, at a depth of 47 fathoms. It is evidently very different from *S. atramentatus*, and needs comparison only with *S. elongatus*, from which it seems to be sufficiently distinct. Length of type 4½ inches. (Named for Prof. Leslie A. Lee and Mr. Thomas Lee, naturalists on board the *Albatross* when the species was discovered.)

Symphurus leei, JORDAN & BOLLMAN, Proc. U. S. Nat. Mus. 1889, 178, Lat. 8° 16′ 30″ N. Long. 79° 37′ 45″ W. (Type, No. 41134. Coll. Prof. L. A. Lee and Mr. Thomas Lee.)

3085. SYMPHURUS PLAGUSIA* (Bloch & Schneider).

(ACEDIA.)

Head 5⅛ to 5⅔; depth 3 1/10 to 3⅔ in length. D. 90 to 95; A. 75 to 80; scales 75 to 85. Body rather elongate. Color brownish, often mottled, usually with more or less distinct darker cross bands, and with longitudinal streaks along the rows of scales, sometimes nearly plain brown; caudal black, including a large part of dorsal and anal, the black continuous as in the form of spots. West Indies to Brazil; Cuba to Rio Janeiro; common. The numerous specimens of this species examined by us are from Havana, Pernambuco, Santos, Rio Janeiro, Curuça, and Victoria. (*plagusia*, an old name, from πλάγιος, oblique.)

Plagusia, BROWNE, Jamaica, 445, No. 1, 1756, Jamaica.
Pleuronectes plagusia, BLOCH & SCHNEIDER, Syst. Ichth., 162, 1801, Jamaica; after BROWNE.
Achirus ornatus, LACÉPÈDE, Hist. Nat. Poiss., IV, 659, 1803, on a specimen "presented by Holland to France."
Plagusia tessellata, QUOY & GAIMARD, Voyage Uranie, Zoologie, 240, 1824, Rio Janeiro.
Plagusia brasiliensis, AGASSIZ, Spix, Pisc. Brasil., 89, tab. 50, 1827, Brazil.

* The synonymy of this species is somewhat doubtful. The original type of *Pleuronectes plagiusa* was sent to Linnæus by Dr. Garden, of Charleston. It would therefore appear probable that this specimen represented the species of this genus which is found on the Carolina coast. But this typical specimen is still preserved in the rooms of the Linnæan Society in London, where it has been examined by Goode and Bean. From their notes (Proc. U. S. Nat. Mus. 1885, 196) we quote:

"The type of this species may have come from Africa or India. There is considerable doubt as to its origin. (See Garden's Correspondence with Linné, p. 314.) D. ca 92, A. ca 80; scales 77. The species is more elongate than our specimens of *Aphoristia plagiusa*, so called, the depth being contained in the total length without caudal 4⅓ times and the head 6 times."

As, however, no species of this genus are yet known from Africa or India, it is rather probable that Garden's fish actually came from Charleston. The greater slenderness of the original type is perhaps due to distortion, and the smaller number than usual of the scales does not afford a marked distinction. The name *Achirus ornatus* is also doubtful in its proper application. The only thing distinctive in the description of Lacépède is that the typical specimen was "given by Holland to France." Many of the species in this Dutch collection seem to have come from Surinam, and this is probably no exception. But Lacépède's description might apply as well to any other species of *Symphurus* as to this. The name *Pleuronectes plagusia*, given by Schneider to the species described by Browne, seems to admit of no doubt, as this is the only one of the group yet known from Jamaica. If, therefore, the name *Symphurus plagiusa* be used for the northern species, or dropped altogether as not identified, the present species will stand as *Symphurus plagusia*. We have compared numerous specimens from Rio Janeiro (representing the nominal species *tessellatus* or *brasiliensis*) with others (*plagusia*=*ornata*) from Havana. There is certainly no permanent difference. The Brazilian specimens are a little more slender on an average, but there are numerous exceptions, and all variations in color are found in both.

Aphoristia ornata, KAUP, Archiv fur Naturgesch. 1858, 106; GÜNTHER, Cat., IV, 490, 1862; POEY, Synopsis, 409, 1868; POEY, Enumeratio, 140, 1875; KNER, Novara Fische, III, 292; D. 90; A. 75; depth 3¼ in length.

Aphoristia plagiusa, JORDAN, Proc. U. S. Nat. Mus. 1886, 53; not *S. plagiusa* of this paper.

Symphurus plagusia, JORDAN & GOSS, Review Flounders and Soles, 324, 1889.

3086. SYMPHURUS PLAGIUSA* (Linnæus).

(TONGUE FISH.)

Head 5; depth 3 to 3⅓. D. 86 to 95; A. 75 to 80; scales 85 to 93. Body not very elongate. Body grayish, with dark cross bands more distinct than in related species; dorsal and anal more or less mottled or spotted with darker; caudal similarly colored, not distinctly black. South Atlantic and Gulf coasts of the United States, from Cape Hatteras to Pensacola and Key West, replacing *S. plagusia* northward, the species as similar as the two names; very common on the sandy shores of our South Atlantic and Gulf States. Our numerous specimens are from Beaufort, Charleston, Pensacola, and Key West. Those from Key West nearly plain gray, as would be expected in fishes taken from the coral sands. (πλάγιος, oblique.)

Pleuronectes plagiusa, LINNÆUS, Syst. Nat., Ed. XII, 455, 1766, on a specimen from Dr. Garden, probably from Charleston, but the locality not quite certain; and of various copyists.

Plagusia fasciata, HOLBROOK MS., DE KAY, New York Fauna: Fishes, 304, 1842, Charleston.

Glossichthys plagiusa, GILL, Cat. Fish East Coast N. Am., 51, 1861.

Plagusia plagiusa, GILL, Cat. Fish. East Coast N. Am., 794, 1873.

Aphoristia plagiusa, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1878, 368; JORDAN, *l. c.*, 1880, 22; JORDAN & GILBERT, *l. c.*, 1882, 305 and 618; JORDAN & GILBERT, Synopsis, 842, 1883; JORDAN, Proc. U. S. Nat. Mus. 1884, 144.

Aphoristia fasciata, JORDAN, Proc. U. S. Nat. Mus. 1886, 53.

Symphurus plagiusa, JORDAN & GOSS, Review Flounders and Soles, 325, 1889.

3087. SYMPHURUS PUSILLUS (Goode & Bean).

Head 5 in total length; depth 3½. D. 78; A. 70; scales 85 to 90–35 (transverse); eye 5½ in head; snout 5½; length of gape of mouth 4⅜. Body slender, lanceolate; mouth small, oblique, curved, its angle under anterior margin of pupil of lower eye; dentition feeble; eyes small, close together, in the same vertical line; tubular nostril midway between lower eye and tip of snout; scales small, strongly ctenoid; jaws and snout scaled. Dorsal beginning above middle of eye, its highest rays 2⅜ in depth of body; greatest height of anal 3; median caudal rays short; ventrals well separated from anal. Color light brown, with 6 or 7 cross bars of slightly

* A specimen of *Symphurus*, nearly 6 inches long, collected at Beaufort, North Carolina, by Prof. O. P. Jenkins, seems referable to *Symphurus pusillus* rather than to the typical *plagiusa*. It is highly mottled in coloration, the body and fins being profusely speckled and blotched with blackish, besides 9 or 10 rather distinct cross bands. D. 85; A. 72; scales about 80. Depth 3¼ in length. Another large specimen, 7 inches long, from the Florida Keys, is in the museum at Cambridge. This has: D. 82; A. 72; scales 76. Depth 3 in length. Color brown, almost plain, except that the fins are mottled, especially posteriorly; caudal fin not black. If these two specimens are really typical of *Symphurus pusillus*, it probably can not be separated as a species from *S. plagiusa*.

darker hue; fins pale, with dusky blotches; blind side white. (Goode & Bean.) Gulf Stream, in deep water. Very close to *Symphurus plagiusa*. (*pusillus*, weak.)

Aphoristia pusilla, Goode & Bean, Proc. U. S. Nat. Mus. 1885, 590, Gulf Stream, Lat. 40° N., in deep water; Goode & Bean. Oceanic Ichthyology, 461, fig. 379, 1896.
Symphurus pusillus, Jordan, Proc. U. S. Nat. Mus. 1889, 651.

3088. SYMPHURUS DIOMEDEANUS (Goode & Bean).

Head 5⅜ in body; depth 3½. D. 96 (including ½ of caudal); A. 79; scales 85; eye 6 in head; snout 5; caudal 10 in total length. Mouth oblique, curved, its angle below front of eye, teeth very feeble; nostril tubular, nearer eye than tip of snout; eyes moderate, equal, very close together, upper eye directly over the lower; scales moderate, somewhat deciduous, ctenoid; jaws and snout with small thin scales. Origin of dorsal above middle of upper eye, highest rays 3½ times depth of body; ventrals well separated from the anal. Color uniform gray, lighter below, the scales above somewhat metallic in luster; the last fourth of dorsal with 3 oblong black blotches somewhat larger than eye, the anal with 4, similar in position; in the young there is a slight brownish marginal line upon each scale, and an appearance of indistinct cloudings of brown upon the colored side. Off Trinidad and Dominica and in the Gulf of Mexico. The specimen here described was collected by the *Albatross* at Station 2414, in the Gulf of Mexico, north of the Tortugas, at a depth of 26 fathoms; its length is 140 mm. Other specimens were dredged by the *Albatross* at Station 2362, in Lat. 22° 08′ 30″ N., Long. 86° 53′ 30″ W., in 25 fathoms, and at Stations 2121 and 2122, between Lat. 10° 37′ 40″ N., Long. 61° 42′ 40″ W., and Lat. 10° 37′ N., Long. 61° 44′ 22″ W., in 31 to 34 fathoms. Specimens were also secured by the *Blake* at Stations XXIV and XXV, off Dominica. (Goode & Bean.) Evidently very close to *Symphurus plagiusa*. (*Diomedea*, the Albatross; from the name of the steamer by which most of the deep-sea explorations of the United States Fish Commission have been accomplished.)

Aphoristia diomedeana, Goode & Bean, Proc. U. S. Nat. Mus. 1885, 589, Gulf of Mexico, Lat. 25° 04′ 30″ N., Long. 82° 59′ 15″ W. (Type, No. 37347. Coll. *Albatross*); Goode & Bean, Oceanic Ichthyology, 460, fig. 378, 1896.

3089. SYMPHURUS WILLIAMSI, Jordan & Culver.

Head 4⅘; depth 3⅜. D. 93; A. 73; scales 92. Body more slender than in *S. plagiusa*, which it much resembles, but less slender than *S. elongatus;* upper eye slightly in advance of lower. Sand color in life; light gray, everywhere finely mottled with light and dark; traces of a few very narrow dark cross bands; fins all mottled; caudal and posterior part of dorsal and anal not black, scarcely darker than anterior part. Known only from Mazatlan, where 2 specimens, the larger about 1½ inches long, were obtained by Mr. T. M. Williams, in tide pools with sandy bottom in very shallow water near the estuary. (Named for Thomas Marion Williams, a student in biology in Stanford University, discoverer of the species.)

Symphurus williamsi, Jordan, Proc. Cal. Ac. Sci. 1895, 506, pl. 55, Mazatlan. (Coll. Hopkins Exped. to Mazatlan.)

Subgenus ACEDIA, Jordan.

3090. SYMPHURUS NEBULOSUS (Goode & Bean).

Head $5\frac{2}{3}$ in total length; depth $4\frac{2}{3}$. D. 119 (to middle of base of caudal); A. 107; V. 5; scales 120; eye $7\frac{1}{2}$ in head; snout 5. Body slender; angle of mouth below front of lower pupil; teeth feeble, very slender, and rather closely placed, apparently equally developed on both sides; eyes small, close together, separated by a single row of scales, the upper one very slightly in advance; tubular nostril nearer eye than tip of snout; scales small, ctenoid, each with a median dark streak (but not keeled, as erroneously stated in the original description);* jaws and snout naked. Origin of dorsal a little behind eyes, highest rays 3 in depth of body; longest anal rays twice length of snout; median caudal rays longest, twice length of snout; pectorals obsolete; ventrals well separated. Color grayish, everywhere mottled with brown; a dark median line on scales. (Goode & Bean.) Gulf stream. A well-marked species. The increased number of fin rays indicates a probability that the number of vertebræ will also be found similarly increased. (*nebulosus*, clouded.)

Aphoristia nebulosa, Goode & Bean, Bull. Mus. Comp. Zool., x, No. 5, 192, 1883, Gulf Stream, off the coast of Carolina; Goode & Bean, Oceanic Ichthyology, 458, fig. 375, 1896.

Symphurus nebulosus, Jordan & Goss, Review Flounders and Soles, 326, 1889.

Order CC. PEDICULATI.

(The Pediculate Fishes.)

Carpal bones notably elongate, forming a kind of arm (pseudobrachium) which supports the broad pectoral. Gill opening reduced to a large or small foramen situated in or near the axil, more or less posterior to the pectorals. Ventral fins jugular if present; anterior dorsal reduced to a few tentacle-like, mostly isolated spines; soft dorsal and anal short; no scales. First vertebra united to cranium by a suture; epiotics united behind supraoccipital; elongate basal pectoral radii (actinosts) reduced in number; no interclavicles; post-temporal broad, flat, simple; upper pharyngeals 2, similar, spatulate, with anterior stem and transverse blade; basis of cranium simple; no air duct to the swim bladder. Marine fishes, chiefly of the tropics and the oceanic abysses. The group is an offshoot from the *Acanthopteri*, its chief modifications being in the elongation of the actinosts and in the position of the gill opening. Its nearest relatives among the spiny-rayed fishes are, perhaps, the *Batrachoididæ*. (*pediculatus*, having a footstalk.)

Analysis of Families of Pediculati.

a. Gill openings in or behind the lower axil of the pectoral; mouth large, terminal.

b. Pseudobranchiæ present; pseudobrachia with 2 actinosts; head broad, depressed, the enormous mouth with very strong teeth; ventrals present.

Lophiidæ, ccxxi.

* The appearance of "keeled scales," described by Goode & Bean, is due to a black line on the skin under the center of each row of scales. There seems to be no real keel, and the species is congeneric with the other species of *Symphurus*.

bb. Pseudobranchiæ none; pseudobrachia with 3 actinosts.

c. Ventrals present; arm angulate, the pseudobrachia elongate. ANTENNARIIDÆ, CCXXII.

cc. Ventrals wanting, arm not angulate, the pseudobrachia moderate. CERATIIDÆ, CCXXIII,

aa. Gill openings in or behind the upper axil of the pectoral; mouth small, usually inferior. OGCOCEPHALIDÆ, CCXXIV.

Family CCXXI. LOPHIIDÆ.

(THE ANGLERS.)

Head wide, depressed, very large. Body contracted, conical, tapering rapidly backward from the shoulders. Mouth exceedingly large, terminal, opening into an enormous stomach; upper jaw protractile; maxillary without supplementary bone; lower jaw projecting; both jaws with very strong, unequal, cardiform teeth, some of the teeth canine-like, most of them depressible; vomer and palatines usually with strong teeth. Gill openings comparatively large, in the lower axil of the pectorals. Pseudobranchiæ present. Gill rakers none. Gills 3. Skin mostly smooth, naked, with many dermal flaps about the head. Spinous dorsal of 3 isolated, tentacle-like spines on the head, and 3 smaller ones behind, which form a continuous fin; second dorsal moderate, similar to the anal; pectoral members scarcely geniculated, each with 2 actinosts and with elongate pseudobrachia; ventrals jugular, I, 5, widely separated, large, much enlarged in the young. Young with the head spinous. Pyloric cæca present. Two genera, with 4 or 5 species, living on sea bottoms, at moderate or great depths; remarkable for their great voracity. (*Pediculati*, part, genus *Lophius*, Günther, Cat., III, 178–182, 1861.)

a. Vertebræ 27 to 31. LOPHIUS, 1058.

aa. Vertebræ 18 or 19 only. LOPHIOMUS, 1059.

1058. LOPHIUS (Artedi) Linnæus.

(FISHING-FROGS.)

Lophius (ARTEDI) LINNÆUS, Syst. Nat., Ed. X, 1, 236, 1758 (*piscatorius*).

Characters of the genus included above. Vertebræ numerous, about 30 in number. (*Lophius*, the ancient name of *L. piscatorius*, from λόφος, a crest.)

3091. LOPHIUS PISCATORIUS, Linnæus.

(COMMON ANGLER; FISHING-FROG; MONKFISH; GOOSEFISH; ALL-MOUTH; BELLOWS-FISH.)

D. I–I–I, III–10; A. 9. Body depressed, tapering, scarcely longer than head. Humeral spine with 3 points, of which the posterior is the longest. Head surrounded with a fringe of barbels; top of head, in young, with many strong spines. Anterior dorsal spine elongate, fleshy at tip. Brownish, mottled, below white; mouth behind the hyoid bone immaculate; pectorals and caudal black at tip; peritoneum black. Length 3 feet. North

Atlantic, on both coasts; generally common, ranging southward along the shore to Cape Hatteras; found in deep water as far south as Barbados, in 209 fathoms, and to the Cape of Good Hope; northward to Norway and Nova Scotia. A well-known fish of singular ugliness of appearance, and of enormous voracity. (Eu.) (*piscatorius*, pertaining to an angler, in allusion to the baited dorsal spines which overhang the cavernous mouth.)

Lophius piscatorius,* LINNÆUS, Syst. Nat., Ed. x, 1, 236, 1758, seas of Europe; after ARTEDI, *Lophius ore cirrhoso*, etc.; GÜNTHER, Cat., III, 179, 1861; GILL, Proc. U. S. Nat. Mus. 1878, 219; JORDAN & GILBERT, Synopsis, 844, 1883.

Lophius americanus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 380, 1837, Philadelphia (Coll. Le Sueur); STORER, Hist. Fish. Mass., pl. 18, fig. 2, 101, 1867.

1059. LOPHIOMUS, Gill.

Lophiomus, GILL, Proc. U. S. Nat. Mus. 1882, 552 (*setigerus*).

This genus is closely allied to *Lophius* in external characters, but it is strikingly distinguished by the reduced number of its vertebræ, which are only 18 or 19, a fact which is associated with its tropical habitat. One species from the Pacific. (*Lophius; ὠμός*, shoulder, in apparent allusion to the trifid humeral spine.)

3092. LOPHIOMUS SETIGERUS (Vahl).

Dorsal III–III–9; A. 5. Head above orbits and laterally with numerous spines and prickles; humeral bone ending in 3 blunt points; numerous cirri scattered along sides of head and body. Vertebræ 18. Color dusky; floor of mouth black posteriorly, but without white spots; pectorals and ventrals pale on basal half, black distally; caudal and anal black, with some white spots; soft dorsal translucent, with black specks; first dorsal spine with its membranaceous tip white, the latter provided with 2 black eye-like spots. Pacific Ocean; not uncommon in rather deep water off coasts of China and Japan. Known on the American coast from 1 speci-

* According to Professor Horace A. Hoffman this fish is called in Athens *Πεσκανδρίτζα* or *Πεσκαντρίτζα*. These names, "probably of Italian origin, meaning fisher; *χλάσχα*, at Chalcis, *σκλεμποῦ*, and *βατραχόψαρο* at Patras. The *βάτραχος ὁ ἁλιευς* (the fisher frog) of Aristotle. (See Aristotle 505a 6b 4, 506b 16, 564b 18, 565b 29, 570b 30, 620b 11 ff, 695b 14, 696a 27, 749a 23, 754a 23 ff, 755a 9, 835b 13, 1527b 41–43, 540b 18.) Aristotle says with regard to the *βάτραχος*: 'Inasmuch as the flat front part is not fleshy, nature has compensated for this by adding to the rear and the tail as much fleshy substance as has been subtracted in front.' The *βάτραχος* is called the angler. He fishes with the hair-like filaments hung before his eyes. On the end of each filament is a little knob just as if it had been placed there for a bait. He makes a disturbance in sandy or muddy places, hides himself and raises these filaments. When the little fishes strike at them he leads them down with the filaments until he brings them to his mouth. The *βάτραχος* is one of the *σελάχη*. All the *σελάχη* are viviparous or ovoviviparous except the *βάτραχος*. The other flat *σελάχη* have their gills uncovered and underneath them, but the *βάτραχος* has its gills on the side and covered with skinny opercula, not with horny opercula like the fish which are not *σελαχώδη*. Some fishes have the gall bladder upon the liver, others have it upon the intestine, more or less remote from the liver and attached to it by a duct. Such are *βάτραχος*, *ἔλλοψ*, *συναγρίς*, *σμύραινα*, and *ξιφίας*. (This has been proved true of *Lophius piscatorius* by a dissection by Dr. C. H. Gilbert.) The *βάτραχος* is the only one of the *σελάχη* which is oviparous. This is on account of the nature of its body, for it has a head many times as large as the rest of its body, and spiny and very rough. For this same reason it does not afterwards admit its young into itself. The size and roughness of the head prevents them both from coming out (i. e., being born alive) and from going in (being taken into the mouth of the parent). The *βάτραχος* is most prolific of the *σελάχη*, but they are scarce because the eggs are easily destroyed, for it lays them in a bunch near the shore." (Hoffman & Jordan, Proc. Ac. Nat. Sci. Phila., 1892, 278.)

men, 2½ inches long, dredged at *Albatross* Station 2805, southwest of Panama. From this specimen, the above description is taken. Comparing this with a larger specimen taken at Tokio by Prof. K. Otaki, we find no differences likely to prove permanent. (*seta*, bristle; *gero*, I bear.)

Lophius setigerus, VAHL,* Skrivt. Naturh., IV, 214, tab. 3, figs. 5 and 6, 1797, China Sea; CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 383, 1837; GÜNTHER, Cat., III, 180, 1861.
Lophius viviparus, BLOCH & SCHNEIDER, Syst. Ichth., 142, 1801, tab. 32, China Sea; after Vahl.
Lophiomus setigerus, GILBERT, Proc. U. S. Nat. Mus. 1890, 454.

Family CCXXII. ANTENNARIIDÆ.

(THE FROG-FISHES.)

Head and body more or less compressed. Mouth vertical or very oblique, opening upward; lower jaw projecting; jaws with cardiform teeth; premaxillaries protractile. Gill openings small, pore-like, in or behind the lower axils of the pectorals. No pseudobranchiæ. Gills 2½ or 3; skin naked, smooth, or prickly. Pectoral members forming an elbow-like angle. Pseudobrachia long, with 3 actinosts. Ventral fins present, jugular, near together. Spinous dorsal of 1 to 3 separated, tentacle-like spines; soft dorsal long, larger than anal. Pyloric cæca none. Genera about 5; species 50. Inhabitants of tropical seas, "living on floating seaweed, and enabled, by filling the capacious stomach with air, to sustain themselves on the surface of the water;" therefore widely dispersed by currents in the sea. (*Pediculati*, pt., Günther, Cat., III, 182 to 200, 1861.)

a. Head compressed; a rostral spine or tentacle, followed by 2 larger spines; palatine teeth developed; dorsal spines disconnected.
b. Skin naked and smooth; ventral fins elongate. PTEROPHRYNE, 1060.
bb. Skin covered with prickles; ventral fins short. ANTENNARIUS, 1061.
aa. Head cuboid; a single rostral spine or tentacle, received in a groove; soft dorsal low. CHAUNAX, 1062.

1060. PTEROPHRYNE, Gill.

(MOUSE-FISH.)

Pterophryne, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 90 (*bougainvillei*).
Pterophrynoides, GILL, Proc. U. S. Nat. Mus., I, 1878, 216 (*histrio*); name a substitute for *Pterophryne*, if the latter be regarded as preoccupied by the earlier *Pterophrynus*.

Body smooth or scarcely granular, short, somewhat compressed, with tumid abdomen; mouth small, oblique; palate with teeth; wrist and pectoral fin slender; ventrals elongated; soft dorsal and anal vertically

**Lophiomus setigerus*, is thus described by Dr. Günther:
"Dorsal III–III, 8 or 9; A. 6 or 7. Teeth arranged in 2 alternate series in the upper jaw, in 3 in the lower; 2 or 3 teeth on each side of the vomer; humeral spine terminating in 3 points; the mouth behind the hyoid bone purplish black, with white spots. Vertebræ 19, the anterior ones very short, the middle and posterior ones nearly equal in length. Coasts of China and Japan." (Günther.)

expanded. Small fishes of fantastic shape in the West Indies and Gulf Stream. (*πτερόν*, wing; *φρύνη*, toad.)

a. "Bait" on first dorsal spine bifurcate at tip. HISTRIO, 3093.
aa. "Bait" on first dorsal spine bulbous, covered with fleshy filaments. GIBBA, 3094.

3093. PTEROPHRYNE HISTRIO (Linnæus).

(MOUSE-FISH; SARGASSUM-FISH.)

Head $2\frac{1}{4}$; depth $1\frac{1}{3}$. D. III-14; A. 7; V. 5. Skin of head and body, as well as dorsal fins, with fleshy tags, which are most numerous on the dorsal spines and abdomen. Wrist slender; ventrals large, nearly $\frac{1}{2}$ as long as head. Dorsal and anal with the posterior rays not adnate to caudal peduncle; first dorsal spine bifurcate at tip. Yellowish, marbled with brown; 3 dark bands radiating from eye; vertical fins barred with brown; belly and sides with small white spots. Tropical parts of the Atlantic; abundant on our Gulf coast and occasional northward to Cape Hatteras or beyond, especially in floating masses of *Sargassum*. Once taken in Europe (Vadsö, Norway) in floating seaweed from the Gulf Stream. Recorded from the coast of Senegambia; its history and synonymy confused with that of the following species. A remarkable fish, excessively variable in coloration. (*histrio*, a harlequin.)

Lophius tumidus, OSBECK, Iter Chinensis, 400, 1757, Open Sea; pre-Linnæan.
Lophius histrio,* LINNÆUS, Syst. Nat., Ed. x, 237, 1758, after various authors, especially *Balistes guaperva* seu *chinensis*, LINNÆUS, Mus. Ad. Fr., 56.
Pterophryne histrio, GILL, Proc. U. S. Nat. Mus. 1878, 216; GOODE & BEAN, Oceanic Ichthyology, 486, 1896.
Antennarius histrio, JORDAN & GILBERT, Synopsis, 846, 1883; COLLETT, Campagnes Hirondelle, 38, 1896.

*Concerning the use of the name *histrio* for this species, Dr. Gill remarks:
"In 1794 (as appears from the dates on the plates), Shaw published a number of his 'Naturalists' Miscellany,' in which he described 3 fishes under the generic name of *Lophius*. These were described as (1) *Lophius striatus* (the Striated Lophius), pl. 175; (2) *Lophius pictus* (the Variegated Lophius), pl. 176, upper figure, and (3) *Lophius marmoratus* (the Marbled Lophius), pl. 176, lower figure. The originals of these are evidently the varieties (*a*, *b*, and *c*) of *Lophius histrio* admitted by Bloch & Schneider. It is quite clear that the first two were based on species of typical *Antennarius* (not *Pterophryne*), while the third is incomprehensible, and, if the figure is at all correct, must represent a factitious fish; it most certainly has nothing to do with *Pterophryne*. The other species, however, notwithstanding the bad figures, are readily identifiable. The *Lophius striatus* (as has recently been recognized by Günther) is the first name of an *Antennarius* peculiar to the Pacific, and quite distinct from the Caribbean *Antennarius scaber* (= *A. histrio* Günther), with which it was at first confounded by Günther. The *Lophius pictus* was evidently based on the species or variety of *Antennarius* which was afterwards named *Antennarius phymatodes* by Bleeker, and it agrees very closely, in the distribution of colors, with a specimen figured by that ichthyologist, and would probably be considered by Günther as a variety of his *Antennarius commersonii*. But whatever may be the value of the forms embraced under the name *Antennarius commersonii* by Günther—whether species or varieties—the name *Antennarius pictus* must be revived from Shaw, either especially for the *Antennarius phymatodes* of Bleeker or for the collection designated as *Antennarius commersonii*. It has thus been demonstrated (1) that the Linnæan name, *Lophius histrio*, was originally created for the common *Pterophryne*, and (2) that the names generally employed for the *Pterophryne* were originally applied to very different forms, and members of even a different genus. Hence if the laws of priority, as formulated by the British and American Associations for the Advancement of Science, are to guide us, there can be no question that the species of *Pterophryne* must hereafter be designated as *Pterophryne histrio*; if, however, it is allowable to go behind even the tenth edition of the Systema Naturæ and to take the oldest binomial name, without other considerations, the designation *tumidus* must be revived. It seems best, however, to follow general usage." (Gill, Proc. U. S. Nat. Mus., I, 1878, 226.)

Chironectes pictus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 393, 1837, Surinam.
Chironectes tumidus, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 397, 1837, "Cabinet du Roi," Sargasso Sea. (Coll. Péron.)
Chironectes arcticus, DÜBEN & KOREN, Kong. Vet. Akad. Abh. Stockholm 1844, 72, Vadsö, Norway, from a specimen carried northward in *Sargassum;* the only European record; fide COLLETT.
Antennarius marmoratus, GÜNTHER, Cat., III, 185, 1861; in part; not of Cuvier.
Chironectes lævigatus, DE KAY, N. Y. Fauna: Fishes, 165, pl. 27, fig. 83, 1842; not of CUVIER.

3094. PTEROPHRYNE GIBBA (Mitchill).

Garman refers to this species certain specimens obtained in Gulf weed about Key West and the Tortugas. These resemble *P. histrio*, but "differ markedly in certain respects. The bait on the first dorsal spine, for instance, is bulbous and covered with slender fleshy filaments in our individuals, but in *P. histrio* it is bifurcate. *P. gibbus* is fairly represented by Cuvier, 1817, in his *Chironectes lævigatus*. The formula for the individuals in hand is D. III, 12; A. 7; V. 5; P. 10; C. 9." (Garman.) West Indies, north to Key West and the Tortugas; not examined by us; probably common, but hitherto confounded with *P. histrio*. (*gibbus*, gibbous.)

Lophius gibbus, MITCHILL, Trans. Lit. and Phil. Soc. N. Y. 1815, I, pl. 4, f. 9, off St. Croix, Lat. 22° N., Long. 64° W. (Coll. Dr. John D. Jaques.)
Chironectes lævigatus, CUVIER, Mém. du Mus., III, 423, pl. 16, fig. 1, 1817, South Carolina (Coll. Bosc); CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 399, 1837.
Pterophryne lævigata, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 90.
? Chironectes sonntagii,* Baron J. W. VON MÜLLER, Reisen in den Vereinigten Staaten, Canada und Mexico, Band I, 180, 1864, in floating seaweed; no exact locality stated.
Pterophrynoides gibbus, GARMAN, Bull. Iowa Lab. Nat. Hist. 1896, 81.

1061. ANTENNARIUS, Lacépède.

Antennarius (COMMERSON) LACÉPÈDE, Hist. Nat. Poiss., I, 421, 1798 (*chironectes*).
Histrio, FISCHER, Zoognosia, 78, 1813 (*histrio*, etc.). (No type; includes all known *Antennariidæ;* description transposed with that of *Lophius* by error.)
Chironectes, CUVIER, Règne Animal, Ed. 2, vol. II, 252, 1829 (*chironectes*); preoccupied in mammals, Illiger, 1811.

Body oblong, compressed, very deep through the occipital region, tapering behind; breast tumid; mouth rather large, more or less oblique, or even vertical; cardiform teeth on jaws, vomer, and palatines; eye small; skin with small granules or spinules, these usually forked, and numerous fleshy slips. First dorsal spine developed as a small rostral tentacle;

* The following is the substance of the long account of *P. sonntagii* (Von Müller):
"D. II, 10 to 12; P. 10 or 11; C. 6 to 7 ('*Strahlenpaare*'); B. 6. Head and body slightly compressed; dorsal spines like little horns, covered over and over with spinous growths. Mouth wide, with numerous rows of small teeth; throat and belly with many fleshy slips. Pectorals produced on a long peduncle like the flippers of a tortoise; ventrals similar, but formed more like feet; anal fin like a rudder. Color clear yellowish green, with greenish brown stripes; a broad dark stripe across breast to root of pectoral; another on the back; another on the side, running backward in the form of a hammer, paler at last on lower part of back; several stripes and spots, more or less dusky, on the tail and other extremities; on the soft underside to the anal intense reddish golden yellow spots; between the dark streaks and the yellow ground color of the body are often white shades and markings; eye fiery orange. Atlantic Ocean or Gulf of Mexico; living in floating seaweed."
This species must be a *Pterophryne*, and it is not evidently different from *Pterophryne gibba*.

second and third dorsal spines strong, covered with skin, with numerous fleshy filaments; soft dorsal high and long; anal short and deep; caudal fin rounded, the peduncle free: pectoral fins wide, with a rather wide wrist, at the lower posterior angle of which are the very small gill openings; ventral fins short. Fantastic-looking fishes, often gaily colored; very numerous in warm seas. (*antenna*, a feeler or tentacle.)

a. Bulbous tip or "bait" of first dorsal spine simple, undivided at tip.
b. Skin smoothish except about eyes; first dorsal spine short, second rough. Body brown, with whitish spots; no ocelli. INOPS, 3095.
bb. Skin with prickles, velvety or shagreen-like.
c. Prickles simple, none of them bifid.
d. Color black; tips of pectorals and ventrals and one or two spots on side white (prickles undescribed). PRINCIPIS, 3096.
dd. Color dusky; dorsal with 3 ocelli; caudal with many spots; first dorsal longer than second; no dermal flaps. TENEBROSUS, 3097.
ddd. Color reddish or grayish, reticulate with heavy black lines; first dorsal spine short. RETICULARIS, 3098.
cc. Prickles or spinules on body mostly bifid.
e. Body without ocelli; first dorsal spine filiform.
f. Mouth immaculate within; body with numerous rosy and dusky tracts, the latter forming bars and concentric streaks below; fins barred. STRIGATUS, 3099.
ff. Mouth largely black within; body blood red, with black spots on sides and below dorsal. SANGUINEUS, 3100.
ee. Body with 3 large ocelli, 1 on dorsal, 1 on caudal, and 1 on middle of side, besides many black spots and streaks; tip of first dorsal spine fringed; mouth largely black within. OCELLATUS, 3101.
aa. Bulbous tip or "bait" on first dorsal spine bifid at tip; skin shagreen-like.
g. Color reddish, with brown spots, those about the eye radiating.
h. Dermal flaps numerous on body; spinules on skin short and stiff, rendering the surface shagreen-like. SCABER, 3102.
hh. Dermal flaps few; spinules on skin longer and slender, rendering the surface velvety. TIGRIS, 3103.
gg. Color uniform black; surface of body rough, shagreen-like; inside of mouth white; first dorsal spine short, little longer than second. NUTTINGII, 3104.
aaa. Bulbous tip or "bait" of first dorsal spine trifid.
i. First dorsal ray twice as long as second and as long as caudal; sides with numerous black ocelli, besides other streaks and dark spots; skin smoothish. MULTIOCELLATUS, 3105.
ii. First dorsal spine barely ½ longer than second; shorter than caudal; sides with dark streaks and reticulations; a large ocellus under middle of soft dorsal; body rough, with shagreen. RADIOSUS, 3106.

3095. ANTENNARIUS INOPS, Poey.

Depth 2¾ with caudal. Skin lustrous, smooth, except for some points behind and below eye; third of the first 3 dorsal rays largest, its membrane not reaching to vent; second ray also large, but shorter, placed between eyes; first spine developed as a fishing rod, filiform, ending in a small, membranaceous lobe, its base close to that of second, and, therefore, distant from end of snout, its spine short, the tip not reaching middle of second spine; short tentacles, like horns, on anterior part of third spine, over the nostrils, and under the mouth; caudal rounded; pectoral so joined that it can not be turned forward as usual in this group, but rising

obliquely backward and upward. Eye slightly longer than snout; mouth brown within. Color brown, with white spots on the body and median fins, 6 of the largest of these each with the center yellowish, the largest from once to twice diameter of eye; spots on dorsal fins small; eye golden. Porto Rico. (Poey.) Not seen by us. The type 70 mm. long. (*inops*, helpless. "I call this species '*inops*' on account of the miserable fishing rod which has fallen to its lot." Poey.)

Antennarius inops, POEY, Anal. Soc. Esp. Hist. Nat., x, 1881, 340, Porto Rico. (Coll. Don Juan Gündlach.)

3096. ANTENNARIUS PRINCIPIS (Cuvier & Valenciennes).

D. III–11; A. 7; P. 10. Anterior dorsal spine twice as long as second, ending in a small, slender lobe; membrane behind third spine extending to root of soft dorsal; last ray of dorsal not reaching caudal. Skin rough, covered with small spines; no cutaneous fringes. Black; tips of pectorals and ventrals white; a small white spot above pectoral. (Günther.) West Indies to Brazil; not seen by us. Günther's specimen, above described, from Para. (*principis*, of the prince. Named for its discoverer, Prince Maurice of Nassau.)

Chironectes principis,* CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 416, 1837, Brazil; on 2 drawings by Prince MAURICE, the second representing the present species to which Günther restricts the name *principis*. The first figure is more like *tigris*.
Antennarius principis, Günther, Cat., III, 193, 1861.

3097. ANTENNARIUS TENEBROSUS (Poey).

D. III–12; A. 7; P. 11. Anterior dorsal spine longer than second, terminating in a simple and slender tentacle; soft dorsal fin terminating at some distance from caudal. Skin rough, covered with small spines, without cutaneous fringes. Blackish brown, marbled with darker and lighter; a series of 3 black, blue-edged ocelli on upper posterior part of dorsal fin; many similar ocelli on caudal fin, irregularly disposed. (Poey.) Cuba. Not seen by us. (*tenebrosus*, dusky.)

Chironectes tenebrosus, POEY, Memorias, I, 219, pl. 17, fig. 1, 1851, Cuba.
Antennarius tenebrosus, GÜNTHER, Cat., III, 197, 1861.

3098. ANTENNARIUS RETICULARIS, Gilbert.

D. III–12; A. 7. First dorsal spine short, very slender and filiform, not reaching tip of second, terminating in a short, fleshy flap; second spine moderately robust, flexible, not curved backward, wholly free and with-

* Concerning this nominal species and *Chironectes mentzelii*, both of which were based on drawings by Prince Maurice, Cuvier & Valenciennes remark:
"We here cite these figures, and we give them specific names only to fix the attention of travelers and to get them to find the species which have served as models for these figures."
The following is the substance of the original description of *Chironectes principis:*
In the first figure, color very deep brown, speckled with black spots on body and fins; dorsal fin with only 1 series of spots. Filament of first spine twice as long as that of the second, and terminating in a little knob or bait, the second spine free from the first and similar in shape.
Second figure of the same form, the second dorsal longer, the first ray ending in a spiral, and the whole body white, with 2 white round spots on each side, one above the other.

out membrane; third spine nearly erect, not free, depressible with difficulty; not curved as in *A. sanguineus;* spines on the body rather coarse and shagreen-like, with expanded, undivided tips. Color in spirits, top of head, including dorsal spines and front of soft dorsal, coral red, the body otherwise light gray, broadly reticulated on sides and below with heavy black lines, which inclose 5 or 6 large pale spots; pectorals, ventrals, and anal with narrow terminal and wide median black bars. Soft dorsal uniformly light. This species closely resembles *A. sanguineus,* but differs from it in the straight, erect spinules, the color, and the character of the plates on the body. Length 1¼ inches. Gulf of California. (Gilbert.) Only the type known. (*reticularis,* netted.)

Antennarius reticularis, GILBERT, Proc. U. S. Nat. Mus. 1891, 566, Gulf of California, at Albatross Station 2825. (Coll. Gilbert.)

3099. ANTENNARIUS STRIGATUS, Gill.

D. III–12; A. 7. First dorsal spine elongate, filiform, twice length of second, with very slender, dermal tip; third spine more robust than second, wholly concealed in the skin, its length equal to that of first spine. Lips, maxillary, and a large transverse area behind second dorsal spine naked, each side of this area with a few spinous tubercles; skin elsewhere covered with fine shagreen-like armature. Color in spirits, olivaceous everywhere on body and on inside of mouth, finely mottled with light olive brown; many irregular blackish areas on head and body, those on lower side of head showing a tendency to form concentric bars; some on sides forming irregular bars downward from back; posterior portion of body not darker than the anterior; terminal parts of all the fins largely blackish, but with distinct black bars; some scattered round blotches on sides, each consisting of a number of smaller black spots on an olive ground; head and body with numerous pinkish and rose-red spots and bars, the latter sinuous, irregular, with wavy margins; a pinkish bar behind maxillary; a broad, saddle-like pinkish blotch across interval between second and third dorsal spines; a third bar from in front of origin of second dorsal downward toward base of pectorals; a fourth across top of caudal peduncle; first dorsal spine narrowly barred with brown. Pacific coast of tropical America, from Cape San Lucas to Panama. Here described from an adult, 10 inches in length, from Panama. This differs considerably from the descriptions of the young (*strigatus, tenuifilis*) given by Gill and Günther.

Two young individuals, types of *A. strigatus,* are thus characterized by Dr. Gill:

"The anterior dorsal spine is very slender and filiform, without appendages; the second is straight and moderate; the third concealed and developed as a hump, obtuse behind. The spines which cover the body are small and mostly bifid. The back and front of the dorsal fin are reddish; the rest light brown, with black stripes which diverge downward above the pectorals, those in front being parallel with the profile and at right angles with those behind; around the pectoral fins and on the flanks, the streaks are generally blended to form a continuous black

area; a black dorsal saddle is in front of the dorsal fin, and a black band covers the posterior half of the caudal fin; the abdomen is broadly reticulated with black, and the brown intervals themselves are frequently striated with the same color; the interior of the mouth is immaculate." (*strigatus*, striped.)

Antennarius strigatus, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 92, Cape San Lucas (Coll. J. Xantus), young; JORDAN & GILBERT, Proc. U. S. Nat. Mus., 1882, 650, adult; JORDAN, Cat. Fishes, 138, 1885.

Antennarius tenuifilis, GÜNTHER, Fishes Centr. Amer., 440, 1869, Panama; young.

3100. ANTENNARIUS SANGUINEUS, Gill.

Anterior dorsal spine very slender, $2\frac{1}{2}$ in length of caudal fin, terminating in a flap extended on each side, laciniated outward; second spine rough, robust, and curved strongly backward at its end; third not free, but apparent as a hump pointed backward, and extending $\frac{2}{3}$ of the distance from its insertion to that of dorsal fin; skin covered with small bifid spines, whose prongs diverge considerably and are acute. Color blood red, except on abdomen, but with several more or less distinct black spots under origin of dorsal fin and on sides; abdomen light or yellowish brown, spotted with black; intervals between caudal and anal rays also marked with black; floor of mouth behind tongue with 2 lateral black bands converging toward the front, while the posterior margin of the tongue itself is also sometimes lined with black. (Gill.) Pacific coast of tropical America, Cape San Lucas to Panama; scarce. (*sanguineus*, bloody.)

Antennarius sanguineus, GILL, Proc. Ac. Nat. Sci. Phila. 1863, 91, Cape San Lucas (Coll. Xantus); JORDAN, Cat. Fishes, 138, 1885.

Antennarius leopardinus, GÜNTHER, Proc. Zool. Soc. London, 1864, 151, Panama.

3101. ANTENNARIUS OCELLATUS (Bloch & Schneider).

Depth $1\frac{2}{3}$ in length. D. II–I–14; A. 8; P. 11; orbit equaling snout, eye much smaller; maxillary $3\frac{1}{2}$ in body; pectoral rays $5\frac{1}{2}$; caudal 4. Body short, oblong, compressed, very deep through occipital region; mouth large, subvertical; teeth small, sharp, cardiform, in wide bands on jaws, vomer, and palatines; maxillary extending downward to below axis of body; a very large knob at symphysis; lower part of head with many large, thick tentacles. First 2 dorsal spines on interorbital space, the first slender, terminating in a fringed lobe, the second shorter and much thicker, behind it a smooth depression; the third spine exceedingly rough and thick, blunt at tip and adnate to body; soft dorsal long and low, its origin in front of middle of body, tips of last rays reaching base of caudal; anal posterior, tips of its rays coterminous with dorsal rays, its height equal to its length; pectorals near middle of body, placed far below axis; ventrals short, the rays thickened, their position under posterior edge of eye. Skin covered with minute bifurcate spines, running upon dorsal, anal, and caudal rays; gill opening in front and below pectoral, its length about equal to snout. Color brown, marbled with

lighter, and with scattered black dots, especially on belly and outer portions of dorsal and caudal; each side with 3 large black spots ocellated with brownish, 1 on dorsal near its base, a second immediately below it on the sides, and a third in the middle of the caudal fin; mouth behind tongue black, with yellow lines. West Indies, north to Florida. Common in the West Indies; the most abundant of the American species. Here described from a specimen from off Pensacola, Florida, about 15 inches in length.

Mr. Garman gives the following note on *Antennarius ocellatus:*

"The species was tolerably figured by Parra, but has not been recognized by some of the subsequent writers. On 5 specimens before me the amount of variations in markings is comparatively small. The 3 large ocelli, on dorsal, caudal, and middle of side, are present on each, as is also the case with the numerous small spots of black on the ventral portions of the body and on the outer portions of dorsal and caudal. The dorsal ocellus lies between the sixth and seventh rays, on the middle of the fin; that on the flank is situated on the vent, and that on the caudal between the fourth and fifth rays, from the top, near the middle of the fin. The black portion of either of these spots is larger than the orbit, which latter is rather small when contrasted with that of other species. The white circle around the black, again, is surrounded by a narrow one of brown. On the caudal, at each side of the ocellus, there are transverse streaks. The first ray of the dorsal is as long as the second, and is covered by scales. The bulb apparently is simple, and bears numerous laciniæ. The second dorsal spine is shorter than the third; both are club shaped. The space behind the second dorsal spine is covered by scales." (*ocellatus*, with eye-like spots.)

Pescador, Parra, Dif. Piezas, Hist. Nat., 1, tab. 1, 1780, Cuba.

Lophius histrio, var. *ocellatus*, Bloch & Schneider, Syst. Ichth., 142, 1801; after Parra.

Antennarius pleurophthalmus, Gill. Proc. Ac. Nat. Sci. Phila. 1863, 92, Key West; Jordan & Gilbert, Synopsis, 846, 1883; Jordan, Cat. Fishes, 138, 1885; Goode & Bean, Oceanic Ichthyology, 487, 1896.

Antennarius ocellatus, Poey, Synopsis, 105, 1868; Garman, Bull. Iowa Lab. Nat. Hist. 1896, 82.

3102. ANTENNARIUS SCABER (Cuvier).

D. III–12; A. 7; P. 9 or 10. Anterior dorsal spine as long as second, and provided with 2 long and thick cutaneous flaps at its tip; third dorsal spine not continuous with the soft dorsal; soft dorsal fin terminating at some distance from the caudal, its last ray not extending to root of caudal, if laid backward; dorsal spines, head, back, and sides of the body with more or less numerous cutaneous fringes, those of dorsal spines sometimes forming a dense cluster; skin very rough, covered with small spines. Ground color yellowish or reddish, with numerous brown spots, those around the eye forming radiating streaks; dorsal and anal fins with 3 series of round brown spots, the middle of which is formed by the largest and most constant spots; sometimes uniform brown. Caribbean Sea. (Günther.) A small specimen from Port Castries, St. Lucia, has the body

light brown, clouded with darker, fins all with round black spots, those of the base of the dorsal somewhat larger than others; ventrals tipped with black. (*scaber*, rough.)

Chironectes scaber, CUVIER, Mém. Mus., III, 425, pl. 16, fig. 2, 1817, Martinique (Coll. Plée); CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 412, 1837.
Lophius spectrum, GRONOW, Cat. Fish.. Ed. Gray, 49, 1854, Antilles; after *Lophius acute scabra*, GRONOW, Zoophyl., 210, 1781.
Antennarius scaber, JORDAN, Proc. U. S. Nat. Mus. 1889, 652, specimen from Port Castries, St. Lucia.
Antennarius histrio, GÜNTHER, Cat., IV, 188, 1861; not *Lophius histrio*, LINNÆUS.

3103. ANTENNARIUS TIGRIS, Poey.

D. III–12; A. 7; P. 11. Anterior dorsal spine longer than second, terminating in 2 long cutaneous flaps; third dorsal spine connected with soft dorsal by a broad membrane, the latter terminating at some distance from the caudal, and its last ray not extending to root of caudal if laid backward. Skin rough, covered with small spines, without cutaneous fringes. Ground color yellow, with numerous brown spots and streaks, the latter radiating from the eye; dorsal fin irregularly spotted, with a series of large round brown spots. (Poey.) Cuba. Not seen by us. According to Mr. Garman, *Antennarius scaber* and *A. tigris* "are closely allied, but if placed side by side the squamation and filaments suffice to distinguish them, great similarity in color notwithstanding. *A. scaber* has coarser scales, with shorter, rougher spines, the scales are farther apart, and the cutaneous flaps appear on the body much as figured by Cuvier. On *A. tigris* there are few of the cutaneous appendages, the scales are closer together, the spines are longer and more slender, giving rise to an appearance more like velvet, and the head and body are more compressed." (*tigris*, tiger.)

Chironectes tigris, POEY, Memorias, I, 217, pl. 17, fig. 2, 1851, Cuba.
Antennarius tigris, GARMAN, Bull. Iowa Lab. Nat. Hist. 1896, 83.

3104. ANTENNARIUS NUTTINGII, Garman.

D. 3+12; A. 7; V. 5; P. 11; C. 9. In form this species is shorter, more massive anteriorly, and less compressed than either *A. ocellatus* or *A. radiosus*. A transerve section across the middle of the body is a nearly equilateral triangle. Caudal region short. Head nearly as wide as high; cheeks swollen; forehead rather broad, converging forward on the edges. Occipital concavity wide and deep, free from scales in a wide space below the ends of the first and second dorsal rays, this bare space being apparently for the reception of the fleshy bait bulb, which latter has 2 elongate lobes. Snout as long as the orbit, broad, truncate; chin vertical; symphyseal knob prominent. Mouth wide, subvertical. Eye small; orbit twice as long, hardly more than ½ the interorbital space. First and second dorsal rays equal in length, not inclusive of the 2 elongate fleshy fringed lobes surmounting the first. The base of the first ray stands forward prominently over the mouth, being free for some distance. The

greater portion of the second ray is free, while the third is connected with the dorsum, by the skin, from base nearly to tip. This last ray is larger than either of its fellows. Soft dorsal large; middle rays longest, as long as the distance from the maxillary to the hind edge of the operculum, or as long as the rays of the caudal fin; fin not reaching back to the bases of the caudal rays, fringed. Hind margin of caudal convex, fringed. Anal moderate, rays prominent in the margin, fin with a blunt angle on the outer edge, subtending, when laid up against the tail, ¼ or more of the length of the caudal rays. The rays on the pectoral fins extend out beyond the margins more noticeably than those of the other fins. Ventrals small, in most instances with 6 points on the outer margin, in one case having but 5. Greatest length of the caudal nearly ¼ of the total length. Length of each maxillary ⅜ of the caudal. Scales short, small, close set, harsh to the touch, having none of the velvety appearance. Uniform black; inside of the mouth black; "bait" white. Great Bahama Banks. Besides the specimens in Nutting's collection there are several others in that of the Mus. Com. Zool. "This species is readily separated from *A. principis* of authors by the short first dorsal spine." ("The specific name is given in honor of Prof. C. C. Nutting, to whom science is so much indebted for the origination and successful accomplishment of the expedition.")

Antennarius nuttingii, GARMAN, Bull. Iowa Lab. Nat. Hist. 1896, 83, pl. II, Great Bahama Banks. (Coll. C. C. Nutting.)

??Chironectes mentzelii,* CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 417, Brazil; on a drawing by Prince MAURICE.

3105. ANTENNARIUS MULTIOCELLATUS (Cuvier & Valenciennes).

(MARTIN PESCADOR.)

Mouth large, vertical. First dorsal spine slender and straight, nearly equal to length of caudal, terminating in 3 simple tentacles; second dorsal spine curved at the middle and extending to the base of the third; third dorsal spine partly embedded in the skin, reaching halfway to dorsal; wrists and pectorals widened; ventrals short. Skin covered with bifid spines. Fawn color, lighter below; many black spots ocellated with white, both on the body and fins; body with several pink areas, 1 of which forms a triangular saddle in front of the dorsal and another a broad ring around the base of the caudal fin; angles of mouth with a pink spot. West Indies, north to Florida Keys; common. This description (after Gill) from the type of *Antennarius annulatus* from Garden Key. According to Mr. Garman, "this species is distinguished by the trifid bulb and the long first dorsal ray, near twice as long as the second and quite as long as the caudal, by the high nape, by the large third dorsal ray, much larger and more swollen than the second, and by the coloration. The eye is very small. The black centers of the largest of the ocelli are smaller than the eye. Besides the ocellus on the soft dorsal that on the anal and the 3

* The following is the substance of the very brief description of this nominal species: "First dorsal filament not longer than the second, and ending in a small bait or knob. Body black, with large marblings."

forming a triangle on the caudal, there are others scattered over the caudal and other fins, and over the sides of the body. Below the eye on the cheek and under the chin and the chest the spots are little more than black dots. Over the sides, a specimen in hand, the type of *A. corallinus*, Poey, is freckled with lighter rounded spots. Behind the pectoral, on the side, there is a small ocellus with a black center. On each side in the same position, a short distance above the pectoral, there is a brown ocellus larger than the orbit, in the center of which there is a white dot. A brown streak passes back from the upper part of the orbit and curves down toward the anal ocellus, another passes back from the middle of the eye and curves down toward the pectoral, and a third below the third dorsal spine runs down and then forward toward the lower end of the maxillary. The forehead is comparatively narrow; behind the second dorsal ray the bare space is hardly large enough to receive the bait."

Poey thus describes *Antennarius corallinus*, which according to Garman, is the same as *A. multiocellatus:*

"D. II–I, 12; A. 7; V. I, 5; C. 17. Two dorsal spines in front of eye, formed like horns, another higher on the nape; gill opening spiral, at the lower base of the pectoral; general form of the fish almost globular; the mouth vertical; tongue marbled with black and white; caudal rounded; eyes very small; pectoral low, reaching middle of body without caudal; ventrals short; vent near anal; first dorsal spine ending in a single short filament. Color reddish with black spots; 2 of these spots eye-like, with a larger black center and the iris of the color of the body, surrounded with a black circle; 1 spot at the base of the soft dorsal at the second third of its length, a very weak one at base of anal; between first spot and pectoral fin 3 small inconspicuous ocellate spots, of which the pupil is a small point; middle of the body with dusky spots; dorsal, anal, and caudal with black points; tubercles about eye and on the cheek, but not spinous. Type 95 mm. long. Cuba." (Poey.) (*multus*, many; *ocellus*, an eye-like spot.)

Chironectes multiocellatus, Cuvier & Valenciennes, Hist. Nat. Poiss., xi, 422, 1837, Martinique. (Coll. M. Garnot.)

Antennarius annulatus, Gill, Proc. Ac. Nat. Sci. Phila. 1863, 91, Garden Key, Florida (Coll. Lieut. Wright); Jordan & Gilbert, Synopsis, 846, 1883.

Antennarius corallinus, Poey, Repertorio, i, 188, 1865, Cuba. (Coll. Poey.)

Antennarius multiocellatus, Günther, Cat., iii, 194, 1861; Garman, Bull. Iowa Lab. Nat. Hist. 1896, 82.

3106. ANTENNARIUS RADIOSUS, Garman.

D. 3+13; A. 8; V. 5; P. 11; C. 9. Resembling *A. tigris*, Poey, in shape, squamation, etc., but differing in coloration and in possession of a much longer dorsal ray. The staff in this ray is very slender, much longer than the second ray, and bears a small, trifid "bait." Second and third dorsal rays shorter than the first, the third well tied down by the skin. Scales uniform, sharp. No cutaneous fringes on large specimens. Grayish or brownish white, darker on nape and dorsal fin, with numerous spots of light color, as large as the orbit, surrounded by more or less complete

edgings of brown, producing a semblance to reticulation, or to spottings by drops of liquid; 7 streaks of brown radiate from the eye; as in *A. tigris*, they are continued upon the head and down toward the ventrals; a large spot of black, white-edged, a little larger than the orbit, ½ on the fin and ½ on the muscles of the body, occupies the space between the eighth and the tenth rays of the soft dorsal fin; the light areas vary in intensity, and lie close together over nearly the whole of body and fins; belly lighter, with faint indications of lines of brownish, radiating from the head; caudal with oblique transverse cloudings of brownish; hindmost ¼ light. The color in life was probably reddish or yellowish. Secured off Key West, in about 50 fathoms. A young individual, of less than an inch, taken opposite Havana, is of lighter gray, and has a large ocellus, of light color in the center, between the black one at the base of the dorsal and the upper end of the humerus. There are small cutaneous fringes on the flanks. (Garman.) (*radiosus*, rayed.)

Antennarius radiosus, GARMAN, Bull. Lab. Nat. Hist. Iowa Univ. 1896, 85, pl. 1, off Key West, in 50 fathoms. (Coll. C. C. Nutting.)

1062. CHAUNAX, Lowe.

Chaunax, LOWE, Trans. Zool. Soc. Lond., III, 1846, 339 (*pictus*).

Head very large, depressed, cuboid. Mouth large, subvertical; jaws and palate with bands of small teeth. Skin with small, sharp spines. Spinous dorsal reduced to a small tentacle above the snout, retractile into a groove; soft dorsal moderate, low; anal short; ventrals small. Gills 2½; no pseudobranchiæ. Muciferous channels very conspicuous, the lateral line prominent, undulate; another series of mucous tubes extending from lower jaws to axil; still another extends backward from snout and maxillary to a point behind eye, when it ceases, uniting with a vertical line which extends from the lateral line to the lower line; these lines thus inclose a quadrate area on the cheek. Gill opening small, well behind pectoral under front of soft dorsal. Deep seas. (χαύναξ, one who gapes.)

a. Dorsal rays 11; anal 5; depth 2½ in length. PICTUS, 3107.
aa. Dorsal rays 13; anal 7; depth 2⅔ in length. NUTTINGII, 3108.

3107. CHAUNAX PICTUS, Lowe.

Head 1⅓; depth 2½. D. I, 11; A. 5; P. 11; V. 4; C. 7. Rostral tentacle short, pedicellate; muciferous channels appearing as chain-like rows of pits. Bright orange above; sides rosy; fins vermilion. Deep waters of the Atlantic; recorded from Madeiro, Soudan, Cape Verdes, Barbados, off Rhode Island, and elsewhere in the Gulf Stream, in 130 to 428 fathoms. A similar species (*Chaunax fimbriatus*), regarded by Günther as the same, occurs in the Japan Seas, Bay of Bengal and the Fiji Islands. (*pictus*, painted.)

Chaunax pictus, LOWE, Trans. Zoöl. Soc. Lond. 1846, 339, Camera de Lobos, Madeira; GÜNTHER, Cat., III, 200, 1861; GOODE, Proc. U. S. Nat. Mus. 1880, 470; JORDAN & GILBERT, Synopsis, 847, 1883; GOODE & BEAN, Oceanic Ichthyology, 487, fig. 398, 1896.

? *Chaunax fimbriatus*, HILGENDORF, Sitzber. Ges. Naturf. Freunde 1879, 80, Sea of Japan.

? *Chaunax nuttingii*, GARMAN, Bull. Lab. Nat. Hist. Iowa Univ. 1896, 85, pl. III, fig. 2, near Sand Key Light, Florida, in 120 fathoms. (Coll. C. C. Nutting.)

3108. CHAUNAX NUTTINGII, Garman.

B. 6. D. II, 13; A. 7; V. 4; P. 14; C. 9. Form resembles that of *Chaunax pictus*, but is shorter, broader, and possessed of more fin rays. Anteriorly it is broad and depressed, posteriorly compressed. From head to soft dorsal on the nape it is arched very little. Head broader than high, flattened or slightly concave on the occiput, nearly vertical on the chin. Snout short, broad, truncate. Eye medium, the length of the scaleless area covering it equals the width of that between the canals on the interorbital space, or about $\frac{3}{5}$ of the space itself; the distance from the maxillary is about the ocular width. The niche in which the first dorsal spine is received is subelliptical and about $\frac{1}{4}$ as long as the eye; the tentacle is little more than $\frac{1}{2}$ as long as the niche, is broad near the base, tapers rapidly and bears a 2-lobed "bait" with slender fringes. Mouth wide, oblique, maxillary about 3 times as long as the eye, widened and rounded at the outer end; intermaxillaries alone forming upper border of mouth. Teeth small, slender, sharp, in villiform bands. Origin of soft dorsal in the middle of the distance from the rostral tentacle to the base of the caudal fin, fourth ray above the gill opening, anterior rays shorter. Vent below the seventh ray of the second dorsal. Pectorals short, broad, rounded. The canals of the lateral system are in the main like those of *C. pictus*, but have stronger curves; they begin to curve outward immediately behind the niche, not remaining parallel or converging as in Lowe's species. Scales very fine, sharp and close together. In life this fish was probably red or yellowish with transverse cloudings or blotches of brownish, it is now dingy brownish white; one of the blotches lies just behind the eye, another lies below the orbit, and apparently 3 transverse bands cross the back through the soft dorsal; orbit blackish; tentacular niche black. The coloration of the individual described indicates a habitat within reach of the effects of sunlight. Florida Keys; the type dredged nearly 8 miles south of Sand Key Light, Florida, in about 120 fathoms. (Garman.) This species is evidently not very different from *Chaunax pictus* and may be the same. (Named for C. C. Nutting, professor of zoology in the University of Iowa, director of the Bahama Expedition of 1893.)

Chaunax nuttingii, Garman, Bull. Lab. Nat. Hist. Iowa Univ. 1896. 86, pl. III, f. 2, Sand Key Light, Florida. (Coll. C. C. Nutting.)

Family CCXXIII. CERATIIDÆ.

(The Sea Devils.)

Head and body compressed. Mouth terminal, more or less oblique. Gill openings small, in the lower part of the axils. No pseudobranchiæ. Spinous dorsal represented by 1 or more tentacles. Pectoral members not geniculated, with short pseudobrachia and 3 actinosts. No ventral fins. Fishes of the open seas, usually inhabiting considerable depths; 13 genera and 15 species known. All are uniform blackish in color.

"The bathybial sea devils," writes Günther, "are degraded forms of *Lophius;* they descend to the greatest depths of the ocean. Their bones

are of an extremely light and thin texture, and frequently other parts of their organization, their integuments, muscles, and intestines are equally loose in texture when the specimens are brought to the surface. In their habits they probably do not differ in any degree from their surface representative, *Lophius*. The number of the dorsal spines is always reduced, and at the end of the series of these species only 1 spine remains, with a simple, very small lamella at the extremity (*Melanocetus johnsonii*, *Melanocetus murrayi*). In other forms sometimes a second cephalic spine, sometimes a spine on the back of the trunk, is preserved. The first cephalic spine always retains the original function of a lure for other marine creatures, but to render it more effective a special luminous organ is sometimes developed in connection with the filaments with which its extremity is provided (*Ceratias bispinosus*, *Oneirodes eschrichtii*). So far as it is known at present these complicated tentacles attain to the highest degree of development in *Himantolophus* and *Ægœonichthys*. In other species very peculiar dermal appendages are developed, either accompanying the spine on the back or replacing it. They may be paired or form a group of 3, are pear-shaped, covered with common skin, and perforated at the top, a delicate tentacle sometimes issuing from the foramen." (*Pediculati*, genus *Ceratias*, Günther, Cat., III, 205, 1861; *Ceratiidæ*, Gill, Proc. U. S. Nat. Mus. 1878, 216.)

a. Mouth moderate.
b. Gills in 2½ pairs.
CERATIINÆ:
c. Cleft of mouth nearly vertical; skin prickly.
d. Cephalic spine single.
e. Dorsal spine present; lateral caruncles present; no teeth on vomer. CERATIAS, 1063.
ee. Dorsal spine wanting; caruncles present.
f. Caruncles remote from soft dorsal. MANCALIAS, 1064.
ff. Caruncles close to soft dorsal. CRYPTOPSARAS, 1065.
ONEIRODINÆ:
cc. Cleft of mouth nearly horizontal; skin smooth; 1 cephalic spine and 1 postcephalic spine. ONEIRODES, 1066.
HIMANTOLOPHINÆ:
bb. Gills in ½ 2½ pairs; body with scattered tubercular scutella; no second dorsal spine.
g. Body and head compressed; mouth oblique; joint of mandible below or behind eye; eye rudimentary.
h. Body oblong, oval; dorsal rays 9; pectoral 12. HIMANTOLOPHUS, 1067.
hh. Body short and deep; dorsal rays 4; pectoral about 17. CORYNOLOPHUS, 1068.
aa. Mouth with enormous gape.
MELANOCETINÆ:
i. Cleft of mouth nearly vertical; pectoral small, in advance of dorsal and of gill opening; second dorsal spine wanting; gills in 2½ pairs.
j. Gular tentacle wanting; no teeth on vomer. LIOCETUS, 1069.
jj. Gular tentacle present; 1 tooth on the vomer. LINOPHRYNE, 1070.
CAULOPHRYNINÆ:
ii. Cleft of mouth nearly horizontal; pectorals below dorsal and behind gill opening; gills in ½ 2½ pairs.
k. Dorsal and anal greatly produced; skin naked; head and body with many luminous filaments. CAULOPHRYNE, 1071.

1063. CERATIAS, Kröyer.

Ceratias, KRÖYER, Naturhist. Tidsskrift. 2 Række, 1, 639, 1844 (*holbolli*).

Head and body much compressed and elevated, oblong, covered with prickly skin. Mouth wide, its cleft nearly vertical; teeth in jaws conic, movable, of moderate size; no teeth on vomer or palatines. Gills $2\frac{1}{2}$; gill arches unarmed. Spinous dorsal reduced to 2 spines, 1 on the head, the other on the back, the basal element of the second spine exserted; the cephalic spine much elongate; soft dorsal and anal short; pectorals very short, broad, of about 20 rays. Caudal fin much produced, fan-shaped, with exserted rays. Pyloric cæca 2, small. Skeleton soft, fibrous. Greenland. (*κερατίας*, one that has horns.)

3109. CERATIAS HOLBOLLI, Kröyer.

D. I–I, 4; A. 4; P. 19; C. 10. Head $2\frac{3}{8}$; depth nearly 2; head deeper than long; eyes small, not more than $\frac{1}{10}$ the length of the head; free rays of the head a little shorter than to the base of the caudal fin; the forked part of the caudal fin shorter than the length of the fish; the length of the pectoral fin equals almost $\frac{1}{10}$ the entire length of the fish, the membranes from the dorsal and anal fins posteriorly extend almost to the base of the caudal fin. Color entirely black. (Kröyer.) North Atlantic; 4 specimens known; 3 from Greenland, 1 from Nova Scotia. (Named for C. Holböll, a Danish naturalist.)

Ceratias holbolli, KRÖYER, Naturh. Tidsskr. 1844, 639, Greenland; GÜNTHER, Cat., III, 205, 1861; JORDAN & GILBERT, Synopsis, 847, 1883; GOODE & BEAN, Oceanic Ichthyology, 489, pl. 117, fig. 394, 1896.

1064. MANCALIAS, Gill.

Mancalias, GILL, Proc. U. S. Nat. Mus., 1, 1878, 227 (*uranoscopus*).
Typhlopsaras, GILL, Forest and Stream, New York, 1883, Nov. 8, 284 (*shufeldti*).

General characters of *Ceratias*, but with the spinous dorsal reduced to a single rostral spine, and 2 fleshy claviform tubercles or caruncles behind it. Pectoral fins narrow, with 10 to 15 slender rays. (*mancus*, defective, "with a quasi-diminutive termination to correspond with *Ceratias*.")

a. Dorsal caruncles placed before dorsal fin a distance 6 times in length of trunk from gill opening to base of caudal. URANOSCOPUS, 3110.

aa. Dorsal caruncles placed nearer dorsal, the distance from dorsal $4\frac{1}{3}$ times in trunk, as above. SHUFELDTI, 3111.

3110. MANCALIAS URANOSCOPUS (Murray).

D. I, 3 or 4; A. 4; C. 8; P. 10. Anterior spine of first dorsal produced in a long filament, ending in a pear-shaped bulb, terminating in a semitransparent whitish spot, this spine originating on posterior part of head, and reaching, when depressed, nearly to the tip of tail; far behind this are 2 short, fleshy tubercles, lying in a depression in front of second dorsal. Teeth moderate, depressible. Skin everywhere with minute embedded conical spines. Eye very small, placed high on the middle of the head.

Color uniform black. (Murray.) To this description Goode & Bean add from the same specimen: "Anal opposite second dorsal, the 4 median caudal rays being much larger than the others and bifid; pectorals small, above the gill opening; the upper jaw is formed by the intermaxillaries, and is armed, together with the lower jaw, with a series of teeth of moderate size, which can be depressed as in *Lophius*. The skin is thickly covered with minute embedded conical spines; the eyes are very small and are placed high upon the middle of the head." Mid-Atlantic, in very deep water; 2 specimens known, the type in 2,400 fathoms, taken between the Canary and Cape Verde islands, the second (26159) in 372 fathoms off the coast of Rhode Island in the Gulf Stream. (*οὐρανοσκόπος*, star-gazing, from the upturned eyes.)

Ceratias uranoscopus, Murray, in Wyville Thompson, The Atlantic, II, 67, fig. 20, 1878, mid-Atlantic, between Canary and Cape Verde Islands in 2,400 fathoms; Günther, Challenger Report, XXII, 54, pl. 11, fig. C, 1887.

Mancalias uranoscopus, Gill, Proc. U. S. Nat. Mus. 1878, 228; Goode, Proc. U. S. Nat. Mus. 1880, 469; Jordan & Gilbert, Synopsis, 848, 1883; Goode & Bean, Oceanic Ichthyology, 490, 1896.

3111. MANCALIAS SHUFELDTI (Gill).

Maxillary $\frac{1}{5}$ the length from gill opening to caudal base; intermaxillary $3\frac{1}{2}$ times in this length. Form more slender than that figured by Günther, with 4 rays in the dorsal, and apparently 15 in pectoral. There are no vomerines; intermaxillary and mandible armed with a narrow band of depressible teeth of various lengths. The skin with a fine granular appearance and everywhere covered with minute prickles. The caruncles only 2 in number and situated as in *Mancalias uranoscopus*, as figured in the *Challenger* fishes. Length of dorsal spine, without the joint bearing the pear-shaped appendage, equaling distance from gill opening to root of tail; the joint bearing the appendage is $\frac{2}{3}$ of this distance; in *Mancalias uranoscopus* (No. 26159) the first dorsal, without the joint bearing the appendage, contains the distance from the gill opening to the root of the tail $1\frac{1}{4}$ times. The joint containing the appendage is $\frac{1}{2}$ as long as the distance from the gill opening to root of tail. Dermal caruncles distant from the dorsal a space equal to $\frac{1}{6}$ of distance from the gill opening to root of tail. In *M. shufeldti* the caruncles are placed at a distance from the dorsal a space contained $4\frac{1}{2}$ times in the distance from the gill opening to the root of the tail. In the specimens described by Goode & Bean as *Mancalias uranoscopus* (No. 26159), the length $3\frac{1}{4}$ inches, the length of the maxillary is $\frac{1}{5}$ of length from gill opening to root of tail, and the intermaxillary $3\frac{1}{2}$ times in same distance. Teeth in jaws depressible, in narrow bands, and of unequal size; vomer toothless. Two small caruncles not far from front of dorsal fin, and instead of being placed opposite each other, according to the usual arrangement, one is placed behind the other. Skin covered with minute granules or papillæ, each one surmounted by a slender prickle, as in *Typhlopsaras*. The pectoral of the individual described contains 15 rays. The pectorals of *T. shufeldti* are imperfect. (Goode & Bean.) Gulf Stream, off the coast of southern New England;

1 specimen known. (Named for Dr. Robert W. Shufeldt, United States Army, the well-known ornithotomist.)

*Typhlopsaras shufeldti,** GILL, Forest and Stream, Nov. 8, 1883, Western Atlantic (Type, No. 33552); JORDAN, Cat. Fishes, 138, 1885.
Ceratius shufeldti, GÜNTHER, Challenger Report, XXII, Deep-Sea Fishes, 54, 1887.
Mançalias shufeldti, GOODE & BEAN, Oceanic Ichthyology, 490, fig. 401, 1896.

1065. CRYPTOPSARAS, Gill.

Cryptopsaras, GILL, Forest and Stream, Nov. 8, 1883, 284 (*couesii*).

Body shortened; back longitudinally convex, eyes small but conspicuous; anterior spine with concealed basal joint and elongated terminal joint; a large intermediate globular and a pair of subpedunculated lateral dorsal appendages or caruncles close to the front of the dorsal fin; pectorals well developed, of about 15 rays. Deep seas. (κρυπτός, concealed; modern Greek ψαρᾶς, fisherman, in reference to the concealed rod bearing the dorsal spine or fishing apparatus.)

3112. CRYPTOPSARAS COUESII, Gill.

The basal joint of the rod-like spine is almost entirely concealed and procumbent, and the distal joint alone free, reaching backward to the dorsal tubercle; the bulb is pyriform, and surmounted by a long whitish filament; dorsal and anal each with 4 spines, the caudal 8 (the 4 middle dichotomous) and the pectorals each about 15 rays. (Gill.) A specimen of *Cryptopsaras* (No. 33558, U. S. Nat. Mus.) was obtained, by the *Albatross*, from Station 2101, in Lat. 38° 18′ 30″ N., Long. 68° 24′ W., at a depth of 1,686 fathoms. The type of *Cryptopsaras couesii* is only 35 mm. long. The caudal is imperfect. The length without caudal is 30 mm. and contains the greatest height 2½ times. The bulb on the dorsal spine when laid backward can be made to reach to the dermal caruncles on the back. The length of the upper jaw is about ¼ of the length without caudal; gill opening nearly midway between front of head and root of tail; mouth placed vertically; intermaxillary teeth occupying about entire length of bone; mandibulary teeth unequal in size; at symphysis of mandible a pair of minute spines closely connected at base and slightly separated at the extremity. Specimen No. 39483 is 58 mm. long; 47 mm. to base of caudal. Greatest height 2⅔ in length without caudal. Gill opening a little nearer end of caudal than to front of head; distal portion of dorsal spine about ½ length without caudal; median dermal caruncle very much

* The following is the original account of *Typhlopsaras:*

"*Typhlopsaras.*—Ceratiines with an elongated trunk, rectilinear back, obsolete or no eyes, far exserted basal joint of the anterior spine and shortened terminal joint, a small intermediate and a pair of pedunculated dorsal appendages some distance in advance of the dorsal fin, and reduced pectoral fin with about 5 or 6 rays.

"*Typhlopsaras shufeldti.*—The first joint of the rod-like spine reaches to the axil of the dorsal fin, and the bulb to the base of the caudal fin, when the spine is bent backward; the bulb is pear-shaped and without any appendages; the dorsal has 4 rays, the anal 4, the caudal 8 (the median 4 of which are forked), and there are 4 or 5 pectoral rays. A single specimen was found. I have dedicated the species to my esteemed friend, Dr. R. W. Shufeldt, U. S. A., the well-known ornithotomist.

"The name, *Typhlopsaras*, is a compound from the Greek τυφλος (blind) and ψαρας (angler), meaning 'blind angler.'"

larger than the two lateral ones; skin covered with minute granules of uniform size; pectoral with 16 rays, its length about $\frac{1}{6}$ that of head; length of upper jaw about $\frac{1}{3}$ of total without caudal; pair of spines at symphysis of mandible replaced by a very small knob; teeth in intermaxillary very small, diminishing in number toward the symphysis, apparently uniserial. On each side of head of vomer 2 or 3 depressible teeth; palatines apparently wanting. We have seen something like traces of similar teeth on the vomer of *Mancalias uranoscopus*, but owing to the condition of the specimen can not be certain about this character. (Goode & Bean.) Gulf Stream, off the coast of New England. (Named for the eminent ornithologist, Dr. Elliott Coues.)

Cryptopsaras couesii, GILL, Forest and Stream, Nov. 8, 1883, 284, Gulf Stream off New England (Coll. Albatross); JORDAN, Cat Fishes, 138, 1885; GOODE & BEAN, Oceanic Ichthyology, 491, fig. 402, 1896.

Ceratias couesii, GÜNTHER, Challenger Report, XXII, 55, 1887.

?Ceratias carunculatus, GÜNTHER, Challenger Report, XXII, 55, pl. 11, fig. *d*, 1887, south of Yezo, Japan, in 345 fathoms; 1½ inches long. (Coll. *Challenger*.)

1066. ONEIRODES, Lütken.

Oneirodes, LÜTKEN, Overs. Kong. Dansk. Vidensk. Selsk. Forhandl. 1871, 56 (*eschrichtii*).

Body compressed, oval, short, covered with smooth skin. Head compressed, very large. Mouth moderate, almost horizontal, the joint of mandible behind eyes. Teeth unequal, depressible; vomer with teeth. Gill arches unarmed; gills in 2½ pairs. Spinous dorsal represented by a cephalic spine, the basal element of which is procumbent and subcutaneous, the tip bulbous, and a second spine about midway between the rostral spine and the soft dorsal; soft dorsal and anal short; no ventrals; no pyloric cæca. Greenland. (ὀνειρώδης, dream-like, in illusion to the small, almost covered, eyes.)

3113. ONEIRODES ESCHRICHTII, Lütken.

D. I–I, 6; A. 8; C. 8. Terminal half of the bulb of the cephalic spine whitish. Cephalic spine with a bulbous termination, surmounted by slender filaments in several transverse rows. Caudal fin shorter than trunk without head. Color black. Deep sea, off Greenland. Known from a single specimen 8 inches long. (Gill.) (Named for D. F. Eschricht, a Danish naturalist, a student of the Cetacea.)

Oneirodes eschrichtii, LÜTKEN, Overs. Dansk. Vidensk. Selsk. Forhandl. 1871, 56, 9–18, pl. 2, deep sea off Greenland; GILL, Proc. U. S. Nat. Mus. 1878, 218; JORDAN & GILBERT, Synopsis, 848, 1883; GOODE & BEAN, Oceanic Ichthyology, 492, 1896.

1067. HIMANTOLOPHUS, Reinhardt.

Himantolophus, REINHARDT, Dansk. Vidensk. Selsk. Nat. 1837, 74 (*grœnlandicus*).

Head large, compressed. Skin thick, with scattered, round, prickly scales. Body oval, compressed. Mouth moderate, the cleft oblique, the joint of the mandible below or behind the eyes. Gills in ½ 2½ pairs; gill arches armed with dentigerous tubercles. Spinous dorsal represented

only by a single long rostral spine, the basal element of which is procumbent and subcutaneous; the extremity with numerous long filaments. Soft dorsal short, with 9 rays; anal short; pectoral rather broad, with 12 rays. Greenland. (*ἱμάς*, a thong; *λόφος*, crest.)

3114. HIMANTOLOPHUS GRŒNLANDICUS, Reinhardt.

Depth 2½ in total length. D. I-9; P. 12. Body oblong oval. Cephalic ray provided with about 11 tentacles. (Gill.) Greenland. "This species has never been fully described, the only existing example being an imperfect one, 23 inches long, obtained off the coast of Greenland, about 1837."

Himantolophus grœnlandicus, Reinhardt, Dansk. Vid. Selsk. Nat. Math. Afh. 1837, 74, Greenland; Gill, Proc. U. S. Nat. Mus. 1878, 218; Jordan & Gilbert, Synopsis. 849, 1883.

1068. CORYNOLOPHUS, Gill.

Corynolophus, Gill, Proc. U. S. Nat. Mus. 1878, 219 (*reinhardti*).

This genus is scarcely distinct* from *Himantolophus*, differing in the short oval form, the short dorsal of about 5 rays, and the broader pectoral with about 17 rays each. (*κορύνη*, club; *λόφος*, crest.)

3115. CORYNOLOPHUS REINHARDTI (Lütken).

Depth 1⅓. D. I-5; P. 17. Body short, oval; cephalic ray ⅔ length of head, with about 8 tentacles, which branch out forming a brush at tip; skin sparsely covered with thorn-like prickles. Greenland. One specimen known, 14 inches long. (Named for Prof. Johann Reinhardt, naturalist at the University of Copenhagen.)

Himantolophus reinhardti, Lütken, Kong. Dansk. Vidensk. Selsk. 1878, 321, Greenland.
Corynolophus reinhardti, Gill, Proc. U. S. Nat. Mus. 1878, 219.

1069. LIOCETUS, Günther.

Liocetus, Günther, Challenger Report, xx., 57, 1887 (*murrayi*).

Mouth enormous, the cleft nearly vertical; pectorals small, in advance of dorsal and of gill opening; second dorsal spine wanting; gills in 2½ pairs; no gular tentacle. This genus is similar to *Melanocetus*, differing in having no teeth on the vomer, a greater projection of the mandible, and a smaller mouth. Deep sea. (*λεῖος*, smooth; *κῆτος*, whale.)

3116. LIOCETUS MURRAYI (Günther).

D. I-13; A. 4; C. 9; P. 14. (Günther.) Extremely similar to *Melanocetus johnsonii*, but without trace of vomerine teeth, while there is no distinction between the two species as regards dentition of jaws; posterior angle

*Dr. Gill, replying to certain strictures as to the validity of this genus, made by Lütken [who calls it a "wanton' subdivision], states that the "differences alleged to exist between *Himantolophus* and *Corynolophus* are very marked. If they do exist as stated there can be no doubt that the two should be kept apart. I know of no reason except the singularity and greatness of the difference specified for doubting the correctness of Reinhardt's observations."

of mandible projecting more and forming a salient point; mouth comparatively less wide, and the maxillary considerably shorter, being about $\frac{2}{3}$ of total length, without caudal, while it is rather more than $\frac{1}{2}$ in the Madeiran species. Eye rudimentary. One cephalic spine, shorter than maxillary; last dorsal ray connected by a short and delicate membrane with caudal fin; most of the caudal rays bifid, the longest shorter than maxillary; pectoral fin as much developed as in *Melanocetus johnsonii*. Entirely black. Total length 44 lines; length of mandible 14 lines; length of maxillary 12 lines; length of caudal fin $10\frac{1}{2}$ lines. A young individual, 44 lines in length, was taken by H. M. S. *Challenger* in the mid-Atlantic, at a depth of 1,850 fathoms (Station 106); another of 13 lines at the depth of 2,450 fathoms (Station 348). (Goode & Bean.) (Named for Dr. John Murray, second director of the civilian staff on board H. M. S. *Challenger*.)

Melanocetus bispinosus, GÜNTHER, Study of Fishes, 473, 1880; name only.

Melanocetus (*Liocetus*) *murrayi*, GÜNTHER, Challenger Report, XXII, 57, pl. 11, fig. A, 1887, mid-Atlantic.

Liocetus murrayi, GOODE & BEAN, Oceanic Ichthyology, 495, fig. 407, 1896.

1070. LINOPHRYNE, Collett.

Linophryne, COLLETT, Proc. Zool. Soc. London 1886, 138 (*lucifer*).

Head enormous; the body slender, compressed, mouth oblique. Spinous dorsal reduced to a single cephalic tentacle, the basal part of which is erect, not procumbent. Teeth in the jaws on the vomer and the upper pharyngeals. Gill openings exceedingly narrow, situated a little below the root of the pectoral. Soft dorsal and anal very short; ventrals none. Abdominal cavity forming a sac, suspended from the trunk. Skin smooth; a long tentacle on the throat. This genus differs from *Melanocetus* in the presence of the gular tentacle. (λίνος, linen, net; φρύνη, a toad.)

3117. LINOPHRYNE LUCIFER, Collett.

D. I–3; A. 2; C. 9; P. 14 or 15. A spinous projection or horn above each orbit. Cephalic tentacle black, with a large ovate bulb, the upper half of which is white; gular tentacle much larger, terminating in 2 tongue-like appendages, which are furnished on the upper edge with a row of round, white papillæ. (Goode & Bean.) Mid-Atlantic, northwest of Madeira, Lat. 36° N., Long. 20° W. One specimen known. (*Lucifer*, an evil spirit; *lux*, light; *fero*, I bear.)

Linophryne lucifer, COLLETT, Proc. Zool. Soc. London 1886, 138, pl. 15, mid-Atlantic, between Madeira and the West Indies (Coll. Capt. P. Andresen. Mus. Univ. Christiania); GÜNTHER, Challenger Report, XXII, 57, 1887; GOODE & BEAN, Oceanic Ichthyology, 496, fig. 408, 1896.

1071. CAULOPHRYNE, Goode & Bean.

Caulophryne, GOODE & BEAN, Oceanic Ichthyology, 496, 1896 (*jordani*).

Head large, compressed; mouth with the cleft nearly horizontal; body short, much compressed. Spinous dorsal reduced to a single cephalic tentacle, which is supported on a short procumbent base. Teeth of unequal size in the intermaxillary and the mandible; vomer, palatines, and upper

pharyngeals toothed. Gill openings narrow, horizontal slits placed below and in front of root of pectorals. Branchiæ in $\frac{1}{2}$ $2\frac{1}{2}$ pairs. Branchial arches armed with dentigerous tubercles. Skin naked. Numerous luminous filaments on head and body. Soft dorsal and anal many-rayed, the rays greatly produced; caudal long, tapering; ventrals none; pectorals very broad, sessile, postmedian, under dorsal fin, with numerous rays. Pyloric appendage reduced to 1 small rudiment. Air bladder absent. (*καυλός*, stem; *φρύνη*, toad, from the many stems or fin rays.)

3118. CAULOPHRYNE JORDANI, Goode & Bean.

Depth about 2 in length without caudal, the greatest height occurring behind the head. Cephalic appendage with a pale tuft at its tip, the length of the distal portion 3 in length of body. The tuft somewhat mutilated, but showing no evidence of a laminated structure; basal portion of cephalic appendage about twice as long as the very small eye; maxillary very slender, narrow, extending about as far backward as intermaxillary; intermaxillary slightly protractile and with about 10 teeth on each side, several of which are nearly twice as large as the rest, its length $2\frac{1}{2}$ in body; mandible as long as head without snout, with 8 teeth on each side, the anterior pair and several other pairs along shaft of bone being greatly enlarged; a pair of enlarged teeth on head of vomer; several similar teeth on palatines; upper pharyngeals armed with several strong teeth; eye very small, inconspicuous, its distance from tip of snout equaling nearly $\frac{1}{3}$ its distance from soft dorsal origin. Intestine shorter than length without caudal. Soft dorsal with 16 rays, all of which, except the last 4, are greatly produced; the second, third, and fourth rays longest, nearly twice as long as body. Anal with 14 rays, all of which, except last 3, are much produced, the fin not quite perfect, yet its anterior rays are longer than body; caudal with 8 rays, the 4 inner ones divided, the rest simple; middle rays of caudal as long as distance from tip of lower jaw to base of pectoral; pectoral comparatively short, with 16 simple articulated rays, the longest about $\frac{1}{2}$ as long as head. About 9 luminous filaments on each side of head, 7 more between nape and dorsal, and about 12 on sides; the filaments nearly twice as long as eye. Head and body black; caudal, cephalic tuft, and most of the rays pale. Gulf Stream. The type of the species (No. 39265) was taken by the steamer *Albatross*, September 19, 1887, in Lat. 39° 27′ N., Long. 71° 15′ W., 1,276 fathoms. (Named for David Starr Jordan.)

Caulophryne jordani, Goode & Bean, Oceanic Ichthyology, 496, pl. 21, fig. 409, 1896, Gulf Stream, off Carolina, in 1,276 fathoms. (Coll. *Albatross*; the plate named *Caulophryne setosus*, by slip in proof reading.)

Family CCXXIV. OGCOCEPHALIDÆ.

(The Bat-Fishes.)

Head very broad and depressed, the snout more or less elevated, the trunk short and slender. Mouth not large, subterminal or inferior, the lower jaw included; teeth villiform or cardiform. Gill openings very

small, above and behind the axils of the pectoral fins. Body and head covered with bony tubercles or spines. Spinous dorsal reduced to a small rostral tentacle, which is retractile into a cavity under a prominent process on the forehead; in 1 genus the rostral tentacle is obsolete; soft dorsal and anal fins small and short; ventrals well developed; pectoral fin well developed, its base strongly angled, with long pseudobrachia and 3 actinosts. Branchiostegals 5; no pseudobranchiæ. Genera 8; species about 30, chiefly American, some of them in the deep sea. (*Pediculati*, part; genera *Malthe* and *Halieutæa*, Günther, III, 200–205, 1861.)

OGCOCEPHALINÆ:

a. Disk with the frontal region elevated and the snout more or less produced forward, the tail stout; orbits lateral; teeth on vomer and palatines; rostral tentacle present.

b. Gills $2\frac{1}{2}$; disk longer than broad. OGCOCEPHALUS, 1072.

bb. Gills $2\frac{1}{2}$; disk broader than long. ZALIEUTES, 1073.

HALIEUTINÆ:

aa. Disk with the frontal region depressed, not elevated above the rest; eyes partly superior; snout rounded, obtuse in front; tail slender.

c. Dorsal fin present.

d. Vomer and palatines with teeth. HALIEUTICHTHYS, 1074.

dd. Vomer and palatines toothless.

e. Disk subcircular; gills $2\frac{1}{2}$.

f. Mouth rather large, subvertical; prickles rather strong. HALIEUTÆA, 1075.

ff. Mouth rather small, terminal; prickles feeble. HALIEUTELLA, 1076.

ee. Disk subtriangular; gills 2; prickles very strong. DIBRANCHUS, 1077.

1072. OGCOCEPHALUS, Fischer.

(SEA-BATS.)

Ogcocephalus, FISCHER, Zoognosia, 78, 1813 (*vespertilio*).
Oncocephalus, GILL, modified spelling.
Malthe, CUVIER, Règne Animal, Ed. I, II, 311, 1817 (*vespertilio*).
Malthæa, corrected spelling.

Body stoutish, tapering backward; head very broad and depressed, triangular in form, the forehead elevated and produced. Eyes large, lateral. Mouth rather small, subinferior under the snout; villiform teeth in bands, on jaws, vomer, and palatines. Skin covered with rough, bony tubercles. Dorsal and anal fins very small; rostral tentacle present, retractile into a cavity under a bony prominence on the forehead; ventrals present, I, 5, well separated; pectorals large, placed horizontally. Gills $2\frac{1}{2}$. No air bladder; no pyloric cæca. Tropical America, in shallow water; small fishes of singular form, often regarded by the ignorant as venomous. (*ὄγκος*, hook; *κεφαλή*, head; properly written *Oncocephalus*, but Fischer chose the above monstrous spelling.)

a. Snout produced, the rostral process pointed, 6 to 10 in length of body. VESPERTILIO, 3119.

aa. Snout short, the rostral process 12 to 15 times in length of body. NASUTUS, 3120.

aaa. Snout short, the rostral tubercle reduced to a button-like tubercle, which is about 25 times in length of body. RADIATUS, 3121.

3119. OGCOCEPHALUS VESPERTILIO (Linnæus).

(Bat Fish; Diablo.)

Head, from tip of upper jaw to gill opening, nearly ½ the length; depth 5 in length from upper jaw to base of caudal; width 1⅓. D. 4; A. 4; rostral process from 6 to 10 (9 in our specimens from Havana); P. 4½; V. 6; C. 4¼. Body stoutish, much depressed, rostral process longer than in the other species, variable in length; mouth small, the maxillary reaching nearly to posterior margin of eye; villiform teeth in bands, on jaws, vomer, and palatines; interorbital flattish, its width less between anterior edge of eyes than posterior edge; rostral groove longer than broad; body covered with bony protuberances, variable in size, and not very definite in position, lower parts with a shagreen-like covering; posterior edge of pectorals much behind middle of body; ventrals long, reaching outward to edge of the disk-like, anterior part of body; origin of dorsal over posterior edge of pectoral; anal under the vertical of tips of dorsal rays, anal reaching nearly to base of caudal. Pale grayish brown above, reddish below; back with round black spots, conspicuous in life, but growing fainter and sometimes disappearing in spirits; belly in life a coppery red; pectorals nearly plain dusky. Length 12 inches. West Indies, north to the Florida Keys; common in shallow water. Here described from a specimen from Havana, Cuba, about 10 inches in length. The length of the snout is subject to great variation, but it is never short and button-like, as in *O. radiatus*. (*vespertilio*, a bat.)

Lophius vespertilio, Linnæus, Syst. Nat., Ed. x, i, 236, 1758, American Seas; after *Lophius fronti unicorni* of Artedi.

Malthæa vespertilio, Cuvier & Valenciennes, Hist. Nat. Poiss., xii, 440, 1837; prominence on snout 10 in length; De Kay, N. Y. Fauna: Fishes, 167, 1842; Günther, Cat., iii, 200, 1861; Jordan & Gilbert, Syopsis, 850, 1883; Jordan & Swain, Proc. U. S. Nat. Mus. 1884, 234.

Lophius rostratus, Shaw, Zool., iv, 383, pl. 163, 1803.

Malthæa longirostris, Cuvier & Valenciennes, Hist. Nat. Poiss., xii, 452, 1837, Bahia (Coll. Blanchet); snout 6 in length.

3120. OGCOCEPHALUS NASUTUS (Cuvier & Valenciennes).

Head 2. D. 4; A. 4. Rostral process short, about 12 to 15 times in length of body. Cavity of nostril tentacle higher than broad; width of body 2 in length; vent behind middle of body. Dusky above, with round black spots, edged with whitish. West Indies. (Lütken); not seen by us; perhaps a variation of *O. vespertilio*. (*nasutus*, long-nosed.)

Malthæa nasuta, Cuvier & Valenciennes, Hist. Nat. Poiss., xii, 452, 1837, Martinique.

Malthe nasuta, Lütken, Nat. For. Vid. Medd. 1865, 4; Jordan & Gilbert, Synopsis, 850, 1883.

Malthæa notata, Cuvier & Valenciennes, Hist. Nat. Poiss., xii, 453, 1837, Surinam; snout 15 in length; body spotted.

3121. OGCOCEPHALUS RADIATUS (Mitchill).

(SHORT-NOSED BAT-FISH.)

Head 2. Dorsal 4; A. 4. Rostral cavity somewhat broader than high, or width equal to height; distance between anterior angles of orbits about equal to that between the posterior angles; eye a little wider than interorbital width; snout, exclusive of rostral tubercle, not produced beyond the rostral cavity, but with a cylindrical button-like tubercle, slightly contracted at base, pointing obliquely upward and forward, its length 2⅝ times in body; posterior edge of pectoral slightly nearer base of caudal than upper jaw; caudal peduncle very thick and heavy; vent about midway between tip of jaw and base of caudal fin. Color brownish, with dark round spots sometimes edged with white; pectorals with a network of white lines dividing the dark color into dark brown spots; tip of caudal blackish, belly coppery red. Length 8 to 12 inches. Coast of Florida and neighboring waters; very common in shallow bays among weeds, especially about the Florida Keys. Here described from a specimen from Cedar Key, Florida, 7 to 8 inches in length. (*radiatus*, rayed.)

Lophius radiatus, MITCHILL, Amer. Monthly Mag., March, 1818, 326, Straits of the Bahamas.

Malthe cubifrons, RICHARDSON, Fauna Bor.-Amer., III, 103, 1836, said to be from Labrador (Coll. J. J. Audubon), but this is certainly an error; Audubon collected also in Carolina and Florida; GÜNTHER, Cat., III, 203, 1861; JORDAN & GILBERT, Synopsis, 850, 1883; JORDAN, Cat. Fishes, 139, 1885.

? *Malthæa truncata*, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 454, 1837, America; snout wholly obsolete; perhaps a species of *Zalieutes*.

Malthæa angusta, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 454, 1837, Dutch Guiana; snout more than 20 in length.

1073. ZALIEUTES, Jordan & Evermann.

Zalieutes, JORDAN & EVERMANN, Check-List Fish. N. and M. A., 511, 1896 (*elater*).

Disk wider than long, about as long as rest of body (including caudal fin); middle line of head elevated, but the forehead not projecting beyond mouth; rostral tentacle present, the cavity about as wide as high; mouth small; minute teeth on vomer and palatines. Gills 2½. Eastern Pacific. The genus is very close to *Malthopsis*, Alcock,* but the latter, like *Ogcocephalus*, has the disk longer than broad, but the gills are reduced to 2. (ζάλη, surge of the sea; ἁλιευτής, fisher.)

3122. ZALIEUTES ELATER (Jordan & Gilbert).

Body very broad and depressed, the disk considerably broader than long, its width 1⅔ times in length of body; back and snout considerably raised above rest of body; greatest depth of body scarcely more than width of mouth. Mouth small, its width ½ greater than diameter of orbit; snout very short, scarcely projecting beyond mouth, its length

* *Malthopsis*, ALCOCK, Ann. and Mag. Nat. Hist. 1891, 26; *Malthopsis luteus*, from the Andaman Sea; *Malthopsis mitriger*, GILBERT & CRAMER, Proc. U. S. Nat. Mus. 1896, 434, with plate, off Hawaiian Islands.

about equal to interorbital width, shorter than its own width in front. Eye rather large, much longer than snout, wider than interorbital area. Process representing first dorsal spine present, small. Skin covered with spines, which are comparatively slender and sharp, their stellate bases inconspicuous, those on snout and middle of back and tail largest, much slenderer and sharper than in *Ogcocephalus vespertilio;* no spines on ocelli of back; belly rough; under side of tail with tubercular plates; tail depressed toward base of fin. Pectorals $\frac{1}{5}$ longer than ventrals, their length $1\frac{3}{8}$ width of mouth; caudal a little longer than pectoral, $4\frac{1}{3}$ in body. Color light olive, above everywhere thickly and uniformly covered with small round spots of dark brown, these about as large as the pupil and about as wide as the lighter interspaces; a conspicuous ocellus, larger than eye, on each side of back, this ocellus with a bright yellow spot in the center, surrounded by a black ring, around which is a pale ring, and finally a fainter dark one; under parts plain white; pectorals spotted; caudal yellowish at base, with a terminal blackish band. Length 4 inches. Pacific coast of Mexico, south to Panama, in water of moderate depth; very rare near the shore, but obtained in abundance by the *Albatross* at Stations 2794 and 2795, near Panama. (*Elater,* the spring beetle, from the resemblance of the ocelli to the eye-like spots on the back of *Elater.*)

Malthe elater, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 365, Mazatlan. (Type, No. 28127. Coll. Dr. J. U. Bastow.)
Ogcocephalus elater, JORDAN, Proc. Cal. Ac. Sci. 1895, 506.

1074. HALIEUTICHTHYS, Poey.

Halieutichthys, POEY, Proc. Ac. Nat. Sci. Phila. 1863, 83 (*reticulatus*).

Disk subcircular, anteriorly cordiform, the head merging into the body, very large and much depressed; cranial portion not elevated; interorbital space low and narrow; eyes partly superior; mouth terminal, horizontal, the jaws subequal, the lower jaw nearly semicircular; teeth fine, on jaws and palate. Gills $2\frac{1}{2}$; no gill rakers; gill openings anterior to pectoral; rostral tentacle very small, retractile; dorsal and anal few-rayed; pectorals large, the carpus slender; caudal rounded; skin above sparsely armed with stellate tubercles; lower surface smooth. (*ἁλιευτής,* fisher; *ἰχθύς,* fish.)

a. Surface of body covered with brownish reticulations; fins not barred with black. ACULEATUS, 3123.

aa. Surface of body blackish, not reticulate; pectorals with a broad black bar mesially, the tip pale; caudal blackish toward the tip. CARIBBÆUS, 3124.

3123. HALIEUTICHTHYS ACULEATUS (Mitchill).

D. I, 4 or 5; A. 4; V. I, 5; P. 16 to 18; C. 9; gills $2\frac{1}{2}$. Disk cordiform, about as wide as long, its length more than $\frac{3}{4}$ that of body. Body covered above with stout conical spines with stellular bases, largest upon the trunk, upon which they are arranged in about 2 irregular longitudinal

rows on each side of the dorsal; upon the disk they are placed above the principal bones of the skeleton, most abundant upon its cranial portion; a single row of stout spines, usually 3-pointed, on the outer margin of disk, a particularly large one at each outer angle; body entirely smooth below; snout very short, obtuse; bridge over the rostral cavity covered in front with a 3-pointed spine, having on each side a simple spine; short, stout, simple spines upon each supraorbital margin, the front of which is immediately above and behind the cavity containing the nostrils; vertex with several similar spines; many spines closely placed upon the humeral area; numerous short tentacles upon margin of disk and on sides of trunk; supraoral cavity elliptical, small (horizontal diameter $\frac{2}{7}$ diameter of orbit), containing a well-developed, club-shaped, very perceptible tentacle; width of opening of anterior nostril, which is in a short tube, $\frac{1}{2}$ that of posterior nostril, which is not tubular; width of mouth much less than distance between pupils and equaling diameter of orbit. Diameter of orbit $8\frac{1}{2}$ times in distance from snout to base of caudal, 6 times in distance from snout to origin of soft dorsal, $6\frac{1}{2}$ times in distance to origin of anal, 3 times in distance to base of ventrals, and 6 times in distance to angle between pectorals and trunk, $4\frac{2}{3}$ times in distance from snout to gill opening, 6 in greatest width of disk, and nearly 2 in that of trunk; width of interorbital area $\frac{2}{3}$ diameter of orbit. First dorsal ray longest, equal to diameter of orbit; anal fin inserted under third ray of dorsal, with 4 rays, the third or longest very slightly longer than the longest dorsal ray; ventral fins inserted nearly under the middle of the disk, with 1 rudimentary and 5 dorsal rays, increasing in length posteriorly, the last and longest contained 5 times in total length; distances between origins of ventrals $6\frac{1}{2}$ in total length; pectorals with peduncles entirely included in common membrane, with blades far back, horizontal, lying close to trunk, composed of 16 rays, the middle or longest $3\frac{3}{5}$ in total length; caudal fin rounded, composed of 9 rays, the external rays, 1 above and 2 below, simple, the others bifid; length of middle ray equal to that of trunk (measured from junction of pectorals to base of caudal rays) and slightly exceeding the longest pectoral ray. Length of intestine contained $1\frac{3}{5}$ times in total length. Color, body covered above with reticulations of brown, the general hue varying from light yellowish gray to grayish brown, the markings being darker upon darker specimens; pectoral and caudal fins with about 3 dark bars; the terminal bars in young very black; body beneath milky white (Goode & Bean.) West Indies, Gulf of Mexico, and Gulf Stream, in water of moderate depth; taken by the *Blake* and the *Albatross* at numerous stations in depths ranging from 10 to 95 fathoms. "As in *Halicutæa, Dibranchus,* and allies, a rostral tentacle is present in this genus. Among specimens belonging to the Museum of Comparative Zoology there is evidence of the existence of a couple of distinct forms in the West Indian waters. The true *H. aculeatus* is much the lighter in the ground colors and has brownish reticulations across the back, 2 or 3 narrowish transverse bands of the same color across the pectorals, and 2 or 3 similar bands appear on the caudal, the posterior being darkest. The margins of the fins are light in color. The rostrum is acute; it ends in a spine which

turns upward, and, seen from above, it is hardly long enough to cover the tentacular niche. Evidently this type belongs to the shallower waters. The localities noted carry its distribution from the Bahamas to the Yucatan Banks, in depths of 40 fathoms and less." (Garman.) (*aculeatus*, with needle-like spines.)

Lophius aculeatus, MITCHILL, Amer. Mon. Mag., II, 1818, 325, Straits of Bahama.
Halieutichthys reticulatus (POEY MS.) GILL, Proc. Ac. Nat. Sci. Phila. 1863, 91, Cuba (Coll. Prof. Felipe Poey); JORDAN & GILBERT, Synopsis, 851, 1883.
Halieutichthys aculeatus, GOODE, Proc. U. S. Nat. Mus. 1879, 109; GOODE & BEAN, Oceanic Ichthyology, 504, pl. 122, fig. 414a and b, 1896; GARMAN, Bull. Lab. Nat. Hist. Iowa Univ. 1896, 87, pl. 4, fig. 1.

3124. HALIEUTICHTHYS CARIBBÆUS, Garman.

D. I–5; A. 4; V. 5; P. 17; C. 9. Color darker than *H. aculeatus;* the reticulations are not present; the outer half of the pectoral, except at the margin, is black; and, excepting the narrow posterior margin, the hinder fifth of the caudal fin is black; the upper surface is clouded brownish without traces of the network pattern common to *H. aculeatus*. On the specimens described, the rostrum is acute, and the spine extends forward to cover the cavity receiving the tentacle so that it is not visible when viewed from above. West Indies. As now known, this species ranges from Jamaica to Barbados in depths of 70 to 150 fathoms or more. (Garman.) (*caribbæus*, from the Caribbean Sea.)

Halieutichthys caribbæus, GARMAN, Bull. Lab. Nat. Hist. Iowa Univ. 1896, 87, pl. 4, fig. 2, Jamaica to Barbados.

1075. HALIEUTÆA, Cuvier & Valenciennes.

Halieutæa, CUVIER & VALENCIENNES, Hist. Nat. Poiss., XII, 455, 1837 (*stellatus*).

Head very large, broad, depressed, its outline nearly circular; cleft of the mouth wide, horizontal; jaws with small cardiform teeth; no teeth on vomer or palatines. Skin everywhere covered with small, stellate spines. Forehead with a transverse bony ridge, beneath which is a tentacle, retractile into a cavity, the only rudiment of the spinous dorsal fin; soft dorsal and anal very short, far back. Gills $2\frac{1}{2}$, the anterior gill arch without laminæ. Branchiostegals 5; vertebræ 17. Pacific Ocean. (ἁλιευτής, one who fishes.)

3125. HALIEUTÆA SPONGIOSA, Gilbert.

D. 6; A. 4; C. 9; V. 4; P. 12 or 13. This species is remarkable for the soft, spongy texture of the body, and the membranaceous or cartilaginous character of its bones. Width of head $1\frac{1}{5}$ in its length; tail long and slender, the vent midway between base of caudal and articulation of mandible; width of base of tail $4\frac{1}{3}$ in its length; mouth little or not at all overpassed by the snout, its width $2\frac{3}{4}$ to 3 in that of head, lower jaw usually not included; gape of mouth oblique, almost wholly anterior. Teeth in wide cardiform bands in the jaws, none of them enlarged; palate toothless. Interorbital width slightly greater than length of snout, 5 in width of

head; eye 1¾ in interorbital width. Rostral tentacle short, with an expanded 3-lobed tip; front of dorsal midway between base of caudal and occiput; caudal long, rounded, the lower rays more shortened than the upper, the longest nearly ½ width of head; anal rays high, closely bound together, the fin slender, shaped like the intromittent organ of *Gambusia*, the length of its base equaling ⅔ diameter of orbit, its longest ray reaching base of caudal; pectorals long, the posterior rays rapidly shortened, the longest ½ width of head; head and body everywhere with broadly conical, tubercular plates, varying in size, marked with strong lines, radiating from the center; the apex sometimes blunt, more often provided with a slender spine, sometimes bifid or trifid; on the tail these spines become longer and are directed backward; plates along edge of disk not compressed nor specially modified. A deep groove-like channel just behind mandible and following curve of latter, becoming continuous with another deeper channel running just below edge of disk to near base of pectorals; a third groove runs backward from nostrils, uniting with the others, these grooves spanned at intervals by pairs of fleshy tentacles with fringed tips, which spring from the edges of the grooves and meet across them; at the bottom of the grooves under each pair of tentacles is a small fleshy tubercle. Fin rays, at least at base, with series of small curved prickles. Color uniform dusky, the tail sometimes lighter; fins blackish, more or less edged with white. One specimen with the body and tail uniformly light. Pacific coast of Mexico in deep water. Numerous specimens, the largest 4½ inches long, from *Albatross* Station 2992, in 460 fathoms. (Gilbert.) (*spongiosus*, spongy.)

Halieutæa spongiosa, GILBERT, Proc. U. S. Nat. Mus. 1890, 124, west of Revillagigedo Islands, at Albatross Station 2992, Lat. 18° 17′ 30″ N., Long. 114° 43′ 15″ W., in 460 fathoms. (Coll. Gilbert.)

1076. HALIEUTELLA, Goode & Bean.

Halieutella, GOODE & BEAN, Proc. Biol. Soc. Washington, 1882, 88 (*lappa*).

Body subcircular, depressed, its width equal to its length, covered with flaccid, inflatable skin. Spines feebler and less numerous than in *Halieutæa*. Head merged in body; forehead with a transverse bony ridge; no perceptible supraoral cavity; no tentacle. Mouth small, terminal; lower jaw slightly curved forward. Teeth in the jaws minute, cardiform, not discernible on palate, though possibly present. Carpus broad, slightly exserted; pectoral fins remote from tail, obliquely placed, with membranes subvertical. Branchial aperture posterior to carpus, upon the disk, and not remote from its margin. Gills 2½. Dorsal fin 5-rayed, inserted at junction of disk with caudal peduncle; anal fin 4-rayed, originating at root of caudal peduncle. (ἁλιευτήρ, a fisherman.)

3126. HALIEUTELLA LAPPA, Goode & Bean.

D. 5; A. 4; C. 9; P. 15; V. 5. Disk subcircular, more than ⅔ as long as the body. Body covered with a loose, flaccid, inflatable skin, which so obscure its proportion, that it is impossible to determine its exact height,

but it is not nearly so much depressed as in the related genera. When the body is inflated the height and length of the disk is nearly equal. Spines rather feeble; about 10 between snout and dorsal fin; about 6 strong spines, with conical bases and stellular tips, on outer margin of disk on each side, the anterior of them being opposite the eye; in front of these spines on the discal margin, and between them and the snout, are several small, simple spines, pointing backward; belly armed with spines similar to those on the back, but weaker; a stellate spine upon tip of snout, with 2 weaker, simple spines on each side; nasal openings midway between eye and tip of snout; mouth small, upon the margin of the disk; upper jaw shorter than diameter of eye. Teeth as described in the generic diagnosis. Dorsal fin inserted at posterior limit of disk, with 5 simple, articulated rays, its longest ray $\frac{1}{3}$ as long as disk; anal fin with 4 simple, articulated rays, inserted directly beneath fourth ray of dorsal, its second and longest ray $\frac{1}{4}$ as long as disk; caudal twice as long as anal, and slightly longer than caudal peduncle, with 9 simple, articulated rays. Carpus inserted at a distance from snout equal to twice length of longest pectoral ray, which is slightly greater than distance of posterior margin of carpus, at its junction with disk, from vent; number of pectoral rays 15; ventral inserted at a point equidistant from the snout and origin of anal, its longest ray (the fourth) equal to $\frac{1}{2}$ distance of anal fin from snout. Color yellowish white. Gulf Stream. A single specimen, $1\frac{1}{4}$ inches long, known. (*lappa*, the burdock, from its prickles.)

Halieutella lappa. Goode & Bean, Proc. Biol. Soc. Washington, II, 1882, 88, Gulf Stream, at Fish Hawk Station 1151, Lat. 39° 58′ 30″ N., Long. 70° 37′ W., in 125 fathoms; Goode & Bean, Oceanic Ichthyology, 500, pl. 122, figs. 512a and 512b, 1896.

1077. DIBRANCHUS, Peters.

Dibranchus, Peters, Monatsber. Kon. Akad. Wiss. Berlin 1876, 736 (*atlanticus*).

Head merged in body, very large, much depressed, forming a broadly ovate disk, with margin laterally prolonged; cranial portion not elevated; the interorbital area low, narrow, with orbits partly superior; supraoval cavity large, protected above by a transverse bony ridge. Mouth terminal, horizontal, wide; lower jaw convex; teeth in cardiform bands, none on vomer or palatines. Gills 2; no gill rakers; gill openings small, anterior to pectorals. Rostral tentacle retractile, trilobate at tip. Skin with numerous strong stellate spines above and below, those at margins of disk especially strong, 3-pointed. Atlantic; distinguished from related genera by the reduction of the gills to 2 pairs. (*δίς*, two; *βράγχος*, gill.)

3127. DIBRANCHUS ATLANTICUS, Peters.

D. 6 or 7; A. 4; C. 9; P. 13 to 15; V. I, 5; Br. 6; gills 2. Disk orbicular, nearly as wide as long, its length about $\frac{1}{2}$ that of body, its lateral outline prolonged on each side, and terminating in a strong spine armed at the tip with a group of irregularly arranged acicular spinelets. Body covered above with numerous stout conical spines with stellular bases, these

largest upon the trunk, where they are arranged approximately in about 4 irregular longitudinal rows upon each side of the dorsal fin; closely set rows of these stout spines mark the outer margin of the disk, and there is also a cluster of 5 to 7 upon each carpal peduncle; outside of these marginal spines, upon each side, is an irregular marginal row of 5 depressed knife-like spines, each tipped with a crown of 3 acicular spinelets; on the anterior margin of the disk the 2 rows coalesce to a greater or less extent and form a bristling array of closely set spines, some pointing dorsally, some laterally, some ventrally; 2 kinds of spines upon the dorsal surface, in addition to the large ones already described, some large, somewhat remote from each other, conical, stellular; others, much more numerous and filling the interspaces, pickle-like, stellular; belly armed with numerous closely-set spines of a similar kind; snout somewhat projecting, armed with 3 many-tipped spines; a spine-armed ridge in front of the eyes, over the top of the snout; in this 4 spines are conspicuous, 1 in front of each eye, and between these a larger pair in front of the supraorbital ridges; from these last mentioned spines extend spine-armed ridges along the upper margins of each orbit; under the snout is a cavity (horizontal diameter ½ that of the orbit) containing a barbel, pedicelled, with thick, club-shaped, trilobate tip; on each side of this cavity are the nasal openings, which are as in *Halieutichthys*. Width of mouth equal to distance between centers of pupils of eyes. Diameter of orbit contained as follows in other dimensions of the body: In total length 9½; in distance from snout to dorsal 6; same to anal 7; the base of ventrals 3; to angle between pectorals and trunk 5½; to gill openings 5; in greatest width of disk 5½; of trunk 4. Width of interorbital area in diameter of orbit ⅔. Dorsal fin with 6 or 7 rays, the longest (third) 1½ times diameter of orbit and 6 times in total length; anal fin inserted entirely behind dorsal, with 4 rays, the longest (third) about as long as longest in dorsal fin; ventral fins inserted nearly under middle of disk, a little nearer vent than to mandibular symphysis, with 1 rudimentary and 5 well-developed rays, increasing in length posteriorly, the last and longest 6½ times in total; distance between ventral organs 7½ in total length. Pectorals with peduncles slightly exserted, bases included in common membrane, composed of 13 to 15 rays, the longest third or fourth $4\frac{2}{3}$ in total. Caudal fin rounded, consisting of 9 rays, all bifid except the 2 external ones; length of middle ray about ½ that of trunk and exceeding that of pectoral, being contained 4⅓ times in total length. Stomach egg-shaped, intestine somewhat longer than body; liver very wide and large. Color uniform reddish, gray above, slightly lighter below. Deep waters of the Atlantic; very abundant, in about 300 fathoms. Known from the west coast of Africa, off the Cape Verdes, off Barbados, and north in the Gulf Stream to Newport. (*atlanticus*, of the Atlantic.)

Dibranchus atlanticus, PETERS, Monatsber. Kon. Akad. Wiss. Berlin 1876, 736, with plate, West Africa, Lat. 10° N., Long. 17° W., in 360 fathoms (Coll. H. M. S. *Gazelle*); GÜNTHER, Challenger Report, XXII, 59, 1887; VAILLANT, Travailleur, etc., 343, 1888; GOODE & BEAN, Oceanic Ichthyology, 501, pl. 122, fig. 413, 1896.

ADDENDA.

Page 12. After *Entosphenus tridentatus* add:

11(a). ENTOSPHENUS CAMTSCHATICUS (Tilesius).

A lamprey taken by Steller in the Bolschaya River, Kamchatka, has not been recorded by subsequent writers. It is reported by Steller as 13½ inches in length; the head ⅛ of an inch; mouth long, with 2 teeth above, 6 below; dorsals 2. Color shining brassy, dark above; sides with dusky serpentine lines. A figure published by Tilesius shows the upper teeth as bifid, and 9 teeth below.

Pallas describes specimens from the sea at Petropaulski as 7 inches long, not marbled nor variegated. The species of Steller is probably an *Entosphenus*. That of Pallas may be the same, or it may be a *Lampetra*, allied to or identical with *L. aurea*.

Petromyzon marinus camtschaticus, TILESIUS, Mém. Acad. St. Petersburg 1809, 240, with plate, Kamchatka.

Petromyzon camtschaticus, TILESIUS, *l. c.*, 241.

Lampetra variegata (STELLER MS.) TILESIUS, *l. c.*, 247, Bolschaya River, Kamchatka.

?Petromyzon marinus camtschaticus, PALLAS, Zoogr. Ross. Asiat., III, 1810, 67, Petropaulski.

Page 14. From the synonymy of *Lampetra wilderi* omit "*Petromyzon branchialis*, Günther, Cat., VIII, 504, 1870," and after the last synonym add: Not *P. branchialis*, Linnæus, which is the larva of some European species, perhaps of *P. marinus*.

Page 25. In the description of *Catulus uter* the teeth should read $\frac{6}{5}\frac{0}{4}$ instead of $\frac{6}{6}\frac{0}{4}$.

Page 27. In the key, under *dd*, read: Root of tail with a conspicuous notch above.

Page 28. The following key to West Coast species of *Galeus* and *Mustelus* will prove helpful.

a. Eye large, spiracle small, the latter not more than ¼ major diameter of orbit.

b. Mouth broad, snout broadly rounded, mandibular angle of 90° or more. Fins less deeply incised; the lower caudal lobe rounded; pectoral and ventral margins nearly straight. GALEUS CALIFORNICUS, 33.

bb. Mouth narrow, snout long, acute, mandibular angle 60° to 65°. Fins deeply incised; lower caudal lobe acute; pectoral and ventral margins concave. MUSTILUS LUNULATUS, 30.

aa. Eye small, spiracle large, the latter ⅔ to ½ the major diameter of orbit. Snout sharp, mouth narrow, the mandibular angle about 70°. Terminal lobe of caudal broad, obliquely truncate posteriorly. Nostrils very large, their width nearly equaling width of interspace. Fins less incised. GALEUS DORSALIS, 32.

Page 37. After *Carcharhinus henlei* add:

45(a). CARCHARHINUS CERDALE, Gilbert, new species.

Body moderately compressed, not elevated, the depth at front of dorsal not more than $\frac{1}{4}$ greater than the oblique anterior margin of dorsal fin, less than distance from the nostril to first gill slit. Head depressed, the snout flattened, long and narrow, acute; length of snout beyond mouth $\frac{1}{6}$ to $\frac{1}{10}$ greater than distance between angles of mouth in all but one (the largest) of our specimens, in which it is slightly less than width of mouth; $\frac{3}{5}$ to $\frac{4}{5}$ greater than distance from tip of lower jaw to a line connecting angles of mouth; $\frac{1}{7}$ to $\frac{1}{12}$ greater than width of snout opposite outer angle of nostrils. Interorbital width equaling distance from tip of snout to front of eye in young, to middle or posterior border of eye in older individuals; less than $\frac{1}{2}$ distance to first gill opening. Middle of eye nearer nostril than angle of mouth by $\frac{1}{2}$ to $\frac{4}{5}$ its diameter; distance from eye to nostril $\frac{1}{2}$ or slightly more than $\frac{1}{2}$ distance from nostril to tip of snout; middle of nostrils much nearer front of mouth than tip of snout; nasal flap with a very narrow, short, acute lobe, placed at end of inner third of flap; outer angle of nostrils nearly at margin of snout, the inner angles separated by a distance equaling or slightly exceeding that between inner angle of nostril and back of eye. Lips very little developed, the lower entirely concealed in closed mouth, the upper visible as a very short fold. Teeth in lower jaw narrow, erect, serrulate on both margins, more coarsely so toward base; the serration more conspicuous in our smallest specimens (450 mm.), and is obsolescent on some of the teeth in adults; teeth in upper jaw broadly triangular, in front of jaw narrower and erect, those in sides of jaw growing at once broader and more and more oblique; the lateral teeth with a strong notch on the outer side; both margins strongly serrate, the serrations increasing toward base, one or more of those below notch sometimes enlarged and cusp-like in adults; teeth about $\frac{13}{13}$. Conspicuous areas of large and of small pores on underside of head. Gill openings of moderate width, the longest equaling distance between eye and nostril, the fifth much shortened, about $\frac{2}{3}$ length of first. Eye small, equaling length of nasal opening, $1\frac{3}{4}$ to 2 in middle gill slit. Pectoral short and broad, the posterior margin not strongly incurved; tip of fin extending to a vertical intersecting dorsal base at origin of its posterior third or fourth; anterior margin of pectoral 3 times length of inner or posterior margin, the latter less than width of base; first dorsal beginning behind a vertical from axil of pectoral a distance about equaling that which separates eye from nostril; free margin of fin gently concave, the anterior angle extending to a point midway between base and tip of posterior lobe, when the fin is depressed; base of first dorsal $2\frac{1}{2}$ to $2\frac{3}{5}$ in interspace between dorsals; base of second dorsal 7 in interspace between dorsals, $2\frac{1}{3}$ in its distance from anterior margin of pit; origin of second dorsal falling over or behind middle of anal base, the fin but slightly concave, with rounded anterior angle, its posterior angle much produced, the posterior margin exceeding base of fin, which about equals length of anterior margin; anal inserted more ante-

riorly than second dorsal, its base longer, its margin much more deeply concave, the length of base contained about $1\frac{1}{3}$ times in its distance from lower caudal lobe; lower caudal pit in advance of the upper; caudal broad throughout, the lower lobe not falcate, slightly less ($\frac{1}{10}$ to $\frac{1}{4}$) than $\frac{1}{2}$ length of upper lobe, which is about $4\frac{1}{4}$ in total length. Shagreen coarse. Color varying from light to dark gray above, the belly and lower part of sides whitish; fins all dusky or grayish, the caudal often with a blackish border; pectoral with or without a black tip, the latter when present not as conspicuous as in *C. æthalorus*, usually not extended into inner face of fin. A specimen 730 mm. long has the claspers undeveloped, extending slightly beyond margin of ventrals. Another specimen, 850 mm. long, has the claspers fully developed, extending beyond the margin of the ventrals for a distance of 50 mm. Strongly resembling *C. æthalorus*, with which it is associated in the Bay of Panama. It is distinguishable at sight by the narrower gill slits, broader and less falcate fins, and by the much less conspicuous black tips to the pectorals. The dentition is very dissimilar in the two, and makes it necessary to arrange them in different subgenera. Abundant at Panama, where numerous specimens were secured. (Gilbert.) (*κερδάλη*, wary, fox-like.)

Carcharhinus, sp. indescr., JORDAN, Proc. U. S. Nat. Mus. 1885, 363.
Carcharhinus cerdale, GILBERT, Fishes of Panama, MS. 1898, Panama. (Type, No. 11884, L. S. Jr. Univ. Mus. Coll. Dr. Gilbert.)

Page 39. *Carcharhinus nicaraguensis* or its marine original form was found in abundance in the Bay of Panama by Dr. Gilbert.

Page 41. After *Carcharhinus oxyrhynchus* add:

54(b). **CARCHARHINUS VELOX**, Gilbert, new species.

Preoral portion of snout slightly more than $1\frac{3}{4}$ times width of mouth, 5 times distance between nostrils, $1\frac{3}{4}$ times width of snout opposite outer angles of nostrils, $1\frac{1}{4}$ times interorbital width, $2\frac{4}{5}$ times distance from chin to line joining angles of mouth. Nostrils transverse in position, the inner angle nearer mouth than tip of snout by a distance slightly less than length of nostril. Front of eye equidistant from nostril and front of mouth, the middle of eye nearer angle of mouth than nostril; diameter of eye less than nostril, slightly more than $\frac{1}{2}$ longest gill slit. Snout very porous. Folds at angle of mouth slightly longer than usual. Gill slits rather wide, the middle one $1\frac{3}{4}$ times diameter of orbit. Teeth of lower jaw very narrow, erect, very minutely serrulate, appearing entire except under a lens. Teeth in upper jaw very oblique, wide at base, with a deep notch on outer margin, the terminal cusp rather narrowly triangular. Pectoral broadly falcate, the anterior margin convex, the distal edge concave, both angles rounded; tip of pectoral reaching a short distance beyond base of first dorsal; anterior margin of pectoral $2\frac{2}{3}$ times the posterior (inner) margin, about $1\frac{1}{8}$ times the distal edge; first dorsal inserted about the diameter of orbit behind a vertical from axil of pectoral, nearer pectoral, therefore, than ventral; anterior margin concave basally, convex on distal half, the anterior angle rounded; free margin concave, largely owing to the much pro-

duced acute posterior lobe; vertical height of fin exceeding length of base, the anterior lobe very high, extending beyond tip of posterior when the fin is declined, equaling $\frac{7}{9}$ length of anterior margin of pectoral; posterior margin of dorsal $3\frac{1}{4}$ in the anterior margin; base of first dorsal contained $2\frac{1}{2}$ times in interspace between dorsals; base of second dorsal $6\frac{2}{7}$ times; margin of second dorsal gently concave; front margin low, the angle broadly rounded, barely reaching posterior end of base when fin is declined; posterior lobe much produced and acute, slightly longer than base of fin, the latter $1\frac{6}{7}$ in the distance from its base to front of caudal pit; upper lobe of caudal $3\frac{2}{3}$ in total length, the lower lobe $2\frac{1}{2}$ in the upper; terminal lobe of caudal $3\frac{2}{3}$ in the upper lobe; anal larger than second dorsal, higher, with deeply incurved margin, its base a little longer, its origin slightly in advance of that of second dorsal, the posterior insertions of the two fins nearly opposite; length of anal base $1\frac{3}{5}$ in its distance from anterior edge of caudal pit. Color bluish above, whitish or grayish below; free margin of pectoral narrowly white, the anterior edge narrowly bordered with black, most evident when seen from the outer surface, the inner surface being dusky; first dorsal unmarked, the second dorsal with dusky anterior lobe; upper edge of caudal black, the lower margin faintly dusky; fins otherwise unmarked. A single specimen, a female 4 feet long, was procured from the Panama market. As preserved, it is partially skinned. The following measurements were taken when the specimen was intact, before preservation. Where not agreeing with dimensions given above, the latter will be found more reliable:

	Inches.
Tip of snout to insertion of dorsal	$16\frac{1}{4}$
Base of first dorsal	$4\frac{3}{8}$
Distance between dorsals	11
Base of second dorsal	$1\frac{3}{4}$
From second dorsal to front of caudal pit	$2\frac{7}{8}$
Front of caudal pit to tip of caudal	$13\frac{3}{4}$
Tip of snout to axil of pectoral	15
Axil of pectoral to front of base of ventrals	$11\frac{1}{8}$
Front of ventrals to front of anal	$6\frac{1}{2}$
Front of anal to front of caudal pit	$4\frac{9}{16}$
Girth at front of first dorsal	$17\frac{3}{4}$

Distinguished from other known sharks of the Pacific coast of America by the excessively long, slender, acute snout, the slender body, and the very long caudal fin. Panama; only the type known. (*velox*, swift.)

Carcharhinus velox, GILBERT, Fishes of Panama, MS. 1898, Panama. (Type, No. 11893, L. S. Jr. Univ. Mus. Coll. Dr. Gilbert.)

Page 42. *Scoliodon longurio* has the teeth serrulate at base only. The base of the first dorsal is $2\frac{1}{4}$ in the interspace between the dorsals.

Page 44. *Sphyrna tiburo* occur also in the Pacific. We have recently secured specimens at Mazatlan, where *S. tudes* and *S. zygæna* are also found.

Page 47. *Carcharias littoralis* reaches a length of 8 feet 7 inches. (Specimens from Beaufort, North Carolina. Coll. H. H. Brimley.)

Page 49. *Lamna cornubica*, the salmon shark, is abundant and destructive to salmon on the coasts of southern Alaska, especially about Kadiak, where it was seen by us.

Page 53. Under *Squalidæ*, read ovoviviparous for "oviparous."

Page 54. *Squalus sucklii* has been but once recorded from Bering Sea. (Bering Island. Coll. Dr. L. Stejneger.) It is very abundant at Kadiak.

Page 60. *Pristis perrotteti* is not authentically known except from the rivers of Africa. Our west coast species is doubtless distinct and should stand as *Pristis zephyreus*. *Pristis pectinatus* occurs northward at least to Beaufort, North Carolina. (Specimen 12½ feet long. Coll. H. H. Brimley.)

Page 61. After the synonymy of *Pristis pectinatus* insert:

80(a). PRISTIS ZEPHYREUS, Jordan & Starks.

(Pez de Espada.)

Snout to nostrils, 3 in length to base of caudal; breadth of saw at anterior end between first 2 pairs of teeth ½ breadth of its base behind the last pairs; teeth on saw trenchant behind, arranged in 22 pairs; hinder teeth wide apart, the interspaces 5 times their base; posterior teeth turned slightly backward, a groove on their posterior edge; front teeth not quite ½ as long as the saw is broad at their base; distance between first and second tooth 3 times base of first. (Other specimens examined for us by Dr. G. W. Rogers show 18 to 21 pairs of teeth.) Eye equal to spiracle, contained 3 times in base of saw just behind last pair of teeth; width of mouth a little greater than base of saw; mouth with about 65 series of blunt teeth; slant height of pectoral in front a little more than half distance from tip of snout to mouth. Dorsals subequal; first dorsal inserted in advance of ventrals, about ½ its base over ventrals; caudal with a lower lobe, which is equal to slant height of pectoral; tail with a keel on side. Color plain olive gray above, light below. Measurements: Length 50 inches; caudal 7 inches; pectoral 7 inches; dorsal front 5½ inches; snout without nostril 11 inches. Type: A skin in L. S. Jr. Univ. Museum. Common in brackish waters at the mouth of the Rio Presidio, where 1 fine specimen was obtained. The species is also recorded (as *Pristis perrotteti*) by Dr. Gilbert from Mazatlan, and by Dr. Günther from Chiapas. Dr. Günther identifies this species with *Pristis perrotteti* described by Müller & Henle, from the Senegal River. In view of the great difference in the fauna of the Gulf of California from that of equatorial Africa this identification may be questioned, especially as there are several details in which the description of *P. perrotteti* differs from our specimen.

Pristis zephyreus, Jordan & Starks, Fishes of Sinaloa, 383, 1895, Mazatlan, Mexico. (Coll. Hopkins Exped. to Sinaloa.)

Page 62. After *Rhinobatus lentiginosus*, add:

81(a). **RHINOBATUS STELLIO**, Jordan & Rutter.

Disk triangular, its greatest width a little less than ½ the distance from snout to dorsal, and equal to distance from snout to a line connecting points of greatest width. Sides of disk straight, tip of snout rounded, posterior point of pectoral more broadly rounded than snout. Length of snout equal to, or a little less than, ½ greatest width of disk, equal to distance between outer points of anterior gill openings; interorbital width 4 to 4⅜ in snout, a little less than length of eye and spiracle, but about equal to length of nostril; internasal width equal to orbit; spiracle ⅓ length of eye, a prominent curved papilla and a slight ridge in its posterior side. Anterior nasal valve with a long slender flap extending across the nostril; 3 broad flaps on posterior side. Rostral ridges separate for their entire length, width between them at base equal to width of spiracle. Mouth nearly straight, its width 2¾ in its distance from snout and equal to distance between inner folds on posterior side of spiracle. Eye 4½ to 5½ in snout. Width of body at axil of pectorals 1⅓ in snout. Dorsal fins about equal in size and shape, the distance between them 2½ times base of first, the distance between the origins of the two fins equal to snout and about equal to distance from axil of pectoral to origin of first dorsal. Sides of tail with a conspicuous fold. Skin above with a fine uniform shagreen, nearly smooth below except near margins of the disk. A series of very small spines above eye and spiracle, 1 or 2 minute spines on shoulder girdle; the largest spines of body situated along median line of back, extending beyond first dorsal; no spine on snout, but in 2 of the 3 specimens there is a pair of minute spineless plates near its tip. Color dusky brown above, about 7 faint dusky bars on the side of the tail behind first dorsal; uniform pale below; large translucent areas on each side of the snout; back with numerous small light spots, much smaller than pupil, arranged symmetrically but not in the same pattern on the 3 type specimens; 2 or 3 pairs between eyes, a few pairs behind eyes near median line, some below eye, where they approach nearest the margin of disk, usually 1 or 2 on median line, sometimes 2 are confluent, about 40 or 45 pairs in all; axil of pectoral in 1 specimen with a dusky blotch on upper side. This species is most nearly related to *Rhinobatus glaucostigma* of the Pacific coast, differing in having a narrow interorbital, narrower body behind disk, and in the very different color. The description is based on 3 specimens, each about 20 inches. Jamaica. (*stellio*, the starry one.)

Rhinobatus stellio, Jordan & Rutter, Proc. Ac. Nat. Sci. 1897, 91, Kingston, Jamaica. (Type, No. 11851, L. S. Jr. Univ. Mus. Coll. Joseph Seed Roberts.)

Page 66. To the synonymy of *Raja* add: *Cephaleutherus*, Rafinesque, Indice, 61, 1810 (*maculatus*).

The genus *Cephaleutherus*, Rafinesque, was, as Dr. Gill has shown (in lit.), probably based on a monstrous example of the genus *Raja*, in which the pectoral fins were not developed on the snout. It should be transferred to the synonymy of *Raja*, leaving *Myliobatis* as the generic name of the Eagle Rays.

Page 74. After *Raja equatorialis* add:

104(a). RAJA ROSISPINIS, Gill & Townsend.

Snout moderately produced, with a soft, moderately narrow, rostral cartilage and a bluntish tip; interorbital space nearly plane; snout with a number of plates having stellate bases about middle, and many smaller asperities, leaving only the borders of the pectorals and ventrals naked; larger spines with stellate bases are interspersed between the disk and the pectoral rays; back with sparse, coarse prickles; a row of about 26 thorn-like spines, with radiating ridges, extends from the interhumeral area to the dorsal fins; 2 spines on each shoulder, 1 spine above antocular region, another above postocular region, and another behind it about ½ the distance; skeleton soft. Bering Sea; only the type known. (*roseus*, rosy; *spinus*, spine.)

Raia rosispinis Gill & Townsend, Proc. Biol. Soc. Wash., xi, 1897 (Sept. 17, 1897), 231, Bering Sea. (Type, No. 48762, U. S. Nat. Mus. Coll. *Albatross.*)

Raia obtusa,* Gill & Townsend, Proc. Biol. Soc. Wash., xi, 1897 (Sept. 17, 1897), 231, Bering Sea. (Type, No. 48763, U. S. Nat. Mus. Coll. *Albatross.*)

104(b). RAJA INTERRUPTA, Gill & Townsend.

Snout moderately produced, with a very soft attenuated rostral cartilage and a blunt tip; interorbital space concave; mouth small; the width equal to ½ preoral area; entire back covered with very small embedded spines, extending nearly uniformly over the disk and snout, leaving only the tip of the latter naked; a row of compressed, acutely curved, smooth spines along middle of back, extending from the interhumeral region to dorsal, but interrupted along the posterior half of disk, where the spines are absent or obsolete; about 4 spines are in the anterior portion and the series recommences on a line with the emargination of the disk; a single spine on each shoulder and occasionally a rudimentary second; no specialized supraorbital spines. Bering Sea; only the type known. (*interruptus*, interrupted.)

Raia interrupta, Gill & Townsend, Proc. Biol. Soc. Wash., xi, 1897 (Sept. 17, 1897), 232, Bering Sea. (Type, No. 48760, U. S. Nat. Mus. Coll. *Albatross.*)

Page 75. *Raja aleutica* and *Raja abyssicola* were described by Gilbert (not Gilbert & Thoburn) in Rept. U. S. Fish. Comm. 1893 (Dec. 9, 1896), 396 and 397, pls. 20 and 21.

* The following is the original description of this nominal species:
Snout not at all produced, but very bluntly rounded; interorbital space narrow; mouth small, rectilinear; minute distant prickles on the snout, the anterior portion of disk and interorbital area, as well as in a broad median band extending on tail to dorsal and commencing at the interhumeral area; a row of scarcely enlarged acute spines above the eye; an uninterrupted row of unguiform spines with smooth bases extending from the interhumeral area to dorsal fin; 2 similar spines arm each shoulder. Bering Sea; only the type known. (Gill and Townsend.) To which we add: Spines in longitudinal series 23 to 25; width of mouth 1⅔ in preoral area; width of disk 1⅓ times its length; tail a little longer than disk; interorbital width 3 in snout; snout from eye 3⅔ in disk to end of base of ventrals. Color plain brown, rather pale. One specimen 11 inches long, a very young male in very bad condition. Evidently the young of *R. rosispinis.*

Page 78. After *Narcine brasiliensis* insert:

112(a). NARCINE ENTEMEDOR, Jordan & Starks.

(ENTEMEDOR.)

Snout 3¾ in length of disk; preocular part of snout equals preoral; interocular space in snout 1½; width of mouth 2¼. Eye much smaller than spiracle; spiracles edged with small tubercles. Length of disk equal to its width; disk equal to length of tail, without caudal fin; tail with a loose fold of skin on each side. First and second dorsals equal, rounded behind; ventrals large, ending midway between posterior edge of disk and caudal fin. Color pale olive brown, a little clouded with darker; second dorsal edged with pale; dots on head dusky. Two specimens taken in the estuary at Mazatlan, and a third procured by Mr. James A. Richardson in the harbor of La Paz. Specimens had also been obtained by Dr. Gilbert, at Panama, in 1883, but having been destroyed by fire, the species has remained undescribed until recently. Length of largest specimen 20 inches. (The Spanish name *Entemedor* seems to be equivalent to *Intimidator*.)

Narcine entemedor, JORDAN & STARKS, Fishes of Sinaloa, 386, 1895, Mazatlan, Mexico. (Type, No. 1699, L. S. Jr. Univ. Mus. Coll. Hopkins Exped. to Sinaloa.)

Page 81. After *Urolophus nebulosus* add:

115(a). UROLOPHUS UMBRIFER, Jordan & Starks.

Disk round, not wider than long, its length greater than tail; snout pointed, not exserted. Snout from eye 4½ in disk; eyes equal to spiracles; mouth 2 in distance to tip of snout; caudal spine inserted in front of middle of tail; skin perfectly smooth. Color brown above, with blackish cross shades or bars, radiating from the shoulder; a dark band behind eyes, and 1 from eyes; caudal fin dark. Mazatlan. One adult female specimen, the uterus containing 4 young. Occasionally taken with *Urolophus mundus*, but much less common. This is probably not identical with Garman's *Urolophus nebulosus*, being perfectly smooth and different in color. (*umbra*, shade; *fero*, I bear.)

Urolophus umbrifer, JORDAN & STARKS, Fishes of Sinaloa, 389, 1895, Mazatlan, Mexico. (Coll. Hopkins Exped. to Sinaloa.)

Page 82. *Urolophus asterias*, Jordan & Gilbert, is identical with *Urolophus mundus* (Gill), as is shown by specimens recently collected by D.. Gilbert at Panama.

120(a). UROLOPHUS ROGERSI, Jordan & Starks.

Disk broader than long by a distance 2½ times the interorbital width; anterior margins of disk nearly straight, the tip of snout projecting; snout from eye 3¾ in length of disk; eye little smaller than spiracles; width of mouth 2½ times in preoral part of snout; caudal spine inserted in front of middle of tail; skin with minute prickles on margin of pectorals and on middle of back, leaving smooth areas near middle of pectorals and

over branchial arches; 16 to 20 large spinules along median line of back and tail. Color plain brown; caudal fin darker, edged with white. This species differs from *Urolophus asterias* in having a wider disk, more acute snout, much smaller prickles, and fewer spinules on back and tail. Mazatlan. Three specimens obtained in the Astillero, the longest 18 inches in entire length. (This species is named for Dr. George Warren Rogers, a scholarly physician, native of Vermont, but long resident in Mazatlan.)

Urolophus rogersi, JORDAN & STARKS, Fishes of Sinaloa, 388, 1895, Mazatlan, Mexico. (Type, No. 1700, L. S. Jr. Univ. Mus. Coll. Hopkins Exped. to Sinaloa.)

Page 88. In the key at top of page for *Rhinopterinæ* read *Myliobatinæ*.

132. AETOBATUS LATICEPS (Gill).

This species is probably not different from *A. narinari* and may be omitted. We find no differences between specimens from Mazatlan and the West Indies. The following description is based upon Mazatlan specimens:

Length of disc 1⅜ in width; proximal ½ of anterior margin of pectoral fins straight, distal ½ convex; posterior margin concave, the end of each ray forming a small scallop; lateral angle sharp. Snout forming an angle, from its tip to division of nasal lobes, 1⅓ times breadth of head; width of snout 1⅕ times distance from its tip to the division of nasal lobes; nasal lobes projecting back over the mouth; width of mouth 1½ its distance to tip of snout; numerous blunt buccal papillæ around upper dental plate and on ridge between nostrils; interorbital 4¾ in disk; eyes smaller than spiracles, which are as long as base of dorsal. Ventrals well rounded, 3⅗ in length of disk; tail 3½ times disk. First caudal spine equals base of dorsal, which is ½ second spine. Color bluish black with many round yellowish spots scattered equally over the back and ventral fins; spots about as large as eye on back, smaller on head, sometimes two spots run together forming an elliptical spot, about 16 spots from eye along anterior margin of pectoral to lateral angle; posterior margin of pectoral very narrowly margined with white; ventral side pearly white. From the description of *Aetobatus laticeps* this species differs in the following respects: Disk not so broad; tail not so long; width of head and snout less; ventrals not truncated behind; pectorals not margined with blackish; spots on ventrals not assuming the form of ocelli. (Jordan.)

Page 87. For the description of *Pteroplatea crebripunctata* in text substitute the following:

Width of disk twice length to posterior end of anal slit; snout forming a regular curve from a little in front of middle of pectorals, a very small blunt projection at tip; anterior margin of disk convex near snout and lateral angles, pectorals concave medially; posterior margin weakly convex; posterior angle broadly rounded; lateral angle sharply rounded; distance from snout to a line drawn through lateral angles, 2½ times in distance to tip of tail. Interorbital a little wider than its distance to tip of snout; eyes twice spiracles; mouth equals snout, 6½ in disk. Tail rat-

like, with a scarcely perceptible fold of skin on its dorsal side. Ground color olive brown, everywhere with small dark points, not so close set as in *Pteroplatea rava*, indistinct grayish spots, ½ as large as iris, scattered over the body among the dark points, these spots more distinct on anterior edge of disk; tail mottled with darker; lower parts light. Markings nowhere so distinct as in *P. rava*. Very common on sandy shores everywhere about Mazatlan, from which locality it was originally described; also taken by Dr. Gilbert.

Page 87. After *Pteroplatea marmorata* add:

130(a). PTEROPLATEA RAVA, Jordan & Starks.

(MANTARAIA COLORADA.)

Length of disk 1⅞ width; snout forming an angle which is almost a right angle; pectorals slightly concave medially; posterior margin of disk weakly convex; posterior angle not broadly rounded, but curved in somewhat suddenly; lateral angles acute. A line drawn through lateral angles would bisect a line from snout to tip of tail. Interorbital 1⅓ in snout; eye 1½ in spiracles; mouth 7 in disk, 1½ in snout; tail straight and slender, with a very slight fold on dorsal side. Ground color light olive brown, thickly set with sharp-cut black points; conspicuous gray or white spots, ½ as large as iris, scattered over the body, around which the black spots form rings; brighter yellowish spots and half-spots around anterior edge of disk; tail mottled above with darker; lower parts chiefly light orange red or rust-colored in life. All the markings are very distinct and clear cut, the reddish of the belly conspicuous. Mazatlan. One specimen 12 inches long. (*ravus*, reddish.)

Pteroplatea rava, JORDAN & STARKS, Fishes of Sinaloa, 390, 1895, Mazatlan, Mexico. (Type, No. 1587, L. S. Jr. Univ. Mus. Coll. Hopkins Exped. to Sinaloa.)

Page 90. After *Myliobatis californicus* add:

134(a). MYLIOBATIS ASPERRIMUS, Gilbert, new species.

Upper surface of head and body, excepting the snout, an area on outer side of spiracle, the pectoral margin and its posterior angle, and the ventral fins thickly covered with minute, usually stellate prickles, of uniform size, most numerous on median portions of head and back; those on basal ½ or ⅔ of pectoral least crowded and arranged in definite longitudinal series, corresponding with the muscle bands; tail very rough throughout, covered with similar stellate prickles and also crossed by numerous narrow grooves, or indented lines, mostly convex forward, somewhat irregular in position and direction, and not corresponding on the two sides. In the type they follow at an average interval of about 10 mm. Lower side of disk mostly smooth, with some prickles on the basal part of pectorals anteriorly, arranged in lengthwise series, and other patches on lower side of head, belly, and base of ventrals. Color dusky brown above, the anterior portion of pectorals with 8 to 10 narrow transverse bars of bluish

white, most of which break up into series of spots toward outer margin of disk, the posterior ones also breaking up toward middle line; the bars and spots fainter anteriorly, becoming whiter and more intense posteriorly; toward outer angles of disk the bars sometimes separated by intermediate series of light round spots, the bars usually failing to meet across the back; posterior portion of disk including base of tail and upper surface of ventrals covered with round white spots not much larger than pupil, some of those immediately succeeding the bars showing a transverse serial arrangement; top of head with one or more pairs of indistinct light spots; margin of snout and of pectorals blackish; spiracular border black; dorsal with a black blotch posteriorly; underside of head and disk bright white; proximal portion of tail blackish above, lighter below, the entire tail becoming black more posteriorly.

Dimensions of type specimen.

	Millimeters.
Length of disk to front of anus	272
Length of disk to posterior edge of pectorals	338
Width of disk	345
Length of tail (not perfect)	1,215
Greater width of head at origin of pectorals	79
Width of cranium between orbits	45
Width of snout opposite front of eye	55
Tip of snout to middle of nasal flap	60
Length of nasal flap	26
Greatest width of nasal flap	35
Diameter of iris	10½
Width of mouth	33
Distance between anterior gill openings	75
Distance between posterior gill openings	45
Distance from anterior to posterior gill openings	45
Length of spiracle	26
Length of fontanelle	60
Greatest width of fontanelle (at anterior end)	23

Rostrofrontal fontanel scarcely constricted anteriorly, the bounding ridges diverging abruptly at their anterior ends. Nasal flap with a shallow median notch, covering the mouth except the median portion of lower dental plate; posterior margin coarsely fringed. Teeth in each jaw in 1 broad median row, and 3 lateral rows, those of median row about 5 times as broad as long anteroposteriorly.

One specimen, a male, with undeveloped claspers which do not nearly reach edge of ventrals, from Panama. (Gilbert.) (*asperrimus*, very rough.)

Myliobatis asperrimus, GILBERT, Fishes of Panama, MS. 1898. (Type, No. 11895, L. S. Jr. Univ. Mus. Coll. Dr. Gilbert.)

134(b). MYLIOBATIS GOODEI, Garman.

Disk about ⅔ as long as broad; lateral angles acute, bluntly rounded at the apices; posterior angles of pectorals nearly right; snout very broad, short, with a slight prominence in front; the fin, or flange, beneath the eye at the side of the head is very wide, much wider than in either *M. freminvillei* or *M. californicus;* eye very small, without a prominence above in either

male or female (immature specimens); tail less than 2 and more than $1\frac{2}{3}$ times length of the disk; dorsal fin smaller than that of *freminvillei;* teeth in 7 series, much shorter and narrower than those of *freminvillei,* third row about 2 and middle row about 4 times as wide as long. Body smooth. Entire length 29 inches; snout to end of ventrals 11.5, vent to end of tail 18.5, and width of disk 17.5 inches. Olivaceous, darker on the center; white below. The Museum of Comparative Zoölogy has a large specimen which agrees well with this description. Compared with *M. freminvillei,* this species has very small eyes, the pectoral below the orbit is wider than the eyeball, and the fin in front of the skull is but little wider than at its sides. In *freminvillei* the eyeball is twice as wide as the fin beneath it, and the fin in front of the skull is much wider than below the eye. Comparing specimens of about the same size, of the same sex, of *freminvillei, californicus,* and *goodei,* the latter is readily distinguished from the former two by the broad flange at the side of the head, the small eyes, the small teeth, and the broader lateral angles of the pectorals. Central America. (Garman); probably on the Atlantic Coast. (Named for George Brown Goode.)

Myliobatis goodei, Garman, Proc. U. S. Nat. Mus. 1885, 39, Central America. (Types, Nos. 9524 male, and 9529 female.)

Page 91. Family XXVIII should stand as *Aodontidæ,* the name *Mantidæ* being used for a family of *Orthoptera.*

Page 92. After *Aodon hypostomus* insert:

58 (a). CERATOBATIS, Boulenger.

Ceratobatis, Boulenger, Ann. Mag. Nat. Hist., ser. 6, vol. xx, August, 1897, 227 (*robertsii*).

Characters of *Dicerobatis,* Blainville, but the teeth restricted to the upper jaw. (*Κεράς,* horn; *Βατίς,* ray.)

138 (a). CERATOBATIS ROBERTSII, Boulenger.

Band of teeth occupying only $\frac{1}{2}$ width of mouth, its width 10 times in its length; teeth tessellated, hexagonal, 2 to 3 times as broad as long, rugose with numerous obtuse ridges; mouth inferior, wide. Pupil vertically elliptic. Body smooth; pectoral fins with nearly straight, slightly convex anterior and slightly concave posterior border; cephalic fins measuring a little less than width of mouth; spiracles behind the eyes; space between last branchial clefts $\frac{1}{4}$ that between first; dorsal fin between the ventrals; tail slender, without spine, nearly twice length of body.

	Millimeters.
Length of disk, without cephalic appendages	350
Width of disk	780
Cephalic fin	90
Width of mouth	105
Diameter of eye	12
Ventral fin	70
Tail	620

Black above, white beneath. Jamaica. One specimen known. This ray grows to a very great size, but specimens are almost impossible to obtain,

owing to the superstitious fear of the fishermen. (Named for Rev. Joseph Seed Roberts.)

Ceratobatis robertsii, BOULENGER, Ann. Mag. Nat. Hist., ser. 6, vol. XX, August, 1897, 227, Jamaica. (Type in British Mus. Coll. J. S. Roberts.)

Page 105. There is no truth in the statement that *Acipenser medirostris* is poisonous. It is a good food-fish, and on the coast of Washington it is somewhat abundant.

Page 116.

ADDITIONAL NOTES ON TACHYSURINÆ.

In the text of this work, pages 116 to 133, the descriptions of the species of *Tachysurinæ* are for the most part too brief to render certain the discrimination of species. The following additional descriptions of these species will be found useful as supplementary to those given in the text. A slight change in the arrangement of the genera has been found desirable, and 3 new species are added.

70. SCIADEICHTHYS, Bleeker.

Dorsal shield much enlarged, formed like an armorial shield; teeth on palate villiform; posterior nasal openings not connected by membrane; band of palatine teeth extended backward.

165. SCIADEICHTHYS TROSCHELI (Gill).

Head $3\frac{1}{4}$ ($3\frac{5}{6}$ in total with caudal); width of head $4\frac{1}{5}$ ($4\frac{5}{8}$ in total); depth 5 ($6\frac{1}{5}$). D. I, 7; P. I, 12; A. 18. Body comparatively robust, broad anteriorly; head not much depressed, broader than high; eye moderate, 7 to 8 times in length of head; width of interorbital space $1\frac{3}{8}$; breadth of mouth $1\frac{3}{8}$; length of snout $3\frac{1}{4}$. Teeth all villiform; band of vomerine teeth simple, trapezoidal, quadrangular, longer than broad, without division on median line; band of palatine teeth very large, each separated in young specimens from the vomerine band by a narrow toothless line; in old specimens the vomerine and palatine bands are wholly confluent; each palatine band with a narrow backward prolongation on the median line; band of premaxillary teeth broad, about six times as long as wide; lower jaw included. Maxillary barbel nearly or quite reaching gill opening; outer mental barbels about $\frac{2}{3}$ head, the inner nearly $\frac{1}{2}$. Dorsal shield much larger than in most species, shaped like an armorial shield, its posterior margin concave, its anterior end acute, wedged into a deep emargination of the occipital process, the two becoming coossified with age; length of antedorsal plate on the median line 5 to 6 in head, a little more than its width; occipital process short and broad, much broader than long, its median line with a broad keel, its edges nearly straight. Shields all coarsely granular, the granulations anteriorly forming radiating striæ. Fontanel large, claviform, broadest posteriorly, its posterior end about midway between tip of snout and front of dorsal, its greatest

breadth about equal to the diameter of the eye, and $\frac{1}{6}$ its length, a short groove extending backward from its obtuse tip; sides of fontanel bony and granulated for its whole length, the granules extending forward to opposite nostrils. Dorsal spine strong, $1\frac{3}{5}$ in head, moderately compressed; pectoral spine $1\frac{1}{3}$ in head. Axillary pore obsolete. Humeral process coarse, granular, broad, nearly $\frac{1}{2}$ length of pectoral spine; base of adipose fin scarcely $\frac{2}{3}$ length of anal, its posterior margin little free; caudal deeply lunate, small, its upper lobe slightly the longer and narrower, $1\frac{2}{3}$ in head; ventrals not quite reaching anal; vent much nearer base of ventrals than anal. Dark brown, with strong bronze luster above, white below; dorsal dusky, especially above; pectorals blackish; anal dark; caudal rather pale; ventrals usually dark toward the tip, their inner side pale; maxillary barbel dusky; mental barbels pale. This species is not rare along the Pacific coast of tropical America, specimens having been observed at Mazatlan, Punta Arenas, and Panama.

Sciades troscheli, Gill, Proc. Ac. Nat. Sci. Phila. 1863, 171, **Panama.** (Coll. Capt. Dow.)
Arius brandtii, Jordan & Gilbert, Bull. U. S. Fish Comm., II, 1882, 39; description from 28230, U. S. Nat. Mus., 24 inches in length.

The following is the original description of *Sciades troscheli*, Gill: "Dorsal I, 7; anal 16; caudal 11, I, 6; 7, I, 11. The greatest height is contained about $4\frac{1}{2}$ times in the length to the base of the caudal fin, and $5\frac{1}{2}$ times in the total. The caudal peduncle, behind the anal, equals the interval between the snout and the eye, and its least height that between the center of the anterior nostril and the eye. The head in front and on the sides is smooth, and a smooth, oblong, triangular area extends nearly to the vertical from the upper angle of the preoperculum; a triangular area on each side is incurved externally to the narrow anterior extremity, and covered with white pisiform granulations. The dorsal buckler is a pentagon, with a semicircular excavation behind and with its surface rugose. The head enters 3 times in the length before the end of the anal fin and more than 4 times in the total; its width equals the interval between the snout and upper angle of preoperculum, and the interocular area equals $\frac{1}{2}$ the head's length. The eye is elliptical, and its diameter is contained $6\frac{1}{2}$ times in the head's length. The distance of the posterior nostril from it equals a diameter. The maxillary barbels extend to about the middle of the pectoral; the outer mental to its base, and the inner mental are $\frac{2}{3}$ as long as the outer. There are 3 villiform patches on the palate which are almost contiguous, and together describe an arch in front; the median patch is small, rather transverse, and widest toward the front; the outer are oblong, subtriangular. The band of the upper jaw is nearly uniform and quite wide; the lower, interrupted at the symphysis, is nearly $\frac{1}{2}$ as wide as the upper, and is narrowed toward its ends. The dorsal spine enters $1\frac{1}{2}$ times in the head's length; has in front, first, minute teeth pointed downward, and then a row of small pisiform tubercles; teeth pointed downward on its hinder border. The first ray is little higher than the spine. The anal commences at a distance from the snout $3\frac{3}{5}$ times as great as that from the base of the caudal

fin; its length enters $6\frac{2}{3}$ in the length, exclusive of the caudal, and when bent back it reaches to the supernumerary caudal rays; the greatest height nearly equals the length. The pectoral fins extend rather beyond the base of the dorsal and exceed $\frac{1}{5}$ of the length, exclusive of the caudal; the spine equals that of the dorsal. The ventrals are inserted midway between the base of the pectoral spines and the axil of the anal, and extend to the origin of the anal. The fins are almost blackish." A single specimen is in the collection of Captain Dow from Panama. The type of *Sciades troscheli* is now lost. At our request, Dr. Gill has again considered this description, in connection with the species now known from the coast. He is positive that his type of *troscheli* had the large dorsal shield characteristic of *brandtii*. Apparently Dr. Eigenmann is right in regarding *troscheli* and *brandtii* as identical.

166. SCIADEICHTHYS EMPHYSETUS (Müller & Troschel).

Head $3\frac{1}{3}$; depth 6. D. I, 7; A. 18. Closely related to *Sciadeichthys troscheli*. Depth little greater than the width. Profile straight, less steep than in *S. troscheli*. Depth of the head $1\frac{3}{4}$ in its length, its width $1\frac{1}{4}$. Top of the head sparsely and coarsely granular, the granulation extending forward only to middle of cheek; fontanel bordered anteriorly by smooth ridges; occipital process coarsely and closely granular, without a prominent keel, its margins convex, its tips emarginate, not coossified with the dorsal plate; dorsal plate shield-shaped, not keeled, its surface irregularly pitted, its margin more finely graven, its length about $3\frac{1}{3}$ in the head. In the specimen examined the dorsal plate seems to have been at some time slightly broken in front, a small, narrow, sharp process of the occipital process fitting into the split. Eye small, 3 in the snout, 11 in the head, 5 in the interocular. Maxillary barbels flattened, reaching to below middle or end of dorsal fin, postmentals not quite to base of pectorals. Upper jaw slightly projecting; all the teeth minute, villiform, the vomerine patch emarginate in front and behind, joined to the subtriangular palatine patches; pterygoid patches long-elliptical. Gill membrane with a narrower free margin than in *troscheli*. Distance of dorsal fin from tip of snout $2\frac{2}{3}$ in the length, the dorsal spine $1\frac{2}{7}$ in the head, granular in front, recurved teeth on its inner margin. Distance between the dorsal and adipose fins $3\frac{1}{7}$ in the length; adipose fin about as long as the dorsal fin. Caudal deeply forked, the upper lobe longer, $3\frac{3}{8}$ in the length; anal little longer than high; ventrals reaching nearly to the anal, about 2 in the head; pectoral spine $1\frac{1}{4}$ in the head, its outer margin granular, the inner rather finely toothed. The skin on the dorsal surface of the head and humeral region finely reticulate with mucous canals. Yellowish brown, lighter below, the fins yellowish, finely punctulate. One specimen 0.51 m. Surinam. (Eigenmann & Eigenmann, Nematognathi, 53.)

167. SCIADEICHTHYS TEMMINCKIANUS (Cuvier & Valenciennes), text, p. 122.

168. SCIADEICHTHYS FLAVESCENS (Cuvier & Valenciennes), text, p. 123.

169. SCIADICHTHYS MESOPS (Cuvier & Valenciennes).

170. SCIADEICHTHYS PROOPS (Cuvier & Valenciennes).

Head 4 to $4\frac{1}{3}$; depth 7. D. I, 7; A. 18; eye $1\frac{1}{4}$ to $1\frac{1}{2}$ in snout, $5\frac{1}{2}$ to 8 in head, $1\frac{3}{4}$ to $2\frac{2}{3}$ in the interorbital, $2\frac{1}{4}$ to $3\frac{3}{4}$ in the interocular. Slender and elongate, broader than deep. Head depressed, its width $1\frac{1}{3}$ in its length, its depth 2, width at mouth 2; anterior portion of the head flat above; top of the head, humeral process, and dorsal plate coarsely granular, the granules arranged in series along the fontanel. Occipital process mucronate, broader than long; dorsal plate large, butterfly-shaped. Opercle striate; fontanel $1\frac{1}{2}$ times as long as the eye, its center in front of the middle of the eye, continued as a shallow groove. Jaws subequal; teeth all villiform, the intermaxillary band very wide and shallow; teeth on the roof of the mouth in 6 contiguous patches. Gill membranes meeting in an angle, forming a broad fold across the isthmus; gill rakers 5 + 10. Pectoral pores large; vertical series of pores. Distance of dorsal spine from the snout $2\frac{4}{5}$ in the length; the dorsal spine granular in front, striate on the sides, weakly serrate behind, its length $1\frac{1}{4}$ to $1\frac{1}{2}$ in head; space between dorsal and adipose fins $2\frac{4}{5}$ to 3 in length, the adipose fin little shorter than the dorsal, the posterior margin free. Caudal deeply forked, its upper lobe longer, 4 to $4\frac{1}{2}$ in the length; anal emarginate, as high as long, 2 to $2\frac{1}{3}$ in head; ventrals 2 in head; pectoral spine roughened or granular in front, serrated behind, $1\frac{1}{5}$ to $1\frac{1}{3}$ in head. Plumbeous above, with blue luster, white below; maxillary barbels dark, the mental barbels white; fins all more or less dotted with brown. Five specimens 0.25 to 0.46 m. long. Pernambuco. (Hartt & Fletcher.) Northern coast of South America to Pernambuco. (Eigenmann & Eigenmann, Nematognathi, 57.)

171. SCIADEICHTYS PASSANY (Cuvier & Valenciennes), text, p. 124.

172. SCIADEICHTHYS ALBICANS (Cuvier & Valenciennes), text, p. 124.

71. SELENASPIS, Bleeker.

Dorsal shield much enlarged, truncate before, in the adult; palatine teeth villiform, the patch extended backward in the adult; posterior nasal openings connected by membrane.

173. SELENASPIS HERZBERGII (Bleeker).

Head $3\frac{2}{3}$ to $3\frac{3}{4}$; depth 5 to 6. D. I, 7; A. 18; eye $1\frac{3}{4}$ to $2\frac{1}{2}$ in snout, $5\frac{1}{2}$ to 8 in head, $2\frac{1}{3}$ to 4 in interocular. Elongate, the width as great or greater than the depth. Width of the head $1\frac{1}{2}$ to $1\frac{1}{3}$ in its length, at the angle of the mouth about 2; depth $1\frac{2}{3}$ to $1\frac{3}{4}$ in its length. Humeral process, dorsal plate, top of head to between the eyes, granular. Occipital

process wider than long, scarcely keeled. Fontanel not continued behind the eyes, and without backward projecting groove; posterior nostrils connected by a membrane. Barbels flattish, those of the maxillary reaching to near the ventrals, to middle of pectorals in older individuals; postmental to or beyond base of pectoral, mental to gill opening. Teeth villiform; vomerine and palatine patches of about equal size and shape in the young; a separate patch behind the palatines is developed later. Gill membranes meeting in an angle, forming a fold across the isthmus; gill rakers 6 + 10. Distance of dorsal spine from snout 2½ to 2⅔ in the length; dorsal and pectoral spines subterete, the outer margins roughened, the sides striate; the dorsal spine slightly serrate behind, a little shorter than the pectoral spine, 1⅔ to 1⅗ in the head; pectoral spine strongly serrate behind; space between dorsal and adipose fins 3⅗ to 4 in the length; adipose fin as long as the dorsal; upper caudal lobe longer, about 4 in the length; anal as high as long, 2 in head; ventrals 1⅗ to 2 in head; pectoral pore minute; sides with vertical series of pores. Color plumbeous above, silvery on sides; fins dusky. The specimens examined measure from 0.14 to 0.38 meters. Para; Curuca; Bahia. (Eigenmann & Eigenmann, Nematognathi, 59.)

174. SELENASPIS DOWII (Gill).

Head 3½; depth 6. D. I, 7; A. 19; eye small, elliptical, 3 in snout, 12 in head, 6 in interocular. A narrow flap of skin across the snout connecting the posterior nasal openings. Width below the dorsal spine a little greater than the depth, less than the width at the humeral process, which equals the greatest width of the head measured at the opercles. Head depressed, its depth at base of occipital process 1⅓ in the greatest width, becoming gradually more depressed forward; width at angle of mouth 1⅗ in length of head, its greatest width about 1⅓ in its length; snout short, 4 in head. Top of head coarsely granular, the granules forming striæ in front, vermiculations posteriorly or, in places, more or less regular striæ. Occipital process truncate, its width at tip greater than its length, the dorsal plate large, saddle-shaped, its bony tubercles forming striæ which are parallel with the strongly convex margin of the "saddle"; opercular bones granular striate, the humeral process with bony tubercles. Fontanel nearly obsolete, the granular bony surface being separated in front by thick skin, which covers an elongate area about 7 times longer than wide. Maxillary barbels reaching beyond humeral process; postmental barbels beyond gill opening, the mentals shorter. Upper jaw produced, equal to the short diameter of the eye. Teeth of the intermaxillaries in a villiform band which is narrowed in front, not produced backward to the angle of the mouth; vomer with a rather broad band confluent with the much wider subquadrate palatine patches which are produced backward in an angle; ovate patches on the pterygoids separate from the palatine teeth; teeth of the lower jaw in a comparatively shallow band, tapering very gradually to the angle of the mouth; the teeth of the jaws minute villiform, those of the palate and

vomer bluntly conical. Gill membranes broadly united, meeting in an angle, joined to the isthmus, but with a free margin; gill rakers 9 + 15. Distance of dorsal from end of snout $2\frac{3}{4}$ in the length. Dorsal spine granular on sides and in front, about $\frac{1}{2}$ the length of the head in height; distance of the adipose fin from the dorsal $3\frac{1}{4}$ in the length, the height of the adipose fin about $2\frac{1}{2}$ in its length, which is contained $2\frac{3}{5}$ in the length of the head. Caudal deeply forked, $5\frac{1}{2}$ in the length; anterior $\frac{2}{3}$ of the anal strongly convex, the posterior $\frac{1}{3}$ slightly emarginate, the highest ray about $2\frac{1}{2}$ in head. Ventrals reaching to anal, about 2 in head, their distance behind the dorsal equal to the length of dorsal and $\frac{1}{2}$ the dorsal plate; pectoral spine granulose on sides, the outer margin with a series of larger granules which become recurved notches toward the tip, the inner edge with recurved hooks, its height $1\frac{3}{5}$ in the length of the head; a small pectoral pore; no evident series of vertical pores. Bluish gray above, becoming white below; the fins brownish with dots. Description from the type of *Arius alatus*, .68 m. long, from Panama; collected by Dr. Steindachner. (Eigenmann & Eigenmann, Nematognathi, 61.)

Selenaspis dowii is thus characterized by Jordan & Gilbert:* Head 4 ($4\frac{3}{5}$ with caudal); depth $6\frac{1}{2}$ ($7\frac{1}{2}$); width of head $5\frac{1}{2}$. D. I, 8; A. 4, 12. Length (29529, U. S. Nat. Mus.) 10 inches. Body elongate, narrow, and slender, the caudal peduncle $1\frac{2}{3}$ in head. Head low and narrow, tapering anteriorly, the snout subtruncate. Eye small, 7 in head, placed rather high; interorbital space little arched, with ridges and depressions, $2\frac{1}{4}$ in head; snout $3\frac{1}{3}$ in head; breadth of mouth $2\frac{1}{3}$ in head. Mouth moderate, with thinnish lips; teeth villiform, bluntish; vomerine teeth forming 2 smallish, rounded patches, separated by a moderate interspace; each patch confluent with the neighboring palatine patch, which is rounded and rather large; the suture marked by a constriction; palatine bands without backward prolongation; premaxillary band of teeth broad. Barbels very long; maxillary barbel extending well beyond tip of pectoral fin; outer mental barbel reaching well past front of pectoral; inner $2\frac{1}{4}$ in head. Dorsal shield comparatively large, not distinctly crescent-shaped, its divisions produced backward, their length about twice the length of the shield on the median line; anterior margin with 2 emarginations, the point fitting into an emargination of the occipital process; dorsal shield without keel. Occipital process very broad and short, its edges nearly straight, its breadth at base considerably greater than its length; its median line with a rather low keel. Fontanel broad and very short, ending obtusely at a point not far behind eye, the distance from this point to tip of snout $1\frac{3}{5}$ in its distance from base of dorsal; each side of fontanel with a conspicuous smooth ridge, the 2 ridges converging anteriorly; shields of head rather finely granulated, few of the granulations forming lines, none of them extending farther forward than posterior margin of eye. Opercle striate. Gill membranes meeting below in a sharp angle, forming a rather broad fold across isthmus. Dorsal spine very short, its length a trifle less than pectoral spine, $2\frac{1}{4}$ in head. Axillary pore obsolete. Humeral process granulated, rather narrowly triangular, a

*Bull. U. S. Fish Comm., II, 1882, 50.

little less than ½ length of pectoral spine, which extends barely ⅔ the distance to the ventral fins; adipose fin long and low, very nearly or quite coterminous with the anal; caudal narrow, rather short, the upper lobe the longer, 1⅔ in head; anal rather low and short; ventrals short, the vent not far behind them. Color dusky above, pale below, the fins all more or less dusky; maxillary barbels dusky, others pale. A single young male was obtained at Panama.

71(a). ASPISTOR, Jordan & Evermann, new genus.

Aspistor, JORDAN & EVERMANN, new genus (*luniscutis*).

This genus differs from *Selenaspis* in the presence of granular teeth on the palate and in the absence of a membranaceous flap connecting the posterior nostrils. (ἀσπιστήρ, a shielded warrior.)

175. ASPISTOR LUNISCUTIS (Cuvier & Valenciennes).

Head 3⅔; depth 5½ to 6. D. I, 7; A. 16 to 19; eye 2 to 3 in snout, 6 to 9 in head, 3 to 4½ in the interocular. Body comparatively stout, the greatest width equaling the greatest depth. Head large, flattish above; profile descending; width of head 1⅙ in its length, width at the mouth 2 to 2⅔, its depth at the base of the occipital process scarcely less than its greatest width; top of head coarsely granular in young, the granules becoming finer and more regularly arranged in the adult; opercles smooth; humeral process with radiating lines of granules. Occipital process variable in shape, broader than long, the posterior margin convex; dorsal plate variable in outline, rounded anteriorly, saddle-shaped, either broader than long or longer than broad; middle of the fontanel above the posterior margin of the eye; the fontanel divided into 3 by 2 bony ridges, the middle portion being more than ½ of its whole length. Sides of head with reticulating mucous canals. No skinny flap connecting the posterior nostrils. Maxillary barbels extending little beyond the base of the pectoral, or shorter; mental barbels short. Upper jaw little produced; teeth in the jaws rather large, conical; teeth of vomer and palatines finely granular, the vomerine patches separated from each other and from the palatine patches in the young, united and covering almost the entire roof of the mouth in the adult; the inner margins of the palatine patches approximated, sometimes a small elliptical patch of teeth between. Gill membranes forming a broad marginal flap across the isthmus. Gill rakers 3 to 4 + 7 to 9. Axillary pore minute or wanting; vertical series of pores present. Distance of dorsal from tip of snout 2½ in the length; the spine 1½ to 1⅔ in the head, granular in front, scarcely serrate behind; distance of adipose fin from the dorsal 3¾ to 4 in the length, the adipose fin twice as long as high, adnate, as long as the dorsal fin; caudal forked, the upper lobe longer, 4½ to 4⅘ in the length; anal fin about as long as high, 2⅓ to 2½ in head; ventrals 1⅔ to 2 in head; pectoral spine stout, 1¼ to 1⅐ in head, granular in front (serrate in the very young), striate on sides, serrate along inner margin. Color purplish brown above, sprinkled with brown dots below; fins about the color of the back. Numerous specimens

examined 0.11 to 0.44 m. long. Porto Alegre; Bahia; Nazareth, near Bahia; Rio Janeiro; Pará; Porto Seguro; São Matheos; Cannavierias. (Eigenmann & Eigenmann, Nematognathi, 63.)

176. SELENASPIS PARKERI (Traill), text, p. 125.

72. NETUMA, Bleeker.

Dorsal shield small, lunate; teeth on palate villiform, the patch on each side with a backward-extending process or angle.

Subgenus NOTARIUS, Gill.

Occipital process constricted at base.

177. NETUMA GRANDICASSIS (Cuvier & Valenciennes).

Head $3\frac{2}{5}$ to $3\frac{3}{4}$; depth $5\frac{3}{8}$ to 6. D. I, 7; A. 18; eye 3 to $3\frac{1}{2}$ in snout, $8\frac{1}{2}$ to 10 in head, 4 to $4\frac{1}{2}$ in interocular. Body cylindrical in front, tapering to a slender caudal peduncle. Head greatly depressed, profile almost straight, descending, the width of the head $1\frac{1}{8}$ to $1\frac{2}{5}$ in its length, its depth $1\frac{1}{5}$ to 2 in its length. Occipital process with a deep constriction where it joins the occiput, shaped like a clover leaflet, much as *Felichthys panamensis*, sometimes broader than long, sometimes much longer than broad, sometimes keeled. Center of the fontanel over the middle of eye, the fontanel not continued backward as a groove; occipital process, top of head, and humeral process granular; interorbital region with 4 ridges, the inner ones bounding the fontanel, the outer ones running obliquely backward from near the posterior nasal opening. Maxillary barbels reaching to the base of the pectoral, mentals to gill opening, postmentals a little longer. Upper jaw projecting a diameter of the eye or more, the lip very wide, especially in front, making the nose pointed; teeth of both jaws rather large, those on the palate somewhat smaller; the depth of the intermaxillary band 7 to 9 in its width; the mandibulary band very shallow; vomerine teeth none in 3 of the examples, a small patch on one side in another specimen, and a small patch on each side in another; palatine patches triangular, produced backward. Gill membranes meeting in an angle, forming a fold across the isthmus; gill rakers 6 + 10. Distance of dorsal spine from snout $2\frac{1}{8}$ to $2\frac{1}{2}$ in the length, the spine broken in the specimen studied. Distance of adipose fin from the dorsal $3\frac{3}{5}$ to 4 in the length; adipose fin at least as long as the dorsal fin, adnate. Caudal fin forked, the upper lobe longer, about 5 in the length, the tips broken; anal fin apparently longer than high, but the rays are somewhat worn off; ventrals small; pectoral pore large, slit-like. Color light brown above, somewhat smutty below from the occurrence of minute scattered dots. We have examined 4 specimens from 0.23 m. to 0.33 m. long, collected by Agassiz & Bourget, Thayer Expedition, at Maranhao, and a fifth, 0.21 m. long, collected by Professor Agassiz at Bahia. (Eigenmann & Eigenmann, Nematognathi, 65.)

178. NETUMA STRICTICASSIS (Cuvier & Valenciennes), text, p. 126.

Subgenus **NETUMA.**

Occipital process not constricted.

179. NETUMA DUBIA (Bleeker), text, p. 126.

180. NETUMA KESSLERI (Steindachner).

Head 3⅛ (4 in total with caudal); depth 6 (7 in total); width of head 4. D. I, 7; A. IV, 13. Length (29252, U. S. Nat. Mus.) 14 inches. Body rather long and low; the head long, broad and much depressed, much broader than deep. Eye very small, about 10 in head, placed well above the mouth. Interorbital space 2 in head; snout 3⅛; breadth of mouth 2. Mouth large, with thickish lips, the upper jaw considerably projecting. Teeth all villiform, rather pointed. Vomerine patches rather large, roundish, usually fully confluent into a trapezoidal band, without division on the median line, and separated by a very narrow groove from the palatine bands. Palatine bands very large, broadly triangular, with a backward prolongation from the inner margin. (Teeth on vomer and palatines all forming one continuous band in old specimens, according to Steindachner.) Bands of teeth in jaws broad, the jaws strong. Barbels rather short and very slender, the maxillary barbels reaching little past base of pectoral; outer mental barbels about reaching gill opening; inner about as long as snout. Antedorsal shield short, crescent-shaped, rough, but without median keel. Occipital process long, narrowly triangular, its edges straight, its length ¼ to ⅓ more than its width at base, its median line sharply keeled. Fontanel broad and shallow, its posterior end obtuse or almost truncate, its tip not prolonged in a groove, its edge bounded by a bony ridge, which is not granulated in front of middle of eye; end of fontanel about midway between tip of snout and front of dorsal, its greatest width about equal to length of eye. Shields of head all very coarsely granular, the roughnesses extending forward about to the eye. Gill membranes forming a broad free fold across isthmus. Dorsal spine moderate, a little more than ½ head, about equal to pectoral spine; humeral process triangular, granular, not quite ⅔ length of pectoral spine; axillary pore obsolete; adipose fin long and low, its posterior margin little free; caudal short and broad, the upper lobe the longer, 1⅔ in head; anal and ventrals rather small, the vent close behind the latter. Color dark brown, with bronze reflections; belly white; fins all dusky in 1 specimen, in the other mostly pale; maxillary barbels dusky, others pale. Two large specimens obtained at Panama by Gilbert. (Jordan & Gilbert, Bull. U. S. Fish Comm., II, 1882, 40.)

181. NETUMA INSCULPTA (Jordan & Gilbert).

Head 4 (4⅜ in total); depth 5⅜ (6¼); width of head 4⅗. D. I, 6; A. IV, 14. Length (29415) 13½ inches. Body moderately elongate, little compressed, the caudal peduncle slender and short. Head shortish, low and

broad, anteriorly depressed. Eye rather large, $6\frac{1}{2}$ in head, placed rather high. Interorbital space flat and nearly smooth, 2 in head; snout 3; breadth of mouth $1\frac{3}{5}$; snout very bluntly rounded, almost truncate in front. Mouth large; teeth all villiform; vomerine bands of teeth large (fully confluent with each other in the type, partly separated in smaller examples), and with the large, club-shaped band on the palatines, from which they are separated by a slight furrow and constriction; palatine band of teeth with a backward prolongation; premaxillary band of teeth large; maxillary barbel long, somewhat compressed, extending to middle of pectoral spine; outer mental barbel reaching base of pectoral spine, inner 2 in head. Dorsal shield short, crescent-shaped, without median keel, its tips produced, its length on the median line about $\frac{2}{3}$ the length of 1 of its halves. Occipital process about as broad at base as long, with a moderate median keel, its lateral margins somewhat concave; fontanel becoming gradually contracted at a point a little nearer base of dorsal than tip of snout, thence forming a narrow groove, which extends to within a diameter of the pupil of the base of the occipital process; this groove sometimes nearly obsolete; greatest width of fontanel about $\frac{2}{3}$ diameter of eye. Granulated striæ extending along the sides of the fontanel to a point opposite or in front of middle of eye. Shield of head finely and evenly granulated, the roughnesses more uniform than usual, and many of them arranged in lines, especially anteriorly; opercle not striate, the skin marked with fine vermiculations; gill membranes forming a broad fold across the isthmus. Dorsal and pectoral spines long, about equal, $1\frac{1}{3}$ in head. No axiliary pores; humeral process very large, triangular, finely granular, about $\frac{1}{2}$ as long as pectoral spine; adipose fin large, without free tip; upper lobe of caudal the longer, $1\frac{1}{5}$ in head; anal and ventrals moderate, the vent close behind the latter. Color rather pale; belly pale; fins and barbels all pale, or but slightly tinged with dusky. A single adult male was obtained by Dr. Gilbert at Panama. Two smaller ones are in the Museum collection, also from Panama. (Jordan & Gilbert, Bull. U. S. Fish Comm., II, 1882, 41.)

182. NETUMA PLANICEPS (Steindachner).

Head 4 ($4\frac{4}{7}$ in total); depth $5\frac{1}{3}$ ($5\frac{2}{3}$); width of head 5. D. I, 7; A. IV, 13. Length (29417) 11 inches. Body comparatively elongate; the head small, rather narrow, depressed anteriorly; the snout rather narrow and moderately rounded. Eye moderate, placed well above mouth, its length $5\frac{1}{2}$ in head. Interorbital space flat and smooth, $2\frac{1}{4}$ in head; snout $3\frac{1}{5}$; breadth of mouth 2. Mouth rather large, with thickish lips; teeth villiform; vomerine bands moderate, confluent with each other and with the much larger ovate palatine bands, a slight constriction or furrow making the divisions; palatine bands each with a backward prolongation; premaxillary band moderate; barbels very short; maxillary barbel scarcely or not reaching to base of pectoral; outer mental barbel scarcely past gill opening below; inner shorter than snout. Dorsal shield short, anteriorly truncate, not keeled, the length on the median line about $\frac{1}{2}$ of 1 of its halves. Occipital process subtriangular, rather narrow, truncate

behind, its margins straight, becoming concave forward, its width at base about equal to its length; fontanel an almost obsolete groove, its posterior end not reaching base of occipital process by about the diameter of the eye, the groove extending forward to a point about midway between tip of snout and base of dorsal spine; anterior to this point is an equilateral triangle, flat, covered with smooth skin, the base of the triangle formed by the smooth, flattish interorbital area. Shields of head rather coarsely granular-striate, the granulations beginning anteriorly about opposite posterior margin of eye; opercle scarcely striate; gill membranes forming a moderate fold across the isthmus. Dorsal spine high, about equal to pectoral spine, and but little shorter than head; no axillary pore; humeral process triangular, granulated, a little more than ⅓ length of pectoral spine; adipose fin rather long; upper lobe of caudal the longer, a little shorter than head; ventrals and anal moderate. Color brownish, not very dark; belly pale, thickly speckled with brown; fins more or less dusky; maxillary barbels black; mental barbels pale. Two specimens were obtained by Dr. Gilbert at Panama. They disagree in several details from Steindachner's description, and it is barely possible that they belong to a different species. The head in Steindachner's types is 3¾ to 3⅘ in length, and the occipital process is narrower and less widened anteriorly. (Jordan & Gilbert, Bull. U. S. Fish Comm., II, 1882, 42.)

183. NETUMA PLATYPOGON (Günther).

Head 3⅔ (4⅔); width of head 4⅔; depth 5⅕ (6⅔). D. I, 7; A. IV, 14. Length (28286) 15½ inches. Body rather elongate, the head not very broad nor much depressed, a little broader than deep. Eye rather large, 5 to 6 in head. Interorbital space slightly more than ⅓ head, a trifle less than width of mouth; length of snout 3⅕ in head. Teeth all pointed; bands of vomerine teeth small, roundish, their boundaries traceable by a slight depression in the young, in the adult fully confluent with each other and with the palatine bands; palatine bands broad, ovate, several times as large as the patches on vomer, continued backward over the pterygoid region; premaxillary band rather broad, 5 to 6 times as broad as long; maxillary barbel reaching past base of pectoral in the young, not to gill opening in the adult, its base a little broader and more compressed than usual; outer mental barbels 2 in head; inner 2½. Dorsal shield very short, lunate, subtruncate in front, its breadth more than 3 times its length on the median line; occipital process long, triangular, with straight margins, its length about 1⅔ times its width in front, its broad median line rather sharply keeled. In the young it is proportionally shorter, little longer than broad. At the beginning of this keel is the end of the long, narrow, groove-like fontanel, which extends forward to a point just behind the eye, where it merges into the flattish and smooth anterior part of the head. Shields of the head all finely granular, the granules rarely forming distinct lines. Dorsal spine long, 1⅛ to 1½ in head, the soft rays projecting beyond the spine; pectoral spine about as long as dorsal, sharply serrate behind, the anterior

serræ not very sharp; axillary pore small or absent; humeral process nearly smooth, rather narrow and short, ½ length of pectoral spine; adipose fin short and rather high, its base barely ⅔ length of base of anal; caudal deeply forked, its upper lobe the longer and slightly falcate, about as long as head; ventrals very short, reaching anal in females, shorter in the males; vent nearer base of ventrals than anal. Color in life very pale olive brown, with bronze and blue reflections, white below; fins all pale, the tip of anal and edges of caudal somewhat dusky; female with fins rather darker, the upper edge of the pectorals and ventrals largely black; in the males these fins are pale, or somewhat brown above; maxillary barbels blackish; lower pale. Generally abundant along the Pacific coast of tropical America. Specimens were observed by Dr. Gilbert at Mazatlan, Libertad, Punta Arenas, and Panama. It reaches a length of about 18 inches, and is seldom eaten. It resembles *Galeichthys gilberti*, but is readily distinguished by the small, pale ventrals, as also by the generic character of the dentition. The males of this species, according to Dr. Steindachner, carry the eggs in their mouths until after hatching. (Jordan & Gilbert, Bull. U. S. Fish Comm., II, 1882, 44.)

184. NETUMA OSCULA (Jordan & Gilbert).

Head 3⅞ (4⅜ in total); depth 6¼ (7⅜); width of head 4⅜. D. I, 7; A. IV, 14. Body moderately elongate, the head short, rather narrow, tapering forward, considerably broader than deep. Eye small, 7½ in head, placed well above the mouth. Interorbital space 1$\frac{9}{10}$ in head; snout 3; breadth of mouth 2⅔. Mouth very small for the genus, with thick lips. Teeth on vomer and palatines villiform, but rather coarse and bluntish. Vomerine patches small, rather longer than broad, separated on the median line, and each also separated by a narrow groove from the large and roundish palatine bands, which have a distinct backward prolongation. Premaxillary band of teeth very broad, barely 3 times as long as wide. Barbels short, the maxillary barbels reaching slightly beyond base of pectorals, the outer mental barbels scarcely past gill opening below; inner mental barbels about as long as snout. Dorsal shield short, crescent-shaped, granulated, but without median keel, its length about ¼ its breadth. Occipital process narrow, its edges almost parallel until abruptly widened at base; the narrow part considerably longer than broad, with curved edges; a well-developed median keel. Fontanel broad and shallow, abruptly contracted at a point midway between tip of snout and end of occipital process, thence continued backward as a narrow groove to a point less than an eye's diameter in front of the base of the occipital process. Greatest width of fontanel about ⅗ eye. Shields of top of head all coarsely and rather sparsely granular, and anteriorly striate. Interorbital space nearly plane, with a few low, smooth ridges. Opercles scarcely rugose. Gill membranes forming a narrow fold across isthmus posteriorly. Dorsal spine very high, 1⅕ in head, a little longer than pectoral spine; humeral process granular, not quite ⅔ length of pectoral spine; no axillary pore; adipose fin adnate posteriorly; caudal long,

its upper lobe the longer, somewhat falcate, $1\frac{1}{10}$ in head; anal rather high. Color brown, with bluish reflections; lower parts dusky, with dark punctulations; fins all blackish; maxillary and outer mental barbels dusky. A single male example 11 inches long was obtained at Panama by Dr. Gilbert. (Jordan & Gilbert, Bull. U. S. Fish Comm., II, 1882, 46.)

185. NETÚMA ELATTURA (Jordan & Gilbert).

Head $3\frac{3}{8}$ ($4\frac{1}{8}$ in total); depth $5\frac{3}{4}$ ($6\frac{3}{4}$); width of head $4\frac{2}{5}$. D. I, 6; A. IV, 14. Length (29408, U. S. Nat Mus.) $12\frac{1}{2}$ inches. Body low, not very elongate, the head rather short and very broad, much broader than deep, the snout depressed and very broadly rounded, almost truncate. Eye moderate, placed rather high, its diameter 7 in head. Interorbital space $2\frac{1}{6}$ in head; snout $3\frac{1}{3}$; breadth of mouth $1\frac{7}{8}$. Mouth large, with thickish lips, the upper jaw considerably projecting. Teeth on vomer and palatines villiform, but bluntly conical, less acute than in most of the species. Vomerine patches oblong, small, separated by a narrow interspace from each other and from the palatine bands, which are roundish and comparatively small, with a backward prolongation. Teeth in jaws in broad bands. Barbels rather short, the maxillary barbels reaching a little past base of pectorals, the outer mental barbels a little past gill opening, the inner a little more than $\frac{1}{3}$ head. Dorsal shield not very short, crescent-shaped, with a distinct median keel, its length on the median line about $\frac{1}{3}$ its breadth. Occipital process short, broadly triangular, with concave sides which spread out abruptly near the base, forming a sort of shoulder, its length scarcely equal to its width at base. Median keel well developed. Fontanel broad and shallow, abruptly narrowed posteriorly at a point a little nearer base of dorsal than tip of snout, but extending as a groove to a point distant less than a diameter of the eye from the base of the occipital process, this groove indistinct in the smaller specimen. Greatest width of fontanel scarcely more than $\frac{1}{2}$ the eye. Shields of head granular-striate, the roughness less coarse than in *A. kessleri*. Interorbital space with 2 prominent ridges and numerous striæ, none of them granular, the granulations chiefly confined to the region behind widest part of fontanel. Opercle striate. Gill membranes forming a moderate fold across isthmus. Dorsal spine low, shorter than pectoral spine, which is $1\frac{3}{5}$ in head, the anterior edges of both bluntly serrate; humeral process broadly triangular, granulated, not $\frac{2}{3}$ length of pectoral spine, much smaller than in *A. insculptus*; no axillary pore; adipose fin long and low, without free posterior margin; lower fins of moderate length; vent much nearer ventrals than anal. Caudal short, the upper lobe longest, $1\frac{3}{5}$ in head (a little more than $\frac{1}{3}$ head in the smaller specimens). Color dusky above, the lower parts soiled with dark points; fins all more or less dusky with dark points; maxillary barbels dusky, others pale. One male individual (29408) was obtained at Panama by Dr. Gilbert; another (30995) at Panama by Mr. Rowell. (Jordan & Gilbert, Bull. U. S. Fish Comm., II, 1882, 45.)

185 (a). NETUMA INSULARUM, Flora Hartley Greene.

Head 3⅓ in length; width of head 4⅔ in length; interorbital space in length 7, interorbital space in head scarcely 2; snout in head 3; breadth of mouth in head 2; eye in head 6⅔. D. I, 7; A. I7. Head much broader than deep; snout depressed and broadly rounded, eye above the level of the mouth. Upper jaw projecting. Teeth on vomer and palatines villiform and bluntly conical, the 2 vomerine patches forming together a band almost as long and slightly broader than the premaxillary band, their sides separated by a narrow interspace; palatine teeth wholly separated from the vomerine teeth and in 2 large triangular patches which extend backward over the pterygoid region; each triangle has a sharp notch in its antero-lateral side; its antero-posterior length is twice its lateral width, teeth of lower jaw in a narrower band than the upper jaw. Maxillary barbel extending to end of first third of the length of the pectoral spine; outer mental barbel to base of pectoral; inner mental barbel past gill opening, 2⅓ in head. Dorsal shield crescent-shaped, without median keel, length on median line 2⅔ in distance between the horns of the crescent; 2 notches on its anterior side to meet the corresponding points from the occipital process. Occipital process broadly triangular, with the outer sides concave and 2 small projections at its posterior end. Median keel evident, rather short. Occipital process much broader at base than long; its length 3⅓ in head; posterior breadth 2 in length of process. Fontanel broad and shallow, narrowed gradually posteriorly to a point halfway between snout and base of dorsal spine. A narrow line runs back from it the distance of a long diameter of the eye. Greatest width of the fontanel equals the short diameter of the eye. Shields of the head granular-striate, the striæ evident and extending to the middle of the interorbital space, and on the side to meet the humeral process at the top of the gill opening. Opercles nearly smooth. Gill membranes forming a fold across the isthmus. Dorsal and pectoral spines crenulate in front and sharply decurved serrate behind. Dorsal shorter than pectoral, which is 1⅓ in head. No axillary pore evident. Adipose fin long and low with posterior margin attached. Vent much nearer ventrals than anal. Color in alcohol, dark blue above, light blue on side, and white below; maxillary barbel dusky; fins all dusky. The type of this species (No. 47577, U. S. Nat. Mus.) was collected by the *Albatross* in the Galapagos Archipelago, being part of the collection studied by Jordan & Bollman in 1889. It was recorded by them (Proc. U. S. Nat. Mus. 1889, 179) as "*Tachysurus elatturus* (var?)." Its relations to *Netuma elattura* are close, but its fins are better developed and there are several differences in details of structure.

Netuma insularum, Flora Hartley Greene in Gilbert, Proc. U. S. Nat. Mus. 1896 (Feb. 5, 1897), 439, Galapagos Archipelago.

69. GALEICHTHYS, Cuvier & Valenciennes.

Dorsal shield small, lunate; teeth on palate villiform, the patches on each side not extending backward over the pterygoid region.

Subgenus GALEICHTHYS.

Shields of head mostly covered by soft skin, hiding the granulations.

163. GALEICHTHYS LENTIGINOSUS (Eigenmann & Eigenmann).

Head 4 to 4⅕; depth 5 to 6. D. I, 6; A. 22. Eye 2½ in snout, 8½ in head, 4½ in interocular, 2¼ in interorbital. Body nearly terete anteriorly, becoming compressed backward; the width, above the pectorals, a little greater than the depth. Head flat, depressed, its depth at base of occipital process 1⅕ in its greatest width, which is about 1¼ in its length. Occipital process somewhat roughened, about twice as long as its greatest width, its margin straight and oblique; the middle of the fontanel above the posterior part of the eye. Head everywhere covered with skin; sides of the head and opercle with vermiculating canals. Snout somewhat pointed; upper jaw very little projecting; lips thick; teeth all villiform; the intermaxillary band strongly curved; vomerine teeth in 2 oval patches joined to the larger patches of the palatines; mandibulary band of teeth separated in front, the outer margins, if continued forward, forming an angle at the symphysis. Maxillary barbels reaching beyond base of pectorals; mental barbels reaching about ⅔ toward the gill opening; the postmentals to the gill opening in 1 specimen, a little before in the other. Gill membranes forming a broad, free margin across the isthmus. Gill rakers 3 + 4. Pectoral pore minute; humeral process pointed behind. Distance of dorsal from snout 2⅓ in the length, the dorsal spine covered with a membrane, its outer margin granular, its height 1⅔ in the head, the first soft ray 1⅔ the length of the fin; distance of adipose fin from the dorsal 3⅔ to 4 in the length, the fin adnate, longer than the dorsal; caudal lunate, the upper lobe longer, somewhat falcate, 4½ to 5 in the length; anal fin twice as long as high, the highest ray 2 to 2¼ in the head; ventrals short and broad, 1⅓ in the head; pectoral spine covered with a membrane, 1½ to 1⅔ in head. Light brown, becoming nearly white below, the sides freckled; fins reddish. Panama. (Eigenmann & Eigenmann, Nematognathi, 50.)

164. GALEICHTHYS PERUVIANUS, Lütken.

Head 3½; depth 4⅛ to 5½. D I, 7; A. 14 to 16. Eye 2 in snout, 7 in head, 4 in the interocular, 2 in the interorbital. Subterete, tapering to a long, slender caudal peduncle; the greatest width about equal to the greatest depth. Head not much depressed; interorbital area flattish, the greatest depth of the head 1¾ in its length, its greatest width 1⅞ to 1⅔; the width at angle of mouth 2 in its length; the surface of the cranial bones longitudinally furrowed, covered with muscle and skin. Occipital process more than 3 times as long as wide; anterior fontanel elongate, its center over the middle of the eye, continued as a deep groove to the base of occipital process; a small opening a pupil's distance behind the anterior fontanel, and a larger one in the occipital bone at the end of the groove. Snout, upper part of the neck, and the opercle sometimes with conspicuous reticulating mucous canals. Snout blunt, decurved. Maxillary barbels extending beyond base of pectoral, mentals about to gill

openings, the postmentals about 1 diameter of the eye farther. Jaws subequal, the upper longer; teeth all fine, villiform; intermaxillary band of teeth very wide, its depth about 8 in its width; 2 small patches on the vomer; palatine patches very wide and shallow, tapering to a point. Gill membranes meeting at an acute angle, forming a fold across the isthmus; gill rakers 3 + 10. Humeral process very thin, covered with skin, more than ½ as long as the pectoral spine, broadly expanded and rounded behind; pectoral pore present. Distance of dorsal spine from snout 2⅔ to 2¾ in the length; the dorsal spine broken off in all the specimens; distance of adipose from the dorsal 3⅜ to 3⅝ in the length; adipose fin as long as the dorsal fin, adnate; caudal fin broadly lunate, the upper lobe longer, falcate, 3¾ in the length; anal fin higher than long, the highest ray 2 to 2⅔ in head; ventrals reaching to the anal, 1½ to 2 in head; pectoral spines broken in all the specimens. Back, top of head, and a band from humeral process to the lower caudal lobe, blue black; a broad conspicuous, bluish-silvery band along the lateral line; lower parts white; fins blackish; ventrals and anal sometimes with light areas. Eleven specimens, 0.25 to 0.35 m. long. Callao, Peru; Haslar Expedition. (Eigenmann & Eigenmann, Nematognathi, 5.)

Subgenus HEXANEMATICHTHYS, Bleeker.

Shields not entirely covered by soft skin, the granulations evident, especially in the male.

187. GALEICHTHYS SEEMANNI (Günther).

D. I, 7; A. 19; P. I, 10. The height of the body is contained 4½ times in the total length (without caudal); the length of the head 2⅓; head much broader than high, its greatest width being equal to its length without snout. Eyes of moderate size, much nearer to the end of the snout than to the operculum; the length of the snout is ⅔ of the width of the interorbital space. The median longitudinal fonticulus on the upper side of the head extends to the base of the occipital process. Teeth on the vomer separated in the middle by a short interspace, forming a pair of small subquadrangular patches which are confluent with those of the palatines. The latter are much longer than broad, elliptical. The band of intermaxillary teeth is 5½ times as broad as long. The maxillary barbels extend nearly to the end of the head, and are about twice as long as the outer ones of the mandible. Crown of the head, and nape finely granular; occipital process broader than long, with a prominent ridge along its middle. The basal bone of the dorsal spine is small, with a few fine granules. Dorsal spine of moderate strength, more than ½ as long as the head, serrated along both edges; the first soft ray is as high as the body. Adipose fin rather shorter than dorsal. The upper caudal lobe is the longer, ⅔ of total length. Porus axillaris present. Ventral fin shorter than pectoral. Sides of the body silvery; basal half of the inner side of the paired fins black. Central America. A fine specimen 12 inches long, from the Haslar collection, collected by Dr. Seemann. (Günther.)

Arius seemanni, GÜNTHER, Cat., v, 147, 1864, Central America.

Jordan (Proc. Ac. Nat. Sci. Phila. 1883, 282) adds the following note on the type of this species: "Fontanel extending backward in a deep and narrow groove which reaches the occipital process. Middle of top of head smooth, much as in *A. platypogon*."

The following account is given by Eigenmann & Eigenmann (Nematognathi, 78):

Head $3\frac{3}{4}$; depth 5. D. I, 7; A. 18. Body about as deep as wide, tapering to a slender peduncle. Head flat, depressed in front, top of the head coarsely granular; opercles smooth or with faint striations; humeral process slightly granular, covered with skin; the greatest depth of the head $1\frac{2}{5}$ in its length, greatest width $1\frac{1}{3}$ to $1\frac{2}{5}$; the width at angles of the mouth $2\frac{1}{5}$. Occipital process wider than long; fontanel open to above the posterior margin of the eye, with a deep backward-extending groove. Interorbital area smooth, without ridges. Eye 2 in snout, 7 in head, $3\frac{3}{5}$ to 4 in the interocular, $2\frac{1}{3}$ in the interorbital. Maxillary barbels reaching slightly beyond base of pectorals, mental barbels $\frac{2}{3}$ toward the gill opening, the postmentals $\frac{1}{2}$ a diameter of the eye behind the gill opening or farther. Upper jaw longer; teeth all villiform; vomerine teeth in 2 small ovate patches, which are separated from each other but joined to the much larger palatine patches. Gill membranes forming a moderate fold across the isthmus. Gill rakers 5 + 12. Pectoral pore large; vertical series of pores present. Distance of dorsal fin from tip of snout $2\frac{1}{3}$ to $2\frac{2}{5}$ in the length, the spine rather stout, $1\frac{2}{3}$ in head, its outer edge granular toothed, its inner edge with short, recurved teeth; distance of the adipose fin from the dorsal $3\frac{1}{2}$ to $3\frac{1}{5}$ in the length; adipose fin slightly longer than high, shorter than the dorsal fin. Caudal 4 in the length; anal emarginate, little longer than high; ventrals $1\frac{3}{4}$ to 2 in the length of the head; pectoral spine $1\frac{1}{3}$ in head, its anterior margin granular toothed, its inner edge with long, straight teeth. Plumbeous, silvery below; fins dusky, inner surface of ventrals and pectorals dark. One female 0.28 m., Panama. One male 0.21 m., Panama.

187(a). GALEICHTHYS GILBERTI, Jordan & Williams.

Head $3\frac{3}{7}$ to 4; width of head $5\frac{1}{3}$; depth 5. D. I, 7; A. IV, 14. Body comparatively elongate, the head depressed but not very broad, somewhat broader than high; eye rather large, 5 to 6 in length of head; width of interorbital space $2\frac{1}{4}$ in head; breadth of mouth $1\frac{3}{4}$; length of snout $3\frac{1}{3}$. Teeth all villiform; bands of vomerine teeth separated by a rather wide interval, each small, roundish, confluent with the neighboring palatine band, the junction marked by a slight constriction; palatine band ovate, broad behind, varying considerably in size and somewhat in form, the width ranging from $\frac{1}{3}$ diameter of eye to $\frac{2}{3}$, being generally largest in adults; band of palatine teeth without backward prolongation; band of maxillary teeth rather broad and short, its length about 5 times its breadth. Maxillary barbel flattened at base, reaching a little past base of pectoral in young, scarcely to gill opening in adult; outer mental barbels 2 in head, inner 3. Gill rakers 4 + 12. Dorsal shield very short, narrowly crescent-shaped, its length on median line not more than $\frac{1}{2}$ that of one

of its sides. Occipital process subtriangular, not quite as long as broad at base, with a strong median keel, its sides slightly curved. A short distance in front of the beginning of the keel is the end of the very narrow groove-like fontanel, which is somewhat widened anteriorly, finally merging into the broad, flat, smooth interorbital area, the boundaries of which are not well defined; shields of head unusually smooth, all finely and very sparsely granular, the granules not forming distinct lines. Some specimens (probably females) about as smooth as in the subgenus *Galeichthys*. Gill membranes forming a rather broad fold across isthmus. Dorsal spine long, usually, but not always, shorter than the pectoral spine, about 1⅘ in head; axillary pore absent; humeral process rather broadly triangular, not much produced backward, less than ½ length of pectoral spine, its surface not granular, covered with skin; adipose fin ½ length of anal, its posterior margin little free; upper lobe of caudal the longer and somewhat falcate, about as long as head; ventrals long, about reaching anal in females, rather shorter in males; vent much nearer base of ventrals than anal. Color olive green, with bluish luster, white below; upper fins dusky olivaceous; caudal yellowish dusky at tip; anal yellowish with a median dusky shade; ventrals yellowish, the basal half of upper side abruptly black; pectorals similarly colored, the black area rather smaller; maxillary barbel blackish; other barbels pale. Length 12 to 18 inches. Coast of Sinaloa; very common; by far the most abundant species at Mazatlan; not recorded from localities farther south.

Arius assimilis, JORDAN & GILBERT, Bull. U. S. Fish Comm., II, 1882, 47, not of GÜNTHER.
Galeichthys gilberti, JORDAN & WILLIAMS, Rept. Fishes Sinaloa, in Proc. Cal. Ac. Sci. 1895, 395, pl. 26, Mazatlan. (Type, No. 29213. Coll. Chas. H. Gilbert.)

188. GALEICHTHYS JORDANI (Eigenmann & Eigenmann).

Head 3⅜ to 3⅝; depth 5½ to 5⅚. D. I, 7; A. 18; eye large, 1⅘ in snout, 5½ in head, 2 in the interorbital, 2¾ to 3 in the interocular. The specimens agree very closely with the description of *assimilis* by Jordan & Gilbert (*gilberti* of the present paper). They differ in the width of the mouth and in having a pectoral pore. Rather robust, the width little less than the depth, caudal peduncle compressed. Head heavy, little broader than high, its height 1½ in its length, its width 1⅔ to 1⅘, width at the angle of the mouth 2 to 2⅛; interorbital area flat and smooth; posterior portion of the head finely and sparsely granular; opercle and humeral process smooth; occipital process about as long as broad, unusually sharply keeled, fontanel extending to above the posterior part of the orbit, continuing as a deep groove to the base of the occipital process; maxillary barbels extending to the pectoral pore, postmentals at least to the gill opening, mental about ⅗ as long as the postmental barbels; snout blunt, decurved; upper jaw a little produced; teeth all villiform, those on the vomer forming 2 small, separate, ovate patches, which are contiguous to the twice or thrice as large palatine patches; gill membranes forming a fold across the isthmus; gillrakers 6 + 9; pectoral pore large; vertical series of pores present; distance of dorsal spine from tip of snout 2⅖ to 2⅔ in the length; the spine of the dorsal and pectoral fins granular on the

basal half of their outer margin; almost the entire inner margins serrate, the spines of equal length, 1½ in head; distance of adipose fin from the dorsal 3⅓ in the length; the adipose more than ½ as long as the dorsal fin, its posterior margin free; caudal deeply forked, the upper lobe longer, somewhat falcate, 3¾ to 4 in length; anal fin about as long as high, deeply emarginate, its highest ray 2⅔ in head; ventral fins not reaching to the anal, 2 in head. Dorsal surface dark blue, with metallic luster, becoming silvery below; lower caudal lobe dusky; basal half of the inner surface of the paired fins black. (Eigenmann & Eigenmann, Nematognathi, 79.) Panama; known only from specimens in the Museum of Comparative Zoology. The specimens from Mazatlan referred to *jordani*, *seemanni*, and *assimilis* by authors belong to *Galeichthys gilberti*.

188(a). GALEICHTHYS AZUREUS, Jordan & Williams.

Head 3¾; width of head 4⅝; depth 9. D. I, 7; P. I. 10: A. IV, 14. Gill rakers 6 + 13. Body robust, its width anteriorly greater than its depth; caudal peduncle short, stout; distance from base of anal fin to base of median caudal rays about ½ length of head; head flat, very broad, its depth at posterior angle of jaw about ⅓ its width; interorbital region flat, smooth anteriorly and granulated posteriorly; fontanel almost obsolete, wide anteriorly and ending in a short groove posteriorly at a point ⅓ distance from tip of snout to posterior edge of occipital process; top of head, occipital process, and antedorsal shield finely granular, granulations mostly arranged in radiating striæ and extending forward to a line with the pupils; nostrils very large and close together, posterior one with a broad [illegible] diagonal, its length 4½ in head, about as long as side, with a very low ridge; dorsal shield crescent-shaped, with [illegible] reaching back on each side of fin, its median length about ½ the length of the side; eye small, about 9 in head; interorbital width almost 2 in head; snout almost 4 in head; breadth of mouth 2⅔ in head; maxillary barbel slender, thick at base, 1⅓ in head; outer mental barbel reaching to posterior angle of jaw, about 2⅔ in head; inner mental barbel about 4 in head; teeth all villiform; premaxillary band narrow, about ⅓ as wide as long; vomerine and palatine bands of teeth fully confluent on each side, forming together a crescent-shaped patch, narrowly divided on the median line of the vomer; form of vomerine bands similar to that of the palatine bands but smaller; palatine band of teeth with [illegible] prolongations; opercle with radiating ridges, humeral process granular, triangular, lower posterior corner prominent; axillary pore very small; gill membranes forming a broad fold across isthmus; dorsal fin short, base not including spine equal to base of adipose dorsal; dorsal spine robust, but little shorter than pectoral spine, about 2 in head, its anterior serræ small and tubercle-like, its posterior edge, as well as that of pectoral, retrorsely serrate; soft rays of dorsal extending but little beyond spine, the longest about ⅔ length of head, adipose dorsal about [illegible] as high as long, caudal lobes unequal, the upper lobe about [illegible] of medium height; distance from vent to base of [illegible]

origin of anal; pectoral spine very strong, its anterior margin with serræ toward the tip, becoming small tubercles toward base; soft rays but little longer than spine, which reaches slightly beyond ½ distance from origin to base of ventrals. Color dark blue, with silvery reflections on sides; belly pale; mental barbels dusky; maxillary barbels light below and black above; paired fins darkest on inner side; other fins almost uniformly dusky. One specimen, 19¼ inches long, taken by the Hopkins Expedition at Mazatlan; probably not distinct from *G. guatemalensis*.

Galeichthys azureus, JORDAN & WILLIAMS, Rept. Fishes Sinaloa, in Proc. Cal. Ac. Sci. 1895, 398, pl. 27, Mazatlan. (Type, No. 1575, L. S. Jr. Univ. Mus. Coll. Hopkins Exped. to Sinaloa.)

189. GALEICHTHYS CÆRULESCENS (Günther).

D. I, 7; A. 17; P. I, 10. The height of the body is contained about 5 times in the total length (without caudal), the length of the head 3½ or 3⅔ times; head much broader than high, its greatest width being ⅔ of its length. Eyes rather small, their diameter being ⅓ of the extent of the snout, ⅔ of their distance from the gill opening, and ⅓ of the width of the interorbital space. The teeth on the palate form a slightly curved band, composed of 2 vomerine patches which are much broader than long, and of a pair of palatine patches which are subcontinuous with, scarcely broader and longer than, those of the vomer. The barbels of the maxillaries extend to the middle, the outer ones of the mandible to the base of the pectoral. Crown of the head granular; occipital process broader than long, subtriangular, subtruncated behind, and slightly raised along the median line; the basal bone of the dorsal spine is subtriangular, small. Dorsal spine of moderate strength, more than ½ as long as the head, granulated in front and slightly serrated behind; the first soft ray is as high as the body; adipose fin shorter than the dorsal; caudal deeply forked, with the upper lobe the longer, its length being nearly equal to that of the head; pectoral spine serrated along its inner edge and on the extremity of its outer edge; it is as long as the head without snout. Ventral fin shorter than pectoral. Sides steel-blue iridescent, blackish toward the back, and silvery below; vertical fins black; inner side of the paired fins blackish. Guatemala. a–b. Five specimens, 12 inches long. Huamuchal. From the collection of Messrs. Godman and Salvin. (Günther, Cat., v, 149.)

The following note on the types of this species is given by Jordan (Proc. Ac. Nat. Sci. Phila. 1883, 282): "Head more depressed than in *A. assimilis*. Fontanel very short, ending abruptly behind and not produced in a groove behind the smooth area of the top of the head, the boundary of the smooth area being rather abruptly convex. Occipital process broader than long, its edges nearly straight. Bands of palatine teeth small, not produced backward on the inner margin. Paired fins black at base above." No recent collector has found this species.

189(a). GALEICHTHYS XENAUCHEN, Gilbert, new species.

Head $3\frac{7}{8}$ in length; depth at front of dorsal $5\frac{1}{4}$; anal 23. Width of head at opercle $1\frac{2}{7}$ in its length; width at front of eyes 2 in head; width of mouth at inner angles $2\frac{2}{5}$ in head; interorbital width $2\frac{1}{10}$; eye very small, 9 in head, $3\frac{1}{3}$ in its distance from tip of snout, $4\frac{5}{9}$ in postocular part of head, $4\frac{1}{3}$ in interorbital width. Teeth all villiform; mandibular bands well separated on middle line, very broad mesially, rapidly tapering to a point laterally, the band produced beyond angle of mouth, its greatest width $2\frac{1}{3}$ times in its length; premaxillary band very convexly curved, following the outline of the snout, its width $5\frac{3}{4}$ in its length; vomerine patches roundish, separated by an evident medial groove, marked off from the palatine patches by a narrower groove and a constriction; the palatine patches are equal in width to the vomerine patches, and less than twice as long, of nearly equal width throughout. Maxillary barbels very slender, reaching slightly beyond the base of the pectoral spine; the mental barbels do not reach edge of gill membrane, the outer pair equaling length of snout and $\frac{1}{2}$ of eye. Nostrils very large, the anterior broadly oval, with widely reflexed rim; the posterior widely elliptical, not concealed by the valve; distance from anterior nostril to tip of snout equaling that from posterior nostril to front of eye. Fontanel wide, with nearly parallel edges on frontal region, abruptly narrowing at front of occiput, where it is continuous with a narrow and shallow groove; the latter fails to reach base of occipital process by a distance equaling $\frac{1}{2}$ diameter of eye. The raised margins of the fontanel continuous with a pair of sharp ridges bounding the groove, these accompanied by a pair of lower ridges on their outer sides and parallel with them; posteriorly these ridges roughened with granules and merging into the granulated area on posterior part of occiput; occipital process granulated, the granules arranged in more or less definite lines radiating backward and downward on each side from median point of base; lateral portions of occiput with an area of radiating striæ separated from the central ridges by a smooth groove-like depression; a narrow granulated area extending forward on each side of fontanel to above back of orbits; occipital process very long and narrow, its width opposite its middle being but $\frac{2}{5}$ of its length, abruptly expanding near base, the basal width being $\frac{1}{2}$ its length plus that of dorsal plate on median line; opercles and humeral plate weakly striate. Gill membranes with a wide, free fold posteriorly; gill rakers weak and short, 1+4 movable ones; no evident axial pore. Dorsal spine slender, with a series of sharp granulations on anterior edge, minutely roughened, not serrate, behind, broken in the type, but its length was about $\frac{2}{3}$ that of head; pectoral spines rather slender, rough granular on outer margins, with short, fine serræ within, both mutilated in the type, but their length was about equal to that of dorsal spine; pectoral extending nearly $\frac{2}{3}$ distance to ventrals, the ventrals nearly to origin of anal; distance from anus to base of ventrals $\frac{2}{5}$ its distance from front of anal; anal fin very long, its base $1\frac{1}{4}$ in head, its longest ray $\frac{1}{3}$ head; distance between dorsals $3\frac{1}{4}$ in length; adipose fin long, highest about opposite the middle, a short, almost verti-

cal, free posterior margin, its vertical height $3\frac{2}{3}$ in its length; the latter over twice its distance from rudimentary caudal rays, greater than base of first dorsal, equal to $\frac{1}{2}$ length of head; caudal with broad lobes, the lower rounded; the upper mutilated in the type, but evidently acute and longer than the lower. Color purplish above, more bluish anteriorly, the lower parts silvery, coarsely punctate with brown; fins all blackish except the lower surface of the paired fins. In appearance most closely allied to species of *Netuma*, having the low, depressed head with the lateral outlines converging forward to the narrow pointed snout, and a long, largely adherent adipose dorsal. The palatine patches are, however, narrow and without backwardly projecting lobes. The species is distinguished from all those known from the Pacific coast of America by the long and extraordinarily narrow occipital process. Type, a female 386 mm. long, from Panama. (ξένος, strange; αὐχήν, nape.)

Galeichthys xenauchen, GILBERT, Fishes of Panama, MS. 1898, Panama.

190. GALEICHTHYS GUATEMALENSIS (Günther).

Head $3\frac{5}{7}$ ($4\frac{3}{4}$ in total); width of head 5 ($6\frac{1}{5}$); depth $6\frac{1}{8}$ (7). D. I, 6; P. I, 10; A. III, 15. Length (28140, U. S. Nat. Mus.) $12\frac{1}{2}$ inches. Body slender, its width anteriorly greater than depth; caudal peduncle compressed, short; distance from end of anal to base of median caudal rays about $\frac{1}{2}$ length of head. Head depressed, not very broad, its depth at posterior margin of branchiostegal membranes less than $\frac{2}{3}$ its width; interorbital region flat, smooth, the smooth area forming a broad equilateral triangle, its base at the interorbital space, the apex at a point $\frac{4}{5}$ the distance from snout to dorsal, the triangle forming the termination of the almost obsolete fontanel; top of head, occipital process, and antedorsal shield finely granular, some of the anterior granulations only arranged in lines, none of them in radiating striæ. Occipital process broadly trapezoidal, its width slightly greater than the length of its side, with a slight or obsolete median carina; its posterior margin truncated; its sides slightly convex posteriorly, concave toward the front. Dorsal shield small, narrow, crescent-shaped, its median length about $\frac{1}{4}$ the length of its side. Eye small, 6 in head; interorbital width $2\frac{2}{3}$ in head; snout 4 in head; breadth of mouth 2. Maxillary barbel very slender, reaching base of pectoral spine; outer mental barbel to well beyond margin of branchiostegal membranes, its length about $\frac{1}{2}$ head; inner mental barbel 3 in head. Teeth all villiform; width of premaxillary band about $\frac{1}{6}$ its length; vomerine and palatine bands of teeth fully confluent on each side, forming together a crescent-shaped patch, narrowly divided on the median line of the vomer; form of vomerine band similar to that of the palatine band; palatine band of teeth without backward prolongation; opercle with radiating ridges; humeral process granular, narrow, produced backward, not quite $\frac{1}{2}$ length of pectoral spines; no axillary pore. Gill membranes forming a narrow fold across isthmus. Dorsal short, its base about equal to that of the adipose dorsal; dorsal spine robust, but little shorter than the pectoral spine, about $\frac{3}{4}$ length of head, its anterior serræ small and tubercle-like; its posterior edge, as well as that of the pectoral, retrorsely serrate; soft rays of dorsal extending much beyond the spine,

the longest about ¾ length of head; adipose dorsal about ½ as high as long, its posterior margin largely free; caudal very widely forked, the upper lobe falcate, nearly ⅓ longer than the lower, as long as head; anal short and low; distance from vent to base of ventrals slightly more than ½ its distance from origin of anal; pectoral spine very strong, much stronger than dorsal spine, its anterior margin with serræ toward the tip, becoming small tubercles toward base; inner edge with strong retrorse serræ, the soft rays longer than spines, reaching ¾ distance to base of ventrals. Color very dark bluish or greenish above; sides with bronze luster; belly silvery; mental barbels white, with black edge; maxillary barbels blackish; fins all blackish, the caudal nearly uniform; paired fins darkest on inner side; sides with vertical series of mucous pores, conspicuous in life. This species is not uncommon at Mazatlan, where several specimens were obtained by Dr. Gilbert. Four specimens from Colima are also in the National Museum. It has not been observed at Panama. The original description of this species is brief and not entirely correct. That it was intended to refer to the species here described we have ascertained by the examination of Dr. Günther's original types in the British Museum. (Jordan & Gilbert, Bull. U. S. Fish Comm., II, 1882, 48.)

191. GALEICHTHYS ASSIMILIS (Günther).

D. I, 7; A. 19; P. I, 10. The height of the body is contained 4⅔ times in the total length (without caudal), the length of head 3⅔; head much broader than high, its greatest width being ¾ of its length. Eyes rather small, situated nearer to the end of snout than to that of operculum; the length of snout is ⅔ of the width of interorbital space. The median longitudinal fonticulus on the upper side of the head does not extend to the base of occipital process. Teeth on vomer but slightly separated in the middle, forming a pair of oblong transverse patches which are confluent with those on the palatine bones; the latter are short, club-shaped. The band of intermaxillary teeth is 5 times as broad as long. All the teeth villiform. The maxillary barbels extend nearly to the end of head; the length of the outer ones of the mandible is ½ or ⅔ that of the head. Crown of the head granular, the granulations being arranged in radiating streaks. Occipital process broader than long, triangular, with its hinder end concave. The basal bone of the dorsal spine of moderate size, crescent-shaped. Dorsal spine of moderate strength, more than ½ as long as head, granulated in front and slightly serrated behind; the first soft ray longer than spine and as high as body; adipose fin shorter than dorsal; caudal deeply forked, with the upper lobe the longer, its length being contained 5½ times in the total; pectoral spine serrated along its inner edge and on the extremity of the outer edge; ventral fin shorter than pectoral. Sides of the body silvery; vertical fins grayish; basal half of the inner side of the paired fins black. Guatemala. A fine specimen, 13 inches long. Lake of Yzabal. From the collection of Messrs. Godman and Salvin. (Günther, Cat., v, 146.)

Jordan (Proc. Ac. Nat. Sci. Phila. 1883, 281) has the following note on the type of this species: "Area between the eyes smooth, extending backward in the form of a rather narrow triangle, which is moderately

obtuse behind. Fontanel narrow and short, ending far in front of occipital process, not extending backward as a groove behind the smooth area of the top of the head; posterior end of fontanel midway between tip of snout and middle of dorsal shield. Occipital process broad, its edges not straight. Band of palatine teeth large, but not produced backward on the inner margin. * * * There is no evidence of the occurrence of the true *A. assimilis* in Pacific waters."

192. GALEICHTHYS SURINAMENSIS (Bleeker), text, p. 129.

193. GALEICHTHYS DASYCEPHALUS (Günther).

Head $4\frac{1}{2}$ ($5\frac{2}{3}$ in total); depth 6 ($7\frac{1}{4}$ in total); width of head $5\frac{1}{3}$. D. I, 7; A. IV, 17. Length (29400) 11 inches. Body elongate, compressed behind, the head small, narrow, and moderately depressed anteriorly, the snout not very blunt. Eye rather large, placed somewhat above level of angle of mouth, its length 5 in head; width of interorbital space $2\frac{1}{5}$ in head; breadth of mouth $2\frac{1}{2}$; length of snout $3\frac{1}{4}$. Teeth villiform, those of vomer and palatines rather coarse, bluntly conic; bands of vomerine teeth separated by a rather broad area, each confluent with the neighboring palatine band, the two forming a small oblong patch much smaller than the eye, the division between the palatine and vomer scarcely appreciable. Palatine bands without backward prolongation. Bands of teeth in jaws short and broad. Maxillary barbel reaching about to middle of pectoral spine; outer mental barbel to base of pectoral; inner slightly more than $\frac{1}{2}$ head. Dorsal shield short, crescent-shaped, a little more than 3 times as broad as long on the median line. Occipital process subtriangular, its sides straight, slightly longer than broad, its median line rather sharply keeled. Close in front of its base begins the deep fontanel, which is narrow and groove-like posteriorly, becoming rather abruptly broader above the opercle, then gradually narrowed anteriorly. Ridges bounding fontanel prominent anteriorly to a point just behind vertical from nostrils, coarsely granular for their whole length, the granules mostly arranged in 1 series. Between these ridges and the eye on each side is another ridge extending obliquely backward and inward from above front of eye, likewise very coarsely granular, the granules mostly in 2 series. Shields of head all rough granular, the granules forming irregular lines. Gill membranes forming a narrow fold across isthmus. Dorsal spine moderate, about equal to pectoral spine, $1\frac{1}{4}$ in head; axillary pore present, small; humeral process broad, scarcely granular, about $\frac{2}{3}$ pectoral spine; adipose fin rather long and low; caudal long, the upper lobe the longer, somewhat longer than head; anal long and high, its outline emarginate, its longest rays a little more than $\frac{1}{2}$ head; ventrals long, the vent nearer their base than that of anal. Color dusky, the entire ventral surface soiled with dark points; fins all largely blackish; barbels black. Two specimens were obtained at Panama by Dr. Gilbert. This species may be known at once by the 4 granulated ridges, which extend the length of the interorbital space. In the female, later taken, the granulations on the head are largely covered by soft skin.

194. GALEICHTHYS LONGICEPHALUS (Eigenmann & Eigenmann).

Head $3\frac{2}{5}$; depth $6\frac{1}{2}$. D. I, 7; A. 20. Elongate, slender, greatest width little greater than the depth. Head long and depressed, its greatest width $1\frac{1}{2}$ in its length, its greatest depth little more than $\frac{1}{2}$ its length. Top of head with faint granules almost entirely concealed by the skin; interorbital area flat and with 4 ridges which are obscurely granular, the inner two bordering the fontanel, the outer ones curved in front extending obliquely backward from near the posterior nasal opening; occipital process as long as broad, its margins concave; fontanel produced as a deep groove to the base of the occipital process; opercle faintly striate; humeral process entirely covered with thick skin, not granular. Eye lateral, well above the angle of the mouth, its diameter $1\frac{1}{2}$ in snout, 6 in head, 3 in interocular; snout depressed and rounded in front. Maxillary barbels extending scarcely beyond base of pectoral, mentals not to gill opening. Upper jaw little projecting; width of the mouth $2\frac{1}{6}$ in the head; intermaxillary teeth long and slender, the depth of the band $4\frac{1}{2}$ in its width; vomerine and palatine teeth obtusely conical, the vomerine patches separate, contiguous to, but not confluent with, the palatine patches. Gill membranes not forming an angle where they meet, with a rather broad, free margin. Gill rakers short and thick, 4 + 5. Pectoral pore small; vertical series of pores present; distance of dorsal fin from tip of snout $2\frac{3}{5}$ in the length, the spine $1\frac{3}{4}$ in the head, its outer margin granular-toothed near its base, its inner margin with short teeth; distance of adipose fin from the dorsal $3\frac{1}{6}$ in the length; adipose fin much longer than high, as long as the dorsal fin; caudal forked, the upper lobe $\frac{1}{3}$ longer than the lower, very nearly as long as head, $3\frac{1}{3}$ in the length; anal fin emarginate, scarcely longer than high, its height $2\frac{1}{3}$ in the head; ventrals reaching almost to the anal, about 2 in head; pectoral spine a little longer than the dorsal spine $1\frac{3}{5}$ in the head; its outer edge roughened, inner edge with rather sharp teeth. Brown above, the sides silvery, entire ventral surface sprinkled with brown dots; a black median line on the back; fins dusky; barbels blackish. One specimen, a male, .29 m. long (No. 4972, M. C. Z.). Panama. Steindachner. (Eigenmann & Eigenmann, Nematognathi, 82.)

195. GALEICHTHYS RUGISPINIS (Cuvier & Valenciennes)

Head $3\frac{1}{2}$ to 4; depth $5\frac{1}{2}$ to 6. D. I, 7; A. 19 to 21. Slender, compressed on the tail. Head broad and depressed, tapering forward; width of the head $1\frac{1}{2}$ to $1\frac{2}{3}$ in its length, at the angle of the mouth $2\frac{1}{3}$ to $2\frac{1}{2}$; depth of head $1\frac{3}{4}$ to 2; profile rather steep. Top of head, humeral process, front and sides of spines, and dorsal plate granular, the granulation not extending forward to above middle of cheeks. Occipital process triangular, about as long as broad, the median ridge not very prominent. Middle of fontanel behind the eye, the posterior portion separated by a bridge, not continued backward as a groove; interorbital region with 4 ridges. Eye small, 3 in snout, 10 in head, $3\frac{1}{2}$ in the interocular. Barbels villiform. Maxillary barbel reaching to or beyond base of pectoral; postmental to gill opening, mental barbels much shorter. Mouth inferior, lower jaw

included, lips thick; teeth villiform, the anterior ones in the jaws longer; depth of the intermaxillary band 4 in its width; palatine patches 1 diameter of eye apart, the width of the patches less than 1 diameter of eye. Gill membranes meeting in an angle, forming a fold across the isthmus. Gill rakers 6 + 11. Pectoral pore none; vertical series of pores present. Distance of dorsal spine from the snout $2\frac{1}{8}$ to $2\frac{1}{2}$ in the length, the spine broken in the specimens examined. Space between dorsal and adipose fins 4 to $4\frac{3}{8}$ in the length. Adipose fin adnate, as long as the anal fin; ventrals $2\frac{1}{2}$ in the head; pectoral spine serrated behind (broken). Two specimens 0.22 m. and 0.26 m. long. Para. Agassiz and Bourget. (Eigenmann & Eigenmann, Nematognathi, 83.)

196. GALEICHTHYS PHRYGIATUS (Cuvier & Valenciennes), text, p. 130.

74. TACHYSURUS, Lacépède.

Teeth on palate granular; dorsal shield small; palatine bands of teeth without backward projecting angle.

197. TACHYSURUS NUCHALIS (Günther).

D. I, 7; A. 21; P. I, 10. The height of the body $\frac{1}{5}$ of the total length (without caudal), the length of head $\frac{1}{4}$. Head as broad as high, its greatest width being $\frac{2}{3}$ its length; its upper surface granulated; occipital process triangular, as long as broad, with the lateral margins slightly concave; it is elevated into an obtuse ridge running along the middle; the longitudinal groove in the middle of the forehead is rather wide, narrow behind, and does not extend to the base of occipital process. Teeth on palate are coarsely granular, and form 2 subtriangular patches of moderate extent, which, sometimes, are subcontinuous with their anterior angles. The maxillary barbels extend nearly to end of pectoral. Dorsal spine of moderate strength, slightly serrated along both edges, $\frac{5}{6}$ length of head; adipose fin small, shorter than dorsal; pectoral spine as long as, but stronger than, that of dorsal; pectoral fin shorter than head. British Guiana. a-c. Six inches long. Purchased of Mr. Scrivener. d-f. Young. Presented by Sir R. Schomburgk. (Günther, Cat., v, 171.)

198. TACHYSURUS FISSUS (Cuvier & Valenciennes).

D. I, 7; A. 20 or 21. Length of head $\frac{1}{5}$ of the total (without caudal). The distance between the end of snout and that of occipital process $\frac{1}{5}$ of the total length (with caudal); basal bone of dorsal spine small. The teeth on the palate form 2 separate subovate patches. The maxillary barbel extends to, or nearly to, the middle of pectoral fin. Adipose fin small. Cayenne. a-b. Presented by Prof. R. Owen. These specimens having had the cavity of the mouth and of the gills extended in an extraordinary manner, I was induced to examine the cause of it, when, to my great surprise, I found them filled with about 20 eggs, rather larger than an ordinary pea, perfectly uninjured, and with the embryos in a forward state of

development. The specimens are males, from 6 to 7 inches long, and in each the stomach was almost empty. Although the eggs might have been put into the mouth of the fish by their captor, this does not appear probable. On the other hand, it is a well-known fact that the American Siluroids take care of their progeny in various ways; and I have no doubt that in this species and in its allies the males carry the eggs in their mouth, depositing them in places of safety, and removing them when they fear the approach of danger or disturbance. (Günther, Cat., v, 172.)

199. TACHYSURUS SPIXII (Agassiz).

Head 3⅜ to 4; depth 5 to 5½. D. I, 7; A. 21. Body compressed, especially toward the caudal fin, the depth greater than the width. Head narrowed forward, its greatest width 1⅓ in its length, its greatest depth 1½; width at the mouth 2⅓ in the length of the head. Top of the head granular in the young, the granules becoming more or less united in the adult, forming fine reticulating ridges, especially on the occipital process, longer than broad, with a blunt median ridge, the margins concave. Fontanel narrow, without interruptions, continued as a deep tapering groove to near the base of the occipital process; interorbital area with 4 ridges; opercles and humeral process roughened, covered with skin; sides of the head, and snout with reticulating mucous canals. Eye 1½ to 2 in the snout, 5 to 6½ in the head, 2¾ to 3 in the interocular. Maxillary barbels varying in extent, from about the middle of the pectoral to the base of the ventrals; postmental barbels extending to the base of pectoral or to near its tip; mentals to edge of gill membrane or to beyond base of pectoral. Upper jaw projecting; lips more or less papillose; teeth on the intermaxillary and the outer ones of the mandible, villiform; the inner series of the mandible and the palate with granular teeth; the palatine patches of teeth small, subovate, sometimes contiguous in front. Gill membranes united, joined to the isthmus, not forming a free margin across it; gill rakers 6 + 11 to 13. Pectoral pore moderate; distance of dorsal spine from snout 2⅜ to 2½ in the length; the spine 1½ to 1½ in head, serrated on its inner margin, granular or almost smooth on its outer margin. Distance of adipose from the dorsal fin 3½ to 3⅔ in the length, the adipose fin shorter than the dorsal fin, free posteriorly; caudal forked, the upper lobe slightly the longer, 4 to 5 in the length; anal fin scarcely longer than high, its highest ray about 2 in head; ventral fin 1⅔ to 2 in head; pectoral spine strong, about as long as the dorsal spine, serrated on its inner margin, granular or scarcely roughened on the outer margin. Color brownish above, sides and ventral surface silvery, sometimes with brown dots. We have examined over 70 specimens measuring from 0 ([illegible]) to [illegible] from Maranhao, Bahia, Rio Janeiro, Para, Santos in São Paulo, Abrolhos, Brazil. The specimens from Para are much darker in color, the lips more papillose, the barbels longer than those of other specimens. The Santos specimens are ashy above, white below the lateral line, with rather large brown dots on sides, becoming fewer below. (Eigenmann & Eigenmann, Nematognathi, 89.)

200. TACHYSURUS MELANOPUS (Günther).

D. I, 7; A. 21; P. I, 10. The height of the body is contained 5 times in the total length (without caudal), the length of the head $4\frac{1}{3}$ times; head somewhat broader than high, its greatest width being $\frac{3}{4}$ of its length; the occiput and nape are finely granulated; occipital process subtriangular, as long as broad, with the lateral margins somewhat concave, and with the median ridge a little elevated. The longitudinal groove in the middle of the crown of the head is indistinct, narrow, linear behind, scarcely extending to the base of the occipital process. The teeth on the palatines are obtusely conical, and form 2 rather small subovate patches, apart from each other, and situated on the front part of the palate. The maxillary barbels do not quite extend to the middle of the pectoral fin. Dorsal spine of moderate strength, scarcely serrated anteriorly, equal in length to the distance of the gill opening from the anterior margin of the orbit, or even somewhat shorter; adipose fin small, the length of its base being less than that of the dorsal; pectoral spine nearly as long and strong as that of the dorsal fin, very strongly serrated anteriorly. Porous axillaries nearly as wide as a nasal opening. The upper (inner) surface of the ventral fins deep black, the lower (outer) white; the inner surface of the pectorals blackish. Rio Motagua (east slope). a–b. From 8 to 9 inches long. From Mr. Salvin's collection. (Günther, Cat., v, 172.) The specimens from the Pacific Coast mentioned in the text (page 132) belong to the following species.

200(a). TACHYSURUS LIROPUS, Susan B. Bristol.

Head $3\frac{3}{7}$ to $3\frac{3}{4}$; depth $4\frac{3}{5}$ to $5\frac{1}{10}$. D. I, 6; A. II, 19; P. I, 9 or 10. Body elongate, its width anteriorally a little less than depth, the posterior portion much compressed; back elevated at front of dorsal; anterior profile from front of dorsal to tip of snout oblique; head flat, very broad, its width $1\frac{1}{3}$ in its length; snout broad, rounded, $1\frac{1}{10}$ to $1\frac{1}{5}$ in interorbital width; eye rather large, laterally placed, its width about $1\frac{3}{5}$ in its length, $4\frac{1}{6}$ to $4\frac{1}{10}$ in head; mouth small, upper jaw considerably projecting, its breadth $2\frac{3}{5}$ to 3 in head; jaws thin; wide bands of minute pointed teeth present on both jaws; vomerine bands widely separated, and indistinguishable from the palatine band, which is small, oblong-ovate, and scarcely prolonged backward; interval separating vomerine bands about $2\frac{1}{2}$ or 3 in eye; the teeth on these bands larger than those on jaws, and very bluntly conical. Interorbital space broad, $2\frac{1}{2}$ to 3 in head. Barbels long and slender, the maxillary barbel extending to, nearly to, or, in some cases, past base of pectoral, $1\frac{1}{4}$ to $1\frac{2}{3}$ in head; outer mental barbel $1\frac{1}{2}$ to $1\frac{2}{3}$ in head; inner mental barbel $2\frac{1}{3}$ to $2\frac{5}{6}$ in head. Antedorsal shield very short, narrowly crescent-shaped, its length on the median line about 2 or 3 in its width; occipital process subtriangular, a little longer than broad at base, its edge slightly concave, its median keel strong. The long, narrow groove of the fontanel beginning abruptly a short distance in front of occipital keel, the distance from its end to base of dorsal $1\frac{2}{3}$ to 2 in the distance to tip of snout. Shields of head rather smooth, finely granular, the granules forming distinct lines anteriorly. The flat

area between eyes triangular, with a median groove extending from fontanel forward to tip of snout, its posterior end a little behind eye, the granulations on each side of it extending forward as far as posterior border of pupil; opercles with no radiating striæ. Gill membranes forming a very narrow fold across the isthmus. Gill rakers 5 + 12. Nostrils 2 on either side, large, placed close together and near tip of snout, the posterior with a large flap; axillary pore well developed; humeral process smooth, very short, 4 to 5½ in pectoral spine. Base of dorsal 2⅜ to 2 9/10 in head; dorsal spine long and very strong, 1⅐ to 1⅛ in head, its upper anterior serræ small and tubercle-like, its upper posterior and its lower edges retrosely serrate; the soft rays extending considerably beyond the spine, 1⅛ to 1¼ in head; adipose fin small, its base 3½ to 4½ in head, its height 1⅜ to 2 in its base; caudal widely forked, the upper lobe, measured from base of caudal to its tip, the longer, about 1¼ in head; base of anal 1⅜ to 1⅔ in head, its longest ray 2⅛ to 2⅜ in head; ventrals reaching ⅚ to ⅞ the distance to origin of anal; vent about midway between origin of ventrals and origin of anal; pectoral spine 1½ in head, serrate, the serræ on inner edge larger and sharper than those on outer, the upper anterior serræ tubercle like, the rays a little longer than spine. Bluish silvery, light yellowish below; top of head and back brown; fins dusky olive, lighter at base, all margined with darker; ventrals pale; adipose fin covered with minute black dots; maxillary barbels dark brown, with bluish silvery luster; other barbels lighter; eye yellowish. Here described from 6 specimens from San Juan Lagoon, mouth of Rio Ahome, Sonora, Mexico (No. 47584, U. S. Nat. Museum). Length 7¼ to 9 inches.

Tachysurus liropus, Bristol, in Gilbert, Proc. U. S. Nat. Mus. 1896 (Feb. 5, 1897), 438, San Juan Lagoon, near mouth of Rio Ahome, Sonora, Mexico.

200(b). **TACHYSURUS EMMELANE**, Gilbert, new species.

Head 3⅖ in length (4 1/10 in total); depth 5 (6 in total). A. 27 (3 + 24). Eye 7 in head, 2½ in its distance from tip of snout, 4 in postorbital part of head, 3⅜ in interorbital width, 2⅕ in frontal width opposite middle of eyes. Mouth of moderate width, gently convex, the distance between its angles (measured internally) 2¼ in head. Teeth in premaxillary and front of mandible finely villiform; posterior mandibular teeth stronger than those in front, bluntly conic, not, however, granular or flat and pavement-like, as are the posterior mandibular teeth in *T. furthii*, *T. melanopus*, and *T. liropus*. Mandibular bands with a wide interspace mesially, each widest near symphysis, rapidly tapering laterally, and extending beyond angle of mouth. The width of the bands is less than in related species, ¼ eye at their widest point. The length of 1 of the mandibular bands is slightly greater (1 1/10) than length of eye. Premaxillary band very short, its length but ⅙ greater than that of 1 of the mandibular bands, extending on each side less than ½ distance from median line to angle of mouth; width of band ⅕ its length. Palatine teeth granular, in small oblanceolate patches, which taper to a point laterally, and are widely separated on medial line, the patches agreeing in size and shape with those in *T. liropus*. Head depressed, tapering and at the same time narrowing

adipose [illegible] as in other species of *Tachysurus;* profile rising in a uniform, gently convex curve to occiput, where it becomes concave, owing to the more rapidly ascending outline of the occipital process. Eye low, but little above angle of mouth, the interorbital space decidedly convex. Barbels slender, the maxillary barbels reaching edge of gill membrane in front of pectoral spine, the outer mental barbels extending beyond gill membrane 1¼ in head, the inner not to edge of membrane. Gill membrane widely attached to isthmus, without free edge. Occipital region with very fine granulations, those on middle of occiput forming parallel series along the fontanel groove, those on median portion of occipital process in series which diverge backward from the median line. The sculptured area extends forward to a vertical which traverses the cheek at a distance of its own diameter behind the eye; anterior edge of granulated area equidistant between tip of snout and front of dorsal plate; fontanel produced backward as a deep, narrow groove, which fails to reach base of occipital process by a distance equaling ½ the length of the process on the median line; the groove widening but little anteriorly; an area behind and on each side of the groove with parallel series of granulations, and marked off from the rest of the head by a shallow trench; base of occipital process similarly indicated by a transverse indented line; occipital process not keeled, very wide at base, becoming abruptly very narrow behind, its posterior ⅓ having parallel margins and being as wide as long, the lateral margins therefore deeply concave; width of process at base equaling its length on median line, plus that of dorsal plate, its hinder edge deeply incised to receive the anterior rounded wedge process of the dorsal plate, the latter finely granulated anteriorly, the lateral wings concealed under the smooth skin; a narrow groove as long as eye occupies the anterior end of the fontanel; no similar groove found in *T. furthii,* a short roundish one present in the type of *T. liropus,* and a continuous one the entire length of fontanel in the specimen which we identify with *T. melanopus;* opercle without radiating ridges; a short, slit-like axillary pore present; humeral process short, the exposed portion not broadly triangular, the surface smooth, or indistinctly rough. Gill rakers 6+13, of moderate length and thickness, the longest below the angle, ⅔ diameter of eye. Dorsal spine with a series of obtuse granulations in front and very weak retrorse serræ behind, its length to tip of calcified portion 1⅔ in head; longest soft ray 1⅓ in head; adipose dorsal not adnate, its anterior insertion about over middle of anal; distance between dorsals equal to length of head; base of adipose dorsal much greater than its height, less than base of first dorsal; pectoral spine strong, ridged and granulated in front, the hinder edge with very strong serræ; length of spine 1¼ in head, the fin projecting beyond tip of spine and reaching ⅔ distance from axil to base of ventrals; ventrals reaching to or nearly to origin of anal; vent midway between base of ventrals and front of anal; base of anal equaling length of pectoral spine; margin of anal gently concave, the longest ray 2⅔ in head; caudal with pointed lobes, the lower longest in the type, 1⅔ in head. Color dark steel blue or brownish above, becoming bright silvery below; posterior ⅔ of anal white

the anterior portion black with a narrow white edge; pectorals and ventrals with anterior (outer) face white or slightly dusky; pectorals with inner face of upper rays black; a black blotch covers all of inner face of ventrals, except terminal half of inner rays; barbels blackish. Closely related to *T. melanopus* and *T. multiradiatus*, differing from the former in the longer anal fin, from the latter in the black markings on lower fins. The description of the type of *T. multiradiatus* (*Bagrus? arioides*) Kner & Steindachner, Abhandl der K. bayer Akad. der Wissen, X, I, 1864, indicates a species with much rougher sculpturing of the head, a longer fontanel groove, narrower occipital process, and more anteriorly inserted adipose dorsal. The type is a single specimen, 280 mm. long, from Panama. (Gilbert.) (ἔν, in μελανη, ink.)

Tachysurus emmelane, GILBERT, Fishes Panama, MS. 1898, Panama.

201. TACHYSURUS FURTHII (Steindachner).

Head 3¼ to 3⅔; depth 5 to 5½. D. I, 7; A. 20. Body compressed posteriorly; profile slightly convex. Head broad, tapering forward, its greatest width 1⅔ to 1½ in its length; width, at the angle of the mouth, 2⅔ to 2⅘ in the head. Top of head densely covered with fine granules. Occipital process about as long as broad, with a median ridge, emarginate on its sides and at tip; interorbital region with 4 smooth ridges, the inner bordering the fontanel, the other extending obliquely backward from near the posterior nasal opening; sides of head and snout with vermiculating mucous pores. Middle of the fontanel over the pupil. Eye strictly lateral, not entirely above the angle of the mouth, its center in front of the posterior end of the mandible, 2 in snout, 6 to 7 in head, 3½ to 4 in the distance between the eyes. Maxillary barbels thin, reaching to the middle of the pectoral or shorter, postmentals beyond base of pectorals, or sometimes not beyond edge of gill membrane; mentals to edge of gill membrane or shorter. Jaws about equal, the upper rather thin; teeth on the intermaxillaries villiform; the mandible with villiform teeth except the inner 2 or 3 series, which are granular; like the palatine patches irregular, suboval, sometimes the anterior end, sometimes the posterior, and sometimes both ends pointed. Gill membranes united, joined to the isthmus without a free margin. Gill rakers long and slender, 4 + 11. Axillary pore small; vertical series of pores present. Distance of dorsal from snout 2⅔ to 2⅗ in the length; the dorsal spine 1⅗ to 1⅜ in the head, on sides and front granular, with small, sharp teeth on its inner margin; the first soft ray little, if any, higher than the spine. Distance of adipose fin from the dorsal 3 to 3⅛ in the length, the fin longer than high, shorter than the dorsal fin. Caudal fin forked, the lobes rounded, 4½ in the length. Anal little longer than high, the highest ray 2 in the length of the head. Ventrals short, 1⅞ to 2⅔ in head. Pectoral spine long and slender, 1¼ to 1⅗ in the head, outer margin granular, inner margin with short teeth. Ashy above, white below. We have examined 15 specimens, the largest measuring 0.29 m. The sexes do not differ externally. Panama. (Eigenmann & Eigenmann, Nematognathi, 90.)

202. TACHYSURUS VARIOLOSUS (Cuvier & Valenciennes), text, p. 132.

203. TACHYSURUS MULTIRADIATUS (Günther), text, p. 132.

75. CATHOROPS, Jordan & Gilbert, text, p. 133.

204. CATHOROPS HYPOPHTHALMUS (Steindachner), text, p. 133.

205. CATHOROPS GULOSUS (Eigenmann & Eigenmann), text, p. 133.

Page 134. After *Ictalurus furcatus* add:

206(a). ICTALURUS ANGUILLA, Evermann & Kendall.

(EEL CAT; WILLOW CAT.)

Head 4; depth $4\frac{1}{3}$; eye 7 in head; snout $2\frac{2}{3}$; interorbital $1\frac{1}{3}$; maxillary (without barbel) 3; free portion of maxillary barbel longer than head; dorsal spine 2 in head; pectoral spine 2; width of mouth 2. D. I, 6; A. 24; vertebræ 42. Head large, broad, and heavy; the mouth unusually broad; cheeks and postocular portion of top of head very prominent; interorbital space flat, a broad, deep groove extending backward to origin of dorsal fin; body stout, compressed posteriorly; back scarcely elevated. Eye small; maxillary barbel long, reaching considerably past gill opening; other barbels short. Origin of dorsal fin equidistant between snout and origin of adipose fin, its distance from snout $2\frac{3}{5}$ in length of body; base of dorsal fin $3\frac{1}{2}$ in head; longest dorsal ray $1\frac{3}{4}$ in head; dorsal spine strong, entire both before and behind; pectoral spine strong, entire in front, a series of strong, retrorse serræ behind; humeral process $2\frac{1}{3}$ in pectoral spine; ventrals barely reaching origin of anal, their length 2 in head; anal fin long and low, the longest rays about $2\frac{1}{3}$ in head; base of fin greater than head, $3\frac{1}{5}$ in body; caudal moderately forked, the middle rays about $2\frac{1}{3}$ in outer rays, which are about $1\frac{2}{3}$ in head. Color uniform pale yellowish or olivaceous; no spots anywhere.

An examination of the 6 cotypes shows that there is not much variation, all the important characters remaining quite constant. The maxillary barbel varies somewhat in length, in some individuals scarcely reaching gill opening, and the number of anal rays varies from 24 to 26.

A comparison of the skull of this species with that of *I. furcatus* and of *I. punctatus* of the same size shows a number of very marked differences. Nearly all the bones in *I. anguilla* are heavier than in the other species; the supraoccipital is broadly triangular, and its upper surface finely grooved, while in each of the other species it is much longer and narrower and the upper surface nearly smooth.

From the blue cat (*Ictalurus furcatus*) this species differs chiefly in the fewer rays in the anal fin, the wider mouth, the shorter, heavier head, the much longer maxillary barbel, and in the cranial characters already given. From the spotted cat (*I. punctatus*) it may be distinguished by its wider mouth, more blunt snout, heavier head, the color, and the cranial characters already mentioned.

The eel cat rarely attains a greater weight than 5 pounds, and usually does not exceed 3 pounds. Its flesh is firm and of excellent flavor. The spawning season appears to be during the spring, as several of the individuals examined were in mature spawning condition.* Lower Mississippi Valley; thus far known only from the Atchafalaya River, Louisiana and the Ohio River at Louisville. (*anguilla*, the generic name of the eel.)

Ictalurus anguilla, EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897 (Feb. 9, 1898), 125, pl. 6, fig. 1, Atchafalaya River, Louisiana. (Type, No. 48788. Coll. Evermann & Chamberlain.)

Recent studies of the catfishes of the Lower Mississippi Valley by Dr. Evermann have shown that the most abundant and most important species of catfish in that region is *Ictalurus furcatus* (Le Sueur), and not *Ameiurus lacustris* (Walbaum), as has hitherto been supposed. The large specimen described by Dr. Bean as *Amiurus ponderosus* is an *Ictalurus* (as shown by the skeleton now in the United States National Museum) and apparently *I. furcatus*. The common names "Great Fork-tailed Cat," "Mississippi Cat," and "Blue Cat" all belong to *I. furcatus*.

Page 138. The species called *Ameiurus dugesii* belongs to the genus *Villarius*, Rutter.

Page 142. After *Ameiurus nigrilabris* add:

77(a). VILLARIUS, Rutter.

Villarius, RUTTER, Proc. Cal. Ac. Sci., ser. 2, vol. VI. 1896, 256 (*pricei*).

Allied to *Ameiurus*, differing in the presence of scattered cilia on the sides. Backward process from occipital short, broad, emarginate, connected by ligament with the first interspinal buckler; in adults the distance between this process and the buckler is equal to the length of the former; in young examples the process overlaps the keel on the underside of the buckler. Head narrow, width of intermaxillary band of teeth $\frac{1}{3}$ of head; caudal deeply forked, the upper lobe the longer; barbels long, those of the maxillary extending past the gill opening. Sides with scattered hair-like cirri; these are very noticeable under a lens, but not readily distinguished by the naked eye. This genus differs from all others of the family in having hair-like cirri on the sides. It differs from *Ictalurus* in having the occipital process and the interspinal buckler widely separated and connected by ligament; from *Ameiurus* in having a narrow head and a deeply forked caudal. Two species known, the following and *Villarius dugesii* (Bean). (*villus*, a hair.)

* This species is well known to the fishermen of the Atchafalaya River, by whom it is usually called the "eel cat," though the name "willow cat" is sometimes applied to it. It was explained by the fishermen that the name "eel cat" was given on account of the long feelers (i. e., barbels) and the name "willow cat" because it is most frequently found about the roots of willow trees. The eel cat is not an abundant species in the Atchafalaya River. During six days (April 19–24) spent at Morgan City several hundred catfish were examined at the three fish houses, and the total number of eel cats seen was fewer than twenty-five. The fishermen report that this proportion is about as great as at any time of the year. Of the four commercial species of catfishes handled on this river the most abundant one is the blue cat (*Ictalurus furcatus*), and the next is the yellow cat or goujon (*Leptops olivaris*); the eel cat comes next and the spotted cat (*Ictalurus punctatus*) last. The blue cat and the yellow cat probably constitute 98 per cent of the entire catch.

220(a). VILLARIUS PRICEI, Rutter.

B. 8; D. I, 6; A. 22 or 23; C. 17; P. I, 9; V. 8. Head 3¼ to 3⅔ in body; eye 5 to 7 in head; snout 2⅔; maxillary 5½ to 6. Maxillary barbel very long, reaching beyond the pectoral spine, in the adult about to its tip when depressed, 3 to 4 times as long as the barbel at nostril. Origin of dorsal midway between snout and middle of base of adipose fin; pectorals inserted halfway between snout and ventrals; longest dorsal ray 6 to 7 times in length of body; spine of dorsal longer than its base, equal to base of adipose fin; longest pectoral ray about half of head, pectoral spine 2½ to 3 in head, with about 12 distinct hooked serræ behind, these fewer and somewhat smaller in the young; base of anal 3 times in its distance from snout, its longest ray equal to length of ventral; caudal deeply forked. Lateral line faint. This species differs from *V. dugesii* (Bean) in having very prominent serrations on the pectoral spines, the types of *dugesii* having the pectoral spines without serræ. We have examined a specimen of *dugesii*, 4 inches long, from Salamanca, Mexico, which is in the type basin; it has the cirri minute and light in color, a row of papillæ along the lateral line, and the pectoral spines with 4 or 5 degenerate serræ. (Named for William Wightman Price, who collected the type specimen.)

Villarius pricei, RUTTER, Proc. Cal. Ac. Sci., ser. 2, vol. VI, 1896, 257, San Bernardino Creek, a tributary of the Yaqui River, southern Arizona. (Type, No. 4826, L. S. Jr. Univ. Mus.)

Page 143. *Leptops olivaris* is known as the *Goujon* in Louisiana, where it is an important food-fish.

Page 146. In *Schilbeodes gyrinus* the anal rays are 14 to 16; not 13.

Page 152. Under *Rhamdia salvini* read "Osbert Salvin" for "Oscar Salvin."

Page 170. *Pantosteus arizonæ*, Gilbert, is described and figured in Proc. U. S. Nat. Mus. 1898, 488, pl. 36.

Page 174. *Catostomus discobolus* is distinct from the true *C. latipinnis*. The two species are confused in the description of *C. latipinnis* given by us. They may each be described as follows:

279. CATOSTOMUS LATIPINNIS, Baird & Girard.

Head 4; depth about 5½; eye high up and small, 5 to 7 in head, 3 to 3½ in snout, 2⅛ to 2¾ in interorbital space; interorbital width 2⅔ in head. D. 14 or 15; A. 7; scales 19 or 20–89 to 102–16 to 18, 46 to 50 transverse rows in front of dorsal fin. Head depressed and flat above, its greatest depth 1⅔ in its length, the depth below lower edge of orbit 3 in its length. Least depth of caudal peduncle 4½ in head, or 3½ in its own length. Fins very large, the dorsal with its upper margin concave; ventrals and pectorals rounded; dorsal as long as its longest ray, $1\frac{1}{10}$ in head, its last ray a little less than ½ the length of the first ray; origin of dorsal fin nearer tip of snout than base of caudal; ventrals not reaching quite to vent, 1⅔ in head. Muzzle not projecting; about 6 rows of short, thick papillæ on upper lip, the smallest above; lower lip large, incised to its base, with

about 12 rows of short, thick papillæ, posteriorly quite small; distance from front of upper lip to back of lower 1½ in snout; jaws with a slight cartilaginous sheath; width of preorbital a little less than ½ its length. Reaches a length of about 2 feet. Lower Colorado River basin. The description by Gilbert & Scofield, based upon specimens from the Gila River at Tempe, Arizona.

279(a). CATOSTOMUS DISCOBOLUS, Cope.

Head 3⅞ to 4½; depth about 5½; eye small, high up, 5½ to 6 in head, 2½ in snout, 2⅔ in interorbital width, which is 6½ in head; width of preorbital less than ½ its length; least depth of caudal peduncle 2½ in its length, or 2 in head; greatest depth of head 1⅗ in its length; depth from lower edge of orbit 3¼ in head. D. 12 or 13; A. 7; scales 19 to 21–101 to 109–17 to 20 52 to 63 in front of dorsal. Muzzle projecting slightly beyond upper lip. Upper margin of dorsal very slightly concave, the length of its base 1 in its longest ray, or 1½ in head; last dorsal ray ½ length of first; origin of dorsal midway between tip of snout and base of caudal. Ventral rounded, 1⅘ in head, not quite reaching vent. Mouth as in *C. latipinnis* except that the posterior tubercles on lower lip are long and not nearly so closely set, there being 9 or 10 rows; jaws with a slight cartilaginous pellicle. Upper portion of the Colorado River basin. Attains the length of a foot or more. The above description by Gilbert & Scofield from specimens from Green River at Green River Station, Wyoming.

Catostomus discobolus, Cope, Hayden's Geol. Surv. Wyo., 435. 1870, Green River, Wyoming; Gilbert & Scofield, Proc. U. S. Nat. Mus. 1898, 490.

Page 175. After *Catostomus griseus* add:

280(a). CATOSTOMUS RETROPINNIS, Jordan.

A doubtful species which is, however, not yet shown to be invalid. Head 4⅛; depth 5⅛; eye 6¾ in head; snout 2; interorbital 2½. D. 11. A. 7. scales 17–108–14. Body slender, head slender, snout very long, caudal peduncle long, its least depth less than snout, 2½ in head; dorsal profile very little elevated; mouth large, wholly inferior, overhung by the pig-like projecting snout; lips thin but very broad, lower lip inclosed nearly to base, with about 6 rows of moderate papillæ; lobes of lower lip very long, about ½ of snout; gill rakers short and weak. Origin of dorsal a little nearer base of caudal than tip of snout, base of dorsal equal to snout; longest dorsal ray a little greater than base of fin; anal fin long and pointed, the fourth ray longest, 1 1/10 in head; caudal lunate, the middle ray 1½ in outer rays; pectoral somewhat falcate, the longest 1½ in head; ventrals rather short, not reaching vent by more than an eye's diameter. Length 14 inches.

Catostomus retropinnis, Jordan, Bull. U. S. Nat. Mus., xii, 178, 1878, Milk River, Montana. (Type, No. 21197. Coll. Dr. Elliott Coues.)

Page 176. After *Catostomus catostomus* add:

282(a). **CATOSTOMUS RIMICULUS,** Gilbert & Snyder.

Head 4⅙ in body; depth 5; depth of caudal peduncle 2¾ in head; eye 7⅓; dorsal rays 11; scales 18–91–13, before dorsal 42. D. 11; A. 7; pectoral 17. Head as deep as wide. Both lips full, the lobe of lower lip broadly rounded behind, the cleft not nearly reaching base of lip, the portion between mandible and apex of cleft with 4 series of tubercles; tubercles coarse and blunt, becoming reduced in size toward margins of lips, but less so than in related species; upper lip with 5 rows of tubercles. Eyes very small, the front of the eye nearly midway of head; interorbital space convex, 2⅘ in head. Scales comparatively smooth, gradually growing smaller posteriorly. Dorsal fin inserted midway between end of snout and base of caudal, first ray preceded by 2 short, simple ones; last ray divided to base; length of base of fin equal to the height, which is contained 6½ times in the body; height of anal twice the length of the base, 5 in body; length of pectorals 4⅜ in body; ventrals 6⅓ in body; caudal 4¼. Color above dusky, the central parts of scales lighter; under parts white; dorsal and caudal fins dusky, others white. This species belongs to the *C. catostomus* type, with very small scales, and is most nearly related to *C. tahoensis*. From the latter it differs in the smaller eye, less deeply cleft lower lip, blunter labial tubercles, larger scales, and the much smaller fontanel, which is reduced in adults to a very narrow linear slit, or more commonly entirely obsolete. Lower portion of the Klamath River basin, northern California. (Diminutive of *rimus*, crevice, from the small fontanel.)

Catostomus rimiculus, Gilbert & Snyder, Bull. U. S. Fish Comm. 1897 (Jan. 6, 1898), 3, Trinity River, Humboldt County, California. (Type, No. 5654, L. S. Jr. Univ, Mus. Coll. Capt. W. E. Dougherty.)

Page 177. *Catostomus rex* is identical with *Deltistes luxatus* and should be added to the synonymy of that species, p. 183.

The type of *Catostomus labiatus* did not come from Klamath Lake, but from the Sacramento River, at Stockton, California. It is identical with *C. occidentalis*. The species from Klamath Lake has been recently described as

285. **CATOSTOMUS SNYDERI,** Gilbert.

Head 4⅛ in length; snout 2 $\frac{3}{10}$ in head, equaling interorbital width; eye 5¾. D. 11; A. 7; scales 13 or 14–69 to 77–10 or 11. Mouth very small, the width between angles but ½ length of snout in our largest specimen; greatest width of lobe of lower lip ⅔ diameter of eye; lower lip deeply incised, with 1 or 2 papillæ between symphysis and base of cleft; upper lip narrow, with 5 or 6 papillæ in a cross series, the uppermost becoming very small; basal portion of the lower lip with coarse tubercles, those toward posterior margin becoming very fine and arranged in evident series separated by grooves. Mucous canals on head forming conspicuous raised ridges with prominent pores, the system much more conspicuously developed than in any related species. Origin of dorsal fin constantly

nearer snout than base of caudal; the dorsal fin short, its base not exceeding the height of the longest ray, usually less. In our specimens the pectorals reach scarcely $\frac{2}{3}$ distance to ventrals and the ventrals scarcely $\frac{2}{3}$ distance to vent; the anal may extend beyond base of rudimentary caudal rays. Scales strongly ridged, their margins crenate; the anterior scales are smaller, but do not appear greatly crowded; the average number of tubes in the lateral line is about 73, the number varying from 69 to 77. Dusky, the lower part of sides with coarse black specks, the under parts white; fins all dusky. (Gilbert.)

A larger specimen has been described as follows: Head $4\frac{1}{5}$; depth 4; eye $6\frac{1}{2}$ in head; snout $2\frac{1}{2}$; maxillary $3\frac{1}{5}$; mandible $2\frac{1}{5}$; interorbital $2\frac{1}{8}$; width of mouth $3\frac{1}{2}$ in head, more than $\frac{1}{2}$ length of snout; greatest width of lower lip $\frac{3}{4}$ diameter of eye. D. II, 11; A. 7; scales 13–70–11. Body rather slender; head long, mouth moderate, horizontal; lips thick papillose, the upper with about 4 or 5 rows of papillæ, lower with about 7; lower lip divided nearly to base, leaving only 1 row of papillæ crossing the symphysis; premaxillary not much projecting and not forming a prominent hump; maxillary rather short, not reaching vertical at front of anterior nostril; eye equally distant between snout and posterior edge of opercle; mucous canals on head forming raised ridges, the pores conspicuous. Fins moderate; origin of dorsal a little nearer snout than base of caudal, sixth spine over insertion of ventrals; pectoral $1\frac{1}{4}$ in head, reaching slightly more than $\frac{2}{3}$ distance to ventrals; ventrals not quite reaching vent, the seventh ray longest, $1\frac{2}{3}$ in head; anal long, pointed, reaching to base of caudal, $1\frac{1}{6}$ in head. Scales crowded anteriorly, about 32 transverse rows in front of dorsal, strongly ridged, the margins crenate. (Evermann & Meek.) Length 1 to 2 feet. Klamath Lakes, Oregon; specimens examined from Upper Klamath Lake, Lost River, and Williamson River. (Named for Mr. John O. Snyder, instructor in Zoology in Stanford University.)

Catostomus snyderi, Gilbert, Bull. U. S. Fish Comm. 1897 (Jan. 6, 1898) 3, Upper Klamath Lake, Oregon (Type, No. 48222. Coll. Gilbert, Cramer & Otaki); Evermann & Meek, Bull. U. S. Fish Comm. 1887, 69.

Page 178. After *Catostomus occidentalis* add:

286(a). CATOSTOMUS TSILTCOOSENSIS, Evermann & Meek.

Head $4\frac{1}{2}$; depth 5; eye $6\frac{1}{2}$ in head; snout 2. D. 13; A. 7; scales 13–65–8, 34 before the dorsal. Pectoral $1\frac{1}{8}$ in head; longest dorsal ray $1\frac{2}{5}$; base of dorsal $1\frac{2}{5}$; longest anal ray $1\frac{1}{4}$; ventral $1\frac{3}{4}$. Body rather slender, subterete; head small, snout long and pointed; mouth inferior, overhung by the projecting snout; lips rather thin, 1 row of large papillæ on upper lip, and about 2 irregular rows of smaller ones behind or inside of it; lower lip incised nearly to base, 1 or 2 rows of small papillæ across the isthmus; lobes of lower lip moderately long and thin, the bases with papillæ merging into plications toward the tips. Eye quite small, the anterior edge of orbit at middle of head. Top of head flat or very slightly convex between the eyes. Fins small; pectorals short and rounded; ventrals short, rounded,

the middle rays but little longer than the others; anal small, somewhat pointed; margin of dorsal somewhat concave; caudal emarginate, not deeply forked. Mucriferous canals on head not strongly developed. Scales moderately large; lateral line nearly straight, not running upward toward nape. This species differs from *C. occidentalis*, to which it is related, in the smaller head, longer, more pointed snout, smaller eye, larger scales, and its much smaller fins. In *C. occidentalis* the pectoral fins are falcate, while in this species they are more rounded; the ventrals also are less pointed. Length a foot or less. Coastal streams of middle western Oregon; known from Tsiltcoos Lake and the Siuslaw River. (*tsiltcoosensis*, from the type locality.)

Catostomus tsiltcoosensis, EVERMANN & MEEK, Bull. U. S. Fish Comm. 1897 (Jan. 6, 1898), 68, fig. 1, Tsiltcoos Lake, Lane County, Oregon. (Type, No. 48479. Coll. Dr. Seth E. Meek.)

Page 180. The species called *Catostomus fecundus* in the text belongs in the genus *Chasmistes*, to which it should be transferred as *Chasmistes fecundus* (Cope & Yarrow).

Page 182. The species of *Chasmistes* are not confined to the Great Basin. One species (*C. brevirostris*) occurs in the Klamath Lakes basin.

Page 183. The species called *Chasmistes luxatus* in the text belongs to a genus distinct from *Chasmistes*, which may be characterized as follows:

93(a). DELTISTES, Seale.

Deltistes, SEALE, Proc. Cal. Ac. Sci., ser. 2, vol. VI, 1896, 269 (*luxatus*).

This genius is close to *Chasmistes*, agreeing with it in every respect except in the peculiar structure of the gill rakers. In *Chasmistes* [illegible] as in *Catostomus*, while in *Deltistes* they are broad, [illegible] the Greek letter Δ (delta), and their edges are unarmed and [illegible]; [illegible] gegals weak, with numerous small teeth. *Deltistes luxatus* (Cope) is the single known species. (δέλτα, the Greek letter corresponding to D.)

After *Chasmistes cujus* add:

297(a). CHASMISTES STOMIAS, Gilbert.

Head 4¼, depth 4¼; eye 7; snout 3⅓; maxillary (measured from free end to tip of snout) 3⅜; mandible 2½. D. II, 11; A. I, 7; scales 13–85–10; interorbital [illegible] width of head at mandibular articulation 2½. Head small, body heavy forward, the back strongly and regularly arched from snout to origin of dorsal fin, thence declined in a nearly straight line to base of caudal; ventral surface nearly straight. Premaxillary spines strongly protruding, forming a prominently projecting snout, mouth rather small, inclined upward at an angle of about 40°, maxillary scarcely reaching vertical from front of anterior nostril; width of mouth 1⅔ in snout or 4⅓ in head; upper lip thin, without papillæ; lower lip thin, interrupted at symphysis, forming narrow lateral lobes, the width of which is about 2⅔ times in their length; faint indications of a few papillæ; mucous canals forming ridges, the pores conspicuous; gill rakers long,

narrowly triangular at the tip when viewed from behind, densely tufted on the anterior edge; fontanel narrow, its length $2\frac{2}{3}$ in the snout, its width about $\frac{1}{7}$ its length. Fins all large; the origin of the dorsal a little nearer tip of snout than base of caudal, the sixth ray over base of ventral, its base $1\frac{3}{5}$ in head, the free edge nearly straight, the last ray $1\frac{3}{8}$ in the first, which is $1\frac{1}{2}$ in head; pectorals scarcely falcate, reaching a little more than $\frac{3}{4}$ distance to base of ventrals, their length $1\frac{1}{4}$ in head; ventrals long, reaching vent, the rays gradually increasing in length from the outer to the seventh and eighth, which are longest, the ninth and tenth being but slightly shorter, the length of the longest ray $1\frac{3}{5}$ in head or about $\frac{1}{5}$ longer than the first; anal long and pointed, the fourth ray longest, reaching base of caudal, $1\frac{1}{6}$ in head; each ray of anal fin with 8 to 12 strong tubercles; caudal lobes about equal, their length $1\frac{2}{3}$ times the middle ray. Length a foot or more. Upper Klamath Lake, Oregon, where it is abundant and of some importance as a food-fish. The Klamath Indian name is K-ahp-tu. (στομίας, large-mouthed.)

Chasmistes stomias, Gilbert, Bull. U. S. Fish Comm. 1897 (Jan. 6, 1898), 5, with plate, Upper Klamath Lake, Oregon (Type, No. 48223. Coll. Gilbert, Cramer & Otaki); Evermann & Meek, Bull. U. S. Fish Comm. 1897, 70.

297(b). CHASMISTES COPEI, Evermann & Meek.

Head $3\frac{3}{4}$; depth 4; eye $6\frac{1}{2}$; snout $2\frac{1}{2}$; maxillary (measured from free end to tip of snout) 3; mandible $2\frac{2}{3}$. D. II, 10; A. I, 7; scales 13–80–12; interorbital width $2\frac{1}{7}$; vertical depth of head at mandibular articulation $2\frac{1}{5}$. Head large, cheek very deep, the depth equal to distance from tip of snout to nostril; body stout, back scarcely elevated, caudal peduncle rather short and stout; ventral surface somewhat convex. Premaxillary spines less protruding than in *C. stomias*, not forming a prominent hump; mouth large, inclined upward at an angle of 45°, maxillary not nearly reaching vertical at front of anterior nostril; width of mouth $1\frac{1}{4}$ in snout, or 4 in head; upper lip thin, without papillæ; lower lip thin, entirely without papillæ, interrupted at symphysis, forming rather broad lateral lobes; pores on head very conspicuous; gill rakers larger than in *C. stomias*, broadly triangular at tip when viewed from behind, densely tufted on anterior edge, each appendage more or less bifid and club-shaped, closely resembling those of *C. horax*; fontanel narrow, its length $2\frac{1}{3}$ in snout, width $\frac{1}{5}$ its length. Fins all small; origin of dorsal a little nearer snout than base of caudal, its sixth ray over base of ventrals, free edge straight, base $2\frac{1}{3}$ in head, last ray a little less than 2 in first, which is 2 in head; pectorals somewhat falcate, reaching slightly more than $\frac{1}{2}$ distance to ventrals, their length $1\frac{2}{3}$ in head; ventrals very short, reaching only $\frac{2}{3}$ distance to vent, free end nearly straight; outer ray longest, $2\frac{1}{3}$ in head, inner shortest, $3\frac{1}{3}$ in head; anal fin short, bluntly pointed, not reaching base of caudal, third and fourth rays longest, $1\frac{1}{4}$ in head; no tubercles on anal rays; caudal lobes equal, length about $1\frac{1}{3}$ times the middle ray. Scales small and crowded anteriorly, about 14 rows downward and backward from front of dorsal to lateral line, 11 vertically upward from base of ventral to lateral line, about 38 oblique series before

dorsal; lateral line nearly straight, with about 80 scales. Entire upper parts of head and body, and sides nearly to level of base of pectorals, dark olivaceous; under parts abruptly whitish or yellowish in alcohol; a dark spot in upper part of axil; dorsal and caudal dark; pectorals dark on inner surface; ventrals and anal plain. From *Chasmistes stomias* this species is readily distinguished by its larger head, larger, more oblique mouth, less prominent snout, and very small fins. The differences in the fins are very great, particularly in the ventrals. It differs from *C. brevirostris*, as characterized by Dr. Gilbert, in its much larger, more oblique mouth, the absence of papillæ on the lips, and shorter fins. Length 2 feet. Upper Klamath Lake, Oregon. Klamath Indian name "Tswam." (Named for the late Prof. Edward Drinker Cope, who wrote the first paper on the fishes of Upper Klamath Lake.)

Chasmistes copei, EVERMANN & MEEK, Bull. U. S. Fish Comm. 1897 (Jan. 6, 1898), 70, fig. 3, Pelican Bay, Upper Klamath Lake, Oregon. (Type, No. 48224. Coll. Meek & Alexander.)

Page 205. *Campostoma pricei* can not be distinguished by us from *C. ornatum*. See Rutter, Proc. Cal. Ac. Sci., ser. 2, vol. VI, 1896, 259.

Page 211. After *Algansea tincella* add:

337(a). ALGANSEA TARASCORUM, Steindachner.

Head $3\frac{2}{5}$; depth $4\frac{2}{5}$; eye less than 5; snout about 4; interorbital 3. D. III, 7; A. III, 6; P. 17; V. 9; scales 84 or 85, 18 or 19. Body stouter than in *A. lacustris*, head shorter, lateral line more decurved and nearer ventral line at middle of body, and scales more numerous. Mouth very oblique, lower jaw not projecting, maxillary not quite reaching vertical at anterior edge of eye. Teeth 4-4, hooked, and with narrow grinding surface. Origin of dorsal in advance of ventrals, equally distant between base of caudal and middle of eye; height of dorsal twice its base. Ventrals not reaching anal fin by an eye's diameter; caudal deeply notched. A dark gray longitudinal band with metallic luster extending from opercle to caudal fin, lying chiefly above lateral line; color otherwise plain. Length $5\frac{1}{2}$ inches. Lake Pátzcuaro, Mexico. (Steindachner.)

Algansea tarascorum, STEINDACHNER, Einige Fischarten Mex., II, pl. 3, figs. 2-2*c*, 1895, Lake Pátzcuaro, Mexico. (Coll. Princess Theresa von Bayern.)

Page 218. To the synonymy of *Pimephales notatus* add:

Spinicephalus fibulatus, LE SUEUR, in VAILLANT, Bull. Soc. Philom., VIII, 1896, 29, pl. 26.

Page 225. After *Ptychocheilus oregonensis* add:

358. PTYCHOCHEILUS GRANDIS (Ayres).

(SACRAMENTO PIKE.)

This species differs from *P. oregonensis* principally in the larger size of the scales above the lateral line, the smaller number of rays in the dorsal fin, and the lighter and slenderer pharyngeal bones. Head $3\frac{1}{8}$ to $3\frac{1}{7}$ in length; depth 5 to $5\frac{1}{10}$; eye $3\frac{2}{5}$ to 4 in head; scales 13 to 16 above lateral

line, 70 to 80 transverse rows along lateral line (16 to 18 above lateral line, 69 to 72 transverse rows in *P. oregonensis*). D. 8; A. 8. In other respects similar to *P. oregonensis*. *Ptychocheilus harfordi* is apparently not distinct from *P. grandis*, being based on a specimen with very small scales. *P. grandis* is confined to waters of California, *P. oregonensis* to Washington, Idaho, and Oregon.

Gila grandis, AYRES, Proc. Cal. Ac. Nat. Sci. 1854, 18, San Francisco.
Ptychocheilus major, AGASSIZ, Am. Jour. Sci. Arts 1855, 229, San Francisco.
Ptychocheilus harfordi, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 72, Sacramento River (Type, No. 27246. Coll. Jordan & Gilbert); JORDAN & GILBERT, Synopsis, 226 1883.

Page 239. After *Leuciscus balteatus* add:

376(a). LEUCISCUS SIUSLAWI, Evermann & Meek.

Head $4\frac{1}{5}$; depth $4\frac{1}{6}$; eye 4; snout $3\frac{1}{2}$; maxillary $3\frac{2}{5}$. D. II, 9; A. II, 12 or 13; scales 11–58–8; teeth 2, 4–5, 2, somewhat hooked. Body rather slender, slightly elevated and somewhat compressed; head small and pointed, cheek not deep; snout pointed, somewhat longer than eye; mouth moderate, somewhat oblique, maxillary just reaching vertical at front of orbit; jaws subequal, the lower sometimes slightly projecting; eye large, not as great as snout. Origin of dorsal fin behind base of ventrals and much nearer base of caudal than tip of snout, the longest ray $1\frac{1}{2}$ in head, greater than base of fin; origin of anal fin under last dorsal ray but 2, its height equal to that of dorsal, its base equal to its longest ray; free edges of dorsal and anal nearly straight; pectoral $1\frac{1}{4}$ to $1\frac{1}{2}$ in head, not reaching insertion of ventrals; ventrals short, $1\frac{1}{3}$ in pectoral, reaching anus; caudal deeply forked; lateral line complete, decurved. Color in spirits, brownish or olivaceous above, middle of side with a broad dark band involving the lateral line anteriorly and posteriorly, but lying chiefly above it mesially; middle of side from gill opening to beneath dorsal fin with a broad rosy band, following closely beneath the lateral line; lower part of sides and under parts silvery, dusted over with fine dark specks; a light yellowish band extending backward from upper posterior border of eye nearly halfway to origin of dorsal fin; cheek with a silvery or golden crescent; top of head dark; opercles dusky silvery; snout dusky; fins plain, dorsal and caudal somewhat dusky. This species is close to *L. balteatus*, but has smaller anal and dorsal fins, a more slender body, smaller and more slender head, and longer, more pointed snout. The extent of variation in proportional measurements and in the number of anal fin rays appears to be much less than in *L. balteatus*. It also resembles *L. cooperi*, but has a much shorter lower jaw and a more pointed snout. Known only from the Siuslaw River and Tsiltcoos Lake, western Oregon, where it is common. (*siuslawi*, of the Siuslaw River.)

Leuciscus siuslawi, EVERMANN & MEEK, Bull. U. S. Fish Comm. 1897 (Jan. 6, 1898), 72, fig. 4, Siuslaw River, Mapleton, Oregon. (Type, No. 48480. Coll. Dr. Meek.)

Page 240. After *Leuciscus elongatus* add:

378(a). LEUCISCUS NACHTRIEBI, Cox.

Head $4\frac{1}{4}$ to $4\frac{1}{2}$; depth 5 ($4\frac{1}{3}$ to $5\frac{1}{2}$); eye 4; snout $4\frac{2}{3}$. D. 8; A. 8. Body rather heavy, not greatly compressed; back slightly elevated, its curve a little greater than that of the belly; caudal peduncle rather stout, its depth $\frac{1}{2}$ the length of the head. Head rather short, not any more compressed than the body, upper surface slightly flattened; snout quite blunt in mature individuals, its length $1\frac{1}{6}$ times width of eye; mouth not very large, but little oblique, lower jaw included; maxillary scarcely reaching to front of orbit; pharyngeal teeth 2, 4–5, 2. Dorsal fin inserted nearer base of caudal than tip of snout, also slightly back of ventrals; caudal fin forked; anal slightly smaller than dorsal; ventrals small, not reaching vent by $\frac{1}{3}$ their length; pectorals inserted rather high, not reaching the ventrals by $\frac{2}{3}$ their length; scales small, 12–72–9, lateral line complete on mature individuals, decurved, the pores extending on head in several lines, 1 passing back of eye, another down to nostril. General color dusky, darkest on back; sides above lateral line dull silvery, below lateral line light silvery; a faint dark dorsal band in some specimens, in others absent; no black lateral band, but some specimens have a very faint dusky shade along lateral line; no light stripe above lateral line; upper portion of opercles with a dusky shade, lower part bright silvery; upper part of head dark-colored; all the above colors typical in the young as well as adults. Length 4 inches. *L. nachtriebi* differs from *L. neogæus* in having a well-developed lateral line, a smaller eye, fewer scales, less oblique mouth, a shorter maxillary, and in being a larger fish and differently colored. It differs from *L. elongatus* in having a smaller mouth, the lower jaw never projecting, head less pointed, a shorter maxillary, finer scales, and the absence of the black lateral band. Lakes of northern Minnesota; at present known from Mille Lacs, Man Trap, Mud and Elbow lakes. (Named for Prof. Henry F. Nachtrieb, State zoologist of Minnesota.)

Leuciscus nachtriebi, Cox, Rept. U. S. Fish Comm. 1894 (Dec. 14, 1896), 615, Mille Lacs Lake, Aikin County, Minnesota. (Type, No. 47688. Coll. Minn. Nat. Hist. Surv.)

Page 241. To the synonymy of *Leuciscus neogæus* add:

Cyprinus burtonianus, Le Sueur in Vaillant, Bull. Soc. Philom., viii, 1896, 28, with plate, Burton Mine, Missouri.

Page 244. *Leucos* and *Myloleucus* can not be maintained as subgenera, the characters of the teeth not being constant.

The following notes on *Rutilus oliraceus* as seen at Emerald Bay, Lake Tahoe, may prove useful.

385. RUTILUS OLIVACEUS (Cope).

(Tahoe Chub.)

This species is very different from *Rutilus symmetricus*, looking like *Leuciscus lineatus*. Very common; reaches 2 to 3 pounds weight; devours eggs of trout. No doubt the records of *Leuciscus lineatus* (*atrarius*) from Lake Tahoe belong to this species. Head 4; depth $4\frac{1}{3}$ to $4\frac{1}{2}$. D. 8; A. 8; scales 11–56–6; teeth always 5–5, with broad grinding surface. Body oblong,

moderately compressed, the back somewhat elevated anteriorly in old examples. Head conical, rounded above; eye moderate, 1⅓ in snout (6 inches long), 5 in head; about as long as maxillary. Mouth terminal, very oblique, the lower jaw included; the snout not prominent; the short maxillary not reaching eye. Dorsal high and pointed; anal short, rather high; pectoral long, reaching ⅔ distance to ventrals, which reach vent; ventrals below front of dorsal, which is behind middle of body. Scales with edges largely exposed; lateral line running low, complete. Dusky olive above and on sides to level of ventrals, with brassy luster everywhere; middle of belly only white, a pale yellowish area between pectorals and ventrals; head brassy, dusky above, closely dotted above and on sides; body everywhere closely dotted with black, except on middle line below; fins all dusky, with dark points. This species is well separated from all the *R. symmetricus* tribe.

Page 247. *Luxilinus occidentalis* is the young of *Lavinia exilicauda*, Baird & Girard (p. 209), and must be placed in the synonymy of that species. *Luxilinus* is a pure synonym of *Lavinia*.

Page 245. Under *Opsopœodus bollmani*, for "Buckland Creek" read "Buckhead Creek."

Page 254. For *Azteca*, line 22, substitute *Aztecula*, Jordan & Evermann, new subgenus. The former name is preoccupied by *Azteca*, Forel, 1878, a genus of ants. The same substitution to be made in the key on page 255 and on page 258.

Page 260. Before *Notropis cayuga* insert:

404(a). NOTROPIS WELAKA, Evermann & Kendall.

Head 4⅓; depth 5; eye 3 in head; snout 3¼. D. 8; A. 8 or 9; scales 6-35-3; teeth 4-4, hooked. Body rather slender, moderately compressed; head short, snout bluntly pointed; mouth moderate, somewhat oblique, lower jaw slightly included, maxillary scarcely reaching front of eye; premaxillaries protractile. Eye large; posterior edge of pupil at middle of longitudinal length of head; interorbital width greater than eye; caudal peduncle long and slender. Dorsal fin inserted well behind base of ventrals, a little nearer base of caudal than tip of snout, its longest rays shorter than head, but slightly longer than longest anal rays; anterior dorsal and anal rays longest; pectoral 1⅓ in head; ventrals reaching origin of anal; caudal deeply notched, the lobes long and pointed. Scales large, lateral line incomplete, developed only on 6 to 10 scales. Back olivaceous; side with a broad black band extending from snout through eye, and ending in a rather distinct black spot on base of caudal, the black spot in some specimens (probably mature males) surrounded by orange; the black line bordered above by a narrow orange or reddish line, less distinct, or even whitish, in females and immature individuals; under parts plain; fins all plain; dorsal and caudal somewhat dusky; dusky specks on body along base of anal and under side of caudal peduncle; lower jaw tipped with dusky. This species resembles *Notropis anogenus*, but differs in having the mouth somewhat larger and less oblique,

the lower jaw more included, the body more slender, the lateral line less developed, the dorsal fin more posterior, and the anal rays more numerous. It was found in considerable abundance in the St. Johns River, near Welaka, Florida. (*welaka*, from the type locality.)

Notropis welaka, EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897 (Feb. 9, 1898), 126, pl. 6, fig. 2, St. Johns River, near Welaka, Florida. (Type, No. 48786. Coll. Dr. W. C Kendall.)

Page 262. After *Notropis blennius* add:

408(a). **NOTROPIS BUCHANANI**, Meek.

Head 4; depth 4. D. 8; A. 8; scales 6-31-2; teeth 4–4. Body rather robust, back considerably elevated, snout blunt, mouth small and nearly horizontal. Snout short, about $\frac{3}{4}$ diameter of eye. Preorbital bone slightly longer than broad. Eye moderate, 3 in head. Lateral line complete, or nearly so; about 12 scales in a series before dorsal fin. Dorsal fin slightly nearer tip of snout than base of caudal; pectorals reaching ventrals; ventrals reaching anal. Color light olivaceous, a faint silvery lateral band; no dark lateral band or black caudal spot. This species belongs to the *N. blennius* type. It is a smaller species, lighter in color, and has fewer scales in the lateral line. Poteau River, Arkansas. (Named for Dr. John L. Buchanan, president of the Arkansas Industrial University.)

Notropis buchanani, MEEK, Bull. U. S. Fish Comm. 1895 (April 13, 1896), 342, small creek near Poteau, Indian Territory. (Type, No. 47532. Coll. Dr. Meek.)

Page 267. Under *Notropis nux; nuece*, not *neche*, is nut in Spanish.

Page 274, line 11, for *luxoides*, read *luxiloides*.

Page 287. After *Notropis lutipinnis* insert:

466(a). **NOTROPIS CHAMBERLAINI**, Evermann, new species.

Head $4\frac{1}{3}$; depth $4\frac{1}{3}$; eye 4; snout 4. D. 7; A. 9; scales 7–39–3, about 15 before the dorsal. General form much like that of *Hybognathus;* body only moderately compressed, dorsal and ventral outlines slightly arched; head rather small, pointed; mouth small, a little oblique, the maxillary scarcely reaching anterior border of orbit, lower jaw slightly included; snout equal to eye; eye in axis of body. Fins all rather small; origin of dorsal slightly behind vertical at insertion of ventrals; free edge of dorsal fin somewhat concave, the anterior rays about equal to length of head; pectoral short, slightly falcate, the longest rays about $1\frac{2}{3}$ in head; ventrals shorter than pectoral, barely reaching vent; anal similar to dorsal, the rays shorter; caudal widely forked, the middle rays $2\frac{1}{2}$ in the outer, the lobes as long as head, the lower lobe slightly longer than the upper. Scales moderately imbricated, the exposed portions not deeper than long; lateral line complete, somewhat decurved. Teeth 2, 4–4, 2 or 1, rather weak, hooked, and with small grinding surface. Intestine short; peritoneum silvery. General color light straw; middle of side with a broad, well-defined silvery band from upper end of gill opening to middle of

base of caudal fin, the anterior half lying wholly above the lateral line, the posterior portion lying partly below it; this silvery band bounded above by a narrow dark border; cheeks and opercles silvery; a darkish band along median line of back; fins all plain straw color or pale lemon. Fourteen examples of this species, 2 to 3 inches in length, were obtained from the Atchafalaya River at Melville, Louisiana, by Mr. Fred M. Chamberlain, for whom the species is named.

Notropis chamberlaini, EVERMANN MS., Atchafalaya River, Melville, Louisiana. (Type, No. 48901.)

Page 291. *Notropis scopifer*, Eigenmann & Eigenmann, is identical with *Notropis hudsonius selene* (Jordan), (p. 269), and should be omitted.

Page 294. After *Notropis dilectus* insert:

487(a). NOTROPIS LOUISIANÆ, Evermann, new species.

Head $4\frac{2}{5}$; depth $5\frac{1}{4}$; eye 3; snout 3. D. 7; A. 11; scales 7–37–3, 19 or 20 before the dorsal. Teeth 1, 4–4, 2, little hooked. Body long and slender, back not arched; head short, but pointed; mouth rather large, oblique, maxillary scarcely reaching orbit, lower jaw somewhat included; eye large, equal to or greater than snout. Fins rather small; origin of dorsal far behind insertion of ventrals, its longest rays $1\frac{2}{3}$ in head; pectorals short, their length equal to height of anal; ventrals very short, 2 in head; caudal deeply forked. Scales firm, moderately imbricated; lateral line complete, gently decurved. Color pale; side with a faint plumbeous band; back and upper part of sides with numerous dark specks chiefly on the margins of the scales, thus forming cross-hatchings; a narrow dark vertebral band on caudal peduncle; peritoneum silvery, with numerous minute round black specks. Length $2\frac{1}{2}$ inches. This species resembles *Notropis dilectus*, but has a much smaller mouth, blunter snout, and in being less silvery along the side. Known only from the Atchafalaya River, Louisiana.

Notropis louisianæ, EVERMANN MS., Atchafalaya River, Melville, Louisiana. (Type, No 48902. Coll. Fred M. Chamberlain.)

Page 348. *Anguilla chrysypa* is abundant in the Gulf of St. Lawrence, according to Dr. Wm. Wakeham.

Page 355. The original type of *Congermuræna* is *C. habenata*, Kaup, a species with blunt or granular teeth. The American species all belong to a distinct genus, *Congrellus*, Ogilby (type *balearica*), distinguished by the villiform teeth. These genera are charactered by Mr. Ogilby in a paper as yet unpublished.

Pages 356 and 357. In *Congermuræna flava* the upper jaw projects far beyond the lower. By a slip in the original description the reverse is said to be the case.

Page 359. *Muræsox conicops* is called Culevra Blanca at Mazatlan, and reaches a length of 7 feet.

Page 368. *Avocettina gilli*, Bean, should probably stand as a species distinct from *Avocettina infans*. The description in the footnote on page

368 is sufficiently full. See Jordan, Proc. Cal. Ac. Sci., ser. 2, vol. VI, 1896, 206, pl. 21.

Page 369. No. 604, *Labichthys elongatus*, is a true *Avocettina*, having the vent far behind the head. **It should stand as—**

602(a). AVOCETTINA ELONGATA (Gill & Ryder).

Page 376. After *Myrichthys tigrinus*, Girard, add:

615(a). MYRICHTHYS XYSTURUS, Jordan & Gilbert.

Teeth all more or less blunt and granular; a band of 3 or 4 series on each side of lower jaw; a band of 2 rows on each side of upper jaw; vomer with a long series divided into 2 for about ½ its length. Anterior nasal tubes conspicuous, turned downward. Eye 2½ in snout; front of eye above middle of gape, the length of which is a little more than ⅓ of head; the angle of mouth well behind eye. Interorbital width about ⅔ length of the rather long and slender snout, which projects much beyond lower jaw, the tip of the latter about reaching middle of snout. Length of head contained 4¼ times in that of trunk; head and trunk together shorter than tail, and contained 2⅔ to 2½ times in total length. Pectoral very small, its length about equal to depth of gill opening. Dorsal beginning close behind nape, much in front of gill opening; fins low; tail pointed, the tip sharp. Color light olive; sides each with 3 series of large round brown spots, those of the 2 upper series of equal size, those of lower scarcely ½ as large, faint, and often obsolete anteriorly; the spots irregular in their arrangement, those of the upper series usually twice as numerous as those of the next; those of the upper series along base of dorsal fin extending partly on the base of the fin; lower series of spots along base of anal, some of them extending on the fin or even entirely upon it; on the belly are sometimes small dark spots, scarcely arranged in series; dorsal fin with a terminal series of dark spots, which are partly confluent, the fin narrowly margined with white; anal reddish, with a lighter margin; pectoral with a blackish blotch; head covered with round black spots, which become smaller and more numerous toward the snout; lower jaw with dark spots; iris light yellow. Pacific coast of Mexico; common among the rocks about Mazatlan. (*ξύστον*, a spike; *οὐρά*, tail.)

A species distinct from *M. tigrinus*, which is known only from the original type figured by Jordan & Davis, and described in the text of Part I of this work, page 376. This specimen, said to be from "Adair Bay, Oregon," may not be American, as there is no such bay in Oregon, and no second specimen of the true *Myrichthys tigrinus* has been found anywhere.

Ophichthys xysturus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1881, 346, Mazatlan, Mexico. (Type No. 28142. Coll. Dr. Gilbert.)

615(b). MYRICHTHYS PANTOSTIGMIUS, Jordan & McGregor.

Head 3⅓ in trunk; head and trunk 1⅓ in tail; cleft of mouth 3 in head; eye 2⅔ in snout, which is 5 in head; pectorals 2 in snout; anterior nasal tube equal to the eye. Color olivaceous, with distinct rows of roundish

blackish spots, some oblong, smaller on head and covering the whole belly; 39 spots in the dorsal row, these spots usually alternating each with its fellow on the other side of dorsal, sometimes opposite; spots of second row usually opposite; spots of third row smaller and more numerous, extending from the cheeks to opposite the vent, thence running along base of anal, not running on fin, most of the spots of this row little more than ½ length of snout; 2 rows of smaller spots along belly from gill opening to front of anal; spots on nape rather large, on head larger and more numerous than in *M. xysturus;* pale color of head reduced to reticulations; chin and throat spotted as much as head; no pale centers to any of the spots; dorsal without spots or with only a few, which come up from back; from beginning to end the dorsal has a broad black margin about ⅛ height of fin; anal mostly pale, but toward tip having some black markings; pectoral with upper half jet-black, a white margin posteriorly, a small black spot in lower corner. This species is distinguishable from all others by the great number of spots of small size and without pale centers; the black edge of dorsal; the black spot on the rather large pectoral, and especially by having the belly spotted as much as the other parts. Clarion Island. One specimen, about a foot long, known. (πᾶς, whole, entire, all; στίγμα, spot.)

Myrichthys pantostigmius, Jordan & McGregor, Rept. U. S. Fish Comm. 1898, pl. 4, Clarion Island. (Type, No. 5710, L. S. Jr. Univ. Mus. Coll. R. C. McGregor.)

Page 377. After *Pisoodonophis cruentifer* add:

618(a). PISOODONOPHIS DASPILOTUS, Gilbert, new species.

Brownish above, gray below, the head and body usually thickly covered with black spots smaller than the eye; these are smaller and more numerous on the head, fewer and fainter on the lighter interior surface, and become indistinct or entirely disappear on the terminal portion of tail. In 1 specimen the head and trunk are spotted and the entire tail unicolor. In another no spots are present, the upper parts being a uniform dark brown, the under parts lighter brown, a few dark freckles only being present on sides of head. In all specimens the snout and lower jaw are blackish. The anus is near the middle of the total length, sometimes nearer the tip of snout, sometimes nearer tip of tail. The cheeks are not greatly swollen. The gape extends behind the eye, its length, measured from tip of lower jaw to angle of mouth, being contained 4⅞ to 4¾ in head. The snout projects beyond the lower jaw for a distance about equaling diameter of orbit. Eye 2 to 2⅛ in snout, 1⅞ to 2⅛ in interorbital width. Tubes of anterior nostrils about ½ diameter of eye, directed downward near tip of snout. Posterior nostrils under front of eye, concealed in the upper lip as usual. Teeth all bluntly conic, in rather wide bands on jaws and vomer; they are usually not disposed in regular series within the bands, but each band has about the width of 1 series, and these are sometimes distinguishable. The mandibular teeth become larger on approaching the symphysis, those at point of mandible and those on head of vomer being much the largest teeth present. The patch on shaft of vomer tapers

backward to a point considerably behind angle of mouth. Origin of dorsal entirely behind tip of pectorals, its distance from snout $\frac{1}{3}$ to $\frac{1}{8}$ greater than length of the head. The tip of the tail is compressed, acute, horny, used for defense. Pectoral very short, from a wide base which slightly exceeds length of gill slit. The fin rapidly narrows downward, the longest portion contained 12 to 14 times in length of head. The width of gill slit is about $\frac{1}{3}$ head.

The following table gives measurements of 4 specimens in millimeters:

Total length.	Head and trunk.	Tail.	Head.	Gape.	Eye.	Interorbital width.	Width at cheeks.	Length of snout.	Projection of snout beyond mandible.	Length of pectoral.	Basal width of pectoral.	Distance from snout to dorsal.	Vertical height of dorsal.	Depth of body.
362	177	185	38	8	3	5	$9\frac{1}{2}$	$6\frac{1}{4}$	$2\frac{3}{4}$	$3\frac{1}{2}$	5	48	3	12
401	203	198	48	$10\frac{1}{2}$	3	7	11	$7\frac{1}{2}$	$3\frac{3}{4}$	4	6	53	4	$14\frac{1}{2}$
492	248	244	52	11	$3\frac{3}{4}$	$7\frac{1}{2}$	$16\frac{1}{2}$	$8\frac{1}{2}$	$3\frac{3}{4}$	$3\frac{3}{4}$	$6\frac{1}{2}$	68	$5\frac{1}{2}$	18
494	255	239	56	12	$4\frac{1}{4}$	7	16	$8\frac{1}{2}$	$4\frac{1}{2}$	4	6	$68\frac{1}{2}$	$5\frac{1}{2}$	16

Four specimens were secured, 3 obtained in brackish water at the mouth of a small stream which empties into Panama Bay, the fourth in a fresh-water pond at Miraflores. There is some reason to suppose that they burrow in the mud.

Pisoodonophis daspilotus, GILBERT, Fishes of Panama, MS. 1898, Panama.

Page 382. *Muræna ophis*, Linnæus, is without much doubt the original *Ophichthus havannensis*. The species would therefore stand as—

626. OPHICHTHUS OPHIS (Linnæus).

Page 396. *Sidera castanea*, Jordan & Gilbert, should be removed from the synonomy of *Lycodontis funebris*. It is apparently a valid species and should be inserted as—

650(a). LYCODONTIS CASTANEUS (Jordan & Gilbert).

(MORENA PRIETA.)

Tail about as long as rest of body, or slightly longer. Head $2\frac{1}{3}$ in length of trunk; cleft of mouth wide, $2\frac{1}{3}$ to $2\frac{1}{2}$ in head. Teeth everywhere uniserial or nearly so, those on sides of mandible small, compressed, close set, subtriangular, directed backward, about 18 in number on each side; mandible with about 4 large canines anteriorly; upper jaw with the teeth partly in 2 series, some of the teeth being movable, the other mostly stronger, caninelike, especially anteriorly; front of vomer with 2 very long, slender canines, behind them a single series of small teeth; teeth all entire. Eye large, slightly nearer tip of snout than angle of mouth, its diameter 2 to $2\frac{1}{2}$ in snout; gill opening $\frac{1}{3}$ wider than orbit; tube of anterior nostril short, less than $\frac{1}{2}$ diameter of orbit; posterior nostril with-

out tube; occiput not especially elevated, the anterior profile scarcely concave (perfectly straight in young 2 feet long). Dorsal fin commencing much in advance of gill opening, becoming unusually high posteriorly, where its vertical height is more than ½ greatest depth of body; the length of the longest ray more than greatest depth of body. Color light brownish chestnut, slightly paler on abdomen; no spots or bands anywhere; fins without dark margins; no dark spot on gill opening or at angle of mouth; no black about eye; head without conspicuous pores. The specimen here described is 44 inches in length; others about 2 feet in length agree very closely. This enormous eel is very common among the rocks about Mazatlan, where it reaches a length of 6 feet. It is close to the West Indian species, *L. funebris*, but the colors are not the same, *funebris* being a greenish black, while *castaneus* is a purplish chestnut, without shades of olive or green. (*castaneus*, chestnut.)

Sidera castanea, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 647, Mazatlan. (Type, Nos. 28246, 29535 and 29591. Coll. C. H. Gilbert.)

After *Lycodontis mordax* add:

649(a). **LYCODONTIS PICTUS** (Ahl).

Head 4 in trunk; tail about as long as body; eye 2½ in snout, situated midway between snout and angle of mouth; cleft of mouth 2⅔ in head; snout 5⅔ in head; anterior nasal tube 5 in snout; gill opening 11 in head. Teeth in each jaw in a single series; palatine series either parallel with these or divergent; no distinct canines; teeth comparatively small; anterior vomerine 1 or 2 in number, bluntish and conical; posterior vomerine teeth rather blunt. Anterior nasal tubes moderate. Dorsal low anteriorly and beginning in front of gill opening. Color brownish gray or purplish, everywhere covered with small purplish black spots, which are not confluent; in the adult the spots are arranged in roundish or ring-like blotches on the sides; fins colored like body, without dark edges; young pale with black ring-shaped markings; variation in color and form of markings numerous. East Indies; everywhere common. East to offshore islands of Mexico. Two specimens, about 3 feet in length, taken at Clarion Island by Mr. R. C. McGregor. (*pictus*, painted.)

Muræna picta, AHL, le Muræna et ophichtho, VI, 6, tab. 2, f. 2; GÜNTHER, Cat. Fish, VIII, 116.
Gymnothorax pictus, BLEEKER, All. Ichth., Muræna, 87, tab. 26, 28, 29, 45.
Muræenophis pantherina, LACÉPÈDE, Hist. Nat. Poiss., V, 628, 1803.
Muræna variegata, QUOY & GAIMARD, Voy. Uranie, Zool., 246, pl. 52, f. 1.
Muræna lita, RICHARDSON, Voy. Erebus and Terror, 84, Moluccas.
Muræna siderea, RICHARDSON, Voy. Erebus and Terror, 85, pl. 48, f. 1-5, Australia.
Muræna pfeifferi, BLEEKER, Nat. Tyds. Ned. Ind., V, 173, Celebes.
Sidera pfeifferi, KAUP, Apodes, 70.

Page 401. After *Muræna argus* insert:

660(a). **MURÆNA CLEPSYDRA**, Gilbert, new species.

Closely related to *M. insularum* and *M. argus*, from the tropical Pacific, differing from both in color. Nostrils tubular, of almost equal length.

Mouth closing completely, the teeth entirely concealed by the lips. Gape straight, horizontal, extending to well behind the eyes, $2\frac{1}{8}$ to $2\frac{3}{4}$ in head. Teeth in jaws large, compressed, and wide at base, tapering uniformly to an acute point, directed backward, close set, everywhere uniserial; those in sides of mandible noticeably smaller than those of upper jaw, the teeth in both jaws increasing in size anteriorly; as many as 18 or 20 teeth may be present in the half of either jaw, but many of them are usually wanting, leaving gaps in the series; a single row of small teeth on shaft of vomer, beginning opposite front of eye; head of vomer with 2 long canines, larger than any of the other teeth, one or both of these usually wanting in larger specimens, having apparently fallen out. Head 2 ($1\frac{11}{12}$ to $2\frac{1}{12}$) in trunk; head and trunk $1\frac{1}{8}$ to $1\frac{1}{4}$ in tail; depth at anus approximately $\frac{1}{2}$ length of head; eye small, its diameter contained 12 to 16 times in head; snout 5 to $5\frac{1}{2}$. Dorsal beginning on the head, its distance from snout $1\frac{1}{8}$ to $1\frac{1}{3}$ in head. Color dark brown, lighter on belly, dull whitish on under side of head; head, body, and fins closely covered with white spots, those on posterior parts larger, with some smaller ones intermingled, the larger spots with a more or less evident central constriction which makes them hourglass shaped; toward the head the spots become very small and crowded, not more than $\frac{1}{2}$ as large as pupil; fins indistinctly light margined; a large elliptical jet-black blotch surrounds the gill slit, distinctly margined by a series of confluent white spots; the longitudinal diameter of the blotch is contained 5 to $5\frac{1}{2}$ times in the length of the head; angle of mouth with a small black blotch, often obscure, preceded by a pale spot on mandible; the throat is marked with a number of parallel lengthwise folds, the bottom of each fold with a dark line.

The following table gives measurements in millimeters of 5 specimens:

Total length.	Head and trunk.	Tail.	Head.	Gape.	Snout.	Eye.	Depth at anus.	Distance from snout to origin of dorsal.
675	311	364	106	45	$20\frac{1}{2}$	7	59	72
630	289	341	96	38	$18\frac{1}{2}$	$6\frac{1}{4}$	$52\frac{1}{2}$	70
612	287	325	98	39	19	$6\frac{1}{2}$	47	64
473	203	270	66	28	13	$5\frac{3}{4}$	40	50
397	177	220	58	21	11	$4\frac{1}{2}$	27	39

This species is abundant at Panama, where it is frequently brought to market. About 25 specimens were seen during our visit, all essentially alike in coloration. The type is 397 millimeters long (see table of measurements), and has the spots on body less numerous than in larger specimens. (*clepsydra*, κλεψύδρα, an hourglass, from its markings.)

Muræna clepsydra, GILBERT, Fishes of Panama. MS. 1898, Panama.

Page 410. It is probable that several species are confounded under the name *Elops saurus*. According to Ogilby the Australian species has only 63 vertebræ.

Page 411. 199. Genus **ALBULA**, Bloch & Schneider.

The proper binomial authority for this generic name, as well as for the names *Synodus*, *Umbra*, and *Anableps*, is Scopoli, as Dr. Gill informs us. These pre-Linnæan names, with others, were first used in binomial nomenclature as names of genera by Scopoli, Introd. His. Nat. 1777, pp. 449 (*Synodus*) and 450 (*Albula*, *Umbra*, *Anableps*). The genera should then stand as follows:

Page 411. 199. **ALBULA** (Gronow) Scopoli.

Page 533. 248. **SYNODUS** (Gronow) Scopoli.

Page 623. 298. **UMBRA** (Krämer) Scopoli.

Page 684. 312. **ANABLEPS** (Artedi) Scopoli.

Page 414. To the description of *Chanos chanos* the following may be added:

The skeletal peculiarities of *Chanos* are numerous and remarkable, many archaic characters persisting. The following account of the skeleton has been prepared by Mr. Starks:

SKELETON OF CHANOS CHANOS.

a. Cranium:

The frontals are very large, covering nearly the whole top of the head, and extending over the dorsoanterior part of the parietals, supraoccipital and the parotic process. On the side of the skull there is an area bounded by the supraoccipital, the opisthotic and the sphenotic, which is not ossified, but is composed of cartilage. Between the frontals, at about their middle, there is a place in which the bone is fibrous and largely cartilaginous; it is easily broken through. The basal cavity under the brain cavity is large. On the upper part of the operculum is a large scale like bone. The suborbitals are well developed and plate-like, extending back nearly to the posterior edge of the preopercle.

b. Vertebral column:

There are 42 vertebræ in the spinal column. The first vertebra is coossified to the skull, and apparently bears no ribs; the second vertebra supports a pair of very small, slender ribs, which articulate directly with the sides of the vertebra; the third vertebra supports the first pair of large ribs; they are articulated with the transverse processes. The first 14 or 15 neural spines and pairs of transverse processes are articulated with the vertebræ by sutures; they are easily separated from the vertebræ by boiling or maceration. The vertebræ gradually increase in size and reach their largest size about $\frac{2}{3}$ of the distance from the anterior to the posterior end of the spinal column, where they are 3 or 4 times the size of the anterior ones. This character is more marked in the adult than in the young.

c. Shoulder girdle:

The shoulder girdle is exceedingly well braced, the post-temporal is widely forked, and strongly articulated to the epiotic processes of the skull. The supraclavicle is long and slender, its posterior face is hollowed out and attached some distance from the upper end of the clavicle, which projects upward. This projecting upper end of the clavicle is braced to the skull by two long bones. The first bone is very slender, at its anterior end it is connected to the exoccipital; near its middle it is connected with the posterior end of the post-temporal, at which point it turns at a sharp angle and runs to the clavicle. The second bone is much larger; it is articulated to the basioccipital. Its posterior edge is nearly straight for its whole length, but its anterior edge is produced and much swollen near its middle, and joins the post-temporal over the first bone, then runs to the upper end of the clavicle. The inner part of the clavicle and the coracoid are thin and pierced by many holes, so that the bone in places is little more than network. The hypercoracoid has a very large foramen; at its posterior edge is a projection which supports a thin bone, probably a dermal bone. The mesocoracoid is well developed. There are 4 actinosts; the first is long, but they rapidly decrease in size to the fourth, which is short and triangular. The first ray of the pectoral is large at the basal end, and hollowed out; it works directly on the hypercoracoid.

d. Branchial apparatus:

The branchial apparatus is peculiar in the adult, in having gill rakers somewhat resembling the filaments of a feather, on both sides of each arch and on the basibranchial. They meet in a middle line between the arches and unite, forming a continuous lattice-work screen, through which nothing but the very smallest bodies can pass. The pharyngeals have no teeth, but have gill rakers similar to those on the arches; they are inclosed in sac-like projections on each side. This description is taken from the skeleton of a large specimen 4 feet long. The gill rakers are not united in young individuals.

e. Other parts:

The septæ between the myotomes are ossified about ½ an inch under the skin, forming long, slender rays of bone. There is an upper series running from the middle of the sides up on the back, and a lower series from the sides down on the belly, and form a sort of a basket around the the body. Those below have a single branch near the middle of each, the ones above have 2 branches each; these branches are lost toward the posterior end. These bones are not present in the young. The large caudal fin is attached very firmly to the hypural, the long rays of each lobe join the hypural at about the same oblique angle, the base of each ray is deeply divided and articulated immovably with the hypural. The middle short rays are all nearly horizontal and are much less firmly fastened. The first interspinal ray of the anal is hollow and cone-shaped, the posterior end of the air bladder runs into it as in the genera *Eucinostomus* and *Calamus*. The scales are very thick and closely imbricated; the skin anteriorly is ¼ inch thick. (Jordan, Fishes of Sinaloa, 404-409.)

Page 417. After *Dorosoma petenense* add:

202(a). SIGNALOSA, Evermann & Kendall.

Signalosa, EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897 (Feb. 9, 1898), 127 (*atchafalayæ*).

Body short, deep, and compressed, the form somewhat elliptical; ventral outline more strongly curved than the dorsal; head rather large, snout sharp and pointed, not tumid; mouth small, oblique, the lower jaw scarcely included; maxillary of 3 pieces, broad and curved, but without notch in the outer margin as in *Dorosoma;* caudal peduncle short and deep. Branchiostegals 5; pseudobranchiæ large; gill rakers short and very numerous, about 340 in number. No teeth; adipose eyelid present; stomach gizzard-like; scutes about 17 + 10. Last ray of dorsal very long and filamentous. This genus is allied to *Dorosoma*, from which it is plainly distinguished by the absence of the notch in the maxillary, the more pointed snout, the less-included lower jaw, the shorter anal fin, larger scales, and the fewer scutes. It differs from *Alosa* in the very numerous gill rakers, the character of the dorsal fin, and in other respects. (*signum*, a flagstaff or pole; *Alosa*, the shad; a reference to the long dorsal ray.)

679(a). SIGNALOSA ATCHAFALAYÆ, Evermann & Kendall.

Head $3\frac{3}{5}$; depth $3\frac{1}{7}$; eye $3\frac{1}{2}$ in head; snout $5\frac{1}{2}$; maxillary $3\frac{1}{6}$. D. I, 12; A. I, 24; scales 42–15; scutes 17 + 10. Body oblong-elliptical, compressed, the back in front of dorsal narrow; ventral edge sharp, serrate; head small, mouth terminal, oblique, lower jaw slightly included; snout rather pointed, not blunt, as in *Dorosoma cepedianum;* maxillary in 3 pieces, long and curved, reaching vertical at front of pupil, the outer edge not notched; no teeth. Caudal peduncle short, compressed, and deep. Origin of dorsal fin over base of ventrals, much nearer tip of snout than base of caudal, the last ray filamentous, about $\frac{1}{4}$ longer than head and nearly reaching base of caudal; the first dorsal ray about 2 in the last one; pectoral $1\frac{1}{4}$ in head, reaching base of ventrals; ventrals short, reaching only halfway to vent, their length $1\frac{1}{5}$ in pectorals; anal rays short, base of fin $1\frac{1}{5}$ in head; scutes moderate; caudal widely forked, the lower lobe the longer; scales large, thin, deciduous, somewhat crowded anteriorly; accessory scales at bases of pectorals and ventrals; base of caudal with small scales. Color bluish black or dark olivaceous on back and sides to level of the jet-black humeral spot; rest of sides and under parts bright silvery; dorsal and caudal dusky; other fins plain. The cotypes from Grand Plains Bayou are 2 females with ripe roe. They are $4\frac{1}{2}$ and $5\frac{1}{2}$ inches long, respectively, and differ from the types only in the deeper body and the much darker coloration of the upper parts.

The amount of variation in this species, shown by the material at hand, is exhibited in the following table:

No.	Head.	Depth.	Eye.	Snout.	Max.	Dorsal.	Anal.	Scutes.	Scales.	Locality.
1	$4\frac{1}{8}$	3	$3\frac{3}{4}$	$5\frac{1}{4}$	$3\frac{1}{5}$	I, 12	I, 24	16 + 11	40–15	Grand Plains
2	4	$2\frac{5}{6}$	$3\frac{1}{4}$	$5\frac{1}{3}$	$3\frac{1}{5}$	I, 12	I, 24	16 + 11	42–15	Bayou, Miss.
3	4	3	$3\frac{2}{3}$	$5\frac{1}{3}$	$3\frac{1}{3}$	I, 12	I, 24	16 + 10	42–14	
4	$3\frac{2}{3}$	$3\frac{1}{7}$	$3\frac{1}{2}$	$5\frac{1}{2}$	$3\frac{1}{5}$	I, 12	I, 24	17 + 10	42–15	Melville, La.
5	$3\frac{3}{5}$	3	4	5	$3\frac{2}{5}$	I, 12	I, 24	16 + 11	43–15	
6	$3\frac{3}{4}$	3	$3\frac{2}{3}$	$5\frac{1}{3}$	3	I, 12	I, 24	16 + 10	41–15	Grand Plains
7	$3\frac{1}{3}$	3	$3\frac{1}{4}$	5	$3\frac{1}{5}$	I, 12	I, 24	17 + 9	41–15	Bayou, Miss.
8	$3\frac{1}{2}$	3	$3\frac{1}{4}$	5	3	I, 12	I, 24	17 + 9	41–15	Black Bayou,
9	$3\frac{1}{4}$	$2\frac{5}{6}$	$3\frac{1}{3}$	$5\frac{1}{4}$	3	I, 12	I, 24	17 + 9	40–15	Miss.

This species appears to be rather common in the larger lowland streams and bayous of Louisiana and Mississippi. It probably does not reach a large size, adult examples being less than 6 inches long. It is not used as food, but is of considerable value as bait in the catfish fishery of the Atchafalaya River and its connecting lakes and bayous. Length 4 to 6 inches. (*atchafalayæ*, from the type locality.)

Signalosa atchafalayæ, Evermann & Kendall, Bull. U. S. Fish Comm. 1897 (Feb. 9, 1898), 127, pl. 7, fig. 4, Atchafalaya River, Melville, Louisiana. (Type, No. 48790. Coll. Fred M. Chamberlain.)

Page 425. The statement that *Pomolobus mediocris* does not ascend rivers to spawn is not correct. This species is known to ascend the St. Johns River, Florida, at least as far as Lake Monroe, during the winter. They usually run somewhat earlier than the shad.

Page 427. After *Alosa*, Cuvier, add:

a. Gill rakers numerous, 93 to 120; upper jaw with sharp, deep notch at tip; lower jaw not projecting. SAPIDISSIMA, 693.

aa. Gill rakers fewer than 76; notch in upper jaw smaller; lower jaw more strongly projecting. ALABAMÆ, 693(a).

Page 428. After synonymy of *Alosa sapidissima* add:

693(a). **ALOSA ALABAMÆ**, Jordan & Evermann.

(Alabama Shad; Gulf Shad.)

Head $4\frac{3}{5}$; depth 3; snout $4\frac{1}{2}$; eye $4\frac{1}{8}$; maxillary $2\frac{1}{8}$. D. 15; A. 20; scales 55, —16 in a crosswise series; scutes 21 + 15; vertebræ 54; gill rakers 56 to 68. Body deep; back gently and evenly arched from tip of snout to origin of dorsal fin, thence descending in a regular curve to base of caudal fin; ventral outline nearly straight from tip of mandible to ventrals, and also from there to base of caudal. Head small, snout pointed; upper lip with a small notch, into which fits the tip of the slightly projecting lower jaw; maxillary narrow; cheek much deeper than long; teeth on tongue and maxillary scarcely perceptible. Origin of dorsal nearer snout than base of caudal, the fin low, the longest ray shorter than the base, or about equal to snout and eye; base of anal somewhat greater

than that of dorsal, or equal to length of pectoral. Gill rakers 68, the longest about equal to length of snout. Peritoneum pale. Color as in *Alosa sapidissima;* the caudal, dorsal, and pectoral fins rather darker tipped. The male differs from the female only in being somewhat more slender. This species differs from *Alosa sapidissima* chiefly in the fewer gill rakers, its sharper, more pointed snout, smaller notch in upper jaw, more projecting mandible, and more slender maxillary. It seems to reach maturity at a much smaller size than the common shad. Streams tributary to the Gulf of Mexico; known from Tuscaloosa, Alabama, and Pensacola, Florida.

Alosa alabamæ, JORDAN & EVERMANN, in EVERMANN, Rept. U. S. Fish Comm. 1895 (Dec. 28, 1896), 203, Black Warrior River, Tuscaloosa, Alabama (Type, Nos. 47689 and 47690. Coll. J. H. Fitts); EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897, 127, pl. 7, figs. 5 and 6.

According to Ogilby, *Kowala* is a genus distinct from *Sardinella.*

Page 436. *Ilisha panamensis* is not separable from *I. furthi.* The latter name has priority.

Page 437. *Opisthopterus lutipinnis* is very abundant on the outer sand beaches about Mazatlan.

Page 445. Species 728, *Stolephorus poeyi,* is a species of *Lycengraulis,* and should stand as—

743(a). LYCENGRAULIS POEYI (Kner & Steindachner).

Numerous specimens lately taken by Dr. Gilbert at Panama. A large species used as food. The teeth are unequal in *Lycengraulis,* but none of them can be properly described as canine-like.

Page 447. After *Stolephorus lucidus* add:

732(a). STOLEPHORUS RASTRALIS, Gilbert & Pierson, new species.

Head 3.16 (3.1 to 3.3); depth 3.8 (3.5 to 4.2); eye 3.4 in head (3.33 to 4). D. 14 (12 to 15); A. 26 to 32. Body much compressed and deep; belly sharply keeled in front of ventrals; dorsal outline much less curved than ventral, the lower profile rising very rapidly from a point opposite middle of pectorals to tip of snout, in shape of head thus closely resembling the species of *Cetengraulis.* Maxillary reaching almost but not quite to gill opening; snout high, compressed, its length ½ to ⅔ diameter of eye. Gill rakers averaging in larger examples 51 + 64, in smaller specimens 44 + 50; the largest about as long as eye. Insertion of dorsal fin variable, but never posterior to a point midway between base of caudal and middle of eye; pectoral fins reaching to or nearly to insertion of ventrals, the latter not to vent. Color olivaceous, the lower part of side with violet reflections; sides of head silvery; a conspicuous silvery lateral band varying in width from about 1⅓ times length of orbit in the largest examples to less than ½ orbit in the smaller specimens; the band is widest before dorsal, tapering to ½ or less than ½ its greatest width on caudal peduncle, where it frequently disappears in the young. In larger specimens the ventral edge of this band is frequently ill-defined anteriorly; top of head with

widely spaced black specks; a dark vertebral streak, more or less of it often consisting of 2 narrow lines; tips of caudal lobes often blackish; fins otherwise unmarked. This species differs from closely allied species in the following characters: From *Stolephorus lucidus*, in the much longer head, more compressed body, well-defined lateral stripe, and smaller eye; from *S. compressus*, in the longer head and wider lateral band; from *S. panamensis* and *S. mundeolus*, in the much more numerous gill rakers, and the more anterior position of the dorsal relative to the anal, the origin of the anal being under the middle of the dorsal, while in *S. panamensis* the origins of the two fins lie in the same vertical. Length 2 to 3 inches. Panama. Many specimens. (Gilbert & Pierson.) (*rastrum*, a rake, from the long gill rakers.)

Stolephorus rastralis, Gilbert & Pierson, Fishes of Panama, MS. 1898, Panama.

732(b). STOLEPHORUS MUNDEOLUS, Gilbert & Pierson, new species.

Head 4.15 (4 to 4.25); depth 3.77 (3.40 to 4.25); eye 3.44 in head (3.12 to 3.70). D. 13 or 14; A. 33 (33 to 35); scales 36 (35 to 39). Dorsal and ventral contours about equally and gradually rounded from the middle region of body to the tip of snout and base of caudal fin. Snout short, high, compressed, blunt at tip, its length 1.8 in eye. Eye very large. Maxillary broad, tapering to a sharp point, which reaches margin of gill opening. Gill rakers 17 to 22 + 21 to 24; the longest 1.5 to 2 in eye. Anterior insertion of dorsal fin varying from a point midway between base of caudal and middle of eye, to a point midway between the caudal and tip of snout. In 10 examples its insertion is before that of the anal. Anal fin long, averaging 33 rays, its origin beneath the anterior third of the dorsal; length of base shorter than in *S. panamensis*, being 3.04 in length, while in the latter its length is contained 2.5 in length. Pectoral long, reaching well beyond the insertion of the ventrals, equaling length of head behind front of pupil; a large axillary scale; ventrals scarcely reaching vent. Uniform light olive, with silvery reflections; a faint, narrow, silvery-gray lateral stripe, sometimes scarcely distinguishable; sides of head plain silvery; upper margin of orbital rim black; dorsal region blackish; a faint, narrow dark line on each side of the light middorsal streak; caudal slightly dusky; fins otherwise unmarked. This species is closely allied to *Stolephorus panamensis* and *S. compressus*, but may be distinguished from the former by its longer head, larger eye, greater depth, fewer scales along the lateral line, and its much shorter anal base; also by the much fainter lateral silvery stripe. The eye is contained 14 to 16 times in length, excluding the caudal, while in *S. panamensis* the length contains the eye 16 to 20 times. From *S. compressus* it differs in the relative length of the head and maxillary. In *S. mundeolus* the maxillary is contained in the head 1.27 times (1.19 to 1.37); in *compressus* 1.48 times (1.30 to 1.81). In *mundeolus* the head is contained 4.15 times in the length; in *compressus* 4.44 times. Length 4 to 6½ inches. Panama; many specimens. (Gilbert.) (*mundeolus*, somewhat shining, from *mundus*, neat or clean.)

Stolephorus mundeolus, Gilbert & Pierson, Fishes of Panama, MS. 1898, Panama.

732(c). STOLEPHORUS NASO, Gilbert & Pierson, new species.

Head 3.3 to 3.5; depth 4.7 to 5.8; eye 4½ to 5 in head. D. 14 or 15; A. 22 to 24; lateral line about 35 (?). Dorsal and ventral outlines weakly arched; body slender, its greatest depth 1.5 in head, compressed; belly carinated in front of ventrals, and sometimes behind them in larger specimens. Head long and slender, its greatest width 1.5 to 1.7 in its length, the lower profile much more oblique than the upper. Snout long, compressed, bluntly rounded, its length exceeding the small eye. Cheek with a very acute posterior angle. Opercle narrow, oblique. Maxillary rather bluntly pointed, failing to reach gill opening by about ½ diameter of pupil. Teeth on the maxillary quite prominent and directed forward. Gill rakers short, 17 + 20 in number, the longest 1.5 in eye. Scales large, thin, deciduous, only a few scattering ones remaining in our specimens. Dorsal fin inserted midway between front or middle of orbit and base of median caudal rays. Origin of anal under or slightly behind middle of dorsal; length of anal base about equal to the distance from front of orbit to base of ventral fin; pectorals not reaching ventrals, their length about ⅓ length of head. Length of ventrals equaling or slightly exceeding distance from tip of snout to middle of pupil. Color light olive, with the usual bright reflections; a large dark patch of brown dots on occiput; a double series of dots along median line posterior to dorsal, this absent in some specimens; large specimens with a bright, well-defined silvery streak, slightly narrowing anteriorly and on caudal peduncle, its greatest width about equaling diameter of eye; in the young, this band is fainter and narrower; a conspicuous series of black dots at base of anal. Characterized by the slender form, well-defined silvery streak, sharply carinated breast, the small eye, and the very long, compressed, deep, and rather bluntly rounded snout. Most closely resembling *S. starksi*, from which it differs in the smaller eye, longer snout, and slightly longer anal. Length 2 to 2¼ inches. Panama; common. (Gilbert & Pierson.) (*naso*, long-nosed.)

Stolephorus naso, Gilbert & Pierson, Fishes of Panama, MS. 1898, Panama.

732(d). STOLEPHORUS STARKSI, Gilbert & Pierson, new species.

Head 3.3 to 3.6; depth 4.8 to 5.5 in length, 1⅓ in head; eye 3 to 3.5 in head. D. 15 or 16; A. 17 to 22; scales about 41. Body long and slender, slightly deeper and more compressed than in *S. ischanus*, which much resembles this species. Dorsal outline very little arched; ventral outline nearly straight from gill opening to insertion of anal fin, the lower profile of head oblique, nearly straight. Belly compressed, keeled for anterior ⅔ of distance anterior to base of ventrals. Head long and pointed, its width 1½ times in its length; maxillary abruptly widened at the mandibular joint, tapering posteriorly to a blunt point, which reaches almost to the gill opening, its length equal to length of base of anal; snout long, sharp, and projecting, abruptly compressed in its terminal portion as seen from above, its length ⅔ diameter of orbit, or slightly more. Branchiostegal membranes united at base for a very short distance. In 4 exam

ples examined as to this point, the gill rakers are as follows: 20+25, 23+24, 21+23, 19+30, the longest contained 1½ to 1⅔ in eye. Scales large, thin, and deciduous, a few only remaining on the specimens at hand. Origin of the dorsal fin equally distant from the base of the caudal fin and tip of snout or front of eye. Anal inserted under beginning of posterior third of base of dorsal; pectorals not reaching ventrals, the latter ⅔ distance to front of anal. Color light olive, with broad, well-defined lateral silvery streak of nearly uniform width, usually narrowing anteriorly and on middle of caudal peduncle, its width in our largest specimen ⅜ diameter of eye; the silvery streak has a slight golden tinge; a narrow dark vertebral line, which widens on the nape; occiput blackish. Vertebræ 40, counted in 1 example only. This species differs from *Stolephorus cultratus* in its slenderer body, shorter snout, wider opercle and smaller teeth; the belly is also not sharply carinate, the dorsal is more anteriorly placed, the ventrals are farther back, and the silvery streak is wider anteriorly. It differs from *S. delicatissimus* in its longer, slenderer head and body, smaller eye, longer, sharper snout, and much wider, better-defined silvery streak. Length 1½ to 2¼ inches. Panama; common. (Gilbert & Pierson.) (Named for Edwin Chapin Starks.)

Stolephorus starksi, Gilbert & Pierson, Fishes of Panama, MS. 1898, Panama.

Page 448. After *Stolephorus spinifer* add:

737(a). STOLEPHORUS SCOFIELDI, Jordan & Culver.

Head 3¾ to 3$\frac{7}{11}$ in length to base of caudal; depth 4¼ to 5; eye 3¾ to 4 in head. D. 12; A. 25 or 26; scales 41 or 42. Body somewhat compressed and elevated, the belly not carinated nor serrated. Teeth in both jaws, and on palatines, a few on vomer; maxillary covered with teeth its entire length and reaching beyond base of mandible, but not to opercular margin. Gill rakers 10+12, the longest a little more than ½ the eye. Origin of dorsal midway between base of median caudal rays and center of eye; anal not quite as long as head, its origin below the middle of dorsal; lower caudal lobe longer than upper; longest ray equaling length of the head; shortest caudal ray 2½ in longest. Pectorals not reaching ventrals, 1¾ in head. Both anal and dorsal fins preceded by a rudimentary spine, not ½ length of first true ray. Color translucent, with a distinct broad silvery stripe as wide as the eye, growing more diffuse at lower anterior edge, narrowing on caudal peduncle, and becoming fan-shaped on the base of caudal; tip of snout black; a distinct median band of black specks extending from tip of snout to base of caudal; no distinct black markings on fins. Close to *Stolephorus delicatissimus*, but with larger head, wider lateral band, and greater number of dorsal and anal rays. Length 3 inches. Found in the Astillero at Mazatlan; not very abundant. (Named for Mr. Norman Bishop Scofield, a member of the Hopkins expedition to Sinaloa.)

Stolephorus scofieldi, Jordan & Culver, Fishes of Sinaloa, 410, 1895, Mazatlan, Mexico. (Type, No. 2941, L. S. Jr. Univ. Mus. Coll. Hopkins Exped. to Sinaloa.)

737(b). STOLEPHORUS ASTILBE, Jordan & Rutter.

Head $4\frac{1}{3}$ in length; depth $4\frac{1}{2}$ to 5. D. 12; A. 19 to 22; eye $3\frac{1}{2}$ in head; pectoral $1\frac{4}{5}$; base of anal $1\frac{1}{6}$. Body rather elongate, not greatly compressed; edge of belly moderately sharp; head sharp; snout projecting beyond lower jaw, shorter than diameter of eye; tip of lower jaw reaching a little past anterior edge of orbit; maxillary reaching gill opening, its end tapering to a sharp point; eye longer than snout, nearly 2 in postorbital part of head; gill rakers $\frac{2}{3}$ eye; a slight keel on top of head. Origin of dorsal midway between base of caudal and eye; scales caducous. Translucent, head silvery; sides without lateral band; a dark spot on top of head; back with black points. This species is similar to *Stolephorus brownii*, but more slender, head shorter, and lateral silvery stripe wanting. Length 3 inches. Jamaica. Numerous specimens obtained. (*ἄ*, not; *στίλβη*, shining.)

Stolephorus astilbe, JORDAN & RUTTER, Proc. Ac. Nat. Sci. Phila. 1897, 95, Jamaica. (Type, No. 4854, L. S. Jr. Univ. Mus. Coll. Joseph Seed Roberts.)

737(c). STOLEPHORUS ROBERTSI, Jordan & Rutter.

Head 3 in length; depth 4. D. 14; A. 23; scales about 35; eye 4 in head; pectoral $2\frac{1}{2}$; base of anal $1\frac{2}{3}$; caudal $1\frac{1}{2}$. Body deep, strongly compressed, abdomen compressed to an edge, head large, compressed, the snout rather sharp, projecting beyond lower jaw, a little shorter than eye; cheek triangular; opercle large; distance from lower angle of cheek to edge of opercle equal to distance from same point to posterior edge of eye; maxillary short, not reaching root of mandible, its end rounded; lower jaw not reaching beyond anterior edge of orbit; gill rakers longer than eye, as long as orbit. Origin of dorsal midway between base of caudal and front of eye; scales caducous. Color translucent; head silvery, punctulate above; a silvery lateral band nearly as broad as eye; caudal with dark points, other fins colorless. This species seems to be related to *Stolephorus opercularis*, but the lateral band is distinct and the opercle is shorter. Jamaica; only the type, 2 inches long, known. (Named for Rev. Joseph Seed Roberts, of Kingston, Jamaica, who collected the type specimen.)

Stolephorus robertsi, JORDAN & RUTTER, Proc. Ac. Nat. Sci. Phila. 1897, 95, Jamaica. (Type, No. 4853, L. S. Jr. Univ. Mus.)

Page 449. *Anchoria* can not be maintained as a distinct genus. The name must be placed as a synonym of *Stolephorus*.

Page 450. Add:

741(a). CETENGRAULIS ENGYMEN, Gilbert & Pierson, new species.

Head 3 to 3.3; depth 4 to 4.9; eye 4 in head. D. 14 or 15; A. 20 to 23; B. 7 (9); vertebræ 41. Body compressed, fusiform, not so deep as in *C. mysticetus* or *C. edentulus*. The dorsal and ventral outlines being about equally and regularly curved in the larger specimens; in the smaller specimens the ventral contour is more nearly straight. Belly trenchant, but

not carinate nor serrate; caudal peduncle moderate, its depth being contained 1.5 times in its length. Head similar to *C. mysticetus;* the snout longer, contained 5.5 to 7 times in head, 1½ times in eye (the snout contained 8 to 9 times in head in *C. mysticetus*). Both jaws bear minute teeth, those on the maxillary largest. Branchiostegal membranes united for only ⅜ to ⅝ of the distance between tip of mandible and mandibular articulation, wholly free from the isthmus. Tip of mandible directly beneath the anterior border of orbit. Gill rakers long, $\frac{9}{10}$ diameter of eye, 20 to 30 on the upper limb, 25 to 30 on the lower limb; in 5 examples as follows: 25 + 30, 27 + 25, 30 + 26, (23 + 29 to 20 + 25), 25 + 30. The origin of the dorsal is midway between base of median caudal rays and a point varying between front and middle of the eye. Insertion of anal below the posterior fourth or third of the dorsal, its length equaling the distance from the posterior border of the eye to insertion of pectoral. The pectoral is short, 2¼ to 2⅓ in head, failing to reach the insertion of the ventrals by ½ or nearly ⅓ its length. Caudal deeply forked, its median rays 2½ to 3 times in head. Color uniformly silvery with a distinct, well-defined lateral silvery band extending from upper angle of gill opening to base of caudal, its greatest width equaling the diameter of orbit, becoming narrower on caudal peduncle. This species differs from *C. mysticetus* in the much narrower union of the gill membranes, the less numerous gill rakers, and in the longer snout. Length 1½ to 2⅓ inches. Panama Bay. Not rare. (Gilbert & Pierson.) (*ἐγγύς*, near; *δμήν*, membrane.)

Page 451. *Lycengraulis* has the teeth large and somewhat unequal, but none of them is properly described as "canine-like."

Page 459. Add:

229(a). ERICARA, Gill & Townsend.

Ericara, Gill and Townsend, Proc. Biol. Soc. Wash., xi, 1897 (Sept. 17, 1897), 232 (*salmonea*).

Alepocephalids with small, perfectly smooth, imbricated cycloid scales, wide cranium, projecting snout, deeply cleft mouth, uniserial and acrodont teeth on vomer and anterior portion of palatines, and dorsal and anal of normal extent and opposite each other. Bering Sea. (*ἐρί*, an intensive particle; *καρά*, head.)

753(a). ERICARA SALMONEA, Gill & Townsend.

D. 17; A. 24. Maxillary extending to vertical of posterior border of orbit; head large; length 8½; depth 5; width 4½. Bering Sea; only the type known, a large example in good condition.

Ericara salmonea, Gill & Townsend, Proc. Biol. Soc. Wash., xi, 1897 (Sept. 17, 1897), 232, Bering Sea, southwest of Pribilof Islands, at Albatross Station 3603, in 1,771 fathoms. (Type, No. 48769, U. S. Nat. Mus. Coll. *Albatross.*)

Page 465. Dr. G. A. Boulenger has kindly sent us the following note regarding the types of *Coregonus richardsonii* which are in the British Museum:

I have examined the types (dry) of *Coregonus richardsonii*. There are about 20 gill rakers on the lower part of the anterior arch, the longest ½ the diameter of the eye. The maxillary extends to below anterior border of eye, and its length is 4 times in length of

head as stated by Günther, therefore a little shorter than in *C. clupeiformis*. Tongue with 4 series of teeth, as in *C. labradoricus*. It seems to agree best with *C. nelsoni* (description), but has fewer scales in lateral line. In short, I can not identify *C. richardsonii* with any of the forms known to me.

Page 471. After *Argyrosomus laurettæ* add:

768(a). ARGYROSOMUS ALASCANUS, Scofield, new species.

Head $4\frac{1}{4}$; depth about 4. D. 12; A. 12; scales 10–85–9. Eye a little shorter than snout, 5 in head, $1\frac{1}{3}$ in interorbital space. Head wedge-shaped, the upper and lower profiles straight and meeting with a sharp angle at the snout. Viewed from above the snout is blunt, almost square, with the narrow, pale rounded tip of the lower jaw slightly projecting. Mouth oblique; the distance from tip of snout to tip of maxillary is equal to the distance from tip of snout to center of pupil; the maxillary from its anterior articulation is contained $3\frac{1}{2}$ in the head, its width 3 in its length, its upper anterior edge closing under the preorbital; mandible $2\frac{1}{3}$ in head, its articulation with the quadrate bone beneath the posterior edge of the eye; width of supplemental bone a little more than $\frac{1}{2}$ width of maxillary. Preorbital broad, its greatest width equaling $\frac{3}{5}$ its length or diameter of pupil; width of supraorbital equals $\frac{2}{7}$ its length. Gill rakers 12 to 14 + 21 to 23, long and slender, the longest $\frac{2}{3}$ diameter of the eye. The tongue, vomer, and palatines without teeth. Distance from tip of snout to nape equaling $\frac{1}{2}$ the distance from nape to the front of the dorsal or $\frac{2}{3}$ length of head. Adipose fin large; ventral scale $\frac{1}{2}$ length of fin; longest dorsal ray $1\frac{1}{2}$ in head; longest anal ray 2 in head; the pectorals reach more than $\frac{1}{2}$ to the ventrals; the ventrals reach $\frac{2}{3}$ to vent; the caudal is forked for a little more than $\frac{1}{2}$ its length. Color dusky above, silvery beneath; the dorsal, adipose fin, tips of caudal rays and upper side of anterior pectoral rays dusky; the rest pale. This species appears nearest related to *Argyrosomus artedi*, from which it differs chiefly in the number of gill rakers.* Length about a foot. Northern Alaska near Bering Straits; 3 specimens known, 1 from salt water at Point Hope, the others from fresh water at Grantley Harbor.

Argyrosomus alascanus, Scofield, in Jordan, Rept. U. S. Fur Seal Investigations, 1898, Point Hope and Grantley Harbor, Alaska. (Coll. Scofield & Seale.)

Page 482. Beginning with line 10 from the bottom, the statement that the small form of the redfish has been traced from the mouth of the Columbia to Wallowa Lake is not true. The remark was meant to apply to the large form. The question as to whether the small form descends to the sea is still unsettled.

*The fin formulæ, etc., of these 3 specimens are as follows:

Locality.	Length.	Gill rakers.		Dorsal.	Anal.	Scales.
Grantley Harbor	$8\frac{1}{8}$ in.	14 + 23	14 + 22	12	14	88
Do	9 in.	14 + 22	12 + 22	12	12	87
Point Hope, Alaska	11 in.	13 + 21	12 + 21	12	12	85

Page 492:

779. SALMO MYKISS, Walbaum.

(MYKISS; SOMKA; KAMCHATKA SALMON TROUT.)

By an unfortunate error the writers have heretofore used the name *Salmo mykiss* for the Cutthroat Trout of the Northwest. It was known that the Cutthroat is the only true or black-spotted trout in Alaska, and it was assumed that its range extended along the coast to all streams in Bering Sea. But our recent explorations have shown that it probably does not occur in Bering Sea, nor is there any undoubted record to the north of Wrangell. If it reaches Kadiak or Sitka or Prince William Sound, it is only rarely, and the streams of the Aleutian Islands and the east coast of Bering Sea contain no species of *Salmo*. The name *Salmo mykiss* must, therefore, be restricted to the Kamchatkan species, while the species of the American rivers heretofore called *Salmo mykiss* must stand as *Salmo clarkii*. We have, therefore, studied with great interest a specimen of the genuine *Salmo mykiss*, the first on record since the times of Pallas, Krasheninnikof, and Steller. The specimen, an adult male, 960 mm. long, was taken by Dr. Leonhard Stejneger, September 15, 1897, in the Kalakhtyrka River, near Petropaulski, Kamchatka. It was called "Sonka" or "Somka" by the natives. It is said to occur rarely and to be found in but few rivers, the Kalakhtyrka among them. It is considered to be superior as food to other *Salmonidæ*, except the King Salmon (*O. tschawytscha*). Head 4 in length; depth $4\frac{1}{4}$. D. 11. A. 10 (developed rays); scales 125–24. Mouth large, the maxillary $1\frac{1}{6}$ in head, being somewhat produced at tip; vomerine teeth few, evidently deciduous, only 3 being present. Eye $8\frac{1}{2}$ in head; snout $2\frac{3}{8}$. Pectoral 2 in head, longest anal ray $2\frac{2}{7}$. Anal fin high and somewhat falcate; ventrals inserted under anterior third of dorsal, reaching about halfway to vent; adipose fin over posterior end of anal; caudal lunate. Color dark grayish above, sides silvery; a few small, faint, round black spots on back and on top of head, these sparse and obscure; a few faint spots on base of dorsal, and some on adipose dorsal; spots on caudal small, but distinct, especially in middle of fin; no trace of red at throat, in example preserved in formalin, and doubtless none in life. The specimen is now a half skin, in good condition.

The following measurements were taken from the fresh specimen by Dr. Stejneger:

	Millimeters.
Total length	960
Total length without caudal	853
Head	215
Tip of nose to anterior end of dorsal	400
Length of base of dorsal	100
Posterior end of dorsal to anterior end of adipose fin	167
Length of base of adipose fin	17
Posterior end of adipose to caudal	81
Posterior end of anal to caudal	81
Length of base of anal	71
Anterior end of anal to posterior end of ventral	165
Height of body in front of dorsal	195
Height of body at posterior end of adipose and anal fins	105

Millimeters.

Height of body at beginning of caudal 77

Ventrals under anterior third of dorsal; adipose fin over posterior end of anal; ventrals reach about ½ distance to vent; 24 scales in transverse series from origin of dorsal to lateral line; 125 scales in lateral line. Color silvery gray on back, black spots obsolete.

This species is evidently a close ally of the Atlantic salmon, belonging to the restricted subgenus *Salmo*. From *Salmo salar* it differs in the slightly larger mouth and rather different coloration and in very little else.

Mykiss, PENNANT, Arctic Zool., Intro., 126, 1792, Kamchatka; after KRASHENINNIKOF, etc.
Salmo mykiss, WALBAUM, Artedi Piscium, 59, 1792, Kamchatka; based on *Mykiss* of PENNANT.
Salmo penshinensis, PALLAS, Zool. Rosso-Asiat., III, 1811, Gulf of Penshin.
Salmo purpuratus, PALLAS, Zool. Rosso-Asiat., III, 374, 1811, Bering Sea.

The correct names of the American Cutthroat Trout and its numerous known varieties are the following:

780. SALMO CLARKII (Richardson).
780(a). SALMO CLARKII LEWISI (Girard).
780(b). SALMO CLARKII GIBBSII (Suckley).
780(c). SALMO CLARKII HENSHAWI (Gill & Jordan).
780(d). SALMO CLARKII VIRGINALIS (Girard).
780(e). SALMO CLARKII SPILURUS (Cope).
780(f). SALMO CLARKII PLEURITICUS (Cope).
780(g). SALMO CLARKII BOUVIERI (Bendire).
780(h). SALMO CLARKII STOMIAS (Cope).
780(i). SALMO CLARKII MACDONALDI, Jordan & Evermann.

Page 500. Before *Salmo irideus* insert the following:

781(b). SALMO GAIRDNERI BEARDSLEEI, Jordan & Seale.

(BLUEBACK TROUT OF LAKE CRESCENT.)

Head 3¼ in length to base of caudal; depth about 4; eye 4⅔ in head, 1⅔ in snout; scales 24–130–20, 130 cross series, those in front of dorsal numerous, about 70 if counted along median line, 60 if the rows along upper side are counted; dorsal with 10 branched rays; anal with 11 branched rays; branchiostegals 11; gill rakers 8 + 13, rather long and slender, the longest nearly ⅓ in length, 7 to 9 in maxillary. Head pointed; mouth rather large; maxillary extending to hinder margin of eye, 1⅓ in head, with about 20 teeth; snout 3⅔ in head; preorbital very narrow, the maxillary almost touching the orbit; posterior suborbitals shorter than eye, about 6 in head; opercle not very broad, equal to eye, its free part 4⅔ in head; interorbital width 3⅔ in head, equal to snout; several large teeth along margin of tongue; no hyoid teeth; teeth on vomer in zigzag series. Origin of dorsal in middle of the length, margin slightly concave, the first ray 1⅘ times the last, the last ray being pointed, slightly greater than base, 2 1/10 in head. Origin of anal midway between origin of dorsal and base of caudal, margin straight, the tip of the last ray slightly exserted; anterior rays 3¼ times posterior, and equal to base of fin, 2¼ in head. Adi-

pose fin high and slender, situated above or anterior to end of anal. Pectorals $1\frac{1}{2}$ in head; ventrals under middle of dorsal, $2\frac{1}{3}$ in head. Caudal broad, nearly truncate, the middle portions abruptly lunate when spread open, with pointed angles, each lobe being somewhat convex on its edge; longest rays $1\frac{1}{3}$ in head. Least depth of caudal peduncle $2\frac{3}{4}$ in head. Pyloric cæca 50 to 60, short and thick, the longest about 3 in head. Color in spirits very dark blue above, sides abruptly brighter, with many scales abruptly silvery; below white, lower jaw white, its margin dusky; cheeks below suborbitals very dark; sides, top of head, dorsal, and caudal fins spotted, the spots all very small; pectorals and ventrals nearly colorless, without spots, and slightly dusky; adipose fin with 2 spots; tips of lower fins faintly tinged with yellowish. Two specimens, each 16 inches long, Nos. 1861 and 1862, L. S. Jr. Univ. Mus. They were taken on March 12 and 16, 1896, in Lake Crescent, by Mrs. George E. Mitchell, of Fairholme, and sent to us by Mr. M. J. Carrigan, of Port Angeles.

A third specimen shows the following characters: Head $3\frac{5}{6}$; depth $3\frac{3}{8}$. D. 12; A. 12 branched rays; branchiostegals 11 or 12; scales 23–123–26, 64 before dorsal; snout $2\frac{5}{6}$; eye $7\frac{3}{8}$; maxillary $1\frac{3}{8}$ in head, its depth 8 in its length. Body robust, little compressed; head large; maxillary moderate, extending beyond eye; opercle moderate, its width $5\frac{3}{8}$ in head. Last ray of dorsal pointed. Caudal subtruncate, lunate mesially, each lobe somewhat convex, pointed at tip. Caudal peduncle short and thick. Series of vomerine teeth long, in double row. Color above dark green, with black spots, which are small and sparse on body, extending to below lateral line; many small spots on head, dorsal, and caudal; spots not more numerous behind than before; sides and belly bright silvery; no red on lower jaw; a faint pink shade along lateral line; pectorals colorless except the upper ray; ventrals and anal colorless; flesh pale; gill rakers removed. This specimen, male, was taken in Lake Crescent. Length $26\frac{1}{2}$ inches; weight in life 14 pounds. This specimen differs from a large *gairdneri* most in the large scales. In addition the head is much larger, and the body deeper.

A fourth, still larger, specimen (No. 1865, L. S. Jr. Univ.), an old spent male, 27 inches long, has been still later received. It shows the following characters: D. 11; A. 12. Head $3\frac{3}{8}$ in length; gill rakers $8+12$, of medium size, rather broad but sharp pointed; opercle $3\frac{1}{4}$ in head; eye 7 in head; branchiostegals 11; maxillary long, reaching beyond eye, $1\frac{3}{8}$ in head, its width $9\frac{1}{2}$ in length. A double row of sharp teeth extending to within a short distance of end, where they are replaced by a single row of slightly larger teeth; teeth on tongue rather large; no teeth on hyoid; teeth on vomer in zigzag series. Scales 137–26. This specimen, a spent male, has the flabby muscles and slimy, half-concealed scales of the spent male salmon. The dark dots are very numerous and small and show very distinctly on back and sides, as also on head and fins; there is a dull red lateral band on head and body—this is about an inch broad, its outlines diffuse; a black blotch on cheek; maxillary dusky with a red blotch toward its tip; lower jaw and branchiostegals dusky; pectoral, ventral, and anal dark; back dark green, belly dusky.

The following account of the life coloration of *Salmo beardslei* is given by Mr. George E. Mitchell:

The Blueback Trout caught in Lake Crescent are on the back a deep dark-blue ultramarine color of a peculiar transparency, dotted with small round black spots from the size of a pin's head to a little larger. The 2 fins on the top of the back are a dark smoky color, also dotted as on back end, and are transparent. The tail is the color and transparency with dots also—same as the top fins. The side fins and the bottom fins are dead white and sometimes faintly tinged with a pinkish hue at the edges; the belly is white. Looking at the fish sideways the sides of the fish show the scales to be iridescent, the red flash predominating. The head has very much the polish of mother-of-pearl around the lower jaws and jowls, red and pale-blue colors predominating; under the eyes a few black spots; on top of head the blue much darker than on top of back—so dark, in fact, that the black spots on it look blacker than the rest. The nearer the shore these fish are caught the lighter the blue on back, the fish often having an impression of the surroundings distinctly marked on them.

The following notes are added by Admiral Beardslee:

HABITS.

The Blueback is a deep-water dweller; those taken by me in late October were caught at depths varying from 30 to 50 feet, on large spoons. They fought hard until brought near the surface, then gave up, and when landed were found puffed up with air. Specimens taken in spring and put in pools in mountain streams with other trout died very soon, while the others lived. The trout caught by Mr. Mitchell, in March, was taken near bottom, by a large spoon, and it is not on record that at so early a date one has previously been caught.

FLESH.

Light lemon color before cooking; devoid of the oily salmon flavor, and very excellent; whitening by cooking.

OVA.

October 28. The eggs in the large fish were in *individual* size, and in size of cluster much smaller than those of a salmon of the same size.

The following extracts from a letter from Mr. Carrigan, dated Port Angeles, April 30, are of much interest:

* * * Answering your direct inquiries: The Beardslees and Crescents are readily distinguishable, and can always be told apart. There are no red spots at the points indicated on the Crescent trout—no markings to suggest the Cutthroat trout.

(Named for Admiral L. A. Beardslee, U. S. N., in recognition of his active and intelligent interest in American game fishes.)

Salmo gairdneri beardsleei, JORDAN & SEALE, Proc. Cal. Ac. Sci., ser. 2, vol. VI, 1896, 209, pl. 23, Crescent Lake, Clallam County, Washington. (Coll. Mrs. George E. Mitchell. Type, No. 1864, L. S. Jr. Univ. Mus.)

780(c). SALMO GAIRDNERI CRESCENTIS, Jordan & Beardslee.

(SPECKLED TROUT OR LAKE CRESCENT.)

Head $3\frac{4}{5}$ in length to base of caudal; depth 5; exposed portion of eye 6 in head, $1\frac{3}{8}$ in snout; scales 32–151–34, 151 cross series, 83 in front of dorsal; dorsal with 10 branched rays, anal with 11; branchiostegals 10; gill rakers 6 + 11, counting rudiments, these very short and thick, the longest but $\frac{3}{16}$ inch in length, $18\frac{1}{2}$ in maxillary; mouth large, maxillary extending much beyond eye, $1\frac{3}{8}$ in head, with about 20 teeth; tongue with

the usual teeth; teeth on vomer in zigzag series; hyoid region of tongue without teeth. Snout 3½ in head; preorbital very narrow, not so wide as maxillary adjacent to it; the posterior suborbitals longer than eye, 5⅓ in head; opercle and subopercle very narrow, scarcely as wide as eye, the free part of opercle 6⅓ in head; interorbital width 4¼ in head. Origin of dorsal in middle of length of body, its margin straight, anterior 2½ times posterior, and slightly longer than base, 2½ in head; last ray of dorsal pointed. Origin of anal midway between origin of dorsal and base of caudal, margin irregular, anterior rays 3 times length of posterior and equal to base of fin, 2⅔ in head. Adipose fin high and slender, situated immediately behind anal; pectoral 1⅘ in head; ventrals under middle of dorsal, 2⅔ in head; caudal broad, slightly emarginate, nearly truncate when spread, its corners not rounded, its longest rays 1½ in head; least depth of caudal peduncle 3⅔ in head. Pyloric cæca about 51, the longest about 1⅝ in head, and very slender. Color in alcohol, very dark steel blue above, becoming paler below, nearly white anteriorly on belly, where only the margins of the scales are punctate; no silvery anywhere; lower jaw dusky, a large black blotch on cheek between suborbital and premaxillary; sides, back, top of head, dorsal and caudal fins with few small dark spots; pectorals dusky, slightly spotted at base; anal slightly dusky, without spots; ventrals dusky with a few spots in middle; adipose fin with a few spots; lower fins all tipped with pale, probably yellowish red in life; spots all very small and faint, not confined to posterior part of body. The specimen before us, No. 1863, L. S. Jr. Univ., is a male, 18¼ inches long. It was taken at Fairholme on Lake Crescent, Clallam County, Washington, March 12, 1896, by Mrs. G. E. Mitchell, of Fairholme. (Named for Crescent Lake, Washington, the type locality.)

Salmo gairdneri crescentis, JORDAN & BEARDSLEE, Proc. Cal. Ac. Sci., ser. 2, vol. VI, 1896, 207, pl. 22, Crescent Lake, Clallam County, Washington. (Coll. Mrs. George E. Mitchell. Type, No. 1863, L. S. Jr. Univ. Mus.)

Page 504. Under *Cristivomer* for "Eastern North America" read "Northern North America." The genus occurs also in the lakes of Alaska and British Columbia.

Page 508. Before *Salvelinus alpinus* insert:

784(a). SALVELINUS KUNDSCHA, Pallas.

This seems to be a species very distinct from *S. malma*. A specimen in the United States National Museum (No. 33814) from Petropaulski has been described by Bean & Bean as follows:

Similar in form to *S. malma*, but the body stouter and less elongate. Head 4¼ to 4⅜ (4⅛ in the Tareinsky Bay specimen); depth 4½ to 4⅝; eye 5⅛ in head, 2 in interorbital, or 1⅓ in snout; maxillary reaching to or beyond vertical through posterior edge of orbit; upper jaw nearly ½ length of head; lower jaw slightly shorter than upper. Hyoid teeth feebly developed. Scales small, 36–195, 122 pores. Fins all short; origin of dorsal about midway between tip of snout and base of upper caudal lobe, the base of the fin nearly as long as the longest ray, or ½ as long as head, its

upper margin very slightly concave, the last ray 2 in the longest; adipose fin over end of anal, its width about ½ its length, which is about equal to eye; pectoral 7 to 7½ in body length; ventral under middle of dorsal, not nearly reaching vent, its length 2 in head; caudal emarginate, its middle rays ⅓ the outer; anal scarcely concave when expanded. Pyloric cæca 22; branchiostegals 12; gill rakers 6 + 10, the longest less than ½ eye. Color bluish gray above, whitish below; the sides with numerous large white spots, some of which are ⅔ as large as eye. (Bean & Bean.)

This species is said to be common from Kamchatka northward, but only 6 specimens are actually extant, 4 obtained at Petropaulski by Dr. Leonhard Stejneger and 1 by Col. N. Grebnitski, and now in the United States National Museum, and 1 obtained from Tarcinsky Bay by Mr. Gerald E. H. Barrett-Hamilton and now in the museum of Stanford University.

Salmo kundscha, PALLAS, Zoogr. Rosso-Asiat., III, 250, 1811, Kamchatka.
Salmo leucomænis, PALLAS, Zoogr. Rosso-Asiat., III, 250, 1811, Kamchatka.
Salmo curilus, PALLAS, Zoogr. Rosso-Asiat., III, 251, 1811, Kuril Islands.

The true *Salvelinus malma* is very common at Unalaska, Kadiak, Komandorski Islands, and Petropaulski. Specimens from these various places are all alike. Head 4¼ to 4⅓; depth 4¼ to 4⅗. Spots grayish, tinged with red, much smaller than eye. Caudal well forked; lower fins short; pectoral reaching halfway to vent. Hyoid teeth present. The head seems much shorter than in examples from the United States. The dwarf form from the little brook (Pyramid Creek) at the head of Captains Harbor agrees fully in form with large examples taken in the sea about Unalaska. The small ones are brighter in color and mature at 4 to 6 inches. The form occurring throughout the northwestern United States, and described on page 508 as *Salvelinus malma*, should apparently be regarded as a species distinct from *S. malma*, and would stand as—

784(a). **SALVELINUS PARKEI** (Suckley).

Page 515. Add this footnote to *Salvelinus oquassa marstoni:*

A specimen of *Salmo marstoni* sent me some days ago indicates a more distinct species than was at first supposed. This is the most slender of our charrs, apparently the swiftest. The male is gorgeous; brilliant red extends upon the back and onto the dorsal and caudal fins as well as upon the other fins. Though quite distinct, the species is nearer to *S. oquassa* than any other. (Garman, in lit., March 24, 1895.)

Page 524. After *Osmerus dentex* add:

794(a). **OSMERUS ALBATROSSIS**, Jordan & Gilbert, new species.

(KADIAK SMELT.)

Head 4⅖; depth 5½. D. 2, 10; A. 1, 20; scales 75; maxillary 2$\frac{1}{10}$; eye 5¼; snout 3⅖; mandible 2; pectorals 1½; ventrals 1½; dorsal 1⅔; caudal 1⅘. Body elongate, moderately compressed; back elevated at nape so that anterior profile is somewhat depressed between and behind eyes; interorbital space 3⅘ in head. Mouth large, lower jaw heavy, strongly projecting; opercle with concentric striæ; pectorals moderate; ventrals long; dorsal high; anal fin low, very long, its longest ray 2⅔ in head; caudal moderate, well forked; ventrals inserted before dorsal.

Scales small, deciduous, those on back still smaller; lateral line distinct. Gill rakers long and slender, about 12 below angle of arch, longest about as long as eye. Tongue with moderate teeth, the anterior 2 to 4 small hooked canines; upper jaw with small sharp teeth similar to those in lower jaw, none of them canine-like; small teeth on palatines and pterygoids; vomer with 2 very small canines scarcely fang-like. Color bluish above with bright reflections; scales margined with dark points; sides silvery with golden and coppery luster; inside of gill openings dusky; fins white, somewhat dotted. About Kadiak Island, Alaska. Two specimens caught in the upward haul of a dredge in Shelikof Straits, north of Karluk, Kadiak Island, Alaska, at *Albatross* Station No. 3675. The depth of the dredge haul was 109 fathoms, but these fishes were no doubt taken from near the surface. One specimen is 8, the other about 7 inches in length. The species is allied to *Osmerus dentex*, the Rainbow Smelt, but differs in the extremely long anal and in the very weak vomerine and lingual canines. The flesh is firm, as in *O. dentex*. (Named for the U. S. Fish Commission steamer *Albatross*.)

Osmerus albatrossis, JORDAN & GILBERT, Rept. Fur. Seal Invest., MS. 1898, Shelikof Straits, north of Karluk, Alaska.

Mesopus should replace *Hypomesus*. It is originally characterized on page 14 (not 168) Proc. Ac. Nat. Sci. Phila. 1862, *Hypomesus* on page 15. The ventrals are inserted below front of dorsal in *Mesopus* as in allied genera, and there are 8 branchiostegals as in allied groups. The feeble teeth distinguish *Mesopus* from *Osmerus*. The statement that the stomach is cæcal in *Argentinidæ* is true of a few genera only, and the character has no high systematic value. In *Mesopus pretiosus* and *Osmerus dentex*, the stomach is siphonal, as in *Salmonidæ*. In *Thaleichthys pacificus*, however, the stomach forms a blind sac. The small number of pyloric cæca and the peculiar structure of the ovaries remain to define *Argentinidæ* as a family distinct from *Salmonidæ*.

Page 525. To the synonymy of *Hypomesus olidus* add:

Osmerus oligodon, KNER.

The species ranges south to Amur River.

Page 530. After *Bathylagus pacificus* add:

804(a). BATHYLAGUS BOREALIS, Gilbert.

Head $4\frac{1}{2}$ to base of caudal; depth $5\frac{2}{3}$; eye $2\frac{1}{2}$ in head; snout $2\frac{3}{4}$ in eye. D. 8; A. 19; ventral 8; pectoral 8. Scales in about 40 rows, judging from the scars; head scaleless. Interorbital width grooved, the groove widening posteriorly, opening onto the flat occipital region, which is not swollen. Width of cartilaginous portion of interorbital space $\frac{1}{3}$ orbit; including the thin membranaceous plates which overarch the orbits, the interorbital width is $\frac{3}{4}$ orbit. The anterior profile of snout declines gently, bringing the mesial portion of premaxillaries on a level with lower margin of pupil. Distance from tip of snout to end of maxillary slightly exceeding length of snout, $2\frac{1}{3}$ in orbit. Opercle with 2 strong ridges diverging downward and backward from behind the eye. Front of dorsal midway

between front of snout and adipose fin; base of dorsal contained $3\frac{1}{3}$ times in length of head. Ventrals inserted under posterior portion of dorsal. Free portion of adipose fin very long and narrow, rising above the base of the second and third anal rays before the last, its tip reaching rudimentary caudal rays when depressed; anal fin rather long, the base $1\frac{2}{3}$ in head, the vent immediately before it. Length of tail much exceeding head, $3\frac{2}{3}$ in total length without caudal. Uniform blackish brown on sides, the head and ventral region blue black. Differing from *B. pacificus* in its much greater depth, longer tail, longer anal fin, and flat occiput. Length 132 mm. Bering Sea, in deep water north of Unalaska; 2 specimens known. (*borealis*, northern.)

Bathylagus borealis, GILBERT, Rept. U. S. Fish Comm. 1893 (Dec. 9, 1896), 402, Bering Sea at Albatross Station 3327, north of Unalaska, in 322 fathoms.

804(b). BATHYLAGUS MILLERI, Jordan & Gilbert, new species.

Distinguished by the posterior insertion of the dorsal fin and the greatly swollen occipital region provided with a median keel. The type is in very poor condition, the skin being largely denuded from head and body. No traces remain of the scales, the pectoral and ventral fins are lost and the others greatly mutilated. Enough remains, however, to demonstrate that it is distinct from all known species and to furnish characters by which the species may be recognized. The interorbital space is converted into a very deep channel by 2 vertical thin lamellæ which arise on either side, and mark off the narrow interorbital space from the contiguous supraocular areas. From the base of these vertical lamellæ arise externally the thin supraocular plates, which extend outward and upward and roof over the orbit. A deep narrow channel is included between the lamellæ and the plates. The floor of the interorbital groove is raised mesially into a sharp ridge, which is continuous anteriorly with the ethmoidal ridge and posteriorly with a ridge running along middle of occiput. On anterior half of occiput this ridge is a high strong keel; posteriorly, it becomes lower and rounded. The occipital region is swollen and prominent, much higher than the interorbital space. It is bounded laterly by 2 strong rounded ridges which originate at the upper posterior margin of the orbit and converge rapidly backward. The occipital cartilage is heavy and strong, not yielding readily to pressure. The width of interorbital space is $\frac{1}{4}$ orbit; the distance between outer margins of orbital plates above middle of eyes is $\frac{2}{3}$ diameter of eye. The opercle is marked with delicate striæ diverging downward and backward, but is without strong ridges. The front of dorsal is midway between adipose fin and gill opening, slightly nearer base of caudal than tip of snout. The fin contains 8 rays. Anal badly mutilated, containing at least 24 rays. The mutilated condition of the type will not permit further description. Length 155 mm. Cortez Banks off San Diego, California, in deep water; known only from the type taken by the *Albatross* at Station 3627, in 776 fathoms. (Named for Walter Miller, professor of classical philology in Leland Stanford Jr. University, in recognition of his intelligent interest in zoological nomenclature.)

Page 531. In key under *h*, for "incomplete" read "complete."

Page 537. We can not separate *Synodus jenkinsi* from *Synodus scituliceps*, and the former name should probably be abandoned.

Page 555. *Macrostoma angustidens* and related species need further study. The synonymy and application of the names *angustidens*, *elongatus*, and *resplendens* are uncertain. *Macrostoma brachychir* is probably a good species.

In *M. caudispinosum* the dorsal has 20, not 36, rays.

Page 580. The generic name *Bonapartia*, Goode & Bean, is preoccupied in ornithology. For its use in fishes the name *Zaphotias* is proposed, taking the same species (*pedaliota*) as type. The genus and its species would then stand as follows:

274. ZAPHOTIAS, Goode & Bean, new generic name.

(*Zaphotias*, having organs which emit light; ζα, intensive particle; φῶς, light.)

872. ZAPHOTIAS PEDALIOTUS (Goode & Bean).

Page 582. *Cyclothone microdon* occurs also in Bering Sea in very deep water.

Page 586. *Astronesthes* is from ἄστρον, star; ἐσθής, vestment.

Page 594. *Plagyodus* (Steller) should probably supersede *Alepisaurus*, in which case the family becomes *Plagyodontidæ*.

Page 603. *Sternoptyx diaphana* is common off both the Japanese and Hawaiian islands.

Page 608. For *Aldrorandia*, Goode & Bean, substitute the earlier name *Halosauropsis*, Collett.

Hallosauropsis, COLLETT, Camp. Sci. Hirondelle, June, 1896, 143 (*macrochir*).

Page 618. Add:

916(a). MACDONALDIA ALTA, Gill & Townsend.

D. 32; A. 31 to end of dorsal, 52 spines, 125 rays. Body comparatively high, greatest height equal to $3\frac{3}{8}$ the distance between vent and tip of snout; pectoral fin with its root twice as far from upper cleft of branchial aperture as from the lateral line, and much nearer to the posterior end of operculum than to lateral line. Bering Sea; only the type known. (*altus*, deep.)

Macdonaldia alta, GILL & TOWNSEND, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17, 1897), 232, Bering Sea, Lat. N. 54° 54′, Long. W. 168° 59′, Albatross Station 3604, Aug. 13, 1895, in 1,401 fathoms. (Type, No. 48774, U. S. Nat. Mus. Coll. *Albatross.*)

916(b). MACDONALDIA LONGA, Gill & Townsend.

D. 33; A. 26 to opposite end of dorsal, 55 spines, 111 rays. Body comparatively slender, with the greatest height about $\frac{1}{5}$ distance between vent and tip of snout; pectoral fin with its root 3 times as far from upper cleft of branchial aperture as from lateral line, and very much nearer

lateral line than end of operculum. Bering Sea; only the type known. (*longus*, long.)

Macdonaldia longa, GILL & TOWNSEND, Proc. Biol. Soc. Wash., XI, 1897 (Sept. 17, 1897), 232, Bering Sea, Albatross Station 3607, 1895, in 900 fathoms. (Type, No. 48775, U. S. Nat. Mus. Coll. *Albatross*.)

Page 627. *Lucius vermiculatus* occurs also in Texas, specimens having been obtained in both the Trinity and Neches rivers near Palestine, by Evermann & Scovell.

Page 632. *Aplocheilus* = *Apocheilichthys* = *Haplocheilus* = *Panchax*, is a genus distinct from *Fundulus*, and should be erased from the synonymy of the latter. The genus is defined by the flat, much produced snout, and the long anal fin.

To the synonymy of *Fundulus* add:

Plancterus, GARMAN, Monogr. Cyprinodonts, in Mem. M. C. Z., XIX, No. 1, 96, 1895 (*kansæ* = *zebrinus*.)

Page 635. In the key, under *aa*, the phrase "inhabiting mountain springs and brooks" applies only to Nos. 943 and 944. It should be transferred and made a part of *s*.

Page 637. *Fundulus punctatus* and *F. vinctus* are wrongly placed by Garman in the synonymy of *F. parvipinnis*.

Page 638. *Fundulus pallidus* is placed by Garman in the synonymy of *F. grandis*, to which it bears but little resemblance.

Page 639. To the synonymy of *Fundulus majalis* add:

Hydrargyra formosa, STORER, Proc. Bost. Soc. Nat. Hist. 1837, 76.

Page 641. To the synonymy of *Fundulus heteroclitus macrolepidotus* add:

Hydrargyra ornata, LE SUEUR, Journ. Ac. Nat. Sci. Phila., I, 1817, 131, Delaware River, near Philadelphia. (Coll. G. Ord.)

Garman regards *Fundulus grandis* as a good species. We have recently compared specimens from Cape Cod with others from Tampa, and reach the same conclusion.

Page 642. Before *Fundulus ocellaris* insert:

932(c). FUNDULUS HETEROCLITUS BADIUS, Garman.

(This is the form found about Grand Manan, named but not characterized by Garman.)

Garman refers *Fundulus ocellaris* to the synonymy of *Fundulus grandis*, which is very doubtful.

Page 643. Garman refers *Fundulus fonticola* also to the synonymy of *F. grandis*, which is not correct. He also wrongly regards *Fundulus bermudæ* as a variety of *heteroclitus*.

Page 644. *Fundulus robustus* is referred, probably by error, by Garman to the synonymy of *F. labialis*, which is certainly incorrectly made a variety of *F. parvipinnis*.

Page 645. Garman refers *Fundulus zebra*, *zebrinus*, and *extensus* to the synonymy of *Fundulus adinia*, all of which is certainly wrong. Such ref-

erences defy all our knowledge of the geographic distribution of these fishes. For example, *F. extensus* is a brackish-water fish of Cape San Lucas; *F. zebra*, which is the basis of *F. zebrinus*, is a fish of the mountain streams of New Mexico, Colorado, and northeastward, while *F. adinia* is found near the mouth of the Rio Grande. There is no doubt that the original *Fundulus zebra* is the species called *zebrinus* by us and *kansæ* by Garman. It came from some point between "Fort Union and Fort Defiance." In other words, it came from the head waters of the Canadian River or the Rio Grande. No species of this type has been recorded from the upper Rio Grande, but the species called *zebrinus* and *kansæ* is in all the upper waters of the Arkansas basin, to which the Canadian River belongs, and doubtless in the streams above Fort Union.

Page 646. To the synonymy of *Fundulus zebrinus* add:

Fundulus kansæ, GARMAN, Monogr. Cyprinodonts, 103, pl. 2, fig. 10, 1895, Kansas.

This species (*F. zebrinus*) is rightly made the type of a new subgenus, or possibly genus, *Plancterus*, by Garman. It has long, convoluted intestines and very small pharyngeals. *Fundulus seminolis* (subgenus *Fontinus*) has short intestines and coarse pharyngeals.

Page 648. *Fundulus stellifer* is wrongly referred by Garman to the synonymy of *F. catenatus*.

Page 649. *Fundulus lineatus* is referred by Garman to the synonymy of *F. sciadicus*, which reference seems to be correct.

Fundulus albolineatus, which Garman also refers to *F. sciadicus*, seems to be a perfectly good species. It is certainly not *F. sciadicus*.

Garman refers *Fundulus confluentus* to the synonymy of *F. grandis*, which is probably not correct.

Page 650. Garman's reference of *Fundulus funduloides* to the synonymy of *F. grandis* may be correct.

The species called *Fundulus dorii* in the text is an *Aplocheilus* and should stand as:

968(a). APLOCHEILUS DOVII (Günther).

Garman recognizes *Zygonectes* as a distinct genus, but its boundaries are not easily defined.

The description of *Fundulus confluentus* should be modified to include the following, taken from the type: Head $3\frac{3}{5}$; depth $4\frac{1}{4}$. D. 11; scales 44 or fewer. A black spot on middle of membrane of last 3 dorsal rays. This species resembles *F. diaphanus* rather than *F. majalis*. *Fundulus ocellaris* seems to be identical with *F. confluentus*.

Page 651. Garman refers *Fundulus macdonaldi* to the synonymy of *F. sciadicus*, which is probably correct, but the reference of *F. floripinnis* to the same synonymy is certainly wrong.

Page 652. Garman refers *F. pulvereus* to the synonymy of *F. grandis*, which is without warrant.

Page 655. To the synonymy of *Fundulus chrysotus* add:

Gambusia arlingtonia, GOODE & BEAN, Proc. U. S. Nat. Mus. 1879, 118, Arlington River, Florida. (Type, No. 21308. Coll. Dr. Goode.)

Zygonectes henshalli, JORDAN, Proc. U. S. Nat. Mus. 1879, 237, San Sebastian River, Florida. (Type, No. 23449. Coll. Dr. James A. Henshall.)

To the synonymy of *Fundulus cingulatus* add:

Zygonectes rubrifrons, JORDAN, Proc. U. S. Nat. Mus. 1879, 237, San Sebastian River, Florida. (Type, No. 23450. Coll. Dr. James A. Henshall.)

Zygonectes auroguttatus, HAY, Proc. U. S. Nat. Mus. 1885, 556, Westville, Florida. (Type, No. 37362. Coll. Mann & Davison.)

An examination of a large amount of material recently collected in Florida by Drs. Evermann and Kendall shows that the synonymy of these species should stand as indicated above.

Examination of the type of *Gambusia arlingtonia* shows it to be the young of the form hitherto known as *Z. henshalli*, which, from an examination of the type and other specimens, proves to be the female of *Fundulus chrysotus*. The dorsal in *Gambusia arlingtonia* is not inserted so far back as the sixth anal ray, but is rather over the third or fourth. Both *G. arlingtonia* and *Z. henshalli*, agree with descriptions of *F. chrysotus* except in coloration. Both are females, as shown by form of anal fin. All specimens examined of the *henshalli* form are females, as shown in part by dissection and by the form of the anal fin. All specimens examined of the form agreeing with descriptions of *F. chrysotus* prove to be males, as shown partly by dissection and by the form of the anal fin. Front series of teeth much enlarged in all; anal fin usually with 11 rays.

The type of *Z. rubrifrons* differs from that of *Z. henshalli* in having a heavier head, really longer snout, mandible more oblique, giving the muzzle a truncated appearance, and the slope of the back to the snout beginning farther forward.

The type of *Z. rubrifrons* agrees with the description of *F. cingulatus*, except in the number of anal rays, there being 10 instead of 8, as given in the description, which is a redescription of the type of *F. cingulatus*. Cuvier & Valenciennes, however, give 10 anal rays in the original description. Specimens in the United States National Museum labeled *Zygonectes cingulatus*, from Pensacola, Florida, agree with the type of *Z. rubrifrons*, with the exception of 1 specimen, which has 11 anal rays.

National Museum specimens collected by Dr. Shufeldt at New Orleans, labelled *Zygonectes chrysotus*, contain both the *Z. henshalli* and *Z. chrysotus* forms, i. e., those with pearly spots and no cross bars (females) and those with dark cross bars (males)—that is, male and female of *Fundulus chrysotus*. Comparison of specimens collected at Tampa and Welaka, Florida, reveal 2 color forms. Most of those from Tampa have the heavier head, truncated muzzle, and outlines of *F. cingulatus*. The 2 color forms are those with dark cross bars, all males as shown by dissection and form of anal fin, and those with no cross bars and no pearly spots, which are all females. The majority of individuals have 10 anal rays each.

Most of the Welaka specimens have more slender and pointed head, preorbital less deep, really shorter snout, and the curve of the body toward the snout beginning farther back than in the preceding, and the majority have each 11 anal rays. The 2 color forms represent the 2 sexes—females with pearly spots and no cross bars, and males with dark cross bars and many with small brown spots.

While a few of the *chysotus* form are found in the Tampa collection, and a few of the *cingulatus* form with the Welaka lot, they can be easily distinguished. A very few of the *cingulatus* form have 11 anal rays and a very few of *chrysotus* 10, but they can be otherwise distinguished. Whereas the females of *F. cingulatus* have no trace of pearly spots the females of *F. chrysotus* almost invariably have them. As a rule, the cross bars in the male, *F. cingulatus,* are narrower and more numerous than in the male of *F. chrysotus,* though young individuals of the latter do not differ in this respect. In *Fundulus cingulatus* there are often faint spots on the scales of the back forming longitudinal lines which seem to be absent in *F. chrysotus.* The teeth in the front row of *F. chrysotus* are larger than in *F. cingulatus.*

Page 658. Garman refers *Fundulus guttatus* to *F. nottii,* which is very doubtful, but he is right in so referring *F. hieroglyphicus.* He also refers *F. dispar* to *F. nottii,* which is probably wrong.

Page 658. *Fundulus guttatus* (Agassiz) can not be separated from *Fundulus nottii* (Agassiz).

Page 659. *Fundulus melapleurus* is, as Garman observes, a *Gambusia,* and should stand as *Gambusia melapleura.*

Adinia guatemalensis and *A. pachycephala* are recklessly referred by Garman to the synonymy of *Fundulus parvipinnis.* They might just as well have been placed at random under any other species of a totally different fauna.

Page 660. Before *Adinia* insert:

300(a). APLOCHEILUS, McClelland.

Snout flat, both jaws much depressed. Bones of mandible firmly united; upper jaw protractile; each jaw with a narrow band of villiform teeth. Body oblong, depressed anteriorly, compressed posteriorly. Dorsal fin short, commencing behind the origin of the anal, which is more or less elongate. Intestinal tract but slightly convoluted; air bladder present. (Günther.)

Aplocheilus, McClelland, Ind. Cypr. As. Res., XIX, 301, 1839 (*chrysostigmus* = *panchax*).
Panchax, Cuvier & Valenciennes, Hist. Nat. Poiss., XVIII, 380, 1846 (*panchax*).
Haplochilus, Günther, Cat., VI, 310, 1866, corrected spelling.

968(a). APLOCHEILUS DOVII (Günther).

For description and synonymy see p. 650.

Page 662. According to Garman the air bladder is present in *Rivulus.* He refers *R. marmoratus* to the synonymy of *R. cylindraceus,* which is probably correct.

Add the following species:

973(a). RIVULUS ISTHMENSIS, Garman.

Head 3¼ in body; eye 3 in head; snout 6. D. 9; A. 11; V. 6; P. 15; scales 32–8. Elongate, compressed posteriorly, depressed forward; head broad, much depressed, flattened on the crown; snout medium, blunt;

interorbital width greater than eye. Origin of dorsal fin over middle of base of anal, $\frac{2}{3}$ distance from snout to base of caudal; origin of anal fin midway between head and caudal, the last ray nearly as far back as that of dorsal; caudal elongate, pointed, as long as head. Light olivaceous, with a dark blotch at base of dorsal and another on back above or in front of first anal ray; apparently a light, transverse streak at base of caudal. Rio San Jose, Costa Rica.

Rivulus isthmensis, GARMAN, The Cyprinodonts, Mem. Mus. Comp. Zool., XIX, No. 1, July, 1895, 140, Rio San Jose, Costa Rica. (Type in M. C. Z.)

Page 663. *Lucania ommata* is wrongly referred to the synonymy of *Heterandria formosa* by Garman.

Page 664. The species called *Lucania goodei* in the text has 2 rows of teeth and is a true *Fundulus*, or rather *Zygonectes*, as Garman has shown. It may stand as *Fundulus goodei*.

Page 665. *Lucania venusta* is wrongly referred by Garman to *L. parva*, to which, however, it is closely related.

Page 668. Garman wrongly refers *Characodon bilineatus* and *C. variatus* to the synonymy of *C. lateralis*.

Page 669. Add:

883(a). CHARACODON EISENI, Rutter.

Head $3\frac{1}{7}$; depth $3\frac{1}{4}$; eye 3. D. 11 to 13; A. 13; scales 30 to 32–12. Snout shorter than eye, lower jaw projecting. About 9 teeth in upper jaw and about 14 in lower; teeth strongly bicuspid, the villiform teeth not developed. Mouth almost vertical when closed, mandible about $\frac{1}{2}$ length of eye; interorbital space flat, the anterior part equal to orbit, wider posteriorly. Insertion of dorsal in middle of total length; anal inserted under fourth ray of dorsal; pectoral reaching past insertion of ventral; tips of depressed dorsal and anal in vertical through middle of caudal peduncle; caudal broad, truncate, length of middle rays equal to length of top of caudal peduncle. Head about $\frac{1}{4}$ of total; greatest depth of body above ventrals; depth of caudal peduncle $\frac{1}{2}$ its length. Color in alcohol, male with a broad indefinite lateral band; female with dark blotches on sides which in 1 of 3 specimens form distinct cross bands. This species is most closely related to *Characodon variatus*, Bean. It differs from that species in having fewer rays and scales, much fewer teeth, larger eye, much more posterior position of dorsal, and in color. Length $1\frac{1}{4}$ inches. Rio Grande de Santiago, Tepic, Mexico.

Characodon eiseni, RUTTER, Proc. Cal. Ac. Sci. 1896, 266, Rio Grande de Santiago, Tepic, Mexico. (Type, No. 4999, L. S. Jr. Univ. Mus. Coll. Dr. Gustav Eisen.)

Page 670. Add the following:

The specimens from Parras, Mexico, referred by Garman to *C. lateralis*, appear to be new. They may be described as follows:

984(a). CHARACODON GARMANI, Jordan & Evermann, new species.

B. 4; D. 12; A. 12; V. 6; P. 17; scales 32–11 or 12; vertebræ 15 + 18. Body compressed, moderately stout, caudal pedicel deep, back gently

arched. Head about $\frac{1}{3}$ of length to base of caudal; very little arched transversely. Snout short, not as long as the eye; chin steep. Mouth medium; upper jaw protractile. Teeth in outer series bicuspid. Eye large, nearly equal to interorbital space, $\frac{1}{8}$ longer than snout, $\frac{2}{7}$ of head. The specimen examined had 4 branchiostegal rays on each side; whether this is normal must be decided from others. Fins small; dorsal origin about $\frac{5}{7}$ of the distance from snout to caudal; anal opposed to dorsal; posterior margin of caudal subtruncate. Olive to reddish brown, with scattered small spots of darker on the back, a darker band with or without spots of dark along the flank, more distinct posteriorly. Fins with fine dots of dark color. Parras, Coahuila, Mexico. (Named for Prof. Samuel Garman of the Museum of Comparative Zoology, in recognition of his valuable studies of the *Cyprinodonts*.)

Caracodon lateralis, GARMAN, The Cyprinodonts, Mem. Mus. Comp. Zool., XIX, No. 1, pl. 1, fig. 9, 1895, **Parras, Coahuila, Mexico; not of GÜNTHER.**

984(b). CHARACODON LUITPOLDII, Steindachner.

Head $4\frac{1}{3}$ to $4\frac{2}{5}$; depth 3 to $3\frac{1}{8}$; eye 4 to $4\frac{2}{3}$ in head; snout 3 to $3\frac{2}{3}$; interorbital $1\frac{2}{3}$ to 2. D. 14; A. 15 or 16; P. 15 or 16; V. 6; scales 40–17. Body moderately slender; caudal peduncle strongly compressed; head short; upper profile slightly arched, somewhat depressed at occiput; ventral outline more convex; bases of anal and dorsal quite oblique. Dorsal rounded, longest ray $1\frac{2}{3}$ in head; anal somewhat smaller. Outer teeth slender, movable, broadened toward front of jaw which is notched; behind these a band of minute teeth, scarcely distinguishable. Two rows of scales below eye; preorbital, jaws, and narrow border of preopercle scaleless. Pectoral shorter than head, not reaching ventrals, which are nearer snout than base of caudal; origin of dorsal nearer base of caudal than gill opening; anal slightly behind dorsal. Color in alcohol, upper half of body light brown or brownish gray, lighter gray or silvery gray below, fading to yellowish white toward ventral line; a silvery gray band along middle of side, not well defined, its width that of 1 or 2 scales. Lake Pátzcuaro, Mexico. (Steindachner.)

Characodon luitpoldii, STEINDACHNER, Einige Fischarten Mexico, 12, pl. 2, figs. 3–3b, 1895, **Lake Pátzcuaro, Mexico.** (Coll. Princess Theresa von Bayern.)

Page 675. Garman refers *Cyprinodon elegans* to the synonymy of *C. eximius* and *C. felicianus* to that of *C. rivcrendi*, both of which seem to be correct.

Page 680. *Gambusia infans* is probably identical with *G. gracilis*, as indicated by Garman.

Page 681. Garman calls our *Gambusia affinis G. patruelis* and makes *G. holbrooki*, the northern form, a distinct species, neither of which views seems to be justifiable.

Page 682. *Gambusia nobilis* and *G. nicaraguensis* are referred by Garman to the synonymy of *G. gracilis*, which is questionable; but his reference to *G. puncticulata* of *G. picturata* is probably correct.

Page 682. After *Gambusia affinis* add:

1000(a). GAMBUSIA TRIDENTIGER, Garman.

Head 4½; depth at anal 4⅓; snout short, not as long as eye, narrow, rounded forward, and blunt. D. 7 or 8; A. 10; V. 6; P. 12; scales 28 to 30–8; vertebræ 14 + 17. Mouth medium, directed obliquely upward; lower jaw longer than the upper, which is short, narrow, and protractile. Teeth in the outer series larger, strongly hooked, pointed, broadened somewhat toward the apex; inner series very small, in bands, tricuspid as in *Pœcilia;* pharyngeal with a shoulder. Eye large, longer than snout, 3 in head. Fins small, excepting the caudal; dorsal smaller than anal and farther back, its origin about midway from occiput to end of caudal, nearly above the hindmost anal ray, 17 or 18 scales from the head; anal origin midway between snout and end of caudal; farther forward on the male, between the ventrals, and the fin is modified to form an intromittent organ about ⅓ length of entire fish; caudal deep, as long as head, rounded on hinder margin. Scales large, median series on flank as wide as eye. Intestine short. Light olivaceous, yellowish or brownish, with 7 or 8 vertical bars of brownish, separated by light or silvery spaces of equal width, on the sides of the caudal portion, edges of scales darker, the centers or median series more or less silvery; belly and lower surface of head silvery or golden; peritoneum black, showing through abdnominal wall; occiput dark; top of snout light; a dark line between anal and caudal; dorsal with a faint spot or group of puncticulations behind the middle near the base; other fins plain to dark tipped. (Garman.) Isthmus of Panama, in fresh water (*tridentiger,* bearing trifid teeth).

Gambusia tridentiger, Garman, Cyprinodonts, Mem. Mus. Comp. Zool., xix, No. 1, 89, pl. 4, fig. 10, 1895, Isthmus of Panama.

Pages 688 and 689. *Heterandria versicolor* and *H. occidentalis* are correctly referred by Garman to the genus *Pœcilia.* It is not improbable that *H. versicolor* is the same as *Pœcilia vivipara,* Bloch & Schneider.

Lebistes is doubtless identical with *Pœcilia,* as is also *Acropœcilia. Acropœcilia tridens* is probably identical with *Pœcilia dominicensis,* as stated by Garman.

Page 691. Garman wrongly refers *Pœcilia butleri* to the synonymy of *P. sphenops.*

Most of the Mexican and Central American species are imperfectly known and imperfectly described. Of these Garman refers the following to the synonymy of *P. sphenops,* whether correctly or not only a study of adequate material can determine: *Pœcilia mexicana, P. thermalis, P. petenensis, P. dovii, P. couchiana, P. plumbeus, P. fasciatus,* and *P. spilurus.*

Pœcilia pavonina is referred, perhaps correctly, to the synonymy of *P. vittata.*

Page 696. Garman thinks that *Pœcilia vandepolli* is identical with *P. reticulata,* Peters, which may be described as follows:

1032. PŒCILIA RETICULATA, Peters.

D. 7 or 8; A. 8 or 9; V. 5; scales 26 to 28–8. Depth of body ⅔ and length of head nearly ¼ of the length to the base of the caudal. Males rather more

slender. Eye longer than snout, not quite $\frac{1}{3}$ of head, $\frac{3}{5}$ of interorbital space. Forehead flat. Dorsal origin somewhat nearer to end of snout than to end of caudal, opposite first ray of anal on females. Anal of male advanced, between the ventrals, which are elongate; anal process as long as the head, without hooks. Caudal large, rather longer than head, obtusely rounded; free portion of tail somewhat elongate, base of anal being $\frac{1}{3}$ of its distance from the caudal; ventrals reaching anal; pectorals as long as the head, not reaching ventrals. Female yellowish olive, scales with a narrow blackish edge, belly silvery, trunk above the belly blackish. Male with 2 brown streaks along the trunk, sometimes confluent into a band, a brown streak along the middle of the side of the tail, a round black spot behind the shoulder, another at the commencement of the caudal streak, and a third at the root of the caudal; 1 or 2 of these spots may be absent. Trinidad; Venezuela (*reticulatus*, netted).

The male from Venezuela differs in color from those from Trinidad. It has large silvery patches between the brown streaks, and a large ovate black spot in the middle of the side of the tail. (Günther.)

NOTE.—The following is the original description: "Grüngelblich mit einem schwarzen Netzwerk, dessen Maschen den Rändern der Schuppen parallel liegen, am Bauche silbrig. Schuppen in 7 Längs- und in 27 Querreihen; obwohl einige derselben durchbohrt erscheinen, ist doch keine deutliche Seitenlinie zu sehen. Ganze Länge 39, Höhe 9, Länge des Kopfes 7 Millimeter. D. 8; A. 10. Caracas; in dem Guayre-Flusse von Gollner gesammelt."

Pœcilia reticulata, PETERS, Monatsb. Berl. Ak. 1859, 412, Caracas; GARMAN, Cyprinodonts, 63, 1895.

Girardinus guppii, GÜNTHER, Cat., VI, 353, 1866, Trinidad; Venezuela; EIGENMANN, Proc. U. S. Nat. Mus. 1891, 65.

Girardinus vandepolli, VAN LINDTH DE JEUDE, Notes from Leyden Museum, IX, 137, 1887, Curaçao, one of the Leeward Islands.

Pœcilia vandepolli arubensis, VAN LINDTH DE JEUDE, Notes from Leyden Museum, IX, 137, 1887, Aruba, one of the Leeward Islands.

Pœcilia branneri, EIGENMANN, Ann. N. Y. Ac. Sci. 1894, 629.

Page 697. Garman refers *Pœcilia elongata*, one of the best marked species of large size, and marine in its habitat, to the synonymy of *P. gillii*. This is certainly wrong, as is also the reference to *P. gillii* of *P. chisoyensis* and *P. boucardi*.

P. melanogaster is probably correctly referred to *P. dominicensis*.

Page 698. Add:

1037(a). PŒCILIA CUNEATA, Garman.

B. 5; D. 8 to 10; A. 10 or 9; V. 6; P. 15 or 16; scales 28 or 29–9. Short and deep; caudal pedicel deep. Head depressed, broad, flat on the crown, equaling depth between dorsal and anal, or $\frac{1}{4}$ of the length to the base of the caudal; snout as long as the eye, broad, truncate; chin short, steep; mouth wide, directed upward; jaws weak, loosely joined, lower short, upper shorter, protractile; outer series of teeth slender, oar-shaped, hooked, movable; inner in bands, small, pointed; eye large, as long as snout, $\frac{1}{3}$ of interorbital space, $\frac{1}{4}$ of head. Dorsal larger than anal, origin midway from head to base of caudal, over third ray of anal, 13 scales behind the occiput. Anal small, acute angled, third ray longest; on the

male the base of the anal is forward of that of the dorsal, the fin is modified to form a sharp-pointed organ in which the rays are less changed than in most species; its length is less than that of the head. Ventrals small, not reaching the anal. Pectorals reaching back over 7 scales. Caudal deep, as long as the head, hind margin rounded. Scales large. Intestine long. Brownish, olive tinted, bases of scales dark, back darker, and top of head darkest; more or less of the hind margin, or ½ of the scale, is whitish to silvery on the scales of the flank; lighter to silvery under head and abdomen; dorsal with 1 to several transverse series of small spots of black; fin sometimes black tipped; a brownish streak extending back and upward on the opercle behind the eye; caudal with small spots of black on the basal half, or with a couple of clouded transverse bands; other fins uniform or punctulate; very small ones are lighter with a faint silvery band along the middle of the flank, but without vertical bars; a large one has numerous small white spots, somewhat like *Fundulus heteroclitus*. Females 2½ and males $1\frac{9}{10}$ inches. Turbo, Gulf of Darien.

Pœcilia cuneata, GARMAN, Cyprinodonts, 62, pl. 5, fig. 3, 1895, Turbo, Gulf of Darien.

Page 704. After *Typhlichihys*, Girard, add:

a No scleral cartilages; no pigment in or about the eye; retinal elements readily separable into ganglionic, inner reticular, and nuclear layers, the nuclear and outer reticular layers rarely distiguishable; diameter of eye about .150 mm.
SUBTERRANEUS, 1047.

aa Scleral cartilages large, forming a hood over front of eye; a mass of pigment in front of eye; pigment layer of retina with more or less pigment; eye a mere vestige, about .040 mm. in diameter. ROSÆ, 1047(a).

Page 706. After *Typhlichthys subterraneus* add:

1047(a). TYPHLICHTHYS ROSÆ, Eigenmann.

Extremely close to *T. subterraneus*, from which it seems to differ only in the less development of the eye. Scleral cartilages large, forming a hood over the front of the eye; a mass of pigment in front of eye; pigment layer of retina with more or less pigment; eye a mere vestige, ½ the size of that of *T. subterraneus*, about .040 mm. in diameter. The types of this species are 2 small, thoroughly dissected specimens, in the Museum of Indiana University, collected from a cave in Jasper County, Missouri, by Miss Ruth Hoppin. (Named for Mrs. Rosa Smith Eigenmann.)

Typhlichthys rosæ, EIGENMANN, Science, N. S., vol. VII, No. 164, 227, February 18, 1898, cave near Sarcoxie, Jasper County, Missouri.

Page 723. *Hemiramphus balao* is a valid species as defined.

Page 729:

Exocœtus volitans, Linnæus, as Lönnberg has shown, is identical with *E. evolans* L. As the genus *Exocœtus*, Syst. Nat., Ed X, 316, is based solely on *Exocœtus volitans*, the name *Exocœtus* must go with this species, taking the place of *Halocypselus*. The ordinary flying fishes must therefore be called *Cypsilurus*. The species with long anal fin may, however, be held as generically distinct from the type of *Cypsilurus*, and for them (*exsiliens, rondeletii*, etc.) the name *Exonautes* has been proposed by Jordan & Evermann, Check List, 322. (Type, *exsiliens*.) (ἔξω, out of; ναυτής, swimmer.)

Our species of *Exonautes* are the following:

1080. EXONAUTES EXSILIENS (Müller).

1081. EXONAUTES RONDELETII (Cuvier & Valenciennes).

1082. EXONAUTES VINCIGUERRÆ (Jordan & Meek).

1083. EXONAUTES SPECULIGER (Cuvier & Valenciennes).

1084. EXONAUTES RUFIPINNIS (Cuvier & Valenciennes).

To these should be added the following:

1084(a). EXONAUTES AFFINIS (Günther).

Head 4⅓ depth 6; eye 3½; snout 3½. D. 11 to 13; A. 11 to 13; scales 6-50 to 52, 35 before dorsal. Interorbital space flat, slightly greater than eye. Pectoral fin extending scarcely beyond dorsal and anal; base of ventral midway between eye and base of caudal, its rays reaching beyond middle of base of anal; dorsal opposite anal, its anterior rays 2½ in head. Pectoral with an oblique white blotch across its lower half, and with a narrow whitish margin; ventrals grayish. Cuba? Atlantic; West Africa. (Günther.) Probably distinct from *E. speculiger*.

Exocœtus affinis, GÜNTHER, Cat., VI, 288, 1866, Cuba?

The species of *Cypsilurus* are the following:

1085. CYPSILURUS HETERURUS (Rafinesque).

1086. CYPSILURUS LUTKENI (Jordan & Evermann).

1087. CYPSILURUS FURCATUS (Mitchill).

1088. CYPSILURUS NIGRICANS (Bennett).

1089. CYPSILURUS XENOPTERUS (Gilbert).

1090. CYPSILURUS LINEATUS (Cuvier & Valenciennes).

Under this species (p. 739) for Corea (in 3 places) read Gorea.

1091. CYPSILURUS CYANOPTERUS (Cuvier & Valenciennes).

This is a good species. The specimens recorded from James Island belong to *C. bahiensis*.

1092. CYPSILURUS BAHIENSIS (Ranzani).

1093. CYPSILURUS CALIFORNICUS (Cooper).

1094. CYPSILURUS CALLOPTERUS (Günther).

1095. CYPSILURUS GIBBIFRONS (Cuvier & Valenciennes).

Page 732. In the key, for "*jj*" read "*hh*," for "*jjj*" read "*hhh*," for "*kk*" read "*ii*," for "*ii*" read "*gg*," and for "*hh*" read "*ff*."

Page 746. According to the studies of Mr. Rutter and Dr. Gilbert all the forms of *Gasterosteus* should probably be reduced to a single species (*Gasterosteus aculeatus*), having 3 or 4 geographic varieties, each running into a number of forms which differ in the degree of armature of the body.

Page 749. After *Gasterosteus bispinosus cuvieri*, insert:

1100(a). GASTEROSTEUS GLADIUNCULUS, Kendall.

Head 3⅓; depth 3⅓; D. II-I, 10; A. I, 8. Head rather long; eye about 3 times in head; opercle not striate; body deep, compressed, with 5 lateral

dermal plates anteriorly counting from pectoral fin, none posteriorly; caudal peduncle short, naked, not keeled; innominate bone lanceolate, its width about 3 times in length; ventral spines rather long, about $1\frac{2}{3}$ times in head, serrated above and below, a strong cusp at base on both upper and lower edge. Color in life, grass green, mottled and finely punctated with black on top of head and back; sides of head and body golden, with dark blotches; breast silvery, ventrals scarlet. In alcohol the back becomes smoky black, the mottling and black dots more distinct, the golden hue of the sides fades, becoming more or less silvery, the dark blotches more pronounced. Coast of Maine and Woods Hole, Massachusetts. (*gladiunculus*, little sword; sticklebacks being called by the boys about Portland, Maine, "Little swordfish.")

Gasterosteus gladiunculus, KENDALL, Proc. U. S. Nat. Mus. 1895, 623, off Seguin Island, Maine. (Type, No. 47589. Coll. *Grampus*.)

Page 754. *Aulostomus maculatus* is pinkish-red in life.

Page 757. *Fistularia tabacaria* has been recorded by Storer from Holmes Hole, Massachusetts, and H. M. Smith records it from Buzzards Bay, near Quisset, and from about Woods Hole.

Page 762. In the key to species of *Siphostoma* read:

eee. Dorsal covering 4 or 5 caudal (not body) rings.
o. Rings 16 to 18 + 29 to 33.
q. Rings 16 + 30 to 33; dorsal 30 to 34, on 3 + 5 rings.

Page 767. It is doubtful if *Siphostoma pelagicum* occurs in America. *S. rousseaui* has probably been sometimes mistaken for it.

Page 768. After *Siphostoma jonesi* add:

1124(a). SIPHOSTOMA ROBERTSI, Jordan & Rutter.

Head $7\frac{1}{3}$ in length; depth $2\frac{1}{3}$ in head; eye $5\frac{2}{3}$ in head. Dorsal 20, on 0 + 4 rings; segments 17 + 32. Snout $2\frac{1}{3}$ in head, with a slight keel; a slight keel on top of head, another above opercle, and 1 on anterior side of opercle, but not reaching posterior edge; shields without spines; lateral keel ending on last body segment; ventral keel on next to last; upper body keel extending nearly to end of dorsal fin, upper caudal beginning below it on first caudal segment; all ridges of body very prominent, the tail with 4 plain ridges; caudal pouch 3 in total length. Color mottled brown, paler below, the membrane connecting the segments pale bluish, forming cross stripes which are especially marked on the egg pouch; prominent pale cross bars on lower side of head; dorsal colorless, except that the base is finely dusted with brown; caudal thickly dusted with brown, except near base. This species is most closely related to *Siphostoma jonesi*, differing in having a shorter dorsal with more rays, and in the lateral keel ending distinct from lower caudal keel. Jamaica; 1 specimen, $4\frac{1}{2}$ inches long, known. (Named for Rev. Joseph Seed Roberts, who collected the type.)

Siphostoma robertsi, JORDAN & RUTTER, Proc. Ac. Nat. Sci. Phila. 1897, 97, Kingston, Jamaica. (Type, No. 4988, L. S. Jr. Univ. Mus.)

1124(b). SIPHOSTOMA STARKSII, Jordan & Culver.

Head 10½; depth 21. Dorsal 38, on 0 + 10 or 11 rings. Rings 13 or 14 + 37 or 38. Head and body in tail 2. Snout 2⅔ in head. Dorsal ½ longer than head. Body rather stout. Head scarcely carinate above. Snout with a slight smooth carina. Two lateral keels, confluent into 1 behind. Belly slightly keeled; no keel on opercle. Color dark olive, much mottled with darker but without distinct markings; yellow below. Male and female common in the fresh waters of Rio Presidio at Mazatlan, among algæ; not seen in salt or brackish water. The pouch of the male teeming with eggs in January. Length 4 to 6 inches. Mazatlan, Mexico. Common in the Rio Presidio in sluggish water, on the bottom, about a mile below the village of Presidio. The species is probably found in brackish and fresh waters rather than in the sea.

Siphostoma starksii, JORDAN & CULVER, Fishes Sinaloa, in Proc. Cal. Ac. Sci. 1895, 416, pl. 30, Rio Presidio, Mazatlan. (Type, No. 2686, L. S. Jr. Univ. Mus. Coll. Hopkins Exped. to Mazatlan.)

1124(c). SIPHOSTOMA SINALOÆ, Jordan & Starks, new species.

Allied to *Siphostoma arctum* Jenkins & Evermann.

Head 8½ in length to base of caudal; depth 3½ in head. Dorsal 26, on 1½ + 5 rings, 14 + 35. Snout 1¼ in head, a strong median ridge above running to between middle of eyes, a ridge on each side from angle of mouth to below eye, occipital and nuchal plates keeled, a slight keel on anterior part of opercle; dorsal keels ceasing in front of the last 4 or 5 rays of dorsal, the lateral ridge running up and continuing as dorsal ridges; belly with a keel on each side. Preanal part of belly 1¾ in postanal part; pectoral shorter than eye, caudal 3 in head. Color olive brown above, abruptly lighter below lateral ridges anteriorly, the edges of the plates dark, forming reticulations on lower parts of body; between every 4 rings is a narrow white cross bar; from each eye is a narrow light bar running upward and backward to occiput; caudal dark. The 2 type specimens, 1 of which was sent to the British Museum, collected by the Hopkins Expedition at Mazatlan. They were erroneously referred to *Siphostoma arctum* in our paper on the Fishes of Sinaloa. Type, No. 2945, L. S. Jr. Univ. Museum.

Page 772. *Corythroichthys*, Kaup, should apparently be recognized as a genus distinct from *Siphostoma*. The species belonging in it are the following:

1134. CORYTHROICHTHYS ALBIROSTRIS, Heckel.

1135. CORYTHROICHTHYS CAYANNENSIS (Sauvage).

1135(a). CORYTHROICHTHYS CAYORUM, Evermann & Kendall.

Head 8⅔; depth 12⅔; snout 3⅓ in head; eye 4⅓. D. 21 rays, on 1½ + 3½ rings; A. 3, on first caudal ring; C. 10; P. 10. Rings 17 + 26 = 43. Body short and stout; head short, snout very short; tail but little longer than head and trunk. Cranial ridges strong; a high, sharp keel on snout, the occipital keel very high, its edge convex, notched near the middle, not

continuous with keel on snout; a strong supraocular ridge, beginning opposite posterior end of nasal keel and continuing backward with 1 hiatus upon upper edge of opercle; just below this on the opercle another longer but scarcely stronger ridge; another short ridge on anterior part of opercle at level of lower part of eye; opercles very convex, as if swollen outward; keels on body and tail all strong; the 2 lateral keels on body terminating on third caudal ring; the 2 lateral keels on tail beginning on the last body ring, thus overlapping the body keels; median keel on side well developed, terminating on sixteenth body ring; ventral keels strong; abdominal keel very strong. Egg sac on first 18 caudal rings. Color yellowish brown, with darker punctulations; tip of snout white; cheek, throat, and under parts of snout white, crossed by about 7 or 8 irregular brownish bars extending downward and backward; opercles brown; fins pale. This species is related to *C. albirostris* of Heckel, differing from it chiefly in the shorter snout, smaller dorsal, and fewer rings. Key West, Florida. (*cayorum*, of the Keys; from Cayo Hueso, Bone Key, the original Spanish name of the island of Key West.)

Corythoichthys cayorum, EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897 (Feb. 9, 1898), 128, pl. 7, fig. 7, near Crawfish Bar, Key West, Florida. (Type, a male 3½ inches long, No. 48784. Coll. Drs. Evermann & Kendall.)

Page 774. *Syngnathus æquoreus* is doubtfully American. Until a comparison of specimens can be made our species may stand as—

1138. SYNGNATHUS HECKELI (Kaup).

Page 792. *Lethostole*, Jordan & Evermann, is identical with *Chirostoma*, and the definition assigned is that of *Chirostoma*.

To the synonymy of *Chirostoma estor* add:

Atherinichthys albus, STEINDACHNER, Anzeiger der Kais. Akad. Wiss. Wien, 1894, 148, Lake Pátzcuaro, Mexico. (Coll. Princess Therese von Bayern.)

Page 793. In *Chirostoma humboldtianum* the scales are serrulate. After this species insert the following:

1155(a). CHIROSTOMA GRANDOCULE (Steindachner).

Head 4; depth 5⅔; eye 3⅔ in head; interorbital width 4⅓; pectoral fin 1⅓; ventral 2¼; caudal 1⅕; anal base 1¼, its greatest height 1⅜. D. V–I, 10; A. I, 20; P. 15 or 16; scales 60 to 62–15 or 16. Upper profile of head merging gradually into that of back, rising slightly toward beginning of second dorsal. Lower jaw slightly projecting; posterior end of upper jaw reaching eye. Teeth on maxillary sharp, brush-like, in 3 or 4 rows, the inner teeth of the maxillary and the outer teeth of lower jaw somewhat enlarged and close set. Cheek narrower than in *C. humboldtianum* and *C. estor*, and with 4 rows of scales. Origin of first dorsal midway between anterior border of eye and base of caudal, the second dorsal ½ diameter of eye nearer base of caudal than hinder border of eye; greatest height of second dorsal scarcely greater than base of fin. Longest anal ray about 1½ in base of fin; dorsal and anal concave on free border; origin of anal nearly an eye's diameter in front of that of second dorsal; caudal deeply incised, the mid-

dle rays about 2 in the longest; caudal peduncle more than $4\frac{2}{5}$ in body, its least depth somewhat more than 2 in greatest depth of body. Scales slightly ctenoid. Side with a broad, sharply defined silvery-gray band. Body much more slender, snout shorter, and eye larger than in *C. humboldtianum* or *C. estor*. Length 5 inches. Lake Pátzcuaro, Mexico.

Atherinichthys grandoculis, STEINDACHNER, Anzeiger der Kais. Akad. d. Wissensch. Wien. 1894, 149, Lake Pátzcuaro, Mexico. (Coll. Princess Therese von Bayern.)

354(a). ESLOPSARUM, Jordan & Evermann.

Eslopsarum, JORDAN & EVERMANN, Check-List Fishes, 330, 1896 (*jordani*).

This genus is close to *Chirostoma*, from which it differs in the large entire scales. To it belong the 2 following species:

1156. ESLOPSARUM BARTONI (Jordan & Evermann).

1157. ESLOPSARUM JORDANI (Woolman).

To the synonymy of this species should be added

Atherinichthys brevis, STEINDACHNER, Anzeiger der Kais. Akad. d. Wissensch. Wien. 1894, 149, Lake Cuitzeo, Mexico. (Coll. Prinzessin Therese von Bayern.)

Page 793. In *Eslopsarum jordani* the anal is I, 16, not I, 6.

Page 795. *Kirtlandia laciniata* has been found to intergrade with *K. vagrans* and should stand as—

1158a. KIRTLANDIA VAGRANS LACINIATA (Swain).

Page 796. Under *d* in the key read:

d. Snout about equal to eye, which is 3 to $3\frac{1}{2}$ in head.

Page 800. An examination of numerous specimens of *Menidia* from various places between Florida and Halifax shows that *M. notata* and *M. menidia* intergrade perfectly. The first will therefore stand as—

1167a. MENIDIA MENIDIA NOTATA (Mitchill).

Page 801. *Menidia guatemalensis* and *Menidia pachylepis* belong in the genus *Thyrina*, Jordan & Culver.

Page 819 *Agonostomus nasutus* has the anal usually II, 10, sometimes II, 9.

Page 821. Add the following:

In the Transactions of the Jamaica Society of Arts for 1855, Mr Richard Hill gives a paper on "Fishes of the Jamaica Shores and Rivers" which has been overlooked by subsequent writers. The list is chiefly a nominal one, but it contains a number of vernacular names not elsewhere given. The only new species are given under the head of *Labrax* (page 142) and *Mugil* (page 143), and these are named rather than described. They are the following:

There is another *Labrax*, common enough in the Kingston market when the rains send strong freshets from the river into the harbor. The fishermen call it the river chub, and confound it with the *mucronatus*. It is a different species; it is marked with

bands like the *Perca fluviatilis* of Europe, and the *Perca granulata* of America. We will call it the *Labrax pluvialis*, rainy weather chub.

Mugil petrosus—rock mullet;
lineatus—short mullet, 1;
albula—short mullet, 2;
curema—long mullet;
equinoculus—horse-eye mullet;
capitulinus—drab mullet, long ears;
plumieri—pond mullet;
liza—callipeva;
Dajaus monticola—mountain mullet;
choirorynchus—hog-nose mullet.

The *Dajaus monticola* inhabits only the mountain streams; the *choirorynchus* or hog-nose mullet is a fish of double the size of the *monticola*, and found in the same waters. The *mugil liza* is the largest of the mullets, from 20 inches to 3 feet long; the callipeva is the name by which it is exclusively known. This is, no doubt, its Indian name. The *equinoculus* and *capitulinus*, known in the market as long mullets, are readily distinguishable from each other by the size of the head, and especially by the size of the eye: the horse-eye mullet has the large eye, the *capitulinus* unusually small. The *plumieri*, Plumier's mullet of Cuvier & Valenciennes, is a long mullet; and the *lineatus* and *albula* what the market people distinguish as short mullets. The callipeva is a river mullet seldom extending further than the embouchure of streams, or into the ponds and marshes. The *curema* is a large mullet found on the sea banks; it is the most highly colored of all the mullets, the back is a golden green and it has scales on the second dorsal fin.

Dajaus choirorynchus is identical with *Agonostomus nasutus*, but the scanty description hardly justifies the substitution of this name for the later one. The other new species we fail to identify. *Labrax pluvialis* we do not recognize.

Page 823. The great *Barracuda* should stand as—

1199. SPHYRÆNA BARRACUDA (Walbaum).

To its synonymy add:

Esox barracuda, WALBAUM, Artedi Piscium, III, 94, 1792; after CATESBY.

Page 827. The ventrals in the *Polynemidæ* are truly thoracic, the long pubic bone being attached to the shoulder girdle. This family is probably nearest allied to the *Sciænidæ*.

Page 833. After *Ammodytes personatus* add:

372(a). **RHYNCHIAS**, Gill, new genus.

Rhynchias, GILL, MS., new genus (*septipinnis*).

This generic name is provisionally given to a species known only from a description of Pallas, and supposed to differ from *Ammodytes* in the presence of ventral fins. It may prove to belong to some different family. (ῥῦγχος, snout.)

1214(a). RHYNCHIAS SEPTIPINNIS (Pallas).

This species has not been recognized by any recent collector, and it is not certain to what family it belongs. The following is the substance of Pallas's description:

D. 43; A. 24; V. 8; P. 16; C. 24. Form of *Ammodytes tobianus*. Head

compressed; snout long, slender, depressed. Maxillary with fine teeth; rictus long. Branchiostegals 4. Body compressed, slender, with transverse streaks. Scales inconspicuous; 1 lateral line. Pectoral large, unarmed. Dorsal short, well backward, lower posteriorly; caudal subbifurcate. Color white, the dorsal edged with darker. Kamchatka. (Pallas.)

If we can trust the description, this fish would seem to represent a distinct genus of *Ammodytidæ*, characterized by the presence of ventral fins, but it may be that the account is erroneous in this regard and that Pallas had in mind *Ammodytes personatus*. (*septem*, seven; *pinna*, fin.)

Ammodytes septipinnis, PALLAS, Rosso-Asiat., III, 1811, Kamchatka.

Page 833. *Ammodytes alascanus* is not separable from *A. personatus*.

Page 839. *Caulolepis longidens* occurs also in the Pacific, specimens having been collected by the *Albatross* at Cortez Banks, off San Diego, California, in 1896.

Page 847. Add:

1230(a). MYRIPRISTIS CLARIONENSIS, Gilbert.

Head $3\frac{1}{6}$ in length; depth $2\frac{5}{7}$. D. X–I, 14; A. IV, 12; scales $3\frac{1}{2}$–41–7. Least depth of caudal peduncle $\frac{1}{2}$ length of snout and eye. Greatest (oblique) diameter of eye $2\frac{1}{2}$ in head. Least interorbital width equaling length of snout, $4\frac{1}{2}$ in head. Mouth less oblique than in related species, the line of upper jaw with a more pronounced double curve. Lower jaw the longer, with well-developed symphyseal knob. Teeth finely villiform, very slightly enlarged toward middle of both jaws; wide patches of similar teeth on head of vomer and on palatine bones. Length of maxillary (measured from front of upper jaw) very slightly (about $\frac{1}{20}$) less than length of snout and eye. Color before immersion in spirits, reddish, the upper parts dusky, especially on top of head and on the margins of the scales; evident horizontal dusky streaks between the rows of scales; opercular membrane blackish; fins all light, without dark markings. Differing from all known American species of *Myripristis* in having $3\frac{1}{2}$ series of scales between the lateral line and the base of the spinous dorsal, instead of $2\frac{1}{2}$. Length $6\frac{1}{4}$ inches. Revillagigedo Islands; only the type known.

Myripristis clarionensis, GILBERT, Proc. U. S. Nat. Mus. 1896, 441, pl. 69, Clarion Island, Revillagigedo Archipelago. (Type, No. 47746. Coll. Dr. Gilbert.)

Page 852. Insert the following description by Jordan & Rutter of *Holocentrus marianus*, based upon a specimen 6 inches long from Jamaica:

Head $2\frac{3}{5}$; depth 3 in length; eye $2\frac{1}{2}$ in head. D. XI, 13; A. IV, 9; scales 4–45–7. Dorsal outline much more curved than ventral; mouth low, but little oblique, the lower jaw projecting and entering upper profile; maxillary to below middle of eye; eye large, lower margin of orbit cut by a line connecting tip of snout and upper base of pectoral; angle of opercle high, higher than top of pupil, with 3 sharp teeth, small teeth along the margin next the subopercle; subopercle long and narrow, dentate near upper end; preopercle very finely serrate, with a strong spine at angle; a single row of scales on opercle along margin of preopercle; suborbital

bones very narrow, finely serrate; premaxillary groove on top of head as long as eye; length of pectoral equals head behind middle of eye; spinous dorsal depressible into a groove, highest (anterior) rays of soft dorsal equal to ventrals, longer than soft rays of anal; third anal spine very long and heavy, as long as pectorals; caudal forked almost to base, the lobes equal, as long as pectorals. Each row of scales with a red band, yellow lines between the rows; fins all yellowish. This is a strongly marked species, very different from *Holocentrus ascensionis*, perhaps the type of a distinct genus, characterized by the large mouth and projecting chin.

Page 856. Dr. Bean reports the Red Mullet or Goat Fish (*Mullus auratus*) as being plentiful at Sandy Hook in September and October.

Page 857. The nominal genus *Mulloides* can not be separated from *Upeneus*.

Page 866. In *Scomber colias* read: Head about 3; depth $4\frac{2}{3}$; first dorsal longer than high.

Page 873. To the synonymy of *Scomberomorus* add:

Polipturus, RAFINESQUE, Anal. de la Nature 1815, 84; substitute for *Scomberomorus*.

Page 874. In line 12 of description of *Scomberomorus maculatus*, for "side" read "part."

Page 878. *Bipinnula*, Jordan & Evermann, is a synonym of *Escolar*, Jordan and Evermann, in Goode and Bean, Oceanic Ichthyology, 519, 1896. The error resulted from Goode & Bean taking our original MS. name *Escolar*, for which we afterwards substituted *Bipinnula*.

This genus and its species will therefore stand as follows:

396. ESCOLAR, Jordan & Evermann.

Escolar, JORDAN & EVERMANN, in GOODE & BEAN, Oceanic Ichthyology, 519, Aug. 23, 1896 (*violaceus*).

Bipinnula, JORDAN & EVERMANN, Fishes North and Middle Amer., 878, Oct. 3, 1896 (*violaceus*).

1267. ESCOLAR VIOLACEUS (Bean).

Page 886. Instead of *Lepidopus caudatus*, which is not yet known to occur in American waters, insert:

1276. LEPIDOPUS XANTUSI, Goode & Bean.

Head $4\frac{2}{3}$ in body; depth 3 in head; eye $5\frac{1}{2}$; interorbital space $8\frac{1}{3}$; snout 3; maxillary $3\frac{1}{3}$. D. 82; A. II, 45. Jaws with long, sharp teeth in front, followed by single rows of weaker ones, arranged in groups of twos and threes. Height of dorsal, near middle of body, 3 in head. Anal preceded by 2 scutes, the first minute, the second wide, strongly keeled, its length $\frac{3}{4}$ the diameter of eye. Pectorals of 12 rays, length 2 in head. Each ventral consists of a flat keeled spine followed by a minute ray. This species is known from 2 small mutilated specimens, both found on the beach near San Jose del Cabo, Cape San Lucas. The type was taken by John Xantus, about 1860, and recorded by Jordan & Gilbert as *Lepidopus caudatus*. The second, of about the same size ($5\frac{1}{2}$ inches), was taken by

Richard C. McGregor, in 1897. From the latter the above account was taken. The species differs from *Lepidopus caudatus* in the much shorter dorsal and longer anal. D. 103; A. 24. (Named for John Xantus de Vesey.)

Lepidopus caudatus, JORDAN & GILBERT, Proc. U. S. Nat. Mus. 1882, 358; not of EUPHRASEN. *Lepidopus xantusi*, GOODE & BEAN, Ocean. Ichth., 519, 1896; same type; no description.

Page 889. *Trichiurus lepturus* is recorded by Storer from Buzzards Bay (1840) and Wellfleet, Massachusetts (1845), and H. M. Smith records it from Woods Hole (1897).

Page 892. The synonymy at top of page under *Tetrapturus imperator* belongs to the footnote on same page.

Page 899. Add:

1286(a). OLIGOPLITES MUNDUS, Jordan & Starks, new species.

Head 4; depth $2\frac{4}{5}$; eye $4\frac{1}{4}$. D. V–I, 19; A. II–I, 20. Body deep and compressed. Length of head about $\frac{1}{6}$ greater than its depth at nape; eye equal to snout and to interorbital; maxillary extending considerably beyond vertical from hinder margin of eye, its length $1\frac{2}{3}$ in head; second suborbital not over $\frac{1}{2}$ as wide as lowest, and much shorter, thus forming a prominent notch in posterior margin of suborbital bones; a slight emargination in opercle in front of pectoral. Teeth small, sharp, in a band in each jaw, narrow in upper. Origin of soft dorsal midway between snout and base of caudal, the anal opposite; the anterior rays of both somewhat produced; second soft ray of each equal to head behind pupil, and equal to pectoral; ventrals equal to $\frac{3}{4}$ of pectorals, their inner margins fastened to body; caudal deeply forked, the middle rays $3\frac{1}{2}$ in longest, which are longer than head. Lateral line nearly straight, but forming a broad angle above pectoral. Color silvery on sides, becoming darker above; fins colorless. This species differs from *Oligoplites altus* in the much larger mouth and in having the suborbital bones notched posteriorly. *Oligoplites saliens* of the West Indies seems to be more elongate in body and with the suborbitals even behind as in *O. altus*. Pacific coast of tropical America.

This description is based on a specimen 11 inches long from San Juan Lagoon, Mexico, at the mouth of Ahome River, collected by the *Albatross*. Three other specimens from Algodones Lagoon, Mexico (*Albatross* Coll.), agree in every respect, except that 1 of them has but 4 free spines in front of dorsal.

Numerous other specimens have been since brought by Dr. Gilbert from Panama.

Oligophites mundus, JORDAN & STARKS, in JORDAN & EVERMANN, Check-List Fishes, 344. 1896, Mazatlan, Mexico; name only.

Page 909. The Californian species *Trachurus symmetricus* is probably a species distinct from *T. picturatus*, described from Madeira. The two forms have never been properly compared.

Page 912. The identity of *Hemicaranx amblyrhynchus* with *Caranx falcatus*, Holbrook, needs proof. The latter species, if distinct, may be described as follows:

1305(a). HEMICARANX FALCATUS (Holbrook).

Head 6 in total length; depth about 3. D. VII–I, 28; A. II–I, 25; C. 19; V. 5; P. 16; lateral line with 50 plates. Body oval, compressed; the head short, the facial outline descending in a gentle curve to snout, which is rounded though narrow. Eye large, in the middle third of the head, the posterior margin rather nearer snout than posterior margin of opercle; nostrils close together, nearly midway between eye and snout, and on a line within the orbit, the posterior larger, subround, the anterior ovoidal. Mouth small; each jaw with a single row of slender, conical teeth; a small patch of minute teeth on the vomer, and a small, narrow group of similar teeth on the palatines; tongue small, narrow, a few minute teeth near its base; pharyngeal bones armed with numerous card-like teeth, longer than those of the jaws. Soft dorsal long and low, the first 3 or 4 rays moderately elevated, the fin scaled at base; pectoral falcate, very long, extending to anterior third of soft dorsal; ventral small, very short, reaching beyond vent; anal shaped like the soft dorsal; caudal very long and widely forked, the upper lobe more than $\frac{1}{3}$ longer than the lower. Lateral line at first almost semicircular; at origin of soft dorsal descending to median plane, then straight; plates beginning with the soft dorsal increasing in size to the thirty-fifth, whence they decrease rapidly; scales minute, those of lateral line elongated quadrilateral, with 1 angle prolonged and rounded. Color, upper part of head and body above lateral line pale brown with slight bluish tint; lower jaw, opercle, and side yellowish; belly silvery, with a slight golden tint; anterior dorsal transparent; posterior transparent but with a yellowish tint; caudal yellowish. Known certainly only from Charleston, South Carolina.

Caranx falcatus, Holbrook, Ichth. South Carolina, 92, pl. 13, fig. 2, Charleston, South Carolina.

Page 914. Add:

1306(a). HEMICARANX ZELOTES, Gilbert, new species.

Head 4 to $4\frac{1}{5}$; depth $2\frac{2}{5}$ to $2\frac{3}{5}$. D. VII–I, 26 to 29; A. II–I, 23 to 25; P. 20 to 22; scutes about 52. Body regularly elliptical, its greatest depth about in middle of its length, exclusive of caudal peduncle. Head small; anterior profile more decurved, and hence the snout is blunter than in *H. atrimanus;* depth of head just behind eye about $\frac{5}{6}$ its length. Jaws subequal, tip of lower slightly projecting; maxillary narrow, not quite reaching anterior margin of pupil, about $3\frac{1}{6}$ in head ($3\frac{1}{8}$ to $3\frac{2}{5}$ in *atrimanus*). A single series of small, close-set, subequal teeth in each jaw; no teeth on vomer, palatines, or tongue. Orbit considerably greater than snout, $3\frac{1}{8}$ to $3\frac{1}{4}$ in head. Interorbital width (taken at anterior margin of orbit) slightly less than orbit. Occiput with an evident carina. Distance from snout to first dorsal spine greater than length of pectoral. Spinous dorsal very low, the highest spine considerably less than orbit (greater than orbit in *atrimanus*); a well-developed antrorse spine before the dorsal; soft dorsal and anal similar, not falcate, the rays decreasing in size from the first; highest ray of soft dorsal 2 to $2\frac{1}{2}$ in head; highest ray of anal about $2\frac{1}{8}$ in head; dorsal and anal depressible into a high sheath of scales, the last 3 or 4 rays

uncovered; caudal fin wide, well forked, the upper lobe the longer, the longest ray not quite $\frac{1}{4}$ total length of body; pectoral fin long, $3\frac{1}{8}$ to $3\frac{1}{2}$ in body ($2\frac{2}{5}$ to $2\frac{4}{7}$ in *atrimanus*); ventrals $2\frac{2}{5}$ to $2\frac{3}{5}$ in head. Scales as in *atrimanus;* lateral line with a very strong curve anteriorly, the height of the curve $2\frac{3}{8}$ to $3\frac{1}{2}$ in its length; its length $2\frac{1}{8}$ to $2\frac{1}{4}$ in the straight portion; entire length of straight portion with scutes, which are very small in front and behind; scutes considerably wider and lower than in *atrimanus*, the widest about $\frac{1}{2}$ diameter of orbit (about $\frac{1}{3}$ diameter of orbit in *atrimanus*). Coloration much as in *H. atrimanus*, but darker, and the fins without yellow; blackish olive above, dusky silvery below; top of head and snout black; spinous dorsal and the broad margins of soft dorsal and anal black; caudal dark, margined with black; pectorals very dark, black inside, the extreme lower rays light; a large jet-black blotch at base, on each side of pectorals, extending for about $\frac{1}{3}$ the whole length of the fin; axil black. Closely related to *Hemicaranx atrimanus*. Like it, it has a large jet-black area at axil and base of pectoral, and differs from it in the following characters: In having a shorter pectoral, shorter ventrals, profile of snout more rounded, a lower spinous dorsal, a shorter maxillary, a higher, shorter curve in lateral line, wider scutes, which are less sharply carinated, and darker fins. Panama. (ζελοτής, an imitator.)

Page 921. *Caranx crysos* and *Caranx pisquetus* are probably distinct species, the former ranging from New York to Florida, the latter from the West Indies to Brazil.

In *Caranx pisquetus* the pectoral fins are very long, as in the Pacific species *Caranx caballus*, from which we can not separate it. The species need further study.

Page 934. After *Vomer setipinnis* add:

1329(a). VOMER SPIXII (Swainson).

Head $2\frac{7}{8}$; depth $1\frac{3}{5}$; D. VI–I, 22; A. I, 18; eye $3\frac{3}{8}$ in head; maxillary $2\frac{1}{3}$; snout $1\frac{1}{3}$; caudal 1. Body very deep, in form much like *Selene orstedii;* profile very steep, almost vertical; snout slightly protruding. Mouth oblique, maxillary reaching to the vertical from front of eye; gill rakers 7 to 27, the longest a little more than $\frac{1}{2}$ eye. Lateral line strongly arched in front, the arch $1\frac{1}{4}$ the straight part; plates of lateral line little differentiated; pectoral falcate, as long or slightly longer than head; ventrals small, under base of pectorals. Color bluish above, sides silvery, fins except ventrals and anal dusky. Here described from specimens from Jamaica about 10 inches in length. These specimens are evidently different from the Northern *Vomer setipinnis* (= *Vomer browni*), the body in specimens of the same length being much deeper. It corresponds to the figur given by Agassiz of *Vomer browni*, this figure being the basis of *Vomer spixii* of Swainson. Probably all West Indian records of *Vomer setipinnis* belong to *Vomer spixii*. (Named for Jean Baptiste Spix, of Munich, naturalist and explorer.)

Platysomus spixii, Swainson, Class. Fishes, III, 250 and 406, 1839, Brazil; after Agassiz & Spix.

Vomer gabonensis, Guichenot, Ann. Soc. Maine et Loire, 1865, 42, Gaboon.

Page 938. To the synonymy of *Chloroscombrus chrysurus* add:

Seriola cosmopolita, CUVIER, Règne Animal, Ed. 2, vol. II, 1829, Gorea; after *Scomber chloris*, BLOCH.

Add the following species:

1334(a). CHLOROSCOMBRUS ECTENURUS, Jordan & Osgood.

Head $3\frac{7}{8}$; depth $2\frac{3}{4}$. D. VIII–I, 27; A. II–I, 26. Snout slightly shorter than eye, which is $3\frac{1}{4}$ in head. Chord of curved part of lateral line $1\frac{3}{8}$ in straight part. Depth of caudal peduncle 2 in its length, measuring from the base of the last dorsal ray to the base of the first caudal ray. Pectorals long and falcate, 3 in length; ventrals short, $2\frac{1}{5}$ in head, extending beyond the vent, which is situated in a groove in which these fins fit. Depth of head equal to or slightly less than its length; maxillary reaching anterior edge of eye, $2\frac{3}{5}$ in head. Lateral line unarmed; curve of ventral outline very slightly more pronounced than that of the dorsal; dorsal and anal fin sheaths well developed. Tips of upper spines and rays dusky; a black blotch at base of upper rays of caudal, and a black axillary and opercular spot. The species is closely related to *Chloroscombrus chrysurus*, the common species of the South Atlantic and Gulf States, which it evidently represents in the West Indies. The species *chrysurus* is deeper in every way, having a deeper body, a deeper head, and a deeper caudal peduncle. In *chrysurus* also the eye is larger, the mouth more nearly vertical, and the arch of the lateral line higher. When specimens of equal size from Florida and Havana are compared the characters are very evident. In 2 specimens, each $7\frac{1}{2}$ inches in length, from Havana and Florida, respectively, the depth of the body of the one is contained $1\frac{1}{4}$ times in that of the other, the depth of the head $1\frac{1}{4}$, the depth of the caudal peduncle $1\frac{1}{6}$, and the length of the eye $1\frac{1}{3}$. The names *chrysurus* (South Carolina), *latus* (Carolina), and *caribæus* (Texas) evidently all belong to the species of the United States coast. The type of *chloris* came from Acará, in Guinea, and *cosmopolita* of Cuvier was originally as a mere substitute for *chloris*. Until the African species can be examined, it is better not to use the name for either of the American forms. Probably *Chloroscombrus chloris*, when studied, will be found distinct from either. If not, that name would take the place of *ectenurus*. West Indies; known from Jamaica and Cuba. (ἐκτενής, extended; οὐρά, tail.)

Chloroscombrus ectenurus, JORDAN & OSGOOD, Proc. Ac. Nat. Sci. Phila. 1897, 101, Jamaica. (Coll. J. S. Roberts.)

The validity of *Chloroscombrus ectenurus* is still doubtful.

Page 942. After *Trachinotus falcatus* add:

1337(a). TRACHINOTUS RHOMBOIDES (Bloch).

Head $3\frac{1}{6}$; depth $1\frac{3}{5}$ in length; eye $3\frac{1}{3}$ in head. D. VI–I, 20; A. II–I 18. Back much elevated, but not angulated at origin of soft dorsal; end of snout not vertical, curved; head slightly concave at occiput. Maxillary to below anterior margin of pupil; eye on level of lower edge of premaxillary and axil of pectoral. Origin of soft dorsal behind tip of pectoral,

its lobe much elongated, extending to middle of caudal; lobe of anal reaching to below base of caudal; caudal lobes equal, $2\frac{1}{3}$ in body; pectoral rounded, $1\frac{1}{2}$ in head; ventrals $2\frac{3}{4}$ in head. Scales minute, large posteriorly near lateral line. Pale olive above, becoming silvery on belly; lobes of vertical fins dusky. This West Indian species is apparently different from the northern *Trachinotus falcatus* with which it has been confounded. *Trachinotus falcatus* seems to be confined to the coasts of the United States. In specimens of the same size the vertical fins are much higher in the West Indian species. (*ρόμβός*, rhomb; *εἴδως*, resemblance.)

Chætodon rhomboides, Bloch, Ichth., 1787, pl. 209, Martinique.

Page 945. After *Trachinotus paloma* insert:

428(a). ZALOCYS, Jordan & McGregor.

Zalocys, Jordan & McGregor, Rept. U. S. Fish Comm. 1898 (*stilbe*).

This genus is closely allied to *Hypodis*, Rafinesque (= *Lichia*, Cuvier), differing in the absence of a procurrent spine before the dorsal, and in the cultrate thoracic region. From *Trachinotus* it is distinguished by the same characters and also by the lower forehead and nonfalcate dorsal and anal fins. *Hypodis* is scarcely different from *Trachinotus*, the only tangible characters being the larger teeth, the low dorsal, and the less elevated forehead. *Porthmeus*, Cuvier (= *Lichia amia* and *L. radigo*) is a well-defined genus, distinguished by the large mouth and projecting lower jaw. (*ζάλη*, surge of the sea; *ἀκύς*, swift.)

1344(a). ZALOCYS STILBE, Jordan & McGregor.

Head $4\frac{1}{3}$; depth $2\frac{1}{2}$. D. VI–I, 26; A. II–I, 23. Body elliptical, deeper than in *Hypodis glaucus;* belly sharply compressed; ventral outline similar to that of dorsal; anterior profile of the head elevated and sharp, the eye being rather below than above its middle; eye 5 in head, with conspicuous adipose eyelid before and behind; posterior nostril much larger than anterior; vertically oblong maxillary broad, without supplemental bone, extending to pupil, $2\frac{5}{6}$ in head. Mouth moderate, oblique; each jaw with bands of villiform teeth; similar teeth on vomer, palatines, and tongue. Preopercle very broad; cheek moderate; suborbital narrow; preorbital very narrow, 4 in eye. No pseudobranchiæ. Gill rakers very long and slender, numerous. No procumbent spine before dorsal; spines low and separate, progressively higher; soft dorsal and anal each with a sheath of scales; first rays of dorsal very slightly elevated, $2\frac{1}{8}$ in head; anal without distinct anterior lobe, longest ray $2\frac{3}{4}$ in head; caudal peduncle long and slender; depth $3\frac{3}{4}$ in head; length below $2\frac{1}{8}$ in head; caudal fin widely forked; lobes long and slender, upper a little the longer, more than $\frac{1}{2}$ longer than the head and $2\frac{3}{4}$ in body; pectoral moderate, $1\frac{1}{8}$ in head; ventrals very small, $6\frac{1}{2}$ in head; snout $3\frac{3}{4}$ in head; premaxillary protractile. Color dark steel blue or blackish above; lower parts soiled white; axil and base of pectoral within jet-black; dorsal and anal each with a narrow whitish edging; caudal black, each lobe with a narrow whitish edging within. Body covered with small smooth scales, much as in *Trachinotus;*

lateral line undulate, very slightly arched anteriorly. Clarion Island; 1 specimen, 16 inches in length, known. (*στίλβη*, shining.)

Zalocys stilbe, JORDAN & MCGREGOR, Rept. U. S. Fish Comm. 1898, pl. 5, Clarion Island, Revillagigedo Archipelago. (Type, No. 11996, L. S. Jr. U. M. Coll. R. C. McGregor.)

Page 965. *Rhombus*, *Palometa*, and *Poronotus* should probably stand as distinct genera. The species placed in *Rhombus* in the text would then stand as follows:

1363. RHOMBUS PARU (Linnæus).

1364. RHOMBUS XANTHURUS (Quoy & Gaimard).

1365. PALOMETA PALOMETA (Jordan & Bollman).

1366. PALOMETA MEDIA (Peters).

1367. PALOMETA SIMILLIMA (Ayres).

1368. PORONOTUS TRIACANTHUS (Peck).

The identity of the South Atlantic Coast *Rhombus alepidotus* with the West Indian *Rhombus paru* is very doubtful.

Page 973. The genus *Acrotus*, Bean, represents a family distinct from *Icosteidæ*.

Family CXXXVI(a). ACROTIDÆ.

Two additional specimens of *Acrotus willoughbyi* have lately come to light—the one from Port Townsend, the other from Monterey.

After *Acrotus willoughbyi* insert:

Family CXXXVI(b). ZAPRORIDÆ.

Body robust, moderately compressed, the back not elevated, the belly not carinate. Body covered with small adherent cycloid scales, which cover the membranes of all the fins except the distal third, as also the gill membranes, lower jaw, cheeks, opercles, and nuchal region. No lateral line; no spinules. Head short, the nape not elevated, the forehead broad and abruptly convex in profile; eye moderate, placed high; preopercle, parietal region, and region about eye with very large open mucous pores. No spines on head; edges of membrane bones of head covered with thick scaly skin. Mouth moderate, terminal, oblique, its cleft mainly anterior; upper jaw protractile, but not movable; maxillary rather narrow, simple; lower jaw very heavy, its thick lip projecting beyond upper jaw. Teeth alike in both jaws, rather strong, blunt, even, close set, forming a uniform cutting edge; no teeth on vomer, palatines, or tongue, the tongue very thick. Lower pharyngeals narrow, with bluntish teeth, those on the edge larger; upper pharyngeals rather large, with small, blunt, velvety teeth; no distinct tooth-like processes in the œsophagus; pseudobranchiæ present; gill rakers very slender and flexible, rather short; gills 4, a large slit behind the fourth; gill membranes separate, free from the isthmus; opercle adnate to shoulder girdle above its angle; coracoids not largely developed. Pectoral fin long, rounded, attached a little nearer

ventral than dorsal outline; ventrals wholly wanting. Dorsal fin beginning above gill opening, composed entirely of simple inarticulate rays or spines, these moderately flexible, attached to the membrane to their tips, and all except the first and last of about equal length. Caudal peduncle short and stout, not contracted, the large caudal subtruncate or rounded at tip, and without procurrent rays; vent nearly median. Anal much shorter than dorsal, somewhat higher, and composed of soft rays, subequal in length. Skeleton rather limp and flexible, but much less so than in *Icosteus*.

445(a). ZAPRORA, Jordan.

Zaprora, JORDAN, Proc. Cal. Ac. Sci. 1896, 202 (*silenus*).

Characters of the genus included above.

This genus bears some resemblance to *Icichthys*, but differs in the stout caudal peduncle, absence of ventrals and lateral line, and in the form and structure of the head. Among the genera known to us it seems to come nearest to *Icichthys*, and it might be placed among the *Icosteidæ* were it not for the presence of pharyngeal teeth. (ζά, an intensive particle; πρώρα, prow.)

1372(a). ZAPRORA SILENUS, Jordan.

Head $5\frac{2}{3}$ in length to base of caudal; depth $4\frac{1}{4}$. D. LVI; A. 27; P. 20 to 22; C. 22; scales about 200–85. Greatest thickness of body about $\frac{2}{3}$ its depth; length of caudal peduncle $1\frac{2}{3}$ in its least depth, which is $1\frac{9}{10}$ in head. Eye $5\frac{1}{3}$ in head; snout $5\frac{1}{3}$; interorbital space 3; maxillary $2\frac{2}{3}$, ending under front of pupil; mandible $2\frac{1}{2}$, its depth $4\frac{2}{3}$; teeth about $\frac{15}{30}$ on each side; lips, snout, and bones about eye naked; rest of head covered with small scales. Lower jaw with a thick lip, slightly fringed on its edge, and with a mesial frenum; the rounded tip entering the profile when the mouth is closed. Three large pores on each ramus of mandible; behind these 3 others in a line on horizontal limb of preopercle; 3 on vertical limb; 2 close together in front of eye; 1 near the nostrils, so similar to them that there seem to be 3 nasal openings; 7 on suborbitals; 4 in 2 rows behind eye; 1 above eye, and before upper edge of preopercle; a horizontal row of 5 along temporal region, the last and largest of all in opercular flap above gill opening; 1 at vertex; 1 between vertex and eye, and 2 on each side of nape. Gill rakers 8 + 20, the longest $\frac{1}{2}$ eye. No trace of lateral line. Scales small, resembling those of a salmon, covering the membranes of all the fins on the basal two-thirds. Pectoral as long as head, its base $2\frac{1}{3}$ in head; longest dorsal spine $1\frac{5}{6}$; caudal $1\frac{1}{10}$; longest anal ray $1\frac{3}{4}$. Color in spirits uniform dusky, without markings on the body, the belly pale, and the side of the head irregularly blotched with lemon yellow, apparently bright in life, and brightest about the pores of the head. Coast of British Columbia; only the type, 29 inches long, known. (σειλένος, a drunken demigod, covered with slime, in allusion to the open mucous pores.)

Zaprora silenus, JORDAN, Proc. Cal. Ac. Sci. 1896, 203, pl. 20, Nanaimo, Vancouver Island. (Type in Provincial Museum at Victoria. Coll. H T. Stainton.)

Page 982. Prof. Harrison Garman records *Elassoma zonatum* from Waccamaw River, Whitesville, North Carolina, and Little Pedee River, South Carolina. Vertebræ 29; scales 34 to 36; D. IV, 9; A. III, 5.

Page 1019. Under *kk* read: "gill membranes narrowly or broadly connected."

Page 1047. Before *Ulocentra* insert:

1436(a). COTTOGASTER CHENEYI, Evermann & Kendall.

Head 4; depth 6; eye 4 in head; snout 4; maxillary 3⅓; interorbital width 5½. D. XI–12; A. II, 8; scales 7–56–6. Body rather stout, heavy forward, compressed behind; head heavy; mouth moderate, slightly oblique, lower jaw included, maxillary reaching front of pupil; premaxillaries protractile. Cheeks, opercles, breast, and nape entirely naked; scales of body large and strongly ctenoid; lateral line complete, straight; median line of belly naked anteriorly, with ordinary scales posteriorly. Fins large; dorsals separated by a space equal to ½ diameter of eye; origin of spinous dorsal a little nearer origin of soft dorsal than tip of snout, its base about equal to length of head; longest dorsal spine 2½ in head, the outline of the fin gently and regularly rounded; soft dorsal higher than spinous portion, the second to tenth rays about equal in length, scarcely 2 in head, the first, eleventh, and twelfth rays but slightly shorter than the others; anal moderate, its origin under base of third dorsal ray, the spines slender, the second a little longer than the first, whose length is 3¼ in head; longest anal rays about 2⅕ in head; caudal lunate, the lobes more produced and pointed than usual among darters; pectorals long and pointed, the middle rays longest, about 1⅕ in head, reaching tips of ventrals; ventrals well separated, not nearly reaching vent, the longest rays 1¼ in head. Color in alcohol, back dark brownish, covered with irregular spots and blotches of darker; side with about 8 or 9 large dark spots lying on the lateral line; belly pale; top of head dark; snout black; lower jaw and throat dark; a broad black line downward from eye to throat; cheek and opercles rusty; spinous dorsal crossed by a median dark line; ventrals blue black; other fins pale, but dusted with rusty specks. An examination of the 14 cotypes shows some variation in the species. In 2 examples there is a well-developed frenum, rendering the premaxillaries nonprotractile, and in a third specimen the frenum is partially developed; in some individuals the origin of the spinous dorsal is exactly midway between the tip of snout and origin of soft dorsal. The females and immature males are less highly colored than the adult male described above. Length 1¾ to 2¼ inches. This species is most closely related to *Cottogaster shumardi*, from which it may be readily distinguished by the shorter snout, the naked cheeks and opercles, the smaller soft dorsal, the smaller anal, and the different coloration. Fifteen examples of this interesting darter were obtained July 18, 1894, by Messrs. Evermann & Bean in the Racket River near Norfolk, St. Lawrence County, New York. It did not seem to be very common, as only 15 examples resulted from numerous hauls of the collecting seine. (Named for Mr. A. Nelson Cheney, State fish-culturist of New York, in recognition

of his valuable contributions to our knowledge of the food and game fishes of that State.)

Cottogaster cheneyi, EVERMANN & KENDALL, Bull. U. S. Fish Comm. 1897 (Feb. 9, 1898), 129, pl. 8, fig. 8, Racket River near Norfolk, New York. (Type, No. 48781. Coll. Evermann & Bean.)

Page 1049. After *Ulocentra gilberti* add:

1488(a). ULOCENTRA MEADIÆ, Jordan & Evermann, new species.

Head $3\frac{1}{5}$; depth $4\frac{2}{3}$; eye $3\frac{1}{2}$ in head; snout $3\frac{2}{3}$; interorbital 5. D. XII–12; A. II, 7; scales 7–48–6. Body rather heavy, somewhat fusiform; head large; snout blunt, decurved, profile rising abruptly to interorbital, thence nearly horizontal to origin of dorsal, from which it descends gently in a straight line to caudal peduncle; opercular spine small but sharp; mouth low, horizontal, rather large, the maxillary reaching vertical at front of orbit; premaxillaries protractile; branchiostegal membranes not connected, free from the isthmus; ventral fins close together, the space separating their bases about $\frac{1}{2}$ diameter of orbit; fins all moderate; distance from tip of snout to origin of spinous dorsal 3 in body; spinous and soft dorsals close together, the space separating them about 2 in orbit; longest soft dorsal rays $1\frac{1}{5}$ in head, about equaling those of anal; the two anal spines of about equal length, the first the stouter; pectorals long, longer than head, their tips passing those of ventrals but not reaching vent; ventrals short, $1\frac{1}{6}$ in head; caudal slightly lunate when expanded. Scales rather large, strongly ctenoid; cheeks and breast naked; opercle scaled above, naked below; nape scaled; lateral line complete, straight; ventral line of body covered with ordinary adherent scales. Color in alcohol, yellowish or olivaceous above and on sides, the back with 6 dark saddle-like blotches, the first just anterior to origin of spinous dorsal, the second under the fifth and sixth spines, the third under the last two spines, the fourth under the sixth and seventh soft rays, the fifth just posterior to the last dorsal ray, and the sixth, which is quite small, upon the caudal peduncle at the base of the caudal fin; sides blotched with dark, 6 to 8 larger dark blotches along side just below lateral line, sometimes more or less continuous with the dark dorsal blotches; a dark blotch at base of middle caudal rays; belly pale; top of head dark; a dark spot at lower posterior angle of eye and a smaller one back of it on upper edge of opercle; a dark band downward from eye; opercle dark; upper lip dark, interrupted by a light line at the symphysis; spinous dorsal pale, with a broad dark band through its lower third; soft dorsal crossed by 3 or 4 irregular lines of dark specks; caudal with about 4 broad dark cross bars; other fins pale. Length 2 inches. This species somewhat resembles *U. gilberti*, but differs from it in the larger head, stouter body, larger scales, naked cheeks, larger mouth, and in other respects. Known only from Indian Creek, basin of Powell River, east Tennessee, where 3 examples were collected October 17, 1893. (Named for Mrs. Meadie Hawkins Evermann.)

Ulocentra meadiæ, JORDAN & EVERMANN, new species, Indian Creek, Cumberland Gap, Tennessee. (Type, No. 48903. Coll. Dr. R. R. Gurley.)

Page 1051. To the synonymy of *Ulocentra simotera* add:

Etheostoma duryi, HENSHALL, Journ. Cin'ti Soc. Nat. Hist., April, 1889, 32, small tributary of Tennessee River at Whiteside, Tennessee. (Type in Mus. Cin'ti Soc. Nat. Hist. Coll. Charles Dury.)

Page 1089. To the synonymy of *Etheostoma cæruleum* add:

Etheostoma formosa, HENSHALL, Journ. Cin'ti Soc. Nat. Hist., April, 1889, 32, small tributary of Tennessee River at Whiteside, Tennessee. (Type in Mus. Cin'ti Soc. Nat. Hist. Coll. Charles Dury.)

Page 1109. Add:

1501(a). APOGON ATRICAUDUS, Jordan & McGregor.

Head $2\frac{1}{2}$; depth 3. D. VI–I, 9; A. II, 8; scales largely ctenoid; eye $3\frac{1}{2}$ in head; second dorsal spine stoutest, about 2 in head; gill rakers 17, moderate. Body similar in shape to *A. retrosellus*. Jaws reaching to posterior border of eye, $1\frac{5}{6}$ in head. Pectoral reaching to opposite front of anal, $1\frac{3}{8}$ in head. Color rosy, darkened with dusky points; more or less olivaceous above; head and throat verging on orange; first dorsal black; second dorsal rosy; caudal dusky, more or less flushed with rosy, other fins paler; no black spot on head or on base of caudal, there being no definite markings anywhere except the dusky red of the tail. West coast of Mexico. Numerous specimens collected at San Benedicto, Socorro, and Clarion islands. Usual length 3 to 4 inches. (*ater*, black; *cauda*, tail.)

Apogon atricaudus, JORDAN & MCGREGOR, Rept. U. S. Fish Comm. 1898, Socorro, Clarion and San Benedicto islands. (Coll. R. C. McGregor.)

Page 1125. *Centropomus affinis* can not be separated from *C. ensiferus*.

Page 1148. To the synonymy of *Epinephelus* add:

Phrynotitan, GILL, Stand. Nat. Hist., III, 255, 1885 (*Batrachus gigas*).

Page 1150. In the key under *dd*, read: Lower jaw strongly projecting.

Page 1156. Add:

1551(a). EPINEPHELUS NIPHOBLES, Gilbert & Starks.

Head $2\frac{2}{5}$ in body; depth $2\frac{1}{2}$. D. XI, 14; A. III, 9; scales 16–116–40; eye 5 in head; maxillary 2; third dorsal spine $2\frac{3}{4}$; middle dorsal rays $2\frac{1}{3}$; highest anal rays 2; third anal spine $3\frac{1}{10}$; pectoral $1\frac{5}{7}$; ventrals $1\frac{2}{3}$; caudal $1\frac{2}{3}$. Form rather robust, moderately compressed; dorsal outline uniformly curved from tip of snout to caudal peduncle; mouth large, the maxillary reaching to below posterior orbital rim; lower jaw strongly projecting; teeth conical and sharp, in 1 or 2 bands at sides of jaws, 3 or 4 in front; upper jaw with a rather strong canine on each side of front; snout longer than eye; nostrils close together, the posterior one the larger, a little in front of the vertical from front of eye, the anterior in a short, wide tube with a flap behind; vertical and horizontal limbs of preopercle meeting at right angles, its edge with blunt serræ, those at angle enlarged; opercle with 3 flat spines before the flap; gill rakers moderate, nearly $\frac{1}{2}$ eye, 8+16 in number. Top of head, orbitals, maxillary, and mandible, naked; fine scales on cheeks and opercles; scales on body ctenoid;

fins without scales. Dorsal beginning a little in front of the vertical from pectoral base, the third spine a little the highest, but the ones behind it not much shortened; soft dorsal higher than spinous, its outline rounded; pectoral rounded behind, reaching to below the base of eighth dorsal spine; third anal spine the longest, not nearly so long as the soft rays, the anal fin similar in shape to the soft dorsal; ventrals reaching past vent, scarcely to front of anal, their ends rounded, as are all the fins; caudal broadly rounded. Color in spirits brownish red, sides with clear-cut, distinct, white spots about as large as pupil, about 6 at base of dorsal, 6 or 7 along lateral line, following its arch, a horizontal series of 4 extending back from opercular flap, about 3 from base of pectoral following curve of ventral outline, 2 at base of anal, 1 behind lower edge of caudal peduncle and 1 above anus; a well-marked streak above maxillary following its outline; lips colored like rest of head; dorsal dusky, with vague white spots; ventrals and anal nearly black, with a reddish tinge; anal with a narrow white border below; pectoral and caudal uniform yellowish. Magdalena Bay, Lower California; only the type, 6 inches long known. (*νιφοβής*, snowed over, from the white spots.)

Epinephelus niphobles, Gilbert & Starks, Proc. U. S. Nat. Mus. 1896, 442, Magdelena Bay, Lower California. (Type, No. 47582. Coll. *Albatross*.)

Page 1164. Species 1558 should probably be called *Alphestes chloropterus* (Cuvier & Valenciennes). The name *afer*, given to a specimen from Guinea, may belong to some other species.

Page 1168. Add:

1560(a). DERMATOLEPIS ZANCLUS, Evermann & Kendall.

Head $2\frac{3}{4}$; depth $2\frac{3}{16}$; eye 8 in head; snout $3\frac{1}{3}$; maxillary 3; mandible 2. D. XI, 19; A. III, 10; scales difficult to count, but about 30–130–35, those above lateral line counted obliquely backward and downward from origin of dorsal, those below from origin of anal upward and forward to lateral line. Branchiostegals 8; gill rakers 8 + 12, short and stout, the longest $1\frac{3}{8}$ in orbit. Body stout, compressed, oblong-elliptical, the dorsal and ventral outlines about equally curved; head moderate, the profile rising from tip of snout to origin of dorsal fin, thence descending in a regular, gentle curve to caudal peduncle; a depression above nostrils and a slight one on nape; interorbital very narrow, equal to orbit; mouth moderate, somewhat oblique; premaxillaries protractile; maxillary broad at tip, reaching vertical at posterior edge of the pupil; supplemental bone well developed; lower anterior edge of maxillary covered by the broad dermal flap of the premaxillary; eye small, high up; nostrils close together and close to eye, the anterior small and round, the posterior oblong-oval, much larger than the other. Small cardiform teeth on each jaw, those in front movable, scarcely canine-like; similar teeth on vomer and a long, narrow band on each palatine. Preopercle coarsely serrate, the serræ short and blunt, more or less obscured by the skin; opercle with a broad dermal border, somewhat produced at lower angle. Fins all large; origin of dorsal slightly in advance of base of pectoral, its distance from tip of snout equal

to length of head; third dorsal spine longest, its length about 2⅔ in head or 2½ times length of first ray; interspinal membranes of the spinous dorsal deeply incised, the anterior portion of each somewhat produced beyond its spine; soft dorsal high, the middle rays longest, 1⅔ in head, the anterior portion of the fin gently convex, the posterior slightly concave; pectoral short, broad, and rounded, barely reaching origin of anal, the length 1½ in head; ventral pointed, the second and third rays longest, 1⅓ in pectoral, the fin somewhat falcate; anal fin strongly falcate, the fourth and fifth rays longest, longer than pectoral, 1¼ in head, 2⅓ times length of last anal ray; second anal spine short, 5½ in head; caudal shallowly lunate, the lobes 1½ in head. Scales small, smooth, and thin, closely but irregularly imbricated; nape, opercles, and cheeks scaled, snout and lower jaw naked; bases of all the fins except the ventrals densely scaled; lateral line beginning at upper angle of opercle, gently arched above pectoral fin, following approximately the curvature of the back and on median line of caudal peduncle. General color of body in life brown, with large, irregular blotches of dirty white on back and upper part of sides, these blotches with small rusty spots; lower part of sides, belly, and caudal peduncle with irregular whitish spots; belly brassy brown; snout and nape with numerous small, round dark spots; cheek with large blotches of whitish overlaid with black and brassy spots; lips whitish, with dark spots; spinous dorsal blotched with white, olivaceous and black; soft dorsal brown, with numerous white spots and a few black ones, the posterior rays tipped with white and orange; anal olivaceous, with irregular white spots, greenish at edge, the produced rays black toward distal ends; pectoral dark olivaceous, with greenish white splotches, the edge yellowish; ventral rays greenish white, the membranes black; inside of mouth white; eye brown. Related to *D. inermis* (Cuvier & Valenciennes), but differing notably from that species in the shorter, stouter gill rakers, the emarginate caudal, the shorter anal spines, and the strongly falcate anal fin. Length 20 inches. Key West; only the type known. (ζάγκλον, a scythe or sickle, from the falcate anal fin.)

Dermatolepis zanclus, Evermann & Kendall, Bull. U. S. Fish Comm. 1897 (Feb. 9, 1898), 129, pl. 8, fig. 9, Key West, Florida. (Type, No. 48843. Coll. Drs. Evermann & Kendall.)

Page 1186. Add:

1576(a). **MYCTEROPERCA HOPKINSI,** Jordan & Rutter.

Head 2⅔; depth 4½. D. XI, 15; A. III, 11; scales about 125; eye 6 in head, 1⅓ in snout. Body long, not much compressed; angle of preopercle sharply serrate; gill rakers 6 + 9, counting rudiments; nostrils close together, the posterior larger, with a horizontal septum across base; profile concave above nostrils; maxillary nearly to posterior margin of eye, 2⅙ in head; lower jaw projecting; 2 anterior canines of upper jaw very strong; third and fourth dorsal spines longest; posterior portion of anal truncate; caudal concave. Pectorals 2, ventrals 2⅓, and caudal 1½ in head. Color of alcoholic specimen nearly uniform brownish, side of jaws paler; soft dorsal, anal, ventrals, and caudal with a narrow pale edging,

these fins otherwise brownish olive, with a subterminal band of black; pectorals pale, darker in middle. Allied to *Mycteroperca calliura*, differing in having fewer gill rakers, more slender body, smaller scales, and a less lunate caudal. Jamaica; only 1 specimen, 6 inches long, known. (Named for Timothy Hopkins.)

Mycteroperca hopkinsi, JORDAN & RUTTER, Proc. Ac. Nat. Sci. Phila. 1897, 105, Jamaica. (Type, No. 5073, L. S. Jr. Univ. Mus. Coll. J. S. Roberts.)

Page 1187. Insert:

1576(a). MYCTEROPERCA BOULENGERI, Jordan & Starks.

Head $2\frac{4}{5}$ in length; depth $2\frac{5}{6}$. D. XI, 14 or 15; A. III, 9 or 10; scales about 90, 20 above and 42 below; snout $3\frac{1}{2}$ in head; maxillary $2\frac{1}{5}$; eye $5\frac{1}{2}$; pectoral $1\frac{3}{4}$; ventral $1\frac{5}{6}$; longest anal ray $1\frac{3}{4}$; caudal $1\frac{3}{5}$; longest dorsal spine $2\frac{1}{2}$; gill rakers short, about 6 + 17, the longest about $\frac{3}{5}$ eye; longest dorsal ray 2 in head. Body short and deep, compressed; head moderate, compressed, its profile not steep, nearly straight, a depression before eye. The supraoccipital and temporal crests are high, the supraoccipital crest extending to the posterior margin of orbit; the temporal crests are parallel to each other, and extending to pupil; interorbital space concave. Upper canines moderate, the lower quite small. Nostrils small, well separated, the anterior slightly larger. Lower jaw very strongly projecting; maxillary reaching opposite posterior edge of pupil. Preopercle slightly notched, the angle slightly salient, with enlarged teeth. Dorsal not deeply notched, the fourth spine not much elevated; second dorsal high, not long, its angle not rounded; caudal scarcely lunate, the upper lobe long, the lower truncate; anal very high, strongly elevated, its posterior border incised, the anterior rounded; pectoral and ventral moderate. Scales smoothish, not very small. Color olive gray, covered everywhere with oblong irregular markings of black, between which the ground color forms rivulations; gray lines radiating from the eye; a black blotch below maxillary; pectoral olive yellow; other fins blackish, clouded with pale; first dorsal with faint small black spots. Mazatlan, Mexico. (Named for George Albert Boulenger, ichthyologist of the British Museum, in recognition of his epoch marking work on the Percoid fishes.)

Mycteroperca boulengeri, JORDAN & STARKS, Fishes Sinaloa, 445, pl. 38, 1895, Mazatlan, Sinaloa, Mexico. (Type, No. 1621, L. S. Jr. Univ. Mus. Coll. Hopkins Exped. to Sinaloa.)

Page 1235. The original type of *Lobotes* is *surinamensis*, not *erate*.

Lobotes erate is a species distinct from *L. surinamensis*, inhabiting the coasts of India and China. *Lobotes farkhari* and *L. incurvus* are probably identical with *L. erate*, and all 3 should be erased from the synonymy of *L. surinamensis*.

Page 1236. After *Lobotes surinamensis* add:

The *Lobotes* of the Pacific coast of Central America is distinguished from the other known species, *L. surinamensis* and *L. erate*, by the small

size of the preopercular serrations, those at the angle not elongated and spine-like, even in the young. The following description is furnished by Dr. Gilbert:

1623(a). LOBOTES PACIFICUS, Gilbert, new species.

(BERRUGATE.)

Head $2\frac{3}{4}$ in length; depth $2\frac{1}{5}$ to $2\frac{1}{10}$ (to base of caudal rays); depth of caudal peduncle $2\frac{1}{2}$ in head. D. XII, 15; A. III, 11; pectoral 15. Scales 11-46 (+6 on base of caudal)-18; vertebræ 12 + 12; Br. 6. Body more elongated than *L. surinamensis,* agreeing in this respect with *L. erate,* the depth less than $\frac{1}{3}$ the length. Upper profile deeply concave at occiput, thence strongly convex to front of dorsal; head shorter and narrower than in *L. surinamensis,* the interorbital width but slightly longer than snout, $3\frac{9}{10}$ to 4 in head ($3\frac{1}{8}$ to $3\frac{2}{3}$ in head in *L. surinamensis*). Eye small, $6\frac{2}{3}$ to $7\frac{1}{3}$ in head, 2 or $2\frac{1}{10}$ in interorbital width. Mandible strongly protruding, but without symphyseal knob; maxillary narrow, not concealed in closed mouth, its tip reaching vertical from middle of pupil, $2\frac{4}{7}$ to $2\frac{9}{10}$ in head. Upper jaw with a moderate villiform band of teeth, in front of which is a single series of conical, close-set canines; lower jaw with a single series, similar to outer series of upper jaw, and behind them a very narrow band of villiform teeth, which grow slightly larger toward symphysis; palate toothless. Posterior margin of preopercle vertical, the angle protruding but little in the young. In 5 young examples, 7 to 11 inches long, the preopercular teeth are fine, acute, short, and inconspicuous, about as in species of *Pomadasis.* They increase but little in size toward the angle, where they are never spine-like; on lower limb they are perceptible only in the immediate vicinity of the angle, the remainder of the horizontal limb being entire. In the adult the vertical limb is finely and evenly toothed, the angle and lower limb slightly roughened or entire; opercle with 2 short spinous points, behind the lower of which a narrow tongue-shaped process of the subopercle extends to near the edge of opercular membrane; humeral process very weakly toothed, contrasting with the strong serrate condition in *L. surinamensis.* Gill rakers short, $2\frac{1}{2}$ in eye in young, comparatively shorter in adults, 6 on vertical limb, all but one of which are broad, firmly fixed tubercles, 14 on horizontal limb, the anterior 2 or 3 tubercular. Spinous dorsal low, with gently rounded outline; notch between dorsals shallow, the eleventh spine $\frac{2}{3}$ the length of the longest, which is contained 2 to $2\frac{1}{8}$ times in head in the young, 3 times in adults; when declined the spines are partially received within a scaly grove; soft dorsal, anal, and caudal with dorsal portions densely scaled and with series of scales running up on membrane to beyond middle of fin; soft dorsal and anal of equal height, forming bluntly rounded lobes, the longest rays of which are about $\frac{1}{3}$ head in adults, $1\frac{1}{2}$ to $1\frac{3}{4}$ in young; third anal spine about $\frac{1}{2}$ length of longest ray; pectorals shorter than ventrals, 2 to $2\frac{1}{8}$ in head; ventrals $1\frac{1}{2}$ in head in young, shorter in adults. Scales less strongly ctenoid than in *L. surinamensis:* tubes of lateral line mostly simple, occasionally with 1 to 3 branches. Color grayish or brownish, with plumbeous or silvery reflections. The youngest examples show faintly the dark streaks so con-

spicuous in young of *L. surinamensis*, viz, a pair running backward from interorbital space; a pair from upper posterior border of eye converging toward front of dorsal, and a broader band from eye downward and backward across cheek; soft dorsal, anal, and caudal uniform blackish, or the caudal with an ill-defined lighter edge; pectorals translucent; ventrals blackish. Abundant at Panama, where it is known as *Berrugate*.

Lobotes auctorum, STEINDACHNER, Ichth. Beitr., IV, 6, 1875; not of GÜNTHER.
Lobotes surinamensis, JORDAN & GILBERT, Bull. U. S. Fish Comm., II, 1882, 110; GILBERT, l. c., 112; JORDAN, Proc. U. S. Nat. Mus. 1885, 378; not *L. surinamensis* of BLOCH.
Lobotes pacificus, GILBERT, Fishes of Panama, 1898 MS., Panama. (Type, No. 5883, L. S. Jr. Univ. Mus. Coll. Gilbert.)

Page 1238. After *Priacanthus cruentatus* add the following:

1625(a). PRIACANTHUS CAROLINUS, Lesson.

This species is very close to *Priacanthus cruentatus*, distinguished by the larger spine on preopercle, which reaches the edges of the opercle and is 2½ in eye; that of *P. cruentatus* not reaching opercle and measuring 4 in eye, its edge less rough. Body a little deeper than that of *P. cruentatus;* depth of the latter 3 in the length. In *P. carolinus* the depth is 2⅔ in the length; caudal truncate. In color and general appearance the 2 species are similar. The distinctness of this species from *P. cruentatus* is very doubtful. Abundant at Clarion Island, where it was taken by Mr. R. C. McGregor. (*carolinus*, from Caroline Islands.)

Priacanthus carolinus, LESSON, Voyage Coquille, Poiss., 204, 1826, **Caroline Islands.**
Priacanthus schlegeli, HILGENDORF, Sitzgber. Ges. Naturf. 1879, 79, **Japan.**

Page 1262. In the first line of the description of *Neomænis vivanus* read: Head 2⅞ to 2¾; depth 2⅓ to 3. D. X, 14; A. III, 8 or 9; eye 4¼ in head; scales (7) 8–72–17, 50 pores.

Page 1264. After *Neomænis vivanus* add:

1639(a). NEOMÆNIS HASTINGSI, Bean.

(Bermuda Silk Snapper.)

Head 3; depth 3; least depth of caudal peduncle 9 in length of type to caudal base. D. X, 14; A. III, 8; V. I, 5; P. 16; scales 8 or 9–65–17. Maxillary reaching scarcely past front of eye, 3⅛ in head. Vomerine teeth in an arrow-shaped patch, with a backward extension which is fully ⅓ as long as the eye; canines in upper jaw very feeble; 2 or 3 posterior teeth of mandible are weak canines; 7 rows of scales on cheeks, 9 rows on gill cover. Least interorbital width equal to eye, which is 1⅓ in snout and 4 in head. Gill rakers 7 + 9, the one in the angle conspicuously longest, about 2 in eye. First dorsal spine 7 in head; fifth and longest spine about 3 in head; last dorsal spine equal to eye in length; longest ray of soft dorsal equal to maxillary, or 3 in head; first anal spine 8, the second and third about 4 in head, the second slightly longer than third; anal base nearly 2½ in head; third and longest anal ray about equal to anal base; pectoral extending to vent; ventral not reaching vent by a space ½ as long as the eye. Colors in life, ground color vermilion, the upper parts over-

laid with coppery brown, lower parts vermilion; 4 or 5 narrow golden stripes below lateral line; caudal dark brown with a narrow black margin; anal dusky, the spines and the membranes of last 2 rays pale; a narrow black blotch at pectoral base; ventral pale, somewhat mingled with dusky; membranes of spinous and soft dorsal uniformly dark; snout copper color; eye lemon yellow; pupil blue black; many scales, especially on front of body, with a minute brown dot at base; brownish spots on scales forming many oblique streaks above lateral line. Some living examples show a faint dark lateral blotch much like that of *N. synagris*, and similarly placed. In spirits the body is pink with the upper parts brownish; the dusky color remains on the anal and the black blotch at base of pectoral; black margin of caudal becoming merged with the general dark color of the fin. (Bean.) Most closely related to *N. vivanus*. Length of type 11½ inches. Bermuda, where numerous specimens were obtained in 1897. (Named for General Russell Hastings of Soncy, Bermuda.)

Neomænis hastingsi, BEAN, Bull. Amer. Mus. Nat. Hist., x, Article III, 45, 1898, Bermuda.

Page 1290. Under *g* read: Anal fin short, its rays III, 7 to III, 13.

Page 1413. In first line of description of *Cynoscion phoxocephalus* for "A. III, 10" read "A. II, 10."

Page 1416. In last line of description of *Sagenichthys ancylodon* for "companion" read "comparison."

Page 1605. Instead of *Chlorichthys* read:

640. THALASSOMA, Swainson.

Thalassoma, SWAINSON, Nat. Hist. Class. Fishes, II, 224, 1839 (*purpureus*).
Julis, GÜNTHER, Cat., IV, 179, 1862; not of CUVIER & VALENCIENNES.

The species of *Thalassoma* (*pavo, unimaculatus, bifasciatus*) examined have 3 anal spines, as is the case with the American species referred to *Chlorichthys*. The first spine, small and hidden in the skin, is easily overlooked. There is therefore no distinction between *Thalassoma* and *Chlorichthys*, and all the American species must be referred to the former genus.

The species will stand as follows:

2014. THALASSOMA LUCASANUM (Gill).
2015. THALASSOMA SOCORROENSE, Gilbert.
2016. THALASSOMA NITIDUM (Günther).
2017. THALASSOMA NITIDISSIMUM (Goode).
2018. THALASSOMA STEINDACHNERI (Jordan).
2019. THALASSOMA BIFASCIATUM (Bloch).
2020. THALASSOMA GRAMMATICUM, Gilbert.
2021. THALASSOMA VIRENS, Gilbert.

Page 1670. In *Pomacanthus* (*P. paru*, species examined by Mr. E. C. Starks) and *Chætodon* the air bladder is wholly contained in the body cavity, while in *Holacanthus* and *Angelichthys* (*A. ciliaris* species examined) it is posteriorly separated from the body cavity. The 2 latter genera con-

stitute the subfamily *Holacanthinæ*, distinct alike from *Chætodontinæ* and *Pomacanthinæ*.

Page 1717. *Ceratacanthus*, including *Osbeckia*, should stand as a valid genus distinguished from *Alutera* by the convex or lanceolate caudal. The species will then stand as follows:

2135. CERATACANTHUS SCHŒPFII (Walbaum).
2136. CERATACANTHUS PUNCTATUS (Agassiz).
2137. CERATACANTHUS SCRIPTUS (Osbeck).
2138. ALUTERA MONOCEROS (Osbeck).

Page 1741, line 17, read: Swainson takes "κάνθα" to mean spine, not "ἄκανθα," which is the correct word for spine. There is no classical warrant for *Cantherines* and *Canthigaster*, unless derived from κάνθος, the ass.

Page 1776. In sixth line from bottom read "increased" for "self."

Page 1786. Note on *Sebastodes rufus*:

This species is ovate in form, like *S. oralis*, from which it differs in color and form of mouth and head. Its depth is $2\frac{2}{3}$ in length, not $3\frac{3}{4}$, as stated (through misprint) by Dr. Eigenmann. A fine specimen before us was taken by Dr. Gilbert off San Diego.

Page 1790. The type number of *Sebastodes hopkinsi* is 2282, not 2286.

Page 1795. The subgenus *Zalopyr* does not include *Sebastodes atrorubens* nor *S. atrovirens*. These 2 species belong in the subgenus *Rosicola*.

Page 1799. In first line under *Sebastodes crameri* for "P. $19\frac{9}{10}$" read "P. 19."

Page 1805. After line 4 add: (*intro*, within; *niger*, black.)

Page 1815. The type of *Sebastodes zacentrus* came from *Albatross* Station 2946, not 2996.

Page 1829. After line 2 insert Subgenus *Sebastosomus*.

Page 1831. Specimens of *Sebastodes taczanowskii* were obtained in 1896 by the *Albatross* at the Kuril Islands, and this species should therefore go in the regular text.

Page 1832. After line 15 insert Subgenus *Sebastomus*.

Page 1833. Before *Sebastodes matsubaræ* (Hilgendorf) insert Subgenus *Zalopyr*.

Page 1833. In fifth line from bottom insert Subgenus *Pteropodus* after "*nebulosus*."

Page 1836. In key at bottom of page, to *a* add: mouth plumbeous within. To *aa* add: mouth black within.

Page 1837, line 2, for "Cardonniera" read "Cardouniera." For "Scorfanudi Funal" read "Scorfana di Funal."

Page 1840. Above "*a*. Breast scaly," insert Parascorpæna (παρά, near; to *Scorpæna*). Before 2236. *Scorpæna agassizii*, Goode & Bean, insert Subgenus *Parascorpæna*, Bleeker.

Page 1850. In last line of synonymy of *Scorpæna mystes* for 1501 read 1601.

LIBRARY
BUREAU OF COMMERCIAL FISHERIES
U. S. FISH AND WILDLIFE SERVICE
P. O. BOX 3830
HONOLULU 12, HAWAII